

Environmental Controls on Isolated Convection during the Amazonian Wet Season

Leandro Alex Moreira Viscardi^{1,2}, Giuseppe Torri², David K. Adams³, and Henrique de Melo Jorge Barbosa^{1,4}

¹Institute of Physics, University of São Paulo, São Paulo, SP, Brazil

²Department of Atmospheric Sciences, University of Hawai‘i at Mānoa, Honolulu, HI, USA

³El Instituto de Ciencias de la Atmósfera y Cambio Climático, Universidad Nacional Autónoma de México, Mexico City, Mexico

⁴Physics Department, University of Maryland Baltimore County, Baltimore, MD, USA

Correspondence: Leandro Viscardi (viscardi@hawaii.edu) and Henrique Barbosa (hbarbosa@umbc.edu)

Abstract.

The Amazon rainforest is a vital component of the global climate system, influencing the hydrological cycle and tropical circulation. However, understanding and modeling the evolution of convection in this region remains a scientific challenge. Here, we assess the environmental conditions associated with shallow, congestus, and isolated deep convection days during the wet season (December to April) employing measurements from the GoAmazon (2014-2015) experiment and large-scale wind field from the constrained variational analysis. Composites of deep days show moister than average conditions below 3 km early in the morning. Analyzing the water budget at the surface through only observations, we estimated the water vapor convergence term as a residual of the water balance closure. Convergence remains nearly zero during the deep days until early afternoon (13 LST), when it becomes a dominant factor in their water budget. At 14 LST, the deep days experience a robust upward large-scale vertical velocity, especially above 4 km, which supports the shallow-to-deep convective transition occurring around 16-17 LST. In contrast, shallow and congestus days exhibit preconvective drier conditions, along with diurnal water vapor divergence and large-scale subsidence that extend from the surface to the lower free troposphere. Moreover, afternoon precipitation exhibits the strongest linear correlation (0.6) with large-scale vertical velocity, nearly double the magnitude observed for other environmental factors, even moisture, at different levels and periods of the day. Precipitation also exhibits a moderate increase with low-level wind shear, while upper-level shear has a relatively minor negative impact on convection.

1 Introduction

In the tropics, deep convection dominates the weather and climate. The formation of cumulus clouds and their evolution into cumulonimbus, and often into mesoscale convective systems (MCS), covers a wide range of spatial and temporal scales. Similarly, complex physical processes from cloud microphysics to gravity wave generation are intrinsically tied to deep convection (Mapes et al., 2006; Mapes and Neale, 2011; Jewtoukoff et al., 2013; Gupta et al., 2023). Representing these convective cloud processes in general circulation models (GCMs) is recognized as a source of bias and uncertainty (Sherwood et al., 2014;

Stevens and Bony, 2013; Ifterly et al., 2018; Maher et al., 2018; Freitas et al., 2020). In particular, the shallow-to-deep (STD) convective transition, whose physical mechanisms are still debated (Kuang and Bretherton, 2006; Khairoutdinov and Randall, 2006; Wu et al., 2009; Waite and Khouider, 2010; Schlemmer and Hohenegger, 2015; Morrison et al., 2022; Barber et al., 2022), has long been problematic for convective parameterizations (Betts, 2002; Betts and Jakob, 2002; Bechtold et al., 2004; Grabowski et al., 2006).

To evaluate model performance and to limit the uncertainty associated with the representation of deep convection in either parameterized or resolved at the cloud-scale, metrics derived from observations are required (Adams et al., 2013, 2017; Schiro et al., 2018; Barber et al., 2022). However, within the deep tropics, data are typically lacking at the resolution needed to evaluate both parameterized convection in GCMs and cloud-resolving models (CRMs), as well as to investigate the physical processes that drive mesoscale convective evolution. Intensive field campaigns, such as the Green Ocean Amazon (GOAmazon) 2014/5 experiment (Martin et al., 2016, 2017), have provided comprehensive measurements from the surface to the clouds, finally providing critical measurements from microphysical scale to large-scale thermodynamic properties critical for studying convection (Giangrande et al., 2016, 2017, 2020; Feng and Giangrande, 2018; Barber et al., 2022).

Convective processes in the Amazon manifest themselves in two primary precipitation modes: isolated convection, which peaks in the late afternoon (around 16-17 LST), and organized convection associated with MCSs, which often peaks during the night into the early morning hours, around 03-04 LST. This distinct cycle is pronounced in the southern (Tota et al., 2000; Machado, 2002; Silva Dias, 2002) and central (Greco et al., 1990; Adams et al., 2013; Ghate and Kollias, 2016; Giangrande et al., 2020) regions. In particular, isolated convection is tied to the diurnal cycle of thermodynamic and dynamical properties (Ghate and Kollias, 2016; Ifterly et al., 2018; Zhuang et al., 2017; Tian et al., 2021). For example, using 3.5 years of water vapor fields from the Amazon Dense Global Navigation Satellite System (GNSS) Meteorological Network, Adams et al. (2013) observed a 4-hr timescale of robust water vapor convergence associated with the STD transition, regardless of convective strength, seasonal variations, or whether convection occurs during day or night. For the vertical moisture distribution, Ghate and Kollias (2016) noted that precipitating days in the dry season start the diurnal cycle by exhibiting a moister layer around 2-5 km. However, this contrasts with later studies, which observed that deep convection days are associated with a moister layer from the surface to mid-levels (~ 5 km) (Zhuang et al., 2017; Tian et al., 2021).

Ifterly et al. (2016) highlighted that convection in the Amazon region has a greater sensitivity to column water vapor, while CAPE turns out to be an inadequate indicator of convection intensity. Further support for these findings comes from the work of Schiro et al. (2016, 2018), who demonstrated the necessity of incorporating deep-inflow mixing to accurately represent the entrainment of dry air in the parcel, which is essential for assessing the sensitivity of convection to instability. Ifterly et al. (2018) reaffirmed the importance of the vertical distribution of humidity for the evolution of convection, but they showed that the relationship between convection and humidity is represented very differently in reanalysis data products. Previous studies have not provided conclusive evidence for the importance of vertical wind shear. For example, Zhuang et al. (2017) proposed that stronger low- and upper-level bulk wind shear predominantly supports the STD transition, but specifically during the dry season (June-September). In contrast, Chakraborty et al. (2018) proposed that increased low-level shear could hinder deep convection, particularly from August to November, especially if it leads to greater entrainment of dry air.

Although the above studies have used a combination of observational and reanalysis data to investigate different aspects of convection, they have not reached a consensus on which variables are most strongly associated with convective triggering or intensity in tropical regions. Here, we investigate the STD transition employing data from the GoAmazon2014/5 rainy season observations. We analyze the diurnal cycle of days designated as shallow, congestus, and deep convection in terms of moisture, instability, and large-scale wind properties to assess the sensitivity of isolated convection in the Amazon to different environmental factors. In addition, we assess the correlation between conditionally averaged precipitation and these environmental factors to further determine their influence on convective intensity.

This paper is organized as follows: First, we present an overview of the GoAmazon2014/5 data and the procedures for convective regime classification are described in Section 2. A comparison of cloud and precipitation properties among the shallow, congestus, and deep days is given in Section 3. Section 4 describes the diurnal cycle of the cloud regimes, while Section 5 presents the analysis of conditionally averaged precipitation based on different environmental controls. A discussion is provided in section 6, and the conclusions are given in section 7.

2 Data and Methods

From January 2014 to December 2015, the GoAmazon2014/5 (Martin et al., 2016, 2017) experiment conducted the first long-term measurements of aerosols, clouds, and atmospheric thermodynamics in the central Amazon region. Nine ground sites and two aircraft were used to examine the environment in and around Manaus, a city ~~border that borders~~ the Rio Negro and surrounded by tropical rainforest. A detailed description of all sites can be found in Martin et al. (2016). This study focuses on the region around the most instrumented site, T3, located 70 km downwind of Manaus, in Manacapuru (3.21°S, 60.60°W), where the AMF-1 and the ARM Mobile Aerosol Observing System (MAOS) were deployed.

2.1 Experimental and large-scale data

To begin with our convective regime classification, described in the next section, we employed ~~the cloud mask~~ a cloud mask based on time-height profiles of the cloud location derived from the Merged RWP-WACR-ARSCL Cloud Mask dataset which combines the Radar Wind Profiler (RWP) data with the original W-band Cloud Radar (WACR) Active Remote Sensing of Cloud (ARSCL) VAP both located at T3 (Giangrande et al., 2017; Feng and Giangrande, 2018). This cloud mask identifies seven types of clouds, three of which correspond to convective clouds: shallow cumulus, with a base and cloud top below 3 km; congestus, with a base below 3 km and a top between 3-8 km; and deep convection, with a base below 3 km and a top > 8 km. From the cloud mask, available every 30 s, we also compute the cloud frequency profile in 12-minute windows to match the S-band radar data described in the following.

The Amazon Protection System in Brazil (Sistema de Proteção da Amazônia, SIPAM), located at the T1 site (3.15°S, 59.99°W), operates an S-band (10 cm), Doppler, single-polarization scanning radar south of downtown Manaus. We used the 12-minute gridded reflectivity product with a horizontal resolution of 2 km and a vertical resolution of 500 m (Schumacher and Funk, 2018a). The S-band domain covers an area with a radius of 202 km. We use reflectivity profiles to assess rain coverage

in our analysis domain, 100x100 km² centered at T3 site, calculated using a reflectivity threshold of 20 dBZ as in Zhuang et al. (2017). The surface precipitation rate is derived from the reflectivity of a constant altitude plan position indicator (CAPPI) at 2.5 km, calibrated with surface measurements from a Joss-Waldvogel disdrometer at the T3 site during the wet season of early 2014 (Schumacher and Funk, 2018b). In examining conditionally averaged precipitation (Section 5), the precipitation rate is averaged over the analysis domain, along with an additional time-average from 14 to 20 LST.

Figure 1 shows the map of S-band precipitation averaged over the wet season (December to April) 2014-2015. The precipitation statistics indicate that some radar beams are partially blocked. We used a threshold of 4 mm day⁻¹ to roughly identify these problematic regions. Beam blockage affects only 7 out of 2,601 pixels in our analysis domain, corresponding to only 0.3%, with no relevant sensitivity to the chosen threshold. Although this contribution is negligible, we removed these blocked pixels from our analysis.




Figure 1. Map of precipitation rate at 2.5 km height averaged over the wet season (December to April) 2014-2015. The square box represents the analysis domain covering an area of 100x100 km² centered at the T3 site. The dotted circle (radius of 202 km) centered over the T1 site describes the S-band radar domain with available measurements.

In addition to rain cover, the analysis of cloud and precipitation properties (section 3) also uses the Aerosol Observing System (AOSMET) surface precipitation measured by the acoustic gauge of a Vaisala WXT520 station (ARM, 2013). A 12-minute average is also applied to match the cloud frequency and S-band radar data.

The balloon-borne sounding system (SONDE) provides the vertical profiles of thermodynamic properties four times per day, at 02, 08, 14, and 20 local standard time (LST) (ARM, 2014f) during the wet season (December-April). For consistency, we only analyze soundings that extend from the surface to at least 8 km, and we linearly interpolate the profiles to a fixed 50 m vertical grid. The planetary boundary layer height is derived from the profile of the bulk Richardson number using a critical threshold of 0.25 (ARM, 2014e). We also calculate additional variables from the thermodynamic profiles. The total column water vapor (CWV) is determined by integrating the [water vapor](#) mixing ratio from the surface to 200 hPa. Similarly,

the partially-integrated CWV is calculated for the layers 1000-850 hPa (\sim 0-1.5 km, CWV_{lower}), 850-700 hPa (\sim 1.5-3 km, CWV_{mid}), and 700-350 hPa (\sim 3-8 km, CWV_{upper}), which are representative of the convective boundary layer, lower free
 110 troposphere, and upper levels, respectively. Convective Inhibition (CIN) and Convective Available Potential Energy (CAPE) are also calculated. For these, a hypothetical air parcel is lifted dry-adiabatically to the Lifting Condensation Level (LCL), then pseudo-adiabatically from there. A mixed-layer parcel from the surface to 100 hPa is used as the initial state for the parcel's ascent Stull (2016).

We used a combination of instruments to characterize the surface water budget. The latent heat flux is taken from the ARM
 115 best-estimate dataset (ARM, 2014a) based on observations from the Eddy Correlation (ECOR) Flux Measurement (ARM, 2014b). Instead of using the S-band radar precipitation, the water balance analysis ~~uses the AOSMET surface precipitation~~ utilizes a combination of rain-gauge source measurements to provide a more robust estimation of the mean surface precipitation and its uncertainties. Specifically, we use the surface AOSMET precipitation (ARM, 2013), tipping bucket (ARM, 2014g), and a laser disdrometer (ARM, 2014c). Here, CWV and liquid water path (LWP) are taken from the microwave radiometer (MWR,
 120 ARM, 2014d). Similarly to Schiro et al. (2016), we exclude data in cases where the brightness temperature surpasses 100 K, and when water accumulates on the MWR lens surface during rainy periods. ~~The water budget derivation, including the water vapor convergence term,~~ Hourly averages are applied for the data utilized in the water budget analysis, and its mathematical derivation is provided in the next section.

Large-scale wind properties and bulk vertical wind shear are analyzed by way of Constrained Variational Analysis (VARANAL)
 125 data developed for the GoAmazon2014/5 experiment (Tang et al., 2016). This large-scale forcing dataset is derived from the European Centre for Medium-Range Weather Forecasts (ECMWF) analysis that is further constrained by S-Band radar precipitation, surface fluxes, and Geostationary Operational Environmental Satellite (GOES-13) radiances. The VARANAL large-scale fields represent an average across an analysis domain centered at the S-band radar location, with a radius of 110 km.

130 Finally, a table summarizing the data is also provided in the supplementary material (table A1).

2.2 Derivation of water budget

The integral form of the continuity equation for the total water r_t can be written as,

$$\frac{\partial}{\partial t} \int_{p_{\text{top}}}^{p_{\text{bottom}}} r_t \frac{dp}{g} + \nabla \cdot \int_{p_{\text{top}}}^{p_{\text{bottom}}} r_t \mathbf{V} \frac{dp}{g} = E - P, \quad (1)$$

where p is pressure, g is gravity acceleration, \mathbf{V} is the horizontal wind vector, and E and P correspond to the water mass fluxes
 135 associated with surface evaporation and precipitation. The total water r_t is the sum of the three different phased-based mixing ratios, i.e., water vapor (r_v), liquid (r_l), and ice (r_i). The second integral is the water mass flux divergence, which is mostly associated with ~~water vapor. As convective clouds develop or dissipate, the time rate of change of total column water—the first integral—greatly contributes to the water mass balance~~ the divergence of water vapor.

For the sake of this analysis, we ~~divide equation by the density of liquid water ρ_l (10^3 kg m^{-3}) and ignore~~ ~~neglect~~ the time
140 variation of the ice term and express all terms in units of mm hr^{-1} . As we show later (section 4.3), the results indicate a minor
contribution of the liquid water term in the water budget, which supports ignoring the ice water term. Thus, the equation can
be rewritten in a simplified form:

$$\frac{\partial}{\partial t} \text{CWV} + \frac{\partial}{\partial t} \text{LWP} - \text{EVAP} + \text{PREC} = \text{CONV}, \quad (2)$$

where EVAP corresponds to the evaporation rate, PREC corresponds to the precipitation rate, and CONV represents the water
145 vapor convergence. Note that we intentionally rearranged the order of the equation to emphasize terms based on GOAmazon
observations on the left-hand side and the residuals, which can only be estimated, on the right-hand side. For this analysis,
we consider only the hourly average of observed variables. Specifically, CWV and LWP are based on the MWR. ECOR latent
heat flux data is utilized for estimating the evaporation term. Precipitation is ~~taken from the AOSMET surface measurements~~
obtained utilizing different sources, namely, aerosol observing system data, tipping bucket, and laser disdrometer (Section 2.1).
150 The water vapor convergence term is estimated using the mean composites for $\partial \text{CWV} / \partial t$, $\partial \text{LWP} / \partial t$, EVAP, and PREC.

Note that we only employed local surface measurements. Surface fluxes depend primarily on the surface properties, which
are uniform around the experimental site, at least up to 10 km away. Surface rain gauges and the microwave radiometer perform
point measurements. However, since we are applying hourly averages for the water budget datasets and the composites average
over many days, the mean precipitation, CWV, and LWP relate to regions associated with the domain average of a cloud size
155 scale. Therefore, our water balance is consistent for a temporal-spatial scale of $\sim 1 \text{ hr}$ and $\sim 10 \text{ km}$.

2.3 Convective regime classification

We define the wet season as the period from December to April following previous studies in the Central Amazon (Machado
et al., 2004; Marengo et al., 2013). To contrast the different atmospheric conditions that lead to different convective regimes, we
apply a classification criterion that identifies shallow, congestus, or deep days. Given our interest in convection that develops
160 in response to the diurnal cycle, days with propagating mesoscale convective systems (MCS) were excluded from our analysis.
This regime classification follows the criteria proposed by Zhuang et al. (2017) and Tian et al. (2021). We first consider the
diurnal period between 10-20 LST. Then, we define three categories of convective regimes: shallow cumulus, congestus, and
deep, based on the maximum development of convective clouds during the diurnal cycle. The cloud mask from the RWP-
WACR-ARSCL is used to identify cloud development throughout each day. In addition, we also employ rain coverage data
165 from the S-band radar to estimate the vertical cloud development based on the echo cloud top. The rain coverage is calculated
as the fraction of reflectivity pixels $> 20 \text{ dBZ}$ (see Section 2.1). The echo-top is defined as the highest level where rain coverage
is greater than 2%. The quantitative criteria are listed below:

- i) Shallow convective days (ShCu): (1) The cloud mask does not show any congestus and deep clouds during the diurnal
cycle (10-20 LST). (2) ~~Precipitation~~ Rain cover above 3 km of altitude must be $< 2\%$ in the analysis domain (100×100
170 km^2 , centered at T3).

- ii) Congestus convective days (Cong): (1) The cloud mask indicates congestus clouds during the diurnal cycle, but none of them develops into deep clouds. (2) ~~Precipitation~~ Rain cover above 8 km (between 3-8 km) of altitude must be $< 2\%$ ($> 2\%$) in the analysis domain.
- iii) Deep convective days (Deep): (1) The cloud mask indicates deep convection during the diurnal cycle. (2) ~~Precipitation~~ Rain cover above 8 km $> 2\%$ in the analysis domain.
- iv) Early morning perturbation: For the 06-10 LST period, we require that no congestus and deep clouds be observed.
- v) Local convection: No convective system with a contiguous area of precipitation $> 10,000 \text{ km}^2$ reaches the analysis domain between 06-20 LST.

Figures 2 and 3 illustrate the diurnal evolution of the occurrence of the different cloud types and rain coverage for the three regimes. The classification essentially depends on the cloud evolution during the diurnal cycle. The exclusion of any relevant early morning disturbance (06-10 LST) associated with important preconvective activity guarantees that nighttime MCSs do not cause significant preconditioning.

The number of days categorized according to the above criteria for the wet season (blue) and dry season (red) is shown in Fig. 4. Deep convection appears to be the dominant category in both seasons, although the ~~convective systems tend to be~~ larger-propagating-convection category occurs more frequently during the wet season. Specifically, we identified 16 days for the ShCu regime, 27 days for the Cong regime, 60 days for the Deep regime, and 111 days for non-local (organized) deep convection. Given our focus on the local shallow-to-deep transition mechanism during the wet season, the results presented in the next section refer only to those 103 days.

3 Cloud and Precipitation Properties

The composite diurnal cycles of the vertical cloud frequency profiles, local surface precipitation rate, and rain coverage for the different convective regimes are shown in Fig. 5. For all convective regimes, daytime convection is usually preceded by some nighttime cloud cover at all levels.

The ShCu regime has a more scattered cloud frequency during the daytime. Low-level clouds dominate the diurnal cycle, with a peak reaching 54% at 1.43 km around 13 LST. After 17 LST, cirrus clouds also contribute to the cloud frequency composite, likely transported from afar. The local surface precipitation rate remains below 0.13 mm hr^{-1} throughout the diurnal cycle, while the rain cover has only a minor contribution of approximately 1% at the end of the day.

The Cong regime also shows a higher cloud frequency below 3 km during the diurnal cycle. The maximum cloud cover is 50% at 1.04 km around 12 LST, earlier and lower than observed for the ShCu regime. Between 3-8 km, the cloud frequency composite has values below 5%. Surface precipitation occurs around 12 LST, corresponding temporally with the period of maximum cloud cover. Nonetheless, the magnitude remains modest, with precipitation values at less than 0.75 mm hr^{-1} throughout the diurnal cycle. Contrary to the local cloud frequency assessed at the T3 site, the domain rain coverage exhibits




Figure 2. Cloud mask (left) and rain coverage (right) for examples of days classified as shallow (ShCu, a-b), congestus (Cong, c-d), and deep (Deep, e-f).

a distinct pattern at mid-levels, indicating values of 1-2%. After the diurnal cycle, the rain coverage composite indicates that congestus clouds may evolve into a deeper phase. We identified 7 cases (26%) in which congestus days fulfilled the Deep criteria (echo top above 8 km) after the diurnal cycle.

205 The Deep regime reveals a more coherent cloud frequency profile associated with more extensive and longer-lived convective clouds during daylight hours. The maximum Deep cloud frequency is also associated with shallow clouds, being 49% at 0.92 km around 11:30 LST. As we discuss next, this is ~~because-associated with~~ both the boundary layer height and the lifting condensation levels ~~are-being~~ lower in the Deep regime. As the day progresses, the Deep cloud frequency also increases throughout the troposphere, with cloud top reaching up to 16 km. After cumulonimbus dissipation (around 18 LST), its anvil structure remains and may become a cirrus cloud that may contribute to the cloud cover of the next day. Surface precipitation typically occurs around noon and in the afternoon as clouds develop. Notably, a substantial rain coverage emerges between
 210 16-17 LST, with the maximum surface precipitation of 3.72 mm hr⁻¹ peaking at 16:24 LST. This occurrence ~~coincides-is~~ consistent with the late afternoon STD transition in the Amazonian wet season.




Figure 3. Convective system. (a) Scattered local system. (b) Non-local propagating system. The dashed black box illustrates the region with a contiguous area of precipitation (> 20 dBZ) not fulfilling the local convection requirement.




Figure 4. Number of days classified in each convective regime during the wet (Dec-Apr) and the dry (Jun-Aug) seasons, from 2014 to 2015. Propagating (Prop) days refer to non-local deep convection, with the early morning perturbation condition being ignored.




Figure 5. Left: Cloud frequency ([cloud counting fraction in %](#), colormap) [as a function of height calculated](#) over 12 minutes window based on the cloud mask and surface precipitation rate (mm/hr, red line) locally measured by the aerosol observing system at the T3 site. Right: Rain cover (%) over the analysis domain (100x100 km², centered at T3) based on the S-band radar. The mean composites distinguish days classified as shallow (a-b, N=16), congestus (c-d, N=27), and deep (e-f, N=60) during the wet season.

4 Environmental Conditions

215 In this section, we specifically evaluate the environmental conditions associated with the STD convective transition. We analyze the local atmospheric conditions, convective properties, surface water balance, and large-scale wind properties associated with shallow, congestus, and deep convective days. The error bars in each figure represent the standard deviation of the composite.

4.1 Atmospheric Conditions

Figure 6 shows the average vertical profiles of potential temperature (Fig. ??(a)), θ , and water vapor mixing ratio (Fig. ??(b)), r_v , for each convective regime at 08 LST. The differences between the Deep and other convective regimes, $\Delta\theta$ and Δr_v , are shown in panels c-d.

The potential temperature differences between the convective regimes are frequently less than 0.5 K below the 8 km level, suggesting no relevant relationship between daytime cloud development and the morning temperature profile. The Deep regime




Figure 6. Atmospheric conditions. (a) Potential temperature (K) and (b) water vapor mixing ratio (g/kg) at 08 LST radiosonde observations. The corresponding convective regime differences (Deep-ShCu and Deep-Cong) for potential temperature ($\Delta\theta$, c) and mixing ratio (Δr_v , d) are also included.

experiences higher moisture content, especially below 3 km. The most remarkable difference occurs around the 2 km level, with Δr_v reaching 1.30 g kg^{-1} and 0.74 g kg^{-1} for Deep-ShCu and Deep-Cong, respectively. Above 3 km, the differences in moisture profile for all regimes is quite similar. have little statistical significance.

These results suggest that the importance of early morning excess humidity to the STD transition in the Amazon is primarily limited to the lower levels, a finding supported by Zhuang et al. (2017); Tian et al. (2021). However, it differs from Ghate and Kollias (2016), which noted an excess of humidity solely above 2 km during precipitating days in the dry season. Moreover, over the these results contrast with studies over tropical oceans, the role of moisture levels changes, with where free-tropospheric humidity playing has been shown to play a more significant role in convective activity, while boundary layer humidity demonstrates minor variability (Bretherton et al., 2004; Holloway and Neelin, 2009).

The column water vapor (CWV) for each radiosonde time is presented in Fig. 7a, while panels b-d show the partial contribution from lower, middle, and upper levels (see Section 2.1). The ShCu regime shows the smaller CWV throughout the day. The difference between Deep and ShCu ranges from 2.3 mm at night to 5.1 mm at 14 LST. Interestingly, CWV for The differences in CWV among Cong and Deep regimes is similar before the diurnal cycle starts, at 2 LST, but is already larger for the Deep by 8 LST. The show little statistical significance from nighttime to early morning, while their difference is maximum at 14 LST when it reaches 2.1 mm.

The $\text{CWV}_{\text{lower}}$ increases from 08 LST to 14 LST for all convective regimes. This increase is possibly due to evapotranspiration, which appears to be the dominant moisture factor during this period (water budget analysis, Section 4.3). A similar, albeit less pronounced, pattern diurnal moistening is also observed in the lower free troposphere CWV_{mid} for all regimes. However, the upper-level CWV is essentially constant for the Deep regime in the 08-14 LST period, while the ShCu and Cong




Figure 7. Column water vapor (CWV, mm) at the radiosonde launch times (02, 08, 14, and 20 LST). (a) Total and its partial contribution in the (b) 1000-850 hPa, (c) 850-700 hPa, and (d) 700-350 hPa layers.

regimes show a decrease of 0.80 and 1.01 mm, respectively. This indicates that the ShCu and Cong regimes might be associated with large-scale subsidence (see Section 4.4), which would explain the drying of the mid-levels, ~~and associated with low-level moisture divergence, which would explain the slower accumulation of moisture below 700 hPa despite slightly larger latent heat~~. Moreover, we note that Deep days show a significant increase in CWV_{upper} from 14 to 20 LST, with a simultaneous decrease of CWV_{lower} . This is likely due to a combination of water vapor convergence (section 4.3) preceding the late afternoon STD transition and the vertical transport of moisture by deep clouds.

4.2 Convective Properties

Figure 8 shows the PBL height and the mixed-layer ~~CIN and CAPE~~ MLCIN and MLCAPE. The PBL height among the convective regimes differs the most at 14 LST, where the convective boundary layer roughly coincides with the LCL or cloud base height (not shown), being about 300 m lower for the Deep (1.09 km) than the ShCu (1.37 km) regime. The most developed convective regimes show a combination of higher MLCAPE and lower MLCIN in the early morning, providing more buoyancy and a lower barrier for convection to develop. ~~In the afternoon, MLCAPE is higher~~ At 14 LST, MLCAPE for the Deep regime (1237 J kg⁻¹), ~~a few hours before the late afternoon STD transition, slightly surpassing the value for the~~ and Cong regime (1111 J kg⁻¹) ~~and significantly exceeding~~ significantly exceed the value for the ShCu regime (671 J kg⁻¹). The change in CAPE-MLCAPE from afternoon to early evening (CAPE-MLCAPE(20 LST) - CAPE-MLCAPE(14 LST)) is negative for the Deep (-390 J kg⁻¹), almost neutral for the Cong (-62 J kg⁻¹), and positive for the ShCu (398 J kg⁻¹) regime. As expected, convection in more advanced stages consumes CAPE-MLCAPE more effectively, reducing atmospheric instability.




Figure 8. (a) Planetary boundary layer (PBL) height (m), (b) 100-hPa mixed-layer convective inhibition (MLCIN, J/kg), and (c) 100-hPa mixed-layer convective available potential energy (MLCAPE, J/kg) at the radiosonde launch times (02, 08, 14, and 20 LST).

260 4.3 Surface Water Balance

To understand how the differences in the surface fluxes change CWV, affecting the STD transition and the accumulated surface precipitation, we analyzed the surface water balance (see Section 2.2).

The water balance results are shown in Fig. 9, with the top panels showing the hourly average rate values (mm/day) and the bottom panels showing the corresponding accumulated values (mm). The separate precipitation and convergence associated with each rain gauge used to estimate the mean surface precipitation and its uncertainty in the water budget results are provided in fig. A1. First, we notice that the ~~variation in ∂_t LWP~~ appears negligible, and it does not contribute significantly to the water budget. ~~While the~~ The daytime of ShCu and Cong days ~~is dominated by~~ shows mostly water vapor divergence. This is primarily due to more significant negative changes in CWV and lower precipitation rates, whereas changes in condensate exert a minor influence on the water budget. On the other hand, the Deep regime exhibits ~~a neutral condition~~ relatively neutral conditions from night to early afternoon. However, the water budget closure requires convergence in the period from 14 to 18 LST. This coincides with the STD transition ~~;~~ and is thus primarily attributed to strong water vapor convergence preceding the late afternoon STD transition (Adams et al., 2013, 2017). At day's end, the ShCu regime is estimated to lose 3.4 mm of water vapor due to divergence, while the Cong regime loses ~~4.3~~ 0.9 mm. In contrast, the Deep regime gains ~~5.4~~ 5.2 mm of water vapor due to convergence.

~~Evaporation approximately balances the water vapor divergence during the~~ For ShCu days, ~~leading to minimal precipitation. On the other hand, the divergence of Cong days~~ relatively high surface evaporation and the absence of precipitation require a strong divergence for water balance closure. The Cong regime shows a significant divergence but is relatively weaker ; resulting in a higher contribution of precipitation, especially in the afternoon compared to the ShCu regime, as their surface evaporation is relatively similar, and congestus precipitation exhibits modest values. For ShCu days, the ~~variation in CWP~~




Figure 9. Surface water balance for (a,d) ShCu regime, (b,e) Cong regime, and (c,f) Deep regime. The upper panels represent the rates of change (mm/day) for water vapor convergence (CONV), evaporation (EVAP), precipitation (PREC), and the time derivatives of column water vapor (∂_t CWV, where $\partial_t = \partial/\partial t$) and liquid water path (∂_t LWP). The lower panels display the accumulated water amounts for each term in the water budget along the day (mm). Note that CWV and LWP changes rely on microwave radiometer observations, evaporation on eddy correlation flux measurements, and precipitation [on is estimated utilizing different sources, namely, the aerosol observing system surface data, tipping bucket, and laser disdrometer](#). The water vapor convergence term is estimated as a residual in the water budget equation [\(Section 2.2\)\(2\)](#).

280 ∂_t CWV tends to be small and negative from nighttime to early morning. Then, it increases and balances around noon. By the end of the day, the accumulation of water vapor [\(fig. 9d\)](#) in the column is negligible. This term is also nearly zero during

Cong days. Conversely, the convergence of vapor and evaporation exceeds the precipitation term on Deep days, resulting in a net accumulation of 1.7 mm in column water vapor, which might ~~affect the convective regimes developing~~ act as a positive feedback for the continuation of nocturnal deep convection into the following day after Deep regime conditions. This particular finding had not been previously addressed in studies utilizing the GoAmazon2014/5 observations.

4.4 Large-Scale Wind Properties

Figure 10 displays the wind speed at 08 LST, 11 LST, and 14 LST for all convective regimes. ~~There is a characteristic low-level jet during the wet season, with a peak between~~ The wind profiles for all regimes peak between the 900-800 hPa layer. ~~Anselmo et al. (2020) also reported an Amazonian, characteristic of a low-level jet ,occurring also reported by Anselmo et al. (2020)~~ , which they observed 10-40% of the time during GoAmazon between March-May 2014-2015. During the morning, the ShCu regime reveals a lower and slightly stronger jet. However, the PBL grows during the day to a height of 1-2 km (see Fig. 8), reaching higher altitudes for ShCu days, and the mixing of free-tropospheric and PBL momentum potentially reduces the wind speed. As a result, the lower jet in the ShCu regime is more significantly affected by the PBL growth. Thus, ~~at both 11 LST or 14 LST, the Deep regime presents a more vigorous and slightly higher~~ ShCu regime reveals a less prominent ~~jet wind peak but at a slightly higher altitude compared to Deep days~~. For the mid- and upper-levels, between ~~600-300~~ 600-350 hPa layer, the ShCu regime shows an additional upper-level jet maxima, while the Cong and Deep regimes exhibit weaker and comparable wind speeds.




Figure 10. Large-scale horizontal wind speed (m/s) at (a) 08 local standard time (LST), (b) 11 LST, and (c) 14 LST for ShCu, Cong, and Deep regimes.

A hodograph at 14 LST is displayed in Fig. 11. The wind turns clockwise with increasing height: north-easterly winds dominate in the boundary layer, approximately below 800 hPa, while easterly winds dominate in the \sim 700-500 hPa layer. The ~~only most~~ only most notable difference in wind direction only most occurs in the upper troposphere, above 400 hPa. Particular, on the Deep days,

the wind stops veering, demonstrating a consistent south-easterly direction. Since the wind profiles at low- and mid-levels are comparable among the convective regimes, the veering of the wind alone only hints at a possible control mechanism for the development from the congestus to the deep phase.




Figure 11. Hodograph representing the u (m/s) and v (m/s) large-scale wind components as a function of pressure at 14 LST for (a) ShCu regime, (b) Cong regime, and (c) Deep regime.

Figure 12 contains the large-scale vertical velocity (ω in hPa hr^{-1}) from the variational analysis at 08 LST, 11 LST, and 305 14 LST. In the early morning, the Deep regime shows moderate upward large-scale vertical velocity below 700 hPa, relatively greater than those associated with the ShCu and Cong regimes. Above 700 hPa, ω becomes positive (subsidence) and exhibits comparable values among the convective regimes. Around noon (11 LST), the Deep regime is dominated by upward large-scale vertical velocity below 600 hPa, while the ShCu and Cong regimes exhibit similar ω profiles with subsidence dominating above the surface. In the afternoon, significant differences are observed between the convective regimes. The Deep regime is 310 dominated by upward large-scale vertical velocity, especially above the 800 hPa level. The ShCu and Cong regimes exhibit important subsidence in the low levels, particularly below 700 hPa, with ShCu-subsidence assuming higher magnitudes. Tian et al. (2021) also reported a significant disparity in the large-scale vertical velocity between Deep and ShCu days, especially in the afternoon and early evening. ~~However, solely through the differences in ω convective profiles presented in their results, it is not possible to adequately determine the levels associated with subsidence on ShCu days.~~

315 To evaluate the vertical wind shear, we used the bulk wind shear, which is defined as the magnitude of the vector difference of the wind at two levels. Figure 13 shows the vertical bulk wind shear for the layers 0-2 km, 0-4 km, and 0-6 km. Additionally, the bulk shear from the level of maximum wind speed below 4 km and the associated pressure level is provided in fig. A2. The 0-2 km layer exhibits a greater dependence on the diurnal cycle, with the Deep days followed by Cong days showing the most substantial wind shear at any time. ~~This result suggests~~ Moreover, the Deep regime shows the largest difference between 320 the maximum wind speed below 4 km and surface wind. These results suggest that low-level vertical wind shear is related to




Figure 12. Large-scale vertical velocity (ω in hPa/hr) at (a) 08 LST, (b) 11 LST, and (c) 14 LST for ShCu, Cong, and Deep regimes.

convection development. For the 0-4 km layer, the wind shear is more similar between the convective regimes. For the 0-6 km layer, only ShCu days exhibit a distinct wind shear, which is more robust and has a standard deviation that can reach more than 1 m s^{-1} . **However, this layer has an extension at least twice that of the cloud top of the shallow cumulus ($< 3 \text{ km}$), suggesting that mid-level wind shear might play a minor role in the STD transition.**




Figure 13. Vertical bulk wind shear (m/s) for the layers (a) 0-2 km (sfc-790 hPa), (b) 0-4 km (sfc-615 hPa), and (c) 0-6 km (sfc-465 hPa).

325 5 Conditionally Averaged Precipitation

In this section, the conditionally averaged precipitation is evaluated as a function of the main variables presumed to control the STD transition. From the previous section, we observed that convective development is primarily associated with humidity at lower levels in the early morning, while in the afternoon, it is more strongly related to upper-level moisture, instability, large-scale vertical velocity, and vertical wind shear.

330 Here, we explore the precipitation response to moisture using low-level and lower free troposphere CWV at 08 LST, upper-level CWV at 14 LST, MLCAPE at 14 LST, mean vertical velocity in the 1000-700 hPa and 700-300 hPa layers at 14 LST, and bulk shear magnitude for the 0-2 and 0-6 km layers at 14 LST as surrogates for vertical wind shear. Precipitation from S-band radar is averaged over the analysis domain ($100 \times 100 \text{ km}^2$ centered at T3) and from 14 to 20 LST. The conditionally averaged precipitation corresponds to the average precipitation observed within distinct bins for each variable, which were
335 defined as 1 mm for CWV, 2.5 J g^{-1} (250 J kg^{-1}) for MLCAPE, 1 hPa hr^{-1} for vertical velocity, and 2 m s^{-1} for bulk shear magnitude. In this analysis, it is important to highlight that we group the ShCu, Cong, and Deep regimes together and consider them collectively as local convective days.

Figure 14 consists of the conditionally averaged precipitation analysis. Large-scale afternoon vertical velocity exhibits the strongest correlation with precipitation: -0.596 in the 700-300 hPa layer and -0.570 in the 1000-700 hPa layer. Above 3 km
340 (700-300 hPa), precipitation increases significantly from nearly zero to approximately 0.8 mm hr^{-1} as vertical motion rises from nearly zero to about -9 hPa hr^{-1} . These correspond to the heaviest conditionally averaged precipitation rates. Interestingly, vertical wind shear, especially the low-level (0-2 km) bulk shear magnitude, exhibits the second strongest correlation with precipitation, although it has a modest value (0.348).

The lower free troposphere CWV at 08 LST demonstrates a similar correlation (0.311) with precipitation. The correlations
345 for low-level CWV at 08 LST (0.274) and upper-level CWV at 14 LST (0.254) are also comparable. Vertical wind shear at higher levels (0-6 km layer) exhibits a weak negative correlation (-0.099) with precipitation. On the other hand, bulk shear in the 0-4 km layer (0.016) and MLCAPE (0.055) exhibit the smallest correlation values, which increase only weakly with precipitation. These results suggest that CAPE is not a good indicator of ~~convection-precipitation~~ in the Amazon, which is consistent with the findings of Iterly et al. (2016); Schiro et al. (2018). Thus, days starting with excess water vapor in the
350 lowest 3 km of the troposphere and exhibiting relatively stronger large-scale vertical velocity and low-level wind shear in the afternoon, have the highest probability of developing deep afternoon convection accompanied by heavy precipitation.

6 Discussion

The previous section presented the wet-season composites of ShCu, Cong, and local Deep convection days for both environmental and cloud properties measured during the GoAmazon2014/5 Campaign. ~~Results showed that isolated deep convection
355 is associated with more extensive and longer-lived clouds throughout the diurnal cycle.~~The cumulonimbus clouds can extend up to 16 km of altitude, and precipitation peaks around 16-17 LST, associated with the STD transition. This is consistent with previous studies in the southern Amazon. For example, Tota et al. (2000) and Machado (2002) analyzed data from the Wet




Figure 14. Conditionally average-averaged precipitation (mm/hr) to (a) low-level column water vapor (CWV, mm) at 08 LST, (b) lower free troposphere CWV at 08 LST, (c) upper levels CWV at 14 LST, (d) 100-hPa mixed-layer convective available potential energy (MLCAPE, J/g) at 14 LST, mean large-scale vertical velocity (ω , hPa/hr) in the (e) 1000-700 hPa and (f) 700-300 hPa layers at 14 LST, and bulk wind shear magnitude (m/s) in the layers (g) 0-2 km and (h) 0-6 km at 14 LST. Precipitation corresponds to the mean S-band radar precipitation from 14 to 20 LST. The bins for averaging each variable are indicated by horizontal bars, which represent 1 mm for CWV, 2.5 J g^{-1} (250 J kg^{-1}) for MLCAPE, 1 hPa hr^{-1} for vertical velocity, and 2 m s^{-1} for wind shear. The conditional averaging analysis is conducted for local convective days (ShCu, Cong, and Deep regimes) indicated by green square markers. Additionally, linear least-squares regression and Pearson correlation are included for each analysis.

Season Atmospheric Mesoscale Campaign (WETAMC), from January to February 1999, and reported two main precipitation modes: isolated convection, which peaks in the afternoon around 16 LST, and organized convection associated with mesoscale convective systems, with a maximum during the night around 04 LST. The diurnal peak associated with deep convection is also documented more recently in Ghate and Kollias (2016) and Tian et al. (2021), who also used the GoAmazon2014/5 observations but did not focus on the wet season as in our study.

Our results show that deep days are associated with moister conditions in the early morning, but only in the lower troposphere, particularly below 3 km. ~~Based on satellite-based observations, radiosonde data, and surface turbulent heat flux measurements distributed across the Amazon Basin before the GoAmazon2014/5 experiment took place, Iitterly et al. (2016)~~ This contrasts with the results from Iitterly et al. (2016). They also noted significant differences between morning atmospheric conditions and convective intensity. ~~However, , but indicated that~~ humidity in the upper ~~and lower troposphere showed the strongest troposphere also exhibits a strong~~ relationship with convective intensity. The work by Ghate and Kollias (2016) focused on GoAmazon2014/5 observations during the dry season and observed the presence of an early morning moist layer, but elevated, between 2 and 5 km. In contrast, Zhuang et al. (2017) and Tian et al. (2021) found that days with deep convection exhibit increased moisture levels extending from the surface to mid-levels, regardless of the season. The discrepancies in the anomaly of the early morning moisture layer with convective intensity among these studies are likely attributed to the data and procedures used to identify convective days. Only Ghate and Kollias (2016); Zhuang et al. (2017); Tian et al. (2021) specifically utilized the GoAmazon2014/5 observations. While Ghate and Kollias (2016) differentiated between days with no precipitation (cumulus days) and those with surface precipitation rates exceeding 0.05 mm h^{-1} (precipitating days), both Zhuang et al. (2017) and Tian et al. (2021) relied on cloud boundaries, albeit with different definitions for identifying convective cloud regimes. Furthermore, all these studies relied on visual inspection to identify and eliminate cases linked to MCSs. Here, we have specifically developed a consistent systematic criterion (item v, section 2.3) based on the contiguous area of the S-band radar precipitation to identify these days and remove them from our analysis. These differences in methods contribute to differences in the selection of convective days. Particularly in Ghate and Kollias (2016), the identification of convective days based on a single surface precipitation threshold might not be distinguishing the varying levels of convective intensity observed in the Amazon. Additionally, relying solely on site measurements ignores the surrounding areas, which could be important for gaining a broader perspective on the convective intensity of a given day. This could explain why only Ghate and Kollias (2016) observed a moist layer extending from 2 km rather than from the surface.

Based on the water vapor convergence derived from the surface water balance, we observed that during ShCu and Cong days, there is a predominant water vapor divergence. In contrast, the Deep regime days are neutral until early afternoon and show a significant water vapor convergence between 14 and 18 LST, coinciding with the STD transition. This aligns well with the convergence timescale preceding the STD transition reported by Adams et al. (2013), based solely on CWV observations. More recently, Zhuang et al. (2017) assessed the accumulated water for the convergence term through diagnostic data derived from ECMWF model runs designed for the GoAmazon2014/5 experiment. During the wet season, they found neutral divergence during shallow days, while water vapor roughly increased linearly throughout the deep convective days. Although Zhuang et al. (2017) and our findings indicate a dominance of water vapor convergence on Deep days, the behavior differs significantly in our analysis, as we observed a relevant divergence during the daytime of ShCu days, and the convergence dominates Deep days only in the afternoon. These differences suggest that the ECMWF model runs may not have accurately ~~capture~~ captured the diurnal cycle of convection, misrepresenting advection and convergence. This is a well-known problem of models that rely on convective parameterizations (Betts, 2002; Betts and Jakob, 2002; Grabowski et al., 2006), and that is why the variational

analysis dataset assimilated the S-Band radar precipitation and surface fluxes from GoAmazon2014/5 experiment (Tang et al., 2016).

We observed that mixed-layer CAPE is lower on ShCu days and higher on Deep days, with the difference increasing from
400 nighttime (2 LST) to early afternoon (14 LST). However, when the conditionally averaged precipitation is considered, we
found that $\frac{\text{CAPE}}{\text{MLCAPE}}$ is only weakly correlated with afternoon precipitation. Iitterly et al. (2016) also found that CAPE
is an unreliable indicator of convective activity in the Amazon. Moreover, Schiro et al. (2018) investigated the relationship
between buoyancy and precipitation and showed that CAPE is weakly related to precipitation when deep-inflow mixing is not
included that represents the entrainment of dry air in the parcel.

405 For the large-scale wind, we observed that precipitation increases significantly with afternoon upward vertical velocity,
especially above 3 km, and moderately with increasing low-level wind shear (0-2 km). At upper levels, vertical wind shear
has a negative impact on convection, although its correlation with precipitation is relatively weak compared to low-level shear.
Using GoAmazon2014/5 observations, Zhuang et al. (2017) reported a similar pattern of bulk wind shear during the wet season.
However, their interpretation differs from ours. They suggested that strong wind shear would favor convection. Nevertheless,
410 during the wet and transition seasons, Deep days are associated with weaker upper-level shear. Thus, they suggested that
wind shear may have no impact or could even hinder convection. Here, we have shown that low-level shear has the greatest
correlation with convection. On the other hand, Chakraborty et al. (2018) suggested that shallow convection is associated with
stronger low-level and weaker upper-level shear intensity during the transition season, which is the opposite of what was shown
by Zhuang et al. (2017) and observed in our study focused on the wet season. This difference could be attributed to variations
415 in the data employed, procedures for identifying convective clouds, and potentially because Chakraborty et al. (2018) did not
evaluate the diurnal cycle. Instead, they considered radiosonde wind measurements within 2 hours before the development of
shallow convection, in scenarios where it did or did not evolve into deep convection. On the other hand, Zhuang et al. (2017)
used radar wind profiler data to specifically assess low-level wind shear. Moreover, our findings generally agree with Tian et al.
(2021) that mid-tropospheric vertical wind shear could significantly suppress the vertical development of convection, although
420 to a lesser extent compared to low-level wind shear.

7 Conclusions

We analyzed measurements from the GoAmazon2014/5 field campaign in the central Amazon with the goal of assessing possible
controlling mechanisms of the shallow-to-deep convective transition. We classified wet season days into shallow (ShCu),
congestus (Cong), and Deep regimes, purposely excluding mesoscale systems to focus on locally-driven deep convection. Un-
425 like previous studies, we used an objective and reproducible method to identify and exclude days dominated by organized
convection. The Deep regime is characterized by early morning moister conditions extending from the surface to the lower free
troposphere. This agrees with some previous studies (Zhuang et al., 2017; Tian et al., 2021), but differs from others (Ghate
and Kollias, 2016). We also found that ShCu and Cong days are characterized by water vapor divergence, while Deep days
start neutrally (from 8 to 13 LST) and develop strong water vapor convergence in the afternoon (from 13 to 17 LST), which

430 peaks coincide with the STD transition (Adams et al., 2013). This is the first analysis of the moisture convergence using GoAmazon2014/5 observations, as previous studies relied on models that use parameterized convection (Zhuang et al., 2017).

In contrast to several previous studies that emphasized the role of humidity in convective activity in the Amazon region (Itterly et al., 2016; Ghate and Kollias, 2016; Schiro et al., 2016), our results indicate that precipitation may exhibit an even stronger correlation with large-scale vertical velocity than with the moisture content in any level or at any time of the day. 435 Nevertheless, it should be noted that ~~this variable~~ vertical velocity is challenging to assess through observations. Furthermore, we have found that precipitation increases with low-level wind shear, displaying a correlation comparable to that observed for moisture content. This correlation has not been evaluated by previous works, which even diverged in their interpretation of the wind shear's role on the STD transition (e.g., Zhuang et al., 2017 and Chakraborty et al., 2018).

Finally, our results suggest that preconvective humidity in the lower troposphere and diurnal large-scale vertical velocity, wa- 440 ter vapor convergence, and low-level wind shear represent the primary environmental factors influencing convective intensity. This indicates that dynamic factors may play a more prominent role in convection during the Amazon wet season than during others. To further disentangle the roles of these environmental controls, we recommend conducting numerical experiments using cloud-resolving models that could, for instance, examine the sensitivity of the STD transition to changes in moisture or wind shear at different atmospheric levels. While numerous studies have explored these recent observations over the Amazon, 445 only a few have utilized high-resolution simulations to investigate the environmental controls of convection (e.g., Cecchini et al., 2022). The VARANAL large-scale data could be used to force cloud-resolving models or even directly evaluate the water budget (but note that it operates on a different spatial and temporal scale than the one analyzed in this study). Likewise, longer-term, high-density observational networks in the Amazon, such as Adams et al. (2015), would be of great value for constraining or evaluating numerical model results.

450 *Data availability.* The observational data used in this research are publicly available at <https://www.arm.gov/research/campaigns/amf2014goamazon>. The variational analysis large-scale forcing data for the GoAmazon2014/5 experiment (Tang et al., 2016) is available at the ARM Archive: http://iop.archive.arm.gov/arm-iop/0eval-data/xie/scm-forcing/iop_at_mao/.

Appendix A: Supplemental Material




Figure A1. Accumulated surface precipitation (mm) for the aerosol observing system surface data, tipping bucket, and laser disdrometer are shown in the upper panels (a-c). The corresponding convergence (mm) term for each instrument is displayed in the lower panels (d-f). The ShCu, Cong, and Deep regime results are shown in the first, second and third columns, respectively.




Figure A2. (a) Vertical bulk wind shear from the level of maximum wind speed below 4 km and (b) the associated pressure level.

Table A1. Data

<u>Name</u>	<u>Description</u>
<u>Cloud mask (Giangrande et al., 2017)</u>	<u>Combines multiple sources of cloud data to provide a high-resolution temporal (30 sec) vertical profile of cloud type, including shallow, congestus, and deep. Cloud frequency profile is also calculated as the cloud counting fraction over 12 minutes.</u>
<u>S-band radar (Schumacher and Funk, 2018a, b)</u>	<u>Volume data of reflectivity and derived precipitation rate at 2 km. Rain cover is calculated as the fraction of reflectivity > 20 dBZ over a 100x100 km² analysis domain. These data have a temporal resolution of 12 minutes, a horizontal resolution of 2 km, and a vertical resolution of 500 m.</u>
<u>Surface precipitation</u>	<u>We employ three source of surface precipitation data, namely, aerosol observing system surface data, tipping bucket, and laser disdrometer. They have only a temporal dimension, for which we use an average over 12 minutes or 1 hour.</u>
<u>Atmospheric State</u>	<u>Radiosonde data has a latency of 6 hours (launches at 02, 08, 14, and 20 LST). We utilized sounding profiles covering at least 8 km of the atmosphere. A derived planetary boundary layer height is also used. Additional variables, such as CWV, MLCAPE, and MLCIN are calculated.</u>
<u>Surface fluxes</u>	<u>Hourly average data from the ARM best-estimate dataset based on observations from the Eddy Correlation Flux Measurement. Surface evaporation is calculated and employed in the water budget analysis.</u>
<u>Water content</u>	<u>Hourly average data of CWV and LWP taken from the microwave radiometer are employed in the water budget analysis.</u>
<u>Large-scale wind field</u>	<u>Based on the variational analysis. It corresponds to a 3-hour average and a domain average of ~100 km.</u>

455 *Author contributions.* Leandro Viscardi: Investigation, Methodology, Data curation, Formal analysis, Visualization, Writing - Original draft preparation. Giuseppe Torri: Investigation, Validation, Resources, Writing - review & editing. David Adams: Investigation, Writing - review & editing. Henrique Barbosa: Conceptualization, Investigation, Resources, Supervision, Writing - review & editing.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We acknowledge the data from the Atmospheric Radiation Measurement (ARM) Program sponsored by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Climate and Environmental Sciences Division.

460 L.A.M.V. acknowledges the Brazilian National Council for Scientific and Technological Development (CNPq) graduate fellowship (grant number 148652/2019-0) for supporting his doctoral studies at the University of São Paulo, Brazil. Additionally, gratitude is extended to the Coordination for the Improvement of Higher Education Personnel (CAPES) fellowship (grant number 88887.571091/2020-00) for supporting 6 months of a visiting graduate student program at the University of Hawai‘i at Mānoa, United States, during his doctoral journey.

465 H.M.J.B acknowledges support by the U.S. Department of Energy, Office of Science, Biological and Environmental Research program under Award Number DE-SC-0023058.

References

- Adams, D. K., Gutman, S. I., Holub, K. L., and Pereira, D. S.: GNSS observations of deep convective time scales in the Amazon, *Geophysical Research Letters*, 40, 2818–2823, <https://doi.org/https://doi.org/10.1002/grl.50573>, 2013.
- Adams, D. K., Barbosa, H. M. J., and Gaitán De Los Ríos, K. P.: A Spatiotemporal Water Vapor–Deep Convection Correlation Metric Derived
470 from the Amazon Dense GNSS Meteorological Network, *Monthly Weather Review*, 145, 279–288, <https://doi.org/10.1175/MWR-D-16-0140.1>, 2017.
- Adams, D. K., Fernandes, R. M. S., Holub, K. L., Gutman, S. I., Barbosa, H. M. J., Machado, L. A. T., Calhlheiros, A. J. P., Bennett, R. A., Kursinski, E. R., Sapucci, L. F., DeMets, C., Chagas, G. F. B., Arellano, A., Filizola, N., Rocha, A. A. A., Silva, R. A., Assuncao, L. M. F., Cirino, G. G., Pauliquevis, T., Portela, B. T. T., Sa, A., De Sousa, J. M., and Tanaka, L. M. S.: THE AMAZON DENSE GNSS METEOROLOGICAL NETWORK A New Approach for Examining Water Vapor and Deep Convection Interactions in the Tropics, *BULLETIN OF THE AMERICAN METEOROLOGICAL SOCIETY*, 96, 2151–2165, <https://doi.org/10.1175/BAMS-D-13-00171.1>, 2015.
475
- Anselmo, E. M., Schumacher, C., and Machado, L. A. T.: The Amazonian Low-Level Jet and Its Connection to Convective Cloud Propagation and Evolution, *Monthly Weather Review*, 148, 4083 – 4099, <https://doi.org/https://doi.org/10.1175/MWR-D-19-0414.1>, 2020.
- ARM, Atmospheric Radiation Measurement.: Meteorological Measurements associated with the Aerosol Observing System (AOSMET). 2013-12-12 to 2015-12-01, ARM Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; MAOS (S1), <https://doi.org/10.5439/1984920>,
480 2013.
- ARM, Atmospheric Radiation Measurement.: ARM Best Estimate Data Products (ARMBEATM). 2014-01-01 to 2015-12-31, ARM Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; AMF1 (M1), <https://doi.org/10.5439/1333748>, 2014a.
- ARM, Atmospheric Radiation Measurement.: Eddy Correlation Flux Measurement System (30ECOR). 2014-04-03 to 2015-12-01, ARM
485 Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; AMF1 (M1), <https://doi.org/10.5439/1025039>, 2014b.
- ARM, Atmospheric Radiation Measurement.: Laser Disdrometer (LD). 2014-09-24 to 2015-08-13, ARM Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; Supplemental Site (S10), <https://doi.org/10.5439/1973058>, 2014c.
- ARM, Atmospheric Radiation Measurement.: Atmospheric Radiation Measurement (ARM) user facility. 2014. Microwave Radiometer (MWRLOS). 2014-01-06 to 2015-12-01, ARM Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; AMF1 (M1),
490 <https://doi.org/10.5439/1046211>, 2014d.
- ARM, Atmospheric Radiation Measurement.: Planetary Boundary Layer Height (PBLHTSONDE1MCFARL). 2014-01-01 to 2015-12-01, ARM Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; AMF1 (M1), <https://doi.org/10.5439/1150253>, 2014e.
- ARM, Atmospheric Radiation Measurement.: Balloon-Borne Sounding System (SONDEWNP). 2014-01-01 to 2015-12-01, ARM Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; AMF1 (M1), <https://doi.org/https://doi.org/10.5439/1595321>, 2014f.
- 495 ARM, Atmospheric Radiation Measurement.: Rain Gauge (RAINTB). 2014-10-14 to 2015-07-03, ARM Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; Supplemental Site (S10), <https://doi.org/10.5439/1224827>, 2014g.
- Barber, K. A., Burleyson, C. D., Feng, Z., and Hagos, S. M.: The Influence of Shallow Cloud Populations on Transitions to Deep Convection in the Amazon, *Journal of the Atmospheric Sciences*, 79, 723–743, <https://doi.org/10.1175/jas-d-21-0141.1>, 2022.
- Bechtold, P., Chaboureau, J.-P., Beljaars, A., Betts, A. K., Köhler, M., Miller, M., and Redelsperger, J.-L.: The simulation of the diurnal
500 cycle of convective precipitation over land in a global model, *Quarterly Journal of the Royal Meteorological Society*, 130, 3119–3137, <https://doi.org/https://doi.org/10.1256/qj.03.103>, 2004.

- Betts, A. K.: Evaluation of the diurnal cycle of precipitation, surface thermodynamics, and surface fluxes in the ECMWF model using LBA data, *Journal of Geophysical Research*, 107, <https://doi.org/10.1029/2001jd000427>, 2002.
- Betts, A. K. and Jakob, C.: Study of diurnal cycle of convective precipitation over Amazonia using a single column model, *Journal of Geophysical Research: Atmospheres*, 107, ACL 25–1–ACL 25–13, <https://doi.org/10.1029/2002jd002264>, 2002.
- Bretherton, C. S., Peters, M. E., and Back, L. E.: Relationships between Water Vapor Path and Precipitation over the Tropical Oceans, *Journal of Climate*, 17, 1517 – 1528, [https://doi.org/https://doi.org/10.1175/1520-0442\(2004\)017<1517:RBWVPA>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0442(2004)017<1517:RBWVPA>2.0.CO;2), 2004.
- Cecchini, M. A., de Bruine, M., Vilà-Guerau de Arellano, J., and Artaxo, P.: Quantifying vertical wind shear effects in shallow cumulus clouds over Amazonia, *Atmospheric Chemistry and Physics*, 22, 11 867–11 888, <https://doi.org/10.5194/acp-22-11867-2022>, 2022.
- Chakraborty, S., Schiro, K. A., Fu, R., and Neelin, J. D.: On the role of aerosols, humidity, and vertical wind shear in the transition of shallow-to-deep convection at the Green Ocean Amazon 2014/5 site, *Atmospheric Chemistry and Physics*, 18, 11 135–11 148, <https://doi.org/10.5194/acp-18-11135-2018>, 2018.
- Feng, Z. and Giangrande, S.: Merged RWP-WACR-ARSCL Cloud Mask and Cloud Type, <https://doi.org/10.5439/1462693>, artwork Size: 1 GB Pages: 1 GB Type: dataset, 2018.
- Freitas, S. R., Putman, W. M., Arnold, N. P., Adams, D. K., and Grell, G. A.: Cascading Toward a Kilometer-Scale GCM: Impacts of a Scale-Aware Convection Parameterization in the Goddard Earth Observing System GCM, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2020gl087682>, 2020.
- Ghate, V. P. and Kollias, P.: On the Controls of Daytime Precipitation in the Amazonian Dry Season, *Journal of Hydrometeorology*, 17, 3079–3097, <https://doi.org/10.1175/jhm-d-16-0101.1>, 2016.
- Giangrande, S. E., Toto, T., Jensen, M. P., Bartholomew, M. J., Feng, Z., Protat, A., Williams, C. R., Schumacher, C., and Machado, L.: Convective cloud vertical velocity and mass-flux characteristics from radar wind profiler observations during GoAmazon2014/5, *J. Geophys. Res.*, 121, 12,891–12,913, 2016.
- Giangrande, S. E., Feng, Z., Jensen, M. P., Comstock, J. M., Johnson, K. L., Toto, T., Wang, M., Burleyson, C., Bharadwaj, N., Mei, F., Machado, L. A. T., Manzi, A. O., Xie, S., Tang, S., Dias, M. A. F. S., de Souza, R. A. F., Schumacher, C., and Martin, S. T.: Cloud characteristics, thermodynamic controls and radiative impacts during the Observations and Modeling of the Green Ocean Amazon (GoAmazon2014/5) experiment, *Atmospheric Chemistry and Physics*, 17, 14 519–14 541, <https://doi.org/10.5194/acp-17-14519-2017>, 2017.
- Giangrande, S. E., Wang, D., and Mechem, D. B.: Cloud regimes over the Amazon Basin: perspectives from the GoAmazon2014/5 campaign, *Atmospheric Chemistry and Physics*, 20, 7489–7507, <https://doi.org/10.5194/acp-20-7489-2020>, 2020.
- Grabowski, W. W., Bechtold, P., Cheng, A., Forbes, R., Halliwell, C., Khairoutdinov, M., Lang, S., Nasuno, T., Petch, J., Tao, W.-K., Wong, R., Wu, X., and Xu, K.-M.: Daytime convective development over land: A model intercomparison based on LBA observations, *Quarterly Journal of the Royal Meteorological Society*, 132, 317–344, <https://doi.org/10.1256/qj.04.147>, 2006.
- Greco, S., Swap, R., Garstang, M., Ulanski, S., Shipham, M., Harriss, R., Talbot, R., Andreae, M., and Artaxo, P.: Rainfall and surface kinematic conditions over central Amazonia during ABLE 2B, *Journal of Geophysical Research: Atmospheres*, 95, 17 001–17 014, <https://doi.org/10.1029/JD095iD10p17001>, 1990.
- Gupta, A. K., Deshmukh, A., Waman, D., Patade, S., Jadav, A., Phillips, V. T. J., Bansemmer, A., Martins, J. A., and Gonçalves, F. L. T.: The microphysics of the warm-rain and ice crystal processes of precipitation in simulated continental convective storms, *Communications Earth & Environment*, 4, <https://doi.org/10.1038/s43247-023-00884-5>, 2023.
- Holloway, C. E. and Neelin, J. D.: Moisture Vertical Structure, Column Water Vapor, and Tropical Deep Convection, *Journal of the Atmospheric Sciences*, 66, 1665 – 1683, <https://doi.org/https://doi.org/10.1175/2008JAS2806.1>, 2009.

- 540 Itterly, K. F., Taylor, P. C., Dodson, J. B., and Tawfik, A. B.: On the sensitivity of the diurnal cycle in the Amazon to convective intensity, *Journal of Geophysical Research: Atmospheres*, 121, 8186–8208, <https://doi.org/10.1002/2016jd025039>, 2016.
- Itterly, K. F., Taylor, P. C., and Dodson, J. B.: Sensitivity of the Amazonian Convective Diurnal Cycle to Its Environment in Observations and Reanalysis, *Journal of Geophysical Research: Atmospheres*, 123, <https://doi.org/10.1029/2018jd029251>, 2018.
- Jewtoukoff, V., Plougonven, R., and Hertzog, A.: Gravity waves generated by deep tropical convection: Estimates from
545 balloon observations and mesoscale simulations, *Journal of Geophysical Research: Atmospheres*, 118, 9690–9707, <https://doi.org/https://doi.org/10.1002/jgrd.50781>, 2013.
- Khairoutdinov, M. and Randall, D.: High-Resolution Simulation of Shallow-to-Deep Convection Transition over Land, *Journal of the Atmospheric Sciences*, 63, 3421–3436, <https://doi.org/10.1175/jas3810.1>, 2006.
- Kuang, Z. and Bretherton, C. S.: A Mass-Flux Scheme View of a High-Resolution Simulation of a Transition from Shallow to Deep Cumulus
550 Convection, *Journal of the Atmospheric Sciences*, 63, 1895–1909, <https://doi.org/10.1175/jas3723.1>, 2006.
- Machado, L., Laurent, H., Dessay, N., and Miranda, I.: Seasonal and diurnal variability of convection over the Amazonia: A comparison of different vegetation types and large scale forcing, *THEORETICAL AND APPLIED CLIMATOLOGY*, 78, 61–77, <https://doi.org/10.1007/s00704-004-0044-9>, 2004.
- Machado, L. A. T.: Diurnal march of the convection observed during TRMM-WETAMC/LBA, *Journal of Geophysical Research*, 107,
555 <https://doi.org/10.1029/2001jd000338>, 2002.
- Maher, P., Vallis, G. K., Sherwood, S. C., Webb, M. J., and Sansom, P. G.: The Impact of Parameterized Convection on Climatological Precipitation in Atmospheric Global Climate Models, *Geophysical Research Letters*, 45, 3728–3736, <https://doi.org/10.1002/2017gl076826>, 2018.
- Mapes, B. and Neale, R.: Parameterizing Convective Organization to Escape the Entrainment Dilemma, *Journal of Advances in Modeling Earth Systems*, 3, <https://doi.org/https://doi.org/10.1029/2011MS000042>, 2011.
- Mapes, B., Tulich, S., Lin, J., and Zuidema, P.: The mesoscale convection life cycle: Building block or prototype for large-scale tropical waves?, *Dynamics of Atmospheres and Oceans*, 42, 3–29, <https://doi.org/10.1016/j.dynatmoce.2006.03.003>, 2006.
- Marengo, J. A., Alves, L. M., Soares, W. R., Rodriguez, D. A., Camargo, H., Riveros, M. P., and Pabló, A. D.: Two Contrasting Severe Seasonal Extremes in Tropical South America in 2012: Flood in Amazonia and Drought in Northeast Brazil, *Journal of Climate*, 26,
565 9137–9154, <https://doi.org/10.1175/jcli-d-12-00642.1>, 2013.
- Martin, S. T., Artaxo, P., Machado, L. A. T., Manzi, A. O., Souza, R. A. F., Schumacher, C., Wang, J., Andreae, M. O., Barbosa, H. M. J., Fan, J., Fisch, G., Goldstein, A. H., Guenther, A., Jimenez, J. L., Pöschl, U., Dias, M. A. S., Smith, J. N., and Wendisch, M.: Introduction: Observations and Modeling of the Green Ocean Amazon (GoAmazon2014/5), *Atmospheric Chemistry and Physics*, 16, 4785–4797, <https://doi.org/10.5194/acp-16-4785-2016>, 2016.
- 570 Martin, S. T., Artaxo, P., Machado, L., Manzi, A. O., Souza, R. A. F., Schumacher, C., Wang, J., Biscaro, T., Brito, J., Calheiros, A., Jardine, K., Medeiros, A., Portela, B., de Sá, S. S., Adachi, K., Aiken, A. C., Albrecht, R., Alexander, L., Andreae, M. O., Barbosa, H. M. J., Buseck, P., Chand, D., Comstock, J. M., Day, D. A., Dubey, M., Fan, J., Fast, J., Fisch, G., Fortner, E., Giangrande, S., Gilles, M., Goldstein, A. H., Guenther, A., Hubbe, J., Jensen, M., Jimenez, J. L., Keutsch, F. N., Kim, S., Kuang, C., Laskin, A., McKinney, K., Mei, F., Miller, M., Nascimento, R., Pauliquevis, T., Pekour, M., Peres, J., Petäjä, T., Pöhlker, C., Pöschl, U., Rizzo, L., Schmid,
575 B., Shilling, J. E., Dias, M. A. S., Smith, J. N., Tomlinson, J. M., Tóta, J., and Wendisch, M.: The Green Ocean Amazon Experiment (GoAmazon2014/5) Observes Pollution Affecting Gases, Aerosols, Clouds, and Rainfall over the Rain Forest, *Bulletin of the American Meteorological Society*, 98, 981–997, <https://doi.org/10.1175/bams-d-15-00221.1>, 2017.

- Morrison, H., Peters, J. M., Chandrakar, K. K., and Sherwood, S. C.: Influences of Environmental Relative Humidity and Horizontal Scale of Subcloud Ascent on Deep Convective Initiation, *Journal of the Atmospheric Sciences*, 79, 337–359, <https://doi.org/10.1175/jas-d-21-0056.1>, 2022.
- Schiro, K. A., Neelin, J. D., Adams, D. K., and Lintner, B. R.: Deep Convection and Column Water Vapor over Tropical Land versus Tropical Ocean: A Comparison between the Amazon and the Tropical Western Pacific, *Journal of the Atmospheric Sciences*, 73, 4043–4063, <https://doi.org/10.1175/jas-d-16-0119.1>, 2016.
- Schiro, K. A., Ahmed, F., Giangrande, S. E., and Neelin, J. D.: GoAmazon2014/5 campaign points to deep-inflow approach to deep convection across scales, *Proceedings of the National Academy of Sciences*, 115, 4577–4582, <https://doi.org/10.1073/pnas.1719842115>, 2018.
- Schlemmer, L. and Hohenegger, C.: Modifications of the atmospheric moisture field as a result of cold-pool dynamics, *Quarterly Journal of the Royal Meteorological Society*, 142, 30–42, <https://doi.org/10.1002/qj.2625>, 2015.
- Schumacher, C. and Funk, A.: GoAmazon2014/5 Three-dimensional Gridded S-band Reflectivity and Radial Velocity from the SIPAM Manaus S-band Radar, <https://doi.org/10.5439/1459573>, 2018a.
- Schumacher, C. and Funk, A.: GoAmazon2014/5 Rain Rates from the SIPAM Manaus S-band Radar, <https://doi.org/10.5439/1459578>, 2018b.
- Sherwood, S. C., Bony, S., and Dufresne, J.-L.: Spread in model climate sensitivity traced to atmospheric convective mixing, *Nature*, 505, 37–42, <https://doi.org/10.1038/nature12829>, 2014.
- Silva Dias, M. A. F.: Cloud and rain processes in a biosphere-atmosphere interaction context in the Amazon Region, *Journal of Geophysical Research*, 107, <https://doi.org/10.1029/2001jd000335>, 2002.
- Stevens, B. and Bony, S.: What Are Climate Models Missing?, *Science*, 340, 1053–1054, <https://doi.org/10.1126/science.1237554>, 2013.
- Stull, R.: *Practical Meteorology: An Algebra-based Survey of Atmospheric Science*, BC Open Textbook Collection, AVP International, University of British Columbia, <https://books.google.com.br/books?id=xP2sDAEACAAJ>, 2016.
- Tang, S., Xie, S., Zhang, Y., Zhang, M., Schumacher, C., Upton, H., Jensen, M. P., Johnson, K. L., Wang, M., Ahlgrim, M., Feng, Z., Minnis, P., and Thieman, M.: Large-scale vertical velocity, diabatic heating and drying profiles associated with seasonal and diurnal variations of convective systems observed in the GoAmazon2014/5 experiment, *Atmospheric Chemistry and Physics*, 16, 14 249–14 264, <https://doi.org/10.5194/acp-16-14249-2016>, 2016.
- Tian, Y., Zhang, Y., Klein, S. A., and Schumacher, C.: Interpreting the diurnal cycle of clouds and precipitation in the ARM GoAmazon observations: Shallow to deep convection transition, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033 766, <https://doi.org/10.1029/2020JD033766>, 2021.
- Tota, J., Fisch, G., Oliveira, P., Fuentes, J., and Siegler, J.: Análise da variabilidade diária da precipitação em área de pastagem para a época chuvosa de 1999 - projeto TRMM/LBA, *Acta Amazonica*, 30, 629–629, <https://doi.org/10.1590/1809-43922000304639>, 2000.
- Waite, M. L. and Khouider, B.: The Deepening of Tropical Convection by Congestus Preconditioning, *Journal of the Atmospheric Sciences*, 67, 2601–2615, <https://doi.org/10.1175/2010jas3357.1>, 2010.
- Wu, C.-M., Stevens, B., and Arakawa, A.: What Controls the Transition from Shallow to Deep Convection?, *Journal of the Atmospheric Sciences*, 66, 1793–1806, <https://doi.org/10.1175/2008jas2945.1>, 2009.
- Zhuang, Y., Fu, R., Marengo, J. A., and Wang, H.: Seasonal variation of shallow-to-deep convection transition and its link to the environmental conditions over the Central Amazon, *Journal of Geophysical Research: Atmospheres*, 122, 2649–2666, <https://doi.org/10.1002/2016jd025993>, 2017.