

# Aerosol uncertainties in tropical precipitation changes for the mid-Pliocene Warm Period

Anni Zhao<sup>1,2</sup>, Ran Feng<sup>3</sup>, Chris M. Brierley<sup>2</sup>, Jian Zhang<sup>1</sup>, and Yongyun Hu<sup>1</sup>

<sup>1</sup>Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University

<sup>2</sup>Department of Geography, University College London

<sup>3</sup>Center for Integrative Geosciences, University of Connecticut

**Correspondence:** Anni Zhao (annizhao@pku.edu.cn) and Yongyun Hu (yyhu@pku.edu.cn)

**Abstract.** The mid-Pliocene Warm Period (mPWP, ~~3.3–3.0~~ 3.3–3.0 Ma) was characterised by an atmospheric CO<sub>2</sub> concentration exceeding 400 ppmv with minor changes in continental and orbital configurations. Simulations of this past climate state have improved with newer models, but still show some substantial differences from proxy reconstructions. There is little information about atmospheric aerosol concentrations during the Pliocene, but previous work suggests that ~~is-~~ it could have been quite different from the modern. Here we apply idealised aerosol scenario experiments to examine the importance of aerosol forcing on mPWP tropical precipitation and the possibility of aerosol uncertainty explaining the mismatch between reconstructions and simulations. The absence of industrial pollutants leads to further warming especially in the Northern Hemisphere. The Intertropical Convergence Zone (ITCZ) becomes narrower and stronger and shifts northward after removal of anthropogenic aerosols. Though not affecting the location of monsoon domain boundary, removal of anthropogenic aerosol alters the amount of rainfall within the domain, increasing summer rain rate over eastern and southern Asia and western Africa. This work demonstrates that uncertainty in aerosol forcing could be the dominant driver in tropical precipitation changes during the mid-Pliocene: causing larger impacts than the changes in topography and greenhouse gases.

*Copyright statement.* TEXT

## 1 Introduction

The mid-Pliocene or mid-Piacenzian Warm Period (mPWP, around 3.3–3.0 Ma, Dowsett et al., 1999; Haywood et al., 2016a) was the most recent time in Earth’s history when atmospheric CO<sub>2</sub> concentration exceeded 400 ppmv (Bartoli et al., 2011; de la Vega et al., 2020) with nearly the same topography and orbital configuration as modern. This period is recognised as the last period with quasi-equilibrium warm climate before Pleistocene when climate had nearly fully responded to the high atmospheric CO<sub>2</sub> concentration (e.g. Haywood et al., 2010). Understanding the environmental process in the mPWP provides a chance to understand how the climate system would respond to a perturbation of radiative forcing and to evaluate the impacts of climate change caused by great CO<sub>2</sub> emissions (e.g. Haywood et al., 2016b; Pagani et al., 2010). The first study reconstructing regional Pliocene climate was conducted by Zubakov and Borzenkova (1988) based on more than 20 sequences from terrestrial

and marine cores. They pointed out that the Pliocene Optimum could be a future analogue, with high CO<sub>2</sub> concentration and climate showing 100-300 kyr cycles with a 4-5 °C amplitude. Proxy data suggested a warming in global mean surface temperature (GMST) of ~~2.5--4.0~~ 2.5-4.0 °C during the mPWP than 1850-1900 CE, based on the estimation of global sea surface temperature (GSST) anomaly and its relationship in changing rate with the GMST (Gulev et al., 2021). The Pliocene climate was ~~characterized~~ characterised by reduced temperature gradients (Haywood et al., 2013, 2020; Dowsett et al., 2013; Foley and Dowsett, 2019) from equator to pole ~~e.g.~~ (e.g. Dowsett and Poore, 1991) and from western to eastern tropical Pacific Ocean which has been recognised as a tropical Pacific warm pool pattern (Wara, 2005; Dowsett, 2007; Zhang et al., 2014; O'Brien et al., 2014).

Climate models have been used to simulate mPWP climate and to investigate the drivers and mechanisms behind the mPWP climate changes shown in proxy data (e.g. Fedorov et al., 2013). The Pliocene Model Intercomparison Project (PlioMIP; Haywood et al., 2010, 2016b) is a coordinated international climate modelling project initiative aimed of understanding the climate and environments of mPWP, exploring model uncertainties and evaluating the potential relevance to future climate change. It estimates a warming of 3.2°C (2.1°C to 4.8°C) compared to the pre-industrial control runs in the PlioMIP2 (Haywood et al., 2020) and 2.7°C (1.8°C to 3.6°C) in the PlioMIP1 (Haywood et al., 2013). PlioMIP simulations suggest enhanced monsoons in West Africa, India and East Asia (Zhang et al., 2013; Berntell et al., 2021; Han et al., 2021; Feng et al., 2022), and precipitation in the South Pacific Convergence Zone (SPCZ) and the ITCZ during the mPWP (Haywood et al., 2020; Han et al., 2021). Yet ongoing climate change since the 20<sup>th</sup> century has not shown those mPWP precipitation changes during which monsoon precipitation is affected by the counteraction between Greenhouse Gases (GHGs) induced increase and aerosol emission induced decrease (Douville et al., 2021). Many climate features, especially the hydroclimate features, of the mPWP differ substantially from the projected features of near future climate (Feng et al., 2022; Bhattacharya et al., 2022; Burton et al., 2023). Although the mPWP climate was thought to reflect earth system responses that operates at a longer time scale (Lunt et al., 2010; Feng et al., 2022), Pliocene climate was also likely driven by different atmospheric aerosol forcing conditions (Unger and Yue, 2014; Sagoo and Storelvmo, 2017). Whether those different forcing conditions play an role in driving the Pliocene hydroclimate remains unknown.

Aerosols are an important driver in climate system. Knowing their atmospheric concentrations and mechanisms behind climate response are key tasks for simulating past and future. Aerosols directly affect absorbed incoming solar radiation and indirectly affect the radiative flux through their role in determining cloud properties. Aerosol-cloud interactions include the changes in cloud albedo through altering droplet size and concentration (Twomey, 1977) and changes in cloud lifetime through altering the precipitation efficiency (Albrecht, 1989). Future changes in the global aerosol burden are dependent on anthropogenic emissions, yet natural sources of aerosol emissions are also expected to change (Tegen and Schepanski, 2018). ~~mPWP simulations now use modern-day or pre-industrial aerosol concentration that may differ from the conditions during the mPWP. It implies that aerosol effects may be one of the possible explanation for the mismatch between reconstructions and simulations.~~ The discrepancy between mPWP simulations and reconstructions (Haywood et al., 2013, 2020) implies that models might miss some important mechanisms (Fedorov et al., 2013) or prescribed forcing could be a source of uncertainty (Feng et al., 2019). The mPWP simulations now use modern-day or pre-industrial aerosol concentration same as the control

runs that may differ from the conditions during mPWP. Lack of including the effect of aerosol forcings and the usage of unrealistic prescribed aerosol concentration in mPWP simulations implies that aerosol effects may be one of the possible explanation for the mismatch between reconstructions and simulations. However, little research has tried to understand aerosol effects in the Pliocene (e.g. Unger and Yue, 2014). More attention in palaeoclimate studies relevant to aerosols has been paid to the Last Glacial Maximum (LGM) period when the global dust cycle was enhanced (Lambert and Albani, 2021). An idealised study suggests the importance of aerosol and chemistry-climate feedbacks in modelling Pliocene climate, as the aerosol cooling compensates 15-100 % of the warming induced by high CO<sub>2</sub> while chemistry-climate feedback warms the climate with the magnitude of 30-250% of the CO<sub>2</sub> induced warming (Unger and Yue, 2014). Sagoo and Storelmo (2017) found that dust indirect effects could explain some of the mismatch between model and data for LGM and mPWP. They used a new empirical parameterisation for ice nucleation on dust particles to investigate radiative forcing induced by different dust loading from low to high. Results showed that increased dust reduces the size of ice crystals in clouds while increasing their amount. This increases the shortwave cloud radiative forcing, thus, cools surface temperature and amplifies polar warmth (Sagoo and Storelmo, 2017). Their mPWP simulation modified by extreme low dust suggests surface temperature warming and polar amplification due to reduced radiative forcing by increased size while decreased amount of ice crystals in clouds (Sagoo and Storelmo, 2017).

In this study, we analyse two Pliocene simulations ~~with different aerosol scenarios~~ that were performed initially to analyse the effects of aerosol-cloud interactions on mPWP seasonally sea ice-free Arctic (Feng et al., 2019) to further investigate the potential effect of aerosol on mPWP climate. Different aerosol scenarios were applied, one with pre-industrial aerosol concentrations and one with present-day aerosol concentrations published in Lamarque et al. (2010). The simulations were performed initially to analyse the effects of aerosol-cloud interactions on mPWP seasonally sea ice-free Arctic (Feng et al., 2019). We compare the changes caused by removing anthropogenic concentrations in Pliocene, in order to raise the importance of considering aerosol effects in modelling Pliocene climate. This investigation also highlights the potential analogue nature of Pliocene climate, not to the present-day polluted climate, but to future climate scenarios which feature removal of anthropogenic pollutants (Lee et al., 2021). We focus on changes in tropical precipitation, because precipitation varies much more than other near-uniform variables like temperature and irradiation in the tropics.

## 2 Methods

### 2.1 Model description

Simulations were performed with the Community Earth System Model version 1.2 (CESM1.2; Hurrell et al., 2013), consisting of Community Atmospheric Model version 5.3 (CAM5.3; Martinez, 2012), Parallel Ocean Program version 2 (POP2; Danabasoglu et al., 2012), Community Land Model version 4 (CLM4; Oleson, 2010) and Community Ice Code version 4 (CICE4; Holland et al., 2012). CESM1 was developed from the Community Climate System Model version 4 (CCSM4, Gent et al., 2011) by adding capabilities related to BGC processes and aerosol effects (Meehl et al., 2013). Differences and similarities between CESM1 and CCSM4 are summarised in ~~Meehl et al. (2013)~~ Meehl et al. (2013). The key differences–updates between

CCSM4 and CESM1 come from a new aerosol scheme, inclusion of aerosol-cloud-interactions, and a more realistic boundary layer and radiation in CAM5 (Meehl et al., 2013; Martinez, 2012). The 3-mode version of modal aerosol module (MAM3; Liu et al., 2012) is used as the aerosol micro-physical scheme for long-term climate simulations, which uses Aiken ~~-,accumulation and coarse-~~ (0.02–0.08  $\mu\text{m}$ ), accumulation (0.08–1  $\mu\text{m}$ ) and coarse (1.0–10  $\mu\text{m}$ ) modes to solve ~~number and size-~~ size and   
95 number concentration of internal condensation and coagulation of different species among modes. Assumptions and limitations of MAM3 are described in Liu et al. (2012). Though CAM5 simulates too strong response in cloud radiative to aerosol changes (Martinez, 2012), its ability to simulate cloud and cloud radiative forcing has been improved since the previous generations (Kay et al., 2012).

## 2.2 Experimental design

100 ~~The simulations-~~ The existed simulations (Feng et al., 2019) were branched from an existing CCSM4-PlioMIP1 simulation that only considers the aerosol direct effect (Rosenbloom et al., 2013). The resolution was set up to  $0.9^\circ \times 1.25^\circ$  for the atmospheric and land components and  $\sim 1^\circ$  for the oceanic and sea ice components. Two pollutant scenarios were applied. One simulation applied pre-industrial emissions (hereafter referred as to Plio\_Pristine, which can be treated as equivalent to a PlioMIP1 simulation). The other prescribed pre-industrial emissions plus industrial pollutants of anthropogenic  $\text{SO}_2$ ,   
105 sulfate and organic compounds estimated for the 2000s from an gridded ( $0.5^\circ \times 0.5^\circ$ ) emission dataset (hereafter referred as to Plio\_Polluted) published in Lamarque et al. (2010). The emission dataset provided consistent gridded anthropogenic (defined as originating from industrial, domestic and agriculture activity sectors) and biomass burning of reactive gases and aerosols covering the historical period from 1850 to 2000 CMIP5 models to use in running chemistry model simulations that would contributed to the assessment in IPCC AR5 (Lamarque et al., 2010). Compared to Plio\_Polluted, Plio\_Pristine simulates   
110 a positive adjusted effective radiative forcing of  $1.29 \text{ W m}^{-2}$ , in which radiative forcing due to aerosol cloud interactions contributes to  $1.11 \text{ W m}^{-2}$  and direct radiative forcing contributes to  $0.18 \text{ W m}^{-2}$  (Feng et al., 2019). The emission dataset (Lamarque et al., 2010) includes reactive gases and aerosols covering the historical period from 1850 to 2000, with the aim of providing consistent gridded emissions for CMIP5 models to run chemistry model simulations that would contributed to the assessment in IPCC AR5. Boundary conditions were set up based on the PlioMIP1 (Haywood et al., 2011). Each simulation   
115 was branched from model year 500 of the CCSM4-PlioMIP1 simulation, and run for another 300 model years. Outputs are taken as the averaged means of the last 50 model years.

## 2.3 Analysis techniques

### 2.3.1 Monsoon analysis

It is not appropriate to use fixed present-day monsoon domain boundary to analyse paleo monsoon properties as the monsoon   
120 domain varies through time. Here we adopt the definition of monsoon domain in Wang and Ding (2008) and Wang et al. (2014). Local summer is defined as May–September (MJJAS) in the Northern Hemisphere or November–March (NDJFM) in the Southern Hemisphere, and local winter is defined as NDJFM in the Northern Hemisphere and MJJAS in the Southern

Hemisphere. Averaged summer rain rate takes the averaged daily rain rate during local summer. Monsoon intensity is the difference in averaged daily precipitation rate between local summer and local winter. Global monsoon domain is defined as at least of 55% of the annual total rainfall comes from summer, and the monsoon intensity is no less than 2 mm per day.

### 2.3.2 Data model comparison

Simulated mPWP annual mean temperature anomalies (~~between~~ Plio\_Pristine ~~and the pre-industrial (hereafter referred as to~~ PI) are compared to the reconstructed mPWP sea surface temperature (SST) anomalies, which were presented as the reconstructed mPWP SSTs dataset in PRISM4 (Foley and Dowsett, 2019) with an interval of 30,000 years minus the observed 1870-1988 SSTs from the NOAA Extended Reconstructed Sea Surface Temperature (ERSST) version 5 dataset (Huang et al., 2017). Salzmann et al. (2008) compared 28 present-day mean annual precipitation anomalies (mid-Pliocene - present-day) from literature for selected regions. Only 10 data points were chosen for data-model comparison in this study as their precise coordinates and anomalies could be found in literature. The 10 points have been regenerated into 6 after the combination of points if their sites were located within a single model grid. Three sites in western coast of USA and 3 in Yunnan, China have been regenerated into new sites respectively by taking their averages. Therefore each three have been reorganised into a single point by taking their average. (~~Feng et al., 2022~~) [Feng et al. \(2022\)](#) published a compilation of mPWP proxy records reflecting the mPWP terrestrial hydroclimate change shown in the difference between precipitation and evaporation, which consists of 62 data points after the exclusion by quality control. Here we follow the treatment of (~~Feng et al., 2022~~) [Feng et al. \(2022\)](#) to combine the co-located sites if they were less than 150 km apart and featured same-signed anomalies. The combination regenerates the site sets 35, 72; 22, 77; 79,61; 58,61,79,101 and 65, 66 [of Feng et al. \(2022\)](#) into new sites located at the center of sets.

## 3 Results

### 3.1 Simulated mPWP climate change relative to the pre-industrial

Fig. 1a shows that the tropical annual mean surface temperature of Plio\_Pristine is overall warmer than the ~~pre-industrial (hereafter referred as to PI)~~ [PI](#). Plio\_Pristine produces warmer high-latitude Pacific and Southern Ocean than the tropics as compared to PI. Eastern Pacific Ocean along the tropics and the upwelling region along the coast line of North America is about 0.5°C warmer than western Pacific. Plio\_Pristine captures both reduced meridional and zonal temperature gradients during the mPWP that have been shown in the PlioMIPs (Haywood et al., 2013, 2020), but the amplifying magnitude in Plio\_Pristine is weaker than the PlioMIPs (Fig. 2a). The underestimation in Northern Hemisphere warming may be explained by that Plio\_Pristine was branched from an earlier CCSM4-PlioMIP1 simulation that had underestimated the warming in the Northern Hemisphere (Rosenbloom et al., 2013). Though Plio\_Pristine underestimates positive temperature anomalies, its simulated change still within the range suggested by the PlioMIPs. 13 out of the 37 sites show a mismatch between simulated surface temperature anomaly and reconstructed SST anomaly ([Foley and Dowsett, 2019](#)) smaller than 1.0°C. Large

mismatch occurs in high-latitude North Atlantic Ocean and the upwelling region near the western coastline of South Africa. Reconstructed SST anomalies show an averaged warming of 4.4°C north to 40°N of North Atlantic Ocean and 7.0°C over the upwelling region, which are 3.5°C and 5.6°C higher than the simulated warming, respectively. The reconstructed ~~zonal-~~meridional temperature gradient along the tropical Pacific Ocean (between sites near Colombia and sites near Southeast Asia) is around 0.5°C greater than simulations (Plio\_Pristine - PI). Data model mismatch suggests that Plio\_Pristine underestimates the magnitude of reduction in both SST gradients during the mPWP as compared to reconstructed SST anomalies (Foley and Dowsett, 2019).

Previous studies of mPWP monsoon change based on data were mainly focused on Asian monsoons due to the rare availability of geological evidence, which suggested a wetter East Asian monsoon during the mPWP (e.g. Xiong et al., 2010; Nie et al., 2014). The PlioMIPs show enhanced monsoon over West and North Africa, India and East Asia that are consistent with proxy data (Li et al., 2018; Feng et al., 2022). Fig. 1b shows that comparing to the PI, Plio\_Pristine produces greater annual mean precipitation (as the sum of stable large-scale and convective precipitation rate) over tropical oceans and land monsoon regions (western and northern Africa, East Asia, Australia and parts of Southeast Asia). Precipitation decreases over subtropical oceans, eastern parts of South America and monsoon areas over North America. Plio\_Pristine produces the right sign of precipitation as proxy data suggest, but in general underestimates the enhancing magnitudes especially over Australian sites where proxy data suggest large positive precipitation anomaly (Fig. 1b and 2b). Though Plio\_Pristine produces less precipitation over tropical Northern Hemisphere as compared to the multi-model means from PlioMIP1 (Haywood et al., 2013) and PlioMIP2 (Haywood et al., 2020), it overall agrees with the findings from the PlioMIPs.

### 3.2 Effects of removing anthropogenic aerosols on temperature

#### ~~Reducing atmospheric aerosols-~~

According to IPCC AR6, human-induced aerosols contributed to a cooling of 0.0°C to 0.8°C between 2010-2019 relative to 1850-1900 (IPCC, 2021). Therefore, removing anthropogenic aerosols is expected to warm the climate. Removal of present-day anthropogenic aerosols from the atmosphere (Plio\_Pristine - Plio\_Polluted) causes a global warming of 0.84 °C (Feng et al., 2019), which is close to the estimated modern cooling contributed by aerosols at ~~0--0.8~~ 0-0.8 °C in the IPCC AR6 (IPCC, 2021). Simulated warming is in line with the theory of aerosol effects, as aerosols contribute a negative forcing on climate which in turn removing aerosols from the atmosphere should lead to a positive forcing.

Spatial pattern of temperature change is consistent with the change in energy flux. Overall, asymmetrical change in radiation flux matches with the larger aerosol concentration change in the Northern Hemisphere (Fig. 3a and 4a). Major change in energy flux after the removal of anthropogenic emissions occurs in the Northern hemisphere especially over lands that leads to further warming. The warming pattern due to asymmetrical emissions concentrating in the Northern Hemisphere agrees with the findings from an earlier study (Samset et al., 2018), in which several models have been applied to test removing all anthropogenic emission under present-day conditions. Annual mean temperature rises about ~~0.4--1.0~~ 0.4-1.0 °C in tropical Pacific Ocean and causes more warming by increasing ~~1.0--1.4~~ 1.0-1.4 °C in subtropical Pacific and the upwelling region in eastern Pacific after removing pollutants (Fig. 5a), which therefore reduces both meridional and zonal temperature gradients in

Pacific Ocean. In future climate projections, removing emitted pollutants is shown to enhance global mean warming, especially in the Northern Hemisphere where the emissions mostly come from. However, ~~removing anthropogenic aerosols causes more warming than~~ the mPWP boundary conditions ~~over~~ causes more warming than removing anthropogenic aerosols nearly global except Northeastern Pacific and high-latitude North Atlantic Ocean (Fig. 5a,c,e). ~~This~~ More warming lead by the mPWP boundaries than removal of aerosols suggests that changes in boundary conditions are relative more important than aerosol effects in mPWP warming.

### 3.3 Effects of removing anthropogenic aerosols on tropical precipitation

Precipitation responds to removal of aerosols in a more complex manner, but in general it responds intensively in deep convective regions (Fig. 5b,d,f). Precipitation enhances in the tropics and reduces in the adjacent subtropics in response to removing emissions. The precipitation change pattern matches more with mid- and high- level clouds (Fig. 6 and A2 ~~-.Precipitation enhances-~~). Removing anthropogenic aerosols leads to stronger vertical velocity at the Equator (Fig. 7a). The meridional circulation in the tropics ~~while reduces in the adjacent subtropics in response to removing emissions, which shows that the local convection becomes narrower and stronger with~~ is strongly enhanced in the Southern Hemisphere and slightly weakened in the Northern Hemisphere (Fig. 7b) after removing anthropogenic aerosols from the atmosphere, which is related to a northward shift of the ITCZ. The narrower ITCZ is consistent with a warmer climate simulated by Plio\_Pristine, and agrees with an earlier study which suggests that future ITCZ would become narrower and weaker under global warming in the CMIP5 simulations (Byrne and Schneider, 2016). Ridley et al. (2015) and Voigt et al. (2017) found that introducing anthropogenic aerosols causes southward shift of the ITCZ. In ~~our case, the~~ opposite, removing anthropogenic emission would expect a northward shift of ~~ITCZ due to removal of anthropogenic aerosols is~~ the ITCZ. Our results are consistent with this finding. The pattern of the magnitudes of tropical precipitation anomaly matches the tropical temperature warming over oceans, which agrees with Berg et al. (2013) as higher temperatures cause stronger convective precipitation. Tropical precipitation increases greater over the tropical Pacific and Indian Ocean where the warming is stronger than the tropical mean warming (Fig. 5), consistent with the positive correlation between tropical precipitation change and spatial deviations of SST warming from the tropical mean (Xie et al., 2010). In the northern tropics, anthropogenic aerosols caused a rainfall reduction through the twentieth century (Ridley et al., 2015). Correspondingly, precipitation should increase after removal of emissions. The positive precipitation change over Asia between Plio\_Pristine and Plio\_Polluted agrees with the argument.

Strong precipitation response to removal of anthropogenic aerosols implies that the choice of prescribed aerosol scenario could affect simulated mPWP monsoon response. Figure 8 shows the change in global monsoon domain and summer rain rate between Plio\_Pristine and PI (panel a) and between Plio\_Pristine and Plio\_Polluted (panel b). Little difference among the boundary of global monsoon domain for the PI, Plio\_Pristine and Plio\_Polluted implies that removal of anthropogenic aerosols has little effect on the location of global monsoon domain. Though not affecting locations, aerosol forcing alters the spatial distribution of land monsoon precipitation. ~~Land regional monsoons are enhanced, agreeing with the observations that reduce aerosol-induced cooling and increase the precipitation over western Africa and eastern and southern Asia during local monsoon seasons as~~ As anthropogenic emission partly caused the global land monsoon precipitation reduction during



1950s to 1980s and weakened western African and eastern and southern Asian monsoon over the 20th century by cooling effect (Douville et al., 2021) ~~–~~, reducing aerosol-induced cooling would increase the precipitation over western Africa and eastern and southern Asia during local monsoon seasons. In our case, land regional monsoons are enhanced after removing anthropogenic aerosol, which is consistent with the expectation. Comparison between Plio\_Pristine and Plio\_Polluted shows the expected change in temperature and monsoon (Fig. 5,8), which is consistent with projected future monsoon enhancement, as global and Asian summer monsoon precipitation is likely to increase by 2050s due to the expected reductions in anthropogenic aerosol emissions (Wilcox et al., 2020). Though this study used idealised aerosol scenario that could not occur during mPWP, it reveals the importance of prescribed aerosol scenario on simulating mPWP monsoon response.

## 230 4 Discussion

### 4.1 Key forcing ~~driving tropical precipitation change~~ contributing to aerosol effect

Aerosols affect the climate directly by altering radiation and indirectly by interacting with clouds via changing cloud albedo and cloud lifetime. ~~In tropical regions, The largest forcing is due to shortwave~~ cloud radiative effects (i.e. indirect effect), ~~which~~ primarily drive the energy flux change after removing pollutants (Fig. 3) ~~– Aerosols play less role in absorbing a).~~  
235 Aerosol change causes direct increase in the net incoming shortwave uptake in the Northern Hemisphere through less reflection and plays less role in affecting shortwave radiation in the Southern Hemisphere ~~– Clear-sky net shortwave flux gradually increases towards north~~ and the tropics (Fig. 3a) due to the asymmetrical change in distribution (Fig. 4a), but its net effect is much smaller than the ~~full-sky condition, while~~ shortwave indirect effect. The direct radiative forcing due to removing anthropogenic aerosols is much smaller (Feng et al. (2019) have found that the effective radiative forcing due to removing anthropogenic aerosols primarily arises from change in aerosol-cloud interactions. In the tropics, clouds contribute to a net forcing greater than  $2 \text{ Wm}^{-2}$  with ~~more shortwave absorption and less longwave emission. In the Northern Hemisphere, aerosol reflection reduces consistently with the decrease in aerosol optical depth~~ less shortwave reflection at top of clouds and less downwelling longwave emission (Fig. 4a), but is still less important than the change in shortwave cloud forcing. Feng et al. (2019) 3b). Feng et al. (2019) focused on change in cloud droplet properties, as shortwave cloud forcing  
240 is dependent on cloud optical depth, whose change is positively correlated with the change in cloud droplet concentration and cloud liquid water content. ~~Figure 6 shows decrease in~~ Removal of pollutants decreases droplet concentration (especially over lands and high latitudes, Fig. A3a) and decrease ~~in~~ cloud liquid path (except tropical Pacific between  $5^{\circ}\text{S}$  and  $5^{\circ}\text{N}$  and western Africa, Fig. A3b) ~~after the removal of pollutants~~. The uniform decrease in droplet concentration and overall decrease in cloud liquid path suggests lower cloud albedo and shorter cloud lifetime due to removing anthropogenic aerosols.

### 250 4.2 Relative importance comparison between mPWP boundary conditions and aerosol forcing

~~The overall uniform warming over~~ Over tropical and subtropical regions, the overall uniform warming induce by mPWP boundaries (Plio\_Pristine - PI and ~~Plio\_Pristine – Plio\_Polluted~~) result in general ~~increase~~ decrease in sea level pressure



and have relatively small effects on surface wind  ~~,while removing anthropogenic aerosols- (Fig. 10a,c,e). Though removal of anthropogenic aerosols also show warming in tropical and subtropical regions, it~~ rises the sea level pressure and have much stronger effects on surface wind that shows seasonal variances (Fig. 10 ~~b,d,f~~ ). Studies have highlighted that anthropogenic aerosols cause the southward shift in the ITCZ, weaken the Hadley circulation and reduce the precipitation in deep convective areas in response to the Northern Hemisphere cooling induced by aerosols (Hwang et al., 2013; Wang et al., 2019). Removal of human-induced aerosol emissions causes more precipitation change than the mPWP boundary condition over subtropical Pacific Ocean and South Atlantic Ocean, Indian Ocean and Southeast Asia. Figure 11 shows the relative importance of removing anthropogenic aerosols and mPWP boundary conditions on mPWP zonal mean precipitation change. ~~The ratio (panel a) indicates that the relative importance of effects of~~ Precipitation change in response to aerosol forcing and mPWP boundary conditions (including high CO<sub>2</sub>) on precipitation ~~is shows~~ complicated  ~~,but a pattern as there is no uniform changing trend.~~ Overall aerosol effect is more important over the tropics ~~(panel b)~~, which could imply the importance of aerosol scenario in simulating Pliocene climate. It highlights the importance of aerosol in mPWP data model comparison, as proxy recorded tropical temperature and hydroclimate conditions can be sensitive to aerosol emission conditions during the Pliocene.

### 4.3 Implications for simulating Pliocene climate

Strong tropical precipitation change in response to removal of anthropogenic aerosols indicates that simulated mPWP precipitation is sensitive to prescribed aerosol scenarios. Previous simulations of the mPWP atmospheric chemistry confirmed that there is a significant increase in the concentrations of aerosol precursors from terrestrial ecosystem emissions (Unger and Yue, 2014) and that simulated mPWP climate is sensitive to indirect effects of dust (Sagoo and Storelvmo, 2017). However, Unger and Yue (2014) only looked at direct radiative effects on mPWP climate induced by feedbacks from physical climate change on reactive composition, yet they did not consider aerosol-cloud interactions. To what extent the terrestrial biogenic aerosols may change mPWP hydrological cycle remains unknown. In Unger and Yue (2014), continental greening and warmer climate during the mPWP leads to greater emissions of aerosols and precursors from more active wildfire and biome productions, yet, this increase is likely much smaller than the anthropogenic pollutant emissions. For example, due to the enhanced wildfire activity, SO<sub>2</sub> increased by 11 Tg yr<sup>-1</sup> in Unger and Yue (2014) mPWP simulations, yet SO<sub>2</sub> from anthropogenic sources is estimated to be ~20 Tg yr<sup>-1</sup> from 1980 to 2000s (Smith et al., 2011). As such, despite that enhanced biogenic emissions may compensate some of the responses seen in the Plio\_Pristine, we argue that the absence of anthropogenic pollutant from the mPWP troposphere remains one of the important factors driving the differences between the mPWP and near future climate, especially the Arctic climate (Feng et al., 2019) and hydroclimate. In addition, the mPWP also likely featured different amount of dust aerosols. ~~Sagoo and Storelvmo (2017) looked in to the effect of dust and demonstrated that increased dust emissions resulted in an enhanced dynamical response that lead to great changes in hydrological circulations. Their~~ Sagoo and Storelvmo (2017)'s Pliocene simulation with idealised extreme low dust scenario shows greater change in precipitation than only consider CO<sub>2</sub> forcing, yet only used idealized idealised scenario ~~(Sagoo and Storelvmo, 2017)~~. Usage of more realistic prescribed aerosol scenario should be considered in future work.

Finally, despite that we ~~analyzed~~ analysed change in mPWP hydrological cycle, we didn't fully account for changes in the moisture budget that explains our simulated hydroclimate changes in the Plio\_Prisine. The anomalous moisture budget can be decomposed in to thermodynamic ~~,dynamic and residual~~ (due to change in specific humidity), dynamic (due to change in mean circulation) and residual (due to transient eddy) components etc (Trenberth and Guillemot, 1995; Held and Soden, 2006).  
290 D'Agostino et al. (2019, 2020) stated that monsoon changes under RCP8.5 are likely driven by thermodynamics and net energy input in response to global warming ~~.As the high~~ , which implies that high CO<sub>2</sub> concentration would affect precipitation by increasing the thermodynamic component of the moisture budget.. As the mPWP simulation also prescribed high atmospheric CO<sub>2</sub> concentration is also a major forcing in mPWP simulations, it implies that thermodynamic change plays an important role in mPWP precipitation change as well. Removal of anthropogenic emissions in this work would logically increase the effects of  
295 thermodynamic and net energy input due to its induced further warming and increased net radiative absorption. Further study could focus on decomposing moisture budget to determine which component plays an important role in generating mPWP precipitation change.

## 5 Conclusions

A set of existing simulations with two idealised aerosol scenarios (Feng et al., 2019) are analysed in this study to investigate the  
300 importance of aerosol forcing on mPWP tropical climate. Our results highlight the importance of prescribed aerosol scenarios in simulating mPWP precipitation change. In contrast to GHGs warm the climate, aerosols cool the climate. As expected, removal of pollutants causes global warming and further warms the Northern Hemisphere due to asymmetrical emissions. Aerosols affect precipitation in a more complicated manner. Precipitation enhances in the tropics especially over deep convective regions while reduces in adjacent subtropics showing a narrower and stronger with a northward shift in the ITCZ. Though not affecting  
305 the location of monsoon domain boundary, removal of anthropogenic aerosol changes the precipitation within the domain and increases summer rain rate over eastern and southern Asia and western Africa. Removal of anthropogenic emissions leads to lower cloud albedo and shorter cloud lifetime in simulated mPWP climate. Aerosols have more impacts on annual and seasonal precipitation over the tropics and subtropics than the mPWP boundary conditions.

In reality, the concentration and composition of aerosol during the mPWP were different to those in the pre-industrial and  
310 present-day, due to ~~mPWP had~~ warmer and wetter climate in the mPWP with greater vegetation cover and natural emissions without human effect, respectively. Climate response to aerosol forcing during the mPWP was expected to be different to those periods. Though we investigate idealised aerosol forcing on mPWP climate that cannot quantify the magnitude of uncertainty in Pliocene aerosol, we still argue that aerosols can be a dominant source of uncertainty as it could contribute to change in certain variables, such as tropical rainfall discussed in this study, as large as from other mPWP boundary conditions. It is  
315 important to consider the differences in aerosol conditions across the periods.

*Code availability.* TEXT

*Data availability.*

*Code and data availability.* The simulation data are hosted at Cheyenne supercomputer with the path: /CCSM/csm/CESM-CAM4-PLIOCENE. Instructions and permissions to relevant experiments will be provided upon request to ran.feng@uconn.edu. The processed data for figures are available at <https://github.com/annizhao1994/mPWPAerosol> and <https://zenodo.org/doi/10.5281/zenodo.10130118>. Source code of CESM1.2 can be downloaded from <https://www2.cesm.ucar.edu/models/cesm1.2/>

*Sample availability.* TEXT

*Video supplement.* TEXT

*Author contributions.* A.Z and C.M.B. performed the bulk of the writing and analysis. R.F contributed data for the manuscript and paid large effort on revision. Y.H and J.Z. revised the manuscript.

*Competing interests.* At least one of the co-author are members of the editorial board of Climate of the Past. The authors have no other competing interests to declare.

*Disclaimer.* TEXT

*Acknowledgements.* We acknowledge the high-performance computing support from Cheyenne (<https://doi.org/10.5065/D6RX99HX>) and Yellowstone (<ark:/85065/d7wd3xhc>) provided by NCAR's Computational and Information Systems Laboratory, sponsored by the National Science Foundation. This material is based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement 1852977. R.F. received funded by NSF grant 2103055 and 2238875. Y.H were founded in part by the National Natural Science Foundation of China, under grant 41888101. J.Z. was founded by National Natural Science Foundation of China grant 42372120.

335 **References**

- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Bartoli, G., Hönisch, B., and Zeebe, R. E.: Atmospheric CO<sub>2</sub> decline during the Pliocene intensification of Northern Hemisphere glaciations, *Paleoceanography*, 26, PA4213, <https://doi.org/10.1029/2010PA002055>, 2011.
- 340 Berg, P., Moseley, C., and Haerter, J. O.: Strong increase in convective precipitation in response to higher temperatures, *Nature Geoscience*, 6, 181–185, <https://doi.org/10.1038/ngeo1731>, 2013.
- Berntell, E., Zhang, Q., Li, Q., Haywood, A. M., Tindall, J. C., Hunter, S. J., Zhang, Z., Li, X., Guo, C., Nisancioglu, K. H., Stepanek, C., Lohmann, G., Sohl, L. E., Chandler, M. A., Tan, N., Contoux, C., Ramstein, G., Baatsen, M. L. J., von der Heydt, A. S., Chandan, D., Peltier, W. R., Abe-Ouchi, A., Chan, W.-L., Kamae, Y., Williams, C. J. R., Lunt, D. J., Feng, R., Otto-Bliesner, B. L., and
- 345 Brady, E. C.: Mid-Pliocene West African Monsoon rainfall as simulated in the PlioMIP2 ensemble, *Climate of the Past*, 17, 1777–1794, <https://doi.org/10.5194/cp-17-1777-2021>, 2021.
- Bhattacharya, T., Feng, R., Tierney, J. E., Rubbelke, C., Burls, N., Knapp, S., and Fu, M.: Expansion and Intensification of the North American Monsoon During the Pliocene, *AGU Advances*, 3, e2022AV000757, <https://doi.org/https://doi.org/10.1029/2022AV000757>, e2022AV000757 2022AV000757, 2022.
- 350 Burton, L. E., Haywood, A. M., Tindall, J. C., Dolan, A. M., Hill, D. J., Abe-Ouchi, A., Chan, W.-L., Chandan, D., Feng, R., Hunter, S. J., Li, X., Peltier, W. R., Tan, N., Stepanek, C., and Zhang, Z.: On the climatic influence of CO<sub>2</sub> forcing in the Pliocene, *Climate of the Past*, 19, 747–764, <https://doi.org/10.5194/cp-19-747-2023>, 2023.
- Byrne, M. P. and Schneider, T.: Narrowing of the ITCZ in a warming climate: Physical mechanisms, *Geophysical Research Letters*, 43, 11,350–11,357, <https://doi.org/https://doi.org/10.1002/2016GL070396>, 2016.
- 355 D’Agostino, R., Bader, J., Bordoni, S., Ferreira, D., and Jungclaus, J.: Northern Hemisphere Monsoon Response to Mid-Holocene Orbital Forcing and Greenhouse Gas-Induced Global Warming, *Geophysical Research Letters*, 46, 1591–1601, <https://doi.org/10.1029/2018GL081589>, 2019.
- D’Agostino, R., Brown, J. R., Moise, A., Nguyen, H., Silva Dias, P. L., and Jungclaus, J.: Contrasting Southern Hemisphere monsoon response: midHolocene orbital forcing versus future greenhouse-gas induced global warming, *Journal of Climate*, pp. 1–56, <https://doi.org/10.1175/JCLI-D-19-0672.1>, 2020.
- 360 Danabasoglu, G., Bates, S. C., Briegleb, B. P., Jayne, S. R., Jochum, M., Large, W. G., Peacock, S., and Yeager, S. G.: The CCSM4 Ocean Component, *J. Clim.*, 25, 1361–1389, 2012.
- de la Vega, E., Chalk, T. B., Wilson, P. A., Bysani, R. P., and Foster, G. L.: Atmospheric CO<sub>2</sub> during the Mid-Piacenzian Warm Period and the M2 glaciation, *Scientific Reports*, 10, 11 002, <https://doi.org/10.1038/s41598-020-67154-8>, 2020.
- 365 Douville, H., Raghavan, K., Renwick, J., Allan, R. P., Arias, P. A., Barlow, M., Cerezo-Mota, R., Cherchi, A., Gan, T. Y., Gergis, J., Jiang, D., Khan, A., Pokam Mba, W., Rosenfeld, D., Tierney, J., and Zolina, O.: Water Cycle Changes, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., , and Zhou, B., Cambridge University Press,
- 370 2021.

- Dowsett, H.: Faunal re-evaluation of Mid-Pliocene conditions in the western equatorial Pacific, *Micropaleontology*, 53, 447–456, <https://doi.org/10.2113/gsmicropal.53.6.447>, 2007.
- Dowsett, H. and Poore, R.: Pliocene sea surface temperatures of the north atlantic ocean at 3.0 Ma, *Quaternary Science Reviews*, 10, 189–204, [https://doi.org/10.1016/0277-3791\(91\)90018-P](https://doi.org/10.1016/0277-3791(91)90018-P), 1991.
- 375 Dowsett, H. J., Barron, J. A., Poore, R. Z., Thompson, R. S., Cronin, T. M., Ishman, S. E., and Willard, D. A.: Middle Pliocene Paleoenvironmental Reconstruction: PRISM 2, U.S. Geological Survey Open File Report 99-535, <http://pubs.usgs.gov/of/1999/of99-535/>, 1999.
- Dowsett, H. J., Foley, K. M., Stoll, D. K., Chandler, M. A., Sohl, L. E., Bentsen, M., Otto-Bliesner, B. L., Bragg, F. J., Chan, W.-L., Contoux, C., Dolan, A. M., Haywood, A. M., Jonas, J. A., Jost, A., Kamae, Y., Lohmann, G., Lunt, D. J., Nisancioglu, K. H., Abe-Ouchi, A., Ramstein, G., Riesselman, C. R., Robinson, M. M., Rosenbloom, N. A., Salzmann, U., Stepanek, C., Strother, S. L., Ueda, H.,  
380 Yan, Q., and Zhang, Z.: Sea Surface Temperature of the mid-Piacenzian Ocean: A Data-Model Comparison, *Scientific Reports*, 3, 2013, <https://doi.org/10.1038/srep02013>, 2013.
- Fedorov, A. V., Brierley, C. M., Lawrence, K. T., Liu, Z., Dekens, P. S., and Ravelo, A. C.: Patterns and mechanisms of early Pliocene warmth, *Nature*, 496, 43–49, <https://doi.org/10.1038/nature12003>, 2013.
- Feng, R., Otto-Bliesner, B. L., Xu, Y., Brady, E., Fletcher, T., and Ballantyne, A.: Contributions of aerosol-cloud interactions to mid-Piacenzian seasonally sea ice-free Arctic Ocean, *Geophysical Research Letters*, 46, 9920–9929, <https://doi.org/https://doi.org/10.1029/2019GL083960>, 2019.
- Feng, R., Bhattacharya, T., Otto-Bliesner, B. L., Brady, E. C., Haywood, A. M., Tindall, J. C., Hunter, S. J., Abe-Ouchi, A., Chan, W.-L., Kageyama, M., Contoux, C., Guo, C., Li, X., Lohmann, G., Stepanek, C., Tan, N., Zhang, Q., Zhang, Z., Han, Z., Williams, C. J. R., Lunt, D. J., Dowsett, H. J., Chandan, D., and Peltier, W. R.: Past terrestrial hydroclimate sensitivity controlled by Earth system feedbacks,  
390 *Nature Communications*, 13, 1306, <https://doi.org/10.1038/s41467-022-28814-7>, 2022.
- Foley, K. M. and Dowsett, H. J.: Community sourced mid-Piacenzian sea surface temperature (SST) data, US Geological Survey data release, <https://doi.org/10.5066/P9YP3DTV>, last access: 21 August 2023, 2019.
- Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z.-L., and Zhang, M.: The community climate system model version 4, *Journal of Climate*, 24, 4973–4991, <https://doi.org/10.1175/2011JCLI4083.1>, 2011.
- 395 Gulev, S. K., Thorne, P., Ahn, J., Dentener, F., Domingues, C., Gerland, S., Gong, D., Kaufman, D., Nnamchi, H., Quaas, J., Rivera, J., Sathyendranath, S., Smith, S., Trewin, B., von Shuckmann, K., and Vose, R.: Changing State of the Climate System, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., , and Zhou, B.,  
400 Cambridge University Press, Cambridge, United Kingdom and New York, U.S.A., 2021.
- Han, Z., Zhang, Q., Li, Q., Feng, R., Haywood, A. M., Tindall, J. C., Hunter, S. J., Otto-Bliesner, B. L., Brady, E. C., Rosenbloom, N., Zhang, Z., Li, X., Guo, C., Nisancioglu, K. H., Stepanek, C., Lohmann, G., Sohl, L. E., Chandler, M. A., Tan, N., Ramstein, G., Baatsen, M. L. J., von der Heydt, A. S., Chandan, D., Peltier, W. R., Williams, C. J. R., Lunt, D. J., Cheng, J., Wen, Q., and Burls, N. J.: Evaluating  
405 the large-scale hydrological cycle response within the Pliocene Model Intercomparison Project Phase 2 (PlioMIP2) ensemble, *Climate of the Past*, 17, 2537–2558, <https://doi.org/10.5194/cp-17-2537-2021>, 2021.


- Haywood, A. M., Dowsett, H. J., Otto-Bliesner, B., Chandler, M. A., Dolan, A. M., Hill, D. J., Lunt, D. J., Robinson, M. M., Rosenbloom, N., Salzmann, U., and Sohl, L. E.: Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 1), *Geoscientific Model Development*, 3, 227–242, <https://doi.org/10.5194/gmd-3-227-2010>, 2010.
- 410 Haywood, A. M., Dowsett, H. J., Robinson, M. M., Stoll, D. K., Dolan, A. M., Lunt, D. J., Otto-Bliesner, B., and Chandler, M. A.: Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 2), *Geoscientific Model Development*, 4, 571–577, <https://doi.org/10.5194/gmd-4-571-2011>, 2011.
- Haywood, A. M., Hill, D. J., Dolan, A. M., Otto-Bliesner, B. L., Bragg, F., Chan, W.-L., Chandler, M. A., Contoux, C., Dowsett, H. J., Jost, A., Kamae, Y., Lohmann, G., Lunt, D. J., Abe-Ouchi, A., Pickering, S. J., Ramstein, G., Rosenbloom, N. A., Salzmann, U., Sohl, L.,
- 415 Stepanek, C., Ueda, H., Yan, Q., and Zhang, Z.: Large-scale features of Pliocene climate: results from the Pliocene Model Intercomparison Project, *Climate of the Past*, 9, 191–209, <https://doi.org/10.5194/cp-9-191-2013>, 2013.
- Haywood, A. M., Dowsett, H. J., and Dolan, A. M.: Integrating geological archives and climate models for the mid-Pliocene warm period, *Nature Communications*, 7, 10646, <https://doi.org/10.1038/ncomms10646>, 2016a.
- Haywood, A. M., Dowsett, H. J., Dolan, A. M., Rowley, D., Abe-Ouchi, A., Otto-Bliesner, B., Chandler, M. A., Hunter, S. J., Lunt, D. J.,
- 420 Pound, M., and Salzmann, U.: The Pliocene Model Intercomparison Project (PlioMIP) Phase 2: Scientific objectives and experimental design, *Climate of the Past*, 12, 663–675, <https://doi.org/10.5194/cp-12-663-2016>, 2016b.
- Haywood, A. M., Tindall, J. C., Dowsett, H. J., Dolan, A. M., Foley, K. M., Hunter, S. J., Hill, D. J., Chan, W.-L., Abe-Ouchi, A., Stepanek, C., Lohmann, G., Chandan, D., Peltier, W. R., Tan, N., Contoux, C., Ramstein, G., Li, X., Zhang, Z., Guo, C., Nisancioglu, K. H., Zhang, Q., Li, Q., Kamae, Y., Chandler, M. A., Sohl, L. E., Otto-Bliesner, B. L., Feng, R., Brady, E. C., von der Heydt, A. S., Baatsen, M. L. J.,
- 425 and Lunt, D. J.: The Pliocene Model Intercomparison Project Phase 2: large-scale climate features and climate sensitivity, *Climate of the Past*, 16, 2095–2123, <https://doi.org/10.5194/cp-16-2095-2020>, 2020.
- Held, I. and Soden, B.: Robust responses of the hydrological cycle to global warming, *Journal Of Climate*, 19, 5686–5699, <https://doi.org/10.1175/JCLI3990.1>, 2006.
- Holland, M. M., Bailey, D. A., Briegleb, B. P., Light, B., and Hunke, E.: Improved sea ice shortwave radiation physics in CCSM4: The impact
- 430 of melt ponds and aerosols on Arctic sea ice, *Journal of Climate*, 25, 1413 – 1430, <https://doi.org/10.1175/JCLI-D-11-00078.1>, 2012.
- Huang, B., Thorne, P. W., Banzon, V. F., Boyer, T., Chepurin, G., Lawrimore, J. H., Menne, M. J., Smith, T. M., Vose, R. S., and Zhang, H.-M.: Extended Reconstructed Sea Surface Temperature, Version 5 (ERSSTv5): Upgrades, Validations, and Intercomparisons, *Journal of Climate*, 30, 8179 – 8205, <https://doi.org/10.1175/JCLI-D-16-0836.1>, 2017.
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J.-F., Large, W. G., Lawrence, D., Lindsay, K.,
- 435 Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: A Framework for Collaborative Research, *Bulletin of the American Meteorological Society*, 94, 1339–1360, <https://doi.org/10.1175/BAMS-D-12-00121.1>, 2013.
- Hwang, Y.-T., Frierson, D. M. W., and Kang, S. M.: Anthropogenic sulfate aerosol and the southward shift of tropical precipitation in the late 20th century, *Geophysical Research Letters*, 40, 2845–2850, <https://doi.org/https://doi.org/10.1002/grl.50502>, 2013.
- 440 IPCC: Summary for Policymakers, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., , and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, U.S.A., 2021.

- 445 Kay, J. E., Hillman, B. R., Klein, S. A., Zhang, Y., Medeiros, B., Pincus, R., Gettelman, A., Eaton, B., Boyle, J., Marchand, R., and Ackerman, T. P.: Exposing Global Cloud Biases in the Community Atmosphere Model (CAM) Using Satellite Observations and Their Corresponding Instrument Simulators, *Journal of Climate*, 25, 5190 – 5207, <https://doi.org/https://doi.org/10.1175/JCLI-D-11-00469.1>, 2012.
- Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liou, S. C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, 450 K., and Van Vuuren, D. P.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: Methodology and application, *Atmospheric Chemistry and Physics*, 10, 7017–7039, <https://doi.org/10.5194/acp-10-7017-2010>, 2010.
- Lambert, F. and Albani, S.: mineral dust in PmiP simulations: a short review, *PAGES*, 29, 86–87, <https://doi.org/doi.org/10.22498/pages.29.2.86>, 2021.
- Lee, J. Y., Marotzke, J., Bala, G., Cao, L., Corti, S., Dunne, J., Engelbrecht, F., Fischer, E., Fyfe, J., Jones, C., Maycock, A., Mutemi, 455 J., Ndiaye, O., Panickal, S., and Zhou, T.: Future Global Climate: Scenario-Based Projections and Near-Term Information., in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., , and Zhou, B., Cambridge University Press, 2021.
- 460 Li, X., Jiang, D., Tian, Z., and Yang, Y.: Mid-Pliocene global land monsoon from PlioMIP1 simulations, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 512, 56–70, <https://doi.org/https://doi.org/10.1016/j.palaeo.2018.06.027>, 2018.
- Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J. F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, 465 D.: Toward a minimal representation of aerosols in climate models: Description and evaluation in the Community Atmosphere Model CAM5, *Geoscientific Model Development*, 5, 709–739, <https://doi.org/10.5194/gmd-5-709-2012>, 2012.
- Lunt, D. J., Haywood, A. M., Schmidt, G. A., Salzmann, U., Valdes, P. J., and Dowsett, H. J.: Earth system sensitivity inferred from Pliocene modelling and data, *Nature Geoscience*, 3, 60–64, <https://doi.org/10.1038/ngeo706>, 2010.
- Martinez, J. M.: Managing scientific uncertainty in medical decision making: the case of the advisory committee on immunization practices., *The Journal of medicine and philosophy*, 37, 6–27, <https://doi.org/10.1093/jmp/jhr056>, 2012.
- 470 Meehl, G. A., Washington, W. M., Arblaster, J. M., Hu, A., Teng, H., Kay, J. E., Gettelman, A., Lawrence, D. M., Sanderson, B. M., and Strand, W. G.: Climate change projections in CESM1(CAM5) compared to CCSM4, *Journal of Climate*, 26, 6287–6308, <https://doi.org/10.1175/JCLI-D-12-00572.1>, 2013.
- Nie, J., Stevens, T., Song, Y., King, J. W., Zhang, R., Ji, S., Gong, L., and Cares, D.: Pacific freshening drives Pliocene cooling and Asian monsoon intensification, *Scientific Reports*, 4, 5474, <https://doi.org/10.1038/srep05474>, 2014.
- 475 O'Brien, C. L., Foster, G. L., Martínez-Botí, M. A., Abell, R., Rae, J. W., and Pancost, R. D.: High sea surface temperatures in tropical warm pools during the Pliocene, *Nature Geoscience*, 7, 606–611, <https://doi.org/10.1038/ngeo2194>, 2014.
- Oleson, K.: Technical Description of the Community Land Model (CLM), NCAR Technical Note, TN-478+STR, <https://doi.org/10.5065/D6RR1W7M>, 2010.
- Pagani, M., Liu, Z., Lariviere, J., and Ravelo, A. C.: High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations, *Nature Geoscience*, 3, 27–30, <https://doi.org/10.1038/ngeo724>, 2010.
- 480 Ridley, H. E., Asmerom, Y., Baldini, J. U., Breitenbach, S. F., Aquino, V. V., Pruffer, K. M., Culleton, B. J., Polyak, V., Lechleitner, F. A., Kennett, D. J., Zhang, M., Marwan, N., Macpherson, C. G., Baldini, L. M., Xiao, T., Peterkin, J. L., Awe, J., and Haug, G. H.: Aerosol forcing




- of the position of the intertropical convergence zone since ad 1550, *Nature Geoscience*, 8, 195–200, <https://doi.org/10.1038/ngeo2353>, 2015.
- 485 Rosenbloom, N. A., Otto-Bliesner, B. L., Brady, E. C., and Lawrence, P. J.: Simulating the mid-Pliocene Warm Period with the CCSM4 model, *Geoscientific Model Development*, 6, 549–561, <https://doi.org/10.5194/gmd-6-549-2013>, 2013.
- Sagoo, N. and Storelvmo, T.: Testing the sensitivity of past climates to the indirect effects of dust, *Geophysical Research Letters*, 44, 5807–5817, <https://doi.org/10.1002/2017GL072584>, 2017.
- Salzmann, U., Haywood, A. M., Lunt, D. J., Valdes, P. J., and Hill, D. J.: A new global biome reconstruction and data-model comparison for the Middle Pliocene, *Global Ecology and Biogeography*, 17, 432–447, <https://doi.org/10.1111/j.1466-8238.2008.00381.x>, 2008.
- 490 Samset, B. H., Sand, M., Smith, C. J., Bauer, S. E., Forster, P. M., Fuglestedt, J. S., Osprey, S., and Schleussner, C. F.: Climate Impacts From a Removal of Anthropogenic Aerosol Emissions, *Geophysical Research Letters*, 45, 1020–1029, <https://doi.org/10.1002/2017GL076079>, 2018.
- Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmospheric Chemistry and Physics*, 11, 1101–1116, <https://doi.org/10.5194/acp-11-1101-2011>, 2011.
- 495 Tegen, I. and Schepanski, K.: Climate Feedback on Aerosol Emission and Atmospheric Concentrations, *Current Climate Change Reports*, 4, 1–10, <https://doi.org/10.1007/s40641-018-0086-1>, 2018.
- Trenberth, K. E. and Guillemot, C. J.: Evaluation of the global atmospheric moisture budget as seen from analyses, *Journal of Climate*, 8, 2255 – 2272, 1995.
- 500 Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the Atmospheric Sciences*, 34, 1149–1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.
- Unger, N. and Yue, X.: Strong chemistry-climate feedbacks in the Pliocene, *Geophysical Research Letters*, 41, 527–533, <https://doi.org/10.1002/2013GL058773>, 2014.
- Voigt, A., Pincus, R., Stevens, B., Bony, S., Boucher, O., Bellouin, N., Lewinschal, A., Medeiros, B., Wang, Z., and Zhang, H.: Fast and slow shifts of the zonal-mean intertropical convergence zone in response to an idealized anthropogenic aerosol, *Journal of Advances in Modeling Earth Systems*, 9, 870–892, <https://doi.org/10.1002/2016MS000902>, 2017.
- 505 Wang, B. and Ding, Q.: Global monsoon: Dominant mode of annual variation in the tropics, *Dynamics of Atmospheres and Oceans*, 44, 165–183, <https://doi.org/https://doi.org/10.1016/j.dynatmoce.2007.05.002>, 2008.
- Wang, C., W.L., L., and Chou, C.: Climate Effects of Anthropogenic Aerosol Forcing on Tropical Precipitation and Circulations, *Journal of Climate*, 32, 5275 – 5287, <https://doi.org/10.1175/JCLI-D-18-0641.1>, 2019.
- 510 Wang, P. X., Wang, B., Cheng, H., Fasullo, J., Guo, Z. T., Kiefer, T., and Liu, Z. Y.: The global monsoon across timescales: coherent variability of regional monsoons, *Climate of the Past*, 10, 2007–2052, <https://doi.org/10.5194/cp-10-2007-2014>, 2014.
- Wara, M. W.: Permanent El Nino-Like Conditions During the Pliocene Warm Period, *Science*, 309, 758–761, <https://doi.org/10.1126/science.1112596>, 2005.
- 515 Wilcox, L. J., Liu, Z., Samset, B. H., Hawkins, E., Lund, M. T., Nordling, K., Undorf, S., Bollasina, M., Ekman, A. M. L., Krishnan, S., Merikanto, J., and Turner, A. G.: Accelerated increases in global and Asian summer monsoon precipitation from future aerosol reductions, *Atmospheric Chemistry and Physics*, 20, 11955–11977, <https://doi.org/10.5194/acp-20-11955-2020>, 2020.
- Xie, S.-P., Deser, C., Vecchi, G. A., Ma, J., Teng, H., and Wittenberg, A. T.: Global Warming Pattern Formation: Sea Surface Temperature and Rainfall, *Journal of Climate*, 23, 966 – 986, <https://doi.org/https://doi.org/10.1175/2009JCLI3329.1>, 2010.


- 520 Xiong, S., Ding, Z., Zhu, Y., Zhou, R., and Lu, H.: A ~6Ma chemical weathering history, the grain size dependence of chemical weathering intensity, and its implications for provenance change of the Chinese loess–red clay deposit, *Quaternary Science Reviews*, 29, 1911–1922, <https://doi.org/https://doi.org/10.1016/j.quascirev.2010.04.009>, 2010.
- Zhang, R., Yan, Q., Zhang, Z. S., Jiang, D., Otto-Bliesner, B. L., Haywood, A. M., Hill, D. J., Dolan, A. M., Stepanek, C., Lohmann, G., Contoux, C., Bragg, F., Chan, C. L., Chandler, M. A., Jost, A., Kamae, Y., Abe-Ouchi, A., Ramstein, G., Rosenbloom, N. A.,
- 525 Sohl, L., and Ueda, H.: Mid-Pliocene East Asian monsoon climate simulated in the PlioMIP, *Climate of the Past*, 9, 2085–2099, <https://doi.org/10.5194/cp-9-2085-2013>, 2013.
- Zhang, Y. G., Pagani, M., and Liu, Z.: A 12-Million-Year Temperature History of the Tropical Pacific Ocean, *Science*, 344, 84–87, <https://doi.org/10.1126/science.1246172>, 2014.
- Zubakov, V. and Borzenkova, I.: Pliocene palaeoclimates: Past climates as possible analogues of mid-twenty-first century climate, *Palaeo-*
- 530 *geography, Palaeoclimatology, Palaeoecology*, 65, 35–49, [https://doi.org/https://doi.org/10.1016/0031-0182\(88\)90110-1](https://doi.org/https://doi.org/10.1016/0031-0182(88)90110-1), 1988.



**Figure 1. Changes in (a) mean annual surface temperature (TS) in °C and (b) precipitation (PR) in mm d<sup>-1</sup> (Plio\_Pristine - PI).** Shaded circles are reconstructed anomalies of (a) SSTs (Foley and Dowsett, 2019) and (b) precipitation (Salzmann et al., 2008). Triangles mark the sign of mPWP hydroclimate changes (mean annual precipitation minus evaporation) published in (Feng et al., 2022) [Feng et al. \(2022\)](#). See section 2.3.2 for details.



**Figure 2. Zonal averaged changes in mean annual (a) surface temperature in  $^{\circ}\text{C}$  and (b) precipitation in  $\text{mm d}^{-1}$ .** In all panels, black solid lines show the simulated anomalies (Plio\_Pristine - PI). Dotted lines represent the multi-model means of the PlioMIP1 (red, Haywood et al., 2013) and PlioMIP2 (blue, Haywood et al., 2020) respectively, with shaded areas showing the model spread, i.e. the standard deviation of the ensemble. Green squares show the site-level reconstructions in Fig. 1, and dots show the corresponding grid-level anomalies with error bars showing the range of the PlioMIPs.




**Figure 3. Zonal-averaged change in (a) radiation flux and (b) cloud forcing in  $W m^{-2}$  after removal of anthropogenic emissions (Plio\_Pristine-P)**


(a) Change in net shortwave flux at top of model and net longwave flux at top of model under full-sky (red, blue) and clear-sky conditions (orange, cyan). (b) Change in shortwave cloud forcing (green) and longwave cloud forcing (brown). The corresponding spatial change in cloud shortwave and longwave forcings shown in panel b are displayed in Fig. A1a,b respectively.

Zonal averaged change in (a) shortwave radiation flux and (b) cloud radiative effect in  $W m^{-2}$  at top of model (TOM) after removal

TOM radiation fluxes are used to show change at top of atmosphere (TOA), because the results are not sensitive to TOM or TOA. (a) Clear-sky shortwave radiation (orange) refers to TOM radiation flux in the absence of clouds, and full-sky shortwave radiation flux (black) is with clouds. Shortwave cloud radiative effect (cyan) is the difference in shortwave radiation flux between full-sky and clear-sky conditions. (b) Cloud radiative effect (black) is the sum of downwelling shortwave (green) and longwave (purple) cloud radiative effect.




**Figure 4. Change in (a) aerosol optical depth 550 nm and (b) cloud condensation nuclei concentration as supersaturation 0.1% ( $\text{cm}^{-3}$ ) after removal of anthropogenic emissions (Plio\_Pristine - Plio\_Polluted). Change in (a) aerosol optical depth 550 nm and (b) cloud condensation nuclei concentration as supersaturation 0.1% ( $\text{cm}^{-3}$ ) after removal of anthropogenic emissions (Plio\_Pristine - Plio\_Polluted).**




**Figure 5. Annual, DJF and JJA mean surface temperature change in °C (a, c, e) and precipitation change in mm d<sup>-1</sup> (b,d,f) after removal of anthropogenic emissions (Plio\_Pristine - Plio\_Polluted) Annual, DJF and JJA mean surface temperature change in °C (a, c, e) and precipitation change in mm d<sup>-1</sup> (b,d,f) after removal of anthropogenic emissions (Plio\_Pristine - Plio\_Polluted).**







**Figure 6. Zonal averaged change in high-, mid- and low- level clouds.**[\(Plio Pristine -Plio Polluted\)](#). The corresponding spatial relative change in cloud fractions are given in Fig. A2.




**Figure 7. Zonal mean changes in (a) vertical velocity ( $10^{-3} \text{ Pa s}^{-1}$ ) and (b) meridional stream function ( $10^9 \text{ kg s}^{-1}$ ) by removing anthropogenic from the mPWP atmosphere (Plio\_Pristine - Plio\_Polluted). Overlaid contours show the climate mean for the Plio\_Pristine simulation (solid for positive values and dashed for negative values).**




**Figure 8.** Changes in monsoon summer rain rate ( $\text{mm d}^{-1}$ ) during the mPWP (Plio\_Pristine - PI, panel a) and by removing anthropogenic from the atmosphere (Plio\_Pristine - Plio\_Polluted, panel b). Black, red and blue contours represent the boundary of the global monsoon domain following Wang et al. (2014) in PI, Plio\_Pristine and Plio\_Polluted respectively.



**Figure 9. Zonal-averaged change in (a) cloud droplet concentration in  $\text{m}^{-2}$  and (b) cloud liquid water path in  $\text{kg m}^{-2}$ . Zonal averaged change (Plio\_Pristine - Plio\_Polluted) in (a) cloud droplet concentration in  $\text{m}^{-2}$  and (b) cloud liquid water path in  $\text{kg m}^{-2}$ . The corresponding spatial relative change are given in Fig. A3.**



**Figure 10.** Changes in annual and seasonal sea level pressure (hPa) and surface wind ( $\text{m s}^{-1}$ ) during the mPWP (Plio\_Pristine - PI, left column) and after ~~removal~~ removal of anthropogenic aerosols (Plio\_Pristine - Plio\_Polluted, right column).



**Figure 11. Relative importance of removing anthropogenic aerosols and mPWP boundary conditions on mPWP zonal mean precipitation change**

(a) Ratio of change in annual, JJA and DJF precipitation change computed as  $((\text{Plio\_Pristine} - \text{Plio\_Polluted}) / (\text{Plio\_Pristine} - \text{PI}))$ . Yellow shading highlights where  $-1 < \text{ratio} < 1$ . (b) Comparison between latitude average mean precipitation change induced by PlioMIP boundary conditions and by aerosol scenarios within the tropics (light colors) and outside of the tropics (darker colors). **Effect of removing anthropogenic aerosols and mPWP boundary conditions on mPWP zonal mean precipitation change and their relative**

Left column shows zonal averaged (a) annual, (c) DJF and (e) JJA mean precipitation change induced by removing anthropogenic aerosols (dashed) and mPWP boundary conditions (solid). Left column shows ratio of change in (b) annual, (d) DJF and (f) JJA precipitation change computed as  $((\text{Plio\_Pristine} - \text{Plio\_Polluted}) / (\text{Plio\_Pristine} - \text{PI}))$ . Yellow shading highlights where  $-1 < \text{ratio} < 1$ .




Figure A1. Change in (a) shortwave cloud forcing and (b) longwave cloud forcing in  $W m^{-2}$  (Plio\_Pristine - Plio\_Polluted).

## Appendix A: Cloud properties







Figure A2. FL Spatial relative change in vertically-integrated (a) high, (b) mid-level, (c) low and (d) total cloud ( $(\text{Plio\_Pristine and Plio\_Polluted}) \cdot 100 / \text{Plio\_Pristine}$ ), and vertical change in (e) cloud fraction as difference between Plio\_Pristine and Plio\_Polluted. Spatial relative change in vertically-integrated (a) high-, (b) mid- and (c) low- level cloud ( $(\text{Plio\_Pristine and Plio\_Polluted}) \cdot 100 / \text{Plio\_Pristine}$ ).



**Figure A3.** Spatial relative change in (a) vertically-integrated droplet concentration and (b) grid-box cloud liquid water path as compared to Plio\_Pristine (i.e.  $(\text{Plio\_Pristine} - \text{Plio\_Polluted}) * 100 / \text{Plio\_Pristine}$ ).