

The emission of CO from tropical ~~rain forest~~ rainforest soils

Hella van Asperen^{1,2}, Thorsten Warneke¹, Alessandro Carioca de Araújo^{3,4}, Bruce Forsberg³, Sávio José Filgueiras Ferreira⁵, Thomas Röckmann⁶, Carina van der Veen⁶, Sipko Bulthuis⁷, Leonardo Ramos de Oliveira³, Thiago de Lima Xavier³, Jailson da Mata³, Marta de Oliveira Sá⁸, Paulo Ricardo Teixeira³, Julie Andrews de França e Silva³, Susan Trumbore², and Justus Notholt¹

¹Institute of Environmental Physics (IUP), University of Bremen, Otto-Hahn-Allee 1, Bremen, 28359, Germany

²Max Planck Institute for Biogeochemistry (MPI-BGC), Hans-Knöll-Strasse 10, Jena, 07745, Germany

³Programa de Grande Escala da Biosfera-Atmosfera na Amazônia (LBA), Instituto Nacional de Pesquisas da Amazônia (INPA), Av. André Araújo, 2936, Petrópolis, AM 69067-375, Manaus, Brazil

⁴Brazilian Agricultural Research Corporation (EMBRAPA), Embrapa Amazônia Oriental, Tv. Dr. Enéas Piheiro, s/n, Marco, PA 66095-903, Caixa postal 48, Belém, Brazil

⁵Coordenação de Dinâmica Ambiental (CODAM), Instituto Nacional de Pesquisas da Amazônia (INPA), Av. André Araújo, 2936, Petrópolis, AM 69067-375, Manaus, Brazil

⁶Institute for Marine and Atmospheric Research Utrecht, Utrecht University, Princetonplein 5, 3584, Utrecht, Netherlands

⁷Max Planck Institute for Chemistry, Hahn-Meitner-Weg 1, Mainz, 55128, Germany

⁸Postgraduate Program in Climate and Environment (CLIAMB) - INPA/UEA, Instituto Nacional de Pesquisas da Amazônia (INPA), Av. André Araújo, 2936, Petrópolis, AM 69067-375, Manaus, Brazil

Correspondence: Hella van Asperen (hasperen@bgc-jena.mpg.de)

Abstract. Soil CO fluxes represent a net balance between biological soil CO uptake and abiotic soil and (senescent) plant CO production. Studies largely from temperate and boreal forests indicate that soils serve as a net sink for CO, but to date uncertainty remains about the role of tropical ~~rain forest~~ rainforest soils. Here we report the first direct measurements of soil CO fluxes in a tropical ~~rain forest~~ rainforest. We compare with estimates of net ecosystem CO fluxes derived from accumulation of CO at night under stable atmospheric conditions. Further, we used laboratory experiments to demonstrate the importance of temperature on net soil CO fluxes. Net soil surface CO fluxes ranged from -0.19 to $3.36 \text{ nmol m}^{-2} \text{ s}^{-1}$, averaging $\sim 1 \text{ nmol CO m}^{-2} \text{ s}^{-1}$. Fluxes varied with season and topographic location, with highest fluxes measured in the dry season in a seasonally inundated valley. Ecosystem CO fluxes estimated from nocturnal canopy air profiles, which showed CO mixing ratios that consistently decreased with height, ranged between 0.3 and $2.0 \text{ nmol CO m}^{-2} \text{ s}^{-1}$. A canopy layer budget method, using the nocturnal increase in CO, estimated similar flux magnitudes (1.1 to $2.3 \text{ nmol CO m}^{-2} \text{ s}^{-1}$). In the wet season, a greater valley ecosystem CO production was observed in comparison to measured soil valley CO fluxes, suggesting a contribution of the valley stream to overall CO emissions. Laboratory incubations demonstrated a clear increase in CO production with temperature that was also observed in field fluxes, though high correlations between soil temperature and moisture limit our ability to interpret the field relationship. At a common temperature ($25 \text{ }^\circ\text{C}$), expected plateau and valley senescent leaf CO production was small (0.012 and $0.002 \text{ nmol CO m}^{-2} \text{ s}^{-1}$) in comparison to expected soil material CO emission ($\sim 0.9 \text{ nmol CO m}^{-2} \text{ s}^{-1}$). Based on our field and laboratory observations, we expect that tropical ~~rain forest~~ rainforest ecosystems are a net source of CO, with thermal degradation-induced soil emissions likely being the main contributor to ecosystem CO emissions. Extrapolating our first observation-based tropical ~~rain forest~~ rainforest soil emission estimate of $\sim 1 \text{ nmol m}^{-2} \text{ s}^{-1}$, a global

tropical ~~rain-forest~~ rainforest soil emission of $\sim 16.0 \text{ Tg CO yr}^{-1}$ is estimated. Nevertheless, total ecosystem CO emissions might be higher, since valley streams and inundated areas might represent local CO emission hot spots. To further improve tropical forest ecosystem CO emission estimates, more in-situ tropical forest soil and ecosystem CO flux measurements are essential.

1 Introduction

Carbon monoxide (CO) is a trace gas in the atmosphere. It is the most important sink for the hydroxyl (OH) radical, which also serves as a sink for methane (CH_4). Thus, an increase in CO emissions will directly affect the atmospheric concentrations of CH_4 , making CO an indirect greenhouse gas, with a possible indirect radiative forcing larger than N_2O (Szopa et al., 2021). Anthropogenic activities, such as ~~(incomplete)~~ combustion of fossil fuel and biomass, contribute strongly to global CO emissions, and CO concentrations in urban areas are usually higher than in rural areas (~~Zheng et al., 2019; Seinfeld and Pandis, 2016~~) (Seinfeld and Pandis, 2016; Zheng et al., 2019). Due to its short atmospheric lifetime of 50 days, spatial differences between regions can be large, and concentrations in the northern hemisphere are generally higher than in the southern hemisphere (~~Szopa et al., 2021; Seinfeld and Pandis, 2016~~) (Seinfeld and Pandis, 2016; Szopa et al., 2021). Besides direct anthropogenic emissions, CO is also produced by atmospheric oxidation sources, such as the in-situ oxidation of methane and hydrocarbons, or can be emitted by (partly) natural sources such as forest fires, ocean emissions, the degradation of chlorophyll, and abiotic degradation of organic matter (~~Sanderson, 2002; Szopa et al., 2021; Seinfeld and Pandis, 2016~~) (Sanderson, 2002; Seinfeld and Pandis, 2016; Szopa et al., 2021). The major natural sinks of carbon monoxide are the tropospheric oxidation with OH ($>80\%$), the uptake by soils ($\sim 10\text{-}15\%$), and the removal in the stratosphere ($\sim 5\%$) (~~Seinfeld and Pandis, 2016; King and Weber, 2007; Bartholomew and Alexander, 1979; Conrad and Seiler, 1982; King and Weber, 2007; Sanderson, 2002; Seinfeld and Pandis, 2016~~) (Bartholomew and Alexander, 1979; Conrad, 1996; Khalil and Rasmussen, 1990; King and Weber, 2007; Sanderson, 2002; Seinfeld and Pandis, 2016).

On the ecosystem level, the sources and sinks of CO are poorly understood. Soils can act as net sources as well as net sinks of CO (Conrad, 1996). Most likely, the main process involved in soil CO uptake is the oxidation of CO to CO_2 , or the reduction to CH_4 by soil bacteria or soil enzymes (~~Bartholomew and Alexander, 1979; Conrad, 1996; Ingersoll et al., 1974; Spratt and Hubbard, 1981; Seinfeld and Pandis, 2016~~) (Bartholomew and Alexander, 1979; Conrad, 1996; Ingersoll et al., 1974; Spratt and Hubbard, 1981; Whalen and Reeburgh, 2001; Yonemura et al., 2000). Soil CO consumption was reported to be poorly related to temperature (Conrad and Seiler, 1985), and more related to soil diffusivity (~~Sanhueza et al., 1994; Conrad and Seiler, 1982; Kisselle et al., 2002~~) (Conrad and Seiler, 1982; Kisselle et al., 2002; Sanhueza et al., 1994). Soil CO emissions are thought to be mostly of non-biological origin, namely photo-degradation (~~Bruhn et al., 2013; Pihlatie et al., 2016; Seinfeld and Pandis, 2016~~) (Bruhn et al., 2013; Derendorp et al., 2011; Lee et al., 2012; Pihlatie et al., 2016; Schade et al., 1999; Tarr et al., 1995) and thermal degradation (~~Conrad and Seiler, 1980; Derendorp et al., 2011; Asperen et al., 2015b; Conrad and Seiler, 1982; Lee et al., 2012; Yonemura et al., 2000~~) (Asperen et al., 2015b; Conrad and Seiler, 1980, 1982; Derendorp et al., 2011; Lee et al., 2012; Yonemura et al., 2000). Besides emissions associated with abiotic degradation of organic matter, living plants are also known to emit small amounts of CO

(Bruhn et al., 2013; Kirchhoff and Marinho, 1990; Tarr et al., 1995)(Bruhn et al., 2013; Kirchhoff and Marinho, 1990; Tarr et al., 1995)

. However, emissions from senescent plant material are 5 to 10 times greater than those observed from photosynthesising leaf material (Derendorp et al., 2011; Schade et al., 1999; Tarr et al., 1995)(Derendorp et al., 2011; Schade et al., 1999; Tarr et al., 1995)

55 .

Soil CO fluxes thus represent the net balance between biological soil CO uptake and abiotic soil and (senescent) plant CO production (Whalen and Reeburgh, 2001; Liu et al., 2018; Potter et al., 1996; Asperen et al., 2015b; Pihlatie et al., 2016; Constant et al., 2008; Asperen et al., 2015b; Constant et al., 2008; Liu et al., 2018; Pihlatie et al., 2016; Potter et al., 1996; Whalen and Reeburgh, 2001)

60 . Besides temperature and radiation, it has been observed that the net flux is dependent on, among others, soil water content, soil organic carbon, land use type, and nutrients (King, 2000; King and Hungria, 2002; Conrad and Seiler, 1985; Funk et al., 1994; Göttsche et al., 2000; Conrad and Seiler, 1985; Funk et al., 1994; Göttsche et al., 2000; King, 2000; King and Hungria, 2002; Moxley and Smith, 1998; Yonemura et al., 2000)

. Due to its dependency on environmental factors, the net CO flux balance might shift diurnally and seasonally. Existing measurements of diurnal cycles mostly show a shift towards uptake during nighttime hours, and emission during daytime hours (Asperen et al., 2015b; Sanhueza et al., 1994; Scharffe et al., 1990; Schade et al., 1999)(Asperen et al., 2015b; Sanhueza et al., 1994; Scharffe et al., 1990)

65 . The few long term CO flux studies found a similar pattern seasonally, with increased uptake during colder periods, and more emission during warmer periods (Constant et al., 2008; Cowan et al., 2018; Pihlatie et al., 2016)(Constant et al., 2008; Cowan et al., 2018)

70 Previous CO flux measurements have been done in boreal ecosystems (Constant et al., 2008; Laasonen, 2021; Pihlatie et al., 2016; Whalen and Reeburgh, 2001), temperate zones (Cowan et al., 2018; Conrad et al., 1988; Göttsche et al., 2000)(Conrad et al., 1988; Cowan et al., 2018; Göttsche et al., 2000), and arid and (sub-)tropical ecosystems (Sanhueza et al., 1994; Kisselle et al., 2002; King and Hungria, 2002; King, 2000; Scharffe et al., 1990; Asperen et al., 2015b; King, 2000; King and Hungria, 2002; Kisselle et al., 2002; Sanhueza et al., 1994; Scharffe et al., 1990)

75 , but we are aware of no previous CO flux measurements from tropical ~~rain forests~~rainforests. Because of this, the net CO flux of tropical forest soils predicted using global models remains highly uncertain even as to sign: ~~while~~While Potter et al. (1996) modelled that tropical soils are likely a source of CO, thereby implying that abiotic emission dominates over soil biological CO uptake, a more recent modelling study suggested that tropical soils are possibly a net sink of CO (Liu et al., 2018). This discrepancy shows the need for in-situ observation of soil and ecosystem CO fluxes in tropical ~~rain forests~~rainforests.

80

In this study, we present results from 2 intensive measurement campaigns in a tropical ~~rain forest in central Amazon~~rainforest in central Amazonia. During wet and dry season campaigns, CO fluxes were estimated in two ways. First, soil chambers enclosing both litter and soil were used to measure net surface CO and CO₂ fluxes. Second, above and below canopy CO and CO₂ mixing ratio patterns were studied to estimate ecosystem CO fluxes from the net change of gases during stable atmospheric nocturnal conditions when mixing with air above the canopy is limited. Both methods demonstrated that tropical ~~rain forests~~rainforests are a net source of CO. Third, using a simple laboratory experiment, we show that soils are the main source

driving these emissions and that abiotic thermal degradation is likely its main driver. Finally, by focusing on different seasons and topographic locations, we attempt to identify the role of additional CO sources in the ecosystem. Based on our observations, we formulate a first observation-based estimate for global tropical ~~rain-forest~~rainforest soil CO emissions.

90

2 Material and methods

2.1 ~~Field-site~~Fieldsite and K34 tower micro-meteorological measurements

This research was performed in a mature rain forest, located ~50 km northwest of Manaus (Brazil) at the Reserva Biológica do Cuieiras (2 °36" 32.67 S, 60 °12"33.48 W), managed by the Instituto Nacional de Pesquisas da Amazônia (INPA), also
95 known as ZF2. The elevation at the site ranges from 40-110 m above sea level and is characterized by a dissected topography with plateaus, steep slopes and valleys. The vegetation on plateaus is terra firma (upland) forest with tree heights of 35-40 m, and with clay rich soils classified as Oxisols and Ultisols. Valleys are periodically inundated, with ~~three-tree~~tree heights of 25-30 m, and with sandy soils, classified as Spodosols (Luizão et al., 2004; Zanchi et al., 2014). The ~~field-site~~fieldsite has a distinct seasonality, with a dry season (months with precipitation <100 mm) lasting ~3 months between June-October, and a
100 wet season from December to May. Annual average precipitation is 2400 mm, and average annual air temperature is 26-28 °C. More information about the ~~field-site~~fieldsite can be found in Araújo et al. (2002); Chambers et al. (2004); Luizão et al. (2004); Quesada et al. (2010); Zanchi et al. (2014).

The K34 tower is a micro-meteorological tower located at ~~field-site~~fieldsite ZF2, run by the project LBA (~~Large Biosphere~~
105 ~~Atmosphere Experiment~~Large-Scale Biosphere-Atmosphere Experiment in Amazonia) since 1999, and is one of the longest running flux towers in a tropical ~~rain-forest~~rainforest. The tower is equipped with micro-meteorological as well as environmental measurements. Unfortunately, due to pandemic challenges, no measurements are available for the campaign periods, but data from earlier years was available to support our analyses.

110 2.2 Available instruments: FTIR-analyzer & ICOS-Analyzer

At the foot of the K34 tower, a Fourier Transform Infrared Spectrometer (ACOEM Spectronus, Trace Greenhouse Gas and Isotope Analyzer, from here on called FTIR-analyzer, ~~Griffith et al. (2012)~~(Griffith et al., 2012)) was installed in an air-conditioned cabin. The FTIR-analyzer simultaneously measures mixing ratios of CO₂, CH₄, N₂O and CO, as well as the $\delta^{13}\text{C}$ of CO₂; all reported mixing ratios in this study are per volume (ppmv or ppbv). The instrument can measure in either
115 static or flow modes. All incoming air samples are internally dried by a Nafion dryer and by a column of magnesium perchlorate, so that H₂O mixing ratios are usually <20 ppm. Measurements were corrected for pressure and temperature variations as well as for inter-species cross-sensitivities, which are related to the overlapping spectral absorption regions of different

trace species (Hammer et al., 2013). The precision (σ) of the FTIR-analyzers CO and CO₂ measurements for 2 min-spectral measurements is 0.45 nmol mol⁻¹ and 0.05 μmol mol⁻¹ respectively (Asperen et al., 2015a; Griffith et al., 2012). For the different methodologies based on concentration differences (explained below), a minimum concentration difference of 2σ was set as a detection limit.

The second available analyzer was an Off-Axis Integrated Cavity Output Spectroscopy gas analyzer (OA-ICOS), namely the Los Gatos Ultraportable Carbon Analyzer, from here on called ICOS-analyzer. The instrument is field portable (weight of 17 kg) with a potential to run on battery power, so that it could be used to measure fluxes at different field locations around the K34 tower. The instrument measures CO₂, CH₄, CO and H₂O at a flow of 0.3 LPM. For this study, the ICOS-analyzer was only used to measure mixing ratios and fluxes of CO₂: since the CO concentrations in a pristine tropical forest are generally low, the mixing ratios fell outside the reliable measurement range of the ICOS-analyzer. For this reason, all reported CO mixing ratios and fluxes are based on measurements from the FTIR-analyzer.

130

2.3 Soil flux chamber measurements

Two intensive campaigns were held in 2020/2021, encompassing 9-days during dry season (DS campaign, 28 September-7 October 2020), and 7-days during wet season (WS campaign, 11-18 May 2021). During both campaigns, a series of soil flux chamber measurements were performed on the plateau and in the valley. A soil chamber was made from a 200 L large bucket (non-transparent), and fitting soil collars were made from stainless steel (15 cm height, 56.5 cm diameter). A strip of closed-pore foam was glued to the inner edge of the chamber, so that no air could pass between the chamber and the collar during measurement. Two holes were made on each side of the chamber at around 50 cm height where a quick connect ¼ inch fitting was installed, serving as the inlet and outlet of the chamber. On the inside of the chamber, a four-inlet vertical sampling tube was placed so that the air sampled (flow rate of 0.3 LPM) was a mix from different heights in the head space (~10 cm, ~25 cm, ~35 cm, and ~50 cm) (Clough et al., 2020). The setup (chamber and tubing) was tested for internal gas emissions under field conditions (high temperature and humidity). For CO, an internal emission of <0.014 nmol s⁻¹ was found; the reported CO fluxes are not corrected for this small possible internal emission.

Five soil collars were installed on the plateau (~50 m from the tower), and another five soil collars were installed in the valley (~50 m from the location of the nighttime valley measurements and valley stream (see section 2.5)), approximately one month before the first (DS) measurement campaign. Soil collars were installed until a depth of 5 cm, and were installed at >1 m from larger trees and bushes, containing only soil and litter. The valley soil collars were just far enough from the valley stream to not be inundated after some days of heavy rain. The litter layer was not removed from the soil in the collars so that the soil surface was representative of the forest floor. During each campaign, each collar was measured 3 times. Each collar was measured for ~35 minutes, during which the air was circulated through the chambers by the internal pump of the ICOS-analyzer, which measured CO₂ simultaneously. Right after chamber closure, a bag sample was sampled from the chamber

150

inlet ~~by use of,~~ using an external pump (KNF, NMP 830 KNDC B). After that, a subsequent bag was sampled every 10 minutes (4 bags in total). Air was stored in 5L inert foil sampling bags (Sigma-Aldrich), which were brought to the FTIR-analyzer and analyzed on the same day. The soil CO flux (F_{CO}) was calculated as follows:

$$155 \quad F_{CO} = \frac{\Delta[CO]}{\Delta t} * \frac{V}{A} \quad (1)$$

wherein $\frac{\Delta[CO]}{\Delta t}$ was calculated with linear regression over the CO mixing ratios of the consecutive 4 bags, and $\Delta[CO]$ converted from mixing ratios (nmol mol^{-1}) to concentrations (nmol m^{-3}) by the ideal gas law (assuming a T_{air} of 25 °C), V is
160 0.20 m^3 , and A is 0.25 m^2 . Requiring a minimum concentration difference of 0.9 nmol mol^{-1} (2σ , FTIR-analyzers precision σ for CO is 0.45 nmol mol^{-1}) between the first and the last sampled bag, the minimal detectable flux of this system is 0.01 $\text{nmol CO m}^{-2} \text{s}^{-1}$. After each measurement, soil temperature T (measured with a manual sensor, type TP-101) and soil volumetric water content (VWC) (AT SMT150) were measured around the collar 5 times, of which the median was taken.

165 2.4 Plateau tower CO mixing ratios and flux estimates

To determine atmospheric CO mixing ratios at different heights in ~~the canopy air and above the canopy~~, inlet lines of Synflex tubing ($\frac{1}{4}$ inch) were installed at the tower at 36 m, 15 m and 5 m height (canopy height is ~ 35 m). Each inlet was equipped with a rain protection cap and a particle filter. Each line extended until to the cabin, where it passed an air cooler (4°C) with several water traps, which prevented condensation droplets from entering the sampling manifold and instrument. After the
170 water traps, the lines led to a sampling manifold, from which one single line entered the FTIR-analyzer. Calibration gases (gas 1 with 381.8 $\mu\text{mol CO}_2 \text{ mol}^{-1}$, and 431.0 nmol CO mol^{-1} , and gas 2 with 501.6 $\mu\text{mol CO}_2 \text{ mol}^{-1}$ and 256.7 nmol CO mol^{-1}) were available, and measured at least 3 times during each campaign. During the campaign-periods, the FTIR-analyzer alternated measuring air from the 3 heights in an half hourly cycle (10 min per height), using a sampling flow of 1.2 LPM. Since the FTIR-analyzer has a large measurement cell (3.5 L) and a corresponding long turn-over time-e-folding time, only the
175 last 2 minutes of each 10 minute measurement window were used.

During the first 5 days of the dry season campaign (28 September to 7 October), a leak was ~~found in the tower present in the~~ 36 m inlet line, ~~so that the data from this inlet between 28 September and 4 October could not be used. For this reason, for some of.~~ To be able to obtain sufficient data for the subsequent analyses, ~~measurements from an extended period were used, namely 5-12 tower measurements continued until the 12th~~ of October. The daytime tower vertical profile measurements were interrupted during campaign days because the instrument was used to measure the different sampling bags, sampled in the ecosystem. To estimate ecosystem CO fluxes from atmospheric CO mixing ratios, only nighttime CO measurements were

used.

185 The measured CO mixing ratios were interpreted using 2 different approaches. Nighttime vertical CO mixing ratio profiles (dCO/dz) were ~~compared between~~ studied over different time windows over each night. To enable a more straightforward comparison, the mixing ratios at 15 m and 36 m were expressed relative to the 5 m height (~~dCO-15m and dCO-36m~~ 'dCO-15m' and 'dCO-36m'): a negative dCO indicates that the CO mixing ratios at 15 m or 36 m are lower than at 5 m height. Vertical profiles per ~1-h time window were calculated, which consisted of 3 measurements per height. Per night, the following time
190 windows were used: 18h-19h, 20h-21h, 22h-23h, 0h-1h, 2h-3h, 4h-5h. The given date of a night indicates the date of the start of the evening, for example '28 September' indicates the night from 28-29 September. Please note that the 'd' is used to indicate a spatial difference (vertical profile, ~~dCO-36m~~ dCO-36m), while the Δ symbol is used to indicate a change over time (introduced below).

195 Next to the analyses of the vertical CO profile, a canopy layer budget method was used, as described by Trumbore et al. (1990) and applied by earlier studies in tropical forests for CH₄ (Carmo et al., 2006):

$$\frac{\Delta CO}{\Delta t} = PCO - k(C(CO) - C(CO_{atm})) \quad (2)$$

200 wherein PCO stands for the production of CO in the canopy layer, C(CO) and C(CO_{atm}) stand for respectively the mixing ratio in the canopy layer and the mixing ratio of the overlying atmosphere, and k represents an exchange coefficient. This equation can also be defined for CO₂, which can then be merged into:

$$\frac{\Delta CO}{\Delta CO_2} = \frac{PCO - k(C(CO) - C(CO_{atm}))}{PCO_2 - k(C(CO_2) - C(CO_2_{atm}))} \quad (3)$$

205

During stable nighttime conditions, when the exchange between the canopy layer and the overlaying atmosphere is low, a similarity between CO₂ and CO mixing ratio patterns and production rates can be assumed, so that Eq. 3 can be simplified to (Carmo et al., 2006):

$$PCO = \frac{\Delta CO}{\Delta CO_2} * PCO_2 \quad (4)$$

210

in which PCO_2 can be inferred from Eddy Covariance flux data. To filter for nighttime stable conditions, the period 18h-4h was chosen, based on an earlier study at this [field-site-fieldsite](#) showing generally stable conditions for these hours (Araújo et al., 2002). For each night of the campaign week, the $\frac{\Delta\text{CO}}{\Delta\text{CO}_2}$ was calculated for different time windows, [namely](#) 18h-20h, 20h-22h, 22h-0h, 0h-2h, 2h-4h. The two heights below the canopy, namely 5 and 15 meters, were both used independently, and values shown are filtered for $R^2 > 0.9$. Due to unavailable micro-meteorological CO_2 flux measurements, it was decided to choose a fixed value for PCO_2 of $7.8 \mu\text{mol m}^{-2} \text{s}^{-1}$, based on a previous study at the same [field-site-fieldsite](#) (Chambers et al., 2004).

220 2.5 Valley CO mixing ratios and flux estimates

To complement the mixing ratio measurements on the plateau, additional measurements were performed in a valley close to the K34 tower. Equipment was placed in a box on a wooden boardwalk, constructed above a stream and a muddy, and sometimes inundated, area. Two 10 m $\frac{1}{4}$ inch teflon lines were extended from the Zarges box and installed ~ 10 m from the boardwalk (~ 2 m from the valley stream), hanging 1 m above the soil surface. The Zarges box contained the ICOS-analyzer, which was continuously sampling air from one teflon line (0.3 LPM, measurement every 10 sec). In addition, a sampling device with a KNF pump (NMP 830 KNDC B, ~ 1 LPM) was placed in the same box, continuously flushing the 2^{nd} sampling line. At fixed times (+0h, +3h, +6h, +9h after start of the measurements), air (~ 8 liters) was sampled into bags (4 bags, 10L inert foil, Sigma-Aldrich). These bags were collected during the following morning, and measured by the FTIR-analyzer on the same day. The starting time of the measurements was usually just around nightfall, between 17:30 and 18:30, and the external battery feeding the ICOS-analyzer usually held approximately 10-12 hours.

The continuous ICOS-analyzer measurements were used to study the general behaviour of the CO_2 mixing ratio trends during the night, while the additional bag measurements were used to determine the CO nighttime increase and the $\frac{\Delta\text{CO}}{\Delta\text{CO}_2}$ ratio (Eq. 4). For PCO_2 , the value of $7.8 \mu\text{mol m}^{-2} \text{s}^{-1}$ was used (Chambers et al., 2004).

235

2.6 Laboratory thermal degradation measurements

To study thermal degradation of ecosystem material, a simple laboratory experiment was set up. Soil (upper 3 cm, not sieved) and senescent leaf material was sampled from a $2 \times 2 \text{ m}^2$ area on the plateau and in the valley. The material was dried at 35°C for 72 h, to assure microbial activity to be negligible (Lee et al., 2012). From each material, 3 sub samples were taken of ~ 2 grams (leaves) and ~ 30 grams (soil). For the experiment, a glass flask (inner diameter = 6.7 cm, height = 15 cm) was placed in a closed loop with the FTIR-analyzer. For this experiment, only glass and stainless steel material was used. Blank measurements showed the set up was not emitting CO. The sample material was distributed equally in the flask. The samples were heated in temperature steps of 5°C (20 – 65°C) by use of a controlled temperature water bath. Temperature time steps were 20

min. During the experiments, air was circulated between the glass flask and the FTIR-analyzer and measured once per minute.

245

The production rate of CO was derived from the measured mixing ratio change over time, and is expressed as $\text{nmol CO gr}_{leaves}^{-1} \text{ min}^{-1}$ (or $\text{nmol CO gr}_{soil}^{-1} \text{ min}^{-1}$). To be able to express senescent leaf CO production rates on ecosystem scale, a literature senescent leaf density value of 117 and 67 g m^{-2} (1.17 and 0.67 t DW ha^{-1}) was taken for respectively plateau and valley, as measured by Luizão et al. (2004) at the same ~~field-site~~fieldsite. To be able to express soil material CO production rates
250 from a 10-cm soil layer on ecosystem scale, a plateau soil bulk density of 1.05 g cm^{-3} was assumed, as measured on the same ~~field-site~~fieldsite (Marques et al., 2013). All experiments were conducted under dark conditions, to exclude photo-degradation fluxes.

3 Results

3.1 Soil CO and CO₂ fluxes

255 On the plateau, CO fluxes determined from the accumulation in the soil chambers were significantly larger in the dry season than in the wet season (Fig. 1, Table 1), and one collar in the wet season even showed uptake during all three measurements. Soil temperature and moisture variation within campaigns was small, so that correlations to CO and CO₂ fluxes were generally not pronounced. When grouping all plateau CO flux measurements (dry and wet season), a correlation with soil temperature ($R^2=0.53$) and an inverse correlation with soil moisture ($R^2=0.57$) was ~~found~~observed (Fig. 2). ~~Plateau~~In addition, plateau
260 soil moisture (VWC) and soil temperature (T) values showed a clear correlation, with higher T accompanied by lower VWC ($R^2=0.70$). Valley CO fluxes were generally higher than plateau CO fluxes. As on the plateau, valley wet season CO fluxes were smaller than dry season valley CO fluxes (Table 1), but only a weak correlation with soil T ($R^2=0.24$) and soil VWC ($R^2=0.13$) was found. Also in the valley, warmer temperatures were accompanied with lower VWC, although the correlation was weak ($R^2=0.15$). For CO₂, dry season fluxes were higher in the valley than on the plateau, while in the wet season
265 the pattern was inverted. ~~Nevertheless, differences~~Plateau CO₂ fluxes showed a small positive relationship with soil VWC, which is opposite of what was observed for CO. Valley CO₂ fluxes showed a positive relationship with soil T and a negative relationship with soil VWC. Differences in CO₂ fluxes between seasons and topographic locations were not significant and the observed relationships between CO₂ fluxes and soil T and soil VWC were weak (Table 1), Fig. 2), and will not be further discussed in this manuscript.

270 3.2 Atmospheric CO mixing ratios and ecosystem CO flux estimates

Dry season CO campaign mixing ratios varied between 127 and 292 ppb (Fig. 3). Mixing ratios between the different heights generally followed a common pattern, indicating that air masses with elevated CO passing the tower are also reaching lower forest levels. Wet season campaign CO mixing ratios ranged between 94 ppb and 250 ppb, and generally showed less variation, fewer peaks, and lower mixing ratios. It is expected that (part of) the elevated mixing ratios and passing peaks in the dry season

275 can be explained by the presence of biomass burning plumes, which can be transported over long distances (Andreae et al., 2012). The CO mixing ratio patterns, and the possible trajectories and dispersion of these biomass burning plumes, are subject of ongoing research and will not be further discussed in this study.

Vertical profiles per 1-h time windows are shown in Fig. 4. In the dry season, 4 out of 7 nights showed constant decreases
280 in CO mixing ratio from 5 to 36m-36m, i.e. consistently negative d-36m-dCO-36m during the whole night. Average nighttime d-36m-dCO-36m values for these nights were -10.5 ppb, -8.1 ppb, -10.9 ppb and -7.0 ppb (6, 7, 9 and 10 October, ~~last column of~~ Fig. 4). In the wet season, vertical nighttime profiles generally showed smaller variation in CO mixing ratios, however still 5 out of 7 nights showed consistently decreasing CO mixing ratios with height over the whole night. In addition, only three 1h-time windows showed a dCO-36m value <0.9 ppb (2σ), which is considered the detection limit of the method. Average
285 dCO-36 values for these nights were generally smaller than in the dry season: -2.5 ppb, -5.6 ppb, -4.6 ppb and -2.3 ppb (~~resp. respectively~~ 13, 15, 16 and 17 May), with the exception of 14 May (-12.9 ppb) (Table 1, Fig. 4). Since no micro-meteorological measurements are available for the campaign periods, we cannot hypothesize why the night of 14 May was divergent.

The canopy layer budget method was applied to the plateau ~~below-canopy inlet heights~~ inlet heights of 5 and 15 m, and the
290 valley inlet height of 1 m. All ΔCO₂ and ΔCO values of the 2h-time windows (plateau) and the 3h-time windows (valley) were higher than the set detection limit of 2σ (ΔCO₂>0.1 ppm, ΔCO>0.9 ppb). For the plateau, calculated $\frac{\Delta CO}{\Delta CO_2}$ ratios with an $R^2 > 0.9$ were selected, which was 29% and 45% of the cases in the dry season, and 41% and 40% of the cases in the wet season for 5 m and 15 m respectively. Dry season $\frac{\Delta CO}{\Delta CO_2}$ ratios were slightly higher than wet season ratios, but differences were not significant (Table 1). Applying Eq. 4 to the 5 m mean nighttime ratios (DS=0.27 and WS= 0.24) gives a plateau ecosystem
295 net production estimate of 2.1 and 1.9 nmol CO m⁻² s⁻¹ for the dry ~~season~~ and wet season respectively. ~~The canopy layer budget was also applied to nighttime valley measurements (inlet at ~1 m height). In the dry season~~ For the valley measurements, correlation coefficients (R^2) between ΔCO and ΔCO₂ ~~reached 0.75~~ were >0.75 in 8 out of 9 nights (4 nights with $R^2 > 0.9$), and in the wet season, 6 out of 7 nights reached $R^2 > 0.75$ (5 nights with $R^2 > 0.9$). The wet season ratios were significantly higher than the dry season ratios, and applying Eq. 4 to the mean ratios leads to estimates of a net valley CO production of 1.1
300 and 2.3 nmol m⁻² s⁻¹ for respectively the dry and the wet season (Table 1).

3.3 Laboratory results

Senescent leaves exposed to different temperatures emitted significant amounts of CO at rates increasing exponentially with higher temperatures. At 25 °C, average emission rates of 0.006 and 0.002 nmol CO g_{leaves}⁻¹ min⁻¹ were measured for plateau and valley samples respectively. The estimated ecosystem CO production rates, based on these average emission rates and literature senescent leaf density (Luizão et al., 2004), are 0.012 and 0.002 nmol CO m⁻² s⁻¹ at 25 °C for respectively the plateau
305 and valley ecosystem (Fig. 5). The plateau soil material showed clear CO production increasing with higher temperatures. The valley soil material also showed CO production, but due to instrument problems, a complete and consistent data set could not be collected. For plateau soil material, an emission of 0.0005 nmol CO g_{soil}⁻¹ min⁻¹ at 25 °C was measured. The estimated

ecosystem emission coming from a plateau soil layer of 10 cm (Marques et al., 2013) at 25 °C was estimated to be ~0.9 nmol
310 CO m⁻² s⁻¹ (Fig. 5).

4 Discussion

Plateau and valley soil chambers, measuring the emission of soil and litter together, generally showed net emission of CO, except for one plateau collar in the wet season. On the plateau, [grouped dry and wet season](#) CO fluxes showed a relation with soil temperature ($R^2=0.53$) as well as with soil VWC ($R^2=0.57$). ~~In the valley, there was less~~ [Correlations for individual campaigns did not fit to this correlation curve \(Fig. 2\): while this can be partly attributed to the limited variation in soil temperature and soil VWC, so that possible dependencies were less pronounced. Correlations between soil temperature moisture values, which diminishes the clarity of correlations, it also suggests that the drivers influencing plateau CO production may differ between seasons. Moreover, the strong correlation between soil T and soil VWC \(\$R^2=0.70\$ \) on the plateau did not permit determination of which factor is driving the soil CO flux variation. In the valley, the variation in soil T and soil VWC was even more limited,](#)
320 [resulting in less pronounced dependencies and overall low correlations \(all \$R^2 < 0.32\$, Fig. 2\).](#)

Abiotic CO production as well as microbial CO uptake should correlate positively with increasing soil temperatures ([Cowan et al., 2018; King, 2000; Lee et al., 2012; Derendorp et al., 2011; Asperen et al., 2015b; Moxley and Smith, 1998](#))([Asperen et al., 2015b](#)). While for microbial CO uptake the relationship is expected to have an optimum temperature (King, 2000), the abiotic thermal
325 degradation fluxes are exponential ([Derendorp et al., 2011; Asperen et al., 2015b; Lee et al., 2012; Conrad and Seiler, 1985](#))-
([Asperen et al., 2015b; Conrad and Seiler, 1985; Derendorp et al., 2011; Lee et al., 2012](#)) so that CO production is expected to become dominant at higher temperatures ([Moxley and Smith, 1998; King, 2000; Asperen et al., 2015b; Cowan et al., 2018](#))
([Asperen et al., 2015b; Cowan et al., 2018; King, 2000; Moxley and Smith, 1998](#)). The role of VWC is more complicated. On the one hand, lower soil VWC leads to higher soil diffusivity, enhancing the CO uptake, thereby shifting the balance to soil
330 CO uptake. For example, at the same ~~field site~~[fieldsite](#), high local CO uptake was observed from termite mounds, which consist of dry porous material ([Asperen et al., 2021; Martius et al., 1993](#))([Asperen et al., 2021; Martius et al., 1993](#)). On the other hand, the availability of soil moisture has a direct effect on microbial CO uptake. Several studies found a parabolic response, with soil CO uptake having an optimum at VWC ~20-30% ([Moxley and Smith, 1998; King, 2000; King and Hungria, 2002](#))
([King, 2000; King and Hungria, 2002; Moxley and Smith, 1998](#)). Based on the supposed decrease in CO uptake when VWC
335 >30%, one would expect a shift towards more positive CO fluxes from the dry season to the wet season, which is opposite of what is observed in our measurements (Fig. 1). Following this line of reasoning, we expect that the observed negative correlation between VWC and soil CO fluxes is an indirect one, driven by the correlation of soil T and soil VWC.

The laboratory experiment, isolating the effect of temperature on CO production of senescent leaves and soil material, indicated a clear exponential increase in CO emissions with temperature, as also reported by earlier studies ([Derendorp et al., 2011; Asperen et al., 2015b; Derendorp et al., 2011; Lee et al., 2012](#)). By combining literature values with our laboratory results, a

simple approximate calculation was done to estimate the CO emission of senescent leaves and soil material at 25°C at the ecosystem scale. Senescent leaves in the amount expected in the surface litter layer (Luizão et al., 2004) were estimated to emit respectively 0.012 and 0.002 nmol CO m⁻² s⁻¹ on the plateau and in the valley, and a 10-cm plateau soil layer (Marques et al., 2013) was estimated to emit 0.93 nmol CO m⁻² s⁻¹. This simple ~~up-sealing~~ upscaling ignores the collocated simultaneous soil CO uptake and, more importantly, this estimate ignores the CO production of the entire soil column below. Soil CO uptake has been shown to be dominant in ecosystems under specific conditions, such as lower temperatures and porous conditions (King, 2000; Kisselle et al., 2002), and therefore in-depth research for this specific ecosystem would be needed to improve this upscaling. Nevertheless, this simple ‘back-of-the-envelope’ calculation already shows the potential of mineral soil to be a strong emitter of CO, and suggests that the observed chamber fluxes, which were measured over soil and litter together, mainly reflect emissions from the soil.

The laboratory CO emissions, the chamber CO fluxes, and the nighttime ecosystem CO increase all demonstrate net production of CO by this ecosystem. All observations were performed in absence of solar radiation, so that a photochemically-induced CO production pathway, such as photodegradation of organic material or the oxidation of VOCs and hydrocarbons (~~Lee et al., 2012; Schade et al., 1999; Tarr et al., 1995; Szopa et al., 2021~~) (Lee et al., 2012; Schade et al., 1999; Szopa et al., 2021; Tarr et al., 2012), is unlikely to have contributed to our fluxes. In addition, the thick canopy of these forests prevent much sunlight from penetrating into the lower canopy or reaching the forest floor. Besides thermal degradation, ozonolysis of unsaturated hydrocarbons can produce CO in absence of radiation (Röckmann et al., 1998). However, CO produced via ozonolysis would be associated with a strong enrichment in δ¹⁸O, which was not observed in additional experiments (see Appendix A). Therefore, we can exclude that ozonolysis plays a major role in our ecosystem. Following this line of reasoning, and supported by the clear observed relationships between temperature and CO fluxes (Figs. 2 and 5), we conclude that thermal degradation is likely the main driver of the observed nighttime CO production increase and measured flux chamber CO fluxes in this central ~~Amazon tropical rain forest.~~ Amazonian tropical rainforest. The possible presence of additional photodegradation-induced CO production during daytime would lead to even higher net total CO fluxes, a process not yet studied for tropical rainforests.

Plateau and valley CO mixing ratios were used to estimate ecosystem CO fluxes, which was done by studying the vertical CO gradient (only on the plateau), as well as by applying a canopy budget method (plateau and valley). Both approaches were only applied to nighttime measurements, when atmospheric conditions are generally more stable and with locally produced gases ‘trapped’ below the canopy, so that mixing ratio changes are more pronounced (Araújo et al., 2002). ~~The~~ Generally, the nighttime vertical CO gradients ~~generally~~ were negative i.e. had higher mixing ratios closer to the forest floor in comparison to above-canopy mixing ratios (Fig. 4). The vertical gradient can be used to estimate an ecosystem flux, by assuming a fixed canopy flushing rate of 90% over a vertical column of 30 m, as measured at a similar ~~field site~~ fieldsite close-by (Simon et al., 2005). Querino et al. (2011) applied the same method and assumptions to vertical CH₄ gradients, measured at the same tower, where it was shown to give comparable flux estimates to on-site Eddy Covariance CH₄ measurements. The vertical CO gradients suggest an ecosystem flux up to ~2 nmol CO m⁻² s⁻¹, with vertical profiles generally indicating larger emissions in the

dry season (Table 1, last [column](#)[rows](#)). On the plateau, the canopy budget method showed no significant differences in $\frac{\Delta CO}{\Delta CO_2}$ -ratios between the ~~wet and the dry~~ [dry and the wet](#) season and, based on the [5m-5 m](#) inlet, plateau CO fluxes were estimated to range between 1.25 and 4.13 nmol CO m⁻² s⁻¹ in the dry season, and between 0.94 and 3.28 nmol CO m⁻² s⁻¹ in the wet
380 season. The valley nighttime $\frac{\Delta CO}{\Delta CO_2}$ -ratios were generally lower in the dry season, and valley CO fluxes were estimated to range between 0.3 and 1.9 nmol CO m⁻² s⁻¹ and between 1.8 and 3.1 nmol CO m⁻² s⁻¹ for respectively the dry and the wet season.

The vertical gradient approach as well as the canopy budget method, and their subsequent ecosystem estimates, are possibly affected by the varying background CO mixing ratios (Fig. 3). We attempted to adjust for these background variations by
385 applying strict filters. As described above, for the ecosystem flux estimate based on dCO/dz, only nights when the ~~d-36m was~~ [dCO-36m was consistently](#) negative over the entire night were selected for [up-sealing](#)[upscaling](#). For the canopy budget method, which is based on the temporal change (ΔCO) below the canopy, only consistent ΔCO changes with a strong correlation to ΔCO_2 were selected ($R^2 > 0.9$), so that CO variations caused by a change in background mixing ratios are removed. The fact that the filtered ratios are relatively constant between heights and nights gives us confidence that this approach indeed selects the
390 CO mixing ratio trends which are caused by the local ecosystem. In addition, the valley CO and CO₂ mixing ratios are possibly affected by nighttime drainage from the plateau ([Tóta et al., 2008](#); [Araújo et al., 2008](#)) ([Araújo et al., 2008](#); [Tóta et al., 2008](#)): Araújo et al. (2008) showed valley nighttime CO₂ pooling at the same [field-site](#)[fieldsite](#), with plateau CO₂ laterally transported below the canopy. If this pooling is happening for CO₂, it is not unlikely that other trace gases, such as CO, are also transported. We therefore expect that the valley $\frac{\Delta CO}{\Delta CO_2}$ -ratios are not affected or, in case only CO₂ is pooling, are underestimated,
395 which would mean that valley CO emissions would be even higher. Thus, possible pooling would not affect our prediction that valleys are a net CO emitter.

The used PCO₂ of 7.8 μmol m⁻² s⁻¹, as reported by Chambers et al. (2004), is a general ecosystem value and not specified for season or topographic location. Different studies at this [field-site](#)[fieldsite](#) have demonstrated differences in CO₂ fluxes
400 between the plateau and the valley. For example, Araújo (2009) performed Eddy Covariance measurements on the plateau and in the valley, and found that the valley PCO₂ was 2/3 of the plateau PCO₂ (7.2 vs 4.8 μmol m⁻² s⁻¹). Comparing plateau and valley, Chambers et al. (2004) found that soil respiration at this [field-site](#)[fieldsite](#) was even twice as high on the plateau, but pointed out that the valley soil respiration fluxes are likely underestimated to an unknown degree. Zanchi et al. (2014) found an opposite pattern, with valley soil respiration being ~1.5 times higher than plateau soil respiration, which is similar as observed
405 in our study in the dry season (Fig. 1). Since the degree and direction of soil and ecosystem CO₂ flux variation between seasons and topographies is unclear, a differentiation in PCO₂ could introduce additional uncertainties. For this reason, for this study, it was decided to use a fixed PCO₂ for all topographies and seasons.

An overview of the *direct* soil CO flux measurements and *indirect* ecosystem CO flux estimates is given in Table 1. A direct
410 comparison of these values should be done with care. First of all, the soil flux measurements are performed during (warmer) daytime hours, while the ecosystem estimates are determined for cooler nighttime conditions, although temperature variations

below the canopy in this ecosystem are generally small ($<7^{\circ}\text{C}$, Araújo et al. (2002)). Secondly, the flux chamber is measuring soil and litter only, while the ecosystem estimates include all possible sources and sinks below the canopy. Most importantly, soil flux values are measured directly, while the ecosystem fluxes are indirect estimates. Nonetheless, for the dry season, the plateau soil CO fluxes and the ecosystem CO flux estimates agree on the sign as well as on the magnitude of the CO fluxes. Moreover, the $\frac{\Delta\text{CO}}{\Delta\text{CO}_2}$ ratio of the plateau nighttime increase shows similar ratios as the soil fluxes (Fig. 1, Table 1). We therefore expect that, in the dry season, the plateau nighttime CO increase is mostly driven by soil emission. For the wet season, the plateau soil CO fluxes were a lot smaller than the estimated ecosystem CO fluxes, and the flux chamber even showed uptake at one location. Because of the decrease in soil CO fluxes, the soil flux $\frac{\Delta\text{CO}}{\Delta\text{CO}_2}$ ratios strongly dropped, which was only weakly observed in the ecosystem $\frac{\Delta\text{CO}}{\Delta\text{CO}_2}$ ratios (Fig. 1). The difference in flux magnitudes and ratios indicates that, in the wet season, the type of soil surface, as measured in our soil chamber, is probably not the main driver of the plateau nighttime CO increase. Possibly, the plateau soil CO fluxes have a large spatial variability, with our soil collars representing cooler/wetter spots than the surrounding area. ~~Additional (nighttime) soil flux chamber measurements~~ In addition, the seasonal shift in correlations between soil T, soil VWC and plateau soil CO fluxes (Fig. 2, left column) suggests that different dominant drivers may come into play during the wet season. A more elaborate flux chamber campaign, with possible nighttime measurements, would be crucial to verify ~~this hypothesis~~ the different hypotheses above.

All three methods on the plateau indicate higher CO emissions in the dry season although the difference is only significant for the direct soil flux measurements. Based on the earlier described relationship between soil temperature and soil CO fluxes (Fig. 2), and the clear increase of CO production with temperature (Fig. 5), we expect that the generally higher soil temperatures in the dry season cause the difference in CO fluxes between the seasons. ~~In~~ While no meteorological data from the K34-tower are available for both campaign periods, measurements from the local airport (~50 km distance) show that the dry season campaign period had clearly higher temperatures and lower relative humidities in comparison to the wet season campaign period (INMET, 2024). In addition, in Appendix B, typical plateau dry and wet season soil temperatures from this ~~field site~~ fieldsite are shown, which indicate that the general diurnal temperature pattern barely drops below the estimated 'soil-CO-uptake-threshold-temperature' of 25.2°C (Fig. 2), even at night or in the wet season (Fig. A1).

Just as on the plateau, the valley soil fluxes were significantly higher in the dry season. In the dry season, valley soil chamber CO fluxes and ecosystem CO flux estimates agree quite well (Table 1), indicating that the nighttime valley CO increase is driven by sources such as our measured valley soil surface. Nevertheless, in the wet season, a clear discrepancy between the soil chamber flux magnitude and the valley ecosystem estimate was observed, indicating that the nighttime CO increase is driven by additional sources, that are not captured in the flux chambers. Since the wet season is characterized by frequent and high amounts of rain fall, the valley stream is frequently flooding its adjacent areas, which was also observed for the area below the valley inlet. Streams and rivers are known to be sources of CO (~~Zuo and Jones, 1997; Campen et al., 2023; Valentine and Zepp, 1993~~ , either produced by photo or by thermal degradation (Campen et al., 2023; Müller, 2015; Valentine and Zepp, 1993; Zhang et al., 2008; Z , so that it is likely that our wet season nighttime measurements were dominated by the nearby valley stream and its inundated

areas. ~~Based on our measurements alone, we do not speculate whether the stream CO fluxes are caused by nighttime thermal degradation CO fluxes, or represent a delayed outgassing of photogradation-produced CO.~~ On the whole, based on our observations, we expect that the valley is a net source of CO, with generally higher *soil* emissions in the dry season, but with likely higher overall *ecosystem* emissions in the wet season, due to the contribution of the valley stream and its inundated areas.

Our measured soil CO fluxes are generally higher compared to the limited previous soil CO flux studies performed in (sub-) tropical ecosystems. Kisselle et al. (2002) observed ~~uptake-negative fluxes (uptake)~~ of -0.31 to -0.07 $\text{nmol m}^{-2} \text{s}^{-1}$ in the dry and wet season, ~~respectively~~ (Brazilian savanna, Cerrado biome, opaque chambers, not burned area). Sanhueza et al. (1994) found that Venezuelan grasslands were a net CO source of 0.6 $\text{nmol m}^{-2} \text{s}^{-1}$, which turned into a small CO sink when ploughed (~ -0.3 $\text{nmol m}^{-2} \text{s}^{-1}$). Venezuelan forest soils were found to be a net sink (~ -4 $\text{nmol m}^{-2} \text{s}^{-1}$), but a net source (~ 0.1 $\text{nmol m}^{-2} \text{s}^{-1}$) after deforestation and 'conversion' into a scrub grass savanna ~~Scharffe et al. (1990)~~ (Scharffe et al., 1990). To our knowledge, only one study before attempted to estimate tropical ~~rain-forest-rainforest~~ CO fluxes, ~~which was done by namely~~ Kirchhoff and Marinho (1990) in a ~~field-site-fieldsite~~ close-by ZF2 (Ducke Forest reserve forest, ~ 50 km). Kirchhoff and Marinho (1990) measured the vertical CO gradient below the canopy, and observed higher concentrations close to the soil surface in comparison to canopy height (dCO of -10 ppb), which is similar to gradients observed in this study. Based on this gradient, they estimated a forest CO flux of ~ 6 $\text{nmol m}^{-2} \text{s}^{-1}$, implying the forest to be a source of CO.

By providing the first *direct* tropical ~~rain-forest-rainforest~~ soil CO flux measurements, and by complementing these observations with nighttime ecosystem mixing ratio measurements, we can confirm the hypothesis of Kirchhoff and Marinho (1990), and can state that tropical ~~rain-forest-rainforest~~ ecosystems are likely a net source of CO. By a simple ~~up-sealing-upscaling~~ (averaging seasons and topographical locations of soil CO fluxes, Table 1), we derive an average tropical ~~rain-forest-rainforest~~ soil emission of ~ 1 $\text{nmol CO m}^{-2} \text{s}^{-1}$. Our nighttime measurements indicate that the general ecosystem CO emission ~~might be~~ is possibly higher than this value, since ~~swampy/inundated areas and~~ valley streams, inundated soils and swampy areas, which are abundant in these ecosystems, might represent local hot spots. Translating our soil (and ecosystem) CO flux estimate to a yearly value gives an estimate of 0.9 $\text{g CO m}^{-2} \text{yr}^{-1}$ and, by assuming a ~~global tropical (evergreen) forest-tropical rainforest~~ area of 17.8×10^6 km^2 (~~Liu et al., 2018~~), (global tropical (evergreen) forest', Liu et al. (2018), Table 4), a total global tropical ~~forest-rainforest~~ emission of ~ 16.0 Tg CO yr^{-1} is estimated.

By an ~~innovate-innovative~~ combination of methods and instruments, we were able to study the CO mixing ratios and fluxes over different temporal and spatial scales, even in a remote challenging ecosystem such as a tropical ~~rain-forest-rainforest~~. In the absence of a mobile CO analyzer, a more logistically challenging bag sampling design had to be employed, which is why the number of CO flux and mixing ratio measurements remains small, especially in the valley. By only focusing on two campaign weeks and two locations in a tropical ~~rain-forest-rainforest~~, we realize that our study presents only a snapshot of the complex CO dynamics of a tropical ~~rain-forest-rainforest~~. Nevertheless, our unique set of measurements shows that tropical forests are a net source of CO, likely dominated by soil CO emissions. In addition, in valley areas, river and water sources are expected to

contribute to the overall ecosystem CO emission. To further improve our understanding of the CO dynamics of a tropical ~~rain forest~~rainforest, more in-situ tropical forest soil and ecosystem CO flux measurements are needed, focusing on the possible complex dependencies between CO fluxes, (soil) temperature and soil moisture, and other environmental variables. Moreover, 485 the role of soil type (e.g. texture, organic matter layer, porosity) and the significance of streams and inundated areas should be investigated. With the recent availability of mobile CO analyzers, we anticipate more ~~in-depth~~in-depth studies, focusing on the different temporal and spatial scales of tropical ~~rain-forest~~rainforest CO fluxes.

5 Conclusions

490 By providing the first direct CO flux measurements of tropical ~~forest~~rainforest soils, we can show that, in this ecosystem, soil CO production generally dominates over soil CO uptake. Complementary measurements of nighttime CO mixing ratios also suggest an overall net ecosystem CO emission, and estimated ecosystem CO fluxes were of the same sign and of similar magnitude as the measured soil CO fluxes. Thus, we can state that tropical ~~rain-forest~~rainforest ecosystems are likely a net source of CO, and we expect that soil emissions are the main contributor to the ecosystem CO emissions.

495

We observed that higher net soil CO emissions were accompanied by higher soil temperatures, and the warmer dry season generally showed larger soil and ecosystem CO emissions. With an additional laboratory experiment, the effect of temperature on CO production of senescent leaves and soil material was studied. The results show the potential of the soil material to be a strong emitter of CO, and indicates that the observed chamber fluxes, which were measured over soil and litter together, are 500 mainly driven by the soil. By excluding a large contribution of the process ozonolysis or a radiation-induced CO production pathway, we expect that the observed CO fluxes are mainly produced by the process of thermal degradation.

By a simple ~~up-scaling~~upscaling, we provide a first observation-based tropical ~~rain-forest~~rainforest soil emission estimate of $\sim 1 \text{ nmol CO m}^{-2} \text{ s}^{-1}$ ($0.9 \text{ g CO m}^{-2} \text{ yr}^{-1}$), which leads to an estimated global tropical ~~rain-forest~~rainforest soil emission 505 of $\sim 16.0 \text{ Tg CO yr}^{-1}$. Total ecosystem CO emissions might still be higher, since valley streams and inundated areas might represent local hot spots. To further improve these tropical forest ecosystem CO emission estimates, and to understand the complex dynamics between soil uptake and emission and its dependencies on environmental variables, more in-situ tropical forest soil and ecosystem CO flux measurements are essential.

Appendix A: CO production via ozonolysis

510 Ozonolysis is the process by which ozone (O_3) can initiate oxidation of unsaturated hydrocarbons via addition to the double bond. In subsequent reaction steps CO can be produced (~~Criegee, 1975; Paulson and Seinfeld, 1992~~)(Criegee, 1975; Paulson and Seinfeld, . Ozonolysis can occur in absence of radiation, and is therefore a potential contributor to our observed ecosystem nighttime

CO increase. Ozone is known to be isotopically strongly enriched in ^{18}O and ^{17}O , with a typical $\delta^{18}\text{O}$ value of 80 - 100 ‰. Röckmann et al. (1998) demonstrated that CO produced via ozonolysis inherits the strong ^{18}O and ^{17}O enrichment of O_3 , because one of the O atoms is transferred from O_3 to CO. By determining the $\delta^{18}\text{O}$ of the CO increase at night, the contribution of ozonolysis can be assessed.

To investigate a potential contribution of ozonolysis to the nighttime CO production, additional measurements were performed in September 2022 in the same valley where the nighttime valley CO increase was observed. A teflon line (~5 m) was placed at ~2 m from the valley stream, at 1 m height. This line was sampled during four time windows: 17:00-17:30 (just before sunset), 21:00-21:30 and 21:30-22:00 (nighttime), and 7:30-8:00 (just after sunrise). Two pressurized air flasks (1L) were sampled per time window, using a manual flask sampler (Heimann et al., 2022). To determine the isotopic composition of CO, these 8 flasks were sent to the isotope laboratory at the Institute of Marine and Atmospheric Research Utrecht (IMAU) of Utrecht University. Unfortunately, several flasks broke during transport, and only 3 flasks (1 night flask, and 2 morning flasks) could be analyzed for CO and its isotopic composition (Table A1). A Keeling plot of these values results in an intercept of -20.0 ‰. Even though only three samples could be analyzed for isotopic composition, and the increase in CO is only 20-30 ppb, the Keeling plot analyses show that the higher nighttime CO concentrations are not accompanied by an enrichment in $\delta^{18}\text{O}$, which would be expected if the CO was produced by ozonolysis of unsaturated non-methane hydrocarbons (Röckmann et al., 1998). Having excluded ozonolysis as a significant contributor, we attribute the nighttime CO production to thermal degradation.

Appendix B: Soil CO flux as a function of soil temperature

Continuous soil temperature measurements were not available for the campaign periods in 2020 and 2021. Fortunately, soil temperature measurements were available for most of the year 2019. Soil temperatures at different depths were monitored in 10-minute intervals (STP01, Hykseflux). From previous measurements at the [field site](#), it was observed that the plateau soil temperature, measured with the manual sensor TP-101, agreed well with the continuous soil temperature measurements at 2 cm [depth](#) (differences generally <0.2 °C).

A simple soil temperature-based diurnal CO flux pattern was estimated, by use of the relationship shown in Fig. 2 ($F_{\text{CO}} = T_{\text{soil}}^{*1.29} - 32.5$), which indicates that soil CO uptake starts to dominate (the net flux turns from positive to negative) when temperatures drop below 25.2 °C. Fig. A1 shows the average soil temperatures of May 2019 and November 2019 (left [y-axis](#)), and the calculated soil CO flux (right [y-axis](#)). These months were chosen because they were close to the campaign months of 2020 and 2021 (May and October), and because they presented an uninterrupted data set for the complete month.

545 While the *average* monthly temperatures did not drop below the threshold of 25.2 °C, individual nights sometimes showed lower temperatures. The standard deviation of the average temperature (not shown) was used to estimate a CO flux range (dotted lines), which shows that ~~as well in the dry season as in the wet season~~ soil CO uptake ~~can take place~~can occur in the wet season as well as in the dry season. The daily averaged dry season (November) and wet season (May) flux was ~~resp-~~respectively 1.22 and 0.63 nmol CO m⁻² s⁻¹, indicating that, based on this simple model, the tropical forest soils are generally
550 a CO source year round.

Data availability. The measured CO and CO₂ mixing ratios and soil chamber fluxes, as presented in this study, have been uploaded to the open-access repository of Zenodo and can be found at <https://doi.org/10.5281/zenodo.10223554> (Asperen, 2023). The soil temperature data, as presented in the Appendix, are available on request from the co-authors AA, MS, PRT and JAFS.

Author contributions. HA designed and performed the field experiment, and wrote the paper, AA, BF, and SF provided access to the logistics
555 and infrastructure of the INPA-LBA ~~field site~~fieldsite, LRO, TLX, JM assisted in the setting up of the research infrastructure before and after the campaign weeks, SB assisted during the field campaign weeks and performed part of the flux measurements, MS, PRT, JAFS processed and provided the soil temperature data from the K34 tower, TR and CV analyzed and evaluated the isotopic flask samples, and TW, JN, TR, and ST reviewed and commented on the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

560 *Acknowledgements.* The study was funded by the DFG-project ‘Methane fluxes from seasonally flooded forests in the Amazon basin’ (project nr. 352322796). We are thankful for the support of the crew of the experimental ~~field site~~fieldsite ZF2, the research station managed by INPA-LBA (National Institute for Amazonian Research (INPA)- ~~The Large Scale~~the Large-Scale Biosphere-Atmosphere ~~Research Program in the Amazon~~Experiment in Amazonia (LBA)). We would like to thank Santiago Botía for providing the flasks for the ozonolysis experiment. We would also like to express our gratitude to the staff of LBA, for providing logistics, advice, and support during different
565 phases of this research.

References

- Andreae, M. O., Artaxo, P., Beck, V., Bela, M., Freitas, S., Gerbig, C., Longo, K., Munger, J. W., Wiedemann, K. T., and Wofsy, S. C.: Carbon monoxide and related trace gases and aerosols over the Amazon Basin during the wet and dry seasons, *Atmospheric Chemistry and Physics*, 12, 6041–6065, <https://doi.org/10.5194/acp-12-6041-2012>, 2012.
- 570 Araújo, A., Nobre, A., Kruijt, B., Elbers, J., Dallarosa, R., Stefani, P., Von Randow, C., Manzi, A., Culf, A., Gash, J., et al.: Comparative measurements of carbon dioxide fluxes from two nearby towers in a central Amazonian rainforest: The Manaus LBA site, *Journal of Geophysical Research: Atmospheres*, 107, LBA–58, 2002.
- Araújo, A. C., Kruijt, B., Nobre, A. D., Dolman, A. J., Waterloo, M. J., Moors, E. J., and de Souza, J. S.: Nocturnal accumulation of CO₂ underneath a tropical forest canopy along a topographical gradient, *Ecological Applications*, 18, 1406–1419, 2008.
- 575 Araújo, A. C. d.: Spatial variation of CO₂ fluxes and lateral transport in an area of terra firme forest in Central Amazonia, Ph.D. thesis, PhD thesis, Vrije Universiteit, Amsterdam, 2009.
- Asperen, H. v.: CO and CO₂ mixing ratios and flux measurements from the Amazon tropical rainforest [Data set], Zenodo, <https://doi.org/10.5281/zenodo.10223554>, 2023.
- Asperen, H. v., Warneke, T., and Notholt, J.: The Use of FTIR-Spectrometry in Combination with Different Biosphere-Atmosphere Flux
580 Measurement Techniques, Towards an Interdisciplinary Approach in Earth System Science: Advances of a Helmholtz Graduate Research School, pp. 77–84, 2015a.
- Asperen, H. v., Warneke, T., Sabbatini, S., Nicolini, G., Papale, D., and Notholt, J.: The role of photo- and thermal degradation for CO₂ and CO fluxes in an arid ecosystem, *Biogeosciences*, 12, 4161–4174, <https://doi.org/10.5194/bg-12-4161-2015>, 2015b.
- Asperen, H. v., Alves-Oliveira, J. R., Warneke, T., Forsberg, B., de Araújo, A. C., and Notholt, J.: The role of termite CH₄ emissions on the
585 ecosystem scale: a case study in the Amazon rainforest, *Biogeosciences*, 18, 2609–2625, 2021.
- Bartholomew, G. and Alexander, M.: Microbial metabolism of carbon monoxide in culture and in soil., *Applied and environmental microbiology*, 37, 932–937, 1979.
- Bruhn, D., Albert, K. R., Mikkelsen, T. N., and Ambus, P.: UV-induced carbon monoxide emission from living vegetation, *Biogeosciences Discussions*, 10, 2013.
- 590 Campen, H. I., Arévalo-Martínez, D. L., and Bange, H. W.: Carbon monoxide (CO) cycling in the Fram Strait, Arctic Ocean, *Biogeosciences*, 20, 1371–1379, 2023.
- Carmo, J. B. d., Keller, M., Dias, J. D., Camargo, P. B. d., and Crill, P.: A source of methane from upland forests in the Brazilian Amazon, *Geophysical Research Letters*, 33, 2006.
- Chambers, J. Q., Tribuzy, E. S., Toledo, L. C., Crispim, B. F., Higuchi, N., Santos, J. d., Araújo, A. C., Kruijt, B., Nobre, A. D., and Trumbore,
595 S. E.: Respiration from a tropical forest ecosystem: partitioning of sources and low carbon use efficiency, *Ecological Applications*, 14, 72–88, 2004.
- Clough, T. J., Rochette, P., Thomas, S. M., Pihlatie, M., Christiansen, J. R., and Thorman, R. E.: Global Research Alliance N₂O chamber methodology guidelines: Design considerations, *Journal of Environmental Quality*, 49, 1081–1091, 2020.
- Conrad, R.: Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS, N₂O, and NO), *Microbiological reviews*,
600 60, 609–640, 1996.
- Conrad, R. and Seiler, W.: Role of microorganisms in the consumption and production of atmospheric carbon monoxide by soil, *Applied and environmental microbiology*, 40, 437–445, 1980.

- Conrad, R. and Seiler, W.: Arid soils as a source of atmospheric carbon monoxide, *Geophysical Research Letters*, 9, 1353–1356, 1982.
- Conrad, R. and Seiler, W.: Influence of temperature, moisture, and organic carbon on the flux of H₂ and CO between soil and atmosphere:
605 Field studies in subtropical regions, *Journal of Geophysical Research: Atmospheres*, 90, 5699–5709, 1985.
- Conrad, R., Schütz, H., and Seiler, W.: Emission of carbon monoxide from submerged rice fields into the atmosphere, *Atmospheric Environment* (1967), 22, 821–823, 1988.
- Constant, P., Poissant, L., and Villemur, R.: Annual hydrogen, carbon monoxide and carbon dioxide concentrations and surface to air exchanges in a rural area (Québec, Canada), *Atmospheric Environment*, 42, 5090–5100, 2008.
- 610 Cowan, N., Helfter, C., Langford, B., Coyle, M., Levy, P., Moxley, J., Simmons, I., Leeson, S., Nemitz, E., and Skiba, U.: Seasonal fluxes of carbon monoxide from an intensively grazed grassland in Scotland, *Atmospheric Environment*, 194, 170–178, 2018.
- Criegee, R.: Mechanism of ozonolysis, *Angewandte chemie international edition in english*, 14, 745–752, 1975.
- Derendorp, L., Quist, J., Holzinger, R., and Röckmann, T.: Emissions of H₂ and CO from leaf litter of *Sequoiadendron giganteum*, and their dependence on UV radiation and temperature, *Atmospheric Environment*, 45, 7520–7524, 2011.
- 615 Funk, D. W., Pullman, E. R., Peterson, K. M., Crill, P. M., and Billings, W.: Influence of water table on carbon dioxide, carbon monoxide, and methane fluxes from taiga bog microcosms, *Global Biogeochemical Cycles*, 8, 271–278, 1994.
- Gödde, M., Meuser, K., and Conrad, R.: Hydrogen consumption and carbon monoxide production in soils with different properties, *Biology and fertility of soils*, 32, 129–134, 2000.
- Griffith, D., Deutscher, N., Caldow, C., Kettlewell, G., Riggensbach, M., and Hammer, S.: A Fourier transform infrared trace gas and isotope
620 analyser for atmospheric applications, *Atmospheric Measurement Techniques*, 5, 2481–2498, 2012.
- Hammer, S., Griffith, D. W. T., Konrad, G., Vardag, S., Caldow, C., and Levin, I.: Assessment of a multi-species in situ FTIR for precise atmospheric greenhouse gas observations, *Atmospheric Measurement Techniques*, 6, 1153–1170, <https://doi.org/10.5194/amt-6-1153-2013>, 2013.
- Heimann, M., Jordan, A., Brand, W. A., Lavrič, J. V., Moossen, H., and Rothe, M.: The atmospheric flask sampling program of MPI-BGC,
625 Version 13, 2022, Max Planck Institute for Biogeochemistry report, 2022.
- Ingersoll, R., Inman, R., and Fisher, W.: Soil's potential as a sink for atmospheric carbon monoxide, *Tellus*, 26, 151–159, 1974.
- INMET: station 8233-Eduardo Gomes, Instituto Nacional de Meteorologia, <https://mapas.inmet.gov.br/>, 2024.
- Khalil, M. and Rasmussen, R.: The global cycle of carbon monoxide: Trends and mass balance, *Chemosphere*, 20, 227–242, 1990.
- King, G. M.: Land use impacts on atmospheric carbon monoxide consumption by soils, *Global biogeochemical cycles*, 14, 1161–1172, 2000.
- 630 King, G. M. and Hungria, M.: Soil-atmosphere CO exchanges and microbial biogeochemistry of CO transformations in a Brazilian agricultural ecosystem, *Applied and environmental microbiology*, 68, 4480–4485, 2002.
- King, G. M. and Weber, C. F.: Distribution, diversity and ecology of aerobic CO-oxidizing bacteria, *Nature Reviews Microbiology*, 5, 107–118, 2007.
- Kirchhoff, V. W. J. H. and Marinho, E. V. d. A.: Surface carbon monoxide measurements in Amazonia, *Journal of Geophysical Research: Atmospheres*, 95, 16933–16943, 1990.
- 635 Kisselle, K. W., Zepp, R. G., Burke, R. A., de Siqueira Pinto, A., Bustamante, M., Opsahl, S., Varella, R. F., and Viana, L. T.: Seasonal soil fluxes of carbon monoxide in burned and unburned Brazilian savannas, *Journal of Geophysical Research: Atmospheres* (1984–2012), 107, LBA–18, 2002.
- Laasonen, A.: Biogenic carbon monoxide fluxes in four terrestrial ecosystems, MSc-thesis, University of Helsinki, 2021.

- 640 Lee, H., Rahn, T., and Throop, H.: An accounting of C-based trace gas release during abiotic plant litter degradation, *Global Change Biology*, 18, 1185–1195, 2012.
- Liu, L., Zhuang, Q., Zhu, Q., Liu, S., Van Asperen, H., and Pihlatie, M.: Global soil consumption of atmospheric carbon monoxide: an analysis using a process-based biogeochemistry model, *Atmospheric Chemistry and Physics*, 18, 7913–7931, 2018.
- Luizão, R. C., Luizão, F. J., Paiva, R. Q., Monteiro, T. F., Sousa, L. S., and Kruijt, B.: Variation of carbon and nitrogen cycling processes
645 along a topographic gradient in a central Amazonian forest, *Global Change Biology*, 10, 592–600, 2004.
- Marques, J., Luizão, F., Teixeira, W., and Araújo, E.: Carbono orgânico em solos sob floresta na Amazônia central, *Congresso Norte Nordeste de Pesquisa e Inovação, CONNEPI*, 7, 2013.
- Martius, C., Waßmann, R., Thein, U., Bandeira, A., Rennenberg, H., Junk, W., and Seiler, W.: Methane emission from wood-feeding termites in Amazonia, *Chemosphere*, 26, 623–632, 1993.
- 650 Moxley, J. and Smith, K.: Carbon monoxide production and emission by some Scottish soils, *Tellus B: Chemical and Physical Meteorology*, 50, 151–162, 1998.
- Müller, D.: Water-atmosphere greenhouse gas exchange measurements using FTIR spectrometry, PhD-thesis, Universität Bremen, 2015.
- Paulson, S. E. and Seinfeld, J. H.: Development and evaluation of a photooxidation mechanism for isoprene, *Journal of Geophysical Research: Atmospheres*, 97, 20703–20715, 1992.
- 655 Pihlatie, M., Rannik, U., Haapanala, S., Peltola, O., Shurpali, N., Martikainen, P. J., Lind, S., Hyvönen, N., Virkajarvi, P., Zahniser, M., and Mammarella, I.: Seasonal and diurnal variation in CO fluxes from an agricultural bioenergy crop, *Biogeosciences*, 13, 5471–5485, <https://doi.org/10.5194/bg-13-5471-2016>, 2016.
- Potter, C. S., Klooster, S. A., and Chatfield, R. B.: Consumption and production of carbon monoxide in soils: a global model analysis of spatial and seasonal variation, *Chemosphere*, 33, 1175–1193, 1996.
- 660 Querino, C. A. S., Smeets, C. J. P. P., Vigano, I., Holzinger, R., Moura, V., Gatti, L. V., Martynowski, A., Manzi, A. O., de Araújo, A. C., and Röckmann, T.: Methane flux, vertical gradient and mixing ratio measurements in a tropical forest, *Atmospheric Chemistry and Physics*, 11, 7943–7953, <https://doi.org/10.5194/acp-11-7943-2011>, 2011.
- Quesada, C., Lloyd, J., Schwarz, M., Patino, S., Baker, T., Czimczik, C., Fyllas, N., Martinelli, L., Nardoto, G., Schmerler, J., et al.: Variations in chemical and physical properties of Amazon forest soils in relation to their genesis, *Biogeosciences*, 7, 1515–1541, 2010.
- 665 Röckmann, T., Brenninkmeijer, C. A., Neeb, P., and Crutzen, P. J.: Ozonolysis of nonmethane hydrocarbons as a source of the observed mass independent oxygen isotope enrichment in tropospheric CO, *Journal of Geophysical Research: Atmospheres*, 103, 1463–1470, 1998.
- Sanderson, M. G.: Emission of carbon monoxide by vegetation and soils, Met office, Hadley Centre technical note 36, 2002.
- Sanhueza, E., Donoso, L., Scharffe, D., and Crutzen, P. J.: Carbon monoxide fluxes from natural, managed, or cultivated savannah grasslands, *Journal of Geophysical Research: Atmospheres*, 99, 16421–16427, 1994.
- 670 Schade, G. W., Hofmann, R.-M., and Crutzen, P. J.: CO emissions from degrading plant matter, *Tellus B*, 51, 889–908, 1999.
- Scharffe, D., Hao, W. M., Donoso, L., Crutzen, P. J., and Sanhueza, E.: Soil fluxes and atmospheric concentration of CO and CH₄ in the northern part of the Guayana Shield, Venezuela, *Journal of Geophysical Research: Atmospheres*, 95, 22475–22480, 1990.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2016.
- Simon, E., Lehmann, B., Ammann, C., Ganzeveld, L., Rummel, U., Meixner, F., Nobre, A. D., Araujo, A., and Kesselmeier, J.: Lagrangian
675 dispersion of ²²²Rn, H₂O and CO₂ within Amazonian rain forest, *Agricultural and forest meteorology*, 132, 286–304, 2005.
- Spratt, H. G. and Hubbard, J. S.: Carbon monoxide metabolism in roadside soils, *Applied and environmental microbiology*, 41, 1192–1201, 1981.

- Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Bernsten, T., Collins, W., Fuzzi, S., Gallardo, L., Kiendler-Scharr, A., Klimont, Z., et al.: Short-lived climate forcers, in: *Climate Change 2021: The Physical Basis. Contribution of Working Group I to the Sixth Assessment report of the IPCC*, vol. 2021, p. 853, 2021.
- 680 Tarr, M. A., Miller, W. L., and Zepp, R. G.: Direct carbon monoxide photoproduction from plant matter, *Journal of Geophysical Research: Atmospheres* (1984–2012), 100, 11 403–11 413, 1995.
- Tóta, J., Fitzjarrald, D. R., Staebler, R. M., Sakai, R. K., Moraes, O. M., Acevedo, O. C., Wofsy, S. C., and Manzi, A. O.: Amazon rain forest subcanopy flow and the carbon budget: Santarém LBA-ECO site, *Journal of Geophysical Research: Biogeosciences*, 113, 2008.
- 685 Trumbore, S. E., Keller, M., Wofsy, S., and Da Costa, J.: Measurements of soil and canopy exchange rates in the Amazon rain forest using ^{222}Rn , *Journal of Geophysical Research: Atmospheres*, 95, 16 865–16 873, 1990.
- Valentine, R. L. and Zepp, R. G.: Formation of carbon monoxide from the photodegradation of terrestrial dissolved organic carbon in natural waters, *Environmental science & technology*, 27, 409–412, 1993.
- Whalen, S. and Reeburgh, W.: Carbon monoxide consumption in upland boreal forest soils, *Soil Biology and Biochemistry*, 33, 1329–1338, 690 2001.
- Yonemura, S., Kawashima, S., and Tsuruta, H.: Carbon monoxide, hydrogen, and methane uptake by soils in a temperate arable field and a forest, *Journal of Geophysical Research: Atmospheres* (1984–2012), 105, 14 347–14 362, 2000.
- Zanchi, F. B., Meesters, A. G., Kruijt, B., Kesselmeier, J., Luizão, F. J., and Dolman, A. J.: Soil CO_2 exchange in seven pristine Amazonian rain forest sites in relation to soil temperature, *Agricultural and Forest Meteorology*, 192, 96–107, 2014.
- 695 Zhang, Y., Xie, H., Fichot, C. G., and Chen, G.: Dark production of carbon monoxide (CO) from dissolved organic matter in the St. Lawrence estuarine system: Implication for the global coastal and blue water CO budgets, *Journal of Geophysical Research: Oceans*, 113, 2008.
- Zheng, B., Chevallier, F., Yin, Y., Ciais, P., Fortems-Cheiney, A., Deeter, M. N., Parker, R. J., Wang, Y., Worden, H. M., and Zhao, Y.: Global atmospheric carbon monoxide budget 2000–2017 inferred from multi-species atmospheric inversions, *Earth System Science Data*, 11, 1411–1436, <https://doi.org/10.5194/essd-11-1411-2019>, 2019.
- 700 Zuo, Y. and Jones, R. D.: Photochemistry of natural dissolved organic matter in lake and wetland waters—production of carbon monoxide, *Water Research*, 31, 850–858, 1997.




Figure 1. Upper and middle row; CO and CO₂ soil fluxes at 5 different locations on the plateau (left, reddish colors), and in the valley (right, blueish colors). Lighter colors indicate dry season (DS), darker colors indicate wet season (WS). Each location was measured 3 times during each campaign, the error bars indicate the standard deviation of the mean of the 3 measurements. CO fluxes are based on bag mixing ratio measurements, CO₂ fluxes are based on ICOS-analyzer measurements. The lower row shows the ratio between the soil CO and CO₂ fluxes (circles and triangles). In addition, the ratio's (average DS and its standard deviation), WS ratios as measured by the canopy layer budget method (tower height of 5 m) are shown in (solid lines, the standard deviations of these averages are indicated with dotted) lines.




Figure 2. CO fluxes (upper row) and CO₂ fluxes (middle row) on the plateau (left column) and in the valley (right column). Dry season measurements are indicated with circles, wet season measurements are indicated with triangles. Correlation coefficients between T (or VWC) and soil CO fluxes (or CO₂ fluxes) are given in the legend, and linear regression lines are shown for the cases where $R^2 > 0.5$. The formula given in the upper left graph indicates the relationship found between the manual soil temperature measurements and the measured CO fluxes. Lower row: correlation between soil T and soil VWC, correlation coefficient given in the legend. In addition, three linear regression lines are shown for plateau measurements: for both campaigns together (DS+WS, $R^2=0.66$) and for the individual campaigns DS ($R^2=0.23$) and WS ($R^2=0.30$). The shift in regression line indicates that plateau environmental drivers might differ between seasons.




Figure 3. Tower CO mixing ratios during the dry season campaign (upper row) and the wet season campaign (lower row). Canopy height is ~ 35 m. During the first 5 days of the dry season campaign, a leak was present in the 36 m inlet line, wherefore data from this height is missing. Despite the variation, a general tendency, with higher CO mixing ratios below the canopy, is visible during both periods.




Figure 4. Vertical Average vertical CO profiles for each dry season (DS, upper row) campaign nights (left figure) and wet season (WS, lower row) campaign nights (right figure). Error bars indicate the standard deviation of dCO_{-15m} and dCO_{-36m} for the six 1h-time windows per night; this average was only calculated if dCO_{-36m} was consistently negative over the whole night. Individual 1h-profiles per night are shown in Fig. A2. The black dotted vertical line indicates the zero line ($dCO=0$). An average nighttime profile was calculated if, and the 36 m height showed consistent lower mixing ratios in comparison to the 5 m height (dark red-green dotted horizontal line) indicates the height of the canopy. The last column show these average profiles, and is plotted on a narrower x-axis range.




Figure 5. Expected CO emission of soil and senescent leaf material, expressed per surface area. For senescent leaf CO emission (left axes, green triangles and squares), the laboratory CO emissions ($\text{nmol gr}_{leaves}^{-1} \text{ min}^{-1}$) were converted to seconds (s^{-1}), and were multiplied by a senescent leaf density of resp., respectively 117 and $67 \text{ gr}_{leaves} \text{ m}^{-2}$ for the plateau and valley ecosystem (Luizão et al., 2004), so that CO production is expressed in ' $\text{nmol m}^{-2} \text{ s}^{-1}$ '. For the estimate of CO emission of a 10-cm soil layer (right axes, dark red diamonds), the laboratory emissions ($\text{nmol gr}_{soil}^{-1} \text{ min}^{-1}$) were converted to seconds (s^{-1}), and were combined with a soil bulk density of 1.05 g cm^{-3} (Marques et al., 2013), so that soil CO production is expressed in ' $\text{nmol m}^{-2} \text{ s}^{-1}$ '. The set-in figure shows a zoom-in of senescent leaves and soil emissions, plotted on the same y-axes y-axis scale, visualizing the expected dominance of soil CO emissions over senescent leaf CO emissions.

Table 1. Overview of the soil chamber CO and CO₂ fluxes and the different ecosystem CO flux estimates. **Upper part:** The two left columns first three rows show the soil CO and CO₂ fluxes (range and mean (stdsd)), and its $\frac{\Delta CO}{\Delta CO_2}$, as measured with the flux chamber technique. The third column shows $\frac{\Delta CO}{\Delta CO_2}$ ratios of these flux chamber measurements (range Each location had 5 collars and mean was measured 3 times per campaign (stdn=15)). **Middle part:** The fourth column-row shows the $\frac{\Delta CO}{\Delta CO_2}$ ratio-ratios of the nighttime increase (range and mean (std)), on which the estimated ecosystem CO flux is based fluxes (fifth column, canopy layer budget estimate, row) are based (range and mean (stdsd)). The second-to-last column shows On the average nighttime vertical d-36m gradient plateau, for each campaign night (mean (std) DS: 9 nights, WS: 7 nights)), and the last column ratios of five 2h-time windows were calculated; the average is based on time windows with ratios with R²>0.90. In the valley, for each campaign night, the ratio of one 9h-time window was used; the average is based on the nights with ratios with R²>0.90 (n=4 in DS, n=5 in WS). **Lower part:** The sixth row shows the vertical profile dCO-36m values, on which the estimated ecosystem CO flux fluxes are based (seventh row). Range and mean (stdsd) are given of the 4 DS nights and the 5 WS nights (same as shown in Fig. 4) based on these d-36m values, with each 6 time windows per night (n=24 & n=30). DS stands for dry season, and WS stands for wet season. Vertical profiles were not determined (n.d.) in the valley.

| | Soil flux Plateau DS | Soil CO ₂ flux Plateau WS | $\Delta CO/\Delta CO_2$ Valley DS | $\Delta CO/\Delta CO_2$ Valley WS |
|---|--|---------------------------------------|---|--|
| Ecosystem flux | dCO-36m | Ecosystem flux | | |
| (chamber) Soil CO flux (nmol m ⁻² s ⁻¹) | (chamber) 0.62 to 2.26 | (chamber) -0.18 to 0.49 | (nighttime increase) -0.03 to 3.36 | (based on $\Delta CO/\Delta CO_2$) 0.19 to 1.14 |
| Flux chamber measurement | (vertical profile) 1.23 (0.52), n=15 | 0.20 (0.19), n=15 | 1.83 (0.97), n=15 | 0.62 (based on dCO-36m) 0.33, n=15 |
| Soil CO₂ flux (μmol m ⁻² s ⁻¹) | | | | |
| | nmol m ⁻² s ⁻¹ | | | |
| | (+) 2.16 to 5.57 | (+) 2.11 to 8.03 | nmol m ⁻² s ⁻¹ 2.38 to 7.93 | nmol 1.59 to 5.21 |
| Flux chamber measurement | nmol m ⁻² s ⁻¹ 3.53 (1.02), n=15 | 4.83 (1.87), n=15 | 4.18 (1.40), n=15 | 3.21 (1.00), n=15 |
| Plateau $\Delta CO/\Delta CO_2$ ratio (-) | 0.62 to 2.26 0.18 to 0.65 | 2.16 to 5.57 -0.05 to 0.11 | 0.18 to 0.65 0.00 to 1.07 | 5m: 0.16 to 0.53, 0.27 (0.09) 0.06 to 0.59 |
| From chamber flux measurement | 5m: -1.25 to -4.13, -2.11 (0.70) 0.35 (0.12), n=15 | -9.1 (1.9) 0.04 (0.04), n=15 | 1.4 (0.3) 0.47 (0.27), n=15 | 0.22 (0.16), n=15 |
| DS | 1.23 (0.52) | 3.53 (1.02) | 0.35 (0.12) | 15m: -0.15 to 0.53, 0.23 (0.12) |
| | 15m: -1.17 to -4.13, -1.79 (0.94) | | | |
| $\Delta CO/\Delta CO_2$ ratio (-) | | | | |
| | 5 m: 0.16 to 0.53 | 5 m: 0.12 to 0.42 | 1 m: 0.04 to 0.25 | 1 m: 0.23 to 0.40 |
| From nighttime increase | 0.27 (0.09), n=13 | 0.24 (0.08), n=14 | 0.14 (0.25), n=4 | 0.30 (0.05), n=5 |
| Plateau | -0.18 | | | |
| | 15 m: -0.15 to 0.49 0.53 | 2.11 15 m: -0.69 to 8.03 0.54 | -0.05 -0.11 | 5m: 0.12 to 0.42, 0.24 (0.08) |
| | 5m: 0.23 (0.12), n=20 | 0.20 (0.25), n=14 | | |
| Ecosystem CO flux (nmol m ⁻² s ⁻¹) | | | | |
| | 5 m: 1.25 to 4.13 | 5 m: 0.94 to 3.28 - 1.87 (0.62), n=14 | 1 m: 0.31 to 1.95 | 1 m: 1.79 to 3.12 |
| Based on $\Delta CO/\Delta CO_2$ ratio | 2.11 (0.70), n=13 | | -5.6 (4.3) -1.09 (1.95), n=4 | 0.9 (0.6) 2.34 (0.39), n=5 |
| WS | 0.20 (0.19) | 4.83 (1.87) | 0.04 (0.04) | 15m: -0.69 to 0.54, 0.20 (0.25) |




Figure A1. Left **y-axis**: Average soil temperature at 2 cm **depth** in the dry season (DS) of 2019 (October) and the wet season (WS) of 2019 (May) (standard deviations of the average soil temperatures are not shown). Right **y-axis**: Modeled soil CO flux for the DS and WS (solid lines) and its standard deviations (dotted lines), based on the soil temperatures (and their standard deviation) at 2 cm depth (relationship shown in Fig. 2). The monthly mean calculated CO flux for the DS and WS is given in the legend, and is in $\text{nmol CO m}^{-2} \text{s}^{-1}$.




Figure A2. Vertical CO profiles for each DS campaign night (upper row) and WS campaign night (lower row). Each line shows the average vertical profile of that 1h-time window, error bars indicate the standard deviation of the average of the 3 measurements in that time window. The black dotted vertical line indicates the zero line ($dCO=0$), and the green dotted horizontal line indicates the height of the canopy. The night of 11-12 May showed a strong CO peak at 15 m after midnight (also visible in Fig. 3), which we cannot explain.

Table A1. Sampling time of flask, measured CO mixing ratios (with sd in brackets), and δ
 ^{18}O -CO of flasks (with sd in brackets).