

# Complex refractive index and single scattering albedo of Icelandic dust in the shortwave spectrum

Clarissa Baldo<sup>1</sup>, Paola Formenti<sup>2</sup>, Claudia Di Biagio<sup>2</sup>, Gongda Lu<sup>1,a</sup>, Congbo Song<sup>1,b</sup>, Mathieu Cazaunau<sup>3</sup>, Edouard Pangui<sup>3</sup>, Jean-Francois Doussin<sup>3</sup>, Pavla Dagsson-Waldhauserova<sup>4,5</sup>, Olafur Arnalds<sup>5</sup>, David Beddows<sup>1</sup>, A. Robert MacKenzie<sup>1</sup>, and Zongbo Shi<sup>1</sup>

<sup>1</sup> School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, United Kingdom

<sup>2</sup> Université Paris Cité and Univ Paris Est Creteil, CNRS, LISA, F-75013 Paris, France

<sup>3</sup> Univ Paris Est Creteil and Université Paris Cité, CNRS, LISA, F-94010 Créteil, France

10 <sup>4</sup> Agricultural University of Iceland, Keldnaholt, Reykjavik, Iceland

<sup>5</sup> Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic

<sup>a</sup> now at: Department of Geography, University College London, London, WC1E 6BT, United Kingdom

<sup>b</sup> now at: National Centre for Atmospheric Science (NCAS), Department of Earth and Environmental Science, University of Manchester, M13 9PL, Manchester, United Kingdom

15

*Correspondence to:* Clarissa Baldo (c.baldo@bham.ac.uk)

**Abstract** Icelandic dust can impact the radiative budget in high-latitude regions directly by affecting light absorption and scattering and indirectly by changing the surface albedo after dust deposition. This tends to produce a positive radiative forcing. However, the limited knowledge of the spectral optical properties of Icelandic dust prevents an accurate assessment of these radiative effects. Here, the spectral single scattering albedo (SSA) and the complex refractive index ( $m = n - ik$ ) of Icelandic dust from five major emission hotspots were retrieved between 370-950 nm using online measurements of size distribution and spectral absorption ( $\beta_{\text{abs}}$ ) and scattering ( $\beta_{\text{sca}}$ ) coefficients of particles suspended in a large-scale atmospheric simulation chamber. The SSA( $\lambda$ ) estimated from the measured  $\beta_{\text{abs}}$  and  $\beta_{\text{sca}}$  increased from 0.90-0.94 at 370 nm to 0.94-0.96 at 950 nm in Icelandic dust from the different hotspots, which falls within the range of mineral dust from northern Africa and eastern Asian. The spectral complex refractive index was retrieved by minimizing the differences between the measured  $\beta_{\text{abs}}$  and  $\beta_{\text{sca}}$  and those computed using the Mie theory for spherical and internally homogeneous particles, using the size distribution data as input. The real part of the complex refractive index ( $n(\lambda)$ ) was found to be 1.60-1.61 in the different samples and independent on wavelength. The imaginary part ( $k(\lambda)$ ) was almost constant with wavelength and was found to be around 0.004 at 370 nm and 0.002-0.003 at 950 nm. The estimated complex refractive index was close to the initial estimates based on the mineralogical composition, also suggesting that the high magnetite content observed in Icelandic dust may contribute to its high absorption capacity in the shortwave spectrum. The  $k(\lambda)$  values retrieved for Icelandic dust are at the upper end of the reported range for low-latitude dust (e.g., from the Sahel). Furthermore, Icelandic dust tends to be more absorbing toward the near-infrared. In Icelandic dust,  $k(\lambda)$  between 660-950 nm was 2-8 times higher than most of the dust

35 samples sourced in northern Africa and eastern Asia. This suggests that Icelandic dust may have a stronger positive direct radiative forcing on climate which has not been accounted for in climate predictions.

## 1 Introduction

Iceland is a major contributor to the dust aerosol loading over high-latitude ( $>60^\circ$  N) regions in the Northern Hemisphere (Arnalds et al., 2016; Baddock et al., 2017; Bullard et al., 2016; Dagsson-Waldhauserova et al., 2019; Groot Zwaaftink et al., 2017; Groot Zwaaftink et al., 2016; Meinander et al., 2022; Prospero et al., 2012). With an area of more than 45,000 km<sup>2</sup>, the Icelandic desert is the largest in the Arctic (Arnalds et al., 2016) and being subject to intense wind erosion, Iceland is one of the most active dust source areas in the world, producing about 4.6-40 Tg of dust per year (Arnalds et al., 2016; Groot Zwaaftink et al., 2016) and 34-135 dust events per year (Dagsson-Waldhauserova et al., 2014; Nakashima and Dagsson-Waldhauserová, 2019). Near glaciers, dust hotspots cover about 500 km<sup>2</sup> and are fed by fluvial-glacial processes acting on volcanic deposits and can produce millions of tons of dust during single dust events (Arnalds et al., 2016).

Icelandic dust can reach several kilometres in altitude, as shown in a study by Dagsson-Waldhauserova et al. (2019). They conducted six winter balloon launches in southern west Iceland during 2013-2016. Vertical profile measurements were collected using the Light Optical Aerosol Counter (LOAC) instrument, which provided aerosol size number distributions, concentrations, typologies, and basic meteorological parameters. LOAC estimates particle typologies for black carbon, mineral dust, volcanic dust, volcanic ash, ice particles, sea salt, and liquid particles, based on specific range referred to as 'specification zone' retrieved during laboratory experiments with different aerosols (Renard et al., 2016). The 'speciation zone' of Icelandic volcanic dust was established based on laboratory simulations using volcanic dust and ash samples collected in Iceland. Volcanic dust particles up to 10  $\mu$ m were detected within 900 m altitude, those up to 5  $\mu$ m mostly within 3.5 km altitude, while submicron particles were detected at altitudes up to 6 km. LOAC measurements were confirmed by in-situ ambient air quality monitoring stations of the Environmental Agency of Iceland for surface, CALIPSO and HYSPLIT model at different altitudes and MODIS true colour images with dust plumes, pointing to the local Icelandic dust sources during the Polar Vortex conditions above Iceland.

Icelandic dust can be also transported over long distances to the high Arctic (Baddock et al., 2017; Groot Zwaaftink et al., 2017; Groot Zwaaftink et al., 2016; Moroni et al., 2018; Prospero et al., 2012). Arctic dust events typically occur during summer/autumn in northeastern Iceland (Dagsson-Waldhauserova et al., 2013; Dagsson-Waldhauserova et al., 2014; Meinander et al., 2022; Nakashima and Dagsson-Waldhauserová, 2019), while subarctic dust events mainly occur during winter/spring in southern Iceland (Dagsson-Waldhauserova et al., 2014; Dagsson-Waldhauserova et al., 2015; Dagsson-Waldhauserova et al., 2019).

A modelling study of Groot Zwaaftink et al. (2016) estimated that Iceland accounts for approximately 2% of the total atmospheric dust load in the Arctic. According to this study, high-latitude dust contributes to about 27% of the dust load, while the remaining amount is dominated by low-latitude dust from Africa and Asia. High-latitude dust is confined at low

altitudes due to the stratified atmosphere in cold climates and dominates dust surface concentrations (Bullard, 2017; Groot Zwaafink et al., 2016; Shi et al., 2022). Dust concentration and dust load have important consequences for the radiative balance in the Arctic, where the radiative effect of light-absorbing particles is enhanced by the presence of snow and ice (Schmale et al., 2021). Icelandic dust absorbs solar – shortwave (SW) radiation (Zubko et al., 2019). Absorbing aerosol particles in the Arctic upper troposphere tend to cool the surface, whereas those located at lower altitudes tend to warm the surface (Schmale et al., 2021). The estimated instantaneous radiative forcing (IRF) of Icelandic dust is  $0.005 \text{ W m}^{-2}$  at the top of the atmosphere (TOA) via dust-radiation interaction, which is lower but still significant compared to the IRF at the bottom of the atmosphere (BOA) via dust deposition onto snow-covered areas,  $0.02 \text{ W m}^{-2}$  (Kylling et al., 2018).

Icelandic dust accounts for around 5.8% of the total dust deposition in the Arctic. Experiments have shown that the deposition of volcanic dust particles from Iceland onto snow and ice can reduce the surface albedo and influence melting (Dragosics et al., 2016; Gunnarsson et al., 2021; Meinander et al., 2014; Möller et al., 2016; Möller et al., 2018; Peltoniemi et al., 2015; Wittmann et al., 2017). Peltoniemi et al. (2015) found that volcanic sand from the Mýrdalssandur hotspot in southern Iceland reduced the snow albedo in the wavelength ( $\lambda$ ) range 400-2500 nm similarly to black carbon (BC). Laboratory measurements conducted on pure volcanic sand (deposited particles) showed a spectral reflectance of less than 0.04, which is close to a black body (0.0) (Dagsson-Waldhauserova, 2014; Peltoniemi et al., 2015).

In addition, Icelandic dust is a source of ice-nucleating particles (e.g., Sanchez-Marroquin et al., 2020; Shi et al., 2022). Sanchez-Marroquin et al. (2020) estimated that during summer days, Icelandic dust is the dominant source of ice-nucleating dust particles over areas near the eastern coast of Greenland. The increase in ice-nucleating particles can reduce the amount of supercooled liquid water and consequently the albedo of mixed-phase clouds (Vergara-Temprado et al., 2018), which have a large impact on the radiative budget in the Arctic (Morrison et al., 2012; Murray et al., 2021). Icelandic dust also contributes to the deposition of dissolved iron (Fe) to the subpolar North Atlantic Ocean (Arnalds et al., 2014; Baldo et al., 2020), which has the potential to alter nutrient biogeochemical cycling and primary production in the surface ocean (Achterberg et al., 2013; Achterberg et al., 2018). The deposition of nutrients such as phosphorus (P) and Fe from mineral dust can also promote the blooming of algae in ice, which consequently reduces the ice albedo and accelerates melting (Cook et al., 2020; Lutz et al., 2015; McCutcheon et al., 2021).

The mineralogical composition of Icelandic dust is considerably different from typical low-latitude dust such as northern African and eastern Asian dust (Baldo et al., 2020), and the absorption of solar radiation by mineral dust is strongly related to the content of total Fe and Fe oxide minerals (Caponi et al., 2017; Derimian et al., 2008; Di Biagio et al., 2019; Engelbrecht et al., 2016; Formenti et al., 2014; Lafon et al., 2006; Moosmuller et al., 2012; Sokolik and Toon, 1999). Icelandic dust primarily consists of amorphous basaltic material, but it is also enriched in Fe contributing to over 10% of the total dust mass, with relatively high magnetite ( $\text{Fe}_3\text{O}_4$ ) content (Baldo et al., 2020). This suggests that the optical properties of Icelandic dust may be also different from that of low-latitude dust.

The complex refractive index, represented by  $m = n - ik$ , links the particle chemical composition and optical properties, with the real part  $n$  defining the scattering properties and the imaginary part  $k$  the absorption properties (Bohren and Hufmann,

1998). In a study of Zubko et al. (2019), the index  $k$  of suspended particles from the Mýrdalssandur hotspot was found to be 0.01 at 647 nm, which is around one order of magnitude higher than the average values reported for mineral dust from northern Africa and eastern Asia by Di Biagio et al. (2019). However, the spectral dependence of  $k$  is unknown to date.

In Baldo et al. (2020), we studied the chemical and mineralogical composition of Icelandic dust from several major emission hotspots and found that it is distinctive and could be highly light-absorbing. Here, we go a step forward and, for the first time, determine the complex refractive index and single scattering albedo (SSA) of Icelandic dust over a broad range of  $\lambda$  (370-950 nm) based on in situ measurements of the size distribution and optical properties of suspended particles generated from natural parent soils.

## 2 Methodology

This section illustrates the approach used to retrieve the SSA and complex refractive index of Icelandic dust in the SW range which was based on the method by Di Biagio et al. (2019). SSA is the ratio of the scattering coefficient ( $\beta_{\text{sca}}$ ) to the extinction coefficient ( $\beta_{\text{ext}}$ ) – the sum of  $\beta_{\text{sca}}$  and the absorption coefficient ( $\beta_{\text{abs}}$ ). The spectral coefficients  $\beta_{\text{abs}}$  and  $\beta_{\text{sca}}$  and the size distribution of suspended dust particles were measured using the multi-instrument atmospheric simulation chamber CESAM – French acronym for Experimental Multiphasic Atmospheric Simulation Chamber (Wang et al., 2011). The complex refractive index was retrieved from the comparison between the spectral coefficients  $\beta_{\text{abs}}$  and  $\beta_{\text{sca}}$  measured and computed with Mie theory assuming that the particles are spherical and internally homogeneous, using the size distribution data as input. Although mineral dust aerosol is neither spherical nor internally homogeneous (e.g., Kandler et al., 2007; Okada et al., 2001), we have simplified the calculations of the optical properties to allow comparison with previous studies (Deguine et al., 2020; Di Biagio et al., 2019; Kylling et al., 2018; Reed et al., 2018; Zubko et al., 2019). A schematic diagram of the method used to retrieve the complex refractive index is shown in Fig. 1.

### 2.1 Experimental setup

The size distribution and optical properties of Icelandic dust were measured by the use of CESAM, a 4.2 m<sup>3</sup> stainless steel atmospheric simulation chamber (Wang et al., 2011). The experiments were conducted using surface sediment samples from five major dust hotspots in Iceland (Baldo et al., 2020) – D3 (Dyngjúsandur), H55 (Hagavatn), Land1 (Landeyjarsandur), Maeli2 (Mælifellssandur) and MIR45 (Mýrdalssandur). Table S1 contains the information on sampling sites. As described in Baldo et al. (2020), 15 g of sediments (previously sieved to <1 mm) was placed in a Büchner flask, this was shaken for 5 min at 70 Hz on a sieve shaker (Retsch AS200), then mixed with nitrogen carrier gas and injected into the CESAM chamber at 10 L min<sup>-1</sup> for 10 min. Once injected in CESAM, the aerosolized dust particles were allowed to mix for around 10 min inside the chamber. A four-blade stainless-steel fan located at the bottom of the chamber ensured that the particle concentration was homogeneous inside the chamber. All experiments were conducted at ambient temperature and relative humidity < 2 %. A particle-free N<sub>2</sub>/O<sub>2</sub> mixture was continuously injected into the chamber to balance the proportion of air flow sampled by the

different instruments. The chamber was manually cleaned between experiments with different samples to avoid any cross-contamination. The background mass concentrations of dust aerosol in the chamber varied from 0.1 to 3.0  $\mu\text{g m}^{-3}$ .

### 2.1.1 Instrumentation

135 The physical and chemical properties of the suspended dust were monitored over the duration of the experiments (approximately 3 hours) using different instruments connected to the chamber (Fig. S1).

The particle number size distribution of dust aerosols (in the units of  $\text{cm}^{-3}$ ) was measured using a scanning mobility particle sizer (SMPS) (TSI Inc., DMA model 3080, CPC model 3772; 2.0 and 0.2  $\text{L min}^{-1}$  sheath–aerosol flow rates; 3-min time resolution), and two optical particle counters (OPCs), GRIMM (Grimm Inc., model 1.109; 655 nm operating  $\lambda$ ; 1.2  $\text{L min}^{-1}$  flow rate; 6 s time resolution) and SkyGRIMM (Grimm Inc., model 1.129; 655 nm operating  $\lambda$ ; 1.2  $\text{L min}^{-1}$  flow rate; 6 s time resolution). The SMPS measured the particle number concentration in the electrical mobility diameter ( $D_m$ ) range from 0.019 to 0.870  $\mu\text{m}$ . The OPC instruments GRIMM and SkyGRIMM measured the particle number concentration in the optical-equivalent diameter ( $D_{op}$ ) range from 0.25 to 32  $\mu\text{m}$ . The SMPS and SkyGRIMM were located at the bottom of the chamber, while the GRIMM was installed at the top of the chamber (about 1 m above the SMPS and the SkyGRIMM) to  
145 verify that the concentration of dust aerosols in the chamber was spatially uniform. The percentage difference between the number concentrations in the size range measured by GRIMM and SkyGRIMM was < 10% for the samples D3 and MIR45, < 5% for the samples Land1 and Maeli2, and < 2% for the sample H55 suggesting that homogeneous conditions were obtained during the experiments.

The scattering coefficient  $\beta_{sca}$  (in the units of  $\text{Mm}^{-1}$ ;  $\text{Mm}^{-1} = 10^{-6} \text{ m}^{-1}$ ) of the suspended particles was measured by a three-  
150 wavelength integrating nephelometer (TSI Inc., model 3563; operating at 450, 550, and 700 nm; 2  $\text{L min}^{-1}$  flow rate, 1 s time resolution). The absorption coefficient  $\beta_{abs}$  (in the units of  $\text{Mm}^{-1}$ ) was retrieved by a seven-wavelength aethalometer (Magee Sci., model AE31; operating at 370, 470, 520, 590, 660, 880, and 950 nm; 2  $\text{L min}^{-1}$  flow rate, 2-min time resolution). As detailed in Baldo et al. (2020), dust aerosol samples were also collected for 1 to 3 hours, each with a flow rate of 7  $\text{L min}^{-1}$ , using custom-made filter samplers. The collected filter samples were analyzed offline to determine the chemical and  
155 mineralogical compositions of the suspended particles (Baldo et al., 2020).

The sampling system consisted of stainless-steel tubes ( $\sim 14\text{--}58$  cm length, 9.5 mm inner diameter) located inside the chamber extracting air. For some instruments, conductive silicone tubing (TSI Inc.,  $\sim 30\text{--}65$  cm length, 5 mm inner diameter) was also used as an external connection between the chamber and the instrument. The sampling lines were designed to be as straight and direct as possible to minimize particle loss. The total length of the sampling line varied  
160 depending on the instrument, 100 cm for the nephelometer, 86 cm for the aethalometer, 79 cm for the SMPS, 58 cm for the SkyGRIMM and the aerosol filter samplers, and 14 cm for the GRIMM.

To allow the comparison of measurements from different instruments, all data in the following analysis were averaged over 12-min intervals from the start of the filter sampling. The data analysis was conducted on the measurements starting after 30 min from the injection peak.

**2.2.1 Calculation of the aerosol geometrical size distribution from SMPS and GRIMM measurements**

The measurements from SMPS and GRIMM were combined into a single geometrical size distribution. Appendix A describes in detail the SMPS and GRIMM corrections. To combine SMPS and GRIMM measurements, first, the SMPS particle number concentration,  $dN(D_m)/d\log D_m$ , and GRIMM particle number concentration,  $dN(D_{op})/d\log D_{op}$ , were both converted into geometrical size distributions,  $dN(D_{g,SMPS})/d\log D_{g,SMPS}$  and  $dN(D_{g,GRIMM})/d\log D_{g,GRIMM}$ , respectively (see Fig. 2). For the SMPS, the volume-equivalent diameters ( $D_g$ ) were retrieved from the mobility diameters  $D_m$  using a range of dynamic shape factors  $\chi$  in Eq. (A1). The interval of  $\chi$  values (from 1.6 to 2.0 by 0.1 steps) was selected according to the aspect ratios observed in fine Icelandic dust particles  $\leq 2.5 \mu\text{m}$  in diameter (Butwin et al., 2020). Iterative calculations were performed to calculate  $D_g$  using Eqs. (A1-3).

For the GRIMM optical diameters  $D_{op}$ , we used the correction factors calculated by Formenti et al. (2021) by applying the Mie theory and assuming that the particles are homogeneous spheres. The  $D_{op}/D_g$  range measured by the GRIMM is from  $0.25 \mu\text{m}$  to  $32 \mu\text{m}$  assuming the complex refractive index of polystyrene latex sphere (PSL) reference particles ( $n = 1.59$  and  $k = 0.000$ ). For the Icelandic dust samples, we selected a range of potential complex refractive indices with the index  $n$  varying from 1.57 to 1.63 by 0.01 steps and the index  $k$  varying from 0.000 to 0.020 by 0.001 steps. The range of  $n$  and  $k$  values was chosen according to the initial guess of the complex refractive index of the Icelandic dust samples examined in this study, estimated on the basis of their mineralogical composition (Baldo et al., 2020).

*Merging of the SMPS and GRIMM size distributions:* The SMPS geometrical size distributions, calculated assuming  $\chi$  from 1.6 to 2.0 by 0.1 steps, and the GRIMM geometrical size distributions, calculated assuming 147 potential complex refractive indices ( $n$  from 1.57 to 1.63 by 0.01 steps, and  $k$  from 0.000 to 0.020 by 0.001 steps), were combined in 735 different geometrical size distributions  $dN(D_g)/d\log D_g$ . Figure 2 shows the good agreement between SMPS and GRIMM in their size overlapping region. Since the size resolution of the SMPS measurements is higher than that of the GRIMM measurements, we assumed the SMPS size distribution as a better representative in the overlapping region.

The merged geometrical size distributions were interpolated at a constant  $d\log D_g$  interval of  $1/64$  using a fitted cubic smoothing spline and normalized so that the total number of particles before and after interpolation is conserved. The `smooth.spline` function from core CRAN R software was used to perform the interpolation following Beddows et al. (2010). The total number of particles was calculated as follows:

$$N_{\text{tot}}(D_g) = \int_{D_{g,\text{min}}}^{D_{g,\text{max}}} \frac{dN(D_g)}{d\log D_g} \cdot d\log D_g \quad (1)$$

After interpolation, the merged geometrical size distributions were corrected for the particle loss along the instrument sampling lines. The particle loss calculator (PLC) software (von der Weiden et al., 2009) was used to calculate the particle

195 loss percentage for each instrument. The details of the method used in the particle loss correction process are described in Appendix B. The geometrical size distributions were first corrected for particle loss along the sampling lines of SMPS and GRIMM to obtain the size distributions of the particles suspended in CESAM. Subsequently, the particle loss correction was applied to the size distribution of particles within the chamber to determine the size distributions of particles sampled by the aethalometer and nephelometer. Furthermore, since the size distributions of the two instruments were similar (the percentage  
200 difference between the loss correction estimated for the aethalometer and that obtained for the nephelometer was  $< 11\%$ ), a common size distribution was calculated as the mean of the size distributions of the aethalometer and nephelometer. The resulting size distribution, referred to as  $N_{\text{sw-OPAs}}(D_g)$ , represents the size distribution detected by the Shortwave Optical Properties Analyzers (SW-OPAs). The size distribution of particles sampled by the SW-OPAs is more representative of transported particles than of local aerosols near the source region, due to the loss of larger particles through the nephelometer and aethalometer sampling lines (Di Biagio et al., 2019; Di Biagio et al., 2017a). In agreement with the size analysis of Di  
205 Biagio et al. (2017a) and Di Biagio et al. (2019), the results of this study would be more suitable for understanding the radiative effects of Icelandic dust in regions affected by its long-range transport.

Based on the combinations of  $\chi$ ,  $n$ , and  $k$  size corrections, a total of 735 geometrical size distributions were calculated. These were used to correct the spectral coefficients  $\beta_{\text{abs}}$  and  $\beta_{\text{sca}}$  measured respectively by the aethalometer and nephelometer, and  
210 to retrieve the complex refractive index of Icelandic dust. An example of the procedure used to process the size distribution data is shown in Fig. S2.

Furthermore, two sensitivity simulations were performed to account for the potential error on measured particle number concentrations –  $\sqrt{N}(D_g)$  according to the Poisson statistics – in the retrieval of the complex refractive index. The observed particle number concentrations plus or minus 1 standard deviation (SD),  $N(D_g) \pm \sqrt{N}(D_g)$ , were used as upper and lower  
215 limits of the SMPS and GRIMM measurements. In Test 1, corrections and calculations were performed using the SMPS and GRIMM data plus 1 SD uncertainty. In Test 2, we used the SMPS and GRIMM data minus 1 SD uncertainty.

Further uncertainty arises from the use of the Mie theory, which assumes homogeneous spherical particles. The particle morphology was not measured during the experiments, but previous research found that Icelandic dust particles can be irregular and present internal pores (Butwin et al., 2020; Richards-Thomas et al., 2020). In particular, the results of Butwin et  
220 al. (2020) indicate that the morphological characteristics of Icelandic dust particles equal or less than  $20 \mu\text{m}$  in diameter tend to be similar to those of mineral dust from global source regions, while larger particles exhibited morphological properties of volcanic ash. In this study, since the particle size distribution effectively measured by the GRIMM corresponds to around the  $\text{PM}_{20}$  fraction, we can assume that Icelandic dust has similar morphological properties to typical mineral dust. Despite mineral dust largely exhibits irregular shapes (Huang et al., 2021), our calculations are based on homogeneous equivalent  
225 spheres. However, previous studies have shown that not considering the asphericity of dust aerosols when converting  $D_{\text{op}}$  into  $D_g$  can lead to an underestimation of the coarse sizes (Formenti et al., 2021; Huang et al., 2021) as aspherical particles have a larger single-particle scattering cross-section  $C_{\text{sca}}$  compared to the volume-equivalent spheres. This effect is greater as

the imaginary index  $k$  of the dust particles increases (Huang et al., 2021). The uncertainty resulting from using the Mie theory was not estimated in this study.

230

*Calculation of the effective diameter:* The effective diameter ( $D_{\text{eff}}$ ) can be used as a measure of the aerosol size distribution as defined by Eq. (2) (Hansen, 1971):

$$D_{\text{eff}} = \frac{\int_{D_{g,\min}}^{D_{g,\max}} D_g^3 \cdot \frac{dN(D_g)}{d\log D_g} \cdot d\log D_g}{\int_{D_{g,\min}}^{D_{g,\max}} D_g^2 \cdot \frac{dN(D_g)}{d\log D_g} \cdot d\log D_g} \quad (2)$$

$D_{\text{eff}}$  differs from the simple mean diameter because the particle surface area is included as a weight factor. Since light scattering is proportional to the particle surface area,  $D_{\text{eff}}$  is relevant for describing the scattering properties of size distributions (Hansen, 1971). In this study,  $D_{\text{eff}}$  was calculated for the aerosol fine fractions ( $D_g \leq 1 \mu\text{m}$ ),  $D_{\text{eff,fine}}$ , and the coarse fractions ( $D_g > 1 \mu\text{m}$ ),  $D_{\text{eff,coarse}}$ , by varying the input parameters  $\chi$ ,  $n$ , and  $k$  in the examined ranges.

235

### 2.2.2 Spectral scattering coefficient measured by the nephelometer

The aerosol scattering coefficient  $\beta_{\text{sca}}$  was measured by the nephelometer at 450, 550, and 700 nm. The nephelometer can only detect scattering angles ( $\theta$ ) from about  $7^\circ$  to  $170^\circ$  rather than from  $0^\circ$  to  $180^\circ$  causing a systematic underestimation of the total scattering known as angular truncation error (Anderson et al., 1996; Anderson and Ogren, 1998). The truncation correction  $C_{\text{trunc}}$  for total scattering was estimated from Mie calculations using  $N_{\text{sw-OPAs}}(D_g)$  as input as described in Appendix C.

240

The average of the  $C_{\text{trunc}}(\lambda)$  values obtained for different  $N_{\text{sw-OPAs}}(D_g)$  was used to correct the data. The  $C_{\text{trunc}}$  decreased with increasing  $\lambda$ , and with the dust residence time in the chamber following the decrease in the contribution of the coarse fraction to the aerosol size distribution.  $C_{\text{trunc}}(\lambda)$  was similar for different Icelandic dust samples and varied between 1.3-1.9. These results are comparable to the range of  $C_{\text{trunc}}$  values (from 1.2 to 1.7) reported by Di Biagio et al. (2019). The uncertainty on  $C_{\text{trunc}}(\lambda)$  was estimated from the relative standard deviation (RSD) of  $C_{\text{trunc}}(\lambda)$  obtained for different  $N_{\text{sw-OPAs}}(D_g)$ . The uncertainty on  $C_{\text{trunc}}(\lambda)$  was in the range 7-17%.

245

The scattering coefficient at a given  $\lambda$  was corrected by multiplying  $\beta_{\text{sca}}$  by  $C_{\text{trunc}}$ . The uncertainty on  $\beta_{\text{sca}}$  at 450, 550, and 700 nm was calculated using the error propagation method considering the photon counting and gas calibration uncertainty (5%), the SD over 12-min intervals, and the uncertainty on  $C_{\text{trunc}}$ . The uncertainty on  $\beta_{\text{sca}}$  corrected for truncation varied between 9% and 19%. Once corrected for truncation,  $\beta_{\text{sca}}$  was extrapolated at the aethalometer operating  $\lambda$  (370, 470, 520, 590, 660, 880, and 950 nm) using the scattering Ångström exponent ( $\text{\AA}_{\text{sca}}$ ) which was obtained from the power-law fitting of  $\beta_{\text{sca}}(\lambda)$  versus  $\lambda$  weighted by the inverse of the variance of  $\beta_{\text{sca}}(\lambda)$ . The uncertainty on  $\beta_{\text{sca}}$  at 370, 470, 520, 590, 660, 880, and 950 nm was estimated as the quadratic combination of the average error on  $\beta_{\text{sca}}$  corrected for truncation and the root-mean-square error (RMSE) of  $\beta_{\text{sca}}$  predicted at 450, 550, and 700 nm. The uncertainty on  $\beta_{\text{sca}}$  predicted was 11%-18%.

255



### 2.2.3 Spectral absorption coefficient measured by the aethalometer

The aerosol absorption coefficient  $\beta_{\text{abs}}$  was retrieved from the aethalometer measurements at 370, 470, 520, 590, 660, 880, and 950 nm. The aethalometer measures the spectral attenuation (ATTN) of light passing through an aerosol-laden quartz filter, which is then converted into the attenuation coefficient (e.g., Weingartner et al., 2003):

$$\text{ATTN} = \ln \frac{I_0}{I} \quad (3)$$

where  $I_0$  is the intensity of light passing through the blank portion of the filter, and  $I$  is the intensity of light passing through the loaded filter. From the measured ATTN the attenuation coefficient ( $\beta_{\text{ATTN}}$ ,  $\text{m}^{-1}$ ) is retrieved as:

$$\beta_{\text{ATTN}}(\lambda) = \frac{\Delta \text{ATTN}(\lambda)}{\Delta t} \cdot \frac{A}{V} \quad (4)$$

where  $A$  is the area of the aerosol collection spot ( $0.00005 \pm 0.00001 \text{ m}^2$ ),  $V$  is the sampling flow rate ( $0.002 \text{ m}^3 \text{ min}^{-1}$ ), and  $\Delta \text{ATTN}$  ( $\text{ATTN}_n - \text{ATTN}_{n-1}$ ) is the variation of ATTN over 2-min time intervals ( $\Delta t$ ) that corresponds to the time resolution of the aethalometer.

The spectral coefficient  $\beta_{\text{ATTN}}$  contains the contributions of light absorption by the aerosol particles deposited on the filter and is affected by scattering by the filter material itself. The aerosol absorption coefficient  $\beta_{\text{abs}}$  was obtained by correcting  $\beta_{\text{ATTN}}$  for measurements artefacts according to the correction scheme from Collaud Coen et al. (2010):

$$\beta_{\text{abs}}(\lambda) = \frac{\beta_{\text{ATTN}}(\lambda) - \alpha(\lambda) \cdot \beta_{\text{sca}}(\lambda)}{R(\lambda) \cdot C_{\text{ref}}(\lambda)} \quad (5)$$

where  $\alpha(\lambda)\beta_{\text{sca}}(\lambda)$  is the scattering effect correction.  $\beta_{\text{sca}}(\lambda)$  is the scattering coefficient weighted by the  $\alpha(\lambda)$  parameter. This term represents the scattered radiation by the aerosol particles deposited on the filter and miscounted as attenuation.  $C_{\text{ref}}(\lambda)$  is the multiple scattering effect correction, representing multiple scattering by the filter fibers.  $R(\lambda)$  is the loading effect correction, which accounts for the reduced aethalometer response with time due to light-absorbing particles accumulating on the filter. The details of the calculations of the parameters in Eq. (5) are described in Appendix D.

The uncertainty on  $\beta_{\text{abs}}(\lambda)$  corrected according to Eq. (5) was calculated using the error propagation method considering the uncertainty on  $\alpha(\lambda)$ ,  $\beta_{\text{sca}}(\lambda)$ ,  $C_{\text{ref}}$ , and  $R(\lambda)$ . The uncertainty on  $\beta_{\text{abs}}(\lambda)$  as 2-min intervals varied between 25%-78%. Ultimately,  $\beta_{\text{abs}}(\lambda)$  was averaged over 12-min intervals, and the final uncertainty was calculated as the quadratic combination of SD and the average systematic error over 12-min intervals. The uncertainty on  $\beta_{\text{abs}}(\lambda)$  as 12-min intervals varied between 27%-75%.

## 2.3 Derivation of the aerosol optical properties

### 2.3.1 Calculation of the aerosol spectral single scattering albedo

The spectral coefficients  $\beta_{\text{sca}}(\lambda)$  and  $\beta_{\text{abs}}(\lambda)$  obtained from the nephelometer and aethalometer measurements were used to calculate 12-min average values of the extinction coefficient  $\beta_{\text{ext}}(\lambda)$  in  $\text{Mm}^{-1}$  at 370, 470, 520, 590, 660, 880, and 950 nm:

$$\beta_{\text{ext}}(\lambda) = \beta_{\text{sca}}(\lambda) + \beta_{\text{abs}}(\lambda) \quad (6)$$

SSA( $\lambda$ ) was also calculated at a 12-min resolution:

$$\text{SSA}(\lambda) = \frac{\beta_{\text{sca}}(\lambda)}{\beta_{\text{sca}}(\lambda) + \beta_{\text{abs}}(\lambda)} \quad (7)$$

The uncertainty on  $\beta_{\text{ext}}(\lambda)$  was 10%-17%, while the uncertainty on SSA( $\lambda$ ) varied between 15%-25%, which were estimated by the error propagation through Eq. (6) and Eq. (7), respectively, considering the uncertainty on  $\beta_{\text{sca}}(\lambda)$  and  $\beta_{\text{abs}}(\lambda)$ .

In addition, the experiment-averaged single scattering albedo  $\text{SSA}_{\text{avg}}$  at 370, 470, 520, 590, 660, 880, and 950 nm was retrieved from the slope ( $m_{\text{RMA}}$ ) of the linear regression between  $\beta_{\text{sca}}(\lambda)$  and  $\beta_{\text{abs}}(\lambda)$  starting from 30 min after the dust injection peak to 2.5 h (Di Biagio et al., 2019; Moosmuller et al., 2012):

$$\text{SSA}_{\text{avg}}(\lambda) = \left(1 + \frac{1}{m_{\text{RMA}}(\lambda)}\right)^{-1} \quad (8)$$

The linear fitting was performed using the reduced major axis (RMA) regression, because both variables (x and y) come from measurements and are subject to errors (Ayers, 2001; Smith, 2009). Overall, a strong correlation between  $\beta_{\text{sca}}(\lambda)$  and  $\beta_{\text{abs}}(\lambda)$  was observed ( $R^2 > 0.99$ ). The uncertainty on  $\text{SSA}_{\text{avg}}(\lambda)$  calculated considering the error on  $m_{\text{RMA}}$  in Eq. (8) was  $\leq 8\%$ .

### 2.3.2 Retrieval of the spectral complex refractive index

The spectral coefficients  $\beta_{\text{abs}}(\lambda)$  and  $\beta_{\text{sca}}(\lambda)$  were computed using the SW-OPA geometrical size distributions  $N_{\text{sw-OPAs}}(D_g)$  for the 735 different combinations of the input parameters (X, n, and k), and hereinafter referred to as  $\beta_{\text{abs,model}}(\lambda)$  and  $\beta_{\text{sca,model}}(\lambda)$ , respectively (see Appendix C). Subsequently, the complex refractive index at 370, 470, 520, 590, 660, 880, and 950 nm was determined through the comparison between  $\beta_{\text{abs,model}}(\lambda)$  and  $\beta_{\text{sca,model}}(\lambda)$  and those measured by the nephelometer and aethalometer referred to as  $\beta_{\text{abs,meas}}(\lambda)$  and  $\beta_{\text{sca,meas}}(\lambda)$ , at individual time points throughout the time series. For each  $\beta_{\text{abs,model}}(\lambda)$  and  $\beta_{\text{sca,model}}(\lambda)$  scenario, we estimated the model error as the percentage difference (%diff) respectively with  $\beta_{\text{abs,meas}}(\lambda)$  and  $\beta_{\text{sca,meas}}(\lambda)$ . To account for the uncertainty on  $\beta_{\text{abs,meas}}(\lambda)$  and  $\beta_{\text{sca,meas}}(\lambda)$ , we used  $\beta_{\text{abs,meas}}(\lambda) \pm 1 \text{ SD}$  and

$\beta_{\text{sca,meas}}(\lambda) \pm 1$  SD as the upper and lower limits of the measurements. We selected  $\beta_{\text{abs,model}}(\lambda)$  and  $\beta_{\text{sca,model}}(\lambda)$  scenarios with the lowest model error corresponding to 0.1-0.5 quantile of %diff. Subsequently, the results for  $\beta_{\text{abs,meas}}(\lambda) \pm 1$  SD and  $\beta_{\text{sca,meas}}(\lambda) \pm 1$  SD (consisting of three datasets for each spectral optical coefficient) were examined, and only the  $\beta_{\text{abs,model}}(\lambda)$  and  $\beta_{\text{sca,model}}(\lambda)$  scenarios common to all six datasets were kept. The indices  $k(\lambda)$  and  $n(\lambda)$  were retrieved from the selected scenarios. Since the comparison between calculation and measurements resulted in multiple solutions for  $k(\lambda)$  and  $n(\lambda)$ , we calculated the mean of the  $k$  solutions and of the  $n$  solutions.

In addition, the experiment-averaged spectral complex refractive index was determined based on the analysis of the linear correlation between  $\beta_{\text{abs,meas}}(\lambda)$  vs  $\beta_{\text{abs,model}}(\lambda)$ , and  $\beta_{\text{sca,meas}}(\lambda)$  vs  $\beta_{\text{sca,model}}(\lambda)$  scenarios starting from 30 min after the dust injection peak to 2.5 h. To retrieve the experiment-averaged real index  $n_{\text{avg}}(\lambda)$  and imaginary index  $k_{\text{avg}}(\lambda)$ , we updated the method applied to determine  $k(\lambda)$  and  $n(\lambda)$  at a 12-min resolution, as described in the preceding paragraph. For each  $\beta_{\text{abs,model}}(\lambda)$  and  $\beta_{\text{sca,model}}(\lambda)$  scenarios, we assumed that the input parameters  $\chi$ ,  $k(\lambda)$  and  $n(\lambda)$  are constant throughout the experimental run. From the analysis of the linear correlation between  $\beta_{\text{abs,meas}}(\lambda) \pm 1$  SD and  $\beta_{\text{abs,model}}(\lambda)$  scenarios, and between  $\beta_{\text{sca,meas}}(\lambda) \pm 1$  SD and  $\beta_{\text{sca,model}}(\lambda)$  scenarios, we selected only the model estimates which showed a high correlation with observations ( $R^2 > 0.70$ ). The modelled and measured spectral coefficients were then compared based on the root mean square error (RMSE) instead of using the %diff at individual time points. The uncertainty on  $k_{\text{avg}}(\lambda)$  and  $n_{\text{avg}}(\lambda)$  was estimated from the RSD of the  $k$  and  $n$  solutions. The uncertainty on  $k_{\text{avg}}(\lambda)$  was up to 99%, while the uncertainty on  $n_{\text{avg}}(\lambda)$  was  $< 2\%$ .

These calculations were repeated in Test 1 and Test 2, the sensitivity simulations to account for the uncertainty resulting from the size distribution measurements which were used to correct the spectral coefficients measured by the aethalometer and nephelometer, and to calculate the  $\beta_{\text{abs,model}}(\lambda)$  and  $\beta_{\text{sca,model}}(\lambda)$  scenarios.

### 3 Results

#### 3.1 Aerosol size distribution and effective diameter

Figure 2 shows examples of geometrical size distributions that were obtained using various dynamic shape factors  $\chi$  and complex refractive indices for the dust particles. For spherical particles ( $\chi = 1$  and  $D_m = D_g$ ), the  $D_g$  range measured by the SMPS was from 0.019  $\mu\text{m}$  to 0.87  $\mu\text{m}$ . For non-spherical particles ( $\chi > 1$ ),  $D_g$  decreased with increasing  $\chi$  as defined by Eq. (A1). After the conversion of  $D_m$  to  $D_g$ , the  $D_g$  range of the SMPS was around 0.015-0.58  $\mu\text{m}$  for  $\chi = 1.6$ , 0.015-0.56  $\mu\text{m}$  for  $\chi = 1.7$ , 0.014-0.53  $\mu\text{m}$  for  $\chi = 1.8$ , 0.014-0.51  $\mu\text{m}$  for  $\chi = 1.9$ , and 0.013-0.49  $\mu\text{m}$  for  $\chi = 2.0$ .

The correction factors used to convert  $D_{\text{op}}$  into  $D_g$  had a considerable impact on  $D_g > 0.6 \mu\text{m}$ .  $D_g$  increased with  $k$ , while the variability of  $n$  mainly affected  $D_g$  at around 1  $\mu\text{m}$ . The minimum  $D_g$  varied between around 0.24-0.25  $\mu\text{m}$ . For  $k \leq 0.003$ , the maximum  $D_g$  sharply increased with  $k$  from 31.5-32.5  $\mu\text{m}$  to 85.6-90.4  $\mu\text{m}$ . For  $k > 0.003$ , the maximum  $D_g$  showed only a small variation reaching up to around 94  $\mu\text{m}$  (Fig. S3).

The overlapping interval between the SMPS and GRIMM data decreased as  $\chi$  increased. The GRIMM size distributions tend to spread out between 0.6-2  $\mu\text{m}$  (Fig. 2), consequently reducing the smoothness of the fitted size distributions within this size range, which is likely due to the larger uncertainty of the correction factors at these  $D_g$  values (Formenti et al., 2021). After  
335 correcting the data for the loss along the instrument sampling lines, the size distribution of the particles suspended in CESAM was in the  $D_g$  range up to 20  $\mu\text{m}$ , while the size distribution of the particles sampled by the SW-OPAs was in the  $D_g$  range up to 9  $\mu\text{m}$ .

The average of the  $D_{\text{eff}}$  values derived from different  $\chi$ -n-k combinations was reported as the  $D_{\text{eff}}$  of the Icelandic dust samples, with the RSD as the uncertainty on  $D_{\text{eff}}$ . The effective diameter of the coarse fractions,  $D_{\text{eff,coarse}}$ , showed a strong  
340 positive correlation with the input parameter k ( $R^2 = 0.6$ - $0.8$ ), while the correlation between  $D_{\text{eff,coarse}}$  and  $\chi$  or n was low ( $R^2 < 0.1$ ). The correlation between the effective diameter of the fine fractions,  $D_{\text{eff,fine}}$ , and all input parameters was low. Figure S4 shows an example of the correlation between  $D_{\text{eff}}$  determined from the SW-OPA size distributions and  $\chi$ , n, and k. The parameter  $D_{\text{eff}}$  was used as an indication of the stability of the size distribution in the chamber. Figure 3 shows an example of how  $D_{\text{eff}}$  changed over time. The  $D_{\text{eff,coarse}}$  decreased with time due to the rapid deposition of the largest particles  
345 in the chamber. In Icelandic dust samples,  $D_{\text{eff,coarse}}$  calculated using the SW-OPA size distributions varied from  $\sim 3.1$ - $3.5 \mu\text{m}$  after around 30 min from the injection peak to  $2.5$ - $2.7 \mu\text{m}$  after around 2.5 h from the injection peak. For the particles suspended in CESAM,  $D_{\text{eff,coarse}}$  varied from  $8.4$ - $11 \mu\text{m}$  (30 min after the injection peak) to  $3.7$ - $4.4 \mu\text{m}$  (2.5 h after the injection peak). The  $D_{\text{eff,fine}}$  remained relatively constant over the duration of the experiments varying between  $0.5$ - $0.7 \mu\text{m}$  in different samples. The uncertainty on  $D_{\text{eff,coarse}}$  was  $< 14\%$ - $19\%$  for the SW-OPA size distributions, and  $44\%$ - $56\%$  for the  
350 particles suspended in CESAM, while the uncertainty on  $D_{\text{eff,fine}}$  was less than  $3\%$ . The  $D_{\text{eff}}$  values for all the samples are shown in Fig. 3, and Figs. S5-9.

The  $D_{\text{eff}}$  results from the sensitivity studies (Tests 1-2) to account for the error on the SMPS and GRIMM measurements were consistent within the uncertainties with the results from the base simulation (Fig. 3 and Figs. S5-9). The difference between the  $D_{\text{eff}}$  results from the sensitivity studies and  $D_{\text{eff}}$  from the base simulation was not significant because it was less  
355 than three times the square root of the sum of their squared uncertainties. For the SW-OPA size distributions,  $D_{\text{eff,coarse}}$  decreased from  $\sim 3.3$ - $3.6 \mu\text{m}$  (30 min after the injection peak) to  $2.8$ - $3.0 \mu\text{m}$  (2.5 h min after the injection peak) in Test 1, and from  $2.8$ - $3.3$  (30 min after the injection peak) to  $1.4$ - $1.9$  (2.5 h min after the injection peak) in Test 2. For the particles suspended in CESAM,  $D_{\text{eff,coarse}}$  decreased from  $\sim 10$ - $12 \mu\text{m}$  (30 min after the injection peak) to  $4.6$ - $5.7 \mu\text{m}$  (2.5 h min after the injection peak) in Test 1, and from  $3.3$ - $5.8$  (30 min after the injection peak) to  $1.6$ - $2.0$  (2.5 h min after the injection peak)  
360 in Test 2. The uncertainty on  $D_{\text{eff,coarse}}$  was  $12\%$ - $17\%$  in Test 1 and  $9\%$ - $22\%$  in Test 2 for the SW-OPA size distributions, and  $41\%$ - $55\%$  in Test 1 and  $9\%$ - $50\%$  in Test 2 for the particles suspended in CESAM. The results of  $D_{\text{eff,fine}}$  were not considerably affected by the sensitivity studies.

### 3.2 Spectral extinction and absorption coefficients, single scattering albedo, and complex refractive index

Figure 4 presents the temporal variation of  $\beta_{\text{ext}}(\lambda)$ ,  $\beta_{\text{abs}}(\lambda)$ , and  $\text{SSA}(\lambda)$  of the Icelandic dust samples, as derived from the base simulation. The coefficients  $\beta_{\text{ext}}(\lambda)$  and  $\beta_{\text{abs}}(\lambda)$  decreased with  $\lambda$ , with the largest variation observed for  $\beta_{\text{abs}}(\lambda)$  between 370 and 590 nm. On the other hand,  $\text{SSA}(\lambda)$  increased marginally with  $\lambda$ , within the error bars. Both  $\beta_{\text{ext}}(\lambda)$  and  $\beta_{\text{abs}}(\lambda)$  decreased with time at all values of  $\lambda$ , while  $\text{SSA}(\lambda)$  was relatively constant. The results of the sensitivity studies for  $\beta_{\text{ext}}(\lambda)$ ,  $\beta_{\text{abs}}(\lambda)$ , and  $\text{SSA}(\lambda)$  were in agreement with the results of the base simulation within their respective uncertainties (Figs. S10-14).

Figure 5 provides an example of the temporal variation of  $n(\lambda)$  and  $k(\lambda)$  in two Icelandic dust samples, including the results from both the base simulation and the sensitivity studies. The parameter  $n(\lambda)$  showed no dependence on  $\lambda$  and time and varied within the range of examined  $n$  values (from 1.57 to 1.63 by 0.01 steps) with an uncertainty of  $< 3\%$  in the base simulation, Test 1, and Test 2 (Figs. S15-19). For the parameter  $k(\lambda)$ , the data at 12-min resolution were too noisy to identify a clear relationship with time or  $\lambda$  (Figs. S15-19). In Test 1,  $k(\lambda)$  was almost constant with time and  $\lambda$ . In Test 2,  $k$  did not show a clear dependence on  $\lambda$ . For the sample MIR45,  $k(\lambda)$  exhibited an increasing trend over time. For D3 and Maeli2,  $k(\lambda)$  increased after around 1.5 h from the dust injection peak. For H55 and Land1,  $k(\lambda)$  did not show a temporal trend. The uncertainty on  $k(\lambda)$  at 12-min resolution varied up to 99% in the base simulation, 198% in Test 1, and 79% in Test 2. We compared  $k(\lambda)$  and  $D_{\text{eff,coarse}}$  to determine the impact of the reduction in the particle coarse fraction over time on the complex refractive index. However, a clear correlation between  $D_{\text{eff,coarse}}$  and  $k(\lambda)$  was not observed. The  $k(\lambda)$  results at a 12-min resolution were noisy, making it challenging to establish a meaningful relationship between the two parameters, as the slopes of the regression lines between  $D_{\text{eff,coarse}}$  and  $k(\lambda)$  were close to zero.

The experiment-averaged single scattering albedo  $\text{SSA}_{\text{avg}}$  increased with  $\lambda$  from 370 to 590 nm, though with some uncertainty, but remained relatively constant between 590 and 950 nm (Fig. 6). The  $\text{SSA}_{\text{avg}}(\lambda)$  varied from 0.93 at 370 nm to 0.96 at 950 nm for the sample D3, from 0.94 to 0.96 for H55, from 0.91 to 0.96 for Land 1, from 0.90 to 0.95 for Maeli2, and from 0.90 to 0.94 for MIR45 (Table 1). The uncertainty on  $\text{SSA}_{\text{avg}}(\lambda)$  was  $\leq 8\%$ . The  $\text{SSA}_{\text{avg}}(\lambda)$  results from the sensitivity studies were within the uncertainty for the base simulation (Table S2).

The experiment-averaged imaginary index,  $k_{\text{avg}}(\lambda)$ , decreased with  $\lambda$ , from 0.006 at 370 nm to 0.002 at 950 nm for the sample D3, from 0.005 to 0.003 for H55, Land1, and Maeli2, and from 0.005 to 0.003 for MIR45, as determined from the base simulation (Table S3).  $k_{\text{avg}}(\lambda)$  was sensitive to the particle size distribution. The variation of  $k_{\text{avg}}$  with respect to  $\lambda$  was less evident in Test 1 compared to the base simulation (Table S3). The  $k_{\text{avg}}(\lambda)$  values varied between 0.003 and 0.001 for H55 and Land1, while they remained relatively constant at around 0.002 for Maeli2 and MIR45. For the sample D3,  $k_{\text{avg}}(\lambda)$  was 0.002 at 370 nm and around 0.001 for all the other  $\lambda$ . In Test 2,  $k_{\text{avg}}$  increased with  $\lambda$  from 0.002 at 370 nm to 0.007 at 950 nm for the sample H55. For all the other Icelandic dust samples, the relationship between  $k_{\text{avg}}$  and  $\lambda$  was not monotonic, and  $k_{\text{avg}}$  varied in the range between 0.002-0.007 for D3, 0.003-0.006 for Land1, 0.002-0.005 for Maeli2, and 0.005-0.007 for MIR45. Overall,  $k_{\text{avg}}(\lambda)$  estimates of Test 1 were lower than those of the base simulation and Test 2 (Table S3). The uncertainty on  $k_{\text{avg}}(\lambda)$  was generally  $< 85\%$  in the base simulation and Test 1, and  $< 70\%$  in Test 2.

The experiment-averaged real index,  $n_{\text{avg}}(\lambda)$ , showed no trend with  $\lambda$ , and  $n_{\text{avg}}(\lambda)$  varied in the range between 1.59-1.62 in the base simulation, 1.58-1.61 in Test 1, and 1.57-1.63 in Test 2 (Table S4). Overall, the uncertainty on  $n_{\text{avg}}(\lambda)$  was  $< 2\%$ .

Table S5 reports a summary of the comparison between  $SSA_{\text{avg}}(\lambda)$  obtained from the measured and computed spectral coefficients for the base simulation, Test 1, and Test 2. Although the RMSE values were generally low, the correlation between the measured and modelled  $SSA_{\text{avg}}(\lambda)$  tends to be higher in the base simulation and Test 1 compared to Test 2 (Table S5). Di Biagio et al. (2019) chose to average the  $k$  values from all three scenarios. Here the increase of  $k_{\text{avg}}$  with  $\lambda$  in Test 2 is hard to explain, suggesting that Test 2 results are not realistic. Based on this, we chose to combine the results from the base simulation and Test 1 to obtain a single set of values for  $k(\lambda)$  (Table 2) and  $n(\lambda)$  (Table 3). While the results obtained from averaging the baseline simulation and Test 1 may have a bias towards high particle number concentration, no significant difference was observed when the results of Test 2 were also included, for both the parameters  $n$  and  $k$ . Although the mean values of  $k$  tended to be higher when Test 2 was included in the average, particularly in the wavelength range of 660-950 nm, the difference between the average with Test 2 included and the average without Test 2 is less than three times the square root of the sum of their squared uncertainties. For these reasons, it was decided to use the more conservative approach.

## 4 Discussion

The Icelandic dust samples examined in this study showed similar spectral optical properties. The spectral single scattering albedo  $SSA(\lambda)$  (Table 1) and imaginary index  $k(\lambda)$  (Table 2) showed opposite trends, as expected.  $SSA(\lambda)$  increased from 0.90-0.94 at 370 nm to 0.94-0.96 at 950 nm in different samples, while  $k(\lambda)$  decreased slightly from 0.004 at 370 nm to 0.002-0.003 at 950 nm. The real index  $n(\lambda)$  was rather constant with  $\lambda$  at 1.60-1.61 (Table 3).

### 4.1 Spectral complex refractive index and single scattering albedo of Icelandic dust comparison with literature data, volcanic ash, and low-latitude dust

The complex refractive index of Icelandic dust is consistent with the initial estimate of Baldo et al. (2020) based on the volume-averaged mineralogical composition and reference complex refractive indices of individual minerals. In Baldo et al. (2020),  $n(\lambda)$  ranged from 1.57-1.63 between 470 and 660 nm, while  $k(\lambda)$  was sensitive to the reference refractive indices of magnetite applied in the calculations, and varied up to one order of magnitude in different absorption scenarios. In this study,  $n(\lambda)$  (Table 3) is comparable to the average value of the initial estimate, and  $k(\lambda)$  (Table 2 and Fig. 7) is within or at the lower end of the range of  $k(\lambda)$  values estimated by Baldo et al. (2020).

The value of  $k$  at 660 nm of the Icelandic dust sample from the Mýrdalssandur hotspot (MIR45, southern Iceland) is lower than that reported in the literature by Zubko et al. (2019). In Zubko et al. (2019), the complex refractive index of Icelandic dust was retrieved based on the measurements of the angular scattered-light intensity and degree of linear polarization of aerosol particles generated from volcanic sand from the Mýrdalssandur area. They estimated a value of 1.60 for  $n$  and 0.01

for  $k$  at 647 nm. (Zubko et al., 2019), the latter is around 3 times higher than the value observed in this study at 660 nm (Table 2).

Icelandic dust shares a similar origin with volcanic ash; however, it is difficult to assess the complex refractive index of Icelandic dust based on what has been observed for Icelandic volcanic ash. This is because the literature reports a wide range of values for  $k$  and  $n$  in the SW-spectrum for Icelandic ash (Fig. 8). The values of  $n(\lambda)$  for Icelandic ash vary from 1.50 to 1.65 (Bukowiecki et al., 2011; Deguine et al., 2020; Derimian et al., 2012; Reed et al., 2018; Weinzierl et al., 2012). Laboratory studies conducted on re-suspended ash samples found that  $k(\lambda)$  of Icelandic ash remains below 0.001 in the UV-visible spectrum and tend to increase towards the near-IR region (Deguine et al., 2020; Reed et al., 2018). The  $k(\lambda)$  values resulting from direct measurements of the Eyjafjallajökull plume in 2010 (e.g., Bukowiecki et al., 2011; Derimian et al., 2012; Weinzierl et al., 2012) are considerably different from the results of laboratory measurements (Deguine et al., 2020; Reed et al., 2018). In Derimian et al. (2012), the range of  $k(\lambda)$  values for Eyjafjallajökull ash varied from around 0.015 at 440 nm to 0.008 at 870 nm, which is similar to what reported by Weinzierl et al. (2012), whereas Bukowiecki et al. (2011) reported an average value of 0.02 between 450-700 nm. Here,  $n(\lambda)$  of Icelandic dust (Table 3) falls within the range of  $n$  values reported for Icelandic ash (Fig. 8a) (Bukowiecki et al., 2011; Deguine et al., 2020; Derimian et al., 2012; Reed et al., 2018; Weinzierl et al., 2012), while the values of  $k(\lambda)$  of Icelandic dust (Table 2) are higher than the  $k(\lambda)$  values of Icelandic ash from laboratory measurements (Deguine et al., 2020; Reed et al., 2018), but lower than ambient observations (Bukowiecki et al., 2011; Derimian et al., 2012; Weinzierl et al., 2012) (Fig. 8b). The difference between observational data and laboratory measurements could be due to several factors, including, for example, the different methods used, and the different type of samples analyzed, as the physical and chemical properties of re-suspended volcanic ash deposited in situ could be different from those of volcanic ash plumes, which could change further during atmospheric transport. On the other hand, our results for the SSA( $\lambda$ ) of Icelandic dust (Table 1 and Fig. 6) are within the range of values observed in Icelandic ash plumes (e.g., Derimian et al., 2012; Toledano et al., 2012). During the 2010 eruptions of Eyjafjallajökull, the values of SSA( $\lambda$ ) of a volcanic ash plume over France (17 April 2010) varied from 0.81 at 440 nm to 0.92 at 870 nm from Aerosol Robotic Network (AERONET) observations (Derimian et al., 2012), whereas SSA( $\lambda$ ) derived from AERONET measurements of the ash plumes over Spain and Portugal (6-12 May 2010) was found to be 0.95 at 440 nm but slightly decreased with  $\lambda$  (Toledano et al., 2012).

In Fig. 9, the complex refractive index and SSA( $\lambda$ ) of Icelandic dust were compared with those of low-latitude dust from major source regions such as northern Africa-Sahara, Sahel, and Eastern Asia analysed by Di Biagio et al. (2019), where the same method as in this study is applied. Therefore, the differences between Icelandic dust and low-latitude dust that we discuss are primarily due to the different physical and chemical properties of the different type of samples. The real index  $n(\lambda)$  of Icelandic dust, at around 1.6 (Table 3), is slightly higher than that of typical low-latitude dust, averaging 1.51 in the spectral range 370-950 nm (Di Biagio et al., 2019) (Fig. 9a). Fig. 9b shows that the mean value of  $k(\lambda)$  of Icelandic dust (Table 2) is at the upper end of the range of typical low-latitude dust (Di Biagio et al., 2019). The values of  $k(\lambda)$  decreased from around 0.002-0.005 at 370 nm to 0.001 at 950 nm in northern African-Saharan dust, from 0.001-0.009 at 370 nm to

0.0003-0.002 at 950 nm in Sahelian dust, and from 0.002-0.004 at 370 nm to 0.0005-0.001 at 950 nm in eastern Asian dust (Di Biagio et al., 2019). In the spectral range between 370 and 590 nm, the difference of  $k(\lambda)$  of Icelandic dust (Table 2) with the  $k(\lambda)$  values of northern African-Saharan dust, Sahelian dust, and eastern Asian dust was less than three times the square root of the sum of their squared uncertainties. The largest difference between Icelandic dust and low-latitude dust samples was observed from 660 to 950 nm (Fig. 9b). In particular, at these  $\lambda$ , the values of  $k(\lambda)$  of the samples D3 and MIR45, representative of the two major dust hotspots in Iceland (e.g., Arnalds et al., 2016), showed a significant difference compared to several low-latitude dust, and were around 2-5 times higher than the  $k(\lambda)$  values of mineral dust from Morocco, Libya, and Algeria in northern Africa-Sahara, and the Taklimakan Desert in eastern Asia, and 5-8 times higher than the  $k(\lambda)$  values of dust from the Bodele Depression in the Sahel (Fig. 9b).

Although  $SSA(\lambda)$  of Icelandic dust (Table 1 and Fig. 6) is within the range of values of low-latitude dust (Di Biagio et al., 2019), an absorption of 4%-6% can still be observed at 950 nm (Fig. 9c), which is relatively high, despite the uncertainty, considering that absorption in mineral dust at these  $\lambda$  is generally around 2% (Di Biagio et al., 2019). The SSA is affected by particle size and chemical composition. The particle size of Icelandic dust is quite comparable to that of low-latitude dust (Di Biagio et al., 2019), according to the  $D_{\text{eff}}$  values for both the coarse and fine fractions of the particles sampled by the SW-OPAs. This suggests that the differences between the two may be mainly related to mineralogy.

Di Biagio et al. (2019) reported that there is a large variability in the SW imaginary index of mineral dust in the literature. The data from their study fall within the lower range of values reported in previous studies, which could be attributed to different type of samples analysed and methodologies employed. However, the range of the real index and single scattering albedo were consistent with the literature.

When considering the imaginary index of Icelandic dust, this is close to the upper end of the imaginary index of northern African and eastern Asian dust estimated by Di Biagio et al. (2019) (Fig. 9b), and falls within the range of values reported in studies conducted on low-latitude dust for example by Formenti et al. (2011) (from the AMMA campaign), Müller et al. (2011) (from the SAMUM campaign), Balkanski et al. (2007) (calculated from mineralogy), Dubovik et al. (2002) (from AERONET measurements), and Sinyuk et al. (2003) (from satellite observations), Volz (1972) (from laboratory experiments), and Patterson et al. (1977) (from laboratory experiments), as shown in Figure S20.

## 4.2 Optical properties and mineralogy

The chemical and mineralogical composition of the Icelandic dust samples examined in this study was determined by Baldo et al. (2020). The Icelandic dust showed a basaltic composition, with relatively high total Fe content, ranging from around 11 wt% in the sample H55 to 13 wt% in MIR45, compared to low-latitude dust (1 wt%-8 wt%; e.g. Di Biagio et al., 2019; Jeong, 2008; Shi et al., 2011). The samples were primarily composed of amorphous basaltic materials ranging from 8 wt% in H55 to 60 wt %-90 wt % in the other Icelandic dust. MIR45 had the highest proportion of amorphous glass. The magnetite fraction (1 wt%-2 wt %) was considerably higher than in low-latitude dust (generally < 0.1 wt% or undetectable in northern African dust, and < 0.8 wt% in eastern Asian dust source regions; Jia et al., 2019; Lazaro et al., 2008; Maher et al., 2009;



495 Moskowitz et al., 2016; Song et al., 2014) contributing to 7 %-15 % of total Fe. Fe oxide minerals accounted for 9%-20% of the total Fe, while most of the iron was contained in other mineral phases (e.g., pyroxene, feldspars and olivine) and in the amorphous glass (Baldo et al., 2020).

The differences between Icelandic dust and low-latitude dust could be related to the presence of different Fe oxide minerals. Hematite ( $\text{Fe}_2\text{O}_3$ ) and goethite ( $\text{FeO}(\text{OH})$ ) are the main Fe oxide in low-latitude dust (Di Biagio et al., 2019; Formenti et al., 2014) and have specific absorption bands in the UV-visible spectrum (Caponi et al., 2017; Derimian et al., 2008; Di Biagio et al., 2019; Dubovik et al., 2002; Engelbrecht et al., 2016; Formenti et al., 2014; Lafon et al., 2006; Moosmuller et al., 2012; Redmond et al., 2010). However, we have shown that Icelandic dust is rich in magnetite (Baldo et al., 2020) which also absorbs light in the UV-visible spectrum, but has relatively high absorption even in the near-IR with  $k > 0.2$  at 950 nm (Table S6). Di Biagio et al. (2019) found a linear relationship between  $\text{SSA}(\lambda)$  and  $k(\lambda)$  and the mass concentrations of Fe oxides and total Fe. In this study, the sample size was too small to reach statistical significance, and the values of  $\text{SSA}(\lambda)$  and  $k(\lambda)$  were similar among different samples (Tables 1-2) and did not show a clear pattern. In addition, since amorphous basaltic material is dominant, but has main absorption bands in the IR region (Deguine et al., 2023; Deguine et al., 2020), we do not expect a good correlation. Overall, the values of  $k(\lambda)$  of Icelandic dust samples (Table 2) appear to be more similar to those of mineral dust from the Sahel region in Niger and Mali (Fig. 9b), which have relatively high Fe oxide content, 5.8 wt% and 3.7 wt%, respectively (Di Biagio et al., 2019). However, it is important to note that the total Fe oxide content in Icelandic dust is lower, ranging from 1.2 wt%-2.6 wt% (Baldo et al., 2020). Additionally, it is worth noting that the Bodele sample was classified as part of the Sahel in Di Biagio et al. (2019), but samples from the Bodele Depression are often dominated by diatoms, with extremely low Fe oxide content.(e.g., Bristow et al., 2009; Shi et al., 2011).

The high magnetite content may contribute to the high absorption capacity of Icelandic dust. In Fig. 7, the  $k(\lambda)$  values estimated in this study tend to be closer to the volume-averaged imaginary indices calculated by Baldo et al. (2020) assuming the spectral refractive index of magnetite from Querry (1985) than to the values predicted using the refractive index of magnetite from Huffman and Stapp (1973). This suggest that the dataset from Querry (1985) may be more representative of the complex refractive index of magnetite in Icelandic dust. Table S6 reports the reference complex refractive indices of mineral components of Icelandic dust in the spectral range 370-950 nm. The average value of  $k(\lambda)$  of magnetite between 660 and 950 nm, 0.18 estimated using data from Querry (1985) and 0.43 using data from Huffman and Stapp (1973), is greater than the reference index of any other mineral components, up to 270 times and 640 times, respectively. This suggests that magnetite may be one of the main minerals contributing to light absorption in Icelandic dust between 660 and 950 nm.

## 5 Concluding remarks: Implications for the radiative effect

Here, we provide for the first time a dataset with the spectral SW- SSA (Table 1) and complex refractive index (Tables 2-3) of mineral dust from five major dust hotspots in Iceland. The spectral imaginary index  $k(\lambda)$  of Icelandic dust (Table 2) is at

the upper end of the range of typical low-latitude dust. However, in Icelandic dust, absorption between 660 and 950 nm can be 2-8 times higher than in several low-latitude dust (Fig. 9b).

Previous research by Kylling et al. (2018) estimated that Icelandic dust produces a positive direct radiative forcing in the Arctic via dust-radiation interaction and deposition onto snow and ice. This estimation was based on the dust load estimates by Groot Zwaaftink et al. (2016) and assumed the same optical properties, in line with the dataset of Balkanski et al. (2007), for all mineral dust from different regions. The dust radiative effect is very sensitive to the complex refractive index, which is a major uncertainty (e.g., Di Biagio et al., 2020; Li et al., 2021). In the dataset used by Kylling et al. (2018),  $k(\lambda)$  was around 0.002 at 555 nm, which is comparable to the values of  $k(\lambda)$  of Icelandic dust at 590 nm observed in this study (Table 2). However,  $k(\lambda)$  in the Icelandic dust samples was found to be 0.002-0.003 in the range of 660 to 950 nm (Table 2), while the results from Balkanski et al. (2007) indicated that  $k(\lambda)$  decreases to below 0.001 in the same  $\lambda$  range. This implies that the positive direct radiative forcing of Icelandic dust in the Arctic could be stronger than previously estimated. Although the 700-1000 nm wavelength band is characterised by lower solar irradiance than the visible band, it still contributes approximately 25% of the total spectral solar irradiance (e.g., Woods et al., 2009). Therefore, the increased absorption of Icelandic dust in this range may still influence the radiative balance and regional climate. The measurements conducted at CESAM on the longwave optical properties of Icelandic dust will provide additional valuable insights into its radiative effect and contribution to Arctic warming. As glaciers retreat and expose more ice-free land as potential dust sources (Meinander et al., 2022), this effect is expected to increase. While these new results on the optical properties of Icelandic dust will contribute to reducing the uncertainty on its radiative effect, additional data on the size distribution and emissions are also crucial to advance our understanding.

## Appendix A: Retrieval of the aerosol size distribution

The aerosol size distribution was obtained by combining the measurements from SMPS and GRIMM into a single geometrical size distribution. Several corrections were applied to the SMPS and GRIMM measurements because of the differences in operating principles and diameter definitions between the instruments.

The SMPS measures the size distribution of submicron particles. The operating principle of the SMPS is the balance between the electrical force on a charged particle due to a constant electric field and the drag force experienced by that particle at terminal velocity which is used to derive the particle size (DeCarlo et al., 2004; Hinds, 1999). The relationship between the electrical mobility diameter  $D_m$  (the diameter of a single-charged spherical particle with the same terminal velocity in a constant electric field as the single-charged particle under consideration), as measured by the SMPS and the volume-equivalent diameter  $D_g$  (the diameter of a spherical particle of the same volume as the particle under consideration) is described by Eq. (A1):

$$D_g = \frac{D_m}{\chi} \cdot \frac{C_c(D_g)}{C_c(D_m)} \quad (A1)$$

where  $\chi$  is the dynamic shape factor,  $C_c$  is the Cunningham slip correction factor assuming a mean free path of 66 nm (air conditions) (Hinds, 1999):

$$D_{g-m} > 100 \text{ nm}, \quad C_c(D_{g-m}) = 1 + \frac{66}{D_{g-m}} \cdot 2.52 \quad (A2)$$

$$D_{g-m} \leq 100 \text{ nm}, \quad C_c(D_{g-m}) = 1 + \frac{66}{D_{g-m}} \cdot (2.34 + 1.05 \cdot e^{-0.39 \frac{D_{g-m}}{66}}) \quad (A3)$$

where  $D_{g-m}$  represents either the volume-equivalent diameter  $D_g$  or the electrical mobility diameter  $D_m$ .

The OPCs measure the size distribution across a wide range of sizes including submicron and supermicron particles. The OPC operating principle is the dependence of angular light scattering on particle size (Hinds, 1999; Wendisch and Brenguier, 2013). The particle size measured by OPCs is therefore referred to as optical-equivalent diameter  $D_{op}$ , that is the diameter of a spherical particle with a given complex refractive index which has the same scattering efficiency ( $Q_{sca}$ ) and cross-sectional area ( $A$ ) as the particle under consideration. In this study, the volume-equivalent diameter  $D_g$  of natural aerosols was retrieved using look up tables from Formenti et al. (2021) which convert  $D_{op}$  into  $D_g$ . These look up tables are calculated for a wide range of complex refractive indices to determine the diameter that would give the same (or closest) single-particle scattering cross-section,  $C_{sca} = Q_{sca} \times A$ , as that calculated using calibration reference PSL particles referring to  $D_{op}$ . For the GRIMM, increasing the imaginary index  $k$ , the scattering cross-section curve tends to become flat at around 1  $\mu\text{m}$  as shown in Figure 2 of Formenti et al. (2021). This leads to large uncertainty when calculating the volume-equivalent

575 diameters  $D_g$  of light-absorbing aerosols in the range between 0.6 and 2  $\mu\text{m}$  (Formenti et al., 2021). Consequently, increasing the value of  $D_{op}$  does not always correspond to increasing  $D_g$ , which may produce negative or very small  $d\log D_g = \log D_{g2} - \log D_{g1}$ , where  $D_{g1}$  and  $D_{g2}$  are the lower and upper cut-off diameters of each size range. This generates unusual spikes in the calculated geometrical size distributions  $dN(D_g)/d\log D_g$ . Two common strategies to address this issue are widening or grouping the OPC bin size (Johnson and Osborne, 2011) or fitting the corrected  $D_g$  curve (Hand and Kreidenweis, 2002; Johnson et al., 2008). Here, the corrected  $D_g$  values which return  $d\log D_g < 0.01$  were replaced with the average of the bin before and after.

## Appendix B: Particle loss correction

580 The particle loss correction accounts for the loss of particles along the instrument sampling lines. The particle loss calculation was performed using the PLC software (von der Weiden et al., 2009). The input parameters for the PLC software included the geometry of the sampling line, the sampling flow rate, the particle density, and the dynamic shape factor  $\chi$ . The particle bulk density was set to  $2.90 \pm 0.01 \text{ g cm}^{-3}$  as measured using a helium pycnometer for particles less than  $63 \text{ }\mu\text{m}$  in diameter (Baldo et al., 2020), a value that is consistent with the results of a recent study by Richards-Thomas et al. (2020).  
 585 However, from the analysis conducted on larger size fractions, less than  $125 \text{ }\mu\text{m}$  in diameter, using the water penetration method, Butwin et al. (2020) found an average density value of  $2.25 \text{ g cm}^{-3}$  for Icelandic dust. For each instrument, the particle loss percentage was calculated assuming  $\chi$  from 1.6 to 2.0 by 0.1 steps.  
 To retrieve the size distribution of the particles suspended in CESAM, the merged geometrical size distributions were divided into two size ranges which were corrected for particle loss in the sampling systems of SMPS and GRIMM,  
 590 respectively. The particle size domain was assigned based on the geometrical size distributions of GRIMM and SMPS used to merge the data:

$$\frac{dN(D_g)}{d \log D_g} = \frac{dN(D_{g,SMPS})}{d \log D_{g,SMPS}} + \frac{dN(D_{g,GRIMM})}{d \log D_{g,GRIMM}} \quad (\text{B1})$$

For the SMPS, the loss of particles with  $D_g < 1 \text{ }\mu\text{m}$  was less than 5%. For the GRIMM, the particle loss was up to 15% in the range between  $1\text{-}2.5 \text{ }\mu\text{m}$ , within 50% in the range between  $2.5\text{-}5 \text{ }\mu\text{m}$  and reached 100% at  $D_g$  around  $20 \text{ }\mu\text{m}$ . The variation in particle loss calculated using the examined  $\chi$  values was generally below 10%, but over 50% at  $D_g$  around  $20 \text{ }\mu\text{m}$ . The size  
 595 distributions were corrected as follows:

$$\left[ \frac{dN(D_g)}{d \log D_g} \right]_{\text{Corr,SMPS-GRIMM}} = \frac{dN(D_{g,SMPS-GRIMM})}{d \log D_{g,SMPS-GRIMM}} \cdot \frac{1}{1 - L(D_{g,SMPS-GRIMM})} \quad (\text{B2})$$

where  $dN(D_{g,SMPS-GRIMM})/d \log D_{g,SMPS-GRIMM}$  is the size distribution measured by the SMPS or GRIMM, and  $L(D_{g,SMPS-GRIMM})$  is the proportion of particle lost as a function of  $D_g$  in the size range measured by the SMPS or GRIMM. Finally, the size distribution of the particles suspended in CESAM was obtained summing up the contributions of the SMPS and GRIMM in their specific size range:

$$\left[ \frac{dN(D_g)}{d \log D_g} \right]_{\text{CESAM}} = \left\{ \begin{array}{l} \left[ \frac{dN(D_g)}{d \log D_g} \right]_{\text{Corr,SMPS}} ; \text{SMPS}_{\min} < D_g < \text{SMPS}_{\max} \\ \left[ \frac{dN(D_g)}{d \log D_g} \right]_{\text{Corr,GRIMM}} ; \text{GRIMM}_{\min} < D_g < \text{GRIMM}_{\max} \end{array} \right\} \quad (\text{B3})$$

600 Subsequently, the size distribution of the particles suspended in CESAM was used to evaluate the size-dependent correction factor in the sampling lines of the aethalometer and nephelometer. For the aethalometer, the particle loss was below 5% at  $D_g$

< 1µm, up to 20% in the range between 1-2.5 µm, below 60% in the range between 2.5-5 µm, and reached 100% at D<sub>g</sub> around 9 µm. The variation in particle loss calculated using the examined χ values was generally below 10%, but over 50% at D<sub>g</sub> around 9 µm. The percentage difference between the loss correction estimated for the aethalometer and that obtained  
605 for the nephelometer was < 11%. The data were corrected as follows:

$$\left[ \frac{dN(D_g)}{d \log D_g} \right]_{\text{Aet-Nep}} = \left[ \frac{dN(D_g)}{d \log D_g} \right]_{\text{CESAM}} \cdot \left( 1 - L(D_g)_{\text{Aet-Nep}} \right) \quad (\text{B4})$$

where  $[dN(D_g)/d \log D_g]_{\text{Aet-Nep}}$  is the size distribution measured by the aethalometer or nephelometer and  $L(D_g)_{\text{Aet-Nep}}$  is the particle loss as function of D<sub>g</sub> estimated for the two instruments. Since the size distributions of the aethalometer and nephelometer were very similar, we decided to estimate a common size distribution that would be representative of the dust particles sampled by these instruments. N<sub>sw-OPAs</sub>(D<sub>g</sub>) is the size distribution of the SW-OPAs which was calculated as the  
610 average of  $[dN(D_g)/d \log D_g]_{\text{Aet}}$  and  $[dN(D_g)/d \log D_g]_{\text{Nep}}$ .

## Appendix C: Truncation correction for total scattering

The nephelometer measurements were corrected for light scattered truncation for scattering angles ( $\theta$ ) between 0-7° and 170°-180° (Anderson et al., 1996; Anderson and Ogren, 1998). The truncation correction  $C_{\text{trunc}}$  for total scattering was  
615 estimated as follows:

$$C_{\text{trunc}}(\lambda, m, N(D_g)) = \frac{\beta_{\text{sca}}(\lambda, m, N(D_g), 0^\circ - 180^\circ)}{\beta_{\text{sca}}(\lambda, m, N(D_g), 7^\circ - 170^\circ)} \quad (\text{C1})$$

where  $\beta_{\text{sca}}(\lambda, m, N(D_g), 0^\circ-180^\circ)$  is the scattering coefficient calculated for  $\theta$  between 0° and 180°, and  $\beta_{\text{sca}}(\lambda, m, N(D_g), 7^\circ-170^\circ)$  is the scattering coefficient calculated for  $\theta$  between 7° and 170°. The Python PyMieScatt package (Sumlin et al., 2018) was used to retrieve  $\beta_{\text{sca}}(\lambda, m, N(D_g), 0^\circ-180^\circ)$  and  $\beta_{\text{sca}}(\lambda, m, N(D_g), 7^\circ-170^\circ)$  at a given  $\lambda$  by Mie calculation for homogeneous spheres by varying the particle size distribution  $N(D_g)$  based on the complex refractive index. The SW-OPA  
620 geometrical size distribution  $N_{\text{sw-OPAs}}(D_g)$  were used in the calculation.

For single homogeneous spherical particles, the scattering intensity function  $|S(X_s, m, \theta)|^2$  at given  $\theta$  can be calculated using the Mie theory, where  $X_s$  is the size parameter ( $X_s = \pi \cdot D_g \cdot \lambda^{-1}$ ) and  $m$  is the complex refractive index of particles of geometrical diameter  $D_g$ . The scattering intensity function can be integrated over  $\theta$  from 0° to 180° to obtain the single-particle total scattering efficiency (e.g., Anderson et al., 1996; Bohren and Hufmann, 1998):

$$Q_{\text{sca}}(\lambda, m, D_g) = \frac{1}{X_s^2} \cdot \int_0^\pi |S(X_s, m, \theta)|^2 \cdot \sin \theta \cdot d\theta \quad (\text{C2})$$

625 Finally,  $\beta_{\text{sca}}$  can be calculated as the integral of  $Q_{\text{sca}}$  over the particle size distribution  $N(D_g)$  multiplied by the particle cross sectional area (e.g., Anderson et al., 1996; Bohren and Hufmann, 1998):

$$\beta_{\text{sca}}(\lambda, m, N(D_g)) = \int_{D_{g,\text{min}}}^{D_{g,\text{max}}} \frac{\pi \cdot D_g^2}{4} \cdot Q_{\text{sca}}(\lambda, m, D_g) \cdot \frac{dN(D_g)}{d\log D_g} \cdot d\log D_g \quad (\text{C3})$$

In this study,  $\beta_{\text{sca}}(\lambda, m, N(D_g), 0^\circ-180^\circ)$  was computed at a given  $\lambda$  by the Mie\_SD function for polydisperse size distributions in the Python PyMieScatt package by inputting the calculated  $N_{\text{sw-OPAs}}(D_g)$ , as the particle size distribution, and the corresponding complex refractive index. To determine  $\beta_{\text{sca}}(\lambda, m, N(D_g), 7^\circ-170^\circ)$ , first the angular scattering intensities  
630  $|S(X_s, m, \theta)|^2$  for  $\theta$  between 7° and 170° were computed by the ScatteringFunction angular function, and then used in Eqs. (C2-3) to obtain  $Q_{\text{sca}}$  and  $\beta_{\text{sca}}$ .

## Appendix D: Calculation of relevant parameters to correct the attenuation coefficient

This section provides a description of the calculation of the parameters in Eq. (5) which were used to retrieve the absorption coefficient  $\beta_{\text{abs}}(\lambda)$  from the attenuation coefficient  $\beta_{\text{ATTN}}(\lambda)$ .

The parameter  $\alpha(\lambda)$  was calculated using the formula from Arnott et al. (2005):

$$\alpha(\lambda) = A^{(d-1)} \cdot c \cdot \lambda^{-\hat{a}_{\text{sca}} \cdot (d-1)} \quad (\text{D1})$$

where  $A$  and  $\hat{a}_{\text{sca}}$  were obtained from the power-law fitting of  $\beta_{\text{sca}}(\lambda)$  ( $\text{m}^{-1}$ ) versus  $\lambda$  (nm) weighted by the inverse of the variance of  $\beta_{\text{sca}}(\lambda)$ . The scattering coefficient  $\beta_{\text{sca}}(\lambda)$  measured by the nephelometer, corrected for truncation, and extrapolated at the aethalometer  $\lambda$  was used in the calculation. We assumed  $c = 3.29 \cdot 10^{-4}$  and  $d = 0.564$  as reported in

Collaud Coen et al. (2010).

The obtained  $\alpha(\lambda)$  increased with time as the particles were accumulating on the filter. Different  $\lambda$  showed similar values of the  $\alpha$  parameters. In the Icelandic dust samples,  $\alpha(\lambda)$  ranged from around 0.005 to 0.02, which is within the range (0.002 to 0.02) reported by Di Biagio et al. (2019) for coarse mineral dust. The uncertainty on  $\alpha(\lambda)$  varied between 17%-76% and was estimated by the error propagation through Eq. (D1) considering the uncertainty on the fitted parameters  $A$  and  $\hat{a}_{\text{sca}}$ . The uncertainty on  $\alpha(\lambda)$  was 35%-76% for the samples D3 and MIR45 which showed the largest errors on the fitted parameters, 29%-45% for the samples Maeli2 and H55, and 17%-27% for the sample Land1.

The multiple scattering effect correction  $C_{\text{ref}}$  was extrapolated at 370, 470, 520, 590, 660, 880, and 950 nm by the linear regression of  $C_{\text{ref}}$  at 450 and 660 nm estimated for mineral dust by Di Biagio et al. (2017b).  $C_{\text{ref}}$  varied from 4.3 at 370 nm to 3.3 at 950 nm. The uncertainty on  $C_{\text{ref}}$  determined by Di Biagio et al. (2017b) was 10%.

The loading effect correction  $R(\lambda)$  was calculated from the spectral attenuation ATTN measured by the aethalometer using the formula from Collaud Coen et al. (2010):

$$R(\lambda) = \left( \frac{1}{f(\lambda)} - 1 \right) \cdot \frac{\text{ATTN}(\lambda)\%}{50\%} + 1 \quad (\text{D2})$$

where  $f(\lambda)$  depends on the aerosol absorption properties and is a function of  $\text{SSA}(\lambda)$ :

$$f(\lambda) = a(1 - \text{SSA}(\lambda)) + 1 \quad (\text{D3})$$

We assumed  $a = 0.74$  as reported in Collaud Coen et al. (2010). As an initial guess for  $R(\lambda)$ , the single scattering albedo in Eq. (D3) was estimated using  $\beta_{\text{sca}}(\lambda)$  measured by the nephelometer, corrected for truncation, and extrapolated at the aethalometer  $\lambda$ , and  $\beta_{\text{abs}}(\lambda)^*$  which is the absorption coefficient corrected only for the scattering effect (Di Biagio et al., 2019):



$$\beta_{\text{abs}}(\lambda)^* = \beta_{\text{ATTN}}(\lambda) - \alpha(\lambda) \cdot \beta_{\text{sca}}(\lambda) \quad (\text{D4})$$

$$\text{SSA}(\lambda)^* = \frac{\beta_{\text{sca}}(\lambda)}{\beta_{\text{sca}}(\lambda) + \beta_{\text{abs}}(\lambda)^*} \quad (\text{D5})$$

R( $\lambda$ ) increased with  $\lambda$  but decreased with time as particles were loaded onto a new filter. R( $\lambda$ ) obtained after the second iteration varied between 0.7 and 1 in different Icelandic dust samples which is within the range reported by Di Biagio et al. (2019) for mineral dust, from 0.5 to 1. The uncertainty on R( $\lambda$ ) was also estimated by applying the error propagation method and was generally  $\leq 10\%$  but varied between 10%-50% within the first 10 min that the particles were loaded onto a new filter.

## Open Research

665 Datasets for this research are included in this paper (and its supplementary information files). Example research codes are available at the following GitHub repository: [https://github.com/ClarissaBaldo/Research\\_codes\\_examples](https://github.com/ClarissaBaldo/Research_codes_examples).

## Author contributions

670 CB, ZS, CDB, PF and JFD designed the experiments and discussed the results. ZS supervised the experimental and data analyses. CB performed the experiments at CESAM with support from ZS, CDB, PF, MC, EP, and JFD, and the data analysis with contributions from ZS, ARM, GL, CS, DB, CDB, and PF. GL refined the research codes. The soil samples used for the experiments were collected by OA and PDW. CB prepared the manuscript with contributions from all co-authors.

## 675 Competing interests

The contact author has declared that neither they nor their co-authors have any competing interests.

## Acknowledgements

Clarissa Baldo is funded by the Natural Environment Research Council (NERC) CENTA studentship (grant no. 680 NE/L002493/1). This paper is partly funded by the NERC highlight topic project (NE/S00579X/1). This project has received funding from the European Union's Horizon 2020 research and innovation programme through the EUROCHAMP-2020 Infrastructure Activity under grant agreement No 730997. Part of this work was supported by the COST Action inDust (CA16202) supported by COST (European Cooperation in Science and Technology). OA and PDW acknowledge financial support from the Czech Science Foundation (project No. 20-06168Y). GL thanks the PhD studentships funded by China 685 Scholarship Council. CNRS-INSU is gratefully acknowledged for supporting the CESAM chamber as a national facility as part of the French ACTRIS Research Infrastructure as well as the AERIS data center ([www.aeris-data.fr](http://www.aeris-data.fr)) for distributing and curing the data produced by the CESAM chamber through the hosting of the EUROCHAMP datacenter (<https://data.eurochamp.org>). The authors also wish to acknowledge the reviewers for their valuable feedback and insightful comments.

690

**Table 1: Experiment-averaged single scattering albedo  $SSA_{avg}(\lambda) \pm$  estimated uncertainty at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm of Icelandic dust for the base simulation.  $SSA_{avg}(\lambda)$  results from the base simulation are consistent with the results from Test 1 and Test 2 (see Table S2).**

Sample ID	$SSA_{avg}(\lambda)$						
	370 nm	470 nm	520 nm	590 nm	660 nm	880 nm	950 nm
D3	$0.93 \pm 0.02$	$0.95 \pm 0.01$	$0.96 \pm 0.01$	$0.96 \pm 0.01$	$0.96 \pm 0.02$	$0.96 \pm 0.01$	$0.96 \pm 0.02$
H55	$0.94 \pm 0.06$	$0.96 \pm 0.08$	$0.96 \pm 0.08$	$0.96 \pm 0.08$	$0.96 \pm 0.08$	$0.96 \pm 0.07$	$0.96 \pm 0.07$
Land1	$0.91 \pm 0.05$	$0.94 \pm 0.03$	$0.95 \pm 0.03$	$0.95 \pm 0.03$	$0.96 \pm 0.04$	$0.96 \pm 0.04$	$0.96 \pm 0.04$
Maeli2	$0.90 \pm 0.03$	$0.93 \pm 0.02$	$0.94 \pm 0.02$	$0.95 \pm 0.02$	$0.95 \pm 0.01$	$0.96 \pm 0.01$	$0.95 \pm 0.02$
MIR45	$0.90 \pm 0.04$	$0.92 \pm 0.03$	$0.93 \pm 0.02$	$0.94 \pm 0.03$	$0.94 \pm 0.03$	$0.94 \pm 0.03$	$0.94 \pm 0.03$


695

**700 Table 2: Experiment-averaged imaginary index  $k_{avg}(\lambda) \pm$  estimated uncertainty at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm of Icelandic dust.  $k_{avg}(\lambda)$  data are the mean of the results of the base simulation and Test 1 in Table S3.**

Sample ID	$k_{avg}(\lambda)$						
	370 nm	470 nm	520 nm	590 nm	660 nm	880 nm	950 nm
D3	$0.004 \pm 0.002$	$0.003 \pm 0.001$	$0.002 \pm 0.001$	$0.002 \pm 0.001$	$0.002 \pm 0$	$0.002 \pm 0$	$0.002 \pm 0$
H55	$0.004 \pm 0.001$	$0.004 \pm 0.002$	$0.003 \pm 0.001$	$0.002 \pm 0.001$	$0.002 \pm 0$	$0.002 \pm 0.001$	$0.002 \pm 0$
Land1	$0.004 \pm 0.001$	$0.004 \pm 0.002$	$0.004 \pm 0.002$	$0.003 \pm 0.001$	$0.003 \pm 0.001$	$0.002 \pm 0.001$	$0.002 \pm 0.001$
Maeli2	$0.004 \pm 0.002$	$0.004 \pm 0.002$	$0.002 \pm 0.001$	$0.003 \pm 0.001$	$0.003 \pm 0.001$	$0.002 \pm 0.000$	$0.002 \pm 0.001$
MIR45	$0.004 \pm 0.001$	$0.004 \pm 0.001$	$0.003 \pm 0.001$	$0.003 \pm 0.001$	$0.003 \pm 0.001$	$0.003 \pm 0.000$	$0.003 \pm 0$


**Table 3: Experiment-averaged real index  $n_{\text{avg}}(\lambda) \pm$  estimated uncertainty at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm of Icelandic dust.  $n_{\text{avg}}(\lambda)$  data are the mean of the results of the base simulation and Test 1 in Table S4.**

Sample ID	$n_{\text{avg}}(\lambda)$						
	370 nm	470 nm	520 nm	590 nm	660 nm	880 nm	X950 nm
D3	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.02$	$1.60 \pm 0.02$	$1.60 \pm 0.01$
H55	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.61 \pm 0.02$	$1.60 \pm 0.01$	$1.60 \pm 0$
Land1	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.61 \pm 0.01$	$1.60 \pm 0.01$	$1.61 \pm 0.01$
Maeli2	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.02$	$1.61 \pm 0.01$	$1.60 \pm 0.01$	$1.61 \pm 0.01$	$1.61 \pm 0.01$
MIR45	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$	$1.61 \pm 0.01$	$1.60 \pm 0.01$	$1.60 \pm 0.01$




**Figure 1: Schematic diagram of the method used to retrieve the shortwave (SW) single scattering albedo (SSA) and complex refractive index ( $m = n - ik$ ) of Icelandic dust for  $\lambda$  from 370 to 950 nm. See main text for details.  $\beta_{sca}(\lambda)$  is the spectral scattering coefficient retrieved from the nephelometer measurements, and  $\theta$  is the light scattering angle.  $\beta_{ATTN}(\lambda)$  and  $\beta_{abs}(\lambda)$  are respectively the spectral attenuation and absorption coefficients retrieved from the aethalometer measurements.  $dN(D_m)/d\log D_m$  is the particle number concentration measured by the SMPS – scanning mobility particle sizer, where  $D_m$  is the electrical mobility diameter.  $dN(D_{op})/d\log D_{op}$  is the particle number concentration measured by the GRIMM – optical particle counter, where  $D_{op}$  is the optical-equivalent diameter.  $dN(D_{g,SMPS-GRIMM})/d\log D_{g,SMPS-GRIMM}$  is the geometrical size distribution retrieved from the SMPS or GRIMM measurements, where  $D_g$  is the geometrical diameter obtained from  $D_m$  or  $D_{op}$  using selected intervals for the dust particle dynamic shape factor ( $\chi$ ), and the real ( $n$ ) and imaginary ( $k$ ) part of the complex refractive index, respectively. The term SW-OPA (Shortwave Optical Properties Analyzers) refers to the aethalometer and nephelometer.**


## Maeli2



**Figure 2:** Geometrical size distributions  $dN(D_g)/d\log D_g$  at 30 min after the injection peak obtained a) using the calibration values of  $\chi$ ,  $n$  and  $k$ ; b-f) for different  $\chi$ - $n$ - $k$  combinations.  $\chi$  is the dynamic shape factor used to convert the mobility diameter  $D_m$  measured by the SMPS into geometrical diameters  $D_g$ ,  $n$  and  $k$  are respectively the real and imaginary part of the complex refractive index used to convert the optical diameter  $D_{op}$  measured by the GRIMM into  $D_g$ . Sample ID: Maeli2. Note: The size ranges shown are before correcting the size distributions for particle loss along the instrument sampling systems. In the end, particles suspended in CESAM were up to 20  $\mu m$  in size, while particles sampled by the SW-OPAs were up to 9  $\mu m$  in size. Details of the particle loss correction can be found in Appendix B




**Figure 3: Effective diameters  $D_{\text{eff}}$  of dust particles sampled by the SW-OPAs and in CESAM from 30 min after the injection peak to 2.5 h. a-b) Base simulation; c-d) Test 1; e-f) Test 2.  $D_{\text{eff}}$  was calculated for particles  $> 1 \mu\text{m}$  ( $D_{\text{eff,coarse}}$ ) and  $\leq 1 \mu\text{m}$  ( $D_{\text{eff,fine}}$ ). Data were reported as 12-min average. In Test 1, corrections and calculations were performed using the SMPS and GRIMM data plus 1 SD uncertainty. In Test 2, we used the SMPS and GRIMM data minus 1 SD uncertainty (see section 2.2.1 in the main text for details). Sample IDs: D3 and Maeli2.**




**Figure 4:** Extinction coefficient  $\beta_{\text{ext}}(\lambda)$ , absorption coefficient  $\beta_{\text{abs}}(\lambda)$ , and single scattering albedo  $\text{SSA}(\lambda)$  of Icelandic dust at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm, from 30 min after the injection peak to 2.5 h. The reported results as 12-min average refer to the base simulation.




Maeli2




**Figure 5:** Real index  $n(\lambda)$  and imaginary index  $k(\lambda)$  at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm, from 30 min after the injection peak to 2.5 h. a-b) Base simulation; c-d) Test 1; e-f) Test 2. In Test 1, corrections and calculations were performed using the SMPS and GRIMM data plus 1 SD uncertainty. In Test 2, we used the SMPS and GRIMM data minus 1 SD uncertainty (see section 2.2.1 in the main text for details). Sample ID: Maeli2.




**Figure 6:** Experiment-averaged single scattering albedo  $SSA_{avg}(\lambda)$  at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm of Icelandic dust samples for the base simulation (Table 1).  $SSA_{avg}(\lambda)$  results from the base simulation are consistent with the results from Test 1 and Test 2 (see Table S2).



**Figure 7:** Comparison between the experiment-averaged imaginary index  $k_{\text{avg}}(\lambda)$  at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm of Icelandic dust (Table 2) and the initial estimates by Baldo et al. (2020) based on the mineralogical composition (low and high absorption case).



**Figure 8.** Comparison between the experiment-averaged a) real index  $n_{\text{avg}}(\lambda)$ , and b) imaginary index  $k_{\text{avg}}(\lambda)$  at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm of Icelandic dust (Tables 2-3) and Icelandic volcanic ash (Bukowiecki et al., 2011; Deguine et al., 2020; Derimian et al., 2012; Reed et al., 2018; Weinzierl et al., 2012). Note that  $n(\lambda)$  and  $k(\lambda)$  data of Deguine et al. (2020) and Reed et al. (2018) were downloaded from <http://eodg.atm.ox.ac.uk/> (last access 05 January 2023). Bukowiecki et al. (2011) reported an average  $k$  value of 0.02 between 450-700 nm, while  $n$  ranged from 1.60 to 1.80 (not shown).



**Figure 9: Comparison between the experiment-averaged a) real index  $n_{avg}(\lambda)$ , b) imaginary index  $k_{avg}(\lambda)$ , and c) single scattering albedo  $SSA_{avg}(\lambda)$  at  $\lambda = 370, 470, 520, 590, 660, 880, 950$  nm of Icelandic dust (Tables 1-3) and mineral dust from major dust source**

765 regions at low-latitude (Di Biagio et al., 2019) including northern Africa-Sahara, the Sahel, and eastern Asia. Note we reported the real index of low-latitude mineral dust as the average value in the range 370-950 nm according to Table 4 of Di Biagio et al. (2019).

## References

- Achterberg, E. P., Moore, C. M., Henson, S. A., Steigenberger, S., Stohl, A., Eckhardt, S., Avendano, L. C., Cassidy, M., Hembury, D., Klar, J. K., Lucas, M. I., Macey, A. I., Marsay, C. M., and Ryan-Keogh, T. J.: Natural iron fertilization by the Eyjafjallajökull volcanic eruption, *Geophys. Res. Lett.*, 40, 921-926, doi: 10.1002/grl.50221, 2013.
- Achterberg, E. P., Steigenberger, S., Marsay, C. M., LeMoigne, F. A. C., Painter, S. C., Baker, A. R., Connelly, D. P., Moore, C. M., Tagliabue, A., and Tanhua, T.: Iron Biogeochemistry in the High Latitude North Atlantic Ocean, *Sci. Rep.*, 8, 1283, doi: 10.1038/s41598-018-19472-1, 2018.
- Anderson, T. L., Covert, D. S., Marshall, S. F., Laucks, M. L., Charlson, R. J., Waggoner, A. P., Ogren, J. A., Caldow, R., Holm, R. L., Quant, F. R., Sem, G. J., Wiedensohler, A., Ahlquist, N. A., and Bates, T. S.: Performance Characteristics of a High-Sensitivity, Three-Wavelength, Total Scatter/Backscatter Nephelometer, *J. Atmos. Ocean. Technol.*, 13, 967-986, doi: 10.1175/1520-0426(1996)013<0967:Pcoahs>2.0.Co;2, 1996.
- Anderson, T. L., and Ogren, J. A.: Determining Aerosol Radiative Properties Using the TSI 3563 Integrating Nephelometer, *Aerosol Sci. Tech.*, 29, 57-69, doi: 10.1080/02786829808965551, 1998.
- Arnalds, O., Olafsson, H., and Dagsson-Waldhauserova, P.: Quantification of iron-rich volcanogenic dust emissions and deposition over the ocean from Icelandic dust sources, *Biogeosciences*, 11, 6623-6632, doi: 10.5194/bg-11-6623-2014, 2014.
- Arnalds, O., Dagsson-Waldhauserova, P., and Olafsson, H.: The Icelandic volcanic aeolian environment: Processes and impacts — A review, *Aeolian Res.*, 20, 176-195, doi: 10.1016/j.aeolia.2016.01.004, 2016.
- Arnott, W. P., Hamasha, K., Moosmüller, H., Sheridan, P. J., and Ogren, J. A.: Towards Aerosol Light-Absorption Measurements with a 7-Wavelength Aethalometer: Evaluation with a Photoacoustic Instrument and 3-Wavelength Nephelometer, *Aerosol Sci. Tech.*, 39, 17-29, doi: 10.1080/027868290901972, 2005.
- Ayers, G. P.: Comment on regression analysis of air quality data, *Atmos. Environ.*, 35, 2423-2425, doi: doi.org/10.1016/S1352-2310(00)00527-6, 2001.
- Baddock, M. C., Mockford, T., Bullard, J. E., and Thorsteinsson, T.: Pathways of high-latitude dust in the North Atlantic, *Earth Planet. Sci. Lett.*, 459, 170-182, doi: 10.1016/j.epsl.2016.11.034, 2017.
- Baldo, C., Formenti, P., Nowak, S., Chevaillier, S., Cazaunau, M., Pangui, E., Di Biagio, C., Doussin, J. F., Ignatyev, K., Dagsson-Waldhauserova, P., Arnalds, O., MacKenzie, A. R., and Shi, Z.: Distinct chemical and mineralogical composition of Icelandic dust compared to northern African and Asian dust, *Atmos. Chem. Phys.*, 20, 13521-13539, doi: 10.5194/acp-20-13521-2020, 2020.
- Balkanski, Y., Schulz, M., Claquin, T., and Guibert, S.: Reevaluation of Mineral aerosol radiative forcings suggests a better agreement with satellite and AERONET data, *Atmos. Chem. Phys.*, 7, 81-95, doi: 10.5194/acp-7-81-2007, 2007.
- Beddows, D. C. S., Dall'osto, M., and Harrison, R. M.: An Enhanced Procedure for the Merging of Atmospheric Particle Size Distribution Data Measured Using Electrical Mobility and Time-of-Flight Analysers, *Aerosol Sci. Tech.*, 44, 930-938, doi: 10.1080/02786826.2010.502159, 2010.
- Bohren, C. F., and Hufmann, D. R.: Absorption and Scattering by a Sphere, in: *Absorption and Scattering of Light by Small Particles*, 82-129, 1998.

- Bristow, C. S., Drake, N., and Armitage, S.: Deflation in the dustiest place on Earth: The Bodélé Depression, Chad, *Geomorphology*, 105, 50-58, doi: 10.1016/j.geomorph.2007.12.014, 2009.
- 805 Bukowiecki, N., Zieger, P., Weingartner, E., Juranyi, Z., Gysel, M., Neiningner, B., Schneider, B., Hueglin, C., Ulrich, A., Wichser, A., Henne, S., Brunner, D., Kaegi, R., Schwikowski, M., Tobler, L., Wienhold, F. G., Engel, I., Buchmann, B., Peter, T., and Baltensperger, U.: Ground-based and airborne in-situ measurements of the Eyjafjallajökull volcanic aerosol plume in Switzerland in spring 2010, *Atmos. Chem. Phys.*, 11, 10011-10030, doi: 10.5194/acp-11-10011-2011, 2011.
- 810 Bullard, J. E., Baddock, M., Bradwell, T., Crusius, J., Darlington, E., Gaiero, D., Gasso, S., Gisladottir, G., Hodgkins, R., McCulloch, R., McKenna-Neuman, C., Mockford, T., Stewart, H., and Thorsteinsson, T.: High-latitude dust in the Earth system, *Rev. Geophys.*, 54, 447-485, doi: 10.1002/2016rg000518, 2016.
- Bullard, J. E.: The distribution and biogeochemical importance of high-latitude dust in the Arctic and Southern Ocean-Antarctic regions, *J. Geophys. Res.-Atmos.*, 122, 3098-3103, doi: 10.1002/2016jd026363, 2017.
- Butwin, M. K., Pfeffer, M. A., von Löwis, S., Støren, E. W. N., Bali, E., and Thorsteinsson, T.: Properties of dust source material and volcanic ash in Iceland, *Sedimentology*, 67, 3067-3087, doi: 10.1111/sed.12734, 2020.
- 815 Caponi, L., Formenti, P., Massabo, D., Di Biagio, C., Cazaunau, M., Pangui, E., Chevaillier, S., Landrot, G., Andreae, M. O., Kandler, K., Piketh, S., Saeed, T., Seibert, D., Williams, E., Balkanski, Y., Prati, P., and Doussin, J. F.: Spectral- and size-resolved mass absorption efficiency of mineral dust aerosols in the shortwave spectrum: a simulation chamber study, *Atmos. Chem. Phys.*, 17, 7175-7191, doi: 10.5194/acp-17-7175-2017, 2017.
- 820 Collaud Coen, M., Weingartner, E., Apituley, A., Ceburnis, D., Fierz-Schmidhauser, R., Flentje, H., Henzing, J. S., Jennings, S. G., Moerman, M., Petzold, A., Schmid, O., and Baltensperger, U.: Minimizing light absorption measurement artifacts of the Aethalometer: evaluation of five correction algorithms, *Atmos. Meas. Tech.*, 3, 457-474, doi: 10.5194/amt-3-457-2010, 2010.
- 825 Cook, J. M., Tedstone, A. J., Williamson, C., McCutcheon, J., Hodson, A. J., Dayal, A., Skiles, M., Hofer, S., Bryant, R., McAree, O., McGonigle, A., Ryan, J., Anesio, A. M., Irvine-Fynn, T. D. L., Hubbard, A., Hanna, E., Flanner, M., Mayanna, S., Benning, L. G., van As, D., Yallop, M., McQuaid, J. B., Gribbin, T., and Tranter, M.: Glacier algae accelerate melt rates on the south-western Greenland Ice Sheet, *Cryosphere*, 14, 309-330, doi: 10.5194/tc-14-309-2020, 2020.
- Dagsson-Waldhauserova, P., Arnalds, O., and Olafsson, H.: Long-term frequency and characteristics of dust storm events in Northeast Iceland (1949–2011), *Atmos. Environ.*, 77, 117-127, doi: 10.1016/j.atmosenv.2013.04.075, 2013.
- 830 Dagsson-Waldhauserova, P.: Variability, origin and physical characteristics of dust aerosol in Iceland (PhD dissertation), Faculty of Physical Sciences Agricultural University of Iceland, Reykjavik, 163 pp., 2014.
- Dagsson-Waldhauserova, P., Arnalds, O., and Olafsson, H.: Long-term variability of dust events in Iceland (1949–2011), *Atmos. Chem. Phys.*, 14, 13411-13422, doi: 10.5194/acp-14-13411-2014, 2014.
- 835 Dagsson-Waldhauserova, P., Arnalds, O., Olafsson, H., Hladil, J., Skala, R., Navratil, T., Chadimova, L., and Meinander, O.: Snow–Dust Storm: Unique case study from Iceland, March 6–7, 2013, *Aeolian Res.*, 16, 69-74, doi: 10.1016/j.aeolia.2014.11.001, 2015.
- Dagsson-Waldhauserova, P., Renard, J.-B., Olafsson, H., Vignelles, D., Berthet, G., Verdier, N., and Duverger, V.: Vertical distribution of aerosols in dust storms during the Arctic winter, *Sci. Rep.*, 9, 16122, doi: 10.1038/s41598-019-51764-y, 2019.



- DeCarlo, P. F., Slowik, J. G., Worsnop, D. R., Davidovits, P., and Jimenez, J. L.: Particle Morphology and Density Characterization by Combined Mobility and Aerodynamic Diameter Measurements. Part 1: Theory, *Aerosol Sci. Tech.*, 38, 1185-1205, doi: 10.1080/027868290903907, 2004.
- 840 Deguine, A., Petitprez, D., Clarisse, L., Gudmundsson, S., Outes, V., Villarosa, G., and Herbin, H.: Complex refractive index of volcanic ash aerosol in the infrared, visible, and ultraviolet, *Appl. Opt.*, 59, 884-895, doi: 10.1364/AO.59.000884, 2020.
- 845 Deguine, A., Petitprez, D., Clarisse, L., Deschutter, L., Fontijn, K., and Herbin, H.: Retrieval of refractive indices of ten volcanic ash samples in the infrared, visible and ultraviolet spectral region, *J. Aerosol Sci.*, 167, 106100, doi: 10.1016/j.jaerosci.2022.106100, 2023.
- Derimian, Y., Karnieli, A., Kaufman, Y. J., Andreae, M. O., Andreae, T. W., Dubovik, O., Maenhaut, W., and Koren, I.: The role of iron and black carbon in aerosol light absorption, *Atmos. Chem. Phys.*, 8, 3623-3637, doi: 10.5194/acp-8-3623-2008, 2008.
- 850 Derimian, Y., Dubovik, O., Tanre, D., Goloub, P., Lapyonok, T., and Mortier, A.: Optical properties and radiative forcing of the Eyjafjallajökull volcanic ash layer observed over Lille, France, in 2010, *J. Geophys. Res.-Atmos.*, 117, D00U25, doi: 10.1029/2011jd016815, 2012.
- 855 Di Biagio, C., Formenti, P., Balkanski, Y., Caponi, L., Cazaunau, M., Pangui, E., Journet, E., Nowak, S., Caqueneau, S., Andreae, M. O., Kandler, K., Saeed, T., Piketh, S., Seibert, D., Williams, E., and Doussin, J. F.: Global scale variability of the mineral dust long-wave refractive index: a new dataset of in situ measurements for climate modeling and remote sensing, *Atmos. Chem. Phys.*, 17, 1901-1929, doi: 10.5194/acp-17-1901-2017, 2017a.
- Di Biagio, C., Formenti, P., Cazaunau, M., Pangui, E., Marchand, N., and Doussin, J. F.: Aethalometer multiple scattering correction C-ref for mineral dust aerosols, *Atmos. Meas. Tech.*, 10, 2923-2939, doi: 10.5194/amt-10-2923-2017, 2017b.
- 860 Di Biagio, C., Formenti, P., Balkanski, Y., Caponi, L., Cazaunau, M., Pangui, E., Journet, E., Nowak, S., Andreae, M. O., Kandler, K., Saeed, T., Piketh, S., Seibert, D., Williams, E., and Doussin, J. F.: Complex refractive indices and single-scattering albedo of global dust aerosols in the shortwave spectrum and relationship to size and iron content, *Atmos. Chem. Phys.*, 19, 15503-15531, doi: 10.5194/acp-19-15503-2019, 2019.
- 865 Di Biagio, C., Balkanski, Y., Albani, S., Boucher, O., and Formenti, P.: Direct Radiative Effect by Mineral Dust Aerosols Constrained by New Microphysical and Spectral Optical Data, *Geophys. Res. Lett.*, 47, e2019GL086186, doi: 10.1029/2019GL086186, 2020.
- Dragosics, M., Meinander, O., Jónsdóttir, T., Dürig, T., De Leeuw, G., Pálsson, F., Dagsson-Waldhauserová, P., and Thorsteinsson, T.: Insulation effects of Icelandic dust and volcanic ash on snow and ice, *Arab. J. Geosci.*, 9, 126, doi: 10.1007/s12517-015-2224-6, 2016.
- 870 Dubovik, O., Holben, B., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanre, D., and Slutsker, I.: Variability of absorption and optical properties of key aerosol types observed in worldwide locations, *J. Atmos. Sci.*, 59, 590-608, doi: 10.1175/1520-0469(2002)059<0590:voaaop>2.0.co;2, 2002.
- Engelbrecht, J. P., Moosmüller, H., Pincock, S., Jayanty, R. K. M., Lersch, T., and Casuccio, G.: Technical note: Mineralogical, chemical, morphological, and optical interrelationships of mineral dust re-suspensions, *Atmos. Chem. Phys.*, 16, 10809-10830, doi: 10.5194/acp-16-10809-2016, 2016.

- 875 Formenti, P., Rajot, J. L., Desboeufs, K., Said, F., Grand, N., Chevaillier, S., and Schmechtig, C.: Airborne observations of mineral dust over western Africa in the summer Monsoon season: spatial and vertical variability of physico-chemical and optical properties, *Atmos. Chem. Phys.*, 11, 6387-6410, doi: 10.5194/acp-11-6387-2011, 2011.
- Formenti, P., Caquineau, S., Chevaillier, S., Klaver, A., Desboeufs, K., Rajot, J. L., Belin, S., and Briois, V.: Dominance of goethite over hematite in iron oxides of mineral dust from Western Africa: Quantitative partitioning by X-ray absorption spectroscopy, *J. Geophys. Res.-Atmos.*, 119, 12740-12754, doi: 10.1002/2014jd021668, 2014.
- 880 Formenti, P., Di Biagio, C., Huang, Y., Kok, J., Mallet, M. C., Boulanger, D., and Cazaunau, M.: Look-up tables resolved by complex refractive index to correct particle sizes measured by common research-grade optical particle counters, *Atmos. Meas. Tech. Discuss.*, doi: 10.5194/amt-2021-403, 2021.
- Groot Zwaaftink, C. D., Grythe, H., Skov, H., and Stohl, A.: Substantial contribution of northern high-latitude sources to mineral dust in the Arctic, *J. Geophys. Res.-Atmos.*, 121, 13678-13697, doi: 10.1002/2016jd025482, 2016.
- 885 Groot Zwaaftink, C. D., Arnalds, O., Dagsson-Waldhauserova, P., Eckhardt, S., Prospero, J. M., and Stohl, A.: Temporal and spatial variability of Icelandic dust emissions and atmospheric transport, *Atmos. Chem. Phys.*, 17, 10865-10878, doi: 10.5194/acp-17-10865-2017, 2017.
- Gunnarsson, A., Gardarsson, S. M., Pálsson, F., Jóhannesson, T., and Sveinsson, Ó. G. B.: Annual and inter-annual variability and trends of albedo of Icelandic glaciers, *Cryosphere*, 15, 547-570, doi: 10.5194/tc-15-547-2021, 2021.
- 890 Hand, J. L., and Kreidenweis, S. M.: A New Method for Retrieving Particle Refractive Index and Effective Density from Aerosol Size Distribution Data, *Aerosol Sci. Tech.*, 36, 1012-1026, doi: 10.1080/02786820290092276, 2002.
- Hansen, J. E.: Multiple Scattering of Polarized Light in Planetary Atmospheres Part II. Sunlight Reflected by Terrestrial Water Clouds, *J. Atmos. Sci.*, 28, 1400-1426, doi: 10.1175/1520-0469(1971)028<1400:Msopli>2.0.Co;2, 1971.
- 895 Hinds, W. C.: *Aerosol Technology : Properties, Behavior, and Measurement of Airborne Particles*, John Wiley & Sons, Incorporated, Somerset, UNITED STATES, 1999.
- Huang, Y., Adebisi, A. A., Formenti, P., and Kok, J. F.: Linking the Different Diameter Types of Aspherical Desert Dust Indicates That Models Underestimate Coarse Dust Emission, *Geophys. Res. Lett.*, 48, e2020GL092054, doi: doi.org/10.1029/2020GL092054, 2021.
- 900 Huffman, D. R., and Stapp, J. L.: Optical measurements on solids of possible interstellar importance, in: *Interstellar dust and related topics*, Springer, 297-301, 1973.
- Jeong, G. Y.: Bulk and single-particle mineralogy of Asian dust and a comparison with its source soils, *J. Geophys. Res.-Atmos.*, 113, D02208, doi: 10.1029/2007jd008606, 2008.
- Jia, J., Wang, Y. J., Xia, D. S., Lu, H., and Gao, F. Y.: Dust Sources of Last Glacial Chinese Loess Based on the Iron Mineralogy of Fractionated Source Samples, *Geophys. Res. Lett.*, 46, 2103-2110, doi: 10.1029/2018gl080909, 2019.
- 905 Johnson, B. T., Osborne, S. R., Haywood, J. M., and Harrison, M. A. J.: Aircraft measurements of biomass burning aerosol over West Africa during DABEX, *J. Geophys. Res.-Atmos.*, 113, D00C06, doi: 10.1029/2007JD009451, 2008.
- Johnson, B. T., and Osborne, S. R.: Physical and optical properties of mineral dust aerosol measured by aircraft during the GERBILS campaign, *Quarterly Journal of the Royal Meteorological Society*, 137, 1117-1130, doi: 10.1002/qj.777, 2011.

- 910 Kandler, K., Benker, N., Bundke, U., Cuevas, E., Ebert, M., Knippertz, P., Rodriguez, S., Schutz, L., and Weinbruch, S.: Chemical composition and complex refractive index of Saharan Mineral Dust at Izana, Tenerife (Spain) derived by electron microscopy, *Atmos. Environ.*, 41, 8058-8074, doi: 10.1016/j.atmosenv.2007.06.047, 2007.
- Kylling, A., Zwaafink, C. D. G., and Stohl, A.: Mineral Dust Instantaneous Radiative Forcing in the Arctic, *Geophys. Res. Lett.*, 45, 4290-4298, doi: 10.1029/2018gl077346, 2018.
- 915 Lafon, S., Sokolik, I. N., Rajot, J. L., Caquineau, S., and Gaudichet, A.: Characterization of iron oxides in mineral dust aerosols: Implications for light absorption, *J. Geophys. Res.-Atmos.*, 111, D21207, doi: 10.1029/2005jd007016, 2006.
- Lazaro, F. J., Gutierrez, L., Barron, V., and Gelado, M. D.: The speciation of iron in desert dust collected in Gran Canaria (Canary Islands): Combined chemical, magnetic and optical analysis, *Atmos. Environ.*, 42, 8987-8996, doi: 10.1016/j.atmosenv.2008.09.035, 2008.
- 920 Li, L., Mahowald, N. M., Miller, R. L., Pérez García-Pando, C., Klose, M., Hamilton, D. S., Gonçalves Ageitos, M., Ginoux, P., Balkanski, Y., Green, R. O., Kalashnikova, O., Kok, J. F., Obiso, V., Paynter, D., and Thompson, D. R.: Quantifying the range of the dust direct radiative effect due to source mineralogy uncertainty, *Atmos. Chem. Phys.*, 21, 3973-4005, doi: 10.5194/acp-21-3973-2021, 2021.
- Lutz, S., Anesio, A. M., Edwards, A., and Benning, L. G.: Microbial diversity on Icelandic glaciers and ice caps, *Front. Microbiol.*, 6, 307, doi: 10.3389/fmicb.2015.00307, 2015.
- 925 Maher, B. A., Mutch, T. J., and Cunningham, D.: Magnetic and geochemical characteristics of Gobi Desert surface sediments: Implications for provenance of the Chinese Loess Plateau, *Geology*, 37, 279-282, doi: 10.1130/g25293a.1, 2009.
- McCutcheon, J., Lutz, S., Williamson, C., Cook, J. M., Tedstone, A. J., Vanderstraeten, A., Wilson, S. A., Stockdale, A., Bonneville, S., Anesio, A. M., Yallop, M. L., McQuaid, J. B., Tranter, M., and Benning, L. G.: Mineral phosphorus drives glacier algal blooms on the Greenland Ice Sheet, *Nat. Commun.*, 12, 570, doi: 10.1038/s41467-020-20627-w, 2021.
- 930 Meinander, O., Kontu, A., Virkkula, A., Arola, A., Backman, L., Dagsson-Waldhauserová, P., Järvinen, O., Manninen, T., Svensson, J., de Leeuw, G., and Leppäranta, M.: Brief communication: Light-absorbing impurities can reduce the density of melting snow, *Cryosphere*, 8, 991-995, doi: 10.5194/tc-8-991-2014, 2014.
- Meinander, O., Dagsson-Waldhauserova, P., Amosov, P., Aseyeva, E., Atkins, C., Baklanov, A., Baldo, C., Barr, S. L., Barzycka, B., Benning, L. G., Cvetkovic, B., Enchilik, P., Frolov, D., Gassó, S., Kandler, K., Kasimov, N., Kavan, J., King, J., Koroleva, T., Krupskaya, V., Kulmala, M., Kusiak, M., Lappalainen, H. K., Laska, M., Lasne, J., Lewandowski, M., Luks, B., McQuaid, J. B., Moroni, B., Murray, B., Möhler, O., Nawrot, A., Nickovic, S., O'Neill, N. T., Pejanovic, G., Popovicheva, O., Ranjbar, K., Romanias, M., Samonova, O., Sanchez-Marroquin, A., Schepanski, K., Semenov, I., Sharapova, A., Shevnina, E., Shi, Z., Sofiev, M., Thevenet, F., Thorsteinsson, T., Timofeev, M., Umo, N. S., Uppstu, A., 940 Urupina, D., Varga, G., Werner, T., Arnalds, O., and Vukovic Vimic, A.: Newly identified climatically and environmentally significant high-latitude dust sources, *Atmos. Chem. Phys.*, 22, 11889-11930, doi: 10.5194/acp-22-11889-2022, 2022.
- Möller, R., Möller, M., Kukla, P. A., and Schneider, C.: Impact of supraglacial deposits of tephra from Grímsvötn volcano, Iceland, on glacier ablation, *J. Glaciol.*, 62, 933-943, doi: 10.1017/jog.2016.82, 2016.
- 945 Möller, R., Möller, M., Kukla, P. A., and Schneider, C.: Modulation of glacier ablation by tephra coverage from Eyjafjallajökull and Grímsvötn volcanoes, Iceland: an automated field experiment, *Earth Syst. Sci. Data*, 10, 53-60, doi: 10.5194/essd-10-53-2018, 2018.

- Moosmuller, H., Engelbrecht, J. P., Skiba, M., Frey, G., Chakrabarty, R. K., and Arnott, W. P.: Single scattering albedo of fine mineral dust aerosols controlled by iron concentration, *J. Geophys. Res.-Atmos.*, 117, D11210, doi: 10.1029/2011jd016909, 2012.
- 950 Moroni, B., Arnalds, O., Dagsson-Waldhauserova, P., Crocchianti, S., Vivani, R., and Cappelletti, D.: Mineralogical and Chemical Records of Icelandic Dust Sources Upon Ny-angstrom lesund (Svalbard Islands), *Front. Earth Sci.*, 6, UNSP 187, doi: 10.3389/feart.2018.00187, 2018.
- Morrison, H., de Boer, G., Feingold, G., Harrington, J., Shupe, M. D., and Sulia, K.: Resilience of persistent Arctic mixed-phase clouds, *Nat. Geosci.*, 5, 11-17, doi: 10.1038/ngeo1332, 2012.
- 955 Moskowitz, B. M., Reynolds, R. L., Goldstein, H. L., Berquo, T. S., Kokaly, R. F., and Bristow, C. S.: Iron oxide minerals in dust-source sediments from the Bodele Depression, Chad: Implications for radiative properties and Fe bioavailability of dust plumes from the Sahara, *Aeolian Res.*, 22, 93-106, doi: 10.1016/j.aeolia.2016.07.001, 2016.
- Müller, T., Schladitz, A., Kandler, K., and Wiedensohler, A.: Spectral particle absorption coefficients, single scattering albedos and imaginary parts of refractive indices from ground based in situ measurements at Cape Verde Island during SAMUM-2, *Tellus Series B-Chemical and Physical Meteorology*, 63, 573-588, doi: 10.1111/j.1600-0889.2011.00572.x, 2011.
- 960 Murray, B. J., Carslaw, K. S., and Field, P. R.: Opinion: Cloud-phase climate feedback and the importance of ice-nucleating particles, *Atmos. Chem. Phys.*, 21, 665-679, doi: 10.5194/acp-21-665-2021, 2021.
- Nakashima, M., and Dagsson-Waldhauserová, P.: A 60 Year Examination of Dust Day Activity and Its Contributing Factors From Ten Icelandic Weather Stations From 1950 to 2009, *Front. Earth Sci.*, 6, UNSP 245, doi: 10.3389/feart.2018.00245, 2019.
- Okada, K., Heintzenberg, J., Kai, K. J., and Qin, Y.: Shape of atmospheric mineral particles collected in three Chinese arid-regions, *Geophys. Res. Lett.*, 28, 3123-3126, doi: 10.1029/2000gl012798, 2001.
- Patterson, E. M., Gillette, D. A., and Stockton, B. H.: Complex index of refraction between 300 and 700 nm for Saharan aerosols, *J. Geophys. Res.*, 82, 3153-3160, doi: 10.1029/JC082i021p03153, 1977.
- 970 Peltoniemi, J. I., Gritsevich, M., Hakala, T., Dagsson-Waldhauserova, P., Arnalds, O., Anttila, K., Hannula, H. R., Kivekas, N., Lihavainen, H., Meinander, O., Svensson, J., Virkkula, A., and de Leeuw, G.: Soot on Snow experiment: bidirectional reflectance factor measurements of contaminated snow, *Cryosphere*, 9, 2323-2337, doi: 10.5194/tc-9-2323-2015, 2015.
- Prospero, J. M., Bullard, J. E., and Hodgkins, R.: High-Latitude Dust Over the North Atlantic: Inputs from Icelandic Proglacial Dust Storms, *Science*, 335, 1078-1082, doi: 10.1126/science.1217447, 2012.
- 975 Magnetite, Refractive index database, available at <https://refractiveindex.info/?shelf=main&book=Fe3O4&> (last access: 02 January 2020), 1985.
- Redmond, H. E., Dial, K. D., and Thompson, J. E.: Light scattering and absorption by wind blown dust: Theory, measurement, and recent data, *Aeolian Res.*, 2, 5-26, doi: 10.1016/j.aeolia.2009.09.002, 2010.
- 980 Reed, B. E., Peters, D. M., McPheat, R., and Grainger, R. G.: The Complex Refractive Index of Volcanic Ash Aerosol Retrieved From Spectral Mass Extinction, *J. Geophys. Res.-Atmos.*, 123, 1339-1350, doi: 10.1002/2017jd027362, 2018.

- Renard, J. B., Dulac, F., Berthet, G., Lurton, T., Vignelles, D., Jégou, F., Tonnelier, T., Jeannot, M., Couté, B., Akiki, R., Verdier, N., Mallet, M., Gensdarmes, F., Charpentier, P., Mesmin, S., Duverger, V., Dupont, J. C., Elias, T., Crenn, V., Sciare, J., Zieger, P., Salter, M., Roberts, T., Giacomoni, J., Gobbi, M., Hamonou, E., Olafsson, H., Dagsson-Waldhauserova, P., Camy-Peyret, C., Mazel, C., Décamps, T., Piringer, M., Surcin, J., and Daugeron, D.: LOAC: a small aerosol optical counter/sizer for ground-based and balloon measurements of the size distribution and nature of atmospheric particles – Part 1: Principle of measurements and instrument evaluation, *Atmos. Meas. Tech.*, 9, 1721-1742, doi: 10.5194/amt-9-1721-2016, 2016.
- 985 Richards-Thomas, T., McKenna-Neuman, C., and Power, I. M.: Particle-scale characterization of volcanoclastic dust sources within Iceland, *Sedimentology*, 68, 1137-1158, doi: 10.1111/sed.12821, 2020.
- 990 Sanchez-Marroquin, A., Arnalds, O., Baustian-Dorsi, K. J., Browse, J., Dagsson-Waldhauserova, P., Harrison, A. D., Maters, E. C., Pringle, K. J., Vergara-Temprado, J., Burke, I. T., McQuaid, J. B., Carslaw, K. S., and Murray, B. J.: Iceland is an episodic source of atmospheric ice-nucleating particles relevant for mixed-phase clouds, *Sci. Adv.*, 6, eaba8137, doi: 10.1126/sciadv.aba8137, 2020.
- 995 Schmale, J., Zieger, P., and Ekman, A. M. L.: Aerosols in current and future Arctic climate, *Nature Climate Change*, 11, 95-105, doi: 10.1038/s41558-020-00969-5, 2021.
- Shi, Y., Liu, X., Wu, M., Zhao, X., Ke, Z., and Brown, H.: Relative importance of high-latitude local and long-range-transported dust for Arctic ice-nucleating particles and impacts on Arctic mixed-phase clouds, *Atmos. Chem. Phys.*, 22, 2909-2935, doi: 10.5194/acp-22-2909-2022, 2022.
- 1000 Shi, Z., Krom, M. D., Bonneville, S., Baker, A. R., Bristow, C., Drake, N., Mann, G., Carslaw, K., McQuaid, J. B., Jickells, T., and Benning, L. G.: Influence of chemical weathering and aging of iron oxides on the potential iron solubility of Saharan dust during simulated atmospheric processing, *Global Biogeochem. Cy.*, 25, GB2010, doi: 10.1029/2010gb003837, 2011.
- Sinyuk, A., Torres, O., and Dubovik, O.: Combined use of satellite and surface observations to infer the imaginary part of refractive index of Saharan dust, *Geophys. Res. Lett.*, 30, doi: 10.1029/2002GL016189, 2003.
- 1005 Smith, R. J.: Use and misuse of the reduced major axis for line-fitting, *Am. J. Phys. Anthropol.*, 140, 476-486, doi: doi.org/10.1002/ajpa.21090, 2009.
- Sokolik, I. N., and Toon, O. B.: Incorporation of mineralogical composition into models of the radiative properties of mineral aerosol from UV to IR wavelengths, *J. Geophys. Res.-Atmos*, 104, 9423-9444, doi: 10.1029/1998jd200048, 1999.
- 1010 Song, Y. G., Chen, X. L., Qian, L. B., Li, C. X., Li, Y., Li, X. X., Chang, H., and An, Z. S.: Distribution and composition of loess sediments in the Ili Basin, Central Asia, *Quat. Int.*, 334, 61-73, doi: 10.1016/j.quaint.2013.12.053, 2014.
- Sumlin, B. J., Heinson, W. R., and Chakrabarty, R. K.: Retrieving the aerosol complex refractive index using PyMieScatt: A Mie computational package with visualization capabilities, *J Quant. Spectrosc. Radiat. Transf.*, 205, 127-134, doi: 10.1016/j.jqsrt.2017.10.012, 2018.
- 1015 Toledano, C., Bennouna, Y., Cachorro, V., de Galisteo, J. P. O., Stohl, A., Stebel, K., Kristiansen, N. I., Olmo, F. J., Lyamani, H., Obregon, M. A., Estelles, V., Wagner, F., Baldasano, J. M., Gonzalez-Castanedo, Y., Clarisse, L., and de Frutos, A. M.: Aerosol properties of the Eyjafjallajökull ash derived from sun photometer and satellite observations over the Iberian Peninsula, *Atmos. Environ.*, 48, 22-32, doi: 10.1016/j.atmosenv.2011.09.072, 2012.

- 1020 Vergara-Temprado, J., Miltenberger, A. K., Furtado, K., Grosvenor, D. P., Shipway, B. J., Hill, A. A., Wilkinson, J. M., Field, P. R., Murray, B. J., and Carslaw, K. S.: Strong control of Southern Ocean cloud reflectivity by ice-nucleating particles, *P. Natl. Acad. Sci. USA*, 115, 2687-2692, doi: 10.1073/pnas.1721627115, 2018.
- Volz, F. E.: Longwave refractive index of atmospheric aerosol substances, *Appl. Optics.*, 11, 755-759, 1972.
- von der Weiden, S. L., Drewnick, F., and Borrmann, S.: Particle Loss Calculator – a new software tool for the assessment of the performance of aerosol inlet systems, *Atmos. Meas. Tech.*, 2, 479-494, doi: 10.5194/amt-2-479-2009, 2009.
- 1025 Wang, J., Doussin, J. F., Perrier, S., Perraudin, E., Katrib, Y., Pangui, E., and Picquet-Varrault, B.: Design of a new multi-phase experimental simulation chamber for atmospheric photosmog, aerosol and cloud chemistry research, *Atmos. Meas. Tech.*, 4, 2465-2494, doi: 10.5194/amt-4-2465-2011, 2011.
- Weingartner, E., Saathoff, H., Schnaiter, M., Streit, N., Bitnar, B., and Baltensperger, U.: Absorption of light by soot particles: determination of the absorption coefficient by means of aethalometers, *J. Aerosol Sci.*, 34, 1445-1463, doi: 10.1016/S0021-8502(03)00359-8, 2003.
- 1030 Weinzierl, B., Sauer, D., Minikin, A., Reitebuch, O., Dahlkötter, F., Mayer, B., Emde, C., Tegen, I., Gasteiger, J., Petzold, A., Veira, A., Kueppers, U., and Schumann, U.: On the visibility of airborne volcanic ash and mineral dust from the pilot's perspective in flight, *Phys. Chem. Earth*, 45-46, 87-102, doi: 10.1016/j.pce.2012.04.003, 2012.
- Wendisch, M., and Brenguier, J.-L.: *Airborne Measurements for Environmental Research : Methods and Instruments*, John Wiley & Sons, Incorporated, Somerset, GERMANY, 2013.
- 1035 Wittmann, M., Zwaafink, C. D. G., Schmidt, L. S., Gudmundsson, S., Palsson, F., Arnalds, O., Björnsson, H., Thorsteinsson, T., and Stohl, A.: Impact of dust deposition on the albedo of Vatnajökull ice cap, Iceland, *Cryosphere*, 11, 741-754, doi: 10.5194/tc-11-741-2017, 2017.
- 1040 Woods, T. N., Chamberlin, P. C., Harder, J. W., Hock, R. A., Snow, M., Eparvier, F. G., Fontenla, J., McClintock, W. E., and Richard, E. C.: Solar Irradiance Reference Spectra (SIRS) for the 2008 Whole Heliosphere Interval (WHI), *Geophys. Res. Lett.*, 36, doi: 10.1029/2008GL036373, 2009.
- Zubko, N., Munoz, O., Zubko, E., Gritsevich, M., Escobar-Cerezo, J., Berg, M. J., and Peltoniemi, J.: Light scattering from volcanic-sand particles in deposited and aerosol form, *Atmos. Environ.*, 215, 116813, doi: 10.1016/j.atmosenv.2019.06.051, 2019.