

Observations-Extensive coverage of ultrathin Tropical Tropopause Layer cirrus clouds from a revealed by balloon-borne lidar observations

Thomas Lesigne¹, François Ravetta¹, Aurélien Podglajen², Vincent Mariage¹, and Jacques Pelon¹

¹Laboratoire Atmosphères, Observations Spatiales (LATMOS/IPSL), Sorbonne Université, UVSQ, CNRS, Paris, France

²Laboratoire de Météorologie Dynamique (LMD/IPSL), École Polytechnique, Institut Polytechnique de Paris, Sorbonne Université, École Normale Supérieure, PSL Research University, CNRS, Paris, France

Correspondence: Thomas Lesigne (thomas.lesigne@latmos.ipsl.fr)

Abstract.

Tropical Tropopause Layer (TTL) clouds have a significant impact on the Earth's radiative budget and regulate the amount of water vapor entering the stratosphere. Estimating the total coverage of tropical cirrus clouds is challenging, since the range of their optical depth spans several order of magnitude, from thick opaque cirrus detrained from convection to sub-visible clouds just below the stratosphere. During the Strateole-2 observation campaign, three microlidars were flown onboard stratospheric superpressure balloons from October 2021 to late January 2022, slowly drifting only a few kilometers above the TTL. These measurements have unprecedented sensitivity to thin cirrus and provide a fine-scale description of cloudy structures both in time and space. Case studies of collocated observations with the space-borne lidar Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) show a very good agreement between the instruments and highlight ~~the unique ability of the microlidar~~ BeCOOL's higher detection sensitivity. Indeed, the microlidar is able to detect optically very thin clouds ~~below CALIOP detection capacity~~ (optical depth $\tau < 2 \cdot 10^{-3}$) that are undetected by CALIOP. Statistics on cloud occurrence show that TTL cirrus appear in ~~more than about~~ 50 % of the microlidar profiles and have a mean geometrical depth of 1 km. Ultrathin TTL cirrus ($\tau < 2 \cdot 10^{-3}$) have a significant coverage (~~+623~~ % of the profiles) and their mean geometrical depth is ~~below 500-m~~ 0.5 km.

15 1 Introduction

In the ~~Tropiestropics~~, the transition between troposphere and stratosphere occurs in a vertically extended layer (14 to 18.5 km) sharing characteristics from both troposphere and stratosphere: the Tropical Tropopause Layer (TTL, Fueglistaler et al. (2009); Randel and Jensen (2013)). Most of the air entering the stratosphere makes its way through the TTL along the ascending branch of the Brewer-Dobson circulation (Brewer, 1949). TTL is then often referred to as the "gate to the stratosphere". On their way
20 up, air masses encounter extremely low temperatures at the cold point tropopause (CPT, ~ 17 km, ~ 190 K) that freeze-dry a great part of their water content and are ultimately responsible for the dryness of the lower stratosphere (Holton et al., 1995). Although water vapor concentration in the stratosphere is very low (~ 5 ppmv), it has a significant radiative impact on the

whole climate system (Solomon et al., 2010) and plays a major role in stratospheric chemistry (Fueglistaler et al., 2009). Yet its evolution is not accurately represented in today's climate models. It is thus necessary to get a better understanding of the various TTL processes (transport, dynamical, radiative and microphysical processes) modulating the amount of water vapor and other trace gases that eventually reaches the stratosphere.

At the heart of the interplay between those processes, TTL clouds have been subject to numerous studies in the past decades. Thanks to their high vertical resolution and unique sensitivity to tenuous clouds, lidar observations have long been used to characterize tropical clouds, operated from the ground (e.g., Platt et al., 1984, 1987), or from research vessels (e.g., Fujiwara et al., 2009), but their spatial coverage is limited and they suffer from being potentially blinded by opaque clouds between the ground and the upper troposphere. Passive space-borne observations (either radiometers (e.g., Prabhakara et al., 1988) or solar occultation measurements (e.g., Wang et al., 1994)) have broaden the picture, providing almost global observations, but they lack sensitivity and resolution to fully capture the TTL cloud coverage. Since the pioneer mission Lidar In-space Technology Experiment (LITE, Winker and Trepte, 1998), space-borne lidars have overcome these limitations. For the past 17 years, the Cloud and Aerosol Lidar with Orthogonal Polarisation (CALIOP) onboard the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) provided continuous observations that fed a great deal of cloud studies (Yang et al., 2010; Martins et al., 2011; Iwasaki et al., 2015)(Yang et al., 2010; Martins et al., 2011; Iwasaki et al., 2015; Sourdeval et al., 2015). This mission recently ended on August 1, 2023. CALIOP's cloud observations have intensively been evaluated against other types of measurements, from ground-base lidars (Thorsen et al., 2011) to geostationary weather satellite (Sèze et al., 2015). Recent airborne campaigns such as the NASA Airborne Tropical Tropopause EXperiment (ATTREX, Jensen et al., 2017) have enabled the in situ characterization of thin TTL cirrus (Krämer et al., 2020). A noteworthy result from aircraft data was the characterization of a systematic relationship between TTL clouds and equatorial and gravity waves (Kim et al., 2016). This finding was later confirmed with space-borne (Chang and L'Ecuyer, 2020) and more recently with balloon-borne observations (Bramberger et al., 2022).

Long-duration stratospheric balloon constitute an invaluable platform to better characterize clouds distribution. Since the balloon is slowly drifting with the air, it is able to capture the fine scale spatial variability of the underlying cloud scene. Here, we introduce the first observations from the Balloon-borne Cirrus and convective overshoot Lidar (BeCOOL, ?). BeCOOL nadir-looking lidar has a viewing geometry comparable to CALIOP, but benefits from a significantly higher signal to noise ratio (SNR) in the TTL and upper troposphere thanks to the long integration time allowed by the low speed of the balloon and the small distance to the observed clouds. BeCOOL was recently flown for the first time onboard three superpressure balloons (SPBs) in the framework of the StratoClim-2 project (Haase et al., 2018; Corcos et al., 2021; Bramberger et al., 2022). The SPBs were launched from Seychelles Island and travelled up to the middle of the pacific ocean at about 20.5 km (50 hPa) between October 2021 and January 2022, gathering 700 nighttime hours of high-resolution lidar profiles.

The article is organised as follow: Section 2 presents the different data sets and the cloud classification. In Sect. 3, three case studies of collocated BeCOOL/CALIOP observations are analyzed to contrast the two instruments, their sampling and detection capability. Section 4 is dedicated to a statistical description of the balloon-borne cloud data and a statistical comparison with space-borne lidar data. Conclusions and perspectives are in section 5.

2 Lidar Data sets

2.1 Balloon-borne lidar data

60 BeCOOL nighttime observations have been gathered during the first Strateole-2 ~~fiels-field~~ campaign, from 20 October 2021 to 26 January 2022. ~~Three balloons carrying the lidar~~ Among the 17 balloons released during the campaign, 3 were carrying BeCOOL microlidar. They were successively launched from Seychelles Islands and drifted eastward at about ~~20~~20.5 km of altitude near the equator, their trajectories are shown on Fig 1.a. The main characteristics of the flights are presented Table 1 and a summary of the technical specifications of BeCOOL microlidar are presented in Table 2. ~~A comprehensive description of the microlidar and the processing of Level-~~

65 Overall, 38631 lidar profiles have been measured at the native 1 minute resolution. Each raw lidar profile is corrected from the radiometric background derived from the tail of the signal, then multiplied by the squared range to account for the geometrical dilution of the laser power. For each BeCOOL instrument, an empirical correction of the lidar overlap function has been constructed from a set of clear sky observations and synthetic pure molecular lidar profiles derived from ERA-5 meteorological
70 fields. Every single lidar profile is first corrected from this overlap effect then normalized upon ERA-5 using a 1 km thick clear sky layer above the uppermost clouds, between 1 and ~~Level-2 products (respectively attenuated backscatter profiles and the retrieved geometrical-~~ km below the lidar, assuming that pure molecular signal prevails in this normalization interval. These normalized lidar profiles are usually called “attenuated backscatter” and referred to as Level 1.

Geometrical and optical properties of ~~cloud and aerosol layers) can be found in Ravetta et al. (2023).~~ The clouds (Level
75 2 data set has been-) are retrieved after averaging ~~10 consecutive 1-minute profiles in order~~ consecutive profiles over 10 minutes to improve the SNR ~~thus the detection of optically thin clouds-~~. Detection of clouds is semi-automated, every profile has been hand-checked after a first round of automated detection. Cloud optical depths are then retrieved using the classical Fernald/Klett inversion method (Fernald et al., 1972; Klett, 1981) which relies on the particular extinction-to-backscatter ratio (lidar ratio). Quite similarly to what is done for CALIOP, this quantity can be iteratively constrained for the thicker semi-transparent clouds
80 that significantly attenuate the lidar beam. Those constrained retrievals need to be further corrected from multiple scattering (Platt, 1973). BeCOOL’s multiple scattering correction factor $\eta = 0.88$ has been determined as the appropriate scaling factor allowing to reconcile BeCOOL’s “apparent” and CALIOP’s “true” optical depth distributions for constrained retrievals. Optical depth of the thinner clouds are retrieved using *a priori* lidar ratio chosen as the most frequent value from CALIOP constrained retrieval over the same period and region at the altitude of the middle of the cloud. The estimated relative uncertainty of
85 these retrievals increases when the optical depth decreases. It is below 10 % for clouds with an optical depth larger than 0.1, below 50 % for clouds with an optical depth larger than 0.01, and much larger (up to 90 %) for the optically thinnest clouds. A comprehensive description of the instrument and the different levels of processing, along with an analysis of their uncertainties, will be detailed in an other article.

Figure 1 shows the trajectories of the three flights and the lidar curtains (time vs altitude) of attenuated backscatter and
90 reveals a large variety of different cloud scenes. Intense surface echos (i.e., the ocean or exceptionally the land surface) are seen in 86 % of the profiles. The lidar beam is fully attenuated by opaque clouds otherwise. In profiles reaching the surface, the

Table 1. Main characteristics of the three Strateole-2 flights carrying BeCOOL microlidar; \bar{z} is the mean altitude above sea level and $|\bar{\mathbf{u}}|$ the mean ground speed of the balloon.

Flight	Strateole-2 ID	Launch Date	End Date	\bar{z}	$ \bar{\mathbf{u}} $	Number of 1 minute profiles
1	ST2_C1_02_STR1	2021-10-20	2021-11-01	20.2 km	$11.2 \text{ m} \cdot \text{s}^{-1}$	3573 <u>3542</u>
2	ST2_C1_08_STR1	2021-11-05	2021-12-29	20.3 km	$7.2 \text{ m} \cdot \text{s}^{-1}$	156171 <u>15468</u>
3	ST2_C1_13_STR1	2021-11-15	2022-01-25	20.3 km	$6.4 \text{ m} \cdot \text{s}^{-1}$	19742 <u>19621</u>

ubiquitous aerosol-rich boundary layer generally occupies the lowest 2.5 km along with frequently occurring low-level clouds (cumulus and stratocumulus). Geometrically thin (a few hundred meters), horizontally extensive mid-level clouds are often found above, below 10 km, mainly between 5 and 8 km; they typically have large backscatter and are likely pure liquid or mixed-phase clouds. Above 10 km are pure ice clouds, cirrus and deep convective clouds. The clouds' vertical structure can be fully resolved up to an optical depth $\tau_{max} \simeq 3$, a threshold value depending on the energetic conditions and optical efficiency of the instrument which both vary with thermal conditions onboard the gondola. Clouds thicker than this appear opaque (the lidar beam is fully attenuated before reaching the bottom of the cloud) and only their upper part can be resolved. Typically, deep convective clouds have a large vertical extent that cannot be accurately inferred from BeCOOL observations.

For the purpose of this study, the area covered by the balloons has been zonally divided into three regions: Indian Ocean (55° E to 95° E), Maritime Continent (95° E to 165° E) and Central Pacific Ocean (165° E to 230° E). There is a striking contrast between very cloudy profiles over Indian Ocean and Maritime Continent, with frequent deep convection, and clear-sky conditions over the Central Pacific Ocean, in the second part of flights 2 and 3.

We built a classification of cloud profiles for the BeCOOL dataset. Clouds are first classified using a set of threshold values on their top and base altitude. ~~Those altitudes have first been determined to retrieve cloud's optical properties where the lidar signal departs from the molecular background signal. For the cloud classification, in order to focus on the main part of clouds, top and base altitudes are slightly modified so that 15 % of the cloud optical depth lay above the new top altitude, and 15 % below the new base altitude.~~ Cirrus clouds are here defined as all clouds with a base altitude lying above 10 km (i.e., temperatures below the glaciation threshold of supercooled droplets at about 240 K), then sub-classed as TTL cirrus if their base altitude is over 14 km. Convective clouds are here defined as opaque clouds (totally attenuating the lidar beam) with top altitude laying above 10 km. Mid-level clouds have a top altitude between 5 and 10 km. A last class gathers clouds that do not fit previous requirements, with top altitude above 10 km and base altitude below, sharing characteristics from both cirrus and mid-level clouds. This classification is somewhat restrictive in the case of deep convection, since mostly the core of convective cells will be flagged in this category, while a large part of the convective anvils will be classified as cirrus as long as BeCOOL's lidar beam goes through and the cloud base is above 10 km.

A profile classification has been built from this cloud classification. "Clear sky" is defined as profiles with no detected cloud above 5 km, since low level clouds and the planetary boundary layer are not considered in this study. "Deep convection" gathers profiles exhibiting any convective clouds, regardless of the presence of cirrus on top of it. "Cirrus only" and "Mid-level clouds




Figure 1. **a:** trajectories of the three balloons carrying BeCOOL during the first Strateole-2 scientific campaign, dashed white boxes show the three studied regions; **b-d:** lidar curtains (time vs altitude, attenuated backscatter) for the three flights, concatenated nights of observation (apart from thin vertical black lines, daytime has been removed for the sake of readability). Overflow regions are color-coded on top of the curtains. Red numbers (1, 2 and 3) highlight the three case studies presented in section 3.

only" stand for profiles where only such type of clouds are detected above 5 km. A last class, "Mixed multilayered scenes" gathers the other profiles, usually exhibiting complex overlay of cirrus and mid-level clouds.

Further classification of cirrus layers is performed based on their optical depth: thin cirrus have an optical depth below 0.1, which is about the detection lower bound for passive radiometer (McFarquhar et al., 2000), sub-visible clouds have an optical depth $\tau < 3 \cdot 10^{-2}$, a classical value from Sassen et al. (1989), and ultrathin cirrus have an optical depth $\tau < 2 \cdot 10^{-3}$ (which is about the detection threshold of CALIOP, see Sect. 4).

125 2.2 Space-borne lidar data

BeCOOL is compared to CALIOP space-borne lidar using the Level 2 Cloud and Aerosols merged product with a 5 km horizontal resolution, version 4.21 (Young et al., 2018). This data set reports optical and geometrical properties of detected clouds or aerosol layers along the satellite track. When CALIOP lidar curtains are displayed, the figures are generated using

the Level 1 version 4.11 attenuated backscatter product (Kar et al., 2018). In this study, only the 532 nm channel is used. The
130 main technical specifications of CALIOP lidar, along with BeCOOL's, are presented in Table 2.

While CALIOP flies at $\sim 8 \text{ km} \cdot \text{s}^{-1}$, achieving its native 1/3 km horizontal resolution from a single lidar shot, BeCOOL flies
one thousand time slower, which allow to integrate individual lidar shots for a whole minute, considerably enhancing the SNR.
This speed difference also implies that CALIOP provides an almost instantaneous description of cloudy structures at synoptic
scale, while temporal and spatial evolution of the underlying scene are entangled in BeCOOL's observations. BeCOOL's laser
135 divergence (667 μrad) is significantly higher than CALIOP's (100 μrad), meaning that BeCOOL's high SNR in near field
decreases toward the surface due to geometric power dilution, whereas this effect can be neglected for CALIOP.

Following Reagan et al. (2002), we assume that the optical depths retrieved at ~~808802~~ nm (for BeCOOL) and 532 nm (for
CALIOP) wavelengths are comparable, i.e. that the scattering particles are larger than 5 – 8 μm such that there is only weak
wavelength dependency of Mie scattering.

140 During the campaign, CALIOP was crossing the equator around 2:30 local time. Originally crossing the equator at 1:30
local time as part of the Afternoon-Constellation (A-train), CALIPSO was moved to a lower orbit in 2018 to join Cloudsat
(Braun et al., 2019). As CALIPSO's fuel reserves were coming to an end, the satellite has been experiencing an orbital drift
which explains the 2:30 AM crossing time during the campaign instead of the usual 1:30 AM.

3 Case studies of BeCOOL/CALIOP collocated observations

145 Three case studies of collocated BeCOOL/CALIOP measurements are now presented in order to compare the two instruments
at coincidence time and highlight their complementarity due to the fundamental differences mentioned in the previous section.
The case studies correspond to different cloud scenes: a thick (anvil) cirrus, a thin cirrus and deep convection. To contextualize
the cloud scene around lidar observations, we use the NOAA/NCEP GPM_MERGIR brightness temperature data in the atmo-
spheric window ($\sim 11 \mu\text{m}$). This product combines observations from 4 geostationary satellites and provides a global coverage
150 with spatial resolution of 4 km and temporal resolution of 30 min (Janowiak et al., 2001).

Figures 2, 3 and 4 present lidar curtains from the two instruments along with the GPM-MERGIR brightness temperature map
closest to the coincidence time, with a size of $5^\circ \times 5^\circ$. ~~In order to display lidar curtains from both instruments at a comparable
resolution,~~ CALIOP's ~~horizontal resolution as been downgraded to 1 curtain resolution~~ below 8.2 km a.s.l. ~~has been degraded
to 1 km horizontally and 60 m vertically,~~ which is the native resolution above ~~-.The step from 30 to 60 m vertical resolution
at 8.2 km a.s.l. is still visible on CALIOP's curtains.~~ BeCOOL's curtains are displayed for the whole nights (roughly 11 h of
155 observations), starting around 18:00 local time and ending around 5:30 local time, covering an horizontal distance of 200 to
500 km along the balloon's track depending on the wind speed. Each CALIOP's curtain is 570 km long (dashed line on the
maps), covered in about 82 s.

As previously stated, CALIOP's observations are almost instantaneous and can be compared with a single brightness tem-
160 perature map from GMP-MERGIR, latitude appearing then as a natural coordinate for CALIOP, while time stays the natural

Table 2. Overview of the main characteristics of BeCOOL and CALIOP lidars.

	BeCOOL	CALIOP
wavelength <u>wavelength</u>	808 <u>802</u> nm	532 nm*
pulse repetition rate	4700 Hz	20 Hz
pulse energy	10 μ J	110 mJ
depolarization channel	no	yes
altitude	20 km	700 km
ground speed	0 to 25 $\text{m} \cdot \text{s}^{-1}$ (mean of 7 $\text{m} \cdot \text{s}^{-1}$)	8 $\text{km} \cdot \text{s}^{-1}$
horizontal resolution Level 1	0 to 1.5 km (mean of 420 m)	333 m
number of lidar shots	$\sim 3 \cdot 10^5$	1
temporal resolution	1 min	5 ms
horizontal resolution Level 2	0 to 15 km (mean of 4.2 km)	5 km
number of lidar shots	$\sim 3 \cdot 10^6$	15
temporal resolution	10 min	0.7 s
vertical sampling	15 m	30 m below 8.2 km a.s.l. 60 m above 8.2 km a.s.l.
laser beam divergence	667 μ rad	100 μ rad
diameter of the illuminated spot		
17 km a.s.l.	2 m	70 m
surface	14 m	70 m

*CALIOP's 1064 nm channel is not used in this study

coordinate for BeCOOL, which can only be compared with successive brightness temperature maps. Hourly maps for the three case studies are presented in the appendix, Fig. B1, B2 and B3.

3.1 First case study: thick cirrus cloud

Figure 2 shows an excellent coincidence that happened on 29 November 2021 over the Pacific Ocean ($\sim 4^\circ$ S, 172° E) for the second BeCOOL flight. The satellite track crossed the balloon track less than 1 km away from it. The balloon covered 430 km during this night. There is a perfect agreement between the two lidars at the coincidence time: they both capture a thick cirrus cloud extending from 12 to 16 km over two mid-level clouds with very small vertical extent around 5 and 7 km. Both lidars' profiles around coincidence time are displayed in appendix on Fig. A1. CALIOP's curtain shows that ~~at coincidence time~~ this thick cirrus is embedded in a larger scale thinner laminar cirrus extending vertically from 14 to 16 km and horizontally all along the 570 km track displayed here. The brightness temperature map at 15:00 UTC reveals the horizontal structure of this thick cirrus, centered on the coincidence spot and with an apparent radius of ~ 100 km. BeCOOL's curtain and the hourly brightness temperatures maps on Fig. B1 allow to follow the temporal evolution of the scene under the balloon: from the beginning of the




Figure 2. First case study: thick cirrus cloud, 2021-11-29. **a:** 11 μm brightness temperature map at 15:00 UTC; **b:** BeCOOL L1 curtain (along the solid line on the map); **c:** CALIOP L1 curtain (along the dashed line on the map); **d, e:** time series of brightness temperature under the balloon and the satellite; **f, g:** time series of optical depth τ above 10 km retrieved from BeCOOL and CALIOP.

night up to 13:00 UTC, a thin and laminar cirrus vertically extending between 15 and 17 km, with an optical depth of $0.2\text{--}0.08$ is overflown, then this cirrus ~~thicken~~ ~~thickens~~ to extend vertically from 12 to 16 km, reaching an optical depth of $0.7\text{--}0.8\text{--}0.5$.
 175 The balloon follows the thick cloud for the second part of the night as they are both advected eastward.

Brightness temperature (BT) below both instruments (Fig. 2.d-e) exhibits high values (almost 300 K) above the thin part of the cloud, between 7:00 and 9:00 UT on Fig. 2.b and between 2 and 3° S on Fig. 2.c. BT drops down to 280 K above the thicker part of the cloud, after 13:00 UT on Fig. 2.b and around 4° S on Fig. 2.c. Cloud's contribution to upward thermal flux increases with optical depth, actually lowering the flux and revealing the thermal contrast between low temperatures at cloud
 180 level and higher temperatures below. BT (Fig. 2.d-e) and total cloud optical depth above 10 km (Fig. 2.f-g) are thus quite anti-correlated: $r = -0.88$ along BeCOOL's track, $r = -0.72$ along CALIOP's. These correlations would be more significant without the presence of mid-level clouds around 5 and 7 km, that are not accounted for in the total cloud optical depth above
 above-10 km but further lower the BT (e.g., Fig. 2.e at 6° S).

At coincidence, the retrieved cirrus' optical depths are $0.6\text{--}0.32$ for BeCOOL and 0.4 for CALIOP. ~~Beyond the optical depth's~~
 185 ~~spectral dependency~~ 0.37 for CALIOP, which is ~~expected to be a second-order effect, there is about 20%~~ of uncertainties in both

retrieval, a quite good agreement. This discrepancy is well within the range of uncertainties and mainly related to the assumed extinction-to-backscatter ratio (lidar ratio) and multiple scattering effects (Ravetta et al., 2023). In strong spatial variability of this thick cirrus case, multiple scattering is the main source of uncertainty. Thus the two retrieved optical depth values can be considered in a fair agreement which can be seen on both lidar curtains and on the lidar profiles displayed on Fig. A1.

190 3.2 Second case study: thin cirrus clouds




Figure 3. Second case study: thin cirrus cloud, 2021-11-24. **a**: 11 μm brightness temperature map at 20:00 UTC; **b**: BeCOOL L1 curtain (along the solid line on the map); **c**: CALIOP L1 curtain (along the dashed line on the map); **d**, **e**: time series of brightness temperature under the balloon and the satellite; **f**, **g**: time series of optical depth τ above 16 km retrieved from BeCOOL and CALIOP. CALIOP L2 operational algorithm partially detects the thin cirrus (white box on **c**, red segment on **g**) after a 80 km horizontal averaging and reports a single optical depth value for this 80 km leg around coincidence time: at this resolution, a single point is detected as cloudy, CALIOP algorithm is missing most of the cloud. Manual retrievals of cloud optical depth using the same lidar ratio for both instrument (blue triangles on **f** and **g**) show an excellent agreement.

The second case study (Fig. 3) happened for the third BeCOOL flight off the east coast of Sumatra island, Indonesia ($\sim 3^\circ\text{N}$, 97°E ; distance at coincidence: 10.4 km) and corresponds to collocated observations of a very thin TTL cirrus, which is only partially reported in CALIOP Level 2 data. BeCOOL curtain on Fig. 3 reveals clearly a thin cloudy layer above 17.5 km,

195 first fading out from the beginning of the night until 14:00 then appearing back from 15:30 UTC and slowly thickening until
20:00 UTC reaching up to ~~about 10^{-2}~~ $5 \cdot 10^{-3}$ optical depth. This horizontally homogeneous, geometrically and optically very
thin cirrus layer appears to fit in the description of Ultrathin Tropical Tropopause Clouds (UTTCs) reported by Peter et al.
(2003). In CALIOP's curtain, this cloud can be identified by the human eye around 17.5 km in the 532 nm total attenuated
backscatter (Fig. 3.c). However, it is only reported in CALIOP L2 for about 10 s around coincidence. It was detected after
an horizontal averaging of 80 km, the last step of the algorithm designed to improve SNR in order to detect tenuous features,
200 but its horizontal extent could likely have been better constrained with even more extensive horizontal averaging. Given the
limitation of CALIOP L2 algorithm for such a case and for the sake of a fair comparison of the instrument capabilities, we
manually retrieved the UTCC optical depth from CALIOP L1. We first improve the SNR applying an horizontal rolling mean
over a 80 km window. Then, we retrieve the cirrus optical depth at three latitudes: 4.5°, 3° and 1.5° N, keeping the same lidar
ratio (21.8 sr) and multiple scattering factor η (0.77) as reported in CALIOP L2 for the central part of this cloud. This cirrus'
205 optical depth is $8.3 \cdot 10^{-4}$ at 5° N, increasing to $4.0 \cdot 10^{-3}$ at 3° N (coincidence) then decreasing to $2.2 \cdot 10^{-3}$ at 1.5° N. At
coincidence, ~~using the same assumed lidar ratio value for BeCOOL but keeping its own multiple scattering factor η of 0.5 gives
an new optical depth of $3.8 \cdot 10^{-3}$. Assuming the same lidar ratio, the optical depths at coincidence~~ the retrieved optical depths
from both observations are thus in excellent agreement. ~~The difference between the optical depth of 10^{-2} reported in BeCOOL
Level-2 at coincidence and the new optical depth value of $3.8 \cdot 10^{-3}$ can fully be explained by the difference in assumed lidar
ratio (48.9 vs 21.8). Furthermore, this example shows that the optical depth of thinnest clouds could be largely overestimated
by BeCOOL retrieval due to an inaccurate lidar ratio choice. This key parameter can only be constrained for thicker clouds
that significantly attenuate the lidar beam (optical depth $\gtrsim 10^{-2}$), but an *a priori* lidar ratio value has always to be assumed for
thinnest features such as this cirrus. Yet, in most cases, we expect an inaccurate choice of lidar ratio to only result in a factor
 ~ 2 difference in τ between CALIOP and our retrieval.~~

215 We can attempt to estimate the horizontal extension of this UTTC assuming ~~an horizontal extension of~~ that it expands a few
hundreds of kilometers along both instrument's track: this cirrus could have an area greater than 10^5 km², which is the order
of magnitude observed by Peter et al. (2003) for UTTCs.

Regarding backscattered power, the contrast between ~~the this~~ cloud and the surrounding clear sky is ~~higher at 808~~ about 3
times higher at 802 than 532 nm (due to the strong wavelength dependency of Rayleigh scattering, emphasized by Peter et al.
220 (2003)). This is why, in addition to a lower absolute noise level, very thin features are more easily detected with BeCOOL. Such
thin layers are sometimes clearer in the depolarization ratio (not shown here) and should have a stronger signature in CALIOP's
second channel (1064 nm) but ~~this channel is unfortunately too noisy, and~~ the operational layer detection algorithm only relies
for now on the 532 nm channel. Further reprocessing of CALIOP's observations are expected to improve the detection and
retrieval of very thin clouds. Vaillant de Guélis et al. (2021) recently introduced a new 2-dimensional multi-channel cloud
225 detection algorithm for CALIOP. Preliminary tests on collocated BeCOOL/CALIOP's observations over very thin clouds show
large improvements in cloud layer detection. Other projects of CALIOP reprocessing rely on machine learning techniques to
detect optically thin clouds (Wang et al., 2019).

3.3 Third case study: convective clouds




Figure 4. Third case study: convective cloud, 2021-11-27. **a:** 11 μm brightness temperature map at 18:30 UTC; **b:** BeCOOL L1 curtain (along the solid line on the map); **c:** CALIOP L1 curtain (along the dashed line on the map); **d, e:** time series of brightness temperature under the balloon and the satellite. White capital letters on **a** and **c** show two overshoots detected by CALIOP.

Figure 4 displays a CALIOP-BeCOOL coincidence which ~~occured~~ occurred on 27 November 2021 off the southeast coast of Borneo island, Indonesia ($\sim 4^\circ\text{S}$, 117°E ; distance at coincidence: 72.7 km) for the third BeCOOL flight. CALIOP overflew several convective cells, capturing two events of convection overshooting the main cloud top (white capital letters on Fig. 4.a-c). The first one (A) has an apparent diameter of 40 km and seems to be fading out at the time of overpass, having appeared 1 to 2 h before (see the hourly maps on Fig. B2). The second one (B) seems to be popping up right under the satellite and has an apparent diameter of 15 km. The core of this cell is characterized by a very strong backscatter and a small penetration depth: the highest part of this cloud is very dense and optically thick. Those two structures clearly overshoot the extensive 17 km cloud top which appears on both curtains and is likely a high-altitude anvil. As shown by the brightness temperature maps, BeCOOL flew all night long around the convective cells, measuring the edges of anvils and revealing the evolution of the multilayered cloud structure. No clear sign of overshoot residual (i.e., cloud above the extensive 17 km deck) appears in BeCOOL data.

240 There could be different reasons for this apparent disagreement in overshoot detection. First, it is worth mentioning that "overshoots" would have different visual aspects in CALIOP and BeCOOL curtains. Assuming a wind difference of $7.5 \text{ m} \cdot \text{s}^{-1}$ between the balloon and the cloud top (mean value over the Maritime Continent along the three flights, according to ERA5 reanalysis), an overshoot with a size of 40 km would be overflowed for about 1.5 hours, a duration comparable to its lifetime (Dauhut et al., 2018; Lee et al., 2019). Thus, it is likely that overshoots in BeCOOL curtains will exhibit a different shape (aspect ratio, ...) compared to CALIOP's, and cannot be identified as clearly. Over the whole campaign, we found no obvious observation in BeCOOL's data of an "overshoot" similar to the protrusion detected by CALIOP in this example. Since convective overshoots are one of the scientific targets of BeCOOL, we investigated further the probability of overflying very deep convection with the balloons using $11 \mu\text{m}$ Brightness Temperature (BT) data. We defined a $T_{\text{overshoot}} = 200 \text{ K}$ BT threshold as a proxy for potentially overshooting convection, based on the comparison between CALIOP and BT maps in this case study.

245 Over the Maritime ~~continent~~ Continent and during the campaign, about 1 % of the pixels ~~have BT of the BT maps have values~~ lower than $T_{\text{overshoot}}$, whereas such cold pixels are 3 times less likely to occur along the balloon tracks (0.3 % of the observations over this area). Such low frequency of observation of "cold" cloud scenes constitutes a "warm" sampling bias for our 3 flights. Nevertheless, extending this analysis to all 17 Strateole-2 campaign balloons, we did not find conclusive evidence of a systematic sampling bias, which tends to discard the hypothesis that a dynamical effect (such as flow divergence) prevents

250 the balloon from flying over overshooting tops. Targeting such relatively rare, sparse and small-scale structures with a limited instrumented fleet may require the use of steerable balloons.

4 Statistical description

4.1 Cloud coverage and scene complexity

Table 3. Main-BeCOOL main profile classification, percentages of 10 minutes averaged profiles. Details on this classification can be found in Sect. 2.1.

	Full Area	Indian Ocean	Maritime Continent	Central Pacific Ocean
Longitude boundaries	55 to 230 °E	55 to 95 °E	95 to 165 °E	165 to 230 °E
Number of 10-min profiles	4050 <u>3878</u>	649 <u>636</u>	1091 <u>1052</u>	2310 <u>2190</u>
Clear sky	8 <u>7</u> %	6 <u>5</u> %	0.2 %	12 <u>10</u> %
Deep convection	5 <u>6</u> %	7 <u>8</u> %	13 <u>15</u> %	1 <u>0.5</u> %
Cirrus only (<i>TTL cirrus only</i>)	57 <u>52</u> % (43 <u>37</u> %)	40 <u>33</u> % (15 <u>11</u> %)	36 <u>30</u> % (21 <u>15</u> %)	71 <u>69</u> % (60 <u>55</u> %)
Mid-level cloud only	3 %	4 <u>2</u> %	0.6 <u>0.2</u> %	4 %
Mixed multilayered scene	28 <u>33</u> %	43 <u>52</u> %	50 <u>55</u> %	13 <u>16</u> %

A summary of BeCOOL profile classification over the Maritime Continent and Central Pacific Ocean is provided in Table 3.

260 A striking contrast appears between the two regions: over the Maritime Continent, convection is detected in up to ~~13~~15 % of

the profiles and clear-sky scenes are almost absent (0.2 %). On the contrary, more frequent clear sky profiles (~~12~~10 %) and far less convective ones (~~0.60~~5 %) are found over the Central Pacific. Over the Central Pacific, ~~71~~69 % of the profiles present only cirrus, ~~60~~55 % only TTL cirrus. Over the Maritime ~~continent~~, ~~Continent, more than~~ half of the profiles correspond to a complex combination of different types of clouds, here reported as "Mixed multilayered scenes", while we only report ~~13~~16 % of such profiles over the Central Pacific Ocean. Although several types of scenes are gathered in this "mixed multilayered" class, a great part of them could be somehow related to different stages of convective activity: developing convection, detrainment and/or precipitation. The campaign took place during La Niña phase of the El Niño Southern Oscillation (soi), the strong contrast in convective activity between the Maritime Continent and Central Pacific Ocean is typical of this ENSO phase (e.g. Gage and Reid, 1987).

Table 4. Frequency of occurrence of cirrus (cloud base > 10 km) in BeCOOL profiles with different thresholds on optical depth, percentage of 10-minute profiles. Bold font stands for TTL cirrus (cloud base > 14 km).


	Full Area		Indian Ocean		Maritime Continent		Central Pacific Ocean	
All cirrus	87 73 %	59 48 %	83 61 %	39 24 %	94 64 %	47 29 %	84 81 %	71 65 %
$\tau > 2 \cdot 10^{-3}$	75 58 %	43 29 %	71 48 %	16 4 %	92 60 %	39 19 %	68 61 %	53 42 %
$\tau > 3 \cdot 10^{-2}$	59 42 %	25 14 %	64 42 %	7 1 %	84 51 %	28 11 %	46 38 %	28 19 %
$\tau > 0.1$	48 25 %	15 3 %	57 30 %	5 0.2 %	77 37 %	21 3 %	32 18 %	15 4 %

Table 4 summarizes the occurrence of cirrus and TTL cirrus for several optical depth thresholds. Regardless of their optical depth, cirrus are detected in ~~87~~73 % of all profiles with a ~~weak~~ ~~small~~ regional contrast: from ~~83~~61 % over Indian Ocean to ~~94~~81 % over the Maritime Continent. The regional contrast is more pronounced for TTL cirrus, which are detected in ~~39~~24 % of the profiles over the ~~Maritime Continent and 71~~Indian Ocean and ~~65~~ % over the Central Pacific Ocean. The thresholds on optical depth show what would be detected by less sensitive instrument: CALIOP ($\tau > 0.002$), human bare eye (visible cirrus, $\tau > 0.03$), passive radiometers ($\tau > 0.1$). The cirrus cloud cover estimate strongly depends on this detection threshold: over the full area, not taking into account the thinnest cirrus clouds (optical depth below $2 \cdot 10^{-3}$) reduces the total cirrus coverage by ~~12~~15%, and ~~16~~19% for the TTL cirrus only. A passive radiometer insensitive to clouds with an optical depth below 0.1 (for example, onboard geostationary satellites) would only detect 1 cirrus out of ~~23~~, and 1 TTL cirrus out of ~~4~~16. Thus, with BeCOOL's sensitivity, the estimated cirrus cover is significantly increased compared to what is derived from space-borne instruments.

4.2 Optical and geometrical properties of mid- and high-level clouds from BeCOOL and comparison with CALIOP

Statistics of cloud properties (optical depth, top and base altitude) have been compiled for all BeCOOL Level 2 profiles ~~from the 2021 Strateole-2 campaign~~. They are compared with CALIOP nighttime profiles measured during the flight period of the microlidars, over the area covered by the balloons (from -10 to 5° N, 50 to 230° E, dashed white box on Fig. 1). All clouds with a reported base altitude below 5 km have been removed from both datasets to focus on free-tropospheric and TTL clouds.

This includes We also excluded deep convective clouds with full attenuation of the beam. Unlike the previous section, we use here the actual lidar-determined top and base altitude instead of the modified altitudes used for the cloud classification. and removed from CALIOP's database all non reliable retrievals, i.e. where the Extinction_QC flag is greater than 2.



(b) Cloud detection rate : % of lidar profiles

	$\tau < 2 \cdot 10^{-3}$	$\tau > 2 \cdot 10^{-3}$
BeCOOL	32 %	74 %
CALIOP	< 1 %	64 %

Figure 5. Statistical comparison of cloud layers properties detected by BeCOOL and CALIOP. **a:** Probability Density Functions of optical depth of all clouds above 5 km, the grey shading highlights the low optical depth, up to $2 \cdot 10^{-3}$, where the distributions diverge. The percentages of detected clouds with an optical depth lower/greater than this $2 \cdot 10^{-3}$ threshold are reported on the figure. **b:** Percentages of lidar profiles showing clouds with an optical depth lower/greater than the $2 \cdot 10^{-3}$ threshold. Both types of clouds can appear in a single lidar profile.

Figure 5 shows histograms of optical depth for clouds detected above 5 km by the two instruments. An excellent agreement between the distributions appears from $2 \cdot 10^{-3}$ to ~ 1 , the frequencies of occurrence both decrease as τ^{-1} . For BeCOOL, this power law is valid down to $\tau \simeq 10^{-4}$ where the distribution reaches a maximum, whereas a clear cut-off appears at a larger

optical depth $\tau \simeq 2 \cdot 10^{-3}$ in CALIOP's distribution, below which the cloud frequency sharply decreases. 2027% of the cloud layers detected by BeCOOL have an optical depth below $2 \cdot 10^{-3}$. They appear in 2632% of the profiles. For CALIOP, such ultrathin clouds only account for 0.5 % of all detected clouds and are reported in less than 1% of the profiles.




Figure 6. Statistical comparison of cloud layers properties detected by BeCOOL and CALIOP. Probability Density Functions of (lines) top altitude, base altitude, and geometrical depth for (columns) all clouds, clouds with an optical depth above/below the $2 \cdot 10^{-3}$ threshold.

295 Figure 6 shows the distributions of cloud top altitude, base altitude and geometrical depth for all clouds, and separately for clouds with an optical depth larger or smaller than a $2 \cdot 10^{-3}$ threshold. For all clouds, BeCOOL's top altitude distribution shows a sharp mode peaking between 17 and 17.5 km, and a wider base altitude distribution peaking between 16.5 and 17 km. The mean geometrical depth is 1.92 km. Considering only the clouds with an optical depth larger than $2 \cdot 10^{-3}$, the top altitude distribution remains almost unchanged, with a slightly smoother mode just below, between 16.5 and 17, base altitude also
 300 shows a smooth mode between 15 and 16.5 does no longer show any clear mode, the mean geometrical depth is 2.252.6 km.

Considering only cirrus with an optical depth below $2 \cdot 10^{-3}$, both top and base altitude distributions shows a very sharp mode, peaking between 17.5 and 18 km for the top and between ~~17 and 17.5~~ 16.5 and 17 km for the base. Correspondingly, the mean geometrical depth is ~~480~~ 490 m. Almost 75 % of those clouds lie within the TTL, and about 50 % have their base above 16.5 km. Hence, not only does BeCOOL perform well in detecting ultrathin TTL clouds (an expected result considering its high SNR in near-field), more importantly, such clouds are detected in ~~almost more than~~ 20 % of the profiles. Now comparing BeCOOL to CALIOP, a very similar total top altitude distribution is seen, yet with a peak shifted towards lower altitudes, between 16.5 and 17 km. The base altitude distribution of CALIOP does not exhibit any sharp mode and does not extend as high as BeCOOL's, it appears quite uniform between 10.5 and 15.5 km and decreases below, the mean geometrical depth is 2.1 km. Those distributions remain unchanged when considering only clouds with an optical depth larger than $2 \cdot 10^{-3}$ as they account for 99.5 % of all clouds. The agreement with BeCOOL's top and base altitude distribution is almost perfect for those clouds, ~~base altitude distributions are in better agreement~~, although CALIOP's still does not extend as high as BeCOOL's. The mean geometrical depth of those clouds is still 2.1 km, which is slightly lower than for BeCOOL, but the distribution does not extend as much to small depths. ~~For CALIOP, according to Yorks et al. (2011), multiple scattering effects due to the large size of the lidar footprint~~ This difference can be attributed to CALIOP's reception channel photomultiplier tubes which exhibit a non-ideal transient response when exposed to high levels that tend to lower the apparent ~~cloud base altitude~~, enhancing the base altitude of dense clouds and enhance their apparent geometrical depth ~~of the~~ (Lu et al., 2013, 2020). This effect appears clearly on CALIOP's profiles of Fig A1 : the base of the mid-level cloud around 7 km is hidden in the decaying "noise tail" while BeCOOL reveals the true geometrical depth of this cloud. This explains the differences between base altitude distributions and geometrical depth, while top altitude distributions show an excellent agreement. In striking contrast with BeCOOL, clouds with an optical depth below $2 \cdot 10^{-3}$ only represent 0.5 % of CALIOP's cloud database, and their top/base altitude distribution is wide and does not show any pronounced mode.

The excellent agreement between the distributions for optical depths larger than $2 \cdot 10^{-3}$ shows that, despite their limited sampling, balloon-borne observations are representative of the area studied. On the contrary, for small optical depths, the comparison highlights BeCOOL's unique ability to detect ultrathin TTL cirrus. As shown in Sect. 3.2, such cirrus can persist throughout the night below the balloon, and appear homogeneous. They usually lay either right underneath the cold point tropopause, or a local temperature minimum, according to collocated temperature profiles from GPS-Radio Occultation (GPS-RO) soundings (not shown). These characteristics make those thin cirrus similar to UTTCs defined by Peter et al. (2003) and Luo et al. (2003) from the airborne measurements.

Mean top altitude \bar{z}_{top} and geometrical depth $\overline{\Delta z}$ of TTL cirrus for different ranges of optical depth τ are summarized in Table 5. ~~The~~ Except for the optically thicker cirrus ($\tau > 0.1$), that tend to reach higher altitudes, the mean top altitude is fairly constant ~~regardless of the value of τ , whereas, as~~. As expected, the geometrical depth is clearly correlated with τ . The depth of the cloud layer is often used as a free parameter for Lagrangian parcel box models of cirrus and stratospheric dehydration (e.g., Fueglistaler and Baker, 2006; Spichtinger and Krämer, 2013; Schoeberl et al., 2014; Poshyvailo et al., 2018; Nützel et al., 2019). Here, BeCOOL observations suggest typical depths of TTL cirrus ranging from ~~less than~~ 0.5 km (optically thinner ones)

Table 5. Mean [BeCOOL mean](#) top altitude \bar{z}_{top} and geometrical depth $\overline{\Delta z}$ of TTL cirrus for different ranges of optical depth τ .

	% of TTL cirrus	% of 10 min profiles	\bar{z}_{top}	$\overline{\Delta z}$
All TTL cirrus	100 %	16.9 48 %	17.1 km	950 1070 m
$\tau < 2 \cdot 10^{-3}$	39 46 %	23 %	17.1 km	430 440 m
$2 \cdot 10^{-3} < \tau < 3 \cdot 10^{-2}$	33 29 %	16.8 16 %	16.9 km	740 1190 m
$3 \cdot 10^{-2} < \tau < 10^{-1}$	15 19 %	16.8 11 %	17.1 km	1360 1890 m
$10^{-1} < \tau$	13 5 %	17.1 3 %	17.5 km	2130 2800 m

335 to about [23](#) km (optically thicker ones), with a mean of ~ 1 km, which is overall compatible with the values used in modeling studies.

4.3 Cirrus and temperature anomalies

This common detection of very thin TTL cirrus layers by BeCOOL raises the question of the processes responsible for their formation. Following recent papers (Kim et al., 2016; Podglajen et al., 2018; Chang and L’Ecuyer, 2020; Bramberger et al.,
340 2022), we investigated the relationship between TTL clouds and temperature anomalies. Those previous studies highlighted the ubiquitous influence of wave-induced temperature anomalies T' and their vertical gradient dT'/dz on cirrus clouds. Following Chang and L’Ecuyer (2020), temperature anomalies have been computed using GPS-Radio Occultation (GPS-RO) temperature profiles from the Constellation Observing System for Meteorology, Ionosphere, and Climate (COSMIC) Data Analysis and Archive Center (CDAAC) of the University Corporation for Atmospheric Research (UCAR). First, for each BeCOOL
345 flight and each night, a background temperature profile has been determined by averaging all GPS-RO profiles within in a 5° latitude \times 10° longitude box centered on the balloon mean position over a 14-day rolling window. Then, for each night, all GPS-RO profiles falling within a 300 km-radius of the balloon and between 3 h before the first lidar observation and 3 h after the last one were selected. Finally, for each lidar observation, the corresponding temperature anomaly profile was computed as the difference between the closest GPS-RO profile in time among the selected ones and the background. We then split the
350 cloudy lidar data points within the TTL into four categories depending on the temperature anomaly, corresponding to wave phases with positive or negative temperature anomaly T' and lapse-rate anomaly dT'/dz . Figure 7 shows the results for the whole campaign ~~and 4 different 1 ranges~~, in a similar fashion as Figure 3 of Chang and L’Ecuyer (2020). Our results are overall consistent with that previous study, placing almost half of the clouds in the wave phase in which both T' and dT'/dz are negative. As explained in Kim et al. (2016), assuming that temperature anomalies are induced by gravity waves with a downward
355 propagating phase, negative anomalies of dT'/dz correspond to positive vertical wind anomalies, thus to cooling conditions that lower the condensation point. Hence, our observations also suggest favorable conditions for TTL cirrus presence in the cold and cooling phase of gravity waves, which might be related to the influence of the wave-induced saturation anomalies on the formation of the ice crystals (Kim et al., 2016) and/or on their subsequent growth and sedimentation ~~Podglajen et al. (2018)~~
[\(Podglajen et al., 2018\)](#).




Figure 7. Fraction of cloudy BeCOOL lidar bins above 14.5 km within the 4 wave phases. a: cloud population for the whole TTL; b-e: population in four different 1-vertical layers. The percentage in parentheses is the portion of clouds in that layer relative to all TTL clouds.

360 5 Conclusions

Three BeCOOL microlidars were flown during the Strateole-2 scientific campaign in the boreal winter 2021-2022. They provide the first long-duration balloon-borne cloud lidar dataset, covering the equatorial region from the Indian Ocean up to the middle of the Pacific Ocean. These observations were compared with space-borne lidar observations from CALIOP.

365 Case studies of collocated BeCOOL/CALIOP observations for two different types of cirrus clouds demonstrated both the agreement between the two lidars for thicker clouds and BeCOOL's enhanced sensitivity to tenuous clouds. A longer integration time and the proximity of BeCOOL to the studied clouds are responsible for its higher sensitivity. A third case study over convective anvils illustrated the low likelihood of observing short-lived, small-scale structures, such as overshooting convective cloud tops, within a limited dataset gathered from freely drifting balloons. Targeting specific uncommon cloud features would require the use of steerable balloons.

370 Occurrence statistics of different cloud types and profile classification reveal that cirrus clouds are ubiquitous over the area overflowed by the balloons, with a wide range of optical depth covering several orders of magnitude. Cirrus clouds are detected in 8773 % of the lidar profiles, with a limited regional variability during the campaign. On the contrary, the deep convective cloud cover varies very significantly between the studied regions, ranging from 1315 % of the observations over the Maritime Continent to less than 1 % over the Central Pacific Ocean. TTL cirrus, i.e. cirrus with a cloud-cloud base above 14 km, are found in 5948 % of all profiles (and 7165 % over the Central Pacific Ocean). Their mean top altitude is 17 km, and does not depend on their optical depth. Their geometrical depth ranges from less than 0.1 to 4 km, with an overall mean of ~ 1 km. Ultrathin TTL cirrus, with optical depth below the detection threshold of CALIOP ($\tau < 2 \cdot 10^{-3}$), are reported in 4623 % of the lidar profiles, and have a mean geometrical depth of about 400440 m.

380 Those very thin TTL clouds are reminiscent of Ultrathin Tropical Tropopause Clouds described by Peter et al. (2003), in particular with respect to their small vertical extension, huge horizontal extension and lateral homogeneity. These-They also share these typical characteristics with laminar cirrus clouds reported notably by Winker and Trepte (1998) from LITE observations and Wang et al. (2019) from reprocessed CALIOP observations. How frequent are those ultrathin cirrus farther

away from the equator is still to be investigated and future BeCOOL flights at higher latitude would be useful to better characterize their coverage.

385 TTL cirrus clouds play a significant role in the dehydration process of air masses entering the stratosphere (e.g., Jensen et al., 1996; Schoeberl et al., 2019). An ongoing study investigates their radiative impact from BeCOOL's measurements. Our observations also confirm the ubiquitous relationship between waves and tropical cirrus clouds found in previous papers (Kim et al., 2016; Chang and L'Ecuyer, 2020; Bramberger et al., 2022): TTL cirrus are more common in the cold and cooling phase of waves. Future work will focus on characterizing the horizontal scales and lifetimes of TTL cirrus combining CALIOP and
390 BeCOOL, in order to elucidate the link between waves and TTL cirrus life cycle.

Data availability. The Strateole-2 BeCOOL data set is available at <https://data.ipsl.fr/catalog/strateole2/>. CALIOP data was downloaded from AERIS/ICARE datacenter (<https://www.icare.univ-lille.fr/calipso/products/>). GPS-RO data can be accessed at COSMIC Data Analysis and Archive Center (<https://cdaac-www.cosmic.ucar.edu/cdaac/index.html>). The merged IR satellite images were collected from the NOAA/NCEP GPM_MERGIR product, available at https://disc.gsfc.nasa.gov/datasets/GPM_MERGIR_1/summary.

395 **Appendix A:** ~~Additional brightness temperature maps~~ Lidar profiles at coincidence for the first case studies

Appendix B: Additional brightness temperature maps for the case studies

Author contributions. TL, FR and AP conceived the study. TL performed the study with scientific support from FR, AP and JP. TL wrote the paper with contributions from FR and AP. VM designed and built BeCOOL microlidar. All authors agreed on the final version.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

400 *Acknowledgements.* The balloon-borne BeCOOL data were collected as part of Strateole-2, which is sponsored by CNES, CNRS/ INSU, NSF, and ESA. TL acknowledges support by a doctoral grant from France's Centre National d'Études Spatiales (CNES). The authors ~~would~~ sincerely thanks three anonymous referees for their helpful comments and suggestions which helped us improve the paper. We would also like to thank the numerous developers that contributed to the free and open-source tools used for the data analysis and visualization, in particular xarray (Hoyer and Hamman, 2017, DOI: <https://doi.org/10.5334/jors.148>) and Matplotlib (Hunter, 2007, DOI: 10.1109/MCSE.2007.55).

First case study : lidar profiles




Figure A1. Attenuated backscatter profiles around coincidence for case study 1 : (left) BeCOOL 10-minute averaged, (right) CALIOP 5 km

405 References

Southern Oscillation Index (SOI) | El Niño/Southern Oscillation (ENSO) | National Centers for Environmental Information (NCEI), <https://www.ncei.noaa.gov/access/monitoring/enso/soi>, last accessed on 2023-08-15.




Figure B1. Hourly brightness temperature maps for case study 1, 2021-11-29. Red dot is the balloon position, solid (dotted) line is the nighttime (daytime) balloon track.




Figure B2. Hourly brightness temperature maps ~~of~~ for case study 2, 2021-11-27. Red dot is the balloon position, solid (dotted) line is the nighttime (daytime) balloon track.

Bramberger, M., Alexander, M. J., Davis, S., Podglajen, A., Hertzog, A., Kalnajs, L., Deshler, T., Goetz, J. D., and Khaykin, S.: First Super-Pressure Balloon-Borne Fine-Vertical-Scale Profiles in the Upper TTL: Impacts of Atmospheric Waves on Cirrus Clouds and the QBO, *Geophysical Research Letters*, 49, e2021GL097596, <https://doi.org/10.1029/2021GL097596>, 2022.

410

Braun, B. M., Sweetser, T. H., Graham, C., and Bartsch, J.: CloudSat's A-Train Exit and the Formation of the C-Train: An Orbital Dynamics Perspective, in: 2019 IEEE Aerospace Conference, pp. 1–10, IEEE, Big Sky, MT, USA, <https://doi.org/10.1109/AERO.2019.8741958>, 2019.




Figure B3. Hourly brightness temperature maps for case study 2, 2021-11-27. Red dot is the balloon position, solid (dotted) line is the nighttime (daytime) balloon track.

- Brewer, A. W.: Evidence for a world circulation provided by the measurements of helium and water vapour distribution in the stratosphere, *Quarterly Journal of the Royal Meteorological Society*, 75, 351–363, <https://doi.org/10.1002/qj.49707532603>, 1949.
- 415 Chang, K.-W. and L'Ecuyer, T.: Influence of gravity wave temperature anomalies and their vertical gradients on cirrus clouds in the tropical tropopause layer – a satellite-based view, *Atmos. Chem. Phys.*, 20, 12 499–12 514, <https://doi.org/10.5194/acp-20-12499-2020>, 2020.
- Corcos, M., Hertzog, A., Plougonven, R., and Podglajen, A.: Observation of Gravity Waves at the Tropical Tropopause Using Superpressure Balloons, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD035 165, <https://doi.org/10.1029/2021JD035165>, 2021.
- 420 Dauhut, T., Chaboureaud, J.-P., Haynes, P. H., and Lane, T. P.: The Mechanisms Leading to a Stratospheric Hydration by Overshooting Convection, *Journal of the Atmospheric Sciences*, 75, 4383–4398, <https://doi.org/10.1175/JAS-D-18-0176.1>, 2018.
- Fernald, F. G., Herman, B. M., and Reagan, J. A.: Determination of Aerosol Height Distributions by Lidar, *J. Appl. Meteor.*, 11, 482–489, [https://doi.org/10.1175/1520-0450\(1972\)011<0482:DOAHDB>2.0.CO;2](https://doi.org/10.1175/1520-0450(1972)011<0482:DOAHDB>2.0.CO;2), 1972.
- Fueglistaler, S. and Baker, M. B.: A modelling study of the impact of cirrus clouds on the moisture budget of the upper troposphere, *Atmospheric Chemistry and Physics*, 6, 1425–1434, <https://doi.org/10.5194/acp-6-1425-2006>, 2006.
- 425 Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Reviews of Geophysics*, 47, <https://doi.org/10.1029/2008RG000267>, 2009.
- Fujiwara, M., Iwasaki, S., Shimizu, A., Inai, Y., Shiotani, M., Hasebe, F., Matsui, I., Sugimoto, N., Okamoto, H., Nishi, N., Hamada, A., Sakazaki, T., and Yoneyama, K.: Cirrus observations in the tropical tropopause layer over the western Pacific, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD011040>, 2009.
- 430 Gage, K. S. and Reid, G. C.: Longitudinal variations in tropical tropopause properties in relation to tropical convection and El Niño-Southern Oscillation events, *Journal of Geophysical Research: Oceans*, 92, 14 197–14 203, <https://doi.org/10.1029/JC092iC13p14197>, 1987.
- Haase, J. S., Alexander, M. J., Hertzog, A., Kalnajs, L., Deshler, T., Davis, S. M., Plougonven, R., Cocquerez, P., and Venel, S.: Around the World in 84 Days, <http://eos.org/science-updates/around-the-world-in-84-days>, 2018.

- 435 Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-troposphere exchange, *Reviews of Geophysics*, 33, 403–439, <https://doi.org/10.1029/95RG02097>, 1995.
- Iwasaki, S., Luo, Z. J., Kubota, H., Shibata, T., Okamoto, H., and Ishimoto, H.: Characteristics of cirrus clouds in the tropical lower stratosphere, *Atmospheric Research*, 164–165, 358–368, <https://doi.org/10.1016/j.atmosres.2015.06.009>, 2015.
- Janowiak, J. E., Joyce, R. J., and Yarosh, Y.: A Real-Time Global Half-Hourly Pixel-Resolution Infrared Dataset and Its Applications, 440 *Bulletin of the American Meteorological Society*, 82, 205–218, [https://doi.org/10.1175/1520-0477\(2001\)082<0205:ARTGHH>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<0205:ARTGHH>2.3.CO;2), 2001.
- Jensen, E. J., Toon, O. B., Pfister, L., and Selkirk, H. B.: Dehydration of the upper troposphere and lower stratosphere by subvisible cirrus clouds near the tropical tropopause, *Geophysical Research Letters*, 23, 825–828, <https://doi.org/10.1029/96GL00722>, 1996.
- Jensen, E. J., Pfister, L., Jordan, D. E., Bui, T. V., Ueyama, R., Singh, H. B., Thornberry, T. D., Rollins, A. W., Gao, R.-S., Fahey, D. W., 445 Rosenlof, K. H., Elkins, J. W., Diskin, G. S., DiGangi, J. P., Lawson, R. P., Woods, S., Atlas, E. L., Navarro Rodriguez, M. A., Wofsy, S. C., Pittman, J., Bardeen, C. G., Toon, O. B., Kindel, B. C., Newman, P. A., McGill, M. J., Hlavka, D. L., Lait, L. R., Schoeberl, M. R., Bergman, J. W., Selkirk, H. B., Alexander, M. J., Kim, J.-E., Lim, B. H., Stutz, J., and Pfeilsticker, K.: The NASA Airborne Tropical Tropopause Experiment: High-Altitude Aircraft Measurements in the Tropical Western Pacific, *Bulletin of the American Meteorological Society*, 98, 129–143, <https://doi.org/10.1175/BAMS-D-14-00263.1>, 2017.
- 450 Kar, J., Vaughan, M. A., Lee, K.-P., Tackett, J. L., Avery, M. A., Garnier, A., Getzewich, B. J., Hunt, W. H., Josset, D., Liu, Z., Lucker, P. L., Magill, B., Omar, A. H., Pelon, J., Rogers, R. R., Toth, T. D., Trepte, C. R., Vernier, J.-P., Winker, D. M., and Young, S. A.: CALIPSO lidar calibration at 532 nm: version 4 nighttime algorithm, *Atmos. Meas. Tech.*, 11, 1459–1479, <https://doi.org/10.5194/amt-11-1459-2018>, 2018.
- Kim, J.-E., Alexander, M. J., Bui, T. P., Dean-Day, J. M., Lawson, R. P., Woods, S., Hlavka, D., Pfister, L., and Jensen, E. J.: Ubiquitous influence of waves on tropical high cirrus clouds: Wave Influence on Tropical High Cirrus, *Geophys. Res. Lett.*, 43, 5895–5901, 455 <https://doi.org/10.1002/2016GL069293>, 2016.
- Klett, J. D.: Stable analytical inversion solution for processing lidar returns, *Appl. Opt.*, 20, 211, <https://doi.org/10.1364/AO.20.000211>, 1981.
- Krämer, M., Rolf, C., Spelten, N., Afchine, A., Fahey, D., Jensen, E., Khaykin, S., Kuhn, T., Lawson, P., Lykov, A., Pan, L. L., Riese, M., 460 Rollins, A., Stroh, F., Thornberry, T., Wolf, V., Woods, S., Spichtinger, P., Quaas, J., and Sourdeval, O.: A microphysics guide to cirrus – Part 2: Climatologies of clouds and humidity from observations, *Atmos. Chem. Phys.*, 20, 12 569–12 608, <https://doi.org/10.5194/acp-20-12569-2020>, 2020.
- Lee, K.-O., Dauhut, T., Chaboureaud, J.-P., Khaykin, S., Krämer, M., and Rolf, C.: Convective hydration in the tropical tropopause layer during the StratoClim aircraft campaign: pathway of an observed hydration patch, *Atmos. Chem. Phys.*, 19, 11 803–11 820, 465 <https://doi.org/10.5194/acp-19-11803-2019>, 2019.
- Lu, X., Hu, Y., Liu, Z., Zeng, S., and Trepte, C.: CALIOP receiver transient response study, p. 887316, San Diego, California, United States, <https://doi.org/10.1117/12.2033589>, 2013.
- Lu, X., Hu, Y., Vaughan, M., Rodier, S., Trepte, C., Lucker, P., and Omar, A.: New attenuated backscatter profile by removing the CALIOP receiver’s transient response, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 255, 107 244, 470 <https://doi.org/10.1016/j.jqsrt.2020.107244>, 2020.
- Luo, B. P., Peter, T., Wernli, H., Fueglistaler, S., Wirth, M., Kiemle, C., Flentje, H., Yushkov, V. A., Khattatov, V., Rudakov, V., Thomas, A., Borrmann, S., Toci, G., Mazzinghi, P., Beuermann, J., Schiller, C., Cairo, F., Don-Francesco, G. D., Adriani, A., Volk, C. M., Strom, J.,

- Noone, K., Mitev, V., MacKenzie, R. A., Carslaw, K. S., Trautmann, T., Santacesaria, V., and Stefanutti, L.: Ultrathin Tropical Tropopause Clouds (UTTCS): II. Stabilization mechanisms, *Atmos. Chem. Phys.*, p. 8, 2003.
- 475 Martins, E., Noel, V., and Chepfer, H.: Properties of cirrus and subvisible cirrus from nighttime Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP), related to atmospheric dynamics and water vapor, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2010JD014519>, 2011.
- McFarquhar, G. M., Heymsfield, A. J., Spinhirne, J., and Hart, B.: Thin and Subvisual Tropopause Tropical Cirrus: Observations and Radiative Impacts, *J. Atmos. Sci.*, 57, 1841–1853, [https://doi.org/10.1175/1520-0469\(2000\)057<1841:TASTTC>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<1841:TASTTC>2.0.CO;2), 2000.
- 480 Nützel, M., Podglajen, A., Garny, H., and Ploeger, F.: Quantification of water vapour transport from the Asian monsoon to the stratosphere, *Atmos. Chem. Phys.*, 19, 8947–8966, <https://doi.org/10.5194/acp-19-8947-2019>, 2019.
- Peter, T., Luo, B. P., Wirth, M., Kiemle, C., Flentje, H., Yushkov, V. A., Khattatov, V., Rudakov, V., Thomas, A., Borrmann, S., Toci, G., Mazinghi, P., Beuermann, J., Schiller, C., Cairo, F., Donfrancesco, G. D., Adriani, A., Volk, C. M., Strom, J., Noone, K., Mitev, V., MacKenzie, R. A., Carslaw, K. S., Trautmann, T., Santacesaria, V., and Stefanutti, L.: Ultrathin Tropical Tropopause Clouds (UTTCS): I.
- 485 Cloud morphology and occurrence, *Atmos. Chem. Phys.*, p. 9, 2003.
- Platt, C. M. R.: Lidar and Radiometric Observations of Cirrus Clouds, *Journal of Atmospheric Sciences*, 30, 1191 – 1204, [https://doi.org/10.1175/1520-0469\(1973\)030<1191:LAROOC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1973)030<1191:LAROOC>2.0.CO;2), 1973.
- Platt, C. M. R., Dille, A. C., Scott, J. C., Barton, I. J., and Stephens, G. L.: Remote Sounding of High Clouds. V: Infrared Properties and Structures of Tropical Thunderstorm Anvils, *Journal of Applied Meteorology and Climatology*, 23, 1296–1308,
- 490 [https://doi.org/10.1175/1520-0450\(1984\)023<1296:RSOHCV>2.0.CO;2](https://doi.org/10.1175/1520-0450(1984)023<1296:RSOHCV>2.0.CO;2), 1984.
- Platt, C. M. R., Scott, S. C., and Dille, A. C.: Remote Sounding of High Clouds. Part VI: Optical Properties of Midlatitude and Tropical Cirrus, *Journal of the Atmospheric Sciences*, 44, 729–747, [https://doi.org/10.1175/1520-0469\(1987\)044<0729:RSOHCV>2.0.CO;2](https://doi.org/10.1175/1520-0469(1987)044<0729:RSOHCV>2.0.CO;2), 1987.
- Podglajen, A., Plougonven, R., Hertzog, A., and Jensen, E.: Impact of gravity waves on the motion and distribution of atmospheric ice particles, *Atmos. Chem. Phys.*, 18, 10 799–10 823, <https://doi.org/10.5194/acp-18-10799-2018>, 2018.
- 495 Poshvyvailo, L., Müller, R., Konopka, P., Günther, G., Riese, M., Podglajen, A., and Ploeger, F.: Sensitivities of modelled water vapour in the lower stratosphere: temperature uncertainty, effects of horizontal transport and small-scale mixing, *Atmospheric Chemistry and Physics*, 18, 8505–8527, <https://doi.org/10.5194/acp-18-8505-2018>, 2018.
- Prabhakara, C., Fraser, R. S., Dalu, G., Wu, M.-L. C., Curran, R. J., and Styles, T.: Thin Cirrus Clouds: Seasonal Distribution over Oceans Deduced from Nimbus-4 IRIS, *Journal of Applied Meteorology and Climatology*, 27, 379–399, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0450(1988)027<0379:TCCSDO>2.0.CO;2)
- 500 [0450\(1988\)027<0379:TCCSDO>2.0.CO;2](https://doi.org/10.1175/1520-0450(1988)027<0379:TCCSDO>2.0.CO;2), 1988.
- Randel, W. J. and Jensen, E. J.: Physical processes in the tropical tropopause layer and their roles in a changing climate, *Nature Geosci*, 6, 169–176, <https://doi.org/10.1038/ngeo1733>, 2013.
- Ravetta, F., Mariage, V., Lesigne, T., and Pelon, J.: Balloonborne lidar for cirrus and convective overshoot studies, in preparation to be submitted to *Atmospheric Measurement Techniques*, 2023.
- 505 Reagan, J., Wang, X., and Osborn, M.: Spaceborne lidar calibration from cirrus and molecular backscatter returns, *IEEE Trans. Geosci. Remote Sensing*, 40, 2285–2290, <https://doi.org/10.1109/TGRS.2002.802464>, 2002.
- Sassen, K., Griffin, M. K., and Dodd, G. C.: Optical Scattering and Microphysical Properties of Subvisual Cirrus Clouds, and Climatic Implications, *Journal of Applied Meteorology and Climatology*, 28, 91–98, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0450(1989)028<0091:OSAMPO>2.0.CO;2)
- 0450(1989)028<0091:OSAMPO>2.0.CO;2, 1989.

- 510 Schoeberl, M. R., Dessler, A. E., Wang, T., Avery, M. A., and Jensen, E. J.: Cloud formation, convection, and stratospheric dehydration, *Earth and Space Science*, 1, 1–17, <https://doi.org/https://doi.org/10.1002/2014EA000014>, 2014.
- Schoeberl, M. R., Jensen, E. J., Pfister, L., Ueyama, R., Wang, T., Selkirk, H., Avery, M., Thornberry, T., and Dessler, A. E.: Water Vapor, Clouds, and Saturation in the Tropical Tropopause Layer, *Journal of Geophysical Research: Atmospheres*, 124, 3984–4003, <https://doi.org/10.1029/2018JD029849>, 2019.
- 515 Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M., Sanford, T. J., and Plattner, G.-K.: Contributions of Stratospheric Water Vapor to Decadal Changes in the Rate of Global Warming, *Science*, 327, 1219–1223, <https://doi.org/10.1126/science.1182488>, 2010.
- Sourdeval, O., Gryspeerdt, E., Krämer, M., Goren, T., Delanoë, J., Afchine, A., Hemmer, F., and Quaas, J.: Ice crystal number concentration estimates from lidar–radar satellite remote sensing – Part 1: Method and evaluation, *Atmos. Chem. Phys.*, 18, 14 327–14 350, <https://doi.org/10.5194/acp-18-14327-2018>, 2018.
- 520 Spichtinger, P. and Krämer, M.: Tropical tropopause ice clouds: a dynamic approach to the mystery of low crystal numbers, *Atmos. Chem. Phys.*, 13, 9801–9818, <https://doi.org/10.5194/acp-13-9801-2013>, 2013.
- Sèze, G., Pelon, J., Derrien, M., Le Gléau, H., and Six, B.: Evaluation against CALIPSO lidar observations of the multi-geostationary cloud cover and type dataset assembled in the framework of the Megha-Tropiques mission, *Quarterly Journal of the Royal Meteorological*
- 525 *Society*, 141, 774–797, <https://doi.org/10.1002/qj.2392>, 2015.
- Thorsen, T. J., Fu, Q., and Comstock, J.: Comparison of the CALIPSO satellite and ground-based observations of cirrus clouds at the ARM TWP sites, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD015970>, 2011.
- Vaillant de Guélis, T., Vaughan, M. A., Winker, D. M., and Liu, Z.: Two-dimensional and multi-channel feature detection algorithm for the CALIPSO lidar measurements, *Atmos. Meas. Tech.*, 14, 1593–1613, <https://doi.org/10.5194/amt-14-1593-2021>, 2021.
- 530 Wang, P.-H., McCormick, M., Poole, L., Chu, W., Yue, G., Kent, G., and Skeens, K.: Tropical high cloud characteristics derived from SAGE II extinction measurements, *Atmospheric Research*, 34, 53–83, [https://doi.org/10.1016/0169-8095\(94\)90081-7](https://doi.org/10.1016/0169-8095(94)90081-7), 1994.
- Wang, T., Wu, D. L., Gong, J., and Tsai, V.: Tropopause Laminar Cirrus and Its Role in the Lower Stratosphere Total Water Budget, *Journal of Geophysical Research: Atmospheres*, 124, 7034–7052, <https://doi.org/10.1029/2018JD029845>, 2019.
- Winker, D. M. and Trepte, C. R.: Laminar cirrus observed near the tropical tropopause by LITE, *Geophysical Research Letters*, 25, 3351–
- 535 3354, <https://doi.org/10.1029/98GL01292>, 1998.
- Yang, Q., Fu, Q., and Hu, Y.: Radiative impacts of clouds in the tropical tropopause layer, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012393>, 2010.
- Yorks, J. E., Hlavka, D. L., Vaughan, M. A., McGill, M. J., Hart, W. D., Rodier, S., and Kuehn, R.: Airborne validation of cirrus cloud properties derived from CALIPSO lidar measurements: Spatial properties, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD015942>, 2011.
- 540 Young, S. A., Vaughan, M. A., Garnier, A., Tackett, J. L., Lambeth, J. D., and Powell, K. A.: Extinction and optical depth retrievals for CALIPSO’s Version 4 data release, *Atmos. Meas. Tech.*, 11, 5701–5727, <https://doi.org/10.5194/amt-11-5701-2018>, 2018.