

# 1 Monthly velocity of the Mont Blanc glaciers derived from Sentinel- 2 2 between 2016-2023

3 Fabrizio Troilo<sup>1,4</sup>, Niccolò Dematteis<sup>3</sup>, Francesco Zucca<sup>4</sup>, Martin Funk<sup>2</sup>, Daniele Giordan<sup>3</sup>.

4 <sup>1</sup>Fondazione Montagna sicura, Glaciers, snow and avalanche research area, Courmayeur, 11013, Italy.

5 <sup>2</sup>ETH-VAW, Versuchsanstalt für Wasserbau, Hydrologie und Glaziologie, Zurich, CH-8092, Switzerland.

6 <sup>3</sup>Research Institute for Geo-Hydrological Protection IRPI, Italian National Research Council, Turin, 10135, Italy.

7 <sup>4</sup>University of Pavia, Department of Earth and Environmental Sciences, Pavia, 27100, Italy.

8 *Correspondence to: Niccolò Dematteis (niccolo.dematteis@irpi.cnr.it); Fabrizio Troilo (ftroilo@fondms.org).*

9 **Abstract.** We investigated the temporal variability of the surface velocity of thirty glaciers in the Mont Blanc massif  
10 (European Alps). We calculated the monthly velocity between 2016 and 2023 using digital image correlation of Sentinel-2  
11 optical imagery. The main objectives of the study are: (i) to characterise the variability of the velocity fields of such glaciers,  
12 referring both to their temporal (seasonal and interannual) and spatial variations; (ii) to investigate relationships between the  
13 morphology of glaciers and their kinematics. We measured monthly velocities varying from 12.7 m yr<sup>-1</sup> to 487.4 m yr<sup>-1</sup>. We  
14 observed an overall decrease in the velocity between 2016 and 2019 and an unexpected rise in 2020-2022, especially visible  
15 in most glaciers on the southern side of the massif. Considering the whole period, half glaciers showed positive acceleration,  
16 which reached values >4 m yr<sup>-2</sup> in three glaciers. In general, the velocity linear trend is higher in the cold season. We found  
17 that smaller glaciers have a more pronounced seasonality, with winter-summer velocity differences of 50-100%. Finally, in  
18 2016, 2018 and 2022, we observed an exceptionally high winter-summer velocity difference in the 0.3 km<sup>2</sup>-wide Charpoua  
19 Glacier, when summer velocities increased by one order of magnitude.

## 20 1 Introduction

21 Glacier flow has been one of the early drivers of glaciological interest and research since it was first studied. Its  
22 understanding and modelling evolved via the observations and findings of Somigliana in the early 1900s, Glen's laboratory  
23 experiments (Glen, 1952), followed by the interpretations of Nye (Nye, 1952) during the 1950s, to cite just a few, and have  
24 explained that the two main mechanisms of glacier flow rely on ice deformation and basal sliding. However, the motion of  
25 Alpine glaciers is largely related to basal sliding (Willis, 1995). Because continuous monitoring of sliding velocities in the  
26 field is extremely difficult and rarely achieved (Vincent and Moreau, 2016), measuring surface flow velocities can be used as  
27 a strong alternative approach. Nonetheless, the continuous monitoring of surface velocities of Alpine glaciers is complex on  
28 specific study sites, and very rarely has it been performed on a spatially distributed scale.

29 The flow of glaciers generally depends on a variety of physical parameters. The main physical parameter influencing ice  
30 velocity is the ice thickness (Jiskoot 2011), which is indirectly related to the glacier trend in mass balance as it determines an  
31 evolution towards an increase or decrease in glacier thickness. Other parameters that influence ice flow are glacier surface  
32 slope, ice properties (temperature, density), bedrock conditions (hard, soft, frozen or thawed ice-bed contact), topography, the  
33 glacier's terminal area type (land, sea, ice shelf), but also air temperature and precipitation and their seasonality that influences  
34 subglacial hydrology (Jiskoot, 2011; Humbert et al., 2005; Cuffey and Paterson, 2010; Benn and Evans, 2014; Bindshadler,  
35 1983).

36 The analysis of glacier surface velocity has a wide array of applications: it is a powerful climate change indicator (Beniston  
37 et al., 2018) and also an important input data for ice thickness models (Millan et al., 2022; Samsonov et al., 2021) and mass  
38 balance models that can also approximate sea-level rise contribution by glaciers (Zekollari et al., 2019). In the field of glacial  
39 hazards, it is used as an indicator for the detection of glacier surges with space-borne measurements (Kamb, 1987; Kääh et al.,  
40 2021), and accelerations that can result in glacier-related hazards using ground-based sensors (Pralong and Funk, 2006;  
41 Giordan et al., 2020). Measurements of the surface velocity of glaciers can be achieved by terrestrial techniques (Dematteis et  
42 al., 2021) such as topographic measurements of stakes or fixed points on the glacier (Stocker-Waldhuber et al., 2019), GNSS  
43 repeated or continuous surveys (Einarsson et al., 2016), digital image correlation of oblique photographs (Evans, 2000; Ahn  
44 and Box, 2010) and terrestrial radar interferometry (Luzi et al., 2007; Allstadt et al., 2015). Considering remote sensing  
45 solutions, glacier surface velocities can be measured by different aerial and space-borne sensors. In recent decades, public  
46 access to satellite optical and radar data (especially from Sentinel and Landsat constellation satellites), as well as the  
47 commercial availability of very high resolution (30 cm to 3 m ground resolution) optical imagery (Deilami and Hashim, 2011)  
48 and radar data (Rankl et al., 2014), have given great input to glaciological research. In particular, Sentinel-2 optical imagery  
49 is widely used in glaciological studies and has been tested in the literature on various environments (Paul et al., 2016; Millan  
50 et al., 2019). Nowadays, the automated processing of ice velocity maps with global coverage from satellite imagery is freely  
51 available online from web-based platforms such as the GoLIVE datasets (Fahnestock et al., 2016), the ITS\_LIVE data portal  
52 (<https://its-live.jpl.nasa.gov/>) or the FAU-Glacier portal (RETREAT, 2021 Ice surface velocities derived from Sentinel-1,  
53 Version 1; <http://retreat.geographie.uni-erlangen.de/search>). The availability of such datasets is very relevant globally, but  
54 their application to Alpine glaciers is limited due to their relatively coarse spatial resolution - e.g., 300x300 m (GoLIVE),  
55 120x120 m (ITS\_LIVE) - which can provide data on just a few of the largest Alpine glaciers. Moreover, for the ITS\_LIVE  
56 dataset, the velocity maps are calculated at a resolution of 240 m and statistically downscaled to 120 m, which has major  
57 limitations for small mountain glaciers. The adopted resolution is a trade-off between computational effort and the best  
58 resolution of the output that must cope with the global availability of the analysis. Limiting the processing of images at a  
59 regional scale decreases the computational effort compared to global products and makes it easier to obtain higher resolution  
60 velocity maps that allow Alpine glaciers to be investigated (Berthier et al., 2005), even though very small glaciers (i.e., width  
61 <250 m) are still difficult to analyse (Millan et al., 2019).

62 Recent studies using different techniques have measured spatio-temporal variations of ice velocity on large valley glaciers  
63 in an Alpine environment, like the Argentière (Vincent and Moreau, 2016) and Miage glaciers (Fyffe, 2012) as well as on  
64 steep glacier snouts (e.g., Planpincieux Glacier by (Giordan et al., 2020) but a spatially distributed analysis at a regional scale  
65 of the variations of velocities over glaciers with different morphological characteristics is, as of today, still lacking in the  
66 Alpine environment. (Millan et al., 2022) calculated the velocities of world glaciers, but they considered a specific analysis  
67 period (2017-2018) not considering velocity variations in time, while (Rabatel et al., 2023a) performed a comparison between  
68 yearly aggregated velocity maps between 2015 and 2021 over three different alpine massifs.

69 Other studies focused on long-term glacier velocity records in the Mont Blanc Massif: at the Argentière Glacier (Vincent  
70 and Moreau, 2016) and Miage Glacier (Smiraglia et al., 2000; Fyffe, 2012). At Argentière Glacier, a unique series of  
71 continuous basal sliding measurements existed from 1997 and was still active as of 2022 (Vincent et al., 2022; Nanni et al.,  
72 2020). The whole series indicates a general decrease in basal sliding velocities (Vincent and Moreau, 2016) since the end of  
73 the 1990s. This general decrease has shown a strong correlation with the negative mass balance of the glacier, which agrees  
74 with the conceptual model from Span and Kuhn (Span and Kuhn, 2003), in which the glacier flow variation is primarily driven  
75 by the mass balance of the accumulation area in the previous year (as it determines glacier thickness variations). Seasonal field  
76 surveys conducted at Argentière Glacier from the 1950s document a longer data series than the basal sliding measurements  
77 started in 1997, and an increase in surface velocities was measured during a period of positive mass balances in the early 1980s


78 (Vincent and Moreau, 2016). The same trend was highlighted by Span and Kuhn (Span and Kuhn, 2003) for at least six other  
79 glaciers: Saint Sorlin in France, Gietro and Corbassiere in Switzerland; Pasterze, Vernagtferner and Odenwinkelkees in  
80 Austria. At Miage Glacier, surface velocities have been measured historically by different authors (Diolaiuti et al., 2005;  
81 Smiraglia et al., 2000; Fyffe, 2012; Lesca, 1974; Pelfini et al., 2007; Deline, 2002) and also show a general velocity decrease  
82 in recent decades (Smiraglia et al., 2000; Fyffe, 2012). Glaciers such as Miage and Argentière, reach a low altitude and have  
83 flat and little crevassed valley tongues, for this reason, they have often been historically chosen for glaciological field surveys  
84 (Span and Kuhn, 2003). Therefore, the knowledge of Alpine glacier kinematics is generally mostly related to this type of  
85 glacier, which can be significantly different compared to the other glaciers analysed in this study.

86 Globally, glacier slowdown linked to a negative mass balance trend was also shown for six different regions around the  
87 globe and dates spanning from 1953 to 2009 by Heid and Kääb (2012b) by an analysis of remotely-sensed optical images.  
88 Specific analysis of velocity trends and glacier mass loss showed generalized decreasing velocity trends over different regions  
89 of High Mountain Asia between 2000 and 2017 and a strong correlation with the negative mass balance trend (Dehecq et al.,  
90 2019).

91 The main purposes of this study are the production of eight-year-long velocity time series of the surface velocity of thirty  
92 glaciers at a massif scale, as well as an integrated analysis of morphological and kinematic features of such glaciers. The  
93 identification of possible trends in the velocity time series is a major objective of the present study. We analysed eight years  
94 (2016-2023) of Sentinel-2 optical imagery to retrieve monthly velocity data on thirty glaciers in the Mont Blanc massif. We  
95 observed different behaviours of surface velocity and identified a relationship between kinematic seasonality and glacier size.

## 96 **2 Area of study**

97 The study area is the Mont Blanc massif. It is located in the western part of the European Alps bordering France, Italy and  
98 Switzerland (Fig. 1) and culminating at 4809 m with the Mont Blanc summit, the highest peak in Central Europe. Many other  
99 peaks in the Mont Blanc massif reach well above 4000 m and the entire area is very highly frequented with famous tourist  
100 resorts such as Courmayeur and Chamonix attracting thousands of tourists every year.



101 **Fig. 1 Study area of the Mont Blanc massif. Background: true colour image (cloud-free Europe mosaic in the upper**  
 102 **left panel), courtesy of the Copernicus Open Access Hub (<https://scihub.copernicus.eu>, last access: 10 September**  
 103 **2023).**  
 104

105 The total surface of glaciers in the Mont Blanc massif is equal to 169 km<sup>2</sup> and totals 116 glaciers, according to the Randolph  
 106 glacier inventory (RGI 7.0) (RGI Consortium, 2023). The inventory refers to 2003 (Pfeffer et al., 2014; Arendt et al., 2017).  
 107 Forty glaciers are very small, covering an area of less than 0.1 km<sup>2</sup>, forty-seven have surfaces comprised between 0,1 and 1  
 108 km<sup>2</sup>, sixteen glaciers fall in the range between 1 km<sup>2</sup> and 3 km<sup>2</sup>, and thirteen glaciers have surface areas of more than 4 km<sup>2</sup>.

109 The geological setting and the geomorphology of the Mont Blanc massif form a high mountain range with its main ridge  
 110 line oriented in a southwest/northeast direction along the French-Italian border. The valley floors flanking the massif have low  
 111 altitudes - in the range of 1000-1500 m - resulting in steep slopes originating from the highest peaks with large vertical  
 112 altitudinal differences. The meteo-climatic local conditions on the massif are of a continental type, but orographic effects on  
 113 the predominant incoming weather fronts produce larger precipitation compared to nearby regions (Gottardi et al., 2012).

114 The Argentière Glacier is the only glacier with regular mass balance measurements included in the World Glacier  
 115 Monitoring Service (WGMS) 'Reference Glaciers' dataset in the Mont Blanc Massif (Zemp et al., 2009). The Argentière  
 116 Glacier has shown a general negative mass balance trend since the early 1990s, in line with mass balances of other Alpine  
 117 glaciers and glaciers from other mountain ranges across the globe. Geodetic mass balance measurements of the Thoula Glacier,  
 118 a small glacier on the border between France and Italy at altitudes between 2900 and 3300 m, represent well the local meteo-

119 climatic conditions that result in slightly less negative mass balance trends compared to other glaciers in the Alps (Zemp et al.,  
120 2021; Zemp et al., 2020; Mondardini et al., 2021).


121 A more spatially distributed analysis of mass balances in the Mont Blanc region has also been outlined in the literature  
122 employing geodetic mass balances of the whole Mont Blanc massif using stereo satellite imagery from the Pléiades and Spot  
123 satellite constellations (Berthier et al., 2014; Beraud et al., 2023). The trend outlined by Berthier et al. (2014) at the massif  
124 scale, reflects data trends comparable to the glaciological mass balances of WGMG reference glaciers in the Alps. Glaciers at  
125 lower altitudes show larger ice volume losses and subsequent substantial glacier front retreats (Paul et al., 2020), while glaciers  
126 at higher altitudes suffer less acute volume loss and shrinkage. Large differences in the glacier frontal position, especially for  
127 the lower altitude terminating glaciers, can be well assessed by the difference of the terminus position in recent satellite imagery  
128 compared to the position outlined on the RGI 7.0.

## 129 **3 Materials and Methods**

### 130 **3.1 General workflow**

131 In this paper, we analysed the Copernicus - ESA Sentinel-2 optical satellite imagery dataset available for the study area. In  
132 addition, we used Airbus Pleiades Stereo derived digital elevation models (DEMs) to retrieve morphometric data of glaciers,  
133 and publicly available modelled ice thickness data from Millan et al. (Millan et al., 2022). We used digital image correlation  
134 (also known as feature tracking) to produce monthly-averaged and multi-year averaged velocity maps to investigate variations  
135 of glacier surface velocity in time and space over the selected glaciers. We can hereby summarize the workflow that was used  
136 (Fig. 2).

137 The input data are a DEM of the area of study, the RGI glacier outlines, the modelled glacier thickness and the stack of  
138 Sentinel-2 images in the reference period. The input DEM is used to obtain morphometric data of the glacier while the RGI  
139 glacier outlines are used, together with the selected satellite imagery, to choose suitable glaciers for surface glacier velocity  
140 analysis. After the glacier selection, we updated the RGI glacier outlines according to the current glacier extensions. Selected  
141 imagery is processed with digital image correlation to obtain glacier velocities. The glacier flowlines of the RGI are then used,  
142 together with the updated glacier outlines and the mapped equilibrium line altitudes (ELAs), to identify sampling areas to  
143 extract velocity time series. ELAs are manually digitized each year by the analysis of Sentinel-2 imagery. Subsequently, the  
144 time series are analysed to identify general trends, seasonal patterns or particular kinematic behaviours. Finally, the velocity  
145 dataset is analysed in relation to the morphological characteristics (in particular their size) of the glacier.



**Fig. 2 Workflow of the present study. The input datasets are evidenced in light blue while the processing steps are indicated in white boxes.**

### 3.2 Sentinel-2 optical satellite imagery

We adopted Sentinel-2 optical images acquired between 2016 and 2023. We chose to start the analysis in 2016 because it was the first full year of acquisition by the satellite. Based on different publications (Kääb et al., 2016; Millan et al., 2019), the geometric misregistration of Sentinel-2 can show up to 1.5 pixel offsets in the horizontal plane even if it can be usually closer to a value of 0.5 pixels, which corresponds to the absolute geolocation specification by ESA. Therefore, an image co-registration process or a correction of stable ground shifts is normally needed for multitemporal analyses.

To select the images, we defined the presented approach: (i) to maximize the geometric and geo-referencing precision, we adopted images acquired from the same orbit and tile (GRANULE T32TLR, relative orbit 108); (ii) to reduce the impact of clouds, we carried out a visual check of all images with a cloud cover percentage lower than 80% (as detected by the Copernicus cloud cover estimation algorithm) on the whole tile, which were 323 in total (150 from Sentinel 2B and 173 from Sentinel 2A). From this dataset, we extracted a subset of 123 cloud-free images on the selected glacier areas via the visual inspection of the individual images. We adopted this manual selection to maximize the quality of the images; in the case of the Mont Blanc massif (like in most mountainous areas worldwide), the local distribution of clouds can be extremely variable; in many cases, this can contribute to a considerable cloud percentage, even though high altitude areas may still be cloud-free.

The number of suitable images that are available per year varies from 10 to 20 with a yearly mean of 15 images in the following distribution: 2016: 10; 2017: 18; 2018: 11; 2019: 14; 2020: 13; 2021: 16; 2022: 20, 2023: 19, 2024: 2. The year 2016, and partially 2017, is influenced by the lack of Sentinel 2B images, which was launched on 7 March 2017.


To apply image correlation, we used the near-infrared band B08 at the processing level L1C as suggested by previous studies (Kääb et al., 2016).

### 169 3.3 Glacier selection

170 To minimize the presence of noisy and unreliable velocity data, we performed a selection of glaciers from the RGI 7.0  
171 dataset. In particular, we did not include in our study: A) glaciers with an area  $< 0.1 \text{ km}^2$ , as those glaciers would be too small  
172 for the reliable extraction of velocity maps with 10 m resolution optical satellite imagery (Millan et al., 2019); B) glaciers  
173 showing strong variations of cast shadow; C) glaciers that lack surface features to be tracked (e.g., ice caps).

174 Selection of point B) was made by creating a stack of images acquired between October and March, when cast shadows  
175 appear on satellite imagery, especially on north-facing slopes. Subsequently, we manually individuated glaciers that are subject  
176 to large variations of shadow on their surface. We used the scene classification map (SCL) class 11 (cast shadows), available  
177 in processing level L2A of the Sentinel-2 images. However, since shadows on glaciers may often be misclassified, we  
178 conducted a manual check to correct potential errors. Selection of point C) is made manually by selecting glaciers that show  
179 very even surfaces on Sentinel-2 images. This is normally noted in ice caps at higher altitudes or flat valley tongues.

180 The glacier selection process identified thirty glaciers with a total glacierised surface of  $85.8 \text{ km}^2$ . Compared to the total  
181 glacierised surface of the massif from RGI 7.0, this represents the covering of 50.8% of the total  $169 \text{ km}^2$  and 25.9% in terms  
182 of number of glaciers; this rises to 39.5% if we consider the subset of seventy-six glaciers having a glacierised surface of more  
183 than  $0.1 \text{ km}^2$ . The selected glaciers are highlighted in Fig. 3 and listed in Table 1. Two of the selected glaciers are located in  
184 Switzerland, ten are located in France and eighteen in Italy. This distribution is mainly due to a small portion of the massif  
185 being located in Switzerland and the presence of more fragmented glacierised bodies on the Italian side. Of the thirty glaciers,  
186 seven have been mapped as sub-areas in comparison to RGI 7.0 individual glacier bodies. A brief description of all the glaciers  
187 we analysed is found in section S1.1, to describe the location and geomorphological setting of the glaciers as well as highlight  
188 when a single glacier complex from the RGI 7.0 was divided into independent glacial bodies because of very distinct kinematic  
189 behaviour.



190  
191 **Fig. 3** Surface glacier velocity map averaged in the 2016-2023 period. Selected glaciers for specific analyses  
192 are outlined in cyan. Background: Sentinel-2 image (B08 band), courtesy of the Copernicus Open Access Hub  
193 (<https://scihub.copernicus.eu>, last access: 10 September 2023).

### 194 3.4 Glaciers' outline delineation and morphometric data calculation

195 Since the RGI 7.0 glacier outlines refer to 2003, we updated them to fit with the present glacier extensions and manually  
196 outlined them from Sentinel-2 imagery. We selected a cloud-free scene acquired on 28 August 2018 that represents well the  
197 conditions of the glaciers in the study period; True Color Image was used for this purpose. The main morphometric data that  
198 were determined for each glacier are summarized in Table 1. In some cases, a morphological indication that some parts of the  
199 glaciers could be considered independently from others and divided into individual kinematic domains was considered (Paul  
200 et al., 2022; Zemp et al., 2021). As the production of velocity maps confirmed distinct behaviours of some glacier parts, those  
201 glaciers were divided and mapped accordingly. The main examples are the tributary glaciers of the larger Miage Glacier  
202 complex, which all have distinct kinematic behaviour, well differentiated from the slow-moving, debris-covered main central  
203 valley tongue. Another example is the Talèfre Glacier, where, over the past twenty years, the western part of the glacier has  
204 become independent from the eastern portion.

205 The determination of morphometric data of sample glaciers was performed using altitudinal data from a 1 m resolution  
206 DEM obtained by the processing of Pleiades stereo pairs acquired in August 2018 (Berthier et al., 2014), while the mean



207 glacier thicknesses were extrapolated from globally modelled ice thickness data (Millan et al., 2022), which has an average  
 208 uncertainty of 30%.

209

210 **Table 1. Table with name and RGI identification codes of glaciers selected for analysis in the present study together**  
 211 **with main morphometric parameters. The elongation is the glacier length divided by its area.**

Glacier name	RGI 7.0 ID	Area (km <sup>2</sup> )	Length (m)	Min alt (m asl)	Max alt (m asl)	Avg slope (°)	Mean ice thickness (m)	Elongation (m <sup>-1</sup> )
A Neuve N	RGI2000-v7.0-G-11-00722	0.27	790	3084	3454	25.0	24	2.95
A Neuve Central	RGI2000-v7.0-G-11-00721	0.89	1800	2664	3554	26.3	26	2.02
Pre de Bard	RGI2000-v7.0-G-11-00716	3.01	3300	2360	3641	21.2	63	1.10
Greuvettaz E	RGI2000-v7.0-G-11-00708	0.20	990	2948	3582	32.8	12	5.03
Greuvettaz W	RGI2000-v7.0-G-11-00707	0.17	840	2704	3291	35.0	12	4.95
Planpincieux	RGI2000-v7.0-G-11-00703	1.01	2050	2627	3650	26.5	45	2.02
Grandes Jorasses	RGI2000-v7.0-G-11-00703	0.48	2110	2701	4206	35.5	15	4.38
Pra Sec	RGI2000-v7.0-G-11-00704	0.12	870	2536	3190	36.8	9	7.34
Rochefort	RGI2000-v7.0-G-11-00701	0.56	1000	2720	3301	30.3	25	1.78
Brenva	RGI2000-v7.0-G-11-00695	6.58	4490	2374	4766	28.0	80	0.68
Thoula	RGI2000-v7.0-G-11-00698	0.58	1080	2880	3416	26.5	25	1.85
Mont Blanc	RGI2000-v7.0-G-11-00688	0.76	2490	2776	3773	21.9	47	3.25
Dome	RGI2000-v7.0-G-11-00688	1.97	3550	2453	4121	25.1	51	1.80
Bionassay (IT)	RGI2000-v7.0-G-11-00688	1.35	2930	2467	3816	24.8	53	2.16


Aiguille Tre la Tete N	RGI2000-v7.0-G-11-00688	0.31	1360	2408	3010	24.0	78	4.34
Freney	RGI2000-v7.0-G-11-00693	1.02	2620	2420	3698	26.0	61	2.58
Brouillard	RGI2000-v7.0-G-11-00691	1.17	2730	2499	3972	28.3	52	2.34
Lex Blanche	RGI2000-v7.0-G-11-00674	2.64	2450	2467	3757	27.8	41	0.93
Petit Mont Blanc	RGI2000-v7.0-G-11-00674	0.56	1770	2863	3580	22.1	26	3.18
Estelette	RGI2000-v7.0-G-11-00673	0.29	950	2716	3214	27.7	24	3.26
Pierre Joseph	RGI2000-v7.0-G-11-00713	0.28	710	2920	3409	34.6	15	2.58
Nant Blanc	RGI2000-v7.0-G-11-00749	0.36	1150	2600	3351	33.1	33	3.17
Charpoua	RGI2000-v7.0-G-11-00751	0.32	1210	2650	3479	34.4	25	3.76
Des Glaciers	RGI2000-v7.0-G-11-00664	1.09	2050	2735	3815	27.8	31	1.88
Talèfre N	RGI2000-v7.0-G-11-00753	2.04	1950	2700	3550	23.6	39	0.96
Des Rognons	RGI2000-v7.0-G-11-00745	4.52	4290	2178	3800	20.7	102	0.95
Mer de Glace	RGI2000-v7.0-G-11-00757	23.56	12090	1774	4025	10.5	104	0.51
Bossons	RGI2000-v7.0-G-11-00773	11.32	6800	1691	4776	24.4	60	0.60
Tacconnaz	RGI2000-v7.0-G-11-00774	4.99	4290	2043	4286	27.6	40	0.86
Bionassay (FR)	RGI2000-v7.0-G-11-00778	4.77	5240	1835	4287	25.1	40	1.10

### 212 3.5 Glaciers' surface velocity calculation

213 Digital image correlation is a common technique used to measure surface displacements using proximal (Evans, 2000; Ahn  
214 and Box, 2010; Schwalbe and Maas, 2017; Dematteis et al., 2024) and remotely sensed imagery (Scambos et al., 1992; Heid  
215 and Kääb, 2012; Marsy et al., 2021; Dematteis and Giordan, 2021). The processing chain performed in the present study uses  
216 the open-source Glacier Image Velocimetry (GIV) toolbox (Van Wyk De Vries and Wickert, 2021). GIV uses frequency-based  
217 correlation, can efficiently process large datasets and has been shown to perform well on glacier surface velocity measurements  
218 at different test sites (Van Wyk De Vries and Wickert, 2021). GIV calculates stable-ground shifts to correct for georeferencing  
219 errors; therefore, we created a stable ground mask composed of non-glacierized terrain surrounding the massif, and fit a 2D  
220 second-degree polynomial to the residual velocities over stable ground in the x- and y- directions. To measure glacier surface  
221 velocities, we adopted the 'multi-pass' option which updates displacement estimates over multiple iterations, refining initial  
222 coarse chip size displacement calculations using progressively smaller chip sizes. The initial chip size is automatically defined  
223 by GIV and cannot be smaller than 32x32 px. The overlap between matching windows was 0.5. Velocities higher than 1500  
224  $\text{m yr}^{-1}$  were considered unrealistic and discarded. The velocity map resolution was set to 40 m, without resampling. Finally,  
225 we smoothed the velocity maps by applying a 3x3 median filter.

226 To produce the time series, given a specific image, we processed the first and second subsequent images (GIV order 2  
227 time-oversampling). The minimum and maximum temporal repeat cycles were respectively 10 and 120 days. Overall, we  
228 calculated 218 image pairs, with an average temporal baseline of 35 days (Sec. S8).

229 Subsequently, the velocities of image pairs were averaged on a monthly basis. We applied the weighted average included  
230 in GIV, where the weights are proportional to the fraction of time included in a given month over the total time gap between  
231 the image pairs. The presence of clouds or snow on the glacier surface made it impossible to extract reliable data in the  
232 following months: i) January 2017 ii) December 2020, January and February 2021 iii) September 2021. The gaps represent  
233 only five months on which we did not retrieve velocity data out of the eight years considered in the study (i.e., <5%). In Fig.  
234 3, we present a velocity map with a resolution of 40 m. This was obtained by averaging all the single monthly velocity maps  
235 in the study period (2016-2023). In Fig. 4, we present an example of the obtained velocity map and the distribution of sampling  
236 areas over several chosen glaciers. All other parameters of the GIV processing were set at default parameters (Supplementary  
237 Materials section S5).



238  
 239  
 240  
 241  
 242  
 243

**Fig. 4** a) Details of glacier surface velocity map averaged in the 2016-2023 period and sampling areas of selected sample glaciers. b, c) show monthly velocity maps of July and February 2018 respectively. Sentinel-2 imagery base map (B08 band), courtesy of the Copernicus Open Access Hub (<https://scihub.copernicus.eu>, last access: 10 September 2023).

### 244 3.6 Velocity time series analysis

245 Sampling areas were identified on the velocity maps to analyse the time series of the selected glaciers. Since the surface  
246 velocity of land-terminating glaciers is expected to be the highest in correspondence with the ELA (Nesje, 1992), we cropped  
247 the RGI 7.0 flowlines at the upper and lower altitudinal limits of the ELAs mapped between 2016 and 2023. Around this  
248 section of the ELA, we buffered an area of 40 m (Fig. 4). The velocity values included in the obtained polygons were averaged  
249 to produce the time series.


250 Subsequently, we applied a quadratic locally weighted scatterplot smoothing (LOWESS) (Cappellari et al., 2013) evaluated  
251 on a rolling window of twelve months. Finally, we calculated the linear trend of the smoothed series using the Huber loss  
252 regressor (Huber, 1992). The Huber fit is robust against outliers; thus, sharp velocity fluctuations should not affect the obtained  
253 linear trends. To evaluate the statistical significance level of the fit, we consider the *t-statistics*, i.e., the ratio between the  
254 coefficient values and their standard errors. According to its definition, the *t-statistics* is expected to be low when the slope of  
255 the linear fit is low and/or when the slope is relatively low compared to the data variability.

256 We performed the principal component analysis (PCA) of the time series to investigate the overall behaviour of all the  
257 considered glaciers, and we weighted the time series according to the glacier area. The PCA is a multivariate analysis technique  
258 that allows a reduction in the dimensionality of a given dataset, increasing interpretability but minimising information loss.  
259 This is achieved by creating new, uncorrelated variables that successively maximise the variance of the dataset (Jolliffe and  
260 Cadima, 2016).

## 261 4 Results and discussion

### 262 4.1 Distributions of monthly velocities

263 On the thirty glacial bodies we investigated in this study, the monthly velocity values range from 30-40 m yr<sup>-1</sup>, typically  
264 reached during winter months, to 350 and 400 m yr<sup>-1</sup>, typically reached in summer/late summer. The highest velocities are  
265 attained by the two large glaciers: the Brenva and the Bossons glaciers. In particular, monthly extreme values vary from 11.8  
266 +/- 9.8 m yr<sup>-1</sup>, reached on January 2022 by the Des Glaciers Glacier, to 487.4 +/- 10.8 m yr<sup>-1</sup>, reached by the Brenva Glacier  
267 in July 2016. The mean surface velocities averaged over the whole period range from 29.9 +/- 10.9 m yr<sup>-1</sup> at the Pierre-Joseph  
268 Glacier to 375.9 +/- 10.9 m yr<sup>-1</sup> at the Brenva Glacier. The standard deviation of the velocity time series of single glaciers  
269 varies from 10.3 m yr<sup>-1</sup> at the Pierre Joseph Glacier to 70.7 m yr<sup>-1</sup> at the Charpoua Glacier. Fig. 5 presents the distributions of  
270 the raw monthly velocity of the considered glaciers.



271  
272 **Fig. 5. Distributions of the glaciers' raw monthly velocity. Glaciers are sorted by their median velocity. The**  
273 **background colours of the glacier names indicate their size: white < 1 km<sup>2</sup>; green 1-4 km<sup>2</sup>; blue >4 km<sup>2</sup>.**

274 **4.2 Velocity time series**

275 Large and thick glaciers have in general high velocities, and two of them (Brenva and Bossons) show the highest values.  
276 The velocity values remain high throughout the whole year (Fig. 6). Brenva displays irregular seasonality variations, while  
277 Bossons has an almost absent seasonal cycle. Their morphology is complex - e.g., the slope varies considerably along their  
278 extent. In the present study, we concentrated the analyses on the middle sectors (near the ELA) of the glaciers, which are the  
279 fastest (Nesje, 1992).

280 Medium-sized glaciers' morphology is less homogeneous, even though most are generally gentler and thicker than average.  
281 We can divide the group into two sets based on the elongation ratio. Medium-sized glaciers which are more elongated generally  
282 feature strong kinematic activity, their velocity is higher than average with marked variability and they often show a  
283 pronounced regular annual cycle, as in the cases of the Bionassay (IT), Brouillard, Mont Blanc and Freney glaciers. The second  
284 set of more "compact" glaciers, show lower mean velocities and have a minor amplitude of the seasonal variability. In  
285 particular, even though some display a regular seasonal cycle (e.g., Planpincieux, Dome, Rochefort), their velocity variability  
286 is much lower (Fig. 6).

287 Small glaciers show lower average velocities, but most show marked and regular seasonality. In this group, some very  
288 small glaciers (e.g., Greuvettaz W, Greuvettaz E, A Neuve N, Pierre Joseph) show a modest, irregular or even non-detectable

289 seasonal cycle since the velocity in winter is close to the measurement uncertainty (Fig. 6). It is worth highlighting that signals  
290 of potential velocity fluctuations could exist but remotely-sensed data are not currently suited for analysing such small glaciers.  
291 On the other hand, Pra Sec and Charpoua, small steep elongated glaciers, feature very low minimum velocities (between  
292  $25 \text{ m yr}^{-1}$  and  $50 \text{ m yr}^{-1}$ ) and very marked peaks during which the velocity increased by one order of magnitude. These high  
293 velocity periods appear in summer/late summer and extreme velocity changes from winter to summer velocities can be noticed.  
294 In 2016, 2018 and 2022, the summer velocity of Charpoua Glacier was exceptionally high compared to the usual. Moreover,  
295 in these years, the spatial distribution of velocities varied, and the highest velocities were registered towards the frontal part of  
296 the glacier, while normally it was higher in the upper sector (Fig. 7). Pra Sec Glacier displays more regular annual speedups  
297 in the period analysed, every summer it had a clear velocity peak (except in 2018), particularly pronounced in 2016, 2020 and  
298 2022. In both glaciers, possible glacier advances are prevented by the steep bedrock cliff at the snout, which causes the  
299 disintegration of the glacier by repeated ice falls from the terminus (Giordan et al., 2020; Pralong and Funk, 2006).





Fig. 6. Time series of monthly glacier surface velocities over the 2016-2024 period. Robust linear trends are represented in dashed black lines. The background colour of the glaciers' names denotes their size: white, green and blue are for small, medium and large glaciers, respectively. The velocity seasonality is indicated with markers: brown diamonds (low seasonality), orange circles (average seasonality) and pink rectangles (high seasonality).






301  
302  
303 **Fig. 7. Charpoua Glacier monthly surface velocity maps showing the spatial variation of the velocity patterns between**  
304 **July 2018 (upper left) and July 2021 (upper right). The lower panel shows the monthly velocity profiles along a**  
305 **longitudinal east-west AB profile (in black on the maps) over the study period. Sentinel-2 imagery base map (B08**  
306 **band), courtesy of the Copernicus Open Access Hub (<https://scihub.copernicus.eu>, last access: 10 September 2023)**

### 307 4.3 Interannual velocity variability and trend

308 The overall behaviour of the glacier velocity can be well represented by the PCA of the time series, weighted by the glacier  
309 area. Fig. 8 shows the 1<sup>st</sup> PC, which explained >50% of the variance. In general, this analysis reflects a common trend of many  
310 time series: a first period between 2016 and 2019 showing decreasing velocities, an anomaly of higher velocities between 2020  
311 and 2022 followed by a new velocity decrease in 2023 (Fig. 6). According to historical observations at the Argentière and  
312 Miage glaciers, the velocity decrease from the early 2000s can be linked to continuous negative mass balances (Vincent and  
313 Moreau, 2016) of most Alpine glacier since the 2000s (Zemp et al., 2021). The results of our study agree with this negative  
314 trend in the first part of the considered period (i.e., 2016–2019), but we detected a rupture in the trend and a velocity rise from  
315 2020 to 2022 occurring in a large number of glaciers under study. An interesting remark about the geographical distribution  
316 of the trends is that the glaciers showing a clear velocity anomaly during 2020-2022 are located on the southeast side of the

317 massif ridgeline along the Italian and Swiss part of the Massif (e.g., A Neuve Central, Bionassay (IT), Brenva, Planpincieux  
318 and Lex Blanche glaciers). This might be linked to different dominant meteorological conditions distributions on the two sides  
319 of the massif. A specific analysis of seasonal meteorologic data in the different areas could give more insights into this  
320 hypothesis, even though whether stations are located far from the glacier accumulation areas and could not get a specific signal  
321 present in the higher altitude sectors.



322  
323

**Fig. 8. 1<sup>st</sup> PC of the LOWESS velocity values of all the time series of glacier monthly velocity.**

324 The glacier velocities shown in Fig. 6 have different linear trends over the full period of the study (Fig. 9). In general, the  
325 linear trends of each glacier calculated using winter, summer months or the whole year have similar behaviour (i.e., they are  
326 always negative or positive), with winter trends having higher values than summer ones (~30% higher on average). As  
327 expected, lower absolute linear trends (i.e.,  $\leq 1 \text{ m yr}^{-1}$ ) have  $t$ -statistics  $< 2$ . Besides, even though Brenva's trend is high ( $2.8 \text{ m}$   
328  $\text{yr}^{-2}$ ), its velocity is very large, thus lowering the  $t$ -statistics. Most trends lie in a relatively tight cluster which crosses the  
329 domain diagonally, except for Charpoua and Planpincieux which have dissimilar seasonal values. In the Charpoua case, this  
330 is probably related to the strong velocity fluctuations occurring only in some years in summer, while the Planpincieux positive  
331 summer trend is likely led by the anomaly 2020-2022, which is particularly strong in this glacier. Notably, the three glaciers  
332 with the highest linear trends (i.e., Freney, Bionassay (IT) and Aiguille de Tré-la-Tete) are all located in Val Veny, on the  
333 southern side of the massif. Bionassay (IT) and Freney have a very similar morphology (Tab. 1), they are both medium-size  
334 ( $1.3 \text{ km}^2$  and  $1.02 \text{ km}^2$ ), relatively elongated ( $2.2 \text{ m}^{-1}$  and  $2.5 \text{ m}^{-1}$ ) glaciers, with slopes of  $\sim 25^\circ$  and elevations between 2400  
335 m and 3800 m. Differently, Aiguille de Tré-la-Tete is a small ( $0.3 \text{ km}^2$ ), much-elongated ( $4.34 \text{ m}^{-1}$ ) and low-altitude glacier  
336 (2408-3010 m). Aiguille de Tré-la-Tete and Bionassay (IT) are both tributaries of the Miage Glacier.

337 An accelerating trend ( $+5 \text{ m yr}^{-2}$  between 2015-2021) has recently been shown for the Brenva Glacier by an analysis of  
338 remotely-sensed optical images (Rabatel et al., 2023b) while detecting decelerating trends on many other glaciers of the Massif.  
339 Our study showed Brenva Glacier an accelerating trend of  $+3 \text{ m yr}^{-2}$  between 2016-2023, which is in good agreement  
340 considering the velocity decrease shown in 2022 and 2023. Besides, (Rabatel et al., 2023b) observed a slight ice thickening  
341 ( $\sim 1 \text{ m}$  between 2000 and 2019) in an upper sector of the Brenva Glacier. They proposed three hypotheses to explain the


342 acceleration of the Brenva: a) a glacier thickening; b) a change in thermal regime; and c) a change in subglacial hydrology,  
343 possibly related to an increased ablation in the upper reaches of the glacier.

344 Even though the hypothesis of glacier thickening could explain the specific case of Brenva, the glacier surface elevation  
345 change across the Mont Blanc massif has been generally negative in the last years, as evidenced by the negative mass balance  
346 of the reference glaciers in the area (Zemp et al., 2021; Zemp et al., 2009). Local anomalies of positive mass balance could  
347 explain an increase of velocity but the lack of measurements at higher altitudes does not allow us to confirm this behaviour at  
348 present. However, the meteorological conditions in recent years have remained approximately constant, which makes unlikely  
349 a general glacier thickening in the region (see S6 “Meteorological conditions”). Localized high rates of accumulation due to  
350 increased avalanche activity and wind accumulation could also contribute to the ice thickening (thus yielding an acceleration),  
351 but cannot be investigated at this stage. It is worth highlighting that the three glaciers highlighting a high winter accelerating  
352 trend in Fig. 9 (Bionassay (IT), Aig Tré la Tete and Freney) are near to each other and located in a small part of the massif on  
353 its southeast side.

354 A change in the thermal regime of the glaciers could explain accelerating trends, but it would explain long-term trends and  
355 not short-period variations such as the one highlighted in 2020-2022, as basal temperature measurements show a warming at  
356 the ice bedrock interface on a decadal to multi-decadal timescale on the Mont Blanc massif (Vincent et al., 2007).

357 A variation of the hydrology of groups of glaciers is a plausible hypothesis for the explanation of accelerating trends, but  
358 it should highlight stronger trends in summer rather than in winter, as basal sliding is enhanced over deformation during  
359 summer. Such a combination of trends is not shown by most glaciers as highlighted in Fig. 9. Only the trends highlighted by  
360 Planpincieux and Mer de Glace glaciers could relate well with this hypothesis, showing almost no trend in winter and an  
361 accelerating trend during the summer months.

362 The distribution of the acceleration trend over different areas of the massif and regarding different types of glaciers suggest  
363 the existence of a long-term meteo-climatic driver of the phenomenon, even though it is not evident limiting the analysis to  
364 the period 2015-2023 (Section S6). In the end, a definitive answer cannot be formulated so far and further research is necessary  
365 to understand the processes involved in this trend.



**Fig 9. Linear trends of the glacier monthly velocity. The x- and y-axes refer to the trends calculated using winter (from November to April) and summer (from June to September) months, respectively. The colours indicate the global trend. The glaciers with linear trends with  $t$ -statistic  $< 2$  have markers with dashed edge lines.**


#### 4.4 Relationship between glacier size, velocity and seasonal behaviour

We examined the relationship between the glacier size, the seasonal velocity behaviour and velocity distribution. To this end, we divided the glaciers into three classes for each feature. Concerning the size, we considered very small glaciers with an area of  $< 1 \text{ km}^2$ , according to (Bahr and Radić, 2012), while medium and large glaciers have areas between  $1 \text{ km}^2$  and  $4 \text{ km}^2$ , and  $> 4 \text{ km}^2$ , respectively. Concerning the velocity distribution, we observed that, besides Brenva and Bossons, which have much higher velocity compared to the rest of the glaciers, a group of sixteen glaciers have the 75<sup>th</sup> percentile of monthly velocity  $< 100 \text{ m yr}^{-1}$  (from Pierre Joseph to Charpoua in Fig. 5). Finally, twelve glaciers have their 25<sup>th</sup> and 75<sup>th</sup> percentiles between  $\sim 100 \text{ m yr}^{-1}$  and  $\sim 200 \text{ m yr}^{-1}$  (from Des Rognons to Freney in Fig. 5).

Concerning the seasonal velocity behaviour, we analysed the raw time series (Fig. 6) and the LOWESS smoothed time series normalised by their median value (Fig. S6). Glaciers such as Freney, Brouillard, and Bionassay (IT) show evident and regular seasonal behaviour and large winter/summer differences; in these cases, summer velocities (occurring between July and October) are 50% to 100% higher than winter ones (occurring between January and April). Another group of glaciers has smaller winter/summer differences or pronounced but irregular variability (e.g., Planpincieux, Pre de Bard, Talefre). A third group does not display evident or regular seasonal behaviour (e.g., Taconnaz, Mer de Glace, Pierre Joseph), with winter/summer differences below 10%.

385 The double-entry heatmaps are presented in Fig. 10, which show a tendency for the smaller glaciers to have more  
 386 pronounced seasonality, while larger glaciers show a more homogeneous velocity throughout the year. Since the glacier area  
 387 is strongly correlated with its thickness (Cuffey and Paterson, 2010), a possible cause of this phenomenon could be related to  
 388 enhanced basal sliding during the accelerating period. In fact, a thicker glacier could be less prone to exhibit enhanced sliding  
 389 because of the larger mass to be uplifted by basal positive water pressures, while shallower glaciers could more easily benefit  
 390 from enhanced sliding by pressure build-up at the ice-bedrock interface by increasing inputs in the hydrological subglacial  
 391 drainage network. Moreover, in winter, the base of thin glaciers could freeze, thus preventing the sliding and determining  
 392 lower winter velocities causing even larger summer-winter velocity differences.

393 On the other hand, as expected, larger (and thicker) glaciers tend to be faster than smaller ones.




394 **Fig. 10** Glacier classification according to their size (SMALL: <1 km<sup>2</sup>; MED: 1-4 km<sup>2</sup>; LARGE: >4 km<sup>2</sup>) and a)  
 395 seasonal velocity behaviour (qualitatively estimated) and b) median velocity (SLOW: <100 m yr<sup>-1</sup>; AVG: 100-300 m  
 396 yr<sup>-1</sup>; FAST: >300 m yr<sup>-1</sup>). The digits within the grid tiles indicate the number of glaciers belonging to each group.  
 397

#### 398 4.5 Uncertainty analysis

399 To estimate the quality of our data, we performed two investigations. First, following the method proposed by (Millan et  
 400 al., 2019), we calculated the median absolute deviation of the velocity obtained on stable terrain for each monthly data. In  
 401 these areas, we applied an outlier spatial filter to the velocity maps according to Rabatel et al. (2023a). The median value of  
 402 the monthly uncertainty was 10.9 m yr<sup>-1</sup>. In their study, Millan et al. (2019) estimated the nominal precision according to the  
 403 temporal baseline between the correlated images, which they found to be between 6–16 m yr<sup>-1</sup> for baselines respectively of 40  
 404 and 20 days, which is the typical range of temporal gaps between images used in our study. Moreover, the value of 10.9 m yr<sup>-1</sup>  
 405 is in close agreement with the uncertainty found by Mouginot et al. (2023), which obtained a root mean squared error of 10.5  
 406 m yr<sup>-1</sup> between glacier velocities measured over the Mer de Glace and Argètiere glaciers using image correlation of Sentinel-  
 407 2 images and GNSS in situ data (<https://glacioclim.osug.fr/>). A comparison with the data from Rabatel et al. (2023) is proposed  
 408 in the supplementary materials (section S4, figure S3) for the glaciers of Brenva, Bionassay (FR), Bossons and Brouillard over  
 409 the timespan that overlaps the two studies (2016-2021).

410 Second, we considered the glacier velocity from Millan et al. (2019), who published mean annual velocity in the period  
 411 2017–2018 on a 50x50 m regular grid. They adopted normalised cross-correlation and chip size refinement (initial size of

412 16x16 px). They estimated an overall uncertainty of glacier surface velocity time series of  $\sim 12 \text{ m yr}^{-1}$  over the Mont Blanc  
413 glaciers, and, specifically at Brenva and Bosson glaciers, an uncertainty of  $15\text{--}20 \text{ m yr}^{-1}$ . We compared these data and ours  
414 along four glacier longitudinal central lines (i.e., in Bossons, Brenva, Freney and Taconnaz), obtaining good agreement (Fig.  
415 11). The largest differences ( $>50 \text{ m yr}^{-1}$ ) were found in a specific sector of the Taconnaz Glacier (Fig. 11d), where the ice flux  
416 is highly channelized in a narrow passage. There, the data of Millan et al. (2019) show a large velocity decrease that seems  
417 unlikely considering the geometry of the site. Our data show a similar but less pronounced velocity decrease. However, the  
418 velocity profiles are similar elsewhere. On average, the surface velocities that we obtained are slightly higher, with a mean  
419 difference of  $0.03 \text{ m yr}^{-1}$  and root mean squared deviation (RMSD) of  $24.0 \text{ m yr}^{-1}$  (Table 2). The slightly higher RMSD  
420 compared to the expected uncertainty can be due to the fact that the precision in glacierized areas is probably larger (less  
421 precise) than in ice-free zones because the surface texture is different and changes (e.g., snow precipitation, surface melt,  
422 glacier movement) occur more rapidly, therefore causing more decorrelation (Millan et al., 2019).



423

424 **Fig 11. Comparison of velocity profiles from Millan et al. (2022) (red) and from this study (black) at (a) Bossons**  
 425 **Glacier, (b) Brenva Glacier, (c) Freney Glacier and (d) Tacconnaz Glacier. The velocity profiles are represented in**  
 426 **yellow in the maps on the left, and the velocity profiles start from the upper to the lower altitudes. Sentinel-2 image**  
 427 **base map (B08 band), courtesy of the Copernicus Open Access Hub (<https://scihub.copernicus.eu>, last access: 10**  
 428 **September 2023).**

429

430 **Table 2. Mean difference and root mean squared deviation (RMSD) between this study and Millan et al. (2022) along velocity**  
 431 **longitudinal profiles.**

	Bossons	Brenva	Freney	Taconnaz	Mean all profiles
Mean difference [m yr <sup>-1</sup> ]	-13.1	-8.8	13.7	8.3	0.03
RMSD [m yr <sup>-1</sup> ]	22.3	25.2	29.6	19.0	24.0

#### 437 4.6 Uses and limits of the proposed methodology

438 The methodology presented in this study allows the detection of monthly changes in glacier velocity, which can be  
 439 precursors of ice avalanches (Pralong et al., 2005; Faillettaz et al., 2008; Giordan et al., 2020). For example, ice avalanches  
 440 from the Pra Sec Glacier occurred in 2020 (Forestry Service of Aosta Valley). In our study, we observed high velocities in  
 441 2020, which could have led to the break-off. At the Charpoua Glacier, an ice avalanche of 45000 +/- 15000m<sup>3</sup> occurred in  
 442 2018 (Lehmann, 2018 - <https://news.unil.ch/display/1536777918113>, accessed online 11 October 2023), when we measured  
 443 velocities >200 m yr<sup>-1</sup>, much higher than usual.

444 In this frame, it would be very relevant to measure and know the typical velocity fluctuations of specific glaciers in stable  
 445 conditions. This could allow an assessment and to what extent a suspect acceleration may be anomalous and potentially  
 446 destabilising, bearing in mind that high-rate monitoring is essential to detect glacial instabilities since the expected sharp  
 447 increase in velocity in the weeks before the failure (Pralong and Funk, 2006) could be hardly detectable from remote sensing  
 448 (e.g., due to scarce visibility, image decorrelation, low resolution).

449 Limits of the methodology presented in this study should also be considered: glaciers moving at slow rates can be surveyed  
 450 using temporal baselines of one year (Millan et al., 2019; Mouginot et al., 2023), but this implies reducing the ability to catch  
 451 short-term velocity fluctuations, like those observed at Charpoua and Pra Sec glaciers. Another known issue pertains to the  
 452 lack of features of the glacier surface that make it impossible to track movements using optical imagery. Satellite optical  
 453 imagery is limited and can be strongly influenced by the presence of clouds that could yield extensive periods without data  
 454 acquisition, even though, in the present study, we had only four of 96 months with no data. Anomalies due to image  
 455 decorrelation for the presence of shadows, snow or morphological surface modifications can occur and an expert-based visual  
 456 check may be required to discriminate anomalous velocities.

#### 457 6 Conclusions

458 We produced ice velocity maps and time series of thirty glaciers of the Mont Blanc massif during the period 2016–2023.  
 459 The proposed results are different compared to the existing publicly available automatically processed velocity datasets that  
 460 have a coarse resolution (i.e., >100 m) and cannot correctly detect the kinematics of most Alpine glaciers, due to their small  
 461 size. Therefore, specific processing and studies are needed to characterize the surface kinematics of Alpine glaciers. In our  
 462 study, we used Sentinel-2 imagery due to its free availability, good ground resolution and high revisit time in the study area to  
 463 obtain monthly surface velocity time series. In addition, we proposed a classification of different groups of glaciers based on  
 464 their morpho-kinematic features. Plus, we observed a significative acceleration trend in many of the studied glaciers in the last  
 465 years (2020–2022), but the causes are still not well understood.



466 From the methodological point of view, the use of the proposed approach can be very useful to process and analyse  
467 available satellite images of other massifs in the Alps and other parts of the World. This approach could stimulate innovative  
468 research on high-resolution spatiotemporal variations of velocities on alpine glaciers and, especially, on understanding the  
469 variations in the motion of mountain glaciers. A large research question remains open and deals with understanding and  
470 measuring the drivers of change in the motion of alpine glaciers. This implicates the complex acquisition of data related to the  
471 possible drivers of the variations such as mass balances, water inputs and temporal variations in the subglacial hydrology of  
472 single glaciers. However, in order to delve further into these investigations, more complete and widespread glacier surface  
473 velocity databases, such as that presented in this study, are needed to build future research on the topic.

#### 474 **Acknowledgements**

475 The authors would like to thank Jean Pierre Fosson, Raffaele Rocco, Valerio Segor, and Guido Giardini for their support  
476 and their will to stimulate cryospheric research activities in the Aosta Valley region and on the Italian side of Mont Blanc  
477 massif. We thank all the staff at Fondazione Montagna Sicura for supporting all the research activities of the research team.  
478 Dr Etienne Berthier supplied a processed 2018 Pleiades stereo DEM on which retrieving altitudinal data that was presented in  
479 this paper. Dr. Antoine Rabatel supplied velocity data for the comparison with their 2023 dataset. We thank Prof. Christian  
480 Vincent for his advice on research activity and for supplying useful additional information, especially on the French glaciers  
481 of Mont Blanc.

#### 482 **Data availability**

483 The following data are available for download at xxxxxxxx.xxx: i) monthly velocity maps, ii) updated outlines (from RGI7)  
484 of the analysed glaciers, iii) sampling areas of extraction of the velocity time series, iv) time series of monthly velocity. The  
485 data description and format are reported in Sec S9. Sentinel-2 imagery is available from the Copernicus Open Access Hub  
486 (<https://scihub.copernicus.eu>, Copernicus, 2022). The GIV toolbox is freely available online  
487 (<https://github.com/MaxVWDV/glacier-image-velocimetry>).

488 **Competing interests.** The authors declare that they have no conflict of interest.

489 **Author contributions.** Fabrizio Troilo: Conceptualization, writing – original draft preparation, investigation,  
490 methodology, data curation, formal analysis, visualization; Niccolò Dematteis: Writing – review & editing, data curation,  
491 methodology, formal analysis, validation, visualization; Francesco Zucca Writing – review & editing, methodology,  
492 supervision, validation; Martin Funk: Writing – review & editing, supervision; Daniele Giordan: Writing – review & editing,  
493 methodology, supervision, validation.

#### 494 **References**

495 Ahn, Y. and Box, J. E.: Glacier velocities from time-lapse photos: technique development and first results from the Extreme  
496 Ice Survey (EIS) in Greenland, *Journal of Glaciology*, 56, 723-734, 2010.  
497 Allstadt, K., Shean, D., Campbell, A., Fahnestock, M., and Malone, S.: Observations of seasonal and diurnal glacier velocities  
498 at Mount Rainier, Washington, using terrestrial radar interferometry, *The Cryosphere*, 9, 2219-2235, 2015.

499 Arendt, A., Bliss, A., Bolch, T., Cogley, J., Gardner, A., Hagen, J.-O., Hock, R., Huss, M., Kaser, G., and Kienholz, C.:  
500 Randolph Glacier inventory—A dataset of Global glacier outlines: Version 6.0: Technical report, Global land ice measurements  
501 from space, 2017.

502 Bahr, D. and Radić, V.: Significant contribution to total mass from very small glaciers, *The Cryosphere*, 6, 763-770, 2012.

503 Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., and  
504 Huss, M.: The European mountain cryosphere: a review of its current state, trends, and future challenges, *The Cryosphere*, 12,  
505 759-794, 2018.

506 Benn, D. I. and Evans, D. J.: *Glaciers & glaciation*, Routledge 2014.

507 Beraud, L., Cusicanqui, D., Rabatel, A., Brun, F., Vincent, C., and Six, D.: Glacier-wide seasonal and annual geodetic mass  
508 balances from Pléiades stereo images: application to the Glacier d'Argentière, French Alps, *Journal of Glaciology*, 69, 525-  
509 537, 2023.

510 Berthier, E., Vadon, H., Baratoux, D., Arnaud, Y., Vincent, C., Feigl, K., Remy, F., and Legresy, B.: Surface motion of mountain  
511 glaciers derived from satellite optical imagery, *Remote Sensing of Environment*, 95, 14-28, 2005.

512 Berthier, E., Vincent, C., Magnússon, E., Gunnlaugsson, Á., Pitte, P., Le Meur, E., Masiokas, M., Ruiz, L., Pálsson, F., and  
513 Belart, J.: Glacier topography and elevation changes derived from Pléiades sub-meter stereo images, *The Cryosphere*, 8, 2275-  
514 2291, 2014.

515 Bindschadler, R.: The importance of pressurized subglacial water in separation and sliding at the glacier bed, *Journal of*  
516 *Glaciology*, 29, 3-19, 1983.

517 Cappellari, M., McDermid, R. M., Alatalo, K., Blitz, L., Bois, M., Bournaud, F., Bureau, M., Crocker, A. F., Davies, R. L., and  
518 Davis, T. A.: The ATLAS3D project—XX. Mass–size and mass– $\sigma$  distributions of early-type galaxies: bulge fraction drives  
519 kinematics, mass-to-light ratio, molecular gas fraction and stellar initial mass function, *Monthly Notices of the Royal*  
520 *Astronomical Society*, 432, 1862-1893, 2013.

521 Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Academic Press 2010.

522 Deilami, K. and Hashim, M.: Very high resolution optical satellites for DEM generation: a review, *European Journal of*  
523 *Scientific Research*, 49, 542-554, 2011.

524 Dematteis, N. and Giordan, D.: Comparison of digital image correlation methods and the impact of noise in geoscience  
525 applications, *Remote Sensing*, 13, 327, 2021.

526 Dematteis, N., Giordan, D., Troilo, F., Wrzesniak, A., and Godone, D.: Ten-Year Monitoring of the Grandes Jorasses Glaciers  
527 Kinematics. Limits, Potentialities, and Possible Applications of Different Monitoring Systems, *Remote Sensing*, 13, 3005,  
528 2021.

529 Dematteis, N., Troilo, F., Scotti, R., Colombarolli, D., Giordan, D., and Maggi, V.: The use of terrestrial monoscopic time-  
530 lapse cameras for surveying glacier flow velocity, *Cold Regions Science and Technology*, 104185, 2024.

531 Einarsson, B., Magnússon, E., Roberts, M. J., Pálsson, F., Thorsteinsson, T., and Jóhannesson, T.: A spectrum of jökulhlaup  
532 dynamics revealed by GPS measurements of glacier surface motion, *Annals of Glaciology*, 57, 47-61, 2016.

533 Evans, A. N.: Glacier surface motion computation from digital image sequences, *IEEE Transactions on Geoscience and Remote*  
534 *Sensing*, 38, 1064-1072, 2000.

535 Fahnestock, M., Scambos, T., Moon, T., Gardner, A., Haran, T., and Klinger, M.: Rapid large-area mapping of ice flow using  
536 Landsat 8, *Remote Sensing of Environment*, 185, 84-94, 2016.

537 Faillietaz, J., Pralong, A., Funk, M., and Deichmann, N.: Evidence of log-periodic oscillations and increasing icequake activity  
538 during the breaking-off of large ice masses, *Journal of Glaciology*, 54, 725-737, 2008.

539 Fyffe, C. L.: *The hydrology of debris-covered glaciers*, University of Dundee, 2012.

540 Giordan, D., Dematteis, N., Allasia, P., and Motta, E.: Classification and kinematics of the Planpincieux Glacier break-offs  
541 using photographic time-lapse analysis, *Journal of Glaciology*, 66, 188-202, 2020.

542 Glen, J.: Experiments on the deformation of ice, *Journal of Glaciology*, 2, 111-114, 1952.

543 Gottardi, F., Obled, C., Gailhard, J., and Paquet, E.: Statistical reanalysis of precipitation fields based on ground network data  
544 and weather patterns: Application over French mountains, *Journal of hydrology*, 432, 154-167, 2012.

545 Heid, T. and Kääb, A.: Evaluation of existing image matching methods for deriving glacier surface displacements globally  
546 from optical satellite imagery, *Remote Sensing of Environment*, 118, 339-355, 2012.

547 Huber, P. J.: Robust estimation of a location parameter, in: *Breakthroughs in statistics: Methodology and distribution*, Springer,  
548 492-518, 1992.

549 Humbert, A., Greve, R., and Hutter, K.: Parameter sensitivity studies for the ice flow of the Ross Ice Shelf, Antarctica, *Journal*  
550 *of Geophysical Research: Earth Surface*, 110, 2005.

551 Jiskoot, H.: Dynamics of Glaciers, *physical Research*, 92, 9083-9100, 2011.

552 Jolliffe, I. T. and Cadima, J.: Principal component analysis: a review and recent developments, *Philosophical transactions of*  
553 *the royal society A: Mathematical, Physical and Engineering Sciences*, 374, 20150202, 2016.

554 Kääb, A., Winsvold, S., Altena, B., Nuth, C., Nagler, T., and Wuite, J.: Glacier Remote Sensing Using Sentinel-2. Part I:  
555 Radiometric and Geometric Performance, and Application to Ice Velocity, *Remote Sensing*, 8, 2016.

556 Kääb, A., Jacquemart, M., Gilbert, A., Leinss, S., Girod, L., Huggel, C., Falaschi, D., Ugalde, F., Petrakov, D., and  
557 Chernomorets, S.: Sudden large-volume detachments of low-angle mountain glaciers—more frequent than thought?, *The*  
558 *Cryosphere*, 15, 1751-1785, 2021.

559 Kamb, B.: Glacier surge mechanism based on linked cavity configuration of the basal water conduit system, *Journal of*  
560 *Geophysical Research: Solid Earth*, 92, 9083-9100, 1987.

561 Luzi, G., Pieraccini, M., Mecatti, D., Noferini, L., Macaluso, G., Tamburini, A., and Atzeni, C.: Monitoring of an alpine glacier  
562 by means of ground-based SAR interferometry, *IEEE Geoscience and Remote Sensing Letters*, 4, 495-499, 2007.

563 Marsy, G., Vernier, F., Trouvé, E., Bodin, X., Castaings, W., Walpersdorf, A., Malet, E., and Girard, B.: Temporal Consolidation  
564 Strategy for Ground-Based Image Displacement Time Series: Application to Glacier Monitoring, *IEEE Journal of Selected*  
565 *Topics in Applied Earth Observations and Remote Sensing*, 14, 10069-10078, 2021.

566 Millan, R., Mouginit, J., Rabatel, A., and Morlighem, M.: Ice velocity and thickness of the world's glaciers, *Nature*  
567 *Geoscience*, 15, 124-129, 2022.

568 Millan, R., Mouginit, J., Rabatel, A., Jeong, S., Cusicanqui, D., Derkacheva, A., and Chekki, M.: Mapping surface flow  
569 velocity of glaciers at regional scale using a multiple sensors approach, *Remote Sensing*, 11, 2498, 2019.

570 Mondardini, L., Perret, P., Frasca, M., Gottardelli, S., and Troilo, F.: Local variability of small Alpine glaciers: Thoula Glacier  
571 geodetic mass balance reconstruction (1991-2020) and analysis of volumetric variations, *Geografia Fisica e Dinamica*  
572 *Quaternaria*, 44, 29-38, 2021.

573 Mouginit, J., Rabatel, A., Ducasse, E., and Millan, R.: Optimization of Cross Correlation Algorithm for Annual Mapping of  
574 Alpine Glacier Flow Velocities; Application to Sentinel-2, *IEEE Transactions on Geoscience and Remote Sensing*, 61, 1-12,  
575 2023.

576 Nesje, A.: Topographical effects on the equilibrium-line altitude on glaciers, *GeoJournal*, 27, 383-391, 1992.

577 Nye, J. F.: The mechanics of glacier flow, *Journal of Glaciology*, 2, 82-93, 1952.

578 Paul, F., Winsvold, S. H., Kääb, A., Nagler, T., and Schwaizer, G.: Glacier remote sensing using Sentinel-2. Part II: Mapping  
579 glacier extents and surface facies, and comparison to Landsat 8, *Remote Sensing*, 8, 575, 2016.

580 Paul, F., Piermattei, L., Treichler, D., Gilbert, L., Girod, L., Kääb, A., Libert, L., Nagler, T., Strozzi, T., and Wuite, J.: Three  
581 different glacier surges at a spot: what satellites observe and what not, *The Cryosphere*, 16, 2505-2526, 2022.

582 Paul, F., Rastner, P., Azzoni, R., Diolaiuti, G., Fugazza, D., Le Bris, R., Nemec, J., Rabatel, A., Ramusovic, M., and Schwaizer,  
583 G.: Glacier shrinkage in the Alps continues unabated as revealed by a new glacier inventory from Sentinel-2, *Earth Syst. Sci.*  
584 *Data*, 12, 1805–1821, 2020.

585 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., and Kienholz,  
586 C.: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *Journal of glaciology*, 60, 537-552, 2014.

587 Pralong, A. and Funk, M.: On the instability of avalanching glaciers, *Journal of Glaciology*, 52, 31-48, 2006.

588 Pralong, A., Birrer, C., Stahel, W. A., and Funk, M.: On the predictability of ice avalanches, *Nonlinear Processes in Geophysics*,  
589 12, 849-861, 2005.

590 Rabatel, A., Ducasse, E., Millan, R., and Mouginit, J.: Satellite-Derived Annual Glacier Surface Flow Velocity Products for  
591 the European Alps, 2015–2021, *Data*, 8, 66, 2023a.

592 Rabatel, A., Ducasse, E., Ramseyer, V., and Millan, R.: State and Fate of Glaciers in the Val Veny (Mont-Blanc Range, Italy):  
593 Contribution of Optical Satellite Products, *Journal of Alpine Research| Revue de géographie alpine*, 2023b.

594 Rankl, M., Kienholz, C., and Braun, M.: Glacier changes in the Karakoram region mapped by multimission satellite imagery,  
595 *The Cryosphere*, 8, 977-989, 2014.

596 RGI Consortium: Randolph Glacier Inventory - A Dataset of Global Glacier Outlines, Version 7 [Data Set], Boulder, Colorado  
597 USA. National Snow and Ice Data Center. <https://doi.org/10.5067/F6JMOVY5NAVZ>. (2023) Date Accessed 03-26-2024.

598 Samsonov, S., Tiampo, K., and Cassotto, R.: SAR-derived flow velocity and its link to glacier surface elevation change and  
599 mass balance, *Remote Sensing of Environment*, 258, 112343, 2021.

600 Scambos, T. A., Dutkiewicz, M. J., Wilson, J. C., and Bindschadler, R. A.: Application of image cross-correlation to the  
601 measurement of glacier velocity using satellite image data, *Remote sensing of environment*, 42, 177-186, 1992.

602 Schwalbe, E. and Maas, H.-G.: The determination of high-resolution spatio-temporal glacier motion fields from time-lapse  
603 sequences, *Earth Surface Dynamics*, 5, 861-879, 2017.

604 Span, N. and Kuhn, M.: Simulating annual glacier flow with a linear reservoir model, *Journal of Geophysical Research:*  
605 *Atmospheres*, 108, 2003.

606 Stocker-Waldhuber, M., Fischer, A., Helfricht, K., and Kuhn, M.: Long-term records of glacier surface velocities in the Ötztal  
607 Alps (Austria), *Earth system science data*, 11, 705-715, 2019.

608 Van Wyk De Vries, M. and Wickert, A. D.: Glacier Image Velocimetry: an open-source toolbox for easy and rapid calculation  
609 of high-resolution glacier velocity fields, *The Cryosphere*, 15, 2115-2132, 10.5194/tc-15-2115-2021, 2021.

610 Vincent, C. and Moreau, L.: Sliding velocity fluctuations and subglacial hydrology over the last two decades on Argentière  
611 glacier, Mont Blanc area, *Journal of Glaciology*, 62, 805-815, 2016.

612 Vincent, C., Le Meur, E., Six, D., Possenti, P., Lefebvre, E., and Funk, M.: Climate warming revealed by englacial temperatures  
613 at Col du Dôme (4250 m, Mont Blanc area), *Geophysical Research Letters*, 34, 2007.

614 Willis, I. C.: Intra-annual variations in glacier motion: a review, *Progress in Physical Geography*, 19, 61-106, 1995.

615 Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the EURO-  
616 CORDEX RCM ensemble, *The Cryosphere*, 13, 1125-1146, 2019.

617 Zemp, M., Hoelzle, M., and Haeberli, W.: Six decades of glacier mass-balance observations: a review of the worldwide  
618 monitoring network, *Annals of Glaciology*, 50, 101-111, 2009.

619 Zemp, M., Nussbaumer, S. U., Gärtner-Roer, I., Bannwart, J., Paul, F., and Hoelzle, M.: Global Glacier Change Bulletin Nr. 4  
620 (2018-2019), *WGMS*, 4, 2021.

621 Zemp, M., Huss, M., Eckert, N., Thibert, E., Paul, F., Nussbaumer, S. U., and Gärtner-Roer, I.: Brief communication: Ad hoc  
622 estimation of glacier contributions to sea-level rise from the latest glaciological observations, *The Cryosphere*, 14, 1043-1050,  
623 2020.

624