

Rate coefficients for the reactions of OH radical with C₃-C₁₁ alkanes determined by the relative rate technique

Yanyan Xin^{1,2#}, Chengtang Liu^{2,*,#}, Xiaoxiu Lun¹, Shuyang Xie², Junfeng Liu², Yujing Mu²

¹ Beijing Forestry University, Beijing, 100083, China

² Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, 100085, China.

[#] These authors contributed equally to this work.

*Correspondence to: Chengtang Liu (ctliu@rcees.ac.cn).

Abstract: Rate coefficients for the reactions of OH radicals with C₃-C₁₁ alkanes were determined using the multivariate relative rate technique. A total of 25 relative rate coefficients at room temperature and 24 Arrhenius expressions in the temperature range of 273-323 K were obtained. Notably, a new room temperature relative rate coefficient for 3-methylheptane that had not been previously reported was determined, and the obtained k_{OH} values-value (in units of $10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) was 7.71 ± 0.35 . Interestingly, whilst results for n-alkanes agreed well with available structure activity relationship (SAR) calculations of Atkinson and Kwok, Neeb, Wilson, Jenkin, and McGillen, the three cyclo-alkanes (cyclopentane, methylcyclopentane, cyclohexane) and one branched alkane (2,2,4-trimethylpentane) were found to be less reactive than predicted by SAR. Conversely, the SAR estimates for 2,3-dimethylbutane were approximately 25% lower than the experimental values, with the exception of those estimated by the Wilson group, ~~highlighting the limited understanding of the oxidative chemistry of these compounds.~~ highlighting that there may be additional factors that govern the reactivity of highly branched alkanes that are not captured by current SAR techniques. Arrhenius expressions (in units of $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) for the reactions of various branched alkanes with OH radical were determined for the first time: 2-methylheptane, $(1.37 \pm 0.48) \times 10^{-11} \exp [-(209 \pm 100)/T]$, and 3-methylheptane, $(3.54 \pm 0.45) \times 10^{-11} \exp [-(374 \pm 49)/T]$. The reactivity relation of saturated alkanes with OH radicals and chlorine atoms was obtained: $\log_{10}[k_{\text{(Cl+alkanes)}}] = 0.569 \times \log_{10}[k_{\text{(OH+alkanes)}}] - 3.111$ ($R^2 = 0.86$). In addition, the rate coefficients for the 24 ~~previous~~ previously studied OH + alkanes reactions were consistent with existing literature values, demonstrating the reliability and efficiency of this method for simultaneous investigation of gas-phase reaction kinetics.

30 **Keywords:** Relative rate coefficients; Atmospheric simulation chamber; Alkanes; OH radical; Arrhenius
31 expressions

32 1. Introduction

33 Volatile organic compounds (VOCs), a category of compounds found ubiquitously in the atmosphere,
34 primarily consist of alkanes, alkenes, aromatics and oxygenated volatile organic compounds (OVOCs)
35 (Lewis et al., 2000; Goldstein and Galbally, 2007; Anderson et al., 2004). Research has shown that
36 alkanes, including straight-chain, branched-chain, and cyclic alkanes within the C₃-C₁₁ range, often
37 constitute a significant portion to VOCs (Liang et al., 2023; Dunmore et al., 2015), and they could be
38 emitted into the atmospheric environment through natural and anthropogenic sources, e.g., C₅-alkanes
39 emitted from gasoline usage and C₆-alkanes and higher homologous VOCs emitted as a consequence of
40 their usage as solvents and from fuel evaporation. (Atkinson, 2000; Guenther, 2002; Atkinson and Arey,
41 2003). In the troposphere, the alkanes are extremely less reactive with NO₃ and ~~and not reacting~~ with ozone,
42 ~~and thus~~ they are degraded and removed from the atmosphere via gas-phase oxidation reactions with OH
43 radicals and chlorine atoms. (Atkinson and Arey, 2003; Shi et al., 2019; Finlayson-Pitts and Pitts, 1997;
44 Atkinson, 2000). These oxidation processes ~~will form a~~ ~~can lead to~~ photochemical smog in the presence
45 of NO_x and light, causing regional photochemical pollution (Fiore et al., 2005; Ling and Guo, 2014).
46 Additionally, degradation products produced by the oxidation of alkanes can form secondary organic
47 aerosol (SOA) through homogeneous nucleation or condensation onto existing primary particles (Sun et
48 al., 2016).

49 Numerous laboratories have conducted research on the kinetics of the reaction between alkanes and
50 OH radicals using ~~the absolute rate method and the relative rate method~~ both absolute and relative rate
51 methods. The absolute rate method (such as flash photolysis and emission flow ~~et al~~ etc.) involves
52 calculating the reaction kinetics parameter k_{OH} for organic compounds with OH radicals during the
53 experimental process by directly measuring changes in OH radical concentration or the concentration of
54 the target compound. Greiner measured the first kinetic data for the reaction of OH radicals with three
55 alkanes at 300 K and 28-149 Pa in the Ar system using the flash photolysis-resonance fluorescence
56 technique (Greiner, 1967). Over the next decade, Gorse et al., Overend et al. and Darnall et al. obtained
57 kinetic data for the reaction of OH radicals with selected alkanes in ~~the~~ carbon monoxide, He and N₂
58 ~~system~~ systems, respectively (Gorse and Volman, 1974; Overend et al., 1975; Darnall et al., 1978). Unlike

the absolute rate method, the relative rate method relies on the recommended rate coefficient for the reaction of a reference compound with OH radicals, with the reference reaction rate coefficient needing to be similar to that of the compound under study to enhance measurement sensitivity. By monitoring the simultaneous decay of the target and reference compounds in the presence of OH radicals due to competitive response mechanisms, the rate coefficient for the reaction of OH radicals with the target compound can be determined (Atkinson and Arey, 2003; Shaw et al., 2018). From 1980s to 2020s, dozens of papers for the rate coefficients of alkanes with OH radical measured by relative rate method have been published. For example, Shaw et al. and Phan and Li obtained rate coefficients of a series of alkanes ~~in the N₂/He system~~ (Phan and Li, 2017; Shaw et al., 2018; Shaw et al., 2020). Anderson et al. obtained the k_{OH} of C₂-C₈ several n-alkanes and cyclic alkanes by the relative technique ~~in the air system~~ at 296 ± 4 K (Anderson et al., 2004). However, the majority of experiments were conducted ~~solely~~ limited to on C₂-C₆ alkanes, ~~and~~ more complex and multifunctional alkanes are often poorly constrained or unmeasured.

Temperature has an important influence on the reaction rate coefficients of alkanes and OH radicals. The reaction rate coefficients of several n-alkanes with OH radicals measured by Greiner increased by about 70% in the range of 300-500 K (Greiner, 1970a). ~~Perry et al's research~~ Perry et al found that the rate coefficients of n-butane ~~multiplied~~ increased by 72% as the temperature rose from 297 K to 420 K (Perry et al., 1976). And the rate coefficients of 10 n-alkanes and cycloalkanes obtained by Donahue et al. also increased ~~in varying degrees at 300-390 K~~ varying the temperature in the range 300 - 390 K (Donahue et al., 1998). However, most reported experimental studies on the reactivity of OH radicals with a series of alkanes focus on temperatures ≥ 290 K (Greiner, 1970a; Perry et al., 1976; Finlayson-pitts et al., 1993; Donahue et al., 1998; Atkinson, 2003; Badra and Farooq, 2015), with relatively few studies at low temperatures (Demore and Bayes, 1999; Li et al., 2006; Wilson et al., 2006; Sprengnether et al., 2009; Crawford et al., 2011). In addition, ~~a further alkane had only two, or fewer, individual OH radical rate coefficient measurements available in the mentioned temperature range, e.g., 3-methylheptane there~~ is another alkane (e.g., 3-methylheptane) for which only two or fewer measurements of OH radical rate coefficients have been reported in the above temperature range, and it is unclear whether the rate coefficients for the reactions of OH radicals with alkanes differ in a mixed system containing oxygen compared to an inert gas system. Therefore, further investigations are required to explore the variations in the rate coefficients for different types of alkanes at various temperatures.

In this study, the rate coefficients for the reactions of 25 different C₃-C₁₁ alkanes with OH radicals

were determined using the multivariate relative rate method, including linear alkanes, cycloalkanes, and methyl-alkanes. To validate the rate coefficients for the reaction between alkanes and OH radicals, multiple comparisons were made with previous literature and structure–activity relationship (SAR) estimated values. Additionally, the rate coefficients of certain straight-chain, branched-chain, and methyl-cycloalkanes were measured at 273–323 K.

2. Methods

2.1 Experiment

2.1.1 Atmospheric simulation chamber




Figure 1. A schematic of the experimental device (MFC: Mass Flow Controller; UV: ultraviolet lamp)

As shown in Fig. 1, the chamber experiments were performed at atmospheric pressure in a climate-controlled box with a temperature range of 263–333 K (accuracy of ± 0.1 K). A 300 L Teflon airbag was suspended in the climate-controlled box to serve as the reaction system. The box was equipped with two Teflon-coated fans for rapid chemical mixing and a 254 nm ultraviolet lamp for photolysis of hydrogen peroxide (H_2O_2) to produce OH radicals. The inner walls of climate-controlled box were constructed with reflective steel plates to enhance ultraviolet light utilization. Bath gas (N_2 or O_2) and NMHCs were introduced into the Teflon bag through mass flow controllers with flow rate of 25 L min^{-1} and 100 mL min^{-1} , respectively, while excess H_2O_2 with respect to VOCs was injected through a three-way valve using a micro syringe. Initial conditions of the different species introduced into the reactor for each experiment are outlined in Table S1 in the Supplementary Material. By varying the presence of H_2O_2 , turning on/off the light, a series of observations were generated ~~such as $\text{N}_2 + \text{NMHCs} + \text{dark reaction}$, $\text{N}_2 + \text{NMHCs} + \text{h}\nu$ (254 nm), and $\text{N}_2 + \text{NMHCs} + \text{H}_2\text{O}_2 + \text{dark reaction}$.~~ Initial conditions of the different species

introduced into the reactor for each experiment are outlined in Table S1 in the Supplementary Material. By varying the presence of H₂O₂, turning on/off the light, a series of observations were generated such as N₂ + NMHCs + dark reaction, N₂ + NMHCs + hν (254 nm), and N₂ + NMHCs + H₂O₂ + dark reaction. to ensure no wall-loss of NMHCs under dark conditions and their stability when exposed to 254 nm light.

2.1.2 Gas sampling and analysis

NMHCs Analyzer (GC-FID) with a time resolution of 1 hour independently developed by the Research Center for Eco-Environmental Sciences (RCEES) was used to analyze 25 C₃-C₁₁ alkanes. The sample gas was enriched by a 60-80 mesh Carbopack B adsorption tube under the condition of 183.15 K, and then the adsorption tube was rapidly heated to 453.15 K. The 25 alkanes were detected by FID at 523.15 K after programmed heating at 253.15 K, 303.15 K and 433.15 K in 30 min (Liu et al., 2016).

Figure S1(a) reveals that the mixed gas diluted with N₂ underwent a 14-hour reaction in a Teflon reactor without light. The k_d values ranged from 1.3 to 4.8 (the units are $\times 10^{-4}$ ppbv/h), implying negligible influence from factors such as alkane loss from reactor walls, self-consumption, or airbag leakage. Figure S1(b) illustrates that the peak height variation for 25 alkanes + 50 μ l of H₂O₂ within 15 hours was less than 3%, indicating the insignificance of dark reactions between H₂O₂ and alkanes. When the same concentration mixed gas was irradiated for 7 hours without H₂O₂, alkane concentration changes were depicted in Fig. S2. The results indicated that minimal impact from alkane photolysis on OH radical reaction rate coefficient determination.



To obtain the reaction rate coefficients of alkanes with OH radicals in 1-2 hour, the alkanes mixture exiting the reactor was collected in more than ten polyvinyl fluoride (PVF) sampling bag (1.0 L) using a transparent vacuum sampling device for GC-FID. Prior to use, the empty sampling bag was flushed with high-purity nitrogen 3 times and placed within the vacuum sampler - a system utilizing an oil-free diaphragm air pump to create a vacuum. The initial concentrations of alkanes sample were collected before the lamp on, and the following sampling process occurred every 10 minutes. Collected samples were subsequently analyzed using a self-developed automated injection system for PVF bag.

2.1.3 Relative rate technique

The rate coefficients were measured by the relative rate method (Atkinson, 1986). The basic principle is that the rate coefficient for the reaction of the reactant used as a reference with OH radicals needs to be

the recommended rate coefficients values, rate constant for the reaction of OH radicals with a reference compound with OH needs to be well established; then, the rate coefficient for the target compound can be determined by monitoring the simultaneous decay of the target and reference compounds in the presence of OH radicals due to the competitive response mechanism. Additionally, an important criterion for the selection of reference compounds, that is, the reference rate coefficient needs to be similar to the one under study in order to improve sensitivity. ~~To ensure that the reactants only react with OH radicals, the OH radicals need to be in excess in the experiment.~~ The research method of this work is based on the multivariate relative rate method published by Shaw et al. (Shaw et al., 2018), taking the mixed system as the research object, broadening the range of compounds that can be examined.

Taking R (reference compounds) and X (target compounds) as examples, the reaction of OH radicals can be described as follows:



$$-\frac{d[R]}{dt} = k_R [OH][R] \quad (R3)$$

$$-\frac{d[X]}{dt} = k_X [OH][X] \quad (R4)$$

$$\ln \left(\frac{[R]_0}{[R]_t} \right) = k_R \cdot \int [OH] dt \quad (R5)$$

$$\ln \left(\frac{[X]_0}{[X]_t} \right) = k_X \cdot \int [OH] dt \quad (R6)$$

$$\ln \left(\frac{[X]_0}{[X]_t} \right) = \frac{k_X}{k_R} \cdot \ln \left(\frac{[R]_0}{[R]_t} \right) \quad (R7)$$

Where $[R]_0$ and $[X]_0$ are the concentrations of reference compounds and target compounds before turning on the light; $[R]_t$ and $[X]_t$ are the corresponding concentrations after turning on the light for time t. k_R and k_X refer to the second-order rate coefficients for the reaction of the reference compounds and target compounds with OH radicals.

2.1.4 Choice of reference k values

It is critical to choose appropriate reference compounds in a kinetics study using the relative rate technique. Some reported values of the rate coefficients for reactions of C_3 - C_{11} alkanes with OH radicals have been measured by different methods in different laboratories, and these measurement results may be

quite different. When these rate coefficients are measured by the relative rate technique, choosing different reference values will lead to a change of the final experimental target rate coefficients. In this work, ~~selecting 3 different commonly used reference compounds (n-Hexane, Cyclohexane, n-Octane)~~ three different commonly used reference compounds (n-hexane, cyclohexane and n-octane) were used to determine the rate coefficients for each reaction at room temperature to check the consistency of kinetic results. The selection of k values for reference compounds and the literature data assessment and comparison gives priority to the available expert-evaluated rate coefficients wherever possible. Here we used the recommended ~~expert-evaluated data~~ evaluated data of database for Version 2.1.0 of McGillen et al. (Database for the Kinetics of the Gas-Phase Atmospheric Reactions of Organic Compounds – Eurochamp Data Center), which is relatively comprehensive and provides rigorously evaluated rate coefficients for many species. Among them, at 298 ± 1 K, the k values (in units of $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) of the three reference compounds selected respectively are ~~expert-evaluated~~ evaluated rate ~~constants~~ coefficients: $k_{\text{OH}+\text{n-Hexane}}=4.97 \times 10^{-12}$, $k_{\text{OH}+\text{Cyclohexane}}=6.69 \times 10^{-12}$, $k_{\text{OH}+\text{n-Octane}}=8.48 \times 10^{-12}$, which is fitted or manually entered data from multiple sources. ~~However, the value of the reference compound at different temperatures (273–323 K) is different than the room temperature.~~ A detailed explanation at different temperatures is ~~reflected~~ presented in Sec. 3.3.

2.1.5 Materials

The air bath gas was obtained by a mix of nitrogen (200 L) and oxygen (50L). H_2O_2 (30%) was provided by Sinopharm Chemical Reagent Co., Ltd. The standard gas (PAMs) is a mixed standard sample of 57 kinds of NMHCs produced by Linde Spectra Environmental Gases (Alpha, NJ). Sampling bag (PVF, 1 L) was provided by Dalian Delin Gas Packing Co., Ltd. The pump is the NMP830 KNDC model produced by KNF, Germany, with a maximum air sampling rate of 23 L/min. The climate-controlled box (ZRG-1000D-C0203) is provided by Shanghai Proline Electronic Technology Co., Ltd.

2.2 Estimation of the rate ~~constant~~ coefficient at 298 K (SAR)

In the past few decades, researchers have been devoted to finding a reasonable theoretical estimation method for the kinetic rate coefficients (Cohen, 1991). Structure-Activity Relationship (SAR) established and developed by Kwok and Atkinson et al. (Kwok and Atkinson, 1995), is the most widely used estimation method of rate coefficients. Based on the relationship between the structure and the reaction

activity of the compounds, this method assumes that the hydrogen extraction reaction mainly occurs in the saturated compounds and the addition reaction mainly occurs in the unsaturated compounds, which is used to estimate the gaseous rate coefficients for the reactions of most VOCs with OH radicals. An advantage of the rate coefficient estimation is that it gives a measure of the rates of attack at different sites in the molecule, which is then useful in predicting the overall temperature dependence. The rate coefficient estimated by SAR method is in good agreement with the experimental data. In this relationship, the calculation of the rate coefficient of the hydrogen atom on the C-H bond is based on the evaluation of the rate coefficient of the -CH₃, -CH₂-, >CH- group. The relationship between the group structure and the rate coefficient is as follows:

$$\begin{aligned}
 k(\text{CH}_3\text{-X}) &= k_{\text{prim}}^0 F(\text{X}) \\
 k(\text{X-CH}_2\text{-Y}) &= k_{\text{sec}}^0 F(\text{X}) F(\text{Y}) \\
 k(\text{X-CH}(\text{Y})\text{Z}) &= k_{\text{tert}}^0 F(\text{X}) F(\text{Y}) F(\text{Z}) \\
 k_{\text{tot}} &= \sum [k(\text{CH}_3\text{-X}) + k(\text{X-CH}_2\text{-Y}) + k(\text{X-CH}(\text{Y})\text{Z})]
 \end{aligned}$$

Where, k_{tot} represents the rate coefficient of each target compound. k_{prim}^0 , k_{sec}^0 , k_{tert}^0 represent the rate-coefficients of each -CH₃, -CH₂- and >CH-. For standard substituent groups such as -CH₃, $F(-\text{CH}_3)=1.00$, X, Y and Z represent substituent groups, $F(\text{X})$, $F(\text{Y})$ and $F(\text{Z})$ refer to the activity coefficient of substituents (X, Y, Z) at different positions on carbon groups. At room temperature, $F(-\text{CH}_2\text{-})=1.23$, $F(>\text{CH-})=1.23$. Based on an extensive review of kinetic literature values for linear alkanes at room temperature, Atkinson and Kwok et al derived the values of k_{prim}^0 , k_{sec}^0 , k_{tert}^0 at room temperature, $k_{\text{prim}}^0=0.136 \times 10^{-12}$, $k_{\text{sec}}^0=0.934 \times 10^{-12}$, $k_{\text{tert}}^0=1.94 \times 10^{-12}$, the unit is cm³ molecule⁻¹ s⁻¹. After that, many researchers continued to update and modify some parameters based on the method of Atkinson and Kwok (Kwok and Atkinson, 1995), and obtained the new base rate coefficients for different positional groups, and also developed independent methods for rate coefficient estimation. Some examples include: ~~some examples include:~~ Neeb, Wilson et al., Jenkin et al., and McGillen et al. (Neeb, 2000; Wilson et al., 2006; Jenkin et al., 2018; McGillen et al., 2020).

3. Result and Discussion

3.1 Results from relative rate experiments at 298 K

The rate coefficients for the reactions involving OH radical with C₃-C₁₁ alkanes in the mixed system were determined at 298±1 K. The concentration curves of target alkanes and the reference compound (n-Hexane) were plotted in Fig. 2. As shown in Fig. 2, the decay of both target and reference compounds correlated well with eq. (7), the intercepts of the linear fits were close to 0 and high correlation coefficients (R²) were observed for most alkanes, exceeding 0.99. Table 1 and Table S2 listed the obtained k_{OH} for C₃-C₁₁ alkanes under three bath gases using the related reference compounds. The error bars (1σ) in Table 1 accounted for reference rate coefficient uncertainty, and experimental parameter uncertainties (pressure, temperature, flow rate, reactant concentration). The results indicated strong agreement (within <15%) between rate coefficients for 25 C₃-C₁₁ straight-chain, branched-chain, and cycloalkanes, using different reference compounds. For example, the k_{OH} obtained for propane with n-hexane, cyclohexane and n-octane as the reference compound were $(1.38\pm0.01)\times10^{-12}$, $(1.25\pm0.03)\times10^{-12}$ and $(1.34\pm0.04)\times10^{-12}$ (the units are cm³ molecule⁻¹ s⁻¹), respectively (within 10%). This suggests that reference compound variation minimally affects results, indicating reliable experimental methods and data. Notably, the rate coefficient for 3-Methylheptane's reaction with OH radicals at room temperature was determined for the first time. As shown in Fig. 3, for the different bath gases, the obtained k_{OH} for C₃-C₁₁ alkanes showed high agreement, errors (dY) between 0.12 and 0.41, the units are 10⁻¹²cm³ molecule⁻¹ s⁻¹. Meanwhile, it can also be observed from the figure that most of the rate coefficients obtained are very similar to the expert-evaluated values of the database by the McGillen et al. However, 2,4-Dimethylpentane is an exception, the k_{OH} value obtained in this study is about 20% lower than the recommended value, but it is similar to expert-evaluated value by Atkinson and Arey (Atkinson and Arey, 2003). Additionally, it can be clearly seen in the figure that the reactivity of linear alkanes (R₁CH₂R₂) with OH radicals increasing as the number of carbon atoms in the hydrocarbon molecules increases, indicating that the increase of R-terminal alkyl chain length will provide additional hydrogen extraction sites. For each additional CH₂ group from C₃-C₁₁, the reaction rate coefficient increases about 0.95-1.81 (the unit is 10⁻¹² cm³ molecule⁻¹ s⁻¹), reflecting the fact that the main way is to extract the H atom from the secondary C-H bond. For branching alkanes, for example, 2,2-Dimethylbutane and 2,3-Dimethylbutane, it is obvious that the addition of CH group increases the reaction rate coefficients with OH radical to a great extent. For cyclic alkanes, such

as cyclopentane, methylcyclopentane, cyclohexane and methylcyclohexane, it can also be seen that the reactivity increases with the increase of cycle size. By comparing the reaction rate coefficients of cyclopentane and cyclohexane (methylcyclopentane and methylcyclohexane), it is found that for cyclic alkanes, each CH₂ group reaction rate increases by about $2.37 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. It can be seen from the reaction rate coefficients of cyclopentane and methylcyclopentane (cyclohexane and methylcyclohexane) that the reaction rate coefficient increases about $2.06 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ for cycloalkanes with each increase addition of methyl.




Figure 2. Typical kinetic data as acquired with the multivariate relative rate technique at 298 K and a fixed reaction time of 70 min for the reaction of C₃-C₁₁ alkanes with the OH radical using n-hexane as reference compound in different bath gases (N₂, Air, O₂). The numbers in parentheses correspond to each substance, followed by the correlation coefficient R². The following data have been displaced for reasons of clarity: (N₂): (1) Methylcyclopentane, (2) Cyclohexane, (3) Cyclopentane, (4) 2-Methylpentane, (5) 2,3-Dimethylbutane, (6) 2,4-Dimethylpentane, (7) Isopentane, (8) 1-pentane, (9) 3-Methylpentane, (10) Isobutane, (11) n-Butane, (12) 2,2-Dimethylbutane, (13) Propane (14) n-Undecane, (15) n-Decane, (16) Nonane, (17) Methylcyclohexane, (18) n-Octane, (19) 3-Methylheptane, (20) 2-Methylheptane, (21) 2,3,4-Trimethylpentane, (22) 1-Heptane, (23) 2-Methylhexane, (24) 3-Methylhexane, (25) 2,2,4-Trimethylpentane vertically displaced by 2.8, 2.6, 2.4, 2.1, 1.8, 1.6, 1.4, 1.1, 0.9, 0.7, 0.5, 0.3, 0.1, 1.2, 1.1, 1, 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.25, 0.1 units, respectively; (Air) Each alkane (in the above order)




Figure 3. Comparison of rate coefficients of C₃-C₁₁ alkanes in different bath gases (N₂, Air, O₂) with expert-evaluated data evaluated data at 298±1 K. The error bar was taken as 1σ.

The obtained k_{OH} values for C₃-C₁₁ alkanes in this work were compared with literature-reported values (Table 1). For several n-alkanes, the average rate coefficient obtained in the air gas are consistent with literature values. For example, the result of n-butane (2.63 ± 0.23), (applicable to all units involved in this paragraph all units in this paragraph are $10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) is highly consistent with the value (2.56 ± 0.25) obtained by Greiner (Greiner, 1970a) and the value (2.72 ± 0.27) obtained by Perry et al. (Perry et al., 1976), with a consistency of 3% or better. Although slightly higher by 7% compared to Talukdar et al. (Talukdar et al., 1994) using absolute techniques (2.46 ± 0.15), when considering the errors, they still exhibit consistency within a certain range taking into account the experimental uncertainties. Compared to the value obtained by DeMore et al. (Demore and Bayes, 1999) using the relative rate method (2.36 ± 0.25) and the expert-evaluated data evaluated data (2.36) of McGillen et al.'s database, these values are higher by 11%.

285 **n-pentane (n-Heptane).** As in the n-butane case, the derived rate coefficients for n-pentane and n-heptane
286 are in excellent agreement (4% or better at 298 K) with previous studies (Donahue et al., 1998; Atkinson,
287 2003; Atkinson and Arey, 2003; Wilson et al., 2006; Crawford et al., 2011; Calvert et al., 2015; Morin et
288 al., 2015).

289 **n-Octane (Nonane).** The reaction rate coefficients of n-Octane and OH radicals are in extremely good
290 agreement with the values reported in the literature (within 5%) (Greiner, 1970a). ~~Same~~ The same applies
291 for Nonane, consistency with previous studies is less than 8% (Greiner, 1970a; Atkinson et al., 1982;
292 Ferrari et al., 1996; Atkinson and Arey, 2003; Li et al., 2006).

293 **n-Decane.** The obtained average k_{OH} for n-decane in the air system was (1.18 ± 0.02) , the unit is 10^{-11} cm^3
294 $\text{molecule}^{-1} \text{ s}^{-1}$. When considering experimental error, these results are consistent with the relative value
295 (1.29 ± 0.10) obtained by Li et al. (Li et al., 2006) and the reviewed value (1.10) of Atkinson and Arey
296 (Atkinson and Arey, 2003), with about a consistency of 6%-9%.

297 **n-Undecane.** The obtained average k_{OH} for n-decane in the air system was (1.33 ± 0.16) , the unit is 10^{-11}
298 $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. It is about 8% higher than the previous research (Atkinson and Arey, 2003;
299 Sivaramakrishnan and Michael, 2009; Calvert et al., 2015).

300 For the cycloalkanes, like cyclopentane, the average rate ~~coefficients are 4.88 ± 0.19~~ , coefficient is
301 4.82 ± 0.27 , ~~4.59 ± 0.14 , respectively~~, the unit is $10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. The results are in excellent
302 agreement (8% or better) with the reviewed value (4.97) of Atkinson and Arey (Atkinson and Arey, 2003)
303 and the relative values $(4.83, 4.84)$ of DeMore et al. (Demore and Bayes, 1999) and Singh et al. (Singh et
304 al., 2013) and the absolute value (5.02) of Droege et al. (Droege and Tully, 1987). And the obtained k_{OH}
305 values for cyclohexane are highly consistent (3% or better) with the absolute values $(7.14 \times 10^{-12}, 7.19 \times 10^{-12})$
306 obtained by Droege and Tully and Sprengnether et al. (Droege and Tully, 1987; Sprengnether et al.,
307 2009). However, this result is slightly higher than the relative value by about 5%-16%. ~~Like~~ For example,
308 the relative values measured by DeMore and Bayes (Demore and Bayes, 1999) or Wilson et al. (Wilson
309 et al., 2006) were 6.70×10^{-12} and 6.38×10^{-12} , respectively. It worth noting that the k_{OH} value for
310 methylcyclopentane in this work is highly consistent (within 3% to 5%) with the absolute data reported
311 by Sprengnether et al. (Sprengnether et al., 2009). However, it is lower by approximately 15% to 18%
312 compared to the relative data obtained by Anderson et al. (Andersen et al., 2003). The k_{OH} values for
313 methylcyclohexane are in excellent agreement (3% or better) with other values reported by Atkinson and
314 Arey (Atkinson and Arey, 2003) and Calvert et al. (Calvert et al., 2015).

Furthermore, for several less studied branched alkanes, such as 2-Methylhexane, 3-Methylhexane, and 2-Methylheptane, there is only one study reported so far. Sprengnether et al. (Sprengnether et al., 2009) conducted a study on 2-Methylhexane and 3-Methylhexane and obtained k_{OH} values at room temperature for the first time, which were 6.69×10^{-12} and 6.30×10^{-12} (the unit is $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$), respectively. The rate coefficients of 2-Methylhexane and 3-Methylhexane obtained in this work are $(6.80 \pm 0.13) \times 10^{-12}$ and $(6.29 \pm 0.11) \times 10^{-12}$, respectively, which are consistent with the values obtained by Sprengnether et al. (Sprengnether et al., 2009). However, the data for 2-Methylheptane in this work is lower by about 17% compared to the value reported by Shaw et al. (Shaw et al., 2018).

Table 1. Comparison of Experimental in this work with the reported in the literature at $298 \pm 1 \text{ K}$.

Alkanes	Reference	This work			Literature measurements
		$k_{OH}/k_{\text{reference}} \pm \theta$	$k_{OH} \pm \theta$ ($\times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	$k_{OH-\text{av}}^a \pm \theta_{\text{av}}$ ($\times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	k_{OH} ($\times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)
Propane	n-Hexane	0.190 \pm 0.033	(9.43 \pm 1.66)		1.11 ^{bcd} 1.09 ^e
	Cyclohexane	0.153 \pm 0.028	(1.03 \pm 0.18)	(1.01 \pm 0.26)	1.91 ^f
	n-Octane	0.136 \pm 0.031	(1.16 \pm 0.26)		(1.15 \pm 0.15) ^g
Isobutane	n-Hexane	0.444 \pm 0.012	(2.21 \pm 0.06)		2.12 ^h 2.22 ⁱ
	Cyclohexane	0.315 \pm 0.008	(2.08 \pm 0.02)	(2.19 \pm 0.13)	(2.34 \pm 0.33) ^j
	n-Octane	0.264 \pm 0.005	(2.24 \pm 0.04)		
n-Butane	n-Hexane	0.516 \pm 0.025	(2.56 \pm 0.12)		(2.36 \pm 0.25) ^b (2.72 \pm 0.27) ^k
	Cyclohexane	0.398 \pm 0.017	(2.66 \pm 0.12)	(2.63 \pm 0.23)	(2.56 \pm 0.25) ^m
	n-Octane	0.345 \pm 0.042	(2.93 \pm 0.36)		(2.46 \pm 0.15) ^d
Isopentane	n-Hexane	0.684 \pm 0.033	(3.40 \pm 0.17)		3.60 ^e 3.65 ^h
	Cyclohexane	0.512 \pm 0.026	(3.43 \pm 0.18)	(3.49 \pm 0.25)	3.50 ^f
	n-Octane	0.442 \pm 0.025	(3.75 \pm 0.22)		
n-pentane	n-Hexane	0.709 \pm 0.042	(3.52 \pm 0.21)		3.80 ^e 3.98 ⁿ
	Cyclohexane	0.527 \pm 0.021	(3.53 \pm 0.14)	(3.59 \pm 0.25)	4.03 ^o
	n-Octane	0.454 \pm 0.029	(3.85 \pm 0.24)		(3.97 \pm 0.20) ^p (4.20 \pm 0.15) ^g

					4.97 ^e
					4.83 ^b
					5.02 ^q
Cyclopentane	n-Hexane	0.951±0.033	(4.72±0.17)		(4.90±0.20) ^p
	Cyclohexane	0.711±0.043	(4.76±0.29)	(4.82±0.27)	
	n-Octane	0.600±0.029	(5.09±0.24)		4.84 ^{b r}
					(2.23±0.15) ^p
2,2-Dimethylbutane	n-Hexane	0.409±0.019	(2.03±0.09)		2.15 ^s
	Cyclohexane	0.301±0.030	(2.02±0.20)	(2.05±0.23)	2.32 ^o
	n-Octane	0.264±0.031	(2.24±0.26)		
					5.78 ^e
2,3-Dimethylbutane	n-Hexane	1.095±0.061	(5.44±0.31)		(6.14±0.25) ^p
	Cyclohexane	0.809±0.039	(5.42±0.26)	(5.62±0.31)	6.03 ^h
	n-Octane	0.728±0.050	(6.05±0.29)		
					5.2 ^e
2-Methylpentane	n-Hexane	0.972±0.022	(4.83±0.11)		(5.25±0.25) ^p
	Cyclohexane	0.722±0.054	(4.83±0.36)	(4.86±0.26)	5.00 ^f
	n-Octane	0.625±0.045	(5.30±0.38)		4.75 ^s
					5.20 ^e
3-Methylpentane	n-Hexane	1.014±0.030	(5.04±0.15)		(5.54±0.25) ^p
	Cyclohexane	0.777±0.059	(5.20±0.40)	(5.08±0.31)	4.93 ^s
	n-Octane	0.669±0.082	(5.67±0.70)		
					(7.65±0.10) ^u
methylcyclopentane	n-Hexane	1.432±0.053	(7.12±0.27)		(8.60±0.30) ^p
	Cyclohexane	1.007±0.023	(6.73±0.15)	(7.31±0.29)	(8.60±2.20) ^t
	n-Octane	0.849±0.017	(7.00±0.24)		
					4.80 ^e
2,4-Dimethylpentane	n-Hexane	0.962±0.012	(4.78±0.06)		5.51 ^s
	Cyclohexane	0.721±0.046	(4.83±0.31)	(4.80±0.20)	(5.76±0.40) ^p
	n-Octane	0.596±0.026	(5.05±0.22)		
					6.97 ^e
					7.14 ^q
Cyclohexane	n-Hexane	1.372±0.054	(6.82±0.27)		6.38 ^h
	Cyclohexane	--	--	(7.20±0.33)	6.70 ^b
	n-Octane	0.872±0.022	(7.39±0.19)		(7.19±0.10) ^u
					(6.85±0.20) ^p
2-Methylhexane	n-Hexane	1.369±0.004	(6.80±0.02)		
	Cyclohexane	0.993±0.022	(6.64±0.15)	(6.80±0.13)	(6.69±0.10) ^u
	n-Octane	0.800±0.031	(6.78±0.26)		
3-Methylhexane	n-Hexane	1.266±0.003	(6.29±0.02)		
	Cyclohexane	0.984±0.046	(6.58±0.31)	(6.29±0.11)	(6.30±0.10) ^u
	n-Octane	0.807±0.122	(6.73±0.74)		

					3.34 ^e
					3.64 ^s
2,2,4-Trimethylpentane	n-Hexane	0.702±0.033	(3.49±0.16)		
	Cyclohexane	0.557±0.032	(3.72±0.21)	(3.58±0.28)	(3.34±0.25) ^p
	n-Octane	0.435±0.065	(3.69±0.55)		(3.71±0.10) ^v
					6.76 ^e
n-Heptane	n-Hexane	1.280±0.066	(6.36±0.33)		6.68 ^y
	Cyclohexane	0.961±0.020	(6.43±0.26)	(6.78±0.36)	6.80 ^h
	n-Octane	0.828±0.029	(7.03±0.25)		(6.70±0.15) ^g
					9.60 ^e
					(9.64±0.30) ^p
Methylcyclohexane	n-Hexane	1.906±0.098	(9.48±0.49)		(11.8±1.00) ^F
	Cyclohexane	1.349±0.012	(9.02±0.08)	(9.25±0.22)	(9.50±0.14) ^D
	n-Octane	1.160±0.016	(9.83±0.14)		(9.29±0.10) ^u
					6.60 ^e
2,3,4-Trimethylpentane	n-Hexane	1.355±0.050	(6.73±0.25)		6.50 ^h
	Cyclohexane	1.008±0.039	(6.74±0.26)	(6.87±0.30)	(6.60±0.26) ^p
	n-Octane	0.861±0.039	(7.30±0.33)		
	n-Hexane	1.532±0.062	(7.62±0.31)		
2-Methylheptane	Cyclohexane	1.061±0.029	(7.09±0.19)	(7.49±0.27)	9.10 ^L
	n-Octane	0.931±0.025	(7.89±0.21)		
	n-Hexane	1.532±0.070	(7.62±0.35)		
3-Methylheptane	Cyclohexane	1.055±0.072	(7.06±0.48)	(7.71±0.35)	--
	n-Octane	0.948±0.036	(8.04±0.31)		
					8.11 ^e
n-Octane	n-Hexane	1.680±0.038	(8.35±0.19)		8.42 ^m
	Cyclohexane	1.157±0.027	(7.74±0.18)	(8.03±0.32)	(8.48±0.10) ^z
	n-Octane	--	--		
					9.70 ^e
Nonane	n-Hexane	2.166±0.079	(10.76±0.39)		10.20 ^A
	Cyclohexane	1.449±0.028	(9.69±0.19)	(10.50±0.26)	10.70 ^w
	n-Octane	1.287±0.017	(10.92±0.14)		(11.30±1.10) ^z
					11.00 ^e
n-Decane	n-Hexane	2.371±0.073	(11.78±0.36)		(12.9±1.00) ^z
	Cyclohexane	1.668±0.022	(11.16±0.15)	(11.81±0.18)	
	n-Octane	1.401±0.006	(11.88±0.05)		
	n-Hexane	2.371±0.073	(11.78±0.36)		12.30 ^e
n-Undecane	Cyclohexane	1.668±0.022	(11.16±0.15)	(12.78±0.53)	12.50 ^B
	n-Octane	1.588±0.056	(13.50±0.60)		(11.90±2.00) ^p

326

327 a: Weighted average $k_{av} = (w_{ref1} k_{ref1} + w_{ref2} k_{ref2} + \dots) / (w_{ref1} + w_{ref2} + \dots)$, where $w_{ref1} = 1/\theta_{ref1}^2$, etc. The error,

328 $\theta_{ref} = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n}}$, $\theta_{av} = (1/\theta_{ref1} + 1/\theta_{ref2} + \dots)^{-0.5}$.

329 b: (Demore and Bayes, 1999); c: (Mellouki et al., 1994); d: (Talukdar et al., 1994); e: (Atkinson and Arey,

2003); f: (Cox et al., 1980); g: (Morin et al., 2015); h: (Wilson et al., 2006); i: (Tully et al., 1986); j: (Edney et al., 1986); k: (Perry et al., 1976); m: (Greiner, 1970a); n: (Donahue et al., 1998); o: (Harris and Kerr, 1988); p: (Calvert et al., 2015); q: (Droege and Tully, 1987); r: (Singh et al., 2013); s: (Badra and Farooq, 2015); u: (Sprengnether et al., 2009); t: (Anderson et al., 2004); v: (Greiner, 1970b); y: (Crawford et al., 2011); z: (Li et al., 2006); L: (Shaw et al., 2020); w: (Atkinson et al., 1982); A: (Ferrari et al., 1996); B: (Sivaramakrishnan and Michael, 2009); D: (Bejan et al., 2018); F: (Ballesteros et al., 2015).

3.2 Comparisons to structure-activity relationships

To assess the accuracy of the estimation technique, multiple comparisons were made between the obtained reaction rate coefficients and the SAR values of different experimental groups (Figure 4). As shown in Figure 4, most n-alkanes are fall into the shaded region, indicating a high level of agreement for k_{OH} rate coefficients of most n-alkanes (experimental values) with the SAR values, particularly for C₃-C₁₁ n-alkanes (about within 10%). Although the measured values of n-butane and n-pentane were lower than the estimated values of Neeb (Neeb, 2000), the similar trend was observed when comparing our experimental data with the SAR values of (Kwok and Atkinson, 1995), and (Jenkin et al., 2018) and (Wilson et al., 2006), (refer to Fig. 4 (a), Fig. 4 (c), and Fig. 4 (d)), suggesting a certain level of reliability in our results.

For branched alkanes, such as monomethyl branched alkanes (2-Methylpentane, 3-Methylpentane, 2-Methylhexane, 3-Methylhexane 2-Methylheptane and 3-Methylheptane), the obtained k_{OH} values all fall within the shadow area. The results indicated a relatively consistent alignment between our experimental data and the SAR estimated data within a certain margin of error, particularly for the SAR values of Neeb and Jenkin et al. (within 8%). Nevertheless, there seemed to be something different for polymethyl branched alkanes, like 2,3-Dimethylbutane, [where](#) the experimental data was about 25% higher than the estimated SAR values of Atkinson and Kwok et al. (1995) and Neeb (2000), especially 53% higher than that of Jenkin et al. (2018). It was also found that the k_{OH} of this compound (at 298 K) could not be accurately estimated by Wilson et al. (Wilson et al., 2006) due to unknown reasons. Furthermore, compared with the SAR values of Atkinson and Kwok et al., the obtained data of 2,2-Dimethylbutane and 2,4-Dimethylpentane were relatively consistent with that, while compared with the estimated data of Neeb, Jenkin et al. and Wilson et al., our results are higher or lower by about 18% and 22%. It is worth noting that the obtained k_{OH} value of 2,2,4-Trimethylpentane was about 23%, 16% and

17%, respectively, lower than the corresponding SAR values of Atkinson and Kwok et al., Neeb, and Jenkin et al. The results indicated that our understanding for the oxidation chemistry of these compounds is still limited, ~~still need a lot of experimental data for alkanes with this structure to confirm.~~ and that further data and analysis for alkanes with this structure are needed.

For cyclic alkanes, such as cyclopentane and cyclohexane, the obtained k_{OH} values in this study were approximately 32% and 15%, respectively, lower than the SAR values of Atkinson and Kwok et al., 1995; b. Neeb 2000; c. Jenkin et al. 2018. On the other hand, the obtained experimental values for methylcyclopentane and methylcyclohexane were similar to SAR values of Neeb and Wilson et al (within 5%) (Neeb, 2000; Wilson et al., 2006), However, compared with the SAR values of Atkinson and Kwok et al. and Jenkin et al., this result is about 15% and 8% lower. The result suggested that the reaction activity of these ~~cycle-chain~~ cyclic alkanes estimated with SAR methods (Kwok and Atkinson, 1995; Jenkin et al., 2018) might be overestimated to varying degrees.

In addition, there are a number of SAR methods that are quite different in their estimation from those of Atkinson, Wilson, et al. and Neeb, et al., for instance, the method of McGillen et al. ~~Figure S3~~ Figure 4. (e) shows a comparison of our measurements with the SAR estimates of McGillen et al. Similar to the results of Kwok and Atkinson, Neeb, and Jenkin et al., the obtained k_{OH} values of cyclopentane and 2,3-Dimethylbutane in this study exceed the shaded area. This further illustrates that there is still a large discrepancy between the experimental values and the SAR estimates for both substances. For cycloalkanes, the SAR estimates of McGillen et al. are still overestimated to varying degrees compared to our measurements, especially for cyclopentane, where the experimentally measured k_{OH} in this work is still about 34% lower than the SAR estimate. And the k_{OH} values for cyclohexane, methylcyclopentane and methylcyclohexane were also lower than the estimated values by about 18%, 12% and 5%, respectively. For the branched alkanes, again the k_{OH} of 2,3-Dimethylbutane is higher than the SAR estimate by about 32% or so. Similarly to the comparison with the Neeb, and Jenkin et al SAR estimates, the experimental measurements we obtained for 2,2,4-Trimethylpentane are also lower than the McGillen et al estimates by about 14%. By comparing the reaction rate coefficients of cyclopentane and cyclohexane, it is found that for cyclic alkanes of Kwok and Atkinson, Neeb, Jenkin et al., and McGillen et al, ~~the cycle size increases~~ the increase in cycle size increases k by about $1.41 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. However, for the SAR estimate of Wilson et al, ~~the cycle size increases by~~ the increase is about $1.12 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. It is worth considering that the presence of ring strain can influence the kinetics of

H-atom abstraction in cyclic alkanes, leading to an overestimation of reaction rate coefficients when using unadjusted F(-CH₂-). And hence, further data and analysis for F(-CH₂-) with these cyclic alkanes are needed.




Figure 4. Measured Alkanes + OH rate coefficients plotted against SAR-derived rate coefficients for all compounds (a. (Kwok and Atkinson, 1995); b. (Neeb, 2000); c. (Jenkin et al., 2018); d. (Wilson et al., 2006); e. (McGillen et al., 2020)). The shaded area demonstrates a 20 % uncertainty in the 1:1 black

gradient line. The alkanes represented by serial number can be identified as follows: (1) Propane; (2) Isobutane; (3) n-Butane; (4) Isopentane; (5) n-pentane; (6) Cyclopentane; (7) 2,2-Dimethylbutane; (8) 2,3-Dimethylbutane; (9) 2-Methylpentane; (10) 3-Methylpentane; (11) Methylcyclopentane; (12) 2,4-Dimethylpentane; (13) Cyclohexane; (14) 2-Methylhexane; (15) 3-Methylhexane; (16) 2,2,4-Trimethylpentane; (17) n-Heptane; (18) Methylcyclohexane; (19) 2,3,4-Trimethylpentane; (20) 2-Methylheptane; (21) 3-Methylheptane; (22) n-Octane; (23) n-nonane; (24) n-Decane; (25) n-Undecane.

3.3 Temperature dependence (273-323 K)

In order to investigate the relationship between temperature and reaction rate coefficient, experiments were carried out in this study in the ~~tropospheric temperature range~~ temperatures relevant to the troposphere (273-323 K), and the Arrhenius formulas was obtained for this temperature range. Also, our data were combined with the literature data (the expert-recommended data from database for Version 2.1.0 of McGillen et al.) to study the kinetic temperature dependence of several alkanes in a wide temperature range. And n-hexane (Arrhenius expression: $k(T)=(2.43\pm0.52)\times10^{-11} \exp [-(481.2\pm60)/T]$ at 240-340 K was used as the reference compound. Since the research results at room temperature show that different bath gases have little effect on the reaction rate coefficient, only the temperature dependence of the reaction rate coefficient under the air system is considered here. Measured values for 24 C₃-C₁₀ alkanes were provided at different temperatures (273-323 K) in Table S3. And the preexponential factor A and activation energy E_a/R obtained by linear regression and non-linear curve fits and along with the values of the literature were listed in Table 2. The value of preexponential factor A increases with the increase of the number of carbon atoms, which is consistent with the law of its reactivity. Additionally, Arrhenius plots were linearly fitted using this data along with literature data. The following is a detailed analysis for several components that are important or temperature dependence data has been less or ~~no~~ not studied, the Arrhenius plots are shown in Figure 4-5, other components are listed in the Supplement (Fig. S9-S21). In addition, typical RR plots for some alkanes at different temperatures are also listed in the Supplement (Fig. S3-S8).

Table 2. Summary of Arrhenius Expression of the Reaction of OH radical with C₃-C₁₁ alkanes in this work and other studies.

Alkanes	Temperature (K)	A-factor ^a ($\times 10^{-11}$)	E _a /R ^b (K)	Technique ^c	Literature measurements
Propane	273-323	2.38±0.90	952±110	RR/DP/GC-FID	this work
	296-908	2.71±0.17	988±31	AR/FP/LIF	(Bryukov et al., 2004)
	227-428	1.29	730	RR/DP/GC	(Demore and Bayes, 1999)
	233-376	1.01	660	AR/FP/LIF	(Talukdar et al., 1994)
	300-390	1.12	692	AR/EB/LIF	(Donahue et al., 1998)
n-Butane	273-323	3.78±0.66	867±52	RR/DP/GC-FID	this work
	235-361	1.68	584	RR/DP/GC	(Demore and Bayes, 1999)
	300-390	1.34	513	AR/EB/LIF	(Donahue et al., 1998)
	231-378	1.18	470	AR/ DF/LIF	(Talukdar et al., 1994)
	294-509	1.88±0.09	617±18	AR/ DF/LIF	(Droege and Tully, 1987)
	298-420	1.76	559	AR/ DF/RF	(Perry et al., 1976)
	298-416	0.629	126	AR-UV	(Gordon and Mulac, 1975)
n-pentane	273-323	0.90±0.05	310±17	RR/DP/GC-FID	this work
	233-364	1.94	494	RR/DP/GC	(Demore and Bayes, 1999)
	300-390	2.97	608	AR/EB/LIF	(Donahue et al., 1998)
n-pentane	224-372	2.45±0.21	516±25	AR/FP/LIF	(Talukdar et al., 1994)
	243-325	--	--	RR/DP/GC	(Harris and Kerr, 1988)
n-Heptane	273-323	3.96±0.37	544±28	RR/DP/GC-FID	this work
	*240-1364	4.48	116, n=1.72		
	*290-1090	1.73	406, n=2	Review	(Atkinson and Arey, 2003)
	241-406	3.38±0.17	497±16	RR/DF/MS	(Wilson et al., 2006)
	240-340	2.25±0.14	293±37	RR/DF/MS	(Crawford et al., 2011)
	*248-896	4.39	138, n=1.7	AR/DF/LIF	(Morin et al., 2015)
	298-500	0.986	600	Theory	(Cohen, 1991)
	*241-1287	6.03	32, n=1.4		(Sivaramakrishnan and Michael, 2009)
	838-1287	248±17	193	AR-UV	
	869-1364	243	180	AR-UV	(Pang et al., 2011)
	*240-1364	3.84	148, n=1.79	recommended	(Mcgillen et al., 2020)

n-Octane	273-323	4.22±0.49	497±34	RR/DP/GC-FID	this work
	*240-1080	3.60	251, n=1.78		
	240-340	2.27±0.21	296±27	RR/DF/MS	(Li et al., 2006)
	284-384	4.52±0.37	538±27	RR/DF/MS	(Wilson et al., 2006)
	*290-1080	2.42	361, n=2.00	Review	(Atkinson and Arey, 2003)
	296-497	2.57	332±65	AR/FP/KS	(Greiner, 1970a)
Nonane	*298-1000	0.986	600, n=2.2	Theory	(Cohen, 1991)
	273-323	5.29±0.63	520±35	RR/DP/GC-FID	this work
	240-340	4.35±0.49	411±32	RR/DF/MS	(Li et al., 2006)
n-Decane	273-323	5.78±0.49	499±25	RR/DP/GC-FID	this work
	240-340	2.26±0.28	160±36	RR/DF/MS	(Li et al., 2006)
Isobutane	273-323	2.29±0.74	739±94	RR/DP/GC-FID	this work
	300-390	0.626	321	AR/EB/LIF	(Donahue et al., 1998)
	213-372	0.572	293	AR/FP/LIF	(Talukdar et al., 1994)
	297-498	0.347	192	AR/FP/GC	(Greiner, 1970a)
	220-407	1.02±0.03	463±10	RR/DF/MS	(Wilson et al., 2006)
Isopentane	273-323	1.12±0.11	443±34	RR/DP/GC-FID	this work
	213-407	1.39±0.12	424±25		
	213-407	1.52	432	RR/DP/GC	(Wilson et al., 2006)
Cyclopentane	273-323	3.67±0.63	619±51	RR/DP/GC-FID	this work
	288-407	2.71	526	RR/DP/GC	(Wilson et al., 2006)
	240-340	2.43±0.50	481±58	RR/DF/MS	(Singh et al., 2013)
	273 - 423	2.57	498	RR/DP/GC	(Demore and Bayes, 1999)
	300-390	1.88	352	AR/EB/LIF	(Donahue et al., 1998)
	295-491	2.29±0.09	457±0.14	AR/FP/LIF	(Droege and Tully, 1987)
Cyclohexane	273-323	3.62±0.59	522±48	RR/DP/GC-FID	this work
	240-340	3.96±0.60	554±42	RR/DF/MS	(Singh et al., 2013)
	288-408	3.40	513	RR/DP/GC	(Wilson et al., 2006)
	273-323	1.65±0.19	262±33	RR/DP/GC-FID	this work
	*230-1344	1.59	439, n=2.15		

Methylcyclopentane	*230-370	--	--	AR/DF/LIF	(Sprengnether et al., 2009)
	*230-1344	1.67	454, n=2.15		McGillen, 2020
Methylcyclohexane	273-323	4.39±0.58	475±29	RR/DP/GC-FID	this work
	273-343	1.85±0.27	195±20	RR/DP/FTIR	(Bejan et al., 2018)
	230-379	1.46±0.07	125±14	AR/ DF/LIF	(Sprengnether et al., 2009)
2,2-Dimethylbutane	273-323	3.53±1.28	899±106	RR/DP/GC-FID	this work
	240-330	3.37	809	Review	(Atkinson and Arey, 2003)
	243-328	--	--	RR/DP/GC	(Harris and Kerr, 1988)
	254-1327	6.14±0.90	1023±76	AR/DF/LIF	(Badra and Farooq, 2015)
2,3-Dimethylbutane	273-323	1.15±0.09	219±24	RR/DP/GC-FID	this work
	*273-1366	1.29	437, n=2.09		
	*240-1220	1.47	407, n=2.00	Review	(Atkinson and Arey, 2003)
	300-498	2.24	321	AR/FP/GC	Greiner,1970
	*250-1366	1.3	427, n=2.08	AR/DF/LIF	(Badra and Farooq, 2015)
	*220-1292	1.6	364, n=1.96	Review	(Sivaramakrishnan and Michael, 2009)
2,4-Dimethylpentane	273-323	2.03±0.17	452±24	RR/DP/GC-FID	this work
	272-410	2.25	408	RR/DP/GC	(Wilson et al., 2006)
	896-1311	14.9±0.8	1533±55	AR/DF/LIF	(Badra and Farooq, 2015)
2-Methylpentane	273-323	2.30±0.29	479±38	RR/DP/GC-FID	This work
	283-387	2.07	413	RR/DP/GC	(Wilson et al., 2006)
3-Methylpentane	273-323	2.44±0.39	511±17	RR/DP/GC-FID	this work
	284-381	2.16	375	RR/DP/GC	(Wilson et al., 2006)
	297-1362	6.43±0.87	834±74	AR/DF/LIF	(Badra and Farooq, 2015)

2-Methylhexane	273-323	1.30±0.08	222±19	RR/DP/GC-FID	this work
	273-385	1.82±0.09	321±16		
	230 - 385	1.21±0.07	171±16	AR/ DF/LIF	(Sprengnether et al., 2009)
3-Methylhexane	273-323	2.53±1.45	575±161	RR/DP/GC-FID	this work
	230-379	1.42±1.52	628±85	AR/ DF/LIF	(Sprengnether et al., 2009)
2-Methylheptane	273-323	1.37±0.48	209±100	RR/DP/GC-FID	this work
3-Methylheptane	273-323	3.54±0.34	456±28	RR/DP/GC-FID	this work
2,2,4-Trimethylpentane	273-323	1.61±0.22	499±40	RR/DP/GC-FID	this work
	240-500	1.62	443	AR/ DF/LIF	(Atkinson, 1986)
	230-385	1.54	456	AR/ DF/LIF	(Atkinson, 2003)
2,3,4-Trimethylpentane	273-323	1.34±0.07	203±15	RR/DP/GC-FID	this work
	287-373	1.3	221	RR/DP/GC	(Wilson et al., 2006)

^{a, b}The error bar was taken as σ .

^cRR: relative rate; AR: absolute rate; DF: discharge flow; DP: direct photolysis; FP: flash photolysis; EB: electron beam; UV: Ultraviolet; GC: gas chromatography; FID: flame ionization detection; LIF: laser induced fluorescence; FTIR: fourier transform infrared spectrometer; MS: mass spectrometry; KS: kinetic-spectroscopy.

* The expression takes the form $k(T) = A \times \exp(E_a/RT) \times (T/300)^n$

A. OH+ n-Octane. Figure 5 (a) exhibits the Arrhenius plot for the reaction between n-Octane and OH radicals, covering a temperature range of 240 to 1080 K. Within the experimental temperature range (273-323 K), our data align well with previous studies. The derived Arrhenius expression is as follow: $k_{\text{n-Octane}}(T) = (4.22 \pm 0.49) \times 10^{-11} \exp [-(497 \pm 34)/T]$ (T=273-323 K). The result agree well with the Arrhenius expression of $(4.52 \pm 0.37) \times 10^{-11} \exp [-(538 \pm 27)/T] \text{ cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$ reported by Wilson et al. (Wilson et al., 2006) between 284 and 384 K, but contrast the expressions of $(2.27 \pm 0.21) \times 10^{-11} \exp [-(296 \pm 27)/T] \text{ cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$ reported by Li et al. between 240 and 340 K (Li

et al., 2006) and $(2.57) \times 10^{-11} \exp[-(332 \pm 65)/T] \text{ cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$ reported by Greiner (Greiner, 1970a) between 296 and 497 K. Fit our data to ~~expert-evaluated data~~ ~~evaluated data~~ (manually entered data from multiple sources), the derived Arrhenius expression in 240-1080 K is as follow:

$k_{\text{n-Octane}}(T) = 3.60 \times 10^{-12} \times \exp(251/T) \times (T/300)^{1.78} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. This result is ~~slightly~~ ~~reasonably~~ consistent with the recommended expression ($k_{\text{n-Octane}}(T) = 2.42 \times 10^{-12} \times \exp(361/T) \times (T/300)^{2.00}$ for and Arey (Atkinson and Arey, 2003). By comparison, our data are highly consistent with the data recommended by experts. The obtained Arrhenius expression more accurately represents the relationship between the reaction rate coefficient of octane and OH radicals and temperature in 273-323 K and a wide temperature range, which has certain reference significance. Further investigations are necessary to understand the discrepancies amongst these studies. Also, the experimental values of n-Octane obtained at different temperatures are in high agreement with the SAR estimates.

B. OH+ n-Heptane. The Arrhenius plot in Fig. 5 (b) displays the reaction between n-Heptane and OH radicals in the air systems, covering a temperature range of 240 to 896 K. As shown in the figure, within the experimental temperature range (273-323 K), our data are highly similar to previous studies. The Arrhenius expression obtained is $k_{\text{n-Heptane}}(T) = (3.96 \pm 0.38) \times 10^{-11} \exp[-(544 \pm 28)/T]$. This result agrees well with the Arrhenius expression of $(3.38 \pm 0.17) \times 10^{-11} \exp[-(497 \pm 16)/T] \text{ cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$ reported by Wilson et al. (Wilson et al., 2006) between 241 and 406 K. By fitting our data and recommended data from multiple sources to the Arrhenius equation, the resulting Arrhenius expression at 240-1364 K is as follow: $k_{\text{n-Heptane}}(T) = 4.48 \times 10^{-12} \times \exp(116/T) \times (T/300)^{1.72} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. This result is in good agreement with the expression ($k_{\text{n-Heptane}}(T) = 4.39 \times 10^{-12} \times \exp(138/T) \times (T/300)^{1.70}$) obtained by Morin et al (Morin et al., 2015) at 248-896 K. Compared to the recommended data from the database of McGillen et al. (Mcgillen et al., 2020) at 240-1464 K, the ~~finger~~ activation energy (E_a/R) obtained in this work is more in line with that of (148), however, the ~~pre-finger~~ factor A obtained (4.48) is about 17% higher than that (3.84).

C. OH+ Isopentane. As Fig. 5 (c), isopentane was extensively studied over a temperature range (213-407 K). As far as we know, at present, only Wilson et al. has reported this compound in the range of 213-407 K (Wilson et al., 2006). Within the experimental temperature range (273-323 K), our data are consistent with Wilson et al. (273-323 K), especially in the low temperature range. The Arrhenius expression obtained at 273-323 K is $k_{\text{Isopentane}}(T) = (1.12 \pm 0.11) \times 10^{-11} \exp[-(443 \pm 34)/T]$. The Arrhenius

expression at 213-407 K obtained by fitting our data and those of Wilson et al. is as follows:
 $k_{\text{Isopentane}}(T) = (1.39 \pm 0.12) \times 10^{-11} \exp [-(424 \pm 25)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. The results are similar to the
relative experimental results of Wilson et al. $(1.52 \pm 0.21) \times 10^{-11} \exp [-(432 \pm 27)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.

D. OH+ 2,3-Dimethylbutane. Figure 5 (d) shows the Arrhenius plot for the reaction of 2,3-Dimethylbutane with OH radicals over the temperature range of 273 K to 1366 K. The temperature-dependent values obtained in this study at high temperature (313-323 K) align closely with those reported by Badra and Farooq (Badra and Farooq, 2015), who used the absolute rate technique, as well as the work of Sivaramakrishnan and Michael with a three-parameter fit (Sivaramakrishnan and Michael, 2009). However, the data obtained at 273-293 K in this work are highly consistent with the reviewed data from Atkinson and Arey (Atkinson and Arey, 2003). In the temperature range studied (273-323 K), the Arrhenius expression obtained in this work is $k_{2,3\text{-Dimethylbutane}}(T) = (1.15 \pm 0.09) \times 10^{-11} \exp [-(219 \pm 24)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. Linear regression applied to our data and high temperature data in the literature (at 273-1366 K) yields the Arrhenius expression as follows:
 $k_{2,3\text{-Dimethylbutane}}(T) = 1.29 \times 10^{-12} \times \exp(437/T) \times (T/300)^{2.09} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. This result agrees well with the Arrhenius expression of $k_{2,3\text{-Dimethylbutane}}(T) = 1.30 \times 10^{-12} \times \exp(427/T) \times (T/300)^{2.08} \text{ cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$ at 250-1366 K reported by Badra and Farooq (Badra and Farooq, 2015).




Figure 5. Arrhenius plots for the reaction of n-Octane (a), n-Heptane (b), Isopentane (c) and 2,3-Dimethylbutane (d) with OH radical in wide temperature range along with available literature data. The error bar was taken as 2σ . The expression for the non-linear curve fits takes the form $k(T) = A \times \exp(E_a/RT) \times (T/300)^n$, and the expression for linear regression takes the form $k(T) = A \times \exp(E_a/RT)$.

E. OH+ Methylcyclopentane (2-Methylhexane).

Figure 6 (a) and (b) illustrate the Arrhenius plot for the reaction of methylcyclopentane (230-1344 K) and 2-methylhexane (273-385) with OH radical. Literature data from Sprengnether et al. (Sprengnether et al., 2009) and Anderson et al. (Anderson et al., 2004) are available for comparison purposes. The rate coefficients of methylcyclopentane at 273-323 K in this work were obtained. Notably, for methylcyclopentane, Anderson et al. (Anderson et al., 2004) reported absolute data that is 26% higher than the relative data obtained in this study at 298 K. However, this difference falls within the margin of error. The absolute data from Sprengnether et al. (Sprengnether et al., 2009) is slightly higher, ranging from 10% to 20%, compared to this study. Fitting our data at 273-323 K yields the Arrhenius expression of $k_{\text{Methylcyclopentane}}(T) = (1.65 \pm 0.19) \times 10^{-11} \exp [-(262 \pm 33)/T]$.

Additionally, they derived an alternative Arrhenius expression to accommodate the curved behavior of the rate coefficient between 230 and 370 K, making it difficult to directly compare with our Arrhenius expression. Besides, in order to obtain temperature-dependent relationships over a wide temperature range, the experimental data obtained at 273-323 K are fitted with multi-party literature data, especially the data from Sivaramakrishnan and Michael at high temperature (859-1344 K), the resulting Arrhenius expression is as follows: $k_{\text{Methylcyclopentane}}(T) = 1.59 \times 10^{-12} \times \exp(439/T) \times (T/300)^{2.15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. The result is highly consistent with the expert-evaluated Arrhenius expression of methylcyclopentane ($k_{\text{Methylcyclopentane}}(T) = 1.67 \times 10^{-12} \times \exp(454/T) \times (T/300)^{2.15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$), indicating that the data obtained has a certain degree of reliability. Fitting our data at 273-323 K yields the Arrhenius expression of $k_{2\text{-Methylhexane}}(T) = (1.30 \pm 0.08) \times 10^{-11} \exp [-(222 \pm 19)/T]$. At present, the research on the temperature dependence of 2-Methylhexane only includes the measured reaction rate coefficient with OH radical of Sprengnether et al. by absolute rate technique at 230-385 K. The Arrhenius expression obtained by fitting our data with Sprengnether et al's data at 230-385 K is as follows: $k_{2\text{-Methylhexane}}(T) = (1.82 \pm 0.09) \times 10^{-11} \exp [-(321 \pm 16)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. Expert-evaluated Arrhenius expression is $k_{2\text{-Methylhexane}}(T) = (1.21 \pm 0.07) \times 10^{-11} \exp [-(171 \pm 16)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. Through comparison, it can be clearly seen that the two are consistent, indicating that the obtained Arrhenius expression has certain reference value. To the best of our knowledge, this is the first investigation of the temperature-dependent kinetics for the reaction of methylcyclopentane and 2-methylhexane with OH radicals utilizing the relative rate technique.

F. OH+ 3-Methylheptane. In Figure 6 (c), the Arrhenius plot presents the reaction between 3-Methylheptane and OH radicals, spanning a temperature range of 273 to 323 K. A linear regression analysis of our data yields the following Arrhenius expressions: $k_{3\text{-Methylheptane}}(T) = (2.72 \pm 0.34) \times 10^{-11} \exp [-(456 \pm 28)/T] \text{ cm}^3 \cdot \text{molecule}^{-1} \cdot \text{s}^{-1}$. We believe this study to be the first investigation of the temperature-dependent kinetics for the reaction between 3-Methylheptane and OH radicals. The only previous study on this reaction, reported by Shaw et al. (Shaw et al., 2020) utilizing the relative rate method in nitrogen at 323 K, demonstrates significantly higher data (>65%) compared to our results. Possible explanations for this discrepancy lie in the different reference compounds used and potential sample loss during sampling in the enrichment tube in Shaw et al.

527 **G. OH+ 3-Methylhexane (Figure 6 (d)).** This is the first temperature-dependence relative data. It

528 can be seen from the figure that this data ~~is this data~~ is significantly lower by approximately 80%

529 compared to the absolute data. The Arrhenius expression at 273-323 K is as follows:

530 $k_{3\text{-Methylhexane}}(T) = (2.53 \pm 1.45) \times 10^{-11} \exp [-(575 \pm 161)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$

531 **H. OH+ 2-Methylheptane (Figure 6 (e)).** There are no previous temperature dependence data on

532 this compound. Similar to 3-Methylhexane, this data is lower by approximately 37% compared to Shaw

533 et al. at room temperature. Within the range of 273-323 K, the obtained Arrhenius expression is as follows:

534 $k_{2\text{-Methylheptane}}(T) = (1.37 \pm 0.48) \times 10^{-11} \exp [-(209 \pm 100)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.




Figure 6. Arrhenius plots for the reaction of Methylcyclopentane (a), 2-Methylhexane (b), 3-Methylheptane (c), 3-Methylhexane (d) and 2-Methylheptane (e) with OH radical along with available literature data. The error bar was taken as 2σ . The expression for the non-linear curve fits takes the form $k(T) = A \times \exp(E_a/RT) \times (T/300)^n$, and the expression for linear regression takes the form $k(T) = A \times \exp(E_a/RT)$.

3.4 Correlation between the rate coefficients of the reaction of alkanes with OH radicals and chlorine atoms

Figure 7 presents a log-log correlation plot between the Cl atoms and OH radical rate coefficients with the series of C₃-C₁₁ studied above. A very clear correlation ($R^2=0.86$) described by the relation $\log_{10}[k_{(\text{Cl}+\text{alkanes})}] = 0.569 \times \log_{10}[k_{(\text{OH}+\text{alkanes})}] - 3.111$ was obtained. Although the correlation between propane and isobutane is relatively weak, the reactivity of saturated alkanes with OH radicals and chlorine atoms is still clearly related ~~to the saturated alkane series~~. In addition, the log-log correlation for the series of saturated alkanes with these two oxidants presented by Calvert et al. (2011) described by the relation $\log_{10}[k_{(\text{Cl}+\text{alkanes})}] = 0.521 \times \log_{10}[k_{(\text{OH}+\text{alkanes})}] - 3.670$ with ($R^2=0.85$) is in better agreement with the log-log correlations obtained in this study for saturated alkanes. This correlation can be utilized to predict rate coefficients for unmeasured reactions, such as the reaction of 2,2,3-trimethylpentane with chlorine atoms. It is currently known that the rate coefficient for the reaction of 2,2,3-trimethylpentane with OH radical at room temperature is $4.84 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, according to the above correlation equation, it can be inferred that the rate coefficient with Cl atoms is $2.72 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.




Figure 7. ~~Double logarithmic plot (log₁₀)~~ Log-log plot of the rate coefficients for the reaction of Cl-atoms versus the reaction of OH radicals with the saturated alkanes (C₃-C₁₁ alkanes studied above). The solid line represents the unweighted least-squares fit to the data. ~~The alkanes represented by serial number can~~

be identified as follows: (1) Propane; (2) Isobutane; (3) n-Butane; (4) Isopentane; (5) n-pentane; (6) Cyclopentane; (7) 2,3-Dimethylbutane; (8) 2-Methylpentane; (9) 3-Methylpentane; (10) Methylcyclopentane; (11) 2,4-Dimethylpentane; (12) Cyclohexane; (13) 2-Methylhexane; (14) 2,2,4-Trimethylpentane; (15) n-Heptane; (16) Methylcyclohexane; (17) n-Octane; (18) n-nonane; (19) n-Decane; (20) n-Undecane.

3.5 Atmospheric lifetime and implications

~~The atmospheric lifetime of alkanes in the troposphere can be estimated~~ The atmospheric lifetime of alkanes in the troposphere, due to their reaction with OH radicals, can be estimated using the following formula:

$$\tau_{\text{alkane}} = 1 / (k_{\text{alkane}+\text{OH}}[\text{OH}])$$

Where τ_{alkane} is the atmospheric lifetime of the alkane due to OH removal, $k_{\text{alkane}+\text{OH}}$ is the rate coefficient for the reaction of the alkane with OH radical at the typical tropospheric temperature of 298 K, and [OH] is the atmospheric concentrations of the hydroxyl radicals. The average tropospheric hydroxyl radical concentration has been previously reported in the literature as 1×10^6 molecules cm^{-3} (Lawrence et al., 2001). Using the $k_{\text{alkane}+\text{OH}}$ (298 K) values determined in the present work, the atmospheric lifetime for 25 alkanes was estimated and listed in Table S4. As can be seen from the table, the atmospheric lifetimes of C₃-C₁₁ alkanes reacting with OH radicals are about 1-11 days. As the carbon chain grows, the atmospheric lifetimes are reduced, especially for long-chain alkanes with carbon atoms of 8-11, the residence time in the atmosphere is only about 1 day. They are emitted into the air and degraded quickly to generate alkyl radicals, which are immediately converted into alkyl peroxy radicals by reacting with abundant O₂ in the atmosphere. Alkyl peroxy radicals will serve to convert NO to NO₂ directly, leading to the production of tropospheric ozone. Longer atmospheric residence time of short-chain alkanes compared to long-chain C₈-C₁₁ alkanes, such as propane, the lifetime is 11 days.

4. Conclusions

The use of the multivariate relative rate method in this study allowed for the simultaneous determination of reaction rate coefficients of C₃-C₁₁ alkanes and OH radicals, which significantly improved the efficiency of determination. A total of 25 relative rate coefficients at room temperature were obtained,

including the determination of a previously unreported room temperature relative rate coefficient for 3-methylheptane. For the studied n-alkanes, the obtained rate coefficients (k_{OH}) were found to be consistent with results estimated by the SAR methods using parameters provided by various **positional** groups, such as Atkinson and Kwok, Neeb, Wilson, Jenkin, and McGillen. However, it is important to note that parameters other than those provided by Wilson group do not appear to reasonably estimate the rate coefficients of 2,3-dimethylbutane. Additionally, SAR estimates for several cyclic alkanes (cyclopentane, methylcyclopentane, cyclohexane) and branched alkanes (2,2,4-trimethylpentane) appear to be overestimated compared to our measurements. This raises reasonable suspicion that these methods may still lack consideration of additional factors, **and a more appropriate empirical ring strain factor needs to be derived based on broader range of experimental data from monocyclic hydrocarbons in the future.** Arrhenius expressions for the reaction of ~~2-Methylhepane~~ 2-Methylheptane and 3-Methylheptane with OH radicals were obtained for the first time in the temperature range of 273-323 K, expanding the existing database. **And the rate coefficients do not change significantly in this temperature range, especially for 2-methylheptane. Whilst the value of preexponential factor A increases with the increase of the number of carbon atoms, which is consistent with the law of its reactivity.** In addition, correlation equations for the rate coefficients of alkanes reacting with OH radicals and chlorine atoms were obtained, and the rate coefficient of 2,2,3-trimethylpentane with chlorine atoms, which has not yet been reported, was deduced. The atmospheric lifetimes of the alkanes were also obtained for further prediction of their environmental impact.

Data availability

Raw data are available upon request.

Author contributions

YM and CL planned the campaign; YX performed the measurements; YX, CL, YM and XL analyzed the data; YX and CL wrote the manuscript draft. SX and JL provided technical support.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (Nos. 22076202, 42077454 and 41975164).

References

- Andersen, M. P. S., Hurley, M. D., Ball, J. C., Schneider, W. F., Wallington, T. J., and Nielsen, O. J.: $\text{CF}_3\text{CH}(\text{ONO})\text{CF}_3$: synthesis, IR spectrum, and use as OH radical source for kinetic and mechanistic studies, *Int. J. Chem. Kinet.*, 35, 159-165, <https://doi.org/10.1002/kin.10116>, 2003.
- Anderson, R. S., Huang, L., Iannone, R., Thompson, A. E., and Rudolph, J.: Carbon kinetic isotope effects in the gas phase reactions of light alkanes and ethene with the OH radical at 296 ± 4 K, *J. Phys. Chem. A*, 108, 11537-11544, <https://doi.org/10.1021/jp0472008>, 2004.
- Atkinson, R.: Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions, *Chem. Rev.*, 86, 69-201, <https://doi.org/10.1021/cr00071a004>, 1986.
- Atkinson, R.: Atmospheric chemistry of VOCs and NO_x , *Atmos. Environ.*, 34, 2063-2101, [https://doi.org/10.1016/S1352-2310\(99\)00460-4](https://doi.org/10.1016/S1352-2310(99)00460-4), 2000.
- Atkinson, R.: Kinetics of the gas-phase reactions of OH radicals with alkanes and cycloalkanes, *Atmos. Chem. Phys.*, 3, 2233-2307, <https://doi.org/10.5194/acp-3-2233-2003>, 2003.
- Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chem. Rev.*, 103, 4605-4638, <https://doi.org/10.1021/cr0206420>, 2003.
- Atkinson, R., Aschmann, S. M., Carter, W. P. L., Winer, A. M., and Pitts Jr., J. N.: Kinetics of the reactions of OH radicals with n-alkanes at 299 ± 2 K, *Int. J. Chem. Kinet.*, 14, 781-788, <https://doi.org/10.1002/kin.550140706>, 1982.
- Badra, J. and Farooq, A.: Site-specific reaction rate constant measurements for various secondary and tertiary H-abstraction by OH radicals, *Combust. Flame*, 162, 2034-2044, <https://doi.org/10.1016/j.combustflame.2015.01.001>, 2015.
- Ballesteros, B., Ceacero-Vega, A. A., Jimenez, E., and Albaladejo, J.: Atmospheric reactions of methylcyclohexanes with Cl atoms and OH radicals: determination of rate coefficients and degradation products, *Environ. Sci. Pollut. Res. Int.*, 22, 4806-4819, <https://doi.org/10.1007/s11356-014-2901-0>, 2015.

641 Bejan, I. G., Winiberg, F. A. F., Mortimer, N., Medeiros, D. J., Brumby, C. A., Orr, S. C., Kelly, J., and
 642 Seakins, P. W.: Gas-phase rate coefficients for a series of alkyl cyclohexanes with OH radicals and Cl
 643 atoms, *Int. J. Chem. Kinet.*, 50, 544-555, <https://doi.org/10.1002/kin.21179>, 2018.
 644 Bryukov, M. G., Knyazev, V. D., Lomnicki, S. M., McFerrin, C. A., and Dellinger, B.: Temperature-
 645 dependent kinetics of the gas-phase reactions of OH with Cl₂, CH₄, and C₃H₈, *J. Phys. Chem. A*, 108,
 646 <https://doi.org/10.1021/jp047340h>, 2004.
 647 Calvert, J. G., Orlando, J. J., Stockwell, W. R., and Wallington, T. J.: The mechanisms of reactions
 648 influencing atmospheric ozone, Oxford University Press, Incorporated, New York, United States, 609 pp.,
 649 ISBN 9780190233037, 2015.
 650 Cohen, N.: Are reaction rate coefficients additive? Revised transition state theory calculations for OH +
 651 alkane reactions, *Int. J. Chem. Kinet.*, 23, 397-417, <https://doi.org/10.1002/kin.550230506>, 1991.
 652 Cox, R. A., Derwent, R. G., and Williams, M. R.: Atmospheric photooxidation reactions. rates, reactivity,
 653 and mechanism for reaction of organic compounds with hydroxyl radicals, *Environ. Sci. Technol.*, 14, 57-
 654 61, <https://doi.org/10.1021/es60161a007>, 1980.
 655 Crawford, M. A., Dang, B., Hoang, J., and Li, Z.: Kinetic study of OH radical reaction with n-heptane
 656 and n-hexane at 240–340K using the relative rate/discharge flow/mass spectrometry (RR/DF/MS)
 657 technique, *Int. J. Chem. Kinet.*, 43, 489-497, <https://doi.org/10.1002/kin.20574>, 2011.
 658 Darnall, K. R., Atkinson, R., and Pitts, J. N.: Rate constants for the reaction of the OH radical with selected
 659 alkanes at 300 K, *J. Phys. Chem.*, 82, 1581-1584, <https://doi.org/10.1021/j100503a001>, 1978.
 660 DeMore, W. and Bayes, K.: Rate constants for the reactions of hydroxyl radical with several alkanes,
 661 cycloalkanes, and dimethyl ether, *J. Phys. Chem. A*, <https://doi.org/10.1021/j100503a001>, 1999.
 662 Donahue, N. M., Anderson, J. G., and Demerjian, K. L.: New rate constants for ten OH alkane reactions
 663 from 300 to 400 K: an assessment of accuracy, *J. Phys. Chem. A*, <https://doi.org/10.1021/j100503a001>, 1998.
 664 Droege, A. T. and Tully, F. P.: Hydrogen-atom abstraction from alkanes by hydroxyl radical. 6.
 665 cyclopentane and cyclohexane, *J. Phys. Chem.*, 91, <https://doi.org/10.1021/j100503a001>, 1987.
 666 Dunmore, R. E., Hopkins, J. R., Lidster, R. T., Lee, J. D., Evans, M. J., Rickard, A. R., Lewis, A. C., and
 667 Hamilton, J. F.: Diesel-related hydrocarbons can dominate gas phase reactive carbon in megacities,
 668 *Atmos. Chem. Phys.*, 15, 9983-9996, <https://doi.org/10.5194/acp-15-9983-2015>, 2015.
 669 Edney, E., Kleindienst, T., and Corse, E.: Room temperature rate constants for the reaction of OH with
 670 selected chlorinated and oxygenated hydrocarbons, *Int. J. Chem. Kinet.*, 18, 1355-1371,

671 <https://doi.org/10.1002/kin.550181207>, 1986.

672 Ferrari, C., Roche, A., Jacob, V., Foster, P., and Baussand, P.: Kinetics of the reaction of OH radicals with
 673 a series of esters under simulated conditions at 295 K, *Int. J. Chem. Kinet.*, 28, 609-614,
 674 [https://doi.org/10.1002/\(sici\)1097-4601\(1996\)28:8<609::aid-kin6>3.0.co;2-z](https://doi.org/10.1002/(sici)1097-4601(1996)28:8<609::aid-kin6>3.0.co;2-z), 1996.

675 Finlaysonpitts, B. J. and Pitts, J. N., Jr.: Tropospheric air pollution: ozone, airborne toxics, polycyclic
 676 aromatic hydrocarbons, and particles, *Science* (New York, N.Y.), 276, <https://doi.org/1045-1052>,
 677 10.1126/science.276.5315.1045, 1997.

678 Finlaysonpitts, B. J., Hernandez, S. K., and Berko, H. N.: A new dark source of the gaseous hydroxyl
 679 radical for relative rate measurements, *J. Phys. Chem.*, 97, 1172-1177,
 680 <https://doi.org/10.1021/j100108a012>, 1993.

681 Fiore, A. M., Horowitz, L. W., Purves, D. W., Levy, H., Evans, M. J., Wang, Y. X., Li, Q. B., and Yantosca,
 682 R. M.: Evaluating the contribution of changes in isoprene emissions to surface ozone trends over the
 683 eastern United States, *J. Geophys. Res.-Atmos.*, 110, 18, <https://doi.org/10.1029/2004jd005485>, 2005.

684 Goldstein, A. H. and Galbally, I. E.: Known and unexplored organic constituents in the earth's atmosphere,
 685 *Environ. Sci. Technol.*, 41, <http://doi.org/1514-1521>, 10.1021/es072476p, 2007.

686 Gordon, S. and Mulac, W.: Reaction of the OH ($X^{\bullet} 71'$) radical produced by the pulse radiolysis of water
 687 vapor *Int. J. Chem. Kinet.*, (Syrnp. 1), 289, 299, 1975.

688 Gorse, R. A. and Volman, D. H.: Photochemistry of the gaseous hydrogen peroxide—carbon monoxide
 689 system. II: Rate constants for hydroxyl radical reactions with hydrocarbons and for hydrogen atom
 690 reactions with hydrogen peroxide, *J. Photochem.*, 3, 115-122, [https://doi.org/10.1016/0047-](https://doi.org/10.1016/0047-2670(74)80011-0)
 691 2670(74)80011-0, 1974.

692 Greiner, N. R.: Hydroxyl-Radical Kinetics by Kinetic Spectroscopy. II. Reactions with C₂H₆, C₃H₈, and
 693 iso-C₄H₁₀ at 300°K, *J. Chem. Phys.*, 46, 3389-3392, <https://doi.org/10.1063/1.1841228>, 1967.

694 Greiner, N. R.: Hydroxyl radical kinetics by kinetic spectroscopy. VI. Reactions with Alkanes in the range
 695 300–500°K, *J. Chem. Phys.*, 53, 1070-&, <https://doi.org/10.1063/1.1674099>, 1970a.

696 Greiner, N. R.: Comparison of the kinetics of alkane H-atom abstraction by methyl and hydroxyl radicals,
 697 *J. Chem. Phys.*, 53, 1285-1287, <https://doi.org/10.1063/1.1674134>, 1970b.

698 Guenther, A.: The contribution of reactive carbon emissions from vegetation to the carbon balance of
 699 terrestrial ecosystems, *Chemosphere*, 49, 837-844, [https://doi.org/10.1016/s0045-6535\(02\)00384-3](https://doi.org/10.1016/s0045-6535(02)00384-3), 2002.

700 Harris, S. J. and Kerr, J. A.: Relative Rate measurements of some reactions of hydroxyl radicals with

alkanes studied under atmospheric conditions, *Int. J. Chem. Kinet.*, 20, 939-955, <https://doi.org/10.1002/kin.550201203>, 1988.

Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R., and Wallington, T. J.: Estimation of rate coefficients and branching ratios for gas-phase reactions of OH with aliphatic organic compounds for use in automated mechanism construction, *Atmos. Chem. Phys.*, 18, 9297-9328, <https://doi.org/10.5194/acp-18-9297-2018>, 2018.

Kwok, E. S. and Atkinson, R.: Estimation of hydroxyl radical reaction rate constants for gas-phase organic compounds using a structure-reactivity relationship: an update, *Atmos. Environ.*, 29, 1685-1695, [https://doi.org/10.1016/1352-2310\(95\)00069-b](https://doi.org/10.1016/1352-2310(95)00069-b), 1995.

Lawrence, M. G., Jöckel, P., and von Kuhlmann, R.: What does the global mean OH concentration tell us?, *Atmos. Chem. Phys.*, 1, 37-49, <https://doi.org/10.5194/acp-1-37-2001>, 2001.

Lewis, A. C., Carslaw, N., Marriott, P. J., Kinghorn, R. M., Morrison, P., Lee, A. L., Bartle, K. D., and Pilling, M. J.: A larger pool of ozone-forming carbon compounds in urban atmospheres, *Nature*, 405, 778-781, <https://doi.org/10.1038/35015540>, 2000.

Li, Z. J., Singh, S., Woodward, W., and Dang, L.: Kinetics study of OH radical reactions with n-octane, n-nonane, and n-decane at 240-340 K using the relative rate/discharge flow/mass spectrometry technique, *J. Phys. Chem. A*, 110, 12150-12157, <https://doi.org/10.1021/jp0638134>, 2006.

Liang, S., Gao, S., Wang, S., Chai, W., Chen, W., and Tang, G.: Characteristics, sources of volatile organic compounds, and their contributions to secondary air pollution during different periods in Beijing, China, *Sci. Total. Environ.*, 858, 159831, <https://doi.org/10.1016/j.scitotenv.2022.159831>, 2023.

Ling, Z. H. and Guo, H.: Contribution of VOC sources to photochemical ozone formation and its control policy implication in Hong Kong, *Environ. Sci. Policy*, 38, 180-191, <https://doi.org/10.1016/j.envsci.2013.12.004>, 2014.

Liu, C. T., Mu, Y. J., Zhang, C. L., Zhang, Z. B., Zhang, Y. Y., Liu, J. F., Sheng, J. J., and Quan, J. N.: Development of gas chromatography-flame ionization detection system with a single column and liquid nitrogen-free for measuring atmospheric C₂-C₁₂ hydrocarbons, *J. Chromatogr. A*, 1427, 134-141, <https://doi.org/10.1016/j.chroma.2015.11.060>, 2016.

McGillen, M. R., Carter, W. P. L., Mellouki, A., Orlando, J. J., Picquet-Varrault, B., and Wallington, T. J.: Database for the kinetics of the gas-phase atmospheric reactions of organic compounds, *Earth Syst. Sci. Data*, 12, 1203-1216, <https://doi.org/10.5194/essd-12-1203-2020>, 2020.

731 Mellouki, A., Téton, S., Laverdet, G., Quilgars, A., and Le Bras, G.: Kinetic studies of OH reactions with
 732 H₂O₂, C₃H₈ and CH₄ using the pulsed laser photolysis-laser induced fluorescence method, *J. Chim. Phys.*
 733 *PCB.*, 91, 473-487, <https://doi.org/10.1051/jcp/1994910473>, 1994.

734 Morin, J., Romanias, M. N., and Bedjanian, Y.: Experimental study of the reactions of OH radicals with
 735 Propane, n-Pentane, and n-Heptane over a wide temperature range, *Int. J. Chem. Kinet.*, 47, 629-637,
 736 <https://doi.org/10.1002/kin.20936>, 2015.

737 Neeb, P.: Structure-reactivity based estimation of the rate constants for hydroxyl radical reactions with
 738 hydrocarbons, *J. Atmos. Chem.*, 35, 295-315, <https://doi.org/10.1023/a:1006278410328>, 2000.

739 Overend, R. P., Paraskevopoulos, G., and Cvetanovic, R. J.: Rates of OH Radical Reactions. I. Reactions
 740 with H₂, CH₄, C₂H₆, and C₃H₈ at 295 K, *Can. J. Chem.*, <https://doi.org/10.1139/v75-482>, 1975.

741 Pang, G. A., Hanson, R. K., Golden, D. M., and Bowman, C. T.: High-Temperature Measurements of the
 742 Rate Constants for Reactions of OH with a Series of Large Normal Alkanes: n-Pentane, n-Heptane, and
 743 n-Nonane, *Z. Phys. Chem.*, 225, 1157-1178, <https://doi.org/10.1524/zpch.2011.0156>, 2011.

744 Perry, R. A., Atkinson, R., and Pitts, J. N.: Rate constants for the reaction of OH radicals with nbutane
 745 over the temperature range 297–420°K, *J. Chem. Phys.*, 64, 5314-5316, <https://doi.org/10.1063/1.432167>,
 746 1976.

747 Phan, M. and Li, Z. J.: Kinetics Study of the Reactions of OH with n-Undecane and n-Dodecane Using
 748 the RR/DF/MS Technique, *J. Phys. Chem. A*, 121, 3647-3654, <https://doi.org/10.1021/acs.jpca.7b01512>,
 749 2017.

750 Shaw, J. T., Rickard, A. R., Newland, M. J., and Dillon, T. J.: Rate coefficients for reactions of OH with
 751 aromatic and aliphatic volatile organic compounds determined by the multivariate relative rate technique,
 752 *Atmos. Chem. Phys.*, 20, 9725-9736, <https://doi.org/10.5194/acp-20-9725-2020>, 2020.

753 Shaw, J. T., Lidster, R. T., Cryer, D. R., Ramirez, N., Whiting, F. C., Boustead, G. A., Whalley, L. K.,
 754 Ingham, T., Rickard, A. R., Dunmore, R. E., Heard, D. E., Lewis, A. C., Carpenter, L. J., Hamilton, J. F.,
 755 and Dillon, T. J.: A self-consistent, multivariate method for the determination of gas-phase rate
 756 coefficients, applied to reactions of atmospheric VOCs and the hydroxyl radical, *Atmos. Chem. Phys.*, 18,
 757 4039-4054, <https://doi.org/10.5194/acp-18-4039-2018>, 2018.

758 Shi, B., Wang, W., Zhou, L., Li, J., Wang, J., Chen, Y., Zhang, W., and Ge, M.: Kinetics and mechanisms
 759 of the gas-phase reactions of OH radicals with three C₁₅ alkanes, *Atmos. Environ.*, 207, 75-81,
 760 <https://doi.org/10.1016/j.atmosenv.2019.03.028>, 2019.

761 Singh, S., de Leon, M. F., and Li, Z. J.: Kinetics Study of the Reaction of OH Radicals with C₅-C₈
 762 Cycloalkanes at 240-340 K using the Relative Rate/Discharge Flow/Mass Spectrometry Technique, J.
 763 Phys. Chem. A, 117, 10863-10872, <https://doi.org/10.1021/jp406923d>, 2013.
 764 Sivaramakrishnan, R. and Michael, J. V.: Rate constants for OH with selected large alkanes: shock-tube
 765 measurements and an improved group scheme, J. Phys. Chem. A, 113, 5047-5060,
 766 <https://doi.org/10.1021/jp810987u>, 2009
 767 Sprengnether, M. M., Demerjian, K. L., Dransfield, T. J., Clarke, J. S., Anderson, J. G., and Donahue, N.
 768 M.: Rate constants of nine C₆-C₉ alkanes with OH from 230 to 379 K: chemical tracers for OH, J. Phys.
 769 Chem. A, 113, 5030-5038, <https://doi.org/10.1021/jp810412m>, 2009.
 770 Sun, J., Wu, F. K., Hu, B., Tang, G. Q., Zhang, J. K., and Wang, Y. S.: VOC characteristics, emissions and
 771 contributions to SOA formation during hazy episodes, Atmos. Environ., 141, 560-570,
 772 <https://doi.org/10.1016/j.atmosenv.2016.06.060>, 2016.
 773 Talukdar, R. K., Mellouki, A., Gierczak, T., Barone, S., Chiang, S. Y., and Ravishankara, A. R.: Kinetics
 774 of the reactions of OH with alkanes, Int. J. Chem. Kinet., 26, 973-990,
 775 <https://doi.org/10.1002/kin.550261003>, 1994.
 776 Tully, F. P., Goldsmith, J. E. M., and Droege, A. T.: Hydrogen-atom abstraction from alkanes by OH. 4.
 777 Isobutane, J. Phys. Chem., 90, 5932-5937, <https://doi.org/10.1021/j100280a095>, 1986.
 778 Wilson, E. W., Hamilton, W. A., Kennington, H. R., Evans, B., Scott, N. W., and DeMore, W. B.:
 779 Measurement and estimation of rate constants for the reactions of hydroxyl radical with several alkanes
 780 and cycloalkanes, J. Phys. Chem. A, 110, <https://doi.org/10.1021/jp055841c>, 2006.