

# Ocean alkalinity enhancement using sodium carbonate salts does not lead to measurable changes in ~~impact~~-Fe dynamics in a mesocosm experiment

David González-Santana<sup>1</sup>, María Segovia<sup>2</sup>, Melchor González-Dávila<sup>1</sup>, Librada Ramírez<sup>2</sup>, Aridane G. González<sup>1</sup>, Leonardo J. Pozzo-Pirotta<sup>2</sup>, Veronica Arnone<sup>1</sup>, Víctor Vázquez<sup>2</sup>, Ulf Riebesell<sup>3</sup>, J. Magdalena Santana-Casiano<sup>1</sup>.


<sup>1</sup>Instituto de Oceanografía y Cambio Global, IOCAG, Universidad de Las Palmas de Gran Canaria, ULPGC, Spain.

<sup>2</sup>Department of Ecology, Faculty of Sciences, University of Malaga, Málaga, Spain.

<sup>3</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany.

*Correspondence to:* David González-Santana (David.gonzalez@fpct.ulpgc.es)

**Abstract.** The addition of carbonate minerals to seawater through an artificial Ocean Alkalinity Enhancement (OAE) process increases the concentrations of hydroxide, bicarbonate, and carbonate ions. This leads to changes in the pH and the buffering capacity of the seawater. Consequently, OAE could have relevant effects on marine organisms and in the speciation and concentration of trace metals that are essential for their physiology. During September and October 2021, a mesocosm experiment was carried out in the coastal waters of Gran Canaria (Spain), consisting on the controlled variation of total alkalinity (TA). Different concentrations of carbonate salts ( $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$ ) previously homogenized were added to each mesocosm to achieve an alkalinity gradient between  $\Delta 0$  to  $\Delta 2400 \mu\text{mol L}^{-1}$ . The lowest point of the gradient was  $2400 \mu\text{mol kg}^{-1}$ , being the natural alkalinity of the medium, and the highest point was  $4800 \mu\text{mol kg}^{-1}$ . Iron (Fe) speciation was monitored during this experiment to analyse whether total dissolved iron (TdFe, unfiltered samples), dissolved iron (dFe, filtered through a  $0.2 \mu\text{m}$  pore size filter), soluble iron (sFe, filtered through a  $0.02 \mu\text{m}$  pore size filter), dissolved labile iron (dFe'), iron-binding ligands (LFe) and their conditional stability constants ( $K'_{\text{FeL}}$ ), because of change due to OAE and the experimental conditions in each mesocosm. Observed iron concentrations were within the expected range for coastal waters, with no significant increases due to OAE. However, there were variations in Fe size fractionation during the experiment. This could potentially be due to chemical changes caused by OAE, but such effect being masked by the stronger biological interactions. In terms of size fractionation, sFe was below  $1.0 \text{ nmol L}^{-1}$ , dFe concentrations were within  $0.5\text{-}4.0 \text{ nmol L}^{-1}$ , and TdFe within  $1.5\text{-}7.5 \text{ nmol L}^{-1}$ . Our results show that over 99% of Fe was complexed, mainly by  $L_1$  and  $L_2$  ligands with  $k'_{\text{FeL}}$  ranging between  $10.92 \pm 0.11$  and  $12.68 \pm 0.32$ , with LFe ranging from  $1.51 \pm 0.18$  to  $12.3 \pm 1.8 \text{ nmol L}^{-1}$ . Our data on iron size fractionation, concentration, and iron-binding ligands substantiate that the introduction of sodium salts in this mesocosm experiment did not modify iron dynamics. As a consequence, phytoplankton remained unaffected by alterations in this crucial element.



## 1. Introduction

Artificial Ocean Alkalinity Enhancement (OAE) ~~is a negative emission technology (NET) that~~ can lead to the atmospheric carbon dioxide removal (CDR) by increasing the ocean carbon uptake and reversing ocean acidification (OA), favoring a higher ocean pH buffering capacity (Kheshgi, 1995; Renforth and Henderson, 2017; Lenton et al., 2018). OAE can be likened to an acceleration of natural chemical weathering processes of minerals. Weathering reactions have played a role in modulating climate on geological time scales (Zeebe, 2012; Colbourn et al., 2015).

OAE has consequences for the  $\text{CO}_2$ -carbonate system in the ocean. The increase in ocean alkalinity raises both total dissolved inorganic carbon (DIC) and pH, subsequently increasing the carbonate ion concentration, as well as decreasing  $\text{pCO}_2$ . This increases the  $\text{pCO}_2$  imbalance between the ocean and atmosphere leading to an enhancement of ocean carbon uptake.

~~Moreover, the increased pH and carbonate ion concentration reverses ocean acidification due to the uptake of anthropogenic  $\text{CO}_2$  from the atmosphere (Lenton et al., 2018).~~ The artificial augmentation of alkalinity could also have consequences for marine ecosystems, both for the biota and for the chemical constituents of seawater, including elements that are essential for the ecosystem (Feng et al., 2016). The consumption of atmospheric  $\text{CO}_2$  and pH increase during OAE may be accompanied by changes in the release of mineral dissolution products such as Si, Ca, Mg, Fe or Ni. The ecological and biogeochemical consequences of OAE will also depend on the minerals used during alkalization (e.g.,  $\text{Mg}_2\text{SiO}_4$ , CaO,  $\text{Na}_2\text{CO}_3$ ) (Renforth and Henderson, 2017; Hartmann et al., 2013; Bach et al., 2019). Hence, it is necessary to perform experiments that may support or dissuade the use and viability of this technique under different experimental conditions. One controlled, i.e. semi-natural,

50 way to perform these types of experiments is by using mesocosms. Mesocosm studies allow rigorous testing of NETs-mCDR at the ecosystem level since the only variable that is artificially modified is the variable of interest (OAE), allowing the enclosed water column to maintain its chemical and biological properties.

OAE could have an impact on trace metals that are essential for marine organisms and this effect has been scarcely investigated so far (Guo et al., 2022; Xin et al., 2023). Among metals, Fe is the most essential micronutrient controlling phytoplankton  
55 growth mainly through nitrate assimilation and photosynthesis (Behrenfeld and Milligan, 2013). Thus, the growth and development of marine phytoplankton is linked to the biogeochemical cycling of Fe (Jickells et al., 2005; Moore et al., 2001), by regulating both the structure and the productivity of marine ecosystems (Tagliabue et al., 2017; Boyd and Ellwood, 2010). The assimilation of iron by marine organisms is not straightforward. Fe bioavailability, i.e. iron available for uptake, and therefore growth, is controlled by many factors. It depends on the different forms (chemical speciation) in which iron is found  
60 in solution (Sutak et al., 2020). Indeed, not all iron species are bioavailable, only those in the dissolved (dFe) phase. Under oxygenated ocean conditions, the most abundant form of iron is the insoluble form of oxidized iron, Fe(III). pH is the main variable that controls Fe speciation in this condition. Fe(II) is thermodynamically unstable and oxidized on timescales of minutes to hours (Santana-Casiano et al., 2005; González-Santana et al., 2021). Higher pH alters iron speciation, decreases Fe solubility and Fe(II) half-life, and thus decreases dissolved iron (dFe) concentrations (Liu and Millero, 2002; Boye et al., 2006; González-Dávila et al., 2006). However, there are several factors that can significantly alter and control iron speciation, such as photooxidation in surface waters, the presence of organic ligands (L) and the decreasing oxygen concentration (Santana-Casiano et al., 2022; González et al., 2019; Benner, 2011; Barbeau, 2006; Moffett, 2021; Hopkinson and Barbeau, 2007). All of these factors can induce a reduction of Fe(III) to Fe(II) depending on the conditions and characteristics of the organic compounds involved. Moreover, organic matter plays an essential role, because it can complex up to 99% of the iron (Fe(III)-  
70 L) forming part of the dFe (Wu and Luther III, 1994; Arnone et al., 2022). The concentration and strength of iron-binding organic ligands ( $L_{Fe}$ ) modulate Fe speciation and its bioavailability. In addition, particle size (physical speciation) is key for Fe cycling, since the colloidal size fraction remains in solution while larger fractions tend to sink down, being excluded from the system. All these processes were demonstrated during a mesocosm experiment, in a Norwegian fjord, that studied the effects of ocean acidification. In this experiment, both low pH and strong organic ligands enhanced the solubility of particulate  
75 and colloidal Fe (Segovia et al., 2017; Lorenzo et al., 2020). This promoted significant higher Fe availability to phytoplankton, directly impacting on their physiology and controlling the phytoplankton community structure in coastal ecosystems (Segovia et al., 2017; Lorenzo et al., 2020; Mausz et al., 2020).

The introduction of carbonate minerals into seawater for OAE (Bach et al., 2019) is expected to increase the concentrations of hydroxide, bicarbonate and carbonate ( $OH^-$ ,  $HCO_3^-$  and  $CO_3^{2-}$ ). This will change metal ion speciation in seawater (Millero et al., 2009). Metals that form complexes with  $OH^-$  and  $CO_3^{2-}$ , such as Fe, may be less abundant in their free forms at higher pH. Moreover, Fe(II) oxidation rate constants are strongly increased under high  $HCO_3^-$  and  $CO_3^{2-}$  concentrations (Santana-Casiano et al., 2005, 2006) making Fe less bioavailable. AOE-OAE can also cause a release of  $Ca^{2+}$  or  $Mg^{2+}$  ions (Bach et al., 2019).


These ions compete with Fe for specific functional groups of the organic ligand, which would affect the redox speciation of this metal (Santana-Casiano et al., 2010).

- 85 In this sense, the impact of OAE on the iron cycle could be that the Fe speciation will change with enhanced alkalinity. In the presence of organic compounds produced by the organisms in the mesocosms, the formation of strong organic complexes (FeL) will dominate the speciation of Fe. The  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  ions produced in ~~AOE-OAE~~ if  $\text{Mg}_2\text{SiO}_4$  or CaO were used, could compete with Fe for the organic ligands, which would affect the Fe redox speciation and the bioavailability. To avoid this effect, alkalization with  $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$  is recommended.
- 90 The aim of this research was to study the evolution of inorganic and organic Fe speciation in seawater under OAE scenarios. We considered if the added material acted as a significant iron source, and its effects on both the iron size fractionation and Fe-binding ligand evolution.

## 2. Material and Methods

### 2.1. Mesocosms experimental design and distribution

- 95 The mesocosm experiment was conducted in the port of Taliarte, Telde, Gran Canaria during a 33-day period from September to October 2021. Nine mesocosms (M1-M9) were deployed in the port of Taliarte containing a natural seawater column of 8.3  $\text{m}^3$  each at 5 m depth, with a sediment trap attached to the bottom of the bag (Riebesell et al., 2013; Taucher et al., 2017) (Fig. 1).



100 **Figure 1. Study site. a) Location in the Gran Canaria port town of Taliarte, Spain (Google Earth Pro); b) Operating mesocosms during the experiment (photograph by Peter Yeung, National Geographic ©); c) Mesocosms experimental design representing the alkalinity gradient (OAE) ( $\mu\text{mol L}^{-1}$ ) in the nine mesocosms: M5 (0), M1 (300), M7 (600), M4 (900), M9 (1200), M3 (1500), M6 (1800), M2 (2100), M8 (2400) (design by Silvan Goldenberg, GEOMAR ©).**

Mesocosms were set-up on September 10<sup>th</sup>, 2021 (day t0). Phase 0 (t1-t3) was established at the beginning of the experiment, prior to the addition of the sodium carbonate salts ( $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$ ), to analyze the conditions of the mesocosms before their alkalization. On day 4 (t4) of the experiment, 40 L of natural seawater with different concentrations of carbonate salts ( $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$ ) previously homogenized were added to each mesocosm to achieve an alkalinity gradient between  $\Delta 0$  (lowest)  $\Delta 2400$  (highest)  $\mu\text{mol L}^{-1}$ . Phase I (t5-t19) began followed by Phase II (t19-t33). Variables were sampled at 1-day intervals. Experimental design representing the alkalinity gradient (OAE) ( $\mu\text{mol L}^{-1}$ ) in the nine mesocosms was: M5 ( $\Delta 0$ ), M1 ( $\Delta 300$ ), M7 ( $\Delta 600$ ), M4 ( $\Delta 900$ ), M9 ( $\Delta 1200$ ), M3 ( $\Delta 1500$ ), M6 ( $\Delta 1800$ ), M2 ( $\Delta 2100$ ), M8 ( $\Delta 2400$ )  $\mu\text{mol L}^{-1}$ .

## 2.2. Sampling strategy

Samples were collected in 1 L acid-cleaned LDPE bottles. Trace metal clean (TMC) samples were subsampled from this bottle within an ISO Class-6 laminar flow hood (Cutter et al., 2017). Unfiltered samples (i.e., TdFe) were first collected. Aliquots were filtered (0.2  $\mu\text{m}$  Sartobran<sup>TM</sup> PES) for sFe, dFe and LFe. Soluble Fe samples were further filtered through acid-cleaned 0.02  $\mu\text{m}$  filters (Whatman<sup>TM</sup> Anotop<sup>TM</sup> 25/0.02) using a peristaltic pump. The TdFe, dFe and sFe samples were acidified with HCl Ultrapure, 2 % v/v to pH  $\sim 1.7$  (Panreac) after sampling and kept in the dark for at least 6-months until analysis. The LFe samples were frozen at  $-20^\circ\text{C}$  until further analysis was performed.

### 2.3. Iron concentrations analyses

The TdFe, dFe and sFe samples were analyzed in duplicates (two analytical peaks; a second duplicate was performed if the standard deviation was >5 %) using flow injection analysis with chemiluminescence detection (FIA) (Obata et al., 1993; Lohan et al., 2006) inside an ISO Class-6 laminar flow hood inside an ISO Class-5 TMC laboratory. Samples were spiked with 0.013 M ultrapure H<sub>2</sub>O<sub>2</sub> (Sigma-Aldrich) 30 min prior to analysis to ensure the complete oxidation of Fe (II) to Fe (III) (Lohan et al., 2006). Each sample was buffered in-line to pH 3.5 with 0.15 M ammonium acetate (Supelco and Sigma-Aldrich, SpA) before Fe(III) was preconcentrated onto the cation exchange resin Toyopearl-AF-Chelate 650 M (Tosohaas) between 60 and 120 s at a flow rate of 1.5 ml min<sup>-1</sup>. Following a rinse step of weak 0.013 M HCl (Honeywell Fluka™, SpA), Fe was eluted from the resin using 0.24 M HCl (Honeywell Fluka™, SpA) and entered the reaction stream where it was mixed with a 0.015 mM luminol solution containing 70 μl L<sup>-1</sup> triethylenetetramine (Sigma-Aldrich), buffered to pH 9.5±0.1 using a 1 M ammonia solution (Supelco, SpA). Iron concentrations were quantified using standard additions (TraceCERT) to low Fe seawater. The limit of detection (three times the standard deviation of the lowest addition) was 0.03±0.02 nmol L<sup>-1</sup> while the precision of three analytical peaks was <2 %. Accuracy was established by repeat quantification of dFe in an inhouse standard, whose concentration was asserted from the repeat measurement of GSC reference samples yielding 1.59±0.03 nmol L<sup>-1</sup>, which agrees with the reported consensus values (1.54±0.12 nmol L<sup>-1</sup>).

### 2.4. Fe-binding ligands

Iron-binding ligand ( $L_{Fe}$ ) concentrations and the conditional stability constants ( $K_{Fe-L}^{cond}$ ) were determined by competitive ligand exchange-adsorptive cathodic stripping voltammetry (CLE-ACSV) with ligand competition against TAC (2-(2-thiazolylazo)-p-cresol; Sigma-Aldrich). The titration procedure used, established by Croot and Johansson (2000), was carried out with a voltammetric system equipped with a hanging mercury drop electrode (VA663 stand Metrohm), an Ag/AgCl reference electrode with a KCl salt bridge (3 M) and a glassy carbon counter electrode.

Three different reagents were required for titration analysis. A 0.01 M stock solution of TAC as complexing agent was prepared in methanol (Sigma-Aldrich). A buffer of 1 M boric acid in 0.3 M ammonia (pH 8.2), the impurities of which were removed from the solution by the addition of 100 μM of MnO<sub>2</sub>, followed by an overnight stirring and filtration through an acid-clean 0.45 μm filter (Campos and van den Berg, 1994). Finally, dFe stock solutions were prepared weekly in suprapure HCl. All reagents were stored in the dark and at 8°C when they were not in use.

In the titration method, 10 mL seawater aliquots were pipetted into 12 PTFE vials to which 100 μL of boric acid (final concentration 10<sup>-2</sup> M) and different concentrations of dFe (from 0 to 20 nM) were added immediately. After 1 h of equilibration, 10 μL of TAC (final concentration 10 μM) was added and allowed to equilibrate overnight. Then, for voltammetric analysis, samples were purged with nitrogen gas for 120 s, followed by the application of a deposition potential of -0.41 V for 120 s. After 10 s of equilibration, a differential pulse scan from -0.43 to -0.65 V was applied, with a step potential of 5 mV, modulation

amplitude of 20 mV, interval time of 0.091 s and modulation time of 0.051 s. Finally, the data processing was performed using  
150 PromCC software (Omanović et al., 2015)


### 3. Results and discussion

The experiment was divided into three distinct phases according to characterized variations in the seawater chemistry and the plankton community development, as indicated above. A phase-0 was established at the beginning of the experiment (t1-t3) to analyze the conditions in the mesocosms prior to treatment additions (OAE). Phase-I (t5-t19) began after alkalization by  
155  $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$  and then phase-II followed (t19-t33).

The experimental setup effectively recreated a scenario of ocean liming by establishing a gradient of total alkalinity (TA) increasing in  $300 \mu\text{mol L}^{-1}$  intervals. Dissolved inorganic carbon (DIC) and TA were steady until day t21 (Fig. S1-supplemental material). On that day, in the highest treatment ( $\Delta 2400 \mu\text{mol L}^{-1}$ ), abiotic precipitation occurred but, this event did not have any impact on the phytoplankton community response. A succinct summary of the phytoplankton abundance results is provided  
160 next and in Fig. S2-supplemental material. During Phase-I, there was a significant increase in picoeukaryotes abundances at  $\Delta 600$ ,  $\Delta 900$ ,  $\Delta 1800$ , and  $\Delta 2100 \mu\text{mol L}^{-1}$ . Additionally, microplankton increased at  $\Delta 900 \mu\text{mol L}^{-1}$ . In contrast, small nanoeukaryotes ( $2\text{-}20 \mu\text{m}$ ) exhibited a decreasing trend from the experiment's onset, with maximal abundance observed before OAE treatment additions (t1-t3). Moving to Phase-II, large nanoeukaryotes ( $> 20 \mu\text{m}$ ) dominated the community, particularly in  $\Delta 1500$  and  $\Delta 1800 \mu\text{mol L}^{-1}$ . Microplankton also increased by the phase's end. *Synechococcus spp.* showed a general increase  
165 in cell density over time, peaking at t27 in  $\Delta 900 \mu\text{mol L}^{-1}$ , but subsequently decreasing in many other treatments. All this data can be consulted for further details and in the articles by Marin-Samper et al. and Ramirez et al., this issue.

#### 3.1. Iron size fractionation

Iron size fractionation (sFe, cFe, dFe and TdFe) was determined during the OAE experiment (Fig. 2). The first metal concentration samples were collected on t1 for dFe and sFe, i.e. phase-0 of the experiment. The initial concentration for dFe  
170 was  $1.74 \pm 0.44 \text{ nmol L}^{-1}$ , ranging from  $1.20 \pm 0.02 \text{ nmol L}^{-1}$  at  $\Delta 1800$  to  $2.39 \pm 0.03 \text{ nmol L}^{-1}$  at  $\Delta 1500$ . For sFe, the initial concentrations were  $0.39 \pm 0.18 \text{ nmol L}^{-1}$ , ranging from  $0.22 \pm 0.01 \text{ nmol L}^{-1}$  at  $\Delta 600$  to  $0.74 \pm 0.03 \text{ nmol L}^{-1}$  at  $\Delta 1800$ .



175 **Figure 2. Temporal evolution of a) total dissolvable iron (TdFe), b) dissolved iron (dFe), c) colloidal iron (cFe = dFe-sFe) and d) soluble iron (sFe) concentrations from day 0 to day 33 of the mesocosm experiment for the nine increased alkalinity mesocosms ranging from 0 (light blue) to 2400  $\mu\text{mol L}^{-1}$  (dark blue). Symbols and color-codes as in figure 1C. The concentration range of the different size fractions is different. Three y-axis ranges have been used; up to 10  $\text{nmol L}^{-1}$  for TdFe, up to 5  $\text{nmol L}^{-1}$  for dFe and cFe, and up to 1  $\text{nmol L}^{-1}$  for sFe.**

180 In the Macaronesia region, dFe concentrations average  $0.80 \pm 0.25 \text{ nmol L}^{-1}$  (varying from 0.46 to  $1.32 \text{ nmol L}^{-1}$ ) (Arnone et al., 2022). However, these concentrations are obtained from samples collected on a research vessel using trace metal clean conditions into a trace metal clean bottle following strict trace metal clean conditions (Cutter et al., 2017). As such, the dFe concentrations just show a slight increase in dFe when no trace metal clean procedures were followed and mesocosms were kept in the open air, but still trying to limit contamination. The observed changes in dFe and sFe concentrations were most likely due to the natural variability between mesocosms. Considering that variability occurs on interannual, seasonal and diurnal time scales, this also forms part of the system's behavior (Schulz and Riebesell, 2013). Indeed, considerable natural variability exists in seawater carbonate chemistry speciation, caused by changes in temperature and biological activities such as photosynthesis, respiration, nutrient utilization, remineralization and calcium carbonate precipitation and dissolution

185



(Segovia et al., 2017). These processes were highly active during our experiment, especially during phase-II (Paul et al; Marin-Samper et al; this issue) thus probably directly affecting metal speciation.

190 The mesocosm with  $\Delta 2100$  was not included in the calculated means due to an outlier in both  $d\text{Fe} = 5.30 \pm 0.10 \text{ nmol L}^{-1}$  and  $s\text{Fe} = 2.86 \pm 0.05 \text{ nmol L}^{-1}$ . The first TdFe samples were taken on day t3 and averaged  $6.26 \pm 2.4 \text{ nmol L}^{-1}$ . The high standard deviation in the results was caused by  $\Delta 300$  and  $\Delta 2100$ . When these two mesocosms were excluded, the average concentration decreased to  $5.21 \pm 0.81 \text{ nmol L}^{-1}$ .

195 On day t4, sodium carbonate salts,  $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$ , were added to the mesocosms and phase-II started. Mesocosms in which  $\Delta\text{TA}$  was greater than  $300 \mu\text{mol L}^{-1}$  showed decreases in dFe. The resulting dFe concentrations were  $0.63 \pm 0.06 \text{ nmol L}^{-1}$  for all the mesocosms except for  $\Delta\text{TA} 0$  and  $300 \mu\text{mol L}^{-1}$  ( $2.56 \pm 0.02$  and  $1.39 \pm 0.08 \text{ nmol L}^{-1}$ , respectively). These concentrations were the lowest observed throughout all the experiment and showed the least variation between mesocosms at any timepoint. The decrease in dFe was most likely caused by a decrease in colloidal sized Fe (cFe). Aggregation processes did not affect the sFe, which remained relatively unchanged. Mesocosms 5 and 1 ( $\Delta\text{TA} 0$  and  $300 \mu\text{mol L}^{-1}$  respectively) showed higher concentrations but were within the analysis range observed during the mesocosm experiments.

200 Two days after the alkalization event, dFe concentrations returned to the original concentration range ( $2.00 \pm 0.39 \text{ nmol L}^{-1}$ ). This change in concentration back to normal did not correlate with the addition of  $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$  required for the OAE. The increase in dFe was not accompanied by an increase in sFe along phase-I. It was the cFe fraction that varied. This suggests that aggregation processes of colloidal particles occurred with other particles with limited adsorption/absorption of sFe. TdFe also increased after the addition of  $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$ . This increase could be due to multiple reasons including 1) the opening of the mesocosms, which could have allowed for aerial input of iron, 2) the addition of Fe during the addition of  $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$  including possible trace amounts in the used reagents, 3) the lack of trace metal clean conditions, 4) the presence of mesozooplankton faecal pellets, and 5) the aggregation of cFe and dFe could have led to the formation of larger size particles ( $>0.2 \text{ mm}$ ) that could have been dissolved during the 6 months of sample acidification, compared to highly refractory particles that may not solubilize in 2% v/v HCl.

210 This mesocosm experiment was carried out in coastal waters where Fe is not a limiting micronutrient (López-García et al., 2021; Arnone et al., 2022). However, Fe redox processes should be considered in future OAE where iron can become a limiting element. For example, Fe(II) is considered the most bioavailable Fe species as it is less energetically costly to uptake by organisms. However, in the current oxic ocean, Fe(II) is rapidly oxidized to Fe(III) (Santana-Casiano et al., 2005). The addition of  $\text{NaHCO}_3$  and  $\text{Na}_2\text{CO}_3$  reduced the cFe concentration. Recent studies of Fe(II) oxidation kinetics have shown that colloidal size particles play a significant role in Fe(II) oxidation kinetics (González-Santana et al., 2021). Indeed, a decrease in colloids drastically accelerates the Fe(II) oxidation rate which subsequently, could lead to a decrease in available Fe(II), reducing Fe bioavailability and Fe acquisition costs in terms of energy (microbial/enzymatic catalysis) (Fujii et al., 2010; Anderson and Morel, 1982).


220 Correlations between the different iron size fractions and their ratios, mesocosm physico-chemical parameters (i.e., pH, alkalinity, dissolved inorganic carbon, nitrates, nitrites, silicates) and biological variables (i.e., fluorescence, community

respiration, net primary production, pico and nanoeukaryotes concentration) did not present any statistical significance (Fig. S3). Initial statistical tests showed a significant inverse correlation ( $p < 0.05$  and  $r > -0.5$ ) between TdFe and DIC and TA, however, when further tests were performed for phase-II, correlation decreased ( $p > 0.28$  and  $r = -0.15$ ). These results show that iron concentrations were not limiting during the experiments and so any of the phytoplankton functional groups were below their Fe demand for growth (Segovia et al. 2017). This is a direct result of high iron concentration as a consequence of the study site being located in coastal waters with high Fe inputs such as sediment resuspension in coastal waters and aerial inputs due to the Canary Islands location which would increase initial Fe concentrations used in the mesocosms compared to open ocean locations. Yet, such Fe concentrations were not toxic either, since phytoplankton growth was not hampered in phase-II. In fact, the contrary occurred, and Chlorophyll-a (Chla) levels increased in parallel with a significant recovery in the Fv/Fm ratio (Ramirez et al. this issue). These improvements in Chla and photophysiology data are primarily attributed to the emergence of blooms of large nanoeukaryotes and picoeukaryotes in  $\Delta 1500$ ,  $\Delta 1800$ ,  $\Delta 600$  and  $\Delta 900$  at t27.

In light of this, we suggest that iron size fractions and phytoplankton are independent. However, we have more than a reasonable doubt about the colloidal fraction being directly linked with mesozooplankton fecal pellets. For instance,  $>200\%$  of a mixture of lithogenic and biogenic Fe seems to be egested via Antarctic mesozooplankton fecal pellets, while only 5% of the body iron content is assimilated (Schmidt et al., 2016; Cabanes et al., 2017). In any case, more work is needed to clarify this particular point.

### 3.2. Iron ligand distribution

Considering that the OAE were not carried out under Trace Metal Clean conditions, according with the GEOTRACES protocol (Cutter et al., 2017), except for both sample collection and measurement in the ULPGC facilities, the OAE experiments have demonstrated non-linearity in the behavior of Fe-binding ligands ( $L_{Fe}$ ) and the conditional stability constant ( $\log K_{Fe \cdot L}^{cond}$ ) across the entire alkalinity gradient (Fig. 3). Fe-binding ligands in the ocean come from different sources such as rupture of cells after grazing (Sato et al., 2007), viral lysis (Poore et al., 2011), transformation of organic matter (Gerringa et al., 2006), phytoplankton exudates (Rico et al., 2013; Santana-Casiano et al., 2014) and sediments (Gerringa et al., 2008). The  $L_{Fe}$  are classified according to the value of the conditional stability constants ( $\log K_{Fe \cdot L}^{cond}$ ). These ligands can be considered to be strong ( $L_1$ ) or weak ( $L_2$ ), where  $\log K_{Fe \cdot L}^{cond} = 12$  divides  $L_1$  and  $L_2$ -type of Fe-binding ligands (Gledhill and Buck, 2012).



**Figure 3. Temporal evolution of a) iron ligands ( $L_{Fe}$ ), and b) Conditional constant ( $K_{cond}$ ), from day 0 to day 33 of the mesocosm experiment for the nine increased alkalinity mesocosms ranging from 0 (light blue) to 2400  $\mu\text{mol L}^{-1}$  (dark blue). Symbols and color-codes as in figure 1C.**

250 Specifically, before OAE (phase-0),  $L_{Fe}$  concentrations ranged from 6.81  $\text{nmol L}^{-1}$  ( $\Delta 300$ ) to 1.51  $\text{nmol L}^{-1}$  ( $\Delta 1800$ ). Throughout the course of the experiments, the mean  $L_{Fe}$  concentration across all mesocosms was  $4.14 \pm 2.01 \text{ nmol L}^{-1}$ , with a  $\log K'_{cond,FeL}$  value of  $11.63 \pm 0.36$ , being  $L_2$ -class in most of the cases (90.5% of the 84 data). Notably, there was a discernible alteration in the behavior of  $L_{Fe}$  in the  $\Delta 1500$  mesocosm, coincident with the emergence of blooms of the Haptophyte *Chrysochromulina tobinni* at t27 (Marin-Samper et al., this issue).  $L_{Fe}$  consistently exhibited higher concentrations than  $dFe$ , with the exception of mesocosm  $\Delta 2100$  at t1,  $\Delta 900$  at t5,  $\Delta 600$  at t5 and t27,  $\Delta 2400$  at t5, and  $\Delta 1200$  at t3 and t5. In these instances, the titration curves were linear, suggesting oversaturation of the natural ligands with the Fe content. Consequently, the percentage of Fe-organically complexed (%FeL) was determined to be  $99.8 \pm 0.2$ . The concentration of labile Fe ( $Fe'$ ) was in the range from 0.04 to 15.98  $\text{pmol L}^{-1}$ , and  $Fe^{3+}$  species, primarily  $Fe(OH)_3$ , was calculated to be within the concentration range of  $1.6 \times 10^{-21}$  to  $3.9 \times 10^{-26} \text{ mol L}^{-1}$ .

260 It is important to remark that the experiment was not conducted under trace metal clean conditions. Consequently, discussing the potential Fe limitation in relation to the measured concentrations becomes a complex task. Additionally, the possibility of Fe contamination during the sampling process for other variables, which may influence Fe concentration and Fe-binding ligands, cannot be definitively ruled out.

The initial concentrations of  $dFe$  and  $L_{Fe}$  at the start of the experiments were found to be higher than those observed in coastal waters previously documented by Arnone et al. (2022) in the Canary Islands. Specifically, the mean  $dFe$  concentration in the region was determined to be  $0.80 \pm 0.16 \text{ nmol L}^{-1}$ , with corresponding mean  $L_{Fe}$  levels of  $1.28 \pm 0.21 \text{ nmol L}^{-1}$ . The initial  $L_{Fe}$  concentrations within the OAE mesocosms ranged from 1.51 to 6.81  $\text{nmol L}^{-1}$ , and their evolution over time fell within the range of 1.51 to 12.30  $\text{nmol L}^{-1}$ . These values were also in line with data from surface waters of the Atlantic Ocean, ranging

from 0.1 to 0.8 nmol L<sup>-1</sup> (Boye et al., 2006; Rijkenberg et al., 2008). The pumping of more coastal water to fill the mesocosms  
270 could explain the observed increase in concentrations.

Measurement of L<sub>Fe</sub> within the mesocosms indicated a consistent nature of Fe-binding ligands, as reflected by the comparable  
values of the log K<sup>cond</sup><sub>Fe·L</sub> throughout the entire study period, within the range of 10.91 – 12.68, corresponding to the L<sub>2</sub> and  
L<sub>1</sub>-class, being weaker ligands (L<sub>2</sub>) more abundant than stronger ligands (L<sub>1</sub>), representing 90.5% of the samples. In the  
surrounding waters of the Canary Islands, log K<sup>cond</sup><sub>Fe·L</sub> values were reported to range between 10.77 and 11.90 (with a mean  
275 value of 11.45±0.29) (Arnone et al., 2022), aligning with surface ocean waters in the region (log K<sup>cond</sup><sub>Fe·L</sub> 8.8 – 12.85) (Boye  
et al., 2006; Bundy et al., 2015; Rijkenberg et al., 2008; Thuróczy et al., 2010; Boye et al., 2003).

The ratio of L<sub>Fe</sub> to dFe serves as an indicator of the saturation state of ligands and whether conditions favor precipitation  
(Thuróczy et al., 2011). The observation of an excess of L<sub>Fe</sub> with values exceeding 1 nmol L<sup>-1</sup> may suggest biological  
production of ligands. Thuróczy et al. (2011) previously reported a ratio of approximately 4.4 for surface waters characterized  
280 by high phytoplankton activity. In the case of our OAE experiment, the production of Fe-binding ligands may be attributed to  
the increase in large nanoplankton and microplankton biomass as described by Ramírez et al. (this issue) during phase-II,  
which was specially observable in the Δ1500 treatment. Additionally, the presence of cyanobacteria, during the OAE can lead  
to the production of siderophores, as indicated by log K<sup>cond</sup><sub>Fe·L</sub> values falling between 11 and 14 (Bundy et al., 2018; Witter et  
al., 2000). Siderophores are organic molecules of microbial origin that bind metals with different affinities. They are thought  
285 to play an important role in iron cycling in the ocean, but relatively few marine siderophores have been identified. Recently,  
new siderophores produced by *Synechococcus sp.* PCC 7002 have been discovered and characterized (Boiteau and Repeta,  
2015) which is of relevance considering that *Synechococcus spp.* peaked during phase-II.

Furthermore, other types of ligands, such as polyphenolic compounds, may also be present, consistent with their log K<sup>cond</sup><sub>Fe·L</sub>  
values (Arreguin et al., 2021; González et al., 2019). Weak Fe-binding ligands (log K<sup>cond</sup><sub>Fe·L</sub> <12) could potentially be produced  
290 during the grazing of phytoplankton and the bacterial remineralization of sinking organic particles (Poorvin et al., 2011; Sato  
et al., 2007).

The observed variability in labile Fe concentrations (Fe<sup>l</sup>) within the range of 0.04 to 15.98 pmol L<sup>-1</sup> aligns with the findings of  
Arnone et al. (2022). Notably, the limited concentration of Fe<sup>3+</sup> precludes the favored precipitation of iron hydroxides. It has  
been established that the presence of organic binding ligands enhances the solubility of Fe, as demonstrated by previous  
295 research (Liu and Millero, 2002; Segovia et al., 2017).

Similar to iron size fraction concentrations, L<sub>Fe</sub> and K<sup>cond</sup><sub>Fe·L</sub> did not present any statistical correlation with any other measured  
physico-chemical or biological parameter. This suggests that the alkalinity addition in the mesocosm experiments did not affect  
the Fe concentration, fractionation, and speciation.

This OAE experiment was based on the addition of sodium carbonate salts. However, one might wonder which are the pros  
300 and cons of this method in comparison to other weathering-based Carbon Dioxide Removal (CDR) techniques. For instance,  
similar alkalinity concentrations could have been obtained by adding calcium or magnesium carbonate salts. However, a  
previous study showed that the addition of calcium or magnesium salts would have led to changes in the iron speciation

(Santana-Casiano et al., 2010). Also open ocean dissolution of the iron-containing mineral olivine was simulated by using a marine carbon cycle model (Hauck et al. (2016). They found that after equilibration with the atmosphere, the addition of olivine did not enough compensate for the effect of ocean acidification and, in the real world experiments, olivine-prompted silicic acid and iron fertilization might have undesired effects since increased dFe does not always have beneficial consequences for the community (Coale, 1991; Behrenfeld and Milligan, 2013).

#### 4. Conclusion

The OAE experiments, while not conducted under stringent trace metal conditions according to the GEOTRACES protocol, have provided valuable insights into the non-linear behavior of Fe-binding ligands ( $L_{Fe}$ ) and the conditional stability constant ( $\log K_{Fe-L}^{cond}$ ) across the entire alkalinity gradient.  $L_{Fe}$  concentrations increased 2-fold just after OAE with a  $\log K_{Fe-L}^{cond}$  throughout within the range of 10.91 – 12.68. This suggests the possible biological production of ligands, particularly during the emergence of nanoeukaryotes blooms and peaks of *Synechococcus* in the  $\Delta 1500$  treatment, as well as zooplankton grazing (Smith et al, this issue). Similarly, dFe concentrations were higher than those one might expect in the Macaronesia region. This could have been a result of the manipulation required to fill in mesocosms in the open environment, where the iron sources such as sediment resuspension, aerial input and water runoffs are combined with our mesocosm manipulation. During the OAE we did not observe any iron size fractionation changes and concentration variations due to different OAE conditions. Nevertheless, the addition of sodium carbonate salts ( $NaHCO_3$  and  $Na_2CO_3$ ) coincided with a decreases in the dFe concentration just after the addition. However, concentrations returned to background level concentrations in 48 h. Soluble iron remained below  $0.8 \text{ nmol L}^{-1}$  through the whole experiment and did not change after the sodium carbonate salts addition. The results suggest that the added sodium carbonate salts may affect cFe, but quickly return to the original concentrations in coastal waters. Nevertheless, the observable decrease is not proportional to the increase in alkalinity. Where other factors such as aggregation due to increases in particles or added mineral salts produce a short term cFe decrease.

The iron size fractionation, concentration and iron-binding ligands data supports the fact that the addition of sodium salts in this mesocosm experiment did not lead to significant changes in the iron cycle, therefore phytoplankton were not affected by changes in this essential element. This may have wider environmental implications in large-scale experiments and implementation of CDRs in larger ocean. However, it's important to consider that while employing this Oceanic Iron Fertilization (OAE) technique in coastal waters might not significantly impact the iron cycle, outcomes could differ under varying conditions. For instance, if the experiment were conducted in oligotrophic open ocean waters, the presence of a bloom exceeding those observed in these experiments could lead to different results. Conversely, an excess of iron might inhibit phytoplankton growth due to potential toxic effects. Therefore, further research into the effects of Carbon Dioxide Removal (CDR) techniques on trace metals is essential to mitigate unintended consequences on the ecosystem.

## References

- 335 Anderson, M. A. and Morel, F. M. M.: The influence of aqueous iron chemistry on the uptake of iron by the coastal diatom *Thalassiosira weissflogii* 1, *Limnol. Oceanogr.*, 27, 789–813, <https://doi.org/https://doi.org/10.4319/lo.1982.27.5.0789>, 1982.
- Arnone, V., González-Santana, D., González-Dávila, M., González, A. G., and Santana-Casiano, J. M.: Iron and copper complexation in Macaronesian coastal waters, *Mar. Chem.*, 240, 104087, <https://doi.org/10.1016/J.MARCHEM.2022.104087>, 2022.
- 340 Arreguin, M. L., González, A. G., Pérez-Almeida, N., Arnone, V., González-Dávila, M., and Santana-Casiano, J. M.: The role of gentisic acid on the Fe(III) redox chemistry in marine environments, *Mar. Chem.*, 234, 104003, <https://doi.org/10.1016/J.MARCHEM.2021.104003>, 2021.
- Bach, L. T., Gill, S. J., Rickaby, R. E. M., Gore, S., and Renforth, P.: CO<sub>2</sub> Removal With Enhanced Weathering and Ocean Alkalinity Enhancement: Potential Risks and Co-benefits for Marine Pelagic Ecosystems, *Front. Clim.*, 1, 476698, <https://doi.org/10.3389/FCLIM.2019.00007/BIBTEX>, 2019.
- 345 Barbeau, K.: Photochemistry of Organic Iron(III) Complexing Ligands in Oceanic Systems, *Photochem. Photobiol.*, 82, 1505–1516, <https://doi.org/https://doi.org/10.1111/j.1751-1097.2006.tb09806.x>, 2006.
- Behrenfeld, M. J. and Milligan, A. J.: Photophysiological Expressions of Iron Stress in Phytoplankton, *Ann. Rev. Mar. Sci.*, 5, 217–246, <https://doi.org/10.1146/annurev-marine-121211-172356>, 2013.
- 350 Benner, R.: Loose ligands and available iron in the ocean, *Proc. Natl. Acad. Sci.*, 108, 893–894, <https://doi.org/10.1073/pnas.1018163108>, 2011.
- Boiteau, R. M. and Repeta, D. J.: An extended siderophore suite from *Synechococcus* sp. PCC 7002 revealed by LC-ICPMS-ESIMS†, *Metallomics*, 7, 877–884, <https://doi.org/10.1039/c5mt00005j>, 2015.
- Boyd, P. W. and Ellwood, M. J.: The biogeochemical cycle of iron in the ocean, *Nat. Geosci.*, 3, 675–682, <https://doi.org/10.1038/ngeo964>, 2010.
- 355 Boye, M., Aldrich, A. P., van den Berg, C. M. G., de Jong, J. T. M., Veldhuis, M., and de Baar, H. J. W.: Horizontal gradient of the chemical speciation of iron in surface waters of the northeast Atlantic Ocean, *Mar. Chem.*, 80, 129–143, [https://doi.org/https://doi.org/10.1016/S0304-4203\(02\)00102-0](https://doi.org/https://doi.org/10.1016/S0304-4203(02)00102-0), 2003.
- Boye, M., Aldrich, A., van den Berg, C. M. G., de Jong, J. T. M., Nirmaier, H., Veldhuis, M. J. W., Timmermans, K. R., and de Baar, H. J. W.: The chemical speciation of iron in the north-east Atlantic Ocean, *Deep. Res. Part I Oceanogr. Res. Pap.*, 53, 667–683, <https://doi.org/10.1016/j.dsr.2005.12.015>, 2006.
- 360 Bundy, R. M., Abdulla, H. A. N., Hatcher, P. G., Biller, D. V., Buck, K. N., and Barbeau, K. A.: Iron-binding ligands and humic substances in the San Francisco Bay estuary and estuarine-influenced shelf regions of coastal California, *Mar. Chem.*, 173, 183–194, 2015.
- 365 Bundy, R. M., Boiteau, R. M., McLean, C., Turk-Kubo, K. A., McIlvin, M. R., Saito, M. A., Van Mooy, B. A. S., and Repeta, D. J.: Distinct Siderophores Contribute to Iron Cycling in the Mesopelagic at Station ALOHA ,

- <https://www.frontiersin.org/articles/10.3389/fmars.2018.00061>, 2018.
- Cabanes, D. J. E., Norman, L., Santos-Echeandía, J., Iversen, M. H., Trimborn, S., Laglera, L. M., and Hassler, C. S.: First evaluation of the role of salp fecal pellets on iron biogeochemistry, *Front. Mar. Sci.*, 3, 289, 2017.
- 370 Campos, M. L. A. and van den Berg, C. M.: Determination of copper complexation in sea water by cathodic stripping voltammetry and ligand competition with salicylaldoxime, *Anal. Chim. Acta*, 284, 481–496, [https://doi.org/10.1016/0003-2670\(94\)85055-0](https://doi.org/10.1016/0003-2670(94)85055-0), 1994.
- Coale, K. H.: Effects of iron, manganese, copper, and zinc enrichments on productivity and biomass in the subarctic Pacific, *Limnol. Oceanogr.*, 36, 1851–1864, <https://doi.org/https://doi.org/10.4319/lo.1991.36.8.1851>, 1991.
- 375 Colbourn, G., Ridgwell, A., and Lenton, T. M.: The time scale of the silicate weathering negative feedback on atmospheric CO<sub>2</sub>, *Global Biogeochem. Cycles*, 29, 583–596, <https://doi.org/10.1002/2014GB005054>, 2015.
- Croot, P. L. and Johansson, M.: Determination of Iron Speciation by Cathodic Stripping Voltammetry in Seawater Using the Competing Ligand 2-(2-Thiazolylazo)-p-cresol (TAC), *Electroanalysis*, 12, 565–576, [https://doi.org/https://doi.org/10.1002/\(SICI\)1521-4109\(200005\)12:8<565::AID-ELAN565>3.0.CO;2-L](https://doi.org/https://doi.org/10.1002/(SICI)1521-4109(200005)12:8<565::AID-ELAN565>3.0.CO;2-L), 2000.
- 380 Cutter, G. A., Casciotti, K., Croot, P., Geibert, W., Heimbürger, L.-E., Lohan, M. C., Planquette, H., and van de Flierdt, T.: Sampling and sample-handling protocols for GEOTRACES Cruises, Version 3.0, 2017.
- Feng, E. Y., Keller, D. P., Koeve, W., and Oschlies, A.: Could artificial ocean alkalization protect tropical coral ecosystems from ocean acidification?, *Environ. Res. Lett.*, 11, 74008, <https://doi.org/10.1088/1748-9326/11/7/074008>, 2016.
- Fujii, M., Rose, A. L., Omura, T., and Waite, T. D.: Effect of Fe(II) and Fe(III) Transformation Kinetics on Iron Acquisition by a Toxic Strain of *Microcystis aeruginosa*, *Environ. Sci. Technol.*, 44, 1980–1986, <https://doi.org/10.1021/es901315a>, 2010.
- 385 Gerringa, L. J. A., Veldhuis, M. J. W., Timmermans, K. R., Sarthou, G., and de Baar, H. J. W.: Co-variance of dissolved Fe-binding ligands with phytoplankton characteristics in the Canary Basin, *Mar. Chem.*, 102, 276–290, <https://doi.org/https://doi.org/10.1016/j.marchem.2006.05.004>, 2006.
- Gerringa, L. J. A., Blain, S., Laan, P., Sarthou, G., Veldhuis, M. J. W., Brussaard, C. P. D., Viollier, E., and Timmermans, K. R.: Fe-binding dissolved organic ligands near the Kerguelen Archipelago in the Southern Ocean (Indian sector), *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 55, 606–621, <https://doi.org/https://doi.org/10.1016/j.dsr2.2007.12.007>, 2008.
- Gledhill, M. and Buck, K. N.: *The Organic Complexation of Iron in the Marine Environment: A Review*, 2012.
- González-Dávila, M., Santana-Casiano, J. M., and Millero, F. J.: Competition between O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> in the oxidation of Fe(II) in natural waters, *J. Solution Chem.*, 35, 95–111, <https://doi.org/10.1007/s10953-006-8942-3>, 2006.
- 395 González-Santana, D., González-Dávila, M., Lohan, M. C., Artigue, L., Planquette, H., Sarthou, G., Tagliabue, A., and Santana-Casiano, J. M.: Variability in iron (II) oxidation kinetics across diverse hydrothermal sites on the northern Mid Atlantic Ridge, *Geochim. Cosmochim. Acta*, 297, 143–157, <https://doi.org/10.1016/j.gca.2021.01.013>, 2021.
- González, A. G., Cadena-Aizaga, M. I., Sarthou, G., González-Dávila, M., and Santana-Casiano, J. M.: Iron complexation by phenolic ligands in seawater, *Chem. Geol.*, 511, <https://doi.org/10.1016/j.chemgeo.2018.10.017>, 2019.
- 400 Guo, J. A., Strzepek, R., Willis, A., Ferderer, A., and Bach, L. T.: Investigating the effect of nickel concentration on

- phytoplankton growth to assess potential side-effects of ocean alkalinity enhancement, *Biogeosciences*, 19, 3683–3697, <https://doi.org/10.5194/bg-19-3683-2022>, 2022.
- Hartmann, J., West, A. J., Renforth, P., Köhler, P., De La Rocha, C. L., Wolf-Gladrow, D. A., Dürr, H. H., and Scheffran, J.: Enhanced chemical weathering as a geoengineering strategy to reduce atmospheric carbon dioxide, supply nutrients, and mitigate ocean acidification, *Rev. Geophys.*, 51, 113–149, <https://doi.org/10.1002/rog.20004>, 2013.
- 405 Hauck, J., Köhler, P., Wolf-Gladrow, D., and Völker, C.: Iron fertilisation and century-scale effects of open ocean dissolution of olivine in a simulated CO<sub>2</sub> removal experiment, *Environ. Res. Lett.*, 11, 24007, <https://doi.org/10.1088/1748-9326/11/2/024007>, 2016.
- Hopkinson, B. M. and Barbeau, K. A.: Organic and redox speciation of iron in the eastern tropical North Pacific suboxic zone, *Mar. Chem.*, 106, 2–17, <https://doi.org/10.1016/j.marchem.2006.02.008>, 2007.
- 410 Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, C., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., LaRoche, J., Liss, P. S., Mahowald, N. M., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean biogeochemistry, and climate, *Science* (80-. ), 308, 67–71, <https://doi.org/10.1126/science.1105959>, 2005.
- 415 Kheshgi, H. S.: Sequestering atmospheric carbon dioxide by increasing ocean alkalinity, *Energy*, 20, 915–922, [https://doi.org/10.1016/0360-5442\(95\)00035-F](https://doi.org/10.1016/0360-5442(95)00035-F), 1995.
- Lenton, A., Matear, R. J., Keller, D. P., Scott, V., and Vaughan, N. E.: Assessing carbon dioxide removal through global and regional ocean alkalization under high and low emission pathways, *Earth Syst. Dynam.*, 9, 339–357, <https://doi.org/10.5194/esd-9-339-2018>, 2018.
- 420 Liu, X. and Millero, F. J.: The solubility of iron in seawater, *Mar. Chem.*, 77, 43–54, [https://doi.org/10.1016/S0304-4203\(01\)00074-3](https://doi.org/10.1016/S0304-4203(01)00074-3), 2002.
- Lohan, M. C., Aguilar-Islas, A. M., and Bruland, K. W.: Direct determination of iron in acidified (pH 1.7) seawater samples by flow injection analysis with catalytic spectrophotometric detection: Application and intercomparison, *Limnol. Oceanogr. Methods*, 4, 164–171, <https://doi.org/10.4319/lom.2006.4.164>, 2006.
- 425 López-García, P., Gelado-Caballero, M. D., Patey, M. D., and Hernández-Brito, J. J.: Atmospheric fluxes of soluble nutrients and Fe: More than three years of wet and dry deposition measurements at Gran Canaria (Canary Islands), *Atmos. Environ.*, 246, 118090, <https://doi.org/10.1016/j.atmosenv.2020.118090>, 2021.
- Lorenzo, M. R., Segovia, M., Cullen, J. T., and Maldonado, M. T.: Particulate trace metal dynamics in response to increased CO<sub>2</sub> and iron availability in a coastal mesocosm experiment, *Biogeosciences*, 17, 757–770, [https://doi.org/10.5194/bg-17-](https://doi.org/10.5194/bg-17-757-2020)
- 430 757-2020, 2020.
- Mausz, M. A., Segovia, M., Larsen, A., Berger, S. A., Egge, J. K., and Pohnert, G.: High CO<sub>2</sub> concentration and iron availability determine the metabolic inventory in an *Emiliania huxleyi*-dominated phytoplankton community, *Environ. Microbiol.*, 22, 3863–3882, <https://doi.org/10.1111/1462-2920.15160>, 2020.
- Millero, F. J., Woosley, R., Ditrolio, B., and Waters, J.: Effect of Ocean Acidification on the Speciation of Metals in Seawater,



- 435 Oceanography, 22, 72–85, 2009.
- Moffett, J. W.: Iron(II) in the world's oxygen deficient zones, Chem. Geol., 120314, <https://doi.org/10.1016/j.chemgeo.2021.120314>, 2021.
- Moore, J. K., Doney, S. C., Glover, D. M., and Fung, I. Y.: Iron cycling and nutrient-limitation patterns in surface waters of the World Ocean, Deep Sea Res. Part II Top. Stud. Oceanogr., 49, 463–507, [https://doi.org/10.1016/S0967-0645\(01\)00109-6](https://doi.org/10.1016/S0967-0645(01)00109-6),  
440 2001.
- Obata, H., Karatani, H., and Nakayama, E.: Automated Determination of Iron in Seawater by Chelating Resin Concentration and Chemiluminescence Detection, Anal. Chem., 65, 1524–1528, <https://doi.org/10.1021/ac00059a007>, 1993.
- Omanović, D., Garnier, C., and Pižeta, I.: ProMCC: An all-in-one tool for trace metal complexation studies, Mar. Chem., 173, 25–39, <https://doi.org/https://doi.org/10.1016/j.marchem.2014.10.011>, 2015.
- 445 Poorvin, L., Sander, S. G., Velasquez, I., Ibisani, E., LeCleir, G. R., and Wilhelm, S. W.: A comparison of Fe bioavailability and binding of a catecholate siderophore with virus-mediated lysates from the marine bacterium *Vibrio alginolyticus* PWH3a, J. Exp. Mar. Bio. Ecol., 399, 43–47, <https://doi.org/https://doi.org/10.1016/j.jembe.2011.01.016>, 2011.
- Renforth, P. and Henderson, G.: Assessing ocean alkalinity for carbon sequestration, Rev. Geophys., 55, 636–674, <https://doi.org/https://doi.org/10.1002/2016RG000533>, 2017.
- 450 Rico, M., López, A., Santana-Casiano, J. M., González, A. G., and González-Dávila, M.: Variability of the phenolic profile in the diatom *Phaeodactylum tricornutum* growing under copper and iron stress, Limnol. Oceanogr., 58, 144–152, <https://doi.org/10.4319/LO.2013.58.1.0144>, 2013.
- Riebesell, U., Czerny, J., von Bröckel, K., Boxhammer, T., Büdenbender, J., Deckelnick, M., Fischer, M., Hoffmann, D., Krug, S. A., Lentz, U., Ludwig, A., Muehe, R., and Schulz, K. G.: Technical Note: A mobile sea-going mesocosm system – new  
455 opportunities for ocean change research, Biogeosciences, 10, 1835–1847, <https://doi.org/10.5194/bg-10-1835-2013>, 2013.
- Rijkenberg, M. J. A., Powell, C. F., Dall'Osto, M., Nielsdottir, M. C., Patey, M. D., Hill, P. G., Baker, A. R., Jickells, T. D., Harrison, R. M., and Achterberg, E. P.: Changes in iron speciation following a Saharan dust event in the tropical North Atlantic Ocean, Mar. Chem., 110, 56–67, <https://doi.org/https://doi.org/10.1016/j.marchem.2008.02.006>, 2008.
- Santana-Casiano, J. M., González-Dávila, M., and Millero, F. J.: Oxidation of nanomolar level of Fe(II) with oxygen in natural  
460 waters, Environ. Sci. Technol., 39, 2073–2079, <https://doi.org/10.1021/es049748y>, 2005.
- Santana-Casiano, J. M., González-Dávila, M., and Millero, F. J.: The role of Fe(II) species on the oxidation of Fe(II) in natural waters in the presence of O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub>, Mar. Chem., 99, 70–82, <https://doi.org/10.1016/j.marchem.2005.03.010>, 2006.
- Santana-Casiano, J. M., González-Dávila, M., González, A. G., and Millero, F. J.: Fe (III) reduction in the presence of catechol in seawater, Aquat. geochemistry, 16, 467–482, 2010.
- 465 Santana-Casiano, J. M., González-Dávila, M., González, A. G., Rico, M., López, A., and Martel, A.: Characterization of phenolic exudates from *Phaeodactylum tricornutum* and their effects on the chemistry of Fe (II)–Fe (III), Mar. Chem., 158, 10–16, 2014.
- Santana-Casiano, J. M., González-Santana, D., Devresse, Q., Hepach, H., Santana-González, C., Quack, B., Engel, A., and

- González-Dávila, M.: Exploring the Effects of Organic Matter Characteristics on Fe(II) Oxidation Kinetics in Coastal Seawater, *Environ. Sci. Technol.*, 56, 2718–2728, <https://doi.org/10.1021/acs.est.1c04512>, 2022.
- 470 Sato, M., Takeda, S., and Furuya, K.: Iron regeneration and organic iron(III)-binding ligand production during in situ zooplankton grazing experiment, *Mar. Chem.*, 106, 471–488, <https://doi.org/https://doi.org/10.1016/j.marchem.2007.05.001>, 2007.
- Schmidt, K., Schlosser, C., Atkinson, A., Fielding, S., Venables, H. J., Waluda, C. M., and Achterberg, E. P.: Zooplankton gut 475 passage mobilizes lithogenic iron for ocean productivity, *Curr. Biol.*, 26, 2667–2673, 2016.
- Schulz, K. G. and Riebesell, U.: Diurnal changes in seawater carbonate chemistry speciation at increasing atmospheric carbon dioxide, *Mar. Biol.*, 160, 1889–1899, <https://doi.org/10.1007/s00227-012-1965-y>, 2013.
- Segovia, M., Lorenzo, M. R., Maldonado, M. T., Larsen, A., Berger, S. A., Tsagaraki, T. M., Lázaro, F. J., Iñiguez, C., García-Gómez, C., Palma, A., Mausz, M. A., Gordillo, F. J. L., Fernández, J. A., Ray, J. L., and Egge, J. K.: Iron availability modulates 480 the effects of future CO<sub>2</sub> levels within the marine planktonic food web, *Mar. Ecol. Prog. Ser.*, 565, 17–33, 2017.
- Sutak, R., Camadro, J.-M., and Lesuisse, E.: Iron Uptake Mechanisms in Marine Phytoplankton , <https://www.frontiersin.org/articles/10.3389/fmicb.2020.566691>, 2020.
- Tagliabue, A., Bowie, A. R., Boyd, P. W., Buck, K. N., Johnson, K. S., and Saito, M. A.: The integral role of iron in ocean biogeochemistry, <https://doi.org/10.1038/nature21058>, 2017.
- 485 Taucher, J., Bach, L. T., Boxhammer, T., Nauendorf, A., Achterberg, E. P., Algueró-Muñiz, M., Arístegui, J., Czerny, J., Esposito, M., Guan, W., Haunost, M., Horn, H. G., Ludwig, A., Meyer, J., Spisla, C., Sswat, M., Stange, P., and Riebesell, U.: Influence of Ocean Acidification and Deep Water Upwelling on Oligotrophic Plankton Communities in the Subtropical North Atlantic: Insights from an In situ Mesocosm Study , <https://www.frontiersin.org/articles/10.3389/fmars.2017.00085>, 2017.
- Thuróczy, C.-E., Gerringa, L. J. A., Klunder, M. B., Middag, R., Laan, P., Timmermans, K. R., and de Baar, H. J. W.: 490 Speciation of Fe in the Eastern North Atlantic Ocean, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 57, 1444–1453, <https://doi.org/https://doi.org/10.1016/j.dsr.2010.08.004>, 2010.
- Thuróczy, C.-E., Gerringa, L. J. A., Klunder, M. B., Laan, P., and de Baar, H. J. W.: Observation of consistent trends in the organic complexation of dissolved iron in the Atlantic sector of the Southern Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 58, 2695–2706, <https://doi.org/https://doi.org/10.1016/j.dsr2.2011.01.002>, 2011.
- 495 Witter, A. E., Hutchins, D. A., Butler, A., and Luther, G. W.: Determination of conditional stability constants and kinetic constants for strong model Fe-binding ligands in seawater, *Mar. Chem.*, 69, 1–17, [https://doi.org/https://doi.org/10.1016/S0304-4203\(99\)00087-0](https://doi.org/https://doi.org/10.1016/S0304-4203(99)00087-0), 2000.
- Wu, J. and Luther III, G. W.: Size-fractionated iron concentrations in the water column of the western North Atlantic Ocean, *Limnol. Oceanogr.*, 39, 1119–1129, <https://doi.org/10.4319/lo.1994.39.5.1119>, 1994.
- 500 Xin, X., Faucher, G., and Riebesell, U.: Phytoplankton Response to Increased Nickel in the Context of Ocean Alkalinity Enhancement, *Biogeosciences Discuss.*, 2023, 1–15, <https://doi.org/10.5194/bg-2023-130>, 2023.
- Zeebe, R. E.: History of Seawater Carbonate Chemistry, Atmospheric CO<sub>2</sub>, and Ocean Acidification,

<https://doi.org/10.1146/annurev-earth-042711-105521>, 40, 141–165, <https://doi.org/10.1146/ANNUREV-EARTH-042711-105521>, 2012.

505

#### Data availability

The raw data supporting the conclusions of this article will be made available by the authors.

510 Author contributions:

Experimental concept and design: UR and JA (IOCAG)

Execution of the experiment: All authors.

Data analysis: DGS, AGG, MGD, JMSC

Manuscript preparation: DGS.

515 Manuscript revisions: All authors.

#### Financial Support

This study was funded by the OceanNETS project (“Ocean-based Negative Emissions Technologies – analysing the feasibility, risks and co-benefits of ocean-based negative emission technologies for stabilizing the climate”, EU Horizon 2020 Research and Innovation Programme Grant Agreement No.: 869357), and the Helmholtz European Partnering project Ocean-CDR (520 “Ocean-based carbon dioxide removal strategies”, Project No.: PIE-0021) with additional support from the AQUACOSM-plus project (EU H2020-INFRAIA Project No.: 871081, “AQUACOSM-plus: Network of Leading European AQUATIC MesoCOSM Facilities Connecting Rivers, Lakes, Estuaries and Oceans in Europe and beyond”).

525 Competing interests

The authors declare that they do not have any competing interests and that all of them agree to the content of this manuscript.

#### Acknowledgements

We would like to thank the Oceanic Platform of the Canary Islands (PLOCAN) and its staff for the use of their facilities and (530 for their help with the logistics and organisation of this experiment. We are also grateful to Andrea Ludwig, and Jana Meyer (KOSMOS Logistics and Coordination), to Jan Hennke, Michael Krudewig, and Anton Theileis (KOSMOS technical staff), to Michael Sswat, Carsten Spisla, Daniel Brüggemann, Silvan Goldenberg, Joaquin Ortiz, Nicolás Sánchez, and Philipp Suessele (KOSMOS Scientific Diving and Maintenance Team).