

Technical Note: A Technique to Convert NO_2 to NO_2^- with S(IV) and its Application to Measuring Nitrate Photolysis

Aaron Lieberman, Julietta Picco, Murat Onder, and Cort Anastasio

5 Department of Land, Air, and Water Resources, University of California - Davis, Davis, California 95616, United States

Corresponding author: Cort Anastasio, e-mail: canastasio@ucdavis.edu

10 Abstract

Nitrate photolysis is a potentially significant mechanism for “renoxifying” the atmosphere, i.e., converting nitrate into nitrogen oxides (nitrogen dioxide (NO_2) and nitric oxide (NO)) and nitrous acid (HONO). Nitrate photolysis in the environment occurs through two channels, which produce: (1) NO_2 and hydroxyl radical ($\cdot\text{OH}$) and (2) nitrite (NO_2^-) and an oxygen atom ($\text{O}(^3P)$). Although the aqueous quantum yields and photolysis rate constants of both

15 channels have been established, field observations suggest that nitrate photolysis is enhanced in the environment. Laboratory studies investigating these enhancements typically only measure one of the two photo-channels, since measuring both channels generally requires separate analytical methods and instrumentation. However, measuring only one channel makes it difficult to assess whether secondary chemistry is enhancing one channel at the expense of the other, or if there is an overall enhancement of nitrate photochemistry. Here, we show that the addition of

20 S(IV), i.e., bisulfite and sulfite, can convert NO_2 to NO_2^- , allowing measurement of both nitrate photolysis channels with the same equipment. By varying the concentration of S(IV) and exploring method parameters, we determine the experimental conditions that quantitatively convert NO_2 and accurately quantify the resulting NO_2^- . We then apply the method to a test case, showing how an $\cdot\text{OH}$ scavenger in solution prevents the oxidation of NO_2^- to NO_2 but does not enhance the overall photolysis efficiency of nitrate.

25

1.0 Introduction


Nitrogen oxides (i.e., nitrogen dioxide (NO_2) and nitric oxide (NO)) and nitrous acid (HONO) are reactive species that play key roles in the formation of tropospheric ozone and hydroxyl radicals ($\cdot\text{OH}$) (Acker et al., 2006; Seinfeld and Pandis, 2006). The fast oxidation of NO_2 to HNO_3 is an important sink of gas-phase NO_x , while the resulting gas-phase nitric acid and aqueous nitrate are traditionally considered relatively stable reservoir species (Stavrakou et al., 2013; Ye et al., 2017). Although nitrate can photolyze to reform NO_x , the lifetime of nitrate is long enough that the small production rates of NO_x and HONO from nitrate photolysis have been considered important only in remote areas (Romer et al., 2018). However, field studies over the past several decades have often

35 shown that atmospheric measurements of HONO and NO_x are higher than modeled values (Li et al., 2014; Zhou et

al., 2002; Romer et al., 2018). This suggests that nitrate photolysis in the atmosphere is faster than originally considered and, therefore, might be a more significant source of HONO and NO_x (Kasibhatla et al., 2018; Andersen et al., 2023; Zhou et al., 2003).

In sunlight (i.e., for wavelengths above 280 nm) aqueous NO₃⁻ photolysis proceeds through two channels:

40



45 The first channel (R1) produces NO₂ and ·O⁻ (which is rapidly protonated to ·OH) and the second produces nitrite (NO₂⁻) and an oxygen atom O(³P). Channel 1 has an average quantum yield of (1.19 ± 0.29)% at 293 K for illumination wavelengths above 300 nm (Chu and Anastasio, 2003; Zellner et al., 1990; Warneck and Wurzinger, 1988; Zepp et al., 1987), as shown in Table S1. The quantum yield for channel 2 (R2) is sometimes erroneously reported to be an order of magnitude smaller than that of channel 1, but in fact the values are comparable: channel 2
50 has an average quantum yield of (0.98 ± 0.11)% at 293 K for wavelengths above 300 nm (Benedict et al., 2017; McFall et al., 2018; Warneck and Wurzinger, 1988; Goldstein and Rabani, 2007) (Table S1).


These two quantum yields have been determined using different analytical methods. Generally, researchers either monitor the production of hydroxyl radical (·OH) from channel 1 or the production of nitrite (NO₂⁻) from channel 2. ·OH is typically quantified using a chemical probe (e.g., benzoic acid) that reacts to form a stable product
55 (e.g., *p*-hydroxybenzoic acid) that is monitored by HPLC (Chu and Anastasio, 2003). In contrast, NO₂⁻ is typically measured via ion chromatography or the more sensitive longpath-Griess method that derivatizes nitrite and measures the highly colored azo-product (Benedict et al., 2017; Ridnour et al., 2000).

Other studies have measured the gas-phase production of NO₂ and/or HONO, which is formed from the protonation of NO₂⁻. However, these gas-phase studies are limited to a specific pH range in order to measure HONO
60 production (Scharko et al., 2014), employ separate instruments to measure HONO and NO₂, and focus on how the production rates of NO₂ and HONO depend on experimental conditions (Frey et al., 2013; Ma et al., 2021; Liang et al., 2021). Although it is possible to measure both NO_x and HONO with commercially available instruments, researchers often engineer their own instrument to measure HONO and operate a second analyzer for the NO₂ channel (Shi et al., 2021; Wang et al., 2021a; Ma et al., 2021). Furthermore, gas-phase studies do not measure
65 quantum yields, but instead examine how the production rates of NO₂ and/or HONO are altered by factors such as the presence of other chemical species.

Typically researchers define an enhancement in nitrate photolysis as an experimentally measured production rate or quantum yield divided by the value under a standard condition (Liang et al., 2021; Wang et al., 2021b; Shi et al., 2021; Zhou et al., 2003). For example, a measured apparent nitrite quantum yield of 8% in the
70 presence of light-absorbing vanillic acid (Wang et al., 2021b) represents an 8-fold enhancement. If we want to fully understand the impact of an enhancement, the quantum yields for both channels must be measured. For instance, if one measures only the NO₂ channel and discovers an enhanced formation rate, it would be unclear whether NO₂⁻

production also increased or if NO_2^- is being converted to NO_2 . Therefore, it would be useful to be able to measure both channels of nitrate photolysis using a single analytical method.

75 One possible method to measure both channels is by reducing NO_2 to NO_2^- after photolysis, such that the total measured NO_2^- is the combination of NO_2 from channel 1 and NO_2^- from channel 2. S(IV) (i.e., sulfite (SO_3^{2-}) and bisulfite (HSO_3^-)) can reduce NO_2 to NO_2^- through the following overall reaction (Lee and Schwartz, 1982; Clifton et al., 1988; Wang et al., 2020; Song et al., 2021):



Although industry has used this reaction to convert NO_2 to NO_2^- , they often operate at very high temperatures, or include additives to enhance the diffusion of NO_2 into the aqueous phase (Shen and Rochelle, 1998; Lian et al., 2022).

85 Our goal is to use S(IV) chemistry to determine both channels of nitrate photolysis by performing two experiments using the same analytical method. In the first run, we measure NO_2^- production directly to quantify channel 2. In the second experiment, we use S(IV) to convert photoproducted NO_2 to NO_2^- so that the measured nitrite represents the sum of both NO_2 and NO_2^- . Then we quantify channel 1 by subtracting the NO_2^- experiment result from the combined ($\text{NO}_2 + \text{NO}_2^-$) experiment result. If this approach is successful, it would simplify and
90 expand our ability to analyze NO_2 and NO_2^- .

2.0 Methods

2.1 Materials

Information about materials and chemicals is in Section S1 of the supplement.

95 **2.2 Sample Illumination**

Illumination solutions were prepared daily, were air saturated, and contained 50 μM NaNO_3 , either 0 or 50 μM 2-propanol, and varying concentrations of S(IV). The pH of the solution was either controlled by a 0.010 M phosphate buffer (pH 5 or 8) or the added S(IV) (pH 8). Samples were illuminated with 313 nm light from a 1000 W Hg/Xe arc lamp with a downstream monochromator (Spectral Energy) and a 310 nm long-pass filter upstream of
100 the sample. 800 μL of aqueous sample in an upright 2 mL HPLC vial (low impurity Type I Class A borosilicate glass, 12 mm O.D. \times 32 mm H, Shimadzu) sealed with a septum cap was illuminated from its side. Samples were illuminated with constant stirring in a custom-built, Peltier-cooled aluminum housing (Paige Instruments) that was held at 20 $^\circ\text{C}$ by a recirculating water bath. Samples were kept sealed throughout the illumination. Dark controls containing the same solution as the illuminated sample but not exposed to light were analyzed periodically
105 throughout each experiment. Nitrite production was never detected in the dark controls. Under our conditions, experiments without S(IV) produced no more than 180 nM NO_2^- , and experiments with S(IV) produced no more than 180 nM $\text{NO}_2 + \text{NO}_2^-$.

2.3 Measurement of Nitrite

110 After illuminating all the samples for a given experiment, we determined nitrite concentrations using the
Griess method, a spectrophotometric technique that forms an azo-dye complex (Doane and Horwath, 2003; Benedict
et al., 2017; McFall et al., 2018). Our experiments had three different sample treatments: (1) no S(IV) in solution,
(2) S(IV) in solution during illumination, and (3) S(IV) added to the solution after illumination. Each treatment
required a slightly different method to efficiently form the azo-dye. The stabilities of the samples are discussed in
115 Section S3 of the supplement.

For samples without S(IV), the Griess method (Pratt et al., 1995; Moorcroft et al., 2001; Ridnour et al.,
2000; Benedict et al., 2017) could be used without adaptation. Within 10 minutes of stopping illumination, we
added 25 μL of a 1% sulfanilamide in 10% HCl (v/v) solution, and let it react for 10 minutes in the dark. We then
added 25 μL of 0.1% N-(1-Naphthyl)ethylenediamine dihydrochloride (NED) solution and allowed it to react for 10
120 minutes to form the azo-dye.

Treatment 2, where S(IV) was present in solution during illumination, required an additional step because
S(IV) interferes with the Griess reagents (SI Section S2). After illumination, we first added hydrogen peroxide
(H_2O_2) to the 800 μL sample to obtain a 2:1 molar ratio of H_2O_2 :S(IV). This was done to oxidize S(IV) to sulfate,
which does not interfere with nitrite determination. Within one minute of adding H_2O_2 , we added 50 μL of 1%
125 sulfanilamide in 30% (v/v) HCl solution and allowed the solution to react for 10 minutes in the dark. Then we
added 50 μL of 0.1% NED solution to the sample and allowed it to react for another 10 minutes in the dark.

Treatment 3 is similar to treatment 2 with one key difference: S(IV) was added to the solution after
illumination. Because NO_2 is volatile and would escape the illumination container if opened, we developed a
method to add the S(IV) without opening the vial. This was done by using a syringe with a hypodermic needle to
130 directly inject 37.5 μL of a 33.3 mM sulfite solution at pH 9 through the septum into the HPLC vial immediately
after the illumination was stopped. The vial was then left to react while stirring for 30 minutes in the dark at room
temperature to completely convert NO_2 to NO_2^- . The samples were then treated exactly as in treatment 2, adding
 H_2O_2 , then 50 μL of sulfanilamide in 30% HCl and then 50 μL of NED.

Once the azo-dye was formed, we measured light absorption at ~ 540 nm in the developed solutions using a
135 TIDAS II spectrophotometer (World Precision Instruments) with a liquid waveguide capillary cell (LWCC; nominal
length of 100 cm, effective path length of 94 cm, 250 μL volume), and a tungsten lamp. The TIDAS contains two
lamps, but the deuterium lamp (200–350 nm) caused an artifact in previous experiments (Benedict et al., 2017), so it
was kept off during our measurements. The absorption spectrum was measured from 350 to 700 nm so we could
correct for any baseline shifts. The peak height between 530 and 550 nm was determined as the difference between
140 the maximum absorbance in this range relative to a baseline drawn from the local absorption minima between 400
and 500 nm and between 550 and 700 nm. The limits of detection for nitrite were 7 nM for sample treatment 1 and
11 nM for sample treatments 2 and 3, as determined using the method of Armbruster and Pry (2008). Fresh
standards of sodium nitrite (0 to 200 nM) were prepared daily and used to calibrate the spectrophotometer. As S(IV)
and H_2O_2 decrease the absorbance response from the spectrophotometer (Figure S2), the same concentrations of
145 S(IV) and H_2O_2 used in the samples were also added to the standards to correct for this matrix effect. Samples and
other solutions were manually injected into the LWCC with a syringe, and 4 mL of Milli-Q water was injected

between samples to eliminate carry over. We cleaned the LWCC both before and after each experiment with 1 mL injections of three separate cleaning solutions: 1 M NaOH, 1 M HCl, and 50% methanol/50% Milli-Q (MQ) water, with pure MQ injected between each cleaning solution.

150 Daily controls included a replicate standard, MQ injection as a check for carry over, and a secondary check standard (Dionex). Analyses were deemed acceptable if the MQ check was below the lowest non-zero standard (10 nM NO₂⁻) and if both the replicate standard and secondary check standard concentrations were within 15% of known values.

155

2.3 Chemical Actinometry and Calculation of Quantum Yield

The photon flux was measured daily using 2-nitrobenzaldehyde (2NB) as a chemical actinometer (Galbavy et al., 2010). Actinometry was performed under the same conditions (container, volume of sample, temperature) as nitrate photolysis. Under low-light-absorbing conditions, the measured rate constant for 2NB loss during 313 nm irradiation ($j_{2NB,313}$) is calculated using

160

$$j_{2NB,313} = 2.303 \times 10^3 (I_{313} l) (\epsilon_{2NB,313} \phi_{2NB,313}), \quad (1)$$

where $I_{313} l$ is the surface-area-normalized photon flux (mol-photon cm⁻² s⁻¹), $\epsilon_{2NB,313} \phi_{2NB,313} = 640 \text{ M}^{-1} \text{ cm}^{-1}$ is the product of the base-10 molar absorption coefficient and quantum yield for 2NB at 313 nm (Anastasio et al., 1994), 2.303 converts ϵ to base-e, and $10^3 \text{ cm}^3 \text{ L}^{-1}$ is for units conversion. Similarly, the formation rate constant for nitrite from nitrate photolysis is:

165

$$j(\text{NO}_3^- \rightarrow \text{NO}_2^-)_{313} = 2.303 \times 10^3 (I_{313} l) (\epsilon_{\text{NO}_3^-,313}) (\phi(\text{NO}_2^-)_{313}), \quad (2)$$

170

where $\phi(\text{NO}_2^-)_{313}$ is the quantum yield of nitrite formation from nitrate photolysis at 313 nm, and $\epsilon_{\text{NO}_3^-,313}$ is the base-10 molar absorption coefficient of nitrate at 313 nm, $5.29 \text{ M}^{-1} \text{ cm}^{-1}$ (Chu and Anastasio, 2003).

The rate of nitrite formation from nitrate photolysis, $d[\text{NO}_2^-]/dt$, is a first-order process:

175

$$\frac{d[\text{NO}_2^-]}{dt} = j(\text{NO}_3^- \rightarrow \text{NO}_2^-)_{313} [\text{NO}_3^-]. \quad (3)$$

Since the experiments were at short time scales where nitrate loss was negligible, the increase of nitrite was linear, and the nitrite formation rate could be determined with simple linear regression. Combining equations 1-3 allows us to solve for the quantum yield of nitrite:

180

$$\phi(\text{NO}_2^-)_{313} = \frac{d[\text{NO}_2^-]}{dt} \times \frac{\epsilon_{2NB,313} \phi_{2NB,313}}{j_{2NB,313} \epsilon_{\text{NO}_3^-,313} [\text{NO}_3^-]}. \quad (4)$$

185 For simplicity, and because all our experiments were performed with 313 nm illumination, we omit the “313” subscript throughout the rest of this manuscript.

2.4 Combined Quantum Yield and $\phi(\text{NO}_2)$ Calculations

190 For experiments with added S(IV), the measured concentration of nitrite represents both the primary nitrite from nitrate photolysis as well as secondary nitrite formed by conversion of NO_2 . Thus, the calculated quantum yield in experiments with S(IV), i.e., $\phi(\text{NO}_2^-)_{\text{S(IV)}}$, is a combination of the quantum yields for both channels 1 and 2:

$$\phi(\text{NO}_2^-)_{\text{S(IV)}} = \phi(\text{NO}_2^-) + f \times \phi(\text{NO}_2). \quad (5)$$

195 Here f is the fraction of NO_2 that reacts with S(IV) to make NO_2^- , as opposed to going down other pathways:

$$f = \frac{k_{\text{HSO}_3^- + \text{NO}_2} \times [\text{HSO}_3^-] + k_{\text{SO}_3^{2-} + \text{NO}_2} \times [\text{SO}_3^{2-}]}{k_{\text{HSO}_3^- + \text{NO}_2} \times [\text{HSO}_3^-] + k_{\text{SO}_3^{2-} + \text{NO}_2} \times [\text{SO}_3^{2-}] + k_{\text{other}}}, \quad (6)$$

200 where $k_{\text{S(IV)+NO}_2}$ is the reaction rate constant of S(IV) and NO_2 , $1.2 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ and $1.7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ for bisulfite and sulfite, respectively (Clifton et al., 1988) and k_{other} is the pseudo-first-order rate constant for all other pathways that consume NO_2 . The concentrations of bisulfite and sulfite are determined based on the total S(IV) in solution, [S(IV)], and their mole fractions, which depend on the two $\text{p}K_{\text{a}}$ values for S(IV) ($\text{p}K_{\text{a}1} = 1.9$, $\text{p}K_{\text{a}2} = 7.2$; Seinfeld and Pandis (2006)). As described below, at $\text{pH} \approx 8$ a S(IV) concentration of 1.5 mM and higher is sufficient to make f equal 1, i.e., S(IV) is essentially the only fate for NO_2 , so it is quantitatively converted to NO_2^- . Under this condition, we calculate the quantum yield for NO_2 formation, $\phi(\text{NO}_2)$, as the difference between the measured nitrite quantum yields in the presence and absence of S(IV):

205

$$\phi(\text{NO}_2) = \phi(\text{NO}_2^-)_{\text{S(IV)}} - \phi(\text{NO}_2^-). \quad (7)$$

3.0 Results

210 3.1 Modification of the Griess Method for Solutions Containing S(IV)

As described in Section S2, we found that the addition of micromolar levels of S(IV) interferes with the determination of nitrite because of two issues: (1) it prevents the formation of the azo-dye derivative and (2) it moves the solution acidity out of the required range. The first issue was solved by oxidizing the S(IV) to S(VI) with H_2O_2 prior to the addition of the Griess reagents (Figure S1). We added H_2O_2 to the samples such that there was a 2:1 molar ratio of H_2O_2 :S(IV), then within 1 minute of the addition of H_2O_2 , we added the sulfanilamide solution and, 10 min later, the NED reagent. After waiting another 10 minutes, we measured the UV-VIS spectra for the entire batch of samples within 20 minutes of capturing the spectrum of the first sample. We also doubled the


standard volumes of both Griess reagents added to the sample solutions to ensure that there were enough reactants to form the azo-dye.

220 The second issue caused by S(IV) was that it pushed the solutions to pH 7. This basicity prevented the conversion of nitrite to the azo-dye because this reaction requires a pH below 2. The standard 10% HCl (v/v) in the sulfanilamide solution only lowered the sample pH to ~ 4 for solutions containing 1.5 mM of sulfite. Per the recommendation by Doane and Horwath (2003), we increased the HCl concentration in the sulfanilamide solution to 30% (v/v), which lowered the pH of the sample-sulfanilamide mixture to less than 2, overcoming the pH issue
225 caused by S(IV).

3.2 Addition of S(IV) to Solution Prior to Illumination

Our goals in this initial set of experiments were to examine whether S(IV) in solution can convert photoproducted NO_2 to NO_2^- and, if so, to determine the concentration of aqueous S(IV) required to make this
230 conversion quantitative, i.e., close to 100%. If S(IV) can quantitatively convert NO_2 to NO_2^- , then the measured nitrite quantum yield at this S(IV) concentration should equal the sum of the quantum yields from both channels of nitrate photolysis.

We started experiments by running a test without S(IV) (50 μM NaNO_3 , 50 μM 2-propanol, and 293 K) to confirm that our result matches the literature. The average $\phi(\text{NO}_2^-)$ from our four replicate experiments without
235 S(IV) is $(1.05 \pm 0.06)\%$, which is statistically no different ($p = 0.36$) from the average of the literature values shown in Table S1, $(0.98 \pm 0.11)\%$. Then we began performing experiments with increasing concentrations of S(IV). As [S(IV)] increases, the apparent nitrite quantum yield increases until it reaches a plateau for S(IV) concentrations at roughly 500 μM and above (Figure 2). The measured quantum yield at the plateau, determined as the average (± 1 σ) of the individual experiments from 500 to 2000 μM S(IV), is $(2.01 \pm 0.05)\%$. This is slightly lower than, but
240 statistically indistinguishable ($p = 0.14$) from the sum of the average literature quantum yields for both channels, $(2.17 \pm 0.52)\%$, which is shown as the upper horizontal line in Figure 2. We then calculate $\phi(\text{NO}_2)$ by taking the difference between the quantum yield determined with S(IV), which measures the sum of the two channels, and the quantum yield for the nitrite channel (Eq. 7). This results in a value of $\phi(\text{NO}_2)$ of $(0.96 \pm 0.12)\%$, which is slightly lower than the average of previous experiments $(1.19 \pm 0.29)\%$, but statistically no different ($p = 0.10$). These
245 results confirm that S(IV) in the reaction solution during illumination can quantitatively convert photochemically produced NO_2 to NO_2^- , allowing the Griess spectrophotometric technique to quantify both channels of nitrate photolysis.



250 **Figure 2.** Measured apparent nitrite quantum yields for the photolysis of 50 μM nitrate solutions (293 K, pH 8) in the presence of different concentrations of S(IV). Hollow circles represent the average ($\pm 1 \sigma$) of individual experiments, which are shown as solid black points. The line through the data is a fit to equation 8. The lower grey area centered at 1.1% is the average of previously determined values of $\phi(\text{NO}_2^-) \pm 1 \sigma$, and the upper grey area centered at 2.2% is the sum of the quantum yields from both channels, $\phi(\text{NO}_2^- + \text{NO}_2) \pm 1 \sigma$, from the literature.

255 Literature values used to calculate these averages are in Table S1.

We can also use our Figure 2 data to estimate the value for k_{other} , the pseudo-first-order rate constant for NO_2 loss due to other pathways, i.e., not reacting with S(IV). Combining equations 5 and 6 yields:

$$260 \quad \phi(\text{NO}_2^-)_{(\text{SIV})} = \phi(\text{NO}_2^-) + \phi(\text{NO}_2) \times \frac{k_{\text{HSO}_3^- + \text{NO}_2} \times [\text{HSO}_3^-] + k_{\text{SO}_3^{2-} + \text{NO}_2} \times [\text{SO}_3^{2-}]}{k_{\text{HSO}_3^- + \text{NO}_2} \times [\text{HSO}_3^-] + k_{\text{SO}_3^{2-} + \text{NO}_2} \times [\text{SO}_3^{2-}] + k_{\text{other}}}. \quad (8)$$

Fitting this equation to our data using Python (Van Rossum and Drake Jr, 1995) yields the solid line in Figure 2 and parameter values of $k_{\text{other}} = 700 \pm 300 \text{ s}^{-1}$, $\phi(\text{NO}_2) = (0.94 \pm 0.07)\%$, and $\phi(\text{NO}_2^-) = (1.10 \pm 0.06)\%$. We can use the

265 value of k_{other} in equation 6 to calculate the percent of NO_2 that is converted to NO_2^- in solutions at a given S(IV) concentration and pH value: values are 96%, 98%, and 99% at 500, 1000, and 1500 μM S(IV), respectively, at pH 8.

3.3 Addition of S(IV) After Illumination

Our experiments above used S(IV) in the illumination solution to convert NO_2 to NO_2^- . While this method

270 works, it has the disadvantage that S(IV) might interfere with other reactive species or reaction pathways during illumination. To avoid this problem, in this section we examine whether we can prevent NO_2 from escaping the sample container and convert it to nitrite by adding S(IV) to the solution after illumination.

We made several changes to the procedure in Section 3.2 to ensure full conversion of NO₂ to NO₂⁻ when adding S(IV) after illumination. We examined the effectiveness of the potential changes based on a single trial where we tested three different treatments of the samples post-illumination: (1) adding 1.5 mM S(IV) to the samples and allowing them to stir for 30 minutes in the dark, (2) adding 1.0 mM S(IV) and stirring for 30 minutes, and (3) adding 1.5 mM S(IV) and stirring for 10 minutes. In each case, we added the S(IV) immediately after the end of sample illumination by injecting a small volume, 25 or 38 μL, of a 33 mM sodium sulfite stock solution through the septum of the HPLC cap with a syringe and hypodermic needle. The goal with this technique was to keep the illumination container sealed so that NO₂ could not escape. Measured values of $\phi(\text{NO}_2^-)_{\text{S(IV)}}$ were (1.97 ± 0.24)%, (1.53 ± 0.19)%, and (1.60 ± 0.45)% for treatments 1, 2, and 3, respectively. The only trial that seemed to completely convert all the NO₂ to NO₂⁻ was the first treatment, i.e., 1.5 mM S(IV) with 30 min of stirring. As such, we used this treatment method going forward.

We also estimated the timescale of NO₂ conversion to nitrite to compare with our experimental results. Based on the volumes in the reaction vial (800 μL of solution and ~1.3 mL of headspace), Henry's Law predicts (at 293 K) that 10% of NO₂ should be in the aqueous phase and 90% in the head space. Based on the kinetic data from Clifton et al. (1988), the lifetime of total NO₂ in the vial is approximately 1 ms. This means that there should have been no difference between the results of trial 1 and 3, which is not the case. It is unclear why there is a discrepancy between the theoretical and experimental timescales for the conversion of NO₂ to NO₂⁻.

Next, we examined whether the addition of S(IV) after illumination produced results that were the same as those for experiments where S(IV) was in the solution during illumination. We performed triplicate experiments measuring the combined NO₂ + NO₂⁻ quantum yield in pH 5 solution containing 50 μM NaNO₃ and 50 μM 2-propanol, and 1.5 mM S(IV) added to the solution after illumination. As shown in Figure S4, the average ± 1σ combined quantum yield from this set of experiments is (2.10 ± 0.08)%. This is statistically no different from the result we obtained above when S(IV) was present in the solution during illumination, (2.00 ± 0.14)% (*p* = 0.32), and no different from the literature value, (2.17 ± 0.52)% (*p* = 0.74; Table S1). This indicates that we can add S(IV) after the photoproduction of nitrogen dioxide has stopped and still convert all the NO₂ to NO₂⁻.

3.4 Applying the S(IV) Method: Impact of an •OH Scavenger

Our final step is to show the utility of determining both NO₂ and NO₂⁻ in a chemical system, by using the example of quantifying the impact of an •OH scavenger on the two channels from nitrate photolysis. Based on past work (Benedict et al., 2017; Roca et al., 2008; McFall et al., 2018), in the absence of a hydroxyl radical scavenger •OH can react with NO₂⁻ to form NO₂:



Because of this reaction, in the absence of an •OH scavenger, the NO₂⁻ quantum yield should be underestimated and the NO₂ quantum yield should be overestimated by an equal amount. In contrast, adding a scavenger suppresses the hydroxyl radical concentration and its impact on both photoproducts, giving the true quantum yield for each

310 channel. However, we expect that the combined quantum yield, i.e., the sum of values for both channels, will be the same regardless of the presence of $\cdot\text{OH}$ scavengers. That is, we expect that an $\cdot\text{OH}$ scavenger will prevent the conversion of NO_2^- to NO_2 but will not alter the overall photochemical efficiency of nitrate photolysis. While the impact of $\cdot\text{OH}$ scavengers on the nitrite channel has been examined previously, we are unaware of any past attempts to measure both channels in the presence and absence of scavengers.

315




Figure 3. Measured quantum yields of nitrite (blue bars), nitrogen dioxide (red bars), and both products (purple bars) from the photolysis of 50 μM NO_3^- at 293 K and pH 5. The hollow bars are experiments without 2-propanol (an $\cdot\text{OH}$ scavenger), while solid bars represent experiments with 50 μM 2-propanol. The grey bar is the sum of the average quantum yields for the two channels from past studies (Table S1). Arrows above the NO_2^- and NO_2 channels indicate the impact of the $\cdot\text{OH}$ scavenger. Error bars are $\pm 1\sigma$. Errors for the NO_2^- and $(\text{NO}_2^- + \text{NO}_2)$ quantum yields were calculated from replicate experiments; these errors were propagated to determine the error for the NO_2 channel result.

325 As shown by the blue arrow in Figure 3, the addition of an $\cdot\text{OH}$ scavenger increases the NO_2^- quantum yield (by 0.14%), as expected since it impedes the oxidation of nitrite by hydroxyl radical (Figure 1). Also consistent with our model above, the red arrow shows that the $\cdot\text{OH}$ scavenger decreases the quantum yield of the NO_2 channel (by 0.14%), a result of the suppression of nitrite oxidation by $\cdot\text{OH}$ to make NO_2 . The NO_2^- quantum yields without S(IV), with and without 2-propanol, are statistically different ($p = 0.04$). However, when S(IV) is added to solution, the presence or absence of an $\cdot\text{OH}$ scavenger has no impact ($p = 0.95$) on the sum of the quantum yields for the two channels. This is what we expect because the NO_2 that was formed from the reaction of $\cdot\text{OH}$ and NO_2^- is converted back to NO_2^- by S(IV), resulting in the same total amount of $\text{NO}_2^- + \text{NO}_2$ in the two sets of experiments. This shows that the addition of a $\cdot\text{OH}$ scavenger does not impact the overall efficiency of nitrate photolysis (i.e., the sum of the quantum yields of the two channels) but prevents the oxidation of NO_2^- to NO_2 .

330

335 Our quantum yields in this set of experiments are in good agreement with previously determined values. As mentioned in section 3.2, our nitrite quantum yield without S(IV) agrees with previously reported $\phi(\text{NO}_2^-)$ values. Our combined quantum yield values, $\phi(\text{NO}_2^-)_{\text{S(IV)}}$, are $(2.10 \pm 0.08)\%$ and $(2.09 \pm 0.16)\%$ with and without an $\cdot\text{OH}$ scavenger, respectively (Figure 3). Our values here agree with the Table S1 average of previously determined combination of both channels, $(2.17 \pm 0.52)\%$ ($p > 0.70$). The NO_2 channel was calculated, using Eq. (7), as the difference in the quantum yield between experiments with S(IV) added after illumination and experiments without S(IV). In this set of experiments, our measured $\phi(\text{NO}_2)$, $(1.05 \pm 0.10)\%$, is similar to the average of the literature values, $(1.19 \pm 0.29)\%$ ($p = 0.47$), as shown in Table S1.

345 4.0 Impacts/Implications

We have demonstrated that S(IV) can convert aqueous NO_2 to NO_2^- , which allows the production of both the gas- and aqueous-phase products of nitrate photolysis to be quantified in the aqueous phase in a sealed container using the same analytical method. Although nitrate photolysis is traditionally considered a minor source of NO_x , recent research has shown that the efficiency of nitrate photolysis can be enhanced by other light-absorbing compounds or its physical environment (Wang et al., 2021b; Mora Garcia et al., 2021; McFall et al., 2018). It is important to understand whether an apparent enhancement impacts only one channel, both channels, or is due simply to a conversion of one product to another. As many field studies have noted that the measured enhancement impacts the NO_2^- channel more than the NO_2 channel, it is likely that different chemicals impact nitrate photolysis in a variety of ways (Andersen et al., 2023; Kasibhatla et al., 2018; Ye et al., 2016). Understanding how different variables impact nitrate photolysis will allow a more comprehensive understanding of nitrogen cycling and should improve model predictions of ambient NO_x and HONO concentrations. Performing experiments with and without S(IV) for a given experimental condition will allow laboratory experiments to determine both channels of nitrate photolysis, which will reveal whether one or both channels are enhanced or if secondary chemistry is converting one product to the other.

360 This new S(IV) method also has applications beyond nitrate photolysis, as it can be used for any system where NO_2 needs to be quantified. This could include studies where NO_2 production occurs in the aqueous phase, such as the decomposition of metallic nitrate compounds (Gallagher et al., 1971; Yuvaraj et al., 2003), or in studies where the production of NO_2 is small enough that it cannot be quantified by commercially available analyzers.

365 Code and data availability.

All data and code can be obtained by emailing the corresponding author at canastasio@ucdavis.edu

Author Contributions

370 AL and CA conceptualized the study, while AL, JP, and MO performed the experiments, and AL performed the coding and data analysis. AL wrote the paper with input from CA.

Competing interests

The authors declare that they have no conflict of interest.

375 Acknowledgments

We appreciate the helpful comments of the anonymous reviewers.

Financial support

This research was supported by the National Science Foundation (grant CHE-2305164) and the California

380 Agricultural Experiment Station (Project CA-D-LAW-6403-RR).

References

- 385 Acker, K., Möller, D., Wieprecht, W., Meixner, F. X., Bohn, B., Gilge, S., Plass-Dülmer, C., and Berresheim, H.: Strong daytime production of OH from HNO₂ at a rural mountain site, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL024643>, 2006.
- Anastasio, C., Faust, B., and Allen, J.: Aqueous phase photochemical formation of hydrogen peroxide in authentic cloud waters, *Journal of Geophysical Research: Atmospheres*, 99, 8231–8248, <https://doi.org/10.1029/94JD00085>, 1994.
- 390 Andersen, S. T., Carpenter, L. J., Reed, C., Lee, J. D., Chance, R., Sherwen, T., Vaughan, A. R., Stewart, J., Edwards, P. M., Bloss, W. J., Sommariva, R., Crilley, L. R., Nott, G. J., Neves, L., Read, K., Heard, D. E., Seakins, P. W., Whalley, L. K., Boustead, G. A., Fleming, L. T., Stone, D., and Fomba, K. W.: Extensive field evidence for the release of HONO from the photolysis of nitrate aerosols, *Science Advances*, 9, eadd6266, <https://doi.org/10.1126/sciadv.add6266>, 2023.
- 395 Armbruster, D. A. and Pry, T.: Limit of Blank, Limit of Detection and Limit of Quantitation, *Clin Biochem Rev*, 29, S49–S52, 2008.
- Benedict, K. B., McFall, A. S., and Anastasio, C.: Quantum Yield of Nitrite from the Photolysis of Aqueous Nitrate above 300 nm, *Environ. Sci. Technol.*, 51, 4387–4395, <https://doi.org/10.1021/acs.est.6b06370>, 2017.
- 400 Chu, L. and Anastasio, C.: Quantum Yields of Hydroxyl Radical and Nitrogen Dioxide from the Photolysis of Nitrate on Ice, *J. Phys. Chem. A*, 107, 9594–9602, <https://doi.org/10.1021/jp0349132>, 2003.
- Clifton, C. L., Altstein, Nisan., and Huie, R. E.: Rate constant for the reaction of nitrogen dioxide with sulfur(IV) over the pH range 5.3–13, *Environ. Sci. Technol.*, 22, 586–589, <https://doi.org/10.1021/es00170a018>, 1988.
- Doane, T. A. and Horwath, W. R.: Spectrophotometric Determination of Nitrate with a Single Reagent, *Analytical Letters*, 36, 2713–2722, <https://doi.org/10.1081/AL-120024647>, 2003.
- 405 Frey, M. M., Brough, N., France, J. L., Anderson, P. S., Traulle, O., King, M. D., Jones, A. E., Wolff, E. W., and Savarino, J.: The diurnal variability of atmospheric nitrogen oxides (NO and NO₂) above the Antarctic Plateau driven by atmospheric stability and snow emissions, *Atmospheric Chemistry and Physics*, 13, 3045–3062, <https://doi.org/10.5194/acp-13-3045-2013>, 2013.
- 410 Galbavy, E. S., Ram, K., and Anastasio, C.: 2-Nitrobenzaldehyde as a chemical actinometer for solution and ice photochemistry, *Journal of Photochemistry and Photobiology A: Chemistry*, 209, 186–192, <https://doi.org/10.1016/j.jphotochem.2009.11.013>, 2010.
- Gallagher, P. K., Schrey, F., and Prescott, B.: The thermal decomposition of aqueous manganese (II) nitrate solution, *Thermochimica Acta*, 2, 405–412, [https://doi.org/10.1016/0040-6031\(71\)85016-5](https://doi.org/10.1016/0040-6031(71)85016-5), 1971.
- 415 Goldstein, S. and Rabani, J.: Mechanism of Nitrite Formation by Nitrate Photolysis in Aqueous Solutions: The Role of Peroxynitrite, Nitrogen Dioxide, and Hydroxyl Radical, *J. Am. Chem. Soc.*, 129, 10597–10601, <https://doi.org/10.1021/ja073609+>, 2007.
- 420 Kasibhatla, P., Sherwen, T., Evans, M. J., Carpenter, L. J., Reed, C., Alexander, B., Chen, Q., Sulprizio, M. P., Lee, J. D., Read, K. A., Bloss, W., Crilley, L. R., Keene, W. C., Pszenny, A. A. P., and Hodzic, A.: Global impact of nitrate photolysis in sea-salt aerosol on NO_x, OH, and O₃ in the marine boundary layer, *Atmospheric Chemistry and Physics*, 18, 11185–11203, <https://doi.org/10.5194/acp-18-11185-2018>, 2018.
- Lee, Y. N. and Schwartz, S. E.: Kinetics of oxidation of aqueous sulfur(IV) by nitrogen dioxide, Brookhaven National Lab., Upton, NY (USA), 1982.

- 425 Li, X., Rohrer, F., Hofzumahaus, A., Brauers, T., Häseler, R., Bohn, B., Broch, S., Fuchs, H., Gomm, S., Holland, F., Jäger, J., Kaiser, J., Keutsch, F. N., Lohse, I., Lu, K., Tillmann, R., Wegener, R., Wolfe, G. M., Mentel, T. F., Kiendler-Scharr, A., and Wahner, A.: Missing Gas-Phase Source of HONO Inferred from Zeppelin Measurements in the Troposphere, *Science*, 344, 292–296, <https://doi.org/10.1126/science.1248999>, 2014.
- Lian, Z., Li, G., Zhang, S., Ma, W., and Zhong, Q.: Mechanism and Kinetic Study of Cyclodextrin Use to Facilitate NO₂ Absorption in Sulfite Solutions, *Environmental Science & Technology*, 56, 7696–7706, <https://doi.org/10.1021/acs.est.2c00838>, 2022.
- 430 Liang, Z., Zhang, R., Gen, M., Chu, Y., and Chan, C. K.: Nitrate Photolysis in Mixed Sucrose–Nitrate–Sulfate Particles at Different Relative Humidities, *J. Phys. Chem. A*, 125, 3739–3747, <https://doi.org/10.1021/acs.jpca.1c00669>, 2021.
- 435 Ma, Q., Zhong, C., Ma, J., Ye, C., Zhao, Y., Liu, Y., Zhang, P., Chen, T., Liu, C., Chu, B., and He, H.: Comprehensive Study about the Photolysis of Nitrates on Mineral Oxides, *Environ. Sci. Technol.*, 55, 8604–8612, <https://doi.org/10.1021/acs.est.1c02182>, 2021.
- McFall, A. S., Edwards, K. C., and Anastasio, C.: Nitrate Photochemistry at the Air–Ice Interface and in Other Ice Reservoirs, *Environ. Sci. Technol.*, 52, 5710–5717, <https://doi.org/10.1021/acs.est.8b00095>, 2018.
- Moorcroft, M. J., Davis, J., and Compton, R. G.: Detection and determination of nitrate and nitrite: a review, *Talanta*, 54, 785–803, [https://doi.org/10.1016/S0039-9140\(01\)00323-X](https://doi.org/10.1016/S0039-9140(01)00323-X), 2001.
- 440 Mora Garcia, S. L., Pandit, S., Navea, J. G., and Grassian, V. H.: Nitrous Acid (HONO) Formation from the Irradiation of Aqueous Nitrate Solutions in the Presence of Marine Chromophoric Dissolved Organic Matter: Comparison to Other Organic Photosensitizers, *ACS Earth Space Chem.*, 5, 3056–3064, <https://doi.org/10.1021/acsearthspacechem.1c00292>, 2021.
- 445 Pratt, P. F., Nithipatikom, K., and Campbell, W. B.: Simultaneous Determination of Nitrate and Nitrite in Biological Samples by Multichannel Flow Injection Analysis, *Analytical Biochemistry*, 231, 383–386, <https://doi.org/10.1006/abio.1995.0067>, 1995.
- Ridnour, L. A., Sim, J. E., Hayward, M. A., Wink, D. A., Martin, S. M., Buettner, G. R., and Spitz, D. R.: A Spectrophotometric Method for the Direct Detection and Quantitation of Nitric Oxide, Nitrite, and Nitrate in Cell Culture Media, *Analytical Biochemistry*, 281, 223–229, <https://doi.org/10.1006/abio.2000.4583>, 2000.
- 450 Roca, M., Zahardis, J., Bone, J., El-Maazawi, M., and Grassian, V. H.: 310 nm Irradiation of Atmospherically Relevant Concentrated Aqueous Nitrate Solutions: Nitrite Production and Quantum Yields, *J. Phys. Chem. A*, 112, 13275–13281, <https://doi.org/10.1021/jp809017b>, 2008.
- 455 Romer, P. S., Wooldridge, P. J., Crouse, J. D., Kim, M. J., Wennberg, P. O., Dibb, J. E., Scheuer, E., Blake, D. R., Meinardi, S., Brosius, A. L., Thames, A. B., Miller, D. O., Brune, W. H., Hall, S. R., Ryerson, T. B., and Cohen, R. C.: Constraints on Aerosol Nitrate Photolysis as a Potential Source of HONO and NO_x, *Environ. Sci. Technol.*, 52, 13738–13746, <https://doi.org/10.1021/acs.est.8b03861>, 2018.
- Scharko, N. K., Berke, A. E., and Raff, J. D.: Release of Nitrous Acid and Nitrogen Dioxide from Nitrate Photolysis in Acidic Aqueous Solutions, *Environ. Sci. Technol.*, 48, 11991–12001, <https://doi.org/10.1021/es503088x>, 2014.
- 460 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics : From Air Pollution to Climate Change*, 2nd ed., John Wiley & Sons, Inc., Hoboken NJ, 1203 pp., 2006.
- Shen, C. H. and Rochelle, G. T.: Nitrogen Dioxide Absorption and Sulfite Oxidation in Aqueous Sulfite, *Environ. Sci. Technol.*, 32, 1994–2003, <https://doi.org/10.1021/es970466q>, 1998.

- 465 Shi, Q., Tao, Y., Krechmer, J. E., Heald, C. L., Murphy, J. G., Kroll, J. H., and Ye, Q.: Laboratory Investigation of Renoxification from the Photolysis of Inorganic Particulate Nitrate, *Environ. Sci. Technol.*, 55, 854–861, <https://doi.org/10.1021/acs.est.0c06049>, 2021.
- Song, H., Lu, K., Ye, C., Dong, H., Li, S., Chen, S., Wu, Z., Zheng, M., Zeng, L., Hu, M., and Zhang, Y.: A comprehensive observation-based multiphase chemical model analysis of sulfur dioxide oxidations in both summer and winter, *Atmospheric Chemistry and Physics*, 21, 13713–13727, <https://doi.org/10.5194/acp-21-13713-2021>, 2021.
- 470 Stavrakou, T., Müller, J.-F., Boersma, K. F., van der A, R. J., Kurokawa, J., Ohara, T., and Zhang, Q.: Key chemical NO_x sink uncertainties and how they influence top-down emissions of nitrogen oxides, *Atmospheric Chemistry and Physics*, 13, 9057–9082, <https://doi.org/10.5194/acp-13-9057-2013>, 2013.
- Van Rossum, G. and Drake Jr, F. L.: Python reference manual, Centrum voor Wiskunde en Informatica Amsterdam, 1995.
- 475 Wang, J., Li, J., Ye, J., Zhao, J., Wu, Y., Hu, J., Liu, D., Nie, D., Shen, F., Huang, X., Huang, D. D., Ji, D., Sun, X., Xu, W., Guo, J., Song, S., Qin, Y., Liu, P., Turner, J. R., Lee, H. C., Hwang, S., Liao, H., Martin, S. T., Zhang, Q., Chen, M., Sun, Y., Ge, X., and Jacob, D. J.: Fast sulfate formation from oxidation of SO₂ by NO₂ and HONO observed in Beijing haze, *Nat Commun*, 11, 2844, <https://doi.org/10.1038/s41467-020-16683-x>, 2020.
- 480 Wang, X., Dalton, E. Z., Payne, Z. C., Perrier, S., Riva, M., Raff, J. D., and George, C.: Superoxide and Nitrous Acid Production from Nitrate Photolysis Is Enhanced by Dissolved Aliphatic Organic Matter, *Environ. Sci. Technol. Lett.*, 8, 53–58, <https://doi.org/10.1021/acs.estlett.0c00806>, 2021a.
- 485 Wang, Y., Huang, D. D., Huang, W., Liu, B., Chen, Q., Huang, R., Gen, M., Mabato, B. R. G., Chan, C. K., Li, X., Hao, T., Tan, Y., Hoi, K. I., Mok, K. M., and Li, Y. J.: Enhanced Nitrite Production from the Aqueous Photolysis of Nitrate in the Presence of Vanillic Acid and Implications for the Roles of Light-Absorbing Organics, *Environ. Sci. Technol.*, 55, 15694–15704, <https://doi.org/10.1021/acs.est.1c04642>, 2021b.
- Warneck, P. and Wurzinger, C.: Product quantum yields for the 305-nm photodecomposition of nitrate in aqueous solution, *J. Phys. Chem.*, 92, 6278–6283, <https://doi.org/10.1021/j100333a022>, 1988.
- Ye, C., Gao, H., Zhang, N., and Zhou, X.: Photolysis of Nitric Acid and Nitrate on Natural and Artificial Surfaces, *Environ. Sci. Technol.*, 50, 3530–3536, <https://doi.org/10.1021/acs.est.5b05032>, 2016.
- 490 Ye, C., Zhang, N., Gao, H., and Zhou, X.: Photolysis of Particulate Nitrate as a Source of HONO and NO_x, *Environ. Sci. Technol.*, 51, 6849–6856, <https://doi.org/10.1021/acs.est.7b00387>, 2017.
- Yuvaraj, S., Fan-Yuan, L., Tsong-Huei, C., and Chuin-Tih, Y.: Thermal Decomposition of Metal Nitrates in Air and Hydrogen Environments, *J. Phys. Chem. B*, 107, 1044–1047, <https://doi.org/10.1021/jp026961c>, 2003.
- 495 Zellner, R., Exner, M., and Herrmann, H.: Absolute OH quantum yields in the laser photolysis of nitrate, nitrite and dissolved H₂O₂ at 308 and 351 nm in the temperature range 278–353 K, *J Atmos Chem*, 10, 411–425, <https://doi.org/10.1007/BF00115783>, 1990.
- Zepp, R. G., Hoigné, J., and Bader, H.: Nitrate-induced photooxidation of trace organic chemicals in water, *Environ. Sci. Technol.*, 21, 443–450, <https://doi.org/10.1021/es00159a004>, 1987.
- 500 Zhou, X., Civerolo, K., Dai, H., Huang, G., Schwab, J., and Demerjian, K.: Summertime nitrous acid chemistry in the atmospheric boundary layer at a rural site in New York State, *Journal of Geophysical Research: Atmospheres*, 107, ACH 13-1-ACH 13-11, <https://doi.org/10.1029/2001JD001539>, 2002.

Zhou, X., Gao, H., He, Y., Huang, G., Bertman, S. B., Civerolo, K., and Schwab, J.: Nitric acid photolysis on surfaces in low-NO_x environments: Significant atmospheric implications, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2003GL018620>, 2003.

505