	Frida S. Hoem <sup>1</sup> , Adrián López-Quirós <sup>2, 3</sup> , Suzanna van de Lagemaat <sup>1</sup> , Johan Etourneau <sup>4,5</sup> , Marie-		
5	Alexandrine Sicre <sup>6</sup> , Carlota Escutia <sup>7</sup> , Henk Brinkhuis <sup>1,8</sup> , Francien Peterse <sup>1</sup> , Francesca Sangiorgi <sup>1</sup> , Peter K. Bijl <sup>1</sup>	(	Deleted: 4.
10	<sup>1</sup> Department of Earth Science, Utrecht University, Utrecht, the Netherlands <sup>2</sup> Department of Stratigraphy and Paleontology, University of Granada, Granada, Spain <sup>3</sup> Department of Geoscience/iCLIMATE Centre, Aarhus University, Aarhus C, Denmark <sup>4</sup> EPHE, PSL Research University, Paris, France. <sup>5</sup> University of Bordeaux, CNRS, Pessac, France <sup>6</sup> LOCEAN, CNRS, Sorbonne Université, Campus Pierre et Marie Curie, Paris, France. <sup>7</sup> UACT. CSIC Universited de Granada, Spain		Formatted: French
	<sup>8</sup> Department of Ocean Systems research OCS, Royal Netherlands Institute for Sea Research (NIOZ), Texel, the Netherlands		Formatted: English (UK)
15	Correspondence to: Frida S. Hoem (Frida spho@gmail.com)	U	Formatted: French
10			
	Abstract. At present, a strong latitudinal sea surface temperature (SST) gradient of ~16°C exists across the Southern Ocean,	(I	Deleted: e.g.,
	maintained by the Antarctic Circumpolar Current (ACC) and a set of complex frontal systems. Together with the Antarctic ice	//	Deleted: biomarker
	masses this system has formed one of the most important global climate regulators. The timing of the onset of the ACC-	///	Deleted: , calibrated to SST,
1	sustem its devialenment towards modern day strength and the consequences for the latitudinal SCT and internet around the	////	Deleted: southern
•	system, its development towards modern-day strength, and the consequences for the latitudinal SST gradient around the	// (	Deleted: overall
20	southern Atlantic Ocean, are still uncertain. Here we present new TEX <sub>86</sub> - <u>derived SST records, from two sites located east of</u>	///	Deleted: rather
	Drake Passage (southwestern South Atlantic) to assist in better understanding two critical time intervals of prominent climate	////	Deleted: Late Eocene to Early Oligocene
	transitions during the Cenozoic: The Late Eocene-Early Oligocene (ODP Site 696) and Middle-Late Miocene (IODP Site	9/1/	Deleted: (~16°C)
1	U1536) transitions. Our results show temperate conditions (20–11°C) during the first time interval, with a weaker latitudinal	ע / /	Deleted: across
	SST gradient (~8°C) across the Atlantic sector of the Southern Ocean compared to present day. We ascribe the regional	///	Deleted: S
25	similarity in SSTs in the Late Eocene_Early Oligocene South Atlantic to a persistent strong subpolar gyre circulation	/	Deleted: G
23	similarly in 5515 in the Late Excellence any ongotene south Auanue to a persistent, strong guopoiat gate encutation,	<u>ا</u> ر *	Deleted: all sites
	connecting Sites 696 and 511, which can only exist in absence of a strong throughflow across the Drake Passage. Surprisingly,		Deleted: records
	the southern South Atlantic record Site 696 show comparable SSTs (~12-14°C) during both the Earliest Oligocene Oxygen	<u> </u>	Deleted: strongly
-	Isotope Step (EOIS, ~33.65 Ma) and the Miocene Climate Optimum (MCO, ~16.5 Ma). Apparently, maximum Oligocene	- / ß	Deleted: around Antarctica
	Antarctic ice volume could coexist with warm ice-proximal surface ocean conditions, while at similar ocean temperatures, the	///	Deleted:
30	Middle Miocene Antarctic ice sheet was likely reduced. Only a few Mid-Late Miocene (discontinuous) high latitude records /	////	Deleted: southern
	exist due to ice advances causing unconformities. Our low-resolution Site U1536 record of Southern South Atlantic SSTs	///	Deleted: meant it experience
I	exist due to fee duvances eating uncontributes. Our few resolution one of 550 record of poundar related to 575	MAS	Deleted: d
1	coored to ~5 C at the onset of the Mildele Mildele Chinate Transition (Mildel, 14 Ma), making it the coldest oceanic region	1///	Deleted: latter part of the
	in the poorly recorded Antarctic realm and the likely main location for deep water formation. The already cold southwestern f	$\mathbb{Z}$	Deleted: , which
	South Atlantic conditions at MMCT with little additional cooling during the Late Miocene contrasts the profound cooling in		Deleted: due to northward expansion of the Southern Ocean frontal systems
35	the lower latitudes and other sectors of the Southern Ocean, due to northward expansion of the Southern Ocean frontal systems,		Formatted: English (UK)

Formatted: English (US)

# Late Cenozoic Sea Surface Temperature evolution of the South Atlantic Ocean

#### 60 1 Introduction

Today, Southern Ocean surface flow is dominated by the strongest ocean surface current on Earth, the Antarctic Circumpolar Current (ACC). This wind-driven, eastward flowing surface current is associated with strong meridional gradients in sea surface temperature (SST) ( $\sim$ 16°C<sub>4</sub> $\sim$ 45–60°S) and oceanographic conditions (Locarnini et al., 2018). Questions remain about the timing and nature of the development of the ACC and concomitant evolution the complex Southern Ocean frontal systems

65 (Fig. 1). A primary prerequisite for the existence of a strong ACC is an unobstructed latitudinal band of (deep ocean) water (Orsi et al., 1995; Barker and Thomas, 2004; Toggweiler et al., 2006), which is largely determined by the tectonic evolution and opening of the Tasmanian Gateway as well as the Drake Passage (Huber et al., 2004).

The tectonic evolution of the Tasmanian Gateway is relatively well constrained; early southern opening of the Tasmanian Gateway started around 49–50 Ma (Huber et al., 2004; Stickley et al., 2004; Bijl et al., 2013) with a change in course of tectonic drift of Australia from the Northeast to the North (Whittaker et al., 2007). Final breakup between Australia and Antarctica started around ~35.5 Ma, with ocean crust formation between southwestern Tasmania and Wilkes Land, Antarctica, and onset of bottom-water currents around 35.5–33.5 Ma (Stickley et al., 2004; Houben et al., 2019), although the strength of this socalled 'proto-ACC' during the Oligocene remains debated. New field data reconstructing \$ST\_and water properties (Bijl et al.,

- 75 2018; Hartman et al., 2018; Salabarnada et al., 2018; Evangelinos et al., 2020, 2022; Sauermilch et al., 2021; Hoem et al., 2021a, b, 2022; <u>Duncan et al., 2022;</u> Hou et al., 2022) allow tracing the migration of frontal systems, which may be reconducted to opening of gateways. Furthermore, <u>high resolution modelling exercises (England et al., 2017; O'Brien et al., 2020; Sauermilch et al., 2021; Nooteboom et al., 2022), show a large effect of the opening and depth of the Southern Ocean gateways on the Australian-Antarctic Gulf oceanographic conditions. A recent SST compilation from around the Tasmanian Gateway</u>
- 80 (Hoem et al., 2022) shows that the latitudinal SST gradient between the subtropical front and the Antarctic Margin progressively increased from ~26 Ma onwards, due to cooling at Antarctic-proximal sites. Since the Tasmanian Gateway was already open, wide, and deep, this Antarctic cooling may have been related to the onset of deep ocean connections through the Drake Passage (Lyle et al., 2007; van de Lagemaat et al., 2021). The plate tectonic configuration of the Drake Passage-Scotia Sea region is more complex than that of the Tasmanian Gateway, with various oceanic basins that opened at different times,
- 85 and the separation of continental fragments from southernmost South America and from the Antarctic Peninsula (Kennett, 1977; Barker and Thomas, 2004; Maldonado et al., 2006; Lagabrielle et al., 2009; Pérez et al., 2019; 2021). The timing and nature of the opening, widening, and deepening of the Drake Passage has also been much debated, and placed between 50 Ma and 16 Ma (Barker et al., 2007; Livermore et al., 2007; Eagles and Jokat, 2014; Maldonado et al., 2014; Pérez et al., 2021; van de Lagemaat et al., 2021). While some deep ocean sedimentary records suggest oceanographic rearrangements occurred that
- 90 were possibly linked to an early Drake Passage opening (Kennett, 1977; Scher and Martin, 2006; López-Quirós et al., 2019; López-Quirós et al., 2021), we lack knowledge of the long-term evolution of South Atlantic oceanographic conditions during this time. By reconstructing the development of the South Atlantic SST gradient, we can interpret phases of Antarctic cooling.

_	
D	eleted: gradients
D	eleted: (~16°C, between ~45–60°S)
D	eleted: between
D	eleted: gradients in
D	eleted: and temperature

Deleted: sea surface temperatures ( Deleted: )

## Deleted: our understanding of the Deleted:

eleteu:

**Deleted:** has been improved through modelling exercises (England et al., 2017; O'Brien et al., 2020; Sauermilch et al., 2021; Nooteboom et al., 2022)

(	Deleted: history of the
(	Deleted: of SST gradients in the
(	Deleted: development
~(	Deleted: could be used to

strengthening of the ACC<sub>4</sub> and shifts in the frontal systems. In turn, the changes in these three processes can be linked to the 110 throughflow of surface and deep waters through the Drake Passage.

Recent drilling efforts in and around the Scotia Sea as part of International Ocean Discovery Program (IODP) Expedition 382

Deleted: Deleted: which are in turn

Deleted: during

(Weber et al., 2021a), and a revisit of previously drilled records in the Weddell Sea during ODP Leg 113 offer improved spatial coverage of sedimentary records across two prominent climate transitions, viz. (1) the Eocene-Oligocene transition (EOT; 33.7 Ma) and (2) the transition from the Miocene Climate Optimum (MCO<sub>3</sub>~16.5 Ma) to the mid-Miocene Climate Transition (MMCT, ~14.7–13.8 Ma) and the late Miocene Cooling (LMC; ~7–5.4 Ma). Both transitions are marked by increases in benthic foraminiferal δ<sup>18</sup>O, suggesting cooling and expansion of the Antarctic ice sheet (Rohling et al., 2022). Given the paucity of carbonaceous sediments in this region, typically employed for paleotemperature reconstructions, we here choose to generate lipid biomarker (TetraEther<u>indeX<sub>2</sub>of tetraethersconsisting of 86 carbon atoms (TEX<sub>86</sub>)) proxy SST reconstructions, based on 120 isoprenoid glycerol dialkyl glycerol tetraether (isoGDGT) distributions from the northern Weddell Sea at ODP
</u>

Site 696 (Late Eocene–Early Oligocene) and southern Scotia Sea IODP Site U1536 (mid–late Miocene) (Fig. 1). We compare our findings to available TEX<sub>86</sub>, alkenone unsaturation index  $(U^{k'}_{37})_{\underline{x}}$  and clumped isotope ( $\Delta_{47}$ ) derived SST records from the South Atlantic region (Fig. 1) for a reconstruction of paleoceanographic conditions <u>over the late Cenozoic</u>.

Deleted: indeX	
Deleted: 3	
Deleted: x	
Deleted: with	
Deleted: ,	
Deleted: L	



#### 135

Figure 1. Present day map of the Southern Ocean showing the location of the drill sites used in this study. Grey areas represent present-day land masses. The colors show average Summer (January) SSTs from 1971\_2000 (Reynolds et al., 2002). The white areas lack modern SST data, as they are covered by ice shelves, The white lines represent the smoothed, simplified position of circumpolar fronts interpreted by Orsi et al. (1995). From north to south: The Subtropical front (STF); the Subantarctic front (SAF); the Polar 140 Front (PF); the southern ACC Front (sACCf); and the Southern Boundary (SBdy) front. SI=Seymour Island, SD=SHALLDRILL, WS= Weddell Sea, SS=Scotia Sea, <u>SOM= South Orkney Microcontinent</u>.

### 2 Material

2.1 Sedimentary drill cores

## 2.1.1 Site 696 lithology, age model and depositional setting

145 ODP Leg 113 Site 696 was drilled in the northern Weddell Sea (61°50.959'S: 42°55.996'W), on the South Orkney Microcontinent (Fig. 1), located south of the Southern Boundary (SBdy) front. The site had a late Eocene paleolatitude of ~67°S (paleolatitude.org Version 2.1; van Hinsbergen et al. (2015), using the paleomagnetic reference frame of Torsvik et al. (2012), and the geological reconstruction of Seton et al. (2012)).

Deleted: -	
Deleted: records	
Deleted:	

Deleted: SOM;

	Lithological descriptions and age constraints are gathered from López-Quirós et al. (2021; Fig. 2, Supplementary Table 1).
155	The age model is based primarily on calcareous nannofossil biostratigraphy (Wei and Wise Jr, 1990; Villa et al., 2008), and
	updated age constraints from organic walled dinoflagellate cysts (dinocysts) published previously (Houben et al., 2013;
	Houben et al., 2019; López-Quirós et al., 2021), which places the studied section of 607.6-548.9 mbsf (Cores 59R-53R) at
1	36.0 to 33.2 Ma (Houben et al., 2013, Supplementary Table 1). In the sediments overlying the lower Oligocene interval, 532-
l I	529.8 mbsf (Cores 52R-51R), no specific age constraint was determined, but dinocyst analysis indicate that the sediment is of
160	Oligocene age. Core 50R was initially dated to 14.3-14.8 Ma (Barker et al., 1988; Gersonde and Burckle, 1990), however, the
	diatom biostratigraphic scheme was adjusted and ages were updated to 17.6 - 15.4 Ma (Carter et al., 2017). More recently a
	thorough review of age-deterministic diatoms (Denticulonsis maccolummii and Actinocyclus ingens) have narrowed the Carter

et al. (2017) age interval of Core 50R to 16.7–16.5 Ma (López-Quirós et al., 2018).

Deleted: -

Deleted: -	
Deleted: In	
Deleted: ,	
<b>Deleted:</b> age deterministic diatoms ( <i>Denticulopsis maccolun</i> and Actinocyclus ingens)	ımii
Deleted: ere	
Deleted: found and	
Deleted: ere	
Deleted: the middle Miocene (	
Deleted: ;	
Deleted:	
Deleted: or more accurate:	
Deleted: ;	
Deleted: .	
Deleted: in prep	
Formatted: Not Highlight	





~~~~(	Deleted: in prep
	Deleted: .
	Deleted: ;
	Deleted: to
	Deleted: to

rhythmically interbedded sandy mudstone facies with glauconite bearing sandstone beds (548.9<u>529.8 mbsf), and 5) pelagic</u> 6

The studied sediment package at Site 696 (607.6–521.08 mbsf) consists of 1) organic-rich sandy mudstone facies (607.6–606.9 mbsf), 2) glaucony-bearing packstone facies (606.9 to 569.7 mbsf), 3) claystone and limestone facies (569.7 548.9 mbsf), 4)

sediments, predominately biosiliceous diatom ooze (522–521 mbsf) (López-Quirós et al., 2019; 2020; 2021). The glauconitic
packstone beds of latest Eocene age (606.9\_569.7 mbsf, ~35.5–34.1 Ma) are attributed to a decline in terrigenous input to Site
696 and increased winnowing (López-Quirós et al., 2019). The glauconitic packstone beds of latest Eocene age (606.9 to 569.7 mbsf, ~35.5–34.1 Ma) are attributed to a decline in terrigenous input to Site 696 and increased winnowing (López-Quirós et al., 2019). The glauconitic packstone beds of latest Eocene age (606.9 to 569.7 mbsf, ~35.5–34.1 Ma) are attributed to a decline in terrigenous input to Site 696 and increased winnowing (López-Quirós et al., 2019), as a result of the Powell Basin at the tip of the Antarctic Peninsula opening and incipient South Orkney Microcontinent subsidence. However, the Antarctic-proximal location makes the site prone to influx of terrigenous material
transported by icebergs, as shown by the unequivocal evidence of iceberg rafted debris (IRD) deposited across the EOT (34.1–

33.6 Ma, 576–568 mbsf) at Site 696 (López-Quirós et al., 2021). Across the EOT, sediments at Site 696 reflect an increasingly distal and deeper environment as the South Orkney Microcontinent continued to deepen (López-Quirós et al., 2021). Eutrophic surface water conditions, indicative of productive and somewhat shallow-water and reduced-oxygen conditions, are indicated by dinocysts (Houben et al., 2019) and sedimentary facies (López-Quirós et al., 2019; 2021). The dominance of large sized

- 205 heterotrophic Protoperidiniacean dinocysts in the earliest Oligocene is suggested to reflect seasonal sea-ice <u>coverage</u> (Houben et al., 2013). During the earliest Oligocene (~33.6–33.2 Ma), the <u>South Orkney Microcontinent shelf subsided, and biological</u> production increased, partially driven by upwelling along the shelf (López-Quirós et al., 2021). Deposition of moderately to intensely bioturbated silty-mudstones during the EOT is attributed to the continued subsidence-related deepening at Site 696 (López-Quirós et al., 2021). Above the claystone/clayey mudstones of the lowermost Oligocene (Core 53R, Fig. 2) we find
- 210 rhythmically interbedded sandy mudstones with glauconite-bearing sandstone beds, a result of reworked sediments, deposited under bottom current activity and possibly slumping (López-Quirós et al., 2020). The nature of this sediment complicates dating of this material. There is likely a break in the sedimentation (hiatus) around 529 mbsf (the top part of Core 51R), with a sharp contact from glauconitic-bearing sandstone to mud bearing diatom ooze 522–521 mbsf (Core 50R), dated to ~16.7–16.5 Ma (López-Quirós et al., 2018).

#### 215 2.1.2 Site U1536: lithology, age model and depositional setting

Site U1536 is located in the Dove Basin, in the southern Scotia Sea (59°26.4608'S, 41°3.6399'W, 3220 m water depth). The site was drilled to study the Neogene flux of icebergs through "Iceberg Alley", the main pathway along which icebergs calved from the margin of the Antarctic ice sheet drift into the warmer waters of the ACC (Weber et al., 2021a). Today, the site is located just south of the southern ACC through the Southern Boundary (SBdy) front and is seasonally covered by

- 220 sea ice (Fig. 1). The rotary drilling at Hole U1536E penetrated down to 643 mbsf. Sediments have moderate to high core disturbance and biscuiting₂ or brecciated core material due to the rough nature of rotary drilling and compaction of gravel rich material. The lithology of the studied interval from Hole U1536E₂ 640–450 mbsf (Cores 33R–13R)₂ consists of silty clays with interbedded diatom ooze (Fig. 3), with an estimated average sedimentation rate of ~4.3 cm/kyr (3.9–6.4 cm/kyr; Pérez et al., 2021). The shipboard bio- and magnetostratigraphic age model was used to date the sediments (Weber et al., 2021b,
- 225 Supplementary Table 2). Sediments between 480–450 mbsf (Cores 16R–13R) have an age of 6–5 Ma based on bio- and magnetostratigraphy. Diatom biostratigraphy between 548–535 mbsf (Cores 24R–22R) indicate ages of 7.7–6.4 Ma. The

7

#### Deleted: ; in prep

Deleted: to

Deleted: The opening of the Powell Basin (at the tip of the Antarctic Peninsula) caused subsidence of the South Orkney Microcontinent, thereby decreasing the delivery of terrigenous sediments to the site (López-Quirós et al., 2019). The Antarcticproximal location makes the site prone to influx of terrigenous material transported by icebergs. Indeed, López-Quirós et al. (2021) show unequivocal evidence for the oldest iceberg rafted debris (IRD) deposited across the EOT (34.1–33.6 Ma, 576–568 mbS) at Site 660.

#### Formatted: Font colour: Text 1

**Deleted:** Across the EOT sediments reflect more distal and deeper environments while the SOM continued to deepen (López-Quirós et al., 2021).

#### Formatted: Font colour: Text 1

Deleted: conditions

Deleted: SOM

#### Deleted: in prep

Deleted: F

sediment directly overlying Reflector-c (Weber et al., 2021b), at 617–570 mbsf (Cores 30R–26R), have an age of 8.4 Ma 245 (Pérez et al., 2021). The sediments below Reflector-c, at 622 mbsf (Core 31R), are dated to ~14.2 Ma, and as the lithologic contact is not recovered, Reflector-c could represent a prolonged time interval of slow sedimentation rates or non-deposition or erosion. The sparse brecciated lithology fragments in the lower cores, below the Reflector-c (566 mbsf), consist of lithified mudstone and gravel-conglomerate-breccia (Weber et al., 2021b; Perez et al., 2021).



#### 250

260

Figure 3. Site U1536 age model and lithology modified after the IODP Expedition 382 shipboard report (Weber et al., 2021b). The depth and age of the stratigraphic discontinuities (seismic reflectors) are derived from Perez et al. (2021) (Supplementary Table 2, Perez et al., 2021). LO = Last occurrence, FO = First occurrence.

#### 3 Methods

#### 255 3.1 Lipid Extraction and glycerol dialkyl glycerol tetraether (GDGT) analysis

Lipid extraction of sediments from Site 696 was performed at the Laboratoire d'Océanographie et du Climat, Expérimentations et Approches Numériques (LOCEAN-Sorbonne Université, Paris, France). First, 71 sediment samples were freeze-dried and crushed to a fine powder. Total lipids were extracted from ~9.5 to 15 g of homogenized sediment using a solvent mixture of 40 ml dichloromethane:methanol (DCM:MeOH; 3:1, v/v). The applar fraction was separated from the polar lipids by passinfpassing the total lipid extract (TLE) over a silica column using 3 ml hexane as eluent, followed by the recovery of the

Moved (insertion) [1]	
Deleted: DCM:MeOH (	
Deleted: 1	
Deleted: f	
Deleted: . T	

Deleted: it means that

Deleted: s

polar fraction by cluting with 3 ml DCM:MeOH (1:1, v/v), The polar lipid fractions were sent to Utrecht University for GDGT analysis. Sediment samples from Site U1536E were processed for GDGT analysis by lipid extraction at Utrecht University, from 10 g freeze-dried and manually powdered sediments using a Milestone Ethos X microwave system and adding

- 270 DCM:MeOH (9:1, v/v). The TLEs were first filtered through a NaSO4 column to remove potential remaining water and sediments. The TLEs were then separated on an activated Al<sub>2</sub>O<sub>3</sub> column into apolar, ketone and polar fractions, using hexane: DCM (9:1, v/v), hexane: DCM (1:1, v/v), and DCM: MeOH (1:1, v:v) as eluents, respectively. All polar fractions were dried under N<sub>2</sub>. A known amount of C<sub>46</sub> GTGT standard was added to the polar fractions from Sites 696 and U1536, which were, subsequently dissolved in hexane: isopropanol (99:1, v/v) to a concentration of  $\sim 2 \text{ mg ml}^{-1}$  and filtered through a 0.45  $\mu$ m
- 275 polytetrafluorethylene filter. After that, the dissolved polar fractions were injected and analysed, by ultra-high performance liquid chromatography/mass spectrometry (UHPLC/MS) according to the method described by Hopmans et al. (2016), using an Agilent 1260 Infinity UHPLC system coupled to an Agilent 6130 single quadrupole mass detector, at Utrecht University. Selected ion monitoring (SIM) was used to identify the GDGTs using their [M+H]<sup>+</sup> ions and integrated using ChemStation software. Samples with very low concentrations (i.e., peak area <3,000 mV/s and/or peak height <3× background signal) of 280 any of the GDGTs included in the TEX86 were excluded from analysis.

Although lipid extractions of Sites 696 and J1536 were performed at different institutions, the latest interlaboratory comparison study (F.Peterse personal communication) that assessed the repeatability and reproducibility of the TEX<sub>86</sub> indicates that differences in sediment extraction and workup procedures do not affect isoGDGT distributions, and thus reconstructed

285 SSTs. Instead, variations in reported TEX<sub>86</sub> values appeared to be mainly introduced by the type of mass spectrometer used for GDGT analysis (Schouten et al., 2013), Since the polar fractions from both sites were analyzed using the same HPLC-MS instrument at Utrecht University, the uncertainty on our TEX<sub>86</sub>-based SSTs mostly represents the analytical uncertainty, which is ±0.3°C based on long-term observation of the in-house standard

#### 3.2 GDGT indices for non-thermal overprints on TEX<sub>86</sub>

290 The TEX<sub>86</sub> SST proxy is based on the temperature dependence of the number of cyclopentane rings in GDGT membrane lipids produced by marine Thaumarchaeota and calculated as defined by Schouten et al. (2002):

 $TEX_{86} = ([GDGT-2]+[GDGT-3]+[cren'])/([GDGT-1]+[GDGT-2]+[GDGT-3]+[cren'])$ 

295 The use of TEX86 as a proxy for SST relies upon the assumption that isoGDGTs in marine sediments are principally derived from membrane lipids of marine pelagic Thaumarchaeota (Schouten et al., 2013). However, in some environments, nonthermal factors may alter the distribution of isoGDGTs stored in the sediment and thus the temperature signal (Supplementary information). We assess potential non-thermal effects on isoGDGT distributions prior to translating TEX<sub>86</sub> into SSTs. We use the branched and isoprenoid tetraether (BIT) index to assess possible overprints of terrestrial GDGT input (Hopmans et al.,

Moved up [1]: lipids over a silica	The aploar fraction was separated from the polar column using 3 ml hexane as eluent.
Deleted: was	
Deleted: ed fron	n the total lipid extracts (TLEs) using
Deleted: over a	silica column
Deleted: further	
Deleted: dichlor	omethane:methanol (
Deleted: ;	
Deleted: dichlor	omethane (DCM)
Deleted: as	
Deleted: z	
Deleted: analyze	ed.

Deleted:	The Site 696
Deleted:	were generated at LOCEAN
Deleted:	those for Site
Deleted:	Utrecht University. An
Deleted:	revealed
Formatte	ed: Subscript
Deleted:	prenoidal
Deleted:	relative abundances
Deleted: a single co biased by t	As the $TEX_{86}$ is based on specific GDGT ratio and not or mpound concentration, the estimated SSTs is therefore no he extraction/separation method performed in the two lab
Formatte	ed: Subscript
Deleted:	
Deleted:	analysed
Formatte	ed: Subscript
Deleted:	can be considered negligible
Deleted:	can be considered negligible

(1)

Deleted: prenoidal

2004), the fractional abundance of the crenarchaeol isomer over that of crenarchaeol (fcren'; O'Brien et al., 2017), the methane index to identify contributions of methanotrophic archaea (MI; Zhang et al., 2011) or methanogenic archaea using the ratio of GDGT-0/crenarchaeol (Blaga et al., 2009), the GDGT-2/GDGT-3 ratio to assess contributions of a deep dwelling GDGT

producing community (Taylor et al., 2013) and the ARing Index to identify GDGT distributions that deviate from modern

330

## analogues ( $\Delta$ RI; Zhang et al., 2016) (Fig. S1C, panel 2–8 and Fig. S2C, panel 2–8, <u>Supplementary Table 3,4</u>).

## 3.3 TEX<sub>86</sub> calibration

The empirical relationship between TEX<sub>86</sub> values and SST has appeared to be not always straightforward, as reflected by continued revisions of the approach of TEX<sub>86</sub> – SST calibrations (Kim et al., 2010; Tierney and Tingley, 2015; Ho and Laepple, 2016; O'Brien et al., 2017; Dunkley Jones et al., 2020). Particularly, the relationship between TEX<sub>86</sub> and SST seems to become obscured in both extreme ends of the core-top calibration: at or above modern SSTs, and in cold polar regions. However, for the time intervals (Late Eocene–Early Oligocene and Middle–Late Miocene) and locations we are targeting, we expect SSTs within the intermediate temperature range. In this study, we used the regionally varying BAYSPAR SST calibration of Tierney

- 340 and Tingley (2015) to reconstruct SST (±4°C standard calibration error) from TEX<sub>86</sub> values. BAYSPAR compares measured TEX<sub>86</sub> values to modern TEX<sub>86</sub> values obtained from surface sediments, to derive linear regression parameters, and propagates uncertainties in the surface sediment data into resulting temperature predictions (Tierney and Tingley, 2015). Even when GDGT-2/GDGT-3 ratios indicate that deeper dwelling GDGT producers do not contribute to the sedimentary signal, the GDGTs could still originate from the subsurface (50–200 m water depth) rather than the sea surface (Schouten et al., 2013).
- 345 However, since the BAYSPAR calibration translates TEX<sub>86</sub> values to SSTs, we will, therefore, refer to the proxy results as such in the remainder of this work. For all new and existing TEX<sub>86</sub> records discussed in this study, we applied a standard deviation of ±20°C and a prior mean of 20°C for the Late Eocene–Early Oligocene interval and 15°C for the Miocene. We also compare the BAYSPAR-derived SST estimates with those based on the exponential function from Kim et al. (2010) and the linear function by O'Brien et al. (2017). The SST records all show similar trends, but BAYSPAR-derived SSTs are usually
  350 cooler compared to those obtained from the functions of Kim et al. (2010) and O'Brien et al. (2017) and can thus be considered conservative estimates (Supplementary Table 3, 4, Fig. S3, S4).

#### 4 Results

### 4.1 Site 696 GDGT distributions and TEX<sub>86</sub>-SST trends

355

A total of 71 samples from ODP Site 696 were analysed for TEX<sub>86</sub> palaeothermometry (Supplementary Table 3). The GDGT pool consist of 90  $\pm$  5% isoGDGTs and 10  $\pm$  5% brGDGTs, resulting in BIT index values <0.2 (Fig.5 b). The isoGDGT distributions indicate that GDGTs are primarily <u>derived from</u>surface<sub>r</sub>dwelling Thaumarchaeota (Supplementary information, Fig. S1). In total, 9 samples had higher  $\Delta$ RI values than the cut off of 0.3 (Fig. S1B) indicating a potential non-thermal overprint

Deleted: W Deleted: used Formatted: Subscript

Deleted: produced by Deleted: ocean

on the GDGT distribution and are excluded from SST analysis (triangles in Fig. 4). The TEX <sub>86</sub> -based SST record from Site	
$696$ (607–521 mbsf, 36–33 Ma and ~16.5 Ma, Fig. 4) thus consists of 62 data points, and predominantly ranges between $12^{\circ}C$	
and 18°C, although the total temperature range is from $4^{\circ}$ C to $25^{\circ}$ C (± $4^{\circ}$ C standard <u>calibration</u> error). There is a general	

365 cooling trend of ~8°C between the Upper Eocene and the Lower Oligocene (from 20°C to 12°C between 610 and 558 mbsf) interval. This is followed by an average ~5°C increase in temperature between ~558 and ~552 mbsf. In the organic rich interval at ~555 mbsf (Core 53R, section 5, 33.5–33.2 Ma), 35 sediment samples were analysed, with J sample at every 3 cm (~15 kyrs, resolution). In this interval, the high resolution TEX<sub>86</sub>-SST record shows high-amplitude variability (total range; 4–21°C, with most datapoints falling within the range between 8–17°C). At ~550 mbsf (Lower Oligocene) SSTs rapidly decreases to 10°C.

370 In the Middle Miocene (~520 mbsf, n=2) SSTs values are ~14°C.

Deleted: distribution, and

Deleted: , but with high sample-to-sample variability
Deleted: ssediments
Deleted: z
Deleted:
Deleted: analyzed
Deleted: at high resolution (
Deleted: ,
Deleted:
Deleted: )
Deleted: T
Deleted:



Figure 4. a) TEX86-SST data (BAYSPAR calibration; Tierney and Tingley, 2015). Datapoints marked by a triangle represent samples (n=9) with a potential non-thermal overprint on the GDGT distribution (outliers) (Supplementary information), b) BITindex values (Hopmans et al., 2004) plotted next to the lithology and age constraints of Site 696, which are modified after Lopez-Quiros et al. (2021), based on Barker et al. (1988) and Lopez-Quiros et al. (2019, 2020), including new constraints of the uppermost interval (Core 50) from López-Quirós et al. (2018).

Deleted: in prep.

### 390 4.2 Site U1536 GDGT distributions and TEX<sub>86</sub>-SST trends

395

A total of 40 sediment samples from IODP Hole U1536E were processed for GDGT analysis (Supplementary Table 4), of which 14 had GDGT concentrations below detection limit (Fig. 5). The Site U1536 GDGT pool consists of variable amounts of both isoGDGTs and brGDGTs, where brGDGTs are relatively more abundant in the middle (500–560 mbsf) and top parts of the record (450 mbsf) (Fig. S2A), resulting in high BIT-index values in these intervals of the record (Fig. 5b). GDGT distributions in 12 sediment samples were outside the range of what is considered reliable for multiple indicator proxies (triangles in Fig. 5a, Supplementary information, Fig. S2). For the remaining 14 samples TEX<sub>86</sub> values were translated into SSTs using the BAYSPAR calibration (Fig. 5a). The obtained record shows temperatures of 5–11°C for the Middle Miocene (619–640 mbsf) and 1.5–5°C for the Upper Miocene (570–450 mbsf).

- Deleted: middle and top





400 Figure 5. Biomarker indices from Site U1536. Left panel shows core recovery, estimated age and lithological composition (Weber et al., 2021a). The crosses (X) indicate samples with GDGT concentrations below detection threshold. a) TEX<sub>86</sub>-SST (BAYSPAR

calibration; Tierney and Tingley, 2015). Triangles mark samples with a potential non-thermal overprint on the GDGT distribution 405 (outliers) (Supplementary information), b) BIT-index (Hopmans et al., 2004) values.

#### 5 Discussion

#### 5.1 Late Eocene-Early Oligocene South Atlantic SST conditions

We have compiled available South Atlantic SST records from Site 1090 (Lie et al., 2009), Site 511 (Houben et al., 2019) and Seymour Island (Douglas et al., 2014), in addition to our new SST record from Site 696 (Fig. 6) to put the SST records into a 410 broader regional context and discuss the surface oceanographic development during the Late Eocene-Early Oligocene. Our SST record from Site 696 (yellow in Fig. 6A) shows warm-temperate conditions (SST range: 22-14°C) during the latest Eocene (~36.5-33.6 Ma) and on average decreasing SSTs (~15-9°C) in the earliest Oligocene (33.6-33.2 Ma). The cooling of South Atlantic surface waters across EOT is in broad agreement with the average Southern Ocean-wide temperature drop (Kennedy-Asser et al., 2020; Tibbett et al., 2023), the increase in benthic foraminifer  $\delta^{18}$ O values as a result of deep sea cooling. 415 a drop in atmospheric pCO<sub>2</sub> levels (https://www.paleo-co2.org; Pearson et al., 2009; Steinthorsdottir et al., 2016; Hoenisch, 2021) and the growth of a continent-wide Antarctic ice sheet across the EOT (e.g. Bohaty et al., 2012). Sites 511 and 1090 also show a step-wise cooling across the EOT (Hutchinson et al., 2021), where the first step occurs around 34.1 Ma and coincides with common IRDs at Site 696, indicating the onset of marine-terminating glaciers in the region (López-Quirós et al., 2021), and the second step coincides with the Earliest Oligocene Oxygen Isotope Step (EOIS, 33.65 Ma; Hutchinson et al.,

- 420 2021). Miospores at Site 696, believed to be of local origin from the South Orkney Microcontinent, changed concomitantly from southern beech, Nothofagus-dominated vegetation to an increase in gymnosperms and cryptogams, accompanied by a rapid rise in taxon diversity after the EOIS (~33.65 Ma, 568.82 mbsf; Thompson et al., 2022). This shift in vegetation to a cooler and dryer climate occurred after the onset of earliest glacial expansions (~34.1 Ma). Sedimentological investigations by López-Quirós et al. (2021) for the same interval showed deepening of the South Orkney Microcontinent shelf and enhancement
- 425 of biological production, possibly due to upwelling along the shelf, leading to low oxygen conditions at the seafloor. The highamplitude SST variability (~4-8°C) in our TEX<sub>86</sub>-SST record would suggest that this upwelling regime was strongly variable. The variability in upwelling conditions could be induced by strong fluctuating ice sheet expansion and retreat and shifts in wind patterns and ocean frontal systems. Low-resolution palynological investigations on Late Eocene - Early Oligocene sediments from the southern South Atlantic (Houben et al., 2019; Hoem et al., 2022) show a highly diverse and variable
- 430 dinocyst assemblage, which includes, Antarctic derived, open ocean, temperate and high nutrient indicative species. respectively, and indeed infers a fluctuation in surface ocean conditions, potentially related to shifts in frontal systems and upwelling regions. Alternatively, the high variability in the TEX<sub>86</sub> signal could be introduced by the input of reworked isoGDGTs. Such inputs were previously found to be high at ice proximal sites where the onset of large-scale Antarctic glaciation across the EOT caused reworking of pre-Eocene deposits, such as e.g., Prydz Bay (Tibbett et al., 20201). However 435 Tibbett et al. (2021) found that this had little impact on the overall TEX<sub>86</sub>-SST trend. Furthermore, we record very low BIT

#### Deleted: with

Deleted: or...Site 696 (yellow in Fig. 6A) shows warm-temperate conditions (SST range: 22-14°C) during the latest Eocene (~36.5-33.6 Ma) and on average decreasing SSTs (~15-9°C) in the earliest Oligocene (33.6-33.2 Ma). The cooling of South Atlantic surface waters across EOT is in broad agreement with the average Southern Ocean-wide temperature drop (Kennedy-Asser et al., 2020: Tibbett et al., 2023), the increase in benthic foraminifer  $\delta^{18}$ O values as a result of deep sea cooling (Bohaty et al., 2012) (11)Deleted: across the EOT Moved down [2]: Step-wise cooling across the EOT (Hutchinson

et al., 2021) is evident at

Deleted: in the Uk'37 record at

#### Moved (insertion) [2]

Deleted: S...tep-wise cooling across the EOT (Hutchinson et al., 2021), where is evident at. T...he first step occurs around 34.1 Ma and coincides with common IRDs at Site 696, suggesting ... [2] Deleted: tion

h	Deleted: onsetn the region (López-Quirós et al., 2021), a [3]				
	Deleted: Earliest Oligocene Oxygen Isotope Step (OIS (, ~33.65 Ma, 568.82 mbsf; ) (hompson et al., 2022). This shift in vegetation to a cooler and dryer climate occurred after the onset of earliest glacial expansions (~34.1 Ma). and corresponds to our SST drop below 12°C and the more common IRD in the sediment at this depth (see also López-Quirós et al., 2021). ([4]				
	Deleted:n our TEX <sub>86</sub> -SST record (~4-8°C)ould suggest that this upwelling regime was strongly variable. The variability in upwelling conditions could be induced by strong fluctuating ice sheet expansion and retreat, and ([5])				
ļ	Deleted: Southern South Atlantic 1				
ķ	Deleted: er ( [6])				
Å	Deleted: Late Eocene-Early Oligocene				
λ	Deleted:ocene ([7])				
Å	Deleted: S				
1	<b>Deleted:</b> showsa highly diverse and variableityin the dinocyst assemblages with varying relative abundance which				

includesing ... Antarctic derived, open ocean, temperate and high 

### Formatted: Subscript

Deleted: could be affecting the TEX<sub>86</sub> signal... Such inputs were previously found to be high at ice proximal sites where T...he onset of large-scale Antarctic glaciation across the EOT caused reworking of pre-Eocene deposits, and input of reworked isoGDGTs were found to be high at ice proximal sites [10] .... Deleted: atas Deleted: t

#### Deleted: the

Formatted: Subscript

	index values (<0.2, Supplementary Table 3, Fig 4b) throughout the record, whereas Eocene sediments around Antarctica are		Deleted:
	commonly high in brGDGTs (e.g., Bijl et al., 2013), because of the vell-developed soils on Antarctica at the time (Inglis et	(	Deleted: abiquitous
	al., 2022). The lack of brGDGTs in our record thus suggests little influence of reworked Eocene GDGTs. We, therefore,	(	Deleted: ubiquitous
	assume that the TEX <sub>86</sub> record from Site 696 represents an in-situ pelagic signal and that the variability in the SST record is	(	Deleted: recurd
545	introduced by unwelling	$\langle \rangle \langle \langle$	Deleted: indicating
545		$\mathbb{N}/\mathbb{X}$	Deleted: soil-derived
	Ι	X/N	Formatted: Subscript
		$\langle \rangle \rangle \langle \rangle$	Deleted: to be recorded at Site 696

## Deleted:

Formatted: English (UK)

## Deleted: ¶

Formatted: English (UK)



The clumped isotope ( $\Delta_{47}$ )-based SST datapoint from Seymour Island (34 Ma, purple <u>star</u> in Fig. 6B) shows a similar temperature (13°C) to that derived from the TEX<sub>86</sub> proxy at the same site (Douglas et al., 2014), as well as from nearby Site 2

- 570 <u>696</u>. The correspondence of the biomarker-derived SSTs with that from *Eurhomalea* (bivalve) △47 (mean annual temperature), adds reliability to the temperature proxies accurately reflecting SST in this region. Interestingly, the SSTs from Seymour Island and Site 696 are very similar to Site 511, even though there was a paleolatitudinal difference of ~12° between Site 511 and Seymour Island (Fig. 6). The subtropical Site 1090 is the warmest site (~19–27°C) in our compilation, which is expected given its lower paleolatitude. However, the absolute SSTs of Sites 696 and 511 are strikingly similar across the EOT (Fig. 6A). The
- 575 southwestern South Atlantic temperature gradient (between Sites 511 and 696) decreased from ~1.8°C in the Late Eocene to ~0°C in the Early Oligocene. A larger throughflow through Drake Passage would increase the temperature gradient between Sites 696 and 511. Today both sites are separated by the strong ACC and an SST gradient of >7°C (Locarnini et al., 2018). We therefore imply that the throughflow changes induced by the opening Drake Passage did not change the South Atlantic Ocean circulation across the EOT. Tectonic evidence suggests that the Drake Passage was narrow, with little deep-water
- 580 connection from the Pacific to the Atlantic around the EOT (Livermore et al., 2007; Eagles and Jokat, 2014; van de Lagemaat et al., 2021), which may explain the lack of regional oceanographic response, Model experiments (Huber et al., 2004; Hill et al., 2013; England et al., 2017; Sauermilch et al., 2021) show that a Southern Ocean without deep gateways featured winddriven clockwise gyres in the South Pacific and South Indian/Atlantic Ocean basins (Fig. 6B) that would advect warm surface waters toward the Antarctic coast. Specifically, eddy resolving ocean model simulations by Sauermilch et al., (2021) for the
- 585 Eocene show that a restricted (depths <600 m) Drake Passage throughflow would sustain the subpolar gyre and lead to SSTs reaching 19°C in the Australian-Antarctic Basin and 15–17°C in the subpolar Pacific and Atlantic. This is very similar to our Site 696 TEX<sub>86</sub>-SST EOT record (~11–18°C). We thus propose that the small differences in SSTs between Sites 511 and 696 are the result of a restricted Drake Passage during the latest Eocene earliest Oligocene, facilitating a persistent wind-driven gyral circulation that connected the southern South Atlantic sites in our compilation.

#### 590 5.2 Middle to Late Miocene

To investigate the cooling step in the <u>Middle to Late</u> Miocene, the new Miocene SST records from the Antarctic-proximal southwest<u>ern South</u> Atlantic Sites 696 and U1536 are compared to records from the Antarctic Peninsula (SHALDRILL IL Core 5D; Tibbett et al., 2022) and Wilkes Land (Site U1356, Sangiorgi et al., 2018). However, there is still a lack of well constrained and overlapping records form the Southern Ocean Middle to Late Miocene due to glacial expansions and erosion causing discontinuous records. Due to the large age uncertainties and gaps in the in the sedimentary records of these sites (Weber et al., 2021a; Perez et al., 2021; Bohaty et al., 2011), we present the TEX<sub>86</sub>-derived SSTs as average temperatures within two broad time intervals corresponding to the age uncertainty. We also compare SST trends from the above-mentioned sites to those from the subantarctic zone; ODP Site 1171 from the southwest Pacific Ocean (Leutert et al., 2020) and subtropical front; ODP Site 1088 (Herbert et al., 2018) in the Southeast Atlantic, and Site ODP 1168 west of Tasmania (Hou et al., 2022).

Deleted: TEX<sub>86</sub>- and Uk'37-based SST reconstructions from ODP Site 511 differ slightly in trends across the EOT, with Uk'37 showing amplified cooling compared to TEX<sub>86</sub>. Uk'37-temperature data reflect surface water temperature following the niche of the haptophyte algae that produce the alkenones on which the Uk'37 is based, while the GDGTs that mostly contribute to the TEX<sub>86</sub> signal stored in the sediments are likely produced in the subsurface (50-200 m water depth) (Schouten et al., 2013). Although Uk'37 is calibrated to mean annual temperature (Müller et al., 1998), in some settings it reflects a seasonally biased temperature (e.g., Ternois et al., 1997). In fact, modern high-latitude alkenone producing haptophyte communities primarily bloom in early spring (Herbert et al., 1998). Thus, at Site 511, the amplified cooling in  $U^{k'_{37}}$  might reflect a shift towards early spring blooming, which could explain its amplified cooling trend across EOT, relative to that in TEX86. As TEX86-based SSTs are consistently higher than Uk' 37-based SSTs at this site, we argue that TEX<sub>86</sub> represents a warmer season than Uk'37 and that it does seem to be reflecting a surface water signal rather than a subsurface s ... [10]

## Deleted: the

Formatted: Font: Italic



845 Further, we compare our new <u>Site U1536</u> SST record to clumped isotope bottom water temperatures (BWT) from South Indian Ocean ODP Site 747 (Leutert et al., 2021) (Fig. 7).





Figure 7. A. TEX86-based SST data from Site 696 and U1536 (this study) compared to Southern Ocean wide SST and BWT records 850 The bars on the left indicate the standard calibration error of the SST proxies. Data of Site U1536 is displayed as two bar plots (red) showing the temperature ranges for the Middle Miocene (16-14 Ma) and late Miocene (7-5.3 Ma), individual data points are shown as red dots. We compare our data to SST records from Wilkes Land Site U1356 (Sangiorgi et al., 2018), SHALDRIL II Core 5D (Tibbett et al., 2022; grey bars indicate the age uncertainty), Site 1168 (west of Tasmania; Hou et al., 2022), Site 1171 (southwest Pacific Ocean; (Leutert et al., 2020), Uk'37- SST data from Site 1088 (Herbert et al., 2016) and clumped isotope bottom water 855 temperature (BWT) data from Leutert et al. (2021) (Site 747). The arrows indicate the temperature gradient between the TEX SST record from the southwest South Atlantic and the subtropical front Site 1168. The black line is the benthic for aminiferal  $\delta^{18}$ O compilation, smoothed by a locally weighted function over 20 kyr (thin blue curve) (CENOGRID; Westerhold et al., 2020). Thick blue curve is the LOESS smoothed (span = 0.2). The stippled vertical line indicates the age for the paleogeographic map below. B. Paleogeographic reconstruction at 16 Ma, based on the GPlates reconstruction of van de Lagemaat et al. (2021) in the paleomagnetic 860 reference frame of Torsvik et al. (2012). All sites from data compilation in A are shown as stars. Dashed black line represents the Miocene surface ocean currents derived from Herold et al. (2012). The data compilation of South Atlantic SSTs (Fig. 7A) shows a 7°C cooling between Site 696 (yellow dots, ~14°±4°C standard4 calibration error, n=2) during the Miocene Climate Optimum (MCO, ~16.5 Ma) and Site U1536 (red, ~7±4°C, n=5) in the Middle Miocene Climate Transition (MMCT, ~14.7-13.8 Ma). Both sites were located at comparable paleolatitudes (albeit 865 with a 2.5° latitudinal difference, Fig 8A) during the Miocene (Fig. 7B). SSTs at Wilkes Land Site U1356 were warmer (17°C

- around 17 Ma) than at Site 696 (~14°C), even though Site U1356 was situated closer to the cooler East Antarctic ice sheet. Additionally, a less pronounced cooling (SST ~16-12°C) occurred across the MMCT at Site U1356 (Sangiorgi et al., 2018), than the 7°C cooling shown in the South Atlantic low-resolution dataset. Thus, the South Atlantic sector was colder with a likely more proximal ice mass at the onset of MMCT than the Wilkes Land Antarctic margin. There was a ~6–10°C temperature
- 870 difference between the southwest South Atlantic Antarctic proximal Sites 696/U1536 and the subantarctic Site 1171 (southwest Pacific Ocean), with Site 1171 clumped isotope (A47) SSTs of 14-12°C and TEX<sub>86</sub> SSTs of 18-13°C during MMCT (Leutert et al., 2020). This suggests a relatively strong SST gradient between the coldest Antarctic-proximal regions and the subantarctic zone in the Middle Miocene, even though the subantarctic zone was likely situated at lower paleolatitudes in the Australian-Antarctic Gulf than in the South Atlantic due to the more southerly position of Australia. The increase in the temperature
- 875 gradient in the complied SST records (Fig. 7A) during the Middle Miocene indicates breakdown of the dominant gyral circulation at the early Oligocene and a subsequent strengthening of the ACC during the Middle-Late Miocene. The alkenonebased SST reconstructions for Site 1088 and TEX<sub>86</sub>-based SST reconstructions for Site 1168, together representing subtropical front conditions, show MCO temperatures between 32-27°C that progressively cooled during the MMCT (~24-14°C) (Hou et al., 2022). This subtropical cooling is weaker than at the Antarctic-proximal sites, which indicates that the cooling was
- 880 amplified at high latitudes. The clumped isotope ( $\Delta_{47}$ ) BWT record from Site 747 in the South Indian Ocean (Leutert et al., 2021) shows strikingly similar temperatures to the reconstructed SSTs at Site U1536, which may suggest that the Weddell Gyre in the southwestern South Atlantic was an important region of deep water formation in the Miocene, like today (e.g., Orsi et al., 1999), which furthermore is in line with what modelling studies suggest for the Miocene (e.g., Herold et al., 2012). The 7°C cooling of the southwestern South Atlantic between the MCO and MMCT occurs during a time of declining  $pCO_2$
- 885 (Foster et al., 2012; Greenop et al., 2014) (Fig. 8D) and increasing benthic foraminiferal  $\delta^{18}$ O (Fig. 7A, 8C), reflecting an increasingly colder climate with larger (in area) ice sheets on the Antarctic continent with oceanward expansion, (Lewis et al., 2008; Shevenell et al., 2008; Holbourn et al., 2018; Levy et al., 2019; Leutert et al., 2020, Given estimates of pCO2 decline (by 100–300 ppm; Sosdian et al., 2018; Super et al., 2018), paleo climate sensitivity ( $1.5-4.5^{\circ}$ C per pCO<sub>2</sub> doubling; Martínez-Botí et al., 2015) and polar amplification factors (2-3; Holland et al., 2003), the 7°C cooling in the southwestern South Atlantic 890 records (Sites 696 and U1536) could have been completely induced by  $pCO_2$  decline in the Middle Miocene. This also means

#### Deleted:

#### Deleted: P Deleted: is

Deleted: Deleted: For the location of all sites in the data compilation in A see Fig. 1.

#### Deleted: for

Formatted: Line spacing: At least 15 pt

Deleted: at a higher latitude (59°S) than Sites 696 and U1536 (~65°S)

#### Deleted: on average

Deleted: that our Deleted: shows

Deleted:

λ	Deleted: in
λ	Deleted: since
λ	Deleted: at
1	Deleted: at
λ	Deleted: albeit
1	Deleted: less
_(	Deleted: an
(	Deleted: cation
~~(	Deleted: of cooling
(	Deleted: southern
(	Deleted: and
(	Deleted: southwestern southern South Atlantic
~~(	Deleted: in
(	Deleted: larger and, more permanent ice sheets on the Antarctic continent
•••••(	Deleted: We, therefore, infer that in the Middle Miocene, the

large(???) degree of cooling in the southern South Atlantic as recordeds at Sites 696 and U1536 indicates that SSTs were sensitive to the decrease in respond sensitively to the pCO2 in the Middle Miocene-induced cooling.

that regional cooling was not amplified through strengthening of the ACC and destruction of the sub polar gyre, which occurred 925 later in the Miocene (Evangelinos et al., 2021).

By the latest Miocene (~6.4-5.2 Ma), temperatures at Site U1536 cooled to a average SST of 3.4°C (n=8) which match those at SHALLDRILL II for, the earliest Pliocene (5.1-4.3 Ma, average\_2.8°C, n=3). The 3-5°C SST decrease at Site U1536 between the MMCT (~14 Ma) and the latest Miocene (~6.4 5.2 Ma) is perhaps surprisingly small compared to the ~10°C

- 930 cooling at the subtropical front Sites 1168 and 1088, at the same time. We surmise that high-latitude cooling was subdued during this time interval because southwestern South Atlantic surface waters (Site U1536) were already cold during the MMCT (Fig. 7A). Instead, the remaining part of the Southern Ocean experienced pronounced cooling during this time interval. Based on the Southern Ocean SST records (Fig. 7A), the southwestern South Atlantic was possibly the coldest region of the Southern Ocean already since at least the Middle Miocene. However, there is a strong lack of Antarctic proximal records (e.g., no records
- from the Ross Sea; Levy et al., 2019) that cover the MMCT to Late Miocene, partly due to glacial advances, to draw a full 935 picture of circum-Antarctic cooling since the MMCT.

Site J1536 SSTs and Site 747 BWTs were low during the MMCT (Leutert et al., 2021), with minor cooling thereafter. Leutert et al. (2021) attributed the subdued post MMCT cooling to the growing Antarctic ice sheet, which could have led to increased 940 stratification and shielding of deeper waters in the Southern Ocean. We here conclude that the southwestern South Atlantic regions already reached cold conditions during the MMCT, because of the proximity to ice sheets, and, as a result, could not cool much more given the global cold climate of the Late Miocene. The cooling of the subtropics (Sites 1088 and 1168) is much more pronounced than the southwestern. South Atlantic, because of the gradual northward expansion of the westerly winds, ACC, and cold subantarctic waters (Leutert et al., 2020) in response to the expansion of the Antarctic ice sheet in the

945 Late Miocene. This process likely also further promoted cooling in other sectors of the Antarctic-proximal Southern Ocean.

#### 5.3 South Atlantic SST gradient evolution

Compiling all available SST records for the two time slices discussed above (Sect. 5.2 and 5.3) yields a unique insight into the long-term temperature trends in the South Atlantic Ocean (Fig. 8). The SST records from the South Atlantic region show unidirectional temperature drops across the EOT, with a small degree of polar amplification where Antarctic-proximal (Site 950 696 and 511) cooled by ~8°C and subtropical records (Site 1090) cooled by ~5°C. The strong, large and persistent South Atlantic subpolar gyre kept the latitudinal SST gradient low in the southernmost part of the South Atlantic across the EOT (Huber et al., 2004; Houben et al., 2019). The latitudinal temperature gradient in the South Atlantic increased during the MMCT (Fig. 7, 8), due to the largest cooling at high latitudes, almost reaching modern temperatures, followed by a subdued cooling during the Late Miocene. Meanwhile the SSTs at the subtropical front (Sites 1090 and 1088) remained relatively stable

from the earliest Oligocene (Site 1090, 33 Ma) until the Late Miocene (Site 1088) (Fig. 8), with minimal cooling until the 955 latest Miocene.

#### Formatted: Font colour: Black

Deleted: The (late) middle Miocene (12.8-11.7 Ma) TEX<sub>86</sub> data from SHALDRILL II Hole 5D (Tibbett et al., 2022) indicate that relatively warm high latitude SSTs (mean: 10.2°± 3.4°C) persisted towards the Late Miocene

$\sum$	Deleted:
$\langle \rangle \rangle$	Deleted: mean
$\langle \rangle \rangle$	Deleted: in
$\langle \rangle \rangle$	Deleted: mean
$\langle \rangle \rangle$	Deleted:
	Deleted: sites
()	Deleted: thisigh-latitude cooling was subdued in ( [24]
1	Deleted: southern
	<b>Deleted:</b> It ishe restemaining part of the Southern Ocean that experienced pronounced cooling inuring this time interval. Based on the Southern Ocean SST records (Fig. 7A), the southwestern South Atlantic record the coldest SSTs andas therefore ([25])
Ì	Deleted: albeit
	Deleted: Both
(/)	Deleted: s
$\langle \rangle \rangle$	Deleted: show a lack of
///	Deleted: reached cold ow temperatures by the uring t
	Deleted: – Late Miocene cooling in SST and BWT, respectively (Leutert et al., 2021), with minor cooling thereafter. Leutert et al. (2021) attributed the lack ofsubtudeost MMCT cooling to the growing Antarctic ice sheet, which could have led to increased stratification and shielding of deeper waters in the Southern Ocean. We here conclude that the southwesternsouthern
	Deleted: further
	Deleted: [old??]
	Deleted: inf the subtropics (Sites 1088 and 1168) is in the Late Miocene
	Deleted: southern
Y	Deleted: biniling all available the ( [30]
>	<b>Deleted:</b> (~8°C) cross the EOT, with a small degree of polar

amplification where between ...ntarctic-proximal (Site 696 and 511) cooled by ~8°C and and ...ubtropical recordss...(Site 1090) cooled by ~5°C. The strong, large and persistent South Atlantic sS...bpolar gG... ... [31]) Deleted: across the EOT

#### Deleted: ??

Deleted: across ...n the South Atlantic increaseds...during the MMCT (Fig. 7, 8), due to the largest cooling at high latitudes, almost reaching high latitude cooling towards ... odern values ... emperatures, followed by a subdued cooling during the Late Miocene. Meanwhile the SSTs at the subtropical front SST records (... [32])



 data from Site 1090 (dark pink; Liu et al., 2009). U<sup>k</sup>'37- and TEX<sub>86</sub>-SST from Site 511 (green; Houben et al., 2019). U<sup>k</sup>'37-SST from
 1060 1088 (pink; Herbert et al., 2016). TEX<sub>86</sub>-SST from Site U1536 (red; this study). Clumped isotope (Δα7) SST estimates from La Meseta Fm., Seymour Island (purple; Douglas et al., 2014). C. Benthic foraminferal δ<sup>18</sup>O compilation, with a locally weighted smooth over 20 kyr (black curve) (CENOGRID; Westerhold et al., 2020), and a LOESS smooth (blue; span = 0.2). D. Published paleo-CO2 data from selected proxies named in legend (https://www.paleo-co2.org), with LOESS smooth (red; span = 0.3). LMC=Late Miocene



14

ммст мсо

12 14

16 18

Cooling, MCO=Miocene climatic optimum, EOIS=Early Oligocene oxygen isotope step, EOT=Eocene-Oligocene transition. The gray shaded areas indicate the time intervals discussed in sections 5.1 and 5.2 as indicated above the figure.

The cooling phases in the South Atlantic across the EOT and from the MCO to the MMCT represent two climatic transitional
 phases, both characterized by declining atmospheric *p*CO<sub>2</sub> concentrations (Pearson et al., 2009; Foster et al., 2012; Greenop et al., 2014; Steinthorsdottir et al., 2016) and increasing benthic foraminiferal δ<sup>18</sup>O values (Westerhold et al., 2020), indicating deep-sea cooling and/or ice sheet expansion (Flower and Kennett, 1993; Zachos et al., 1996) (Fig. 8). The EOT marks the first installation of a continent-wide Antarctic ice sheet (Deconto and Pollard, 2003; Coxall et al., 2005), with a volume between 60 and 130% of that of present<sub>3</sub>day ice sheet (Bohaty et al., 2012). The MCO is considered as a global warm phase, with warm-

1075 temperate ice proximal conditions (Sangiorgi et al., 2018) and a profoundly reduced Antarctic ice volume (Shevenell et al., 2008; Foster et al., 2012), and the MMCT is a strong and stepwise transition towards a larger Antarctic ice sheet (Rohling et al., 2022). Surprisingly, although outhwestern South Atlantic records (Sites 696 and U1536) are of low resolution with notable age uncertainties, they do suggest similar Antarctic-proximal SSTs (~12–14°C) for the early Oligocene, when a large, predominately terrestrial ice sheet with marine terminating glaciers was installed, as for the MCO, when ice sheets were profoundly reduced. Keeping in mind the higher-than-modern Antarctic paleotopography in the Oligocene (Wilson et al., 2009; Duncan et al., 2022), with a gradual subsidence during the Miocene (Paxman et al., 2019), this still puts both climate phases into perspective: apparently the Oligocene Antarctic ice sheet could be strongly reduced despite a relatively cold ice-proximal South Atlantic Ocean.

#### 1085 6 Conclusions

090

1095

Our lipid biomarker records from IODP Site U1536 and ODP Site 696 have generated new insights for the understanding of the SST evolution of the South Atlantic Ocean, viz.:

- The EOT in the South Atlantic is characterized by a relatively small latitudinal SST gradient of ~5 degrees between the subtropical front and the western Weddell Sea and a regional decrease in SST (4–6°C) as global pCO<sub>2</sub> declined.
- The South Atlantic SST gradient remains constant, across the EOT, which we ascribe to a gyral circulation that connects all South Atlantic sites, and can persist in the absence of a strong throughflow through Drake Passage.
  - Southwestern, South Atlantic SSTs at the earliest Oligocene glaciation were similar to those of the warm MCO, implying that Antarctic proximal SSTs are not the only determining factor for the extent of the Antarctic ice sheet.
  - The southwestern, South Atlantic experienced cold polar climate conditions (SSTs of ~5°C) already during the MMCT. This made it the coldest oceanic region around Antarctica and the likely region of deep-water formation.
  - Due to the already relatively cold conditions in the <u>southwestern</u> South Atlantic in the Middle Miocene, it experienced little further cooling during the Late Miocene. This is in contrast to subtropical sites and other sectors of the Southern

Deleted: we	
Deleted: S	
Deleted: .	
Deleted: . These Sections are	
Deleted: on top of	

Deleted: the	
Deleted:	
Deleted:	
Deleted: the MMCT a strong, stepwise transition	
Deleted: southern	

Deleted:

Deleted: decrease	
Deleted: s show no prominent increase in the SST gradient	
Deleted: the persisten	
Deleted: t	
Deleted: ,	
Deleted: hern	
Deleted: ern	

Deleted: southern

Ocean, which experienced profound cooling due to northward expansion of the Southern Ocean frontal systems as the Antarctic ice sheet expanded in the Late Miocene.

## 120 Supplement

The supplement related to this article including the full TEX<sub>26</sub>-SST datasets from Site 696 (Supplementary Table 3) and Site 4, U1536 (Supplementary Table 4) is available online at: XYZ

### Author contributions

FSH, PKB and FS designed the research. FSH, PKB, CE, ALQ and JE collected the samples. ALQ ansd CE provided depositional information, core description, facies analyses and age constraint for ODP Site 696. SvdL advised on the Drake Passage tectonic evolution and provided (paleo) geographic maps for figures 1, 6 and 7. JE, MAS and FSH processed samples for organic geochemistry. FSH, PKB, FP and FS interpreted the data. FSH wrote the paper with input from all authors.

#### **Competing interest**

The authors declare that they have no conflict of interest.

1130

#### Acknowledgements

This work used International Ocean Discovery Program (IODP) archived samples and data. We thank the the great scientist and crew on Expedition 382, also helping with data interpritation and discussions of results from Site U1536. We thank Mariska Hoorweg for technical support at the Utrecht University GeoLab. FSH and PKB acknowledges funding from the NWO polar

programme (grant number ALW.2016.001) and ERC starting grant 802835 "OceanNice". CE and ALQ acknowledges funding provided by the Spanish Ministry of Science and Innovation (grants CTM2014-60451-C2-1/2-P and CTM2017-89711-C2-1/2-P, co-funded by the European Union through FEDER funds) and FJC2021-047046-I (MCIN/AEI/10.13039/501100011033 and NextGenerationEU/PRTR). SvdL acknowledges funding by NWO Vici (grant nr. 865.17.001) awarded to Douwe van Hinsbergen.

#### 1140 References

Barker, P. and Thomas, E.: Origin, signature and palaeoclimatic influence of the Antarctic Circumpolar Current, Earth-Science Reviews, 66, 143-162, 2004.

Barker, P. F., Filippelli, G. M., Florindo, F., Martin, E. E., and Scher, H. D.: Onset and role of the Antarctic Circumpolar Current, Deep Sea Research Part II: Topical Studies in Oceanography, 54, 2388-2398, 2007.

1145 Barker, P. F., Kennett, J. P., O'Connell, S., Berkowitz, S., Bryant, W. R., Burckle, L. H., Egeberg, P. K., Futterer, D. K., Gersonde, R. E., and Golovchenko, X.: Proceedings of the Ocean Drilling Program, Initial Reports, Vol. 113. Weddell Sea,

24

## Formatted: Font: Font colour: Auto, Check spelling and grammar

- Formatted: Subscript
- Formatted: Normal, Line spacing: single

**Commented [HF(1]:** Link to supplement provided by Copernicus

Formatted: Font: Font colour: Auto, Check spelling and grammar

Formatted: Font: Font colour: Auto, Check spelling and grammar

Formatted: Font: Font colour: Auto, Check spelling and grammar

Deleted:

Antarctica. Covering Leg 113 of the cruises of the drilling vessel JOIDES Resolution, Valparaiso, Chile, to East Cove, Falkland Islands, Sites 689-697, 25 December 1986-11 March 1987, <u>1988.</u>

1150 Bijl, P. K., Bendle, J. A., Bohaty, S. M., Pross, J., Schouten, S., Tauxe, L., Stickley, C. E., McKay, R. M., Röhl, U., and Olney, M.: Eocene cooling linked to early flow across the Tasmanian Gateway, Proceedings of the National Academy of Sciences, 110, 9645-9650, 2013.

 Bijl, P. K., Houben, A. J., Hartman, J. D., Pross, J., Salabarnada, A., Escutia, C., and Sangiorgi, F.: Paleoceanography and ice sheet variability offshore Wilkes Land, Antarctica-Part 2: Insights from Oligocene-Miocene dinoflagellate cyst
 assemblages, Climate of the Past, 14, 1015-1033, 2018.

Blaga, C. I., Reichart, G.-J., Heiri, O., and Sinninghe Damsté, J. S.: Tetraether membrane lipid distributions in water-column particulate matter and sediments: a study of 47 European lakes along a north–south transect, Journal of Paleolimnology, 41, 523-540, 2009.

 Bohaty, S. M., Kulhanek, D. K., Wise Jr, S. W., Jemison, K., Warny, S., and Sjunneskog, C.: Age assessment of Eocene–
 Pliocene drill cores recovered during the SHALDRIL II expedition, Antarctic Peninsula, Tectonic, Climatic, and Cryospheric Evolution of the Antarctic Peninsula, 63-113, 2011.

Bohaty, S. M., Zachos, J. C., and Delaney, M. L.: Foraminiferal Mg/Ca evidence for southern ocean cooling across the eocene–oligocene transition, Earth and Planetary Science Letters, 317, 251-261, 2012.

Carter, A., Riley, T. R., Hillenbrand, C.-D., and Rittner, M.: Widespread Antarctic glaciation during the late Eocene, Earth and Planetary Science Letters, 458, 49-57, 2017.

Coxall, H. K., Wilson, P. A., Pälike, H., Lear, C. H., and Backman, J.: Rapid stepwise onset of Antarctic glaciation and deeper calcite compensation in the Pacific Ocean, Nature, 433, 53-57, 2005.

Cramwinckel, M. J., Huber, M., Kocken, I. J., Agnini, C., Bijl, P. K., Bohaty, S. M., Frieling, J., Goldner, A., Hilgen, F. J., and Kip, E. L.: Synchronous tropical and polar temperature evolution in the Eocene, Nature, 559, 382, 2018.

1170 DeConto, R. M. and Pollard, D.: Rapid Cenozoic glaciation of Antarctica induced by declining atmospheric CO2, Nature, 421, 245-249, 2003.

Douglas, P. M., Affek, H. P., Ivany, L. C., Houben, A. J., Sijp, W. P., Sluijs, A., Schouten, S., and Pagani, M.: Pronounced zonal heterogeneity in Eocene southern high-latitude sea surface temperatures, Proceedings of the National Academy of Sciences, 111, 6582-6587, 2014.

175 Duncan, B., McKay, R., Levy, R., Naish, T., Prebble, J., Sangiorgi, F., Krishnan, S., Hoem, F., Clowes, C., and Dunkley Jones, T.: Climatic and tectonic drivers of late Oligocene Antarctic ice volume, Nature Geoscience, 15, 819-825, 2022.

Dunkley Jones, T., Eley, Y. L., Thomson, W., Greene, S. E., Mandel, I., Edgar, K., and Bendle, J. A.: OPTiMAL: A new machine learning approach for GDGT-based palaeothermometry, Climate of the Past, 16, 2599-2617, 2020.

Eagles, G. and Jokat, W.: Tectonic reconstructions for paleobathymetry in Drake Passage, Tectonophysics, 611, 28-50, 2014.

1180 England, M. H., Hutchinson, D. K., Santoso, A., and Sijp, W. P.: Ice-atmosphere feedbacks dominate the response of the climate system to Drake Passage closure, Journal of Climate, 30, 5775-5790, https://doi.org/10.1175/JCLI-D-15-0554.1, 2017.

Flower, B. and Kennett, J.: Middle Miocene ocean-climate transition: High-resolution oxygen and carbon isotopic records from Deep Sea Drilling Project Site 588A, southwest Pacific, Paleoceanography, 8, 811-843, 1993.

1185 Foster, G. L., Lear, C. H., and Rae, J. W.: The evolution of pCO2, ice volume and climate during the middle Miocene, Earth and Planetary Science Letters, 341, 243-254, 2012.

Gersonde, R. and Burckle, L. H.: 43. NEOGENE DIATOM BIOSTRATIGRAPHY OF ODP LEG 113, WEDDELL SEA (ANTARCTIC OCEAN), 1990.

Goldner, A., Herold, N., and Huber, M.: Antarctic glaciation caused ocean circulation changes at the Eocene–Oligocene transition, Nature, 511, 574-577, 2014.

Greenop, R., Foster, G. L., Wilson, P. A., and Lear, C. H.: Middle Miocene climate instability associated with highamplitude CO2 variability, Paleoceanography, 29, 845-853, 2014.

Hartman, J. D., Sangiorgi, F., Salabarnada, A., Peterse, F., Houben, A. J., Schouten, S., Brinkhuis, H., Escutia, C., and Bijl, P. K.: Paleoceanography and ice sheet variability offshore Wilkes Land, Antarctica-Part 3: Insights from Oligocene-Miocene TEX86-based sea surface temperature reconstructions, Climate of the Past, 14, 1275-1297, https://doi.org/10.5194/cp-14-

1275-2018, 2018.

Herbert, T., Schuffert, J., Thomas, D., Lange, C., Weinheimer, A., Peleo-Alampay, A., and Herguera, J. C.: Depth and seasonality of alkenone production along the California margin inferred from a core top transect, Paleoceanography, 13, 263-271, 1998.

1200 Herold, N., Huber, M., Müller, R., and Seton, M.: Modeling the Miocene climatic optimum: Ocean circulation, Paleoceanography, 27, 2012.

1195

Hill, D. J., Haywood, A. M., Valdes, P. J., Francis, J. E., Lunt, D. J., Wade, B. S., and Bowman, V. C.: Paleogeographic controls on the onset of the Antarctic circumpolar current, Geophysical Research Letters, 40, 5199-5204, 2013.

Ho, S. L. and Laepple, T.: Flat meridional temperature gradient in the early Eocene in the subsurface rather than surface l205 ocean, Nature Geoscience, 9, 606-610, 2016.

Hoem, F. S., Valero, L., Evangelinos, D., Escutia, C., Duncan, B., McKay, R. M., Brinkhuis, H., Sangiorgi, F., and Bijl, P.
 K.: Temperate Oligocene surface ocean conditions offshore of Cape Adare, Ross Sea, Antarctica, Climate of the Past, 17, 1423-1442, https://doi.org/10.5194/cp-17-1423-2021, 2021a.

Hoem, F. S., Sauermilch, I., Hou, S., Brinkhuis, H., Sangiorgi, F., and Bijl, P. K.: Late Eocene–early Miocene evolution of the southern Australian subtropical front: a marine palynological approach, Journal of Micropalaeontology, 40, 175-193, https://doi.org/10.5194/jm-40-175-2021, 2021b.

Hoem, F. S., Sauermilch, I., Aleksinski, A. K., Huber, M., Peterse, F., Sangiorgi, F., and Bijl, P. K.: Strength and variability of the Oligocene Southern Ocean surface temperature gradient, Communications Earth & Environment, 3, 322, 2022.

215	Hoem, F. S.: A Song of Ice and a Warm Southern Ocean: The paleoceanographic evolution of the Oligocene–Miocene Southern Ocean, Doctoral dissertation, Utrecht University, 2022	Deleted:
	Holbourn, A. E., Kuhnt, W., Clemens, S. C., Kochhann, K. G., Jöhnck, J., Lübbers, J., and Andersen, N.: Late Miocene climate cooling and intensification of southeast Asian winter monsoon, Nature communications, 9, 1-13, 2018.	Deleted: Paleo-CO2 data archive (Version 1) [Data set]. Zenodo., last access: 02.01.2023.¶
	Holland, M. M. and Bitz, C. M.: Polar amplification of climate change in coupled models, Climate dynamics, 21, 221-232, 2003.	
1220	Hopmans, E. C., Weijers, J. W., Schefuß, E., Herfort, L., Damsté, J. S. S., and Schouten, S.: A novel proxy for terrestrial organic matter in sediments based on branched and isoprenoid tetraether lipids, Earth and Planetary Science Letters, 224, 107-116, 2004.	
	Hopmans, E. C., Schouten, S., and Damsté, J. S. S.: The effect of improved chromatography on GDGT-based palaeoproxies, Organic Geochemistry, 93, 1-6, 2016.	
1225	Hou, S., Lamprou, F., Hoem, F. S., Hadju, M. R. N., Sangiorgi, F., Peterse, F., and Bijl, P. K.: Lipid biomarker-based sea (sub) surface temperature record offshore Tasmania over the last 23 million years, Climate of the Past Discussions, 1-33, 2022.	
1230	Houben, A. J., Bijl, P. K., Pross, J., Bohaty, S. M., Passchier, S., Stickley, C. E., Röhl, U., Sugisaki, S., Tauxe, L., and van de Flierdt, T.: Reorganization of Southern Ocean plankton ecosystem at the onset of Antarctic glaciation, Science, 340, 341-344, 2013.	
	Houben, A. J., Bijl, P. K., Sluijs, A., Schouten, S., and Brinkhuis, H.: Late Eocene Southern Ocean cooling and invigoration of circulation preconditioned Antarctica for full-scale glaciation, Geochemistry, Geophysics, Geosystems, 2019.	
	Huber, M., Brinkhuis, H., Stickley, C. E., Döös, K., Sluijs, A., Warnaar, J., Schellenberg, S. A., and Williams, G. L.: Eocene circulation of the Southern Ocean: Was Antarctica kept warm by subtropical waters?, Paleoceanography, 19, 2004.	
1235	Hutchinson, D. K., Coxall, H. K., Lunt, D. J., Steinthorsdottir, M., De Boer, A. M., Baatsen, M., von der Heydt, A., Huber, M., Kennedy-Asser, A. T., and Kunzmann, L.: The Eocene–Oligocene transition: a review of marine and terrestrial proxy data, models and model–data comparisons, Climate of the Past, 17, 269-315, 2021.	
240	Inglis, G. N., Toney, J. L., Zhu, J., Poulsen, C. J., Röhl, U., Jamieson, S. S., Pross, J., Cramwinckel, M. J., Krishnan, S., and Pagani, M.: Enhanced terrestrial carbon export from East Antarctica during the early Eocene, Paleoceanography and Paleoclimatology, 37, e2021PA004348, 2022.	Formatted: English (UK)
Ι	Kennedy, A. T., Farnsworth, A., Lunt, D., Lear, C. H., and Markwick, P.: Atmospheric and oceanic impacts of Antarctic glaciation across the Eocene–Oligocene transition, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences, 373, 20140419, 2015.	
1245	Kennedy-Asser, A. T., Lunt, D. J., Valdes, P. J., Ladant, JB., Frieling, J., and Lauretano, V.: Changes in the high-latitude Southern Hemisphere through the Eocene–Oligocene transition: a model–data comparison, Climate of the Past, 16, 555-573, 2020.	
	Kennett, J. P.: Cenozoic evolution of Antarctic glaciation, the circum-Antarctic Ocean, and their impact on global paleoceanography, Journal of geophysical research, 82, 3843-3860, 1977.	
	27	

Kim, J.-H., Van der Meer, J., Schouten, S., Helmke, P., Willmott, V., Sangiorgi, F., Koç, N., Hopmans, E. C., and Damsté, J. S. S.: New indices and calibrations derived from the distribution of crenarchaeal isoprenoid tetraether lipids: Implications for past sea surface temperature reconstructions, Geochimica et Cosmochimica Acta, 74, 4639-4654, 10.1016/j.gca.2010.05.027, 2010.

Lagabrielle, Y., Goddéris, Y., Donnadieu, Y., Malavieille, J., and Suarez, M.: The tectonic history of Drake Passage and its possible impacts on global climate, Earth and Planetary Science Letters, 279, 197-211, 2009.

Leutert, T. J., Auderset, A., Martínez-García, A., Modestou, S., and Meckler, A. N.: Coupled Southern Ocean cooling and Antarctic ice sheet expansion during the middle Miocene, Nature Geoscience, 13, 634-639, 2020.

1260 Leutert, T. J., Modestou, S., Bernasconi, S. M., and Meckler, A. N.: Southern Ocean bottom-water cooling and ice sheet expansion during the middle Miocene climate transition, Climate of the Past, 17, 2255-2271, 2021.

 Levy, R. H., Meyers, S. R., Naish, T. R., Golledge, N. R., McKay, R. M., Crampton, J. S., DeConto, R. M., De Santis, L., Florindo, F., Gasson, E. G. W., Harwood, D. M., Luyendyk, B. P., Powell, R. D., Clowes, C., and Kulhanek, D. K.: Antarctic ice-sheet sensitivity to obliquity forcing enhanced through ocean connections, Nature Geoscience, 12, 132-137, 10.1038/s41561-018-0284-4, 2019.

Lewis, A. R., Marchant, D. R., Ashworth, A. C., Hedenäs, L., Hemming, S. R., Johnson, J. V., Leng, M. J., Machlus, M. L., Newton, A. E., and Raine, J. I.: Mid-Miocene cooling and the extinction of tundra in continental Antarctica, Proceedings of the National Academy of Sciences, 105, 10676-10680, 2008.

Livermore, R., Hillenbrand, C. D., Meredith, M., and Eagles, G.: Drake Passage and Cenozoic climate: an open and shut l270 case?, Geochemistry, Geophysics, Geosystems, 8, 2007.

Locarnini, M., Mishonov, A., Baranova, O., Boyer, T., Zweng, M., Garcia, H., Seidov, D., Weathers, K., Paver, C., and Smolyar, I.: World ocean atlas 2018, volume 1: Temperature, 2018.

 <u>López-Quirós, A., Escutia, C., Etourneau, J., Rodríguez-Tovar, F.J., Bijl, P.K., Lobo, F.J., Bohoyo, F., Evangelinos, D.,</u> <u>Salabarnada, A. 2018. Eocene-Miocene paleoceanographic changes in Drake Passage (Antarctica). POLAR 2018: Where the</u> <u>Poles come together. Abstract Proceedings, 487. Publisher: WSL Institute for Snow and Avalanche Research SLF. ISBN</u> <u>978-0- 948277-54-2.</u>

López-Quirós, A., Escutia, C., Sánchez-Navas, A., Nieto, F., Garcia-Casco, A., Martín-Algarra, A., Evangelinos, D., and Salabarnada, A.: Glaucony authigenesis, maturity and alteration in the Weddell Sea: An indicator of paleoenvironmental conditions before the onset of Antarctic glaciation, Scientific reports, 9, 1-12, 2019.

1280 López-Quirós, A., Sánchez-Navas, A., Nieto, F., and Escutia, C.: New insights into the nature of glauconite, American Mineralogist: Journal of Earth and Planetary Materials, 105, 674-686, 2020.

López-Quirós, A., Escutia, C., Etourneau, J., Rodríguez-Tovar, F. J., Roignant, S., Lobo, F. J., Thompson, N., Bijl, P. K., Bohoyo, F., and Salzmann, U.: Eocene-Oligocene paleoenvironmental changes in the South Orkney Microcontinent (Antarctica) linked to the opening of Powell Basin, Global and Planetary Change, 204, 103581, 2021.

[285 Lyle, M., Gibbs, S., Moore, T. C., and Rea, D. K.: Late Oligocene initiation of the Antarctic Circumpolar Current: Evidence from the South Pacific, Geology, 35, 691-694, 10.1130/g23806a.1, 2007. Deleted: López-Quirós, A., Nieto, F., Garcia-Casco, A., Hoem, F.S., Harwood, D., Morales-Ocaña, C., Bohoyo, F., Escutia, C. Bijl, P.K. Insights into the development of Jane Basin, northern Weddell Sea (Antarctica): Diagenetic constraints on marine sediments from ODP Site 696. In prep.<sup>6</sup>

Maldonado, A., Bohoyo, F., Galindo-Zaldívar, J., Hernández-Molina, J., Jabaloy, A., Lobo, F., Rodríguez-Fernández, J., Suriñach, E., and Vázquez, J.: Ocean basins near the Scotia–Antarctic plate boundary: influence of tectonics and paleoceanography on the Cenozoic deposits, Marine Geophysical Researches, 27, 83-107, 2006.

1295 Maldonado, A., Bohoyo, F., Galindo-Zaldívar, J., Hernández-Molina, F. J., Lobo, F. J., Lodolo, E., Martos, Y. M., Pérez, L. F., Schreider, A. A., and Somoza, L.: A model of oceanic development by ridge jumping: opening of the Scotia Sea, Global and Planetary Change, 123, 152-173, 2014.

Martínez-Botí, M. A., Foster, G. L., Chalk, T. B., Rohling, E. J., Sexton, P. F., Lunt, D. J., Pancost, R. D., Badger, M. P., and Schmidt, D. N.: Plio-Pleistocene climate sensitivity evaluated using high-resolution CO2 records, Nature, 518, 49-54, 2015.

1300 Müller, P. J., Kirst, G., Ruhland, G., Von Storch, I., and Rosell-Melé, A.: Calibration of the alkenone paleotemperature index U37K' based on core-tops from the eastern South Atlantic and the global ocean (60 N-60 S), Geochimica et Cosmochimica Acta, 62, 1757-1772, 1998.

Nooteboom, P. D., Baatsen, M., Bijl, P. K., Kliphuis, M. A., van Sebille, E., Sluijs, A., Dijkstra, H. A., and von der Heydt,
 A. S.: Improved Model-Data Agreement With Strongly Eddying Ocean Simulations in the Middle-Late Eocene,
 Paleoceanography and paleoclimatology, 37, e2021PA004405, 2022.

O'Brien, C. L., Robinson, S. A., Pancost, R. D., Damsté, J. S. S., Schouten, S., Lunt, D. J., Alsenz, H., Bornemann, A., Bottini, C., and Brassell, S. C.: Cretaceous sea-surface temperature evolution: Constraints from TEX86 and planktonic foraminiferal oxygen isotopes, Earth-Science Reviews, 172, 224-247, 2017.

O'Brien, C. L., Huber, M., Thomas, E., Pagani, M., Super, J. R., Elder, L. E., and Hull, P. M.: The enigma of Oligocene l310 climate and global surface temperature evolution, Proceedings of the National Academy of Sciences, 117, 25302-25309, 2020

Orsi, A. H., Whitworth III, T., and Nowlin Jr, W. D.: On the meridional extent and fronts of the Antarctic Circumpolar Current, Deep Sea Research Part I: Oceanographic Research Papers, 42, 641-673, 1995.

Orsi, A. H., Johnson, G. C., and Bullister, J. L.: Circulation, mixing, and production of Antarctic Bottom Water, Progress in 1315 Oceanography, 43, 55-109, 1999.

Paxman, G. J., Jamieson, S. S., Hochmuth, K., Gohl, K., Bentley, M. J., Leitchenkov, G., and Ferraccioli, F.: Reconstructions of Antarctic topography since the Eocene–Oligocene boundary, Palaeogeography, palaeoclimatology, palaeoecology, 535, 109346, 2019.

Pearson, P. N., Foster, G. L., and Wade, B. S.: Atmospheric carbon dioxide through the Eocene–Oligocene climate transition, Nature, 461, 1110-1113, 2009.

Pérez, L. F., Hernández-Molina, F. J., Lodolo, E., Bohoyo, F., Galindo-Zaldívar, J., and Maldonado, A.: Oceanographic and climatic consequences of the tectonic evolution of the southern scotia sea basins, Antarctica, Earth-Science Reviews, 198, 102922, 2019.

Pérez, L. F., Martos, Y. M., García, M., Weber, M. E., Raymo, M. E., Williams, T., Bohoyo, F., Armbrecht, L., Bailey, I.,
 and Brachfeld, S.: Miocene to present oceanographic variability in the Scotia Sea and Antarctic ice sheets dynamics: Insight from revised seismic-stratigraphy following IODP Expedition 382, Earth and Planetary Science Letters, 553, 116657, 2021.

Rohling, E. J., Foster, G. L., Gernon, T. M., Grant, K. M., Heslop, D., Hibbert, F. D., Roberts, A. P., and Yu, J.: Comparison and synthesis of sea-level and deep-sea temperature variations over the past 40 million years, Reviews of Geophysics, e2022RG000775, 2022.

1330 Salabarnada, A., Escutia, C., Röhl, U., Nelson, C. H., McKay, R., Jiménez-Espejo, F., Bijl, P., Hartman, J., Strother, S., and Salzmann, U.: Paleoceanography and ice sheet variability offshore Wilkes Land, Antarctica–Part 1: Insights from late Oligocene astronomically paced contourite sedimentation, Climate of the Past, 14, 991-1014, 10.5194/cp-14-991-2018, 2018.

Sangiorgi, F., Bijl, P. K., Passchier, S., Salzmann, U., Schouten, S., McKay, R., Cody, R. D., Pross, J., van de Flierdt, T.,
 Bohaty, S. M., Levy, R., Williams, T., Escutia, C., and Brinkhuis, H.: Southern Ocean warming and Wilkes Land ice sheet retreat during the mid-Miocene, Nat Commun, 9, 317, 10.1038/s41467-017-02609-7, 2018.

Sauermilch, I., Whittaker, J. M., Klocker, A., Munday, D. R., Hochmuth, K., Bijl, P. K., and LaCasce, J. H.: Gateway-driven weakening of ocean gyres leads to Southern Ocean cooling, Nature communications, 12, 1-8, 2021.

Scher, H. D. and Martin, E. E.: Timing and climatic consequences of the opening of Drake Passage, science, 312, 428-430, 1340 2006.

- Schouten, S., Hopmans, E. C., Schefuß, E., and Damste, J. S. S.: Distributional variations in marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient sea water temperatures?, Earth and Planetary Science Letters, 204, 265-274, 2002.
- Schouten, S., Hopmans, E. C., and Damsté, J. S. S.: The organic geochemistry of glycerol dialkyl glycerol tetraether lipids: 1345 A review, Organic geochemistry, 54, 19-61, 2013.

Seton, M., Müller, R. D., Zahirovic, S., Gaina, C., Torsvik, T., Shephard, G., Talsma, A., Gurnis, M., Turner, M., and Maus, S.: Global continental and ocean basin reconstructions since 200 Ma, Earth-Science Reviews, 113, 212-270, 2012.

Shevenell, A. E., Kennett, J. P., and Lea, D. W.: Middle Miocene ice sheet dynamics, deep-sea temperatures, and carbon cycling: A Southern Ocean perspective, Geochemistry, Geophysics, Geosystems, 9, 2008.

1350 Sijp, W. P., Anna, S., Dijkstra, H. A., Flögel, S., Douglas, P. M., and Bijl, P. K.: The role of ocean gateways on cooling climate on long time scales, Global and Planetary Change, 119, 1-22, 2014.

Sijp, W. P., von der Heydt, A. S., and Bijl, P. K.: Model simulations of early westward flow across the Tasman Gateway during the early Eocene, Climate of the Past, 12, 807-817, 2016.

Sinninghe Damsté, J. S., Ossebaar, J., Abbas, B., Schouten, S., and Verschuren, D.: Fluxes and distribution of tetraether
 lipids in an equatorial African lake: constraints on the application of the TEX86 palaeothermometer and BIT index in
 lacustrine settings, Geochimica et Cosmochimica Acta, 73, 4232-4249, 2009.

Sinninghe Damsté, J. S.: Spatial heterogeneity of sources of branched tetraethers in shelf systems: The geochemistry of tetraethers in the Berau River delta (Kalimantan, Indonesia), Geochimica et Cosmochimica Acta, 186, 13-31, 2016.

Sosdian, S. M., Greenop, R., Hain, M., Foster, G. L., Pearson, P. N., and Lear, C. H.: Constraining the evolution of Neogene
 ocean carbonate chemistry using the boron isotope pH proxy, Earth and Planetary Science Letters, 498, 362-376, 2018.

Deleted: Scher, H. D., Bohaty, S. M., Smith, B. W., and Munn, G. H.: Isotopic interrogation of a suspected late Eocene glaciation, Paleoceanography, 29, 628-644, 2014.¶ Scher, H. D., Whittaker, J. M., Williams, S. E., Latimer, J. C., Kordesch, W. E. C., and Delaney, M. L.: Onset of Antarctic Circumpolar Current 30 million years ago as Tasmanian Gateway aligned with westerlies, Nature, 523, 580, 10.1038/nature14598, 2015.¶

Steinthorsdottir, M., Porter, A. S., Holohan, A., Kunzmann, L., Collinson, M., and McElwain, J. C.: Fossil plant stomata indicate decreasing atmospheric CO 2 prior to the Eocene–Oligocene boundary, Climate of the Past, 12, 439-454, 2016.

Stickley, C. E., Brinkhuis, H., Schellenberg, S. A., Sluijs, A., Röhl, U., Fuller, M., Grauert, M., Huber, M., Warnaar, J., and Williams, G. L.: Timing and nature of the deepening of the Tasmanian Gateway, Paleoceanography, 19, 2004.

Super, J. R., Thomas, E., Pagani, M., Huber, M., O'Brien, C., and Hull, P. M.: North Atlantic temperature and pCO2 coupling in the early-middle Miocene, Geology, 46, 519-522, 2018.

1375 Taylor, K. W., Huber, M., Hollis, C. J., Hernandez-Sanchez, M. T., and Pancost, R. D.: Re-evaluating modern and Palaeogene GDGT distributions: Implications for SST reconstructions, Global and Planetary Change, 108, 158-174, 2013.

Thompson, N., Salzmann, U., López-Quirós, A., Bijl, P. K., Hoem, F. S., Etourneau, J., Sicre, M.-A., Roignant, S., Hocking, E., and Amoo, M.: Vegetation change across the Drake Passage region linked to late Eocene cooling and glacial disturbance after the Eocene–Oligocene Transition, Climate of the Past, 18, 209-232, 2022.

380 Tibbett, E. J., Scher, H. D., Warny, S., Tierney, J. E., Passchier, S., and Feakins, S. J.: Late Eocene record of hydrology and temperature from Prydz Bay, East Antarctica, Paleoceanography and Paleoclimatology, 36, e2020PA004204, 2021.

Tibbett, E. J., Warny, S., Tierney, J. E., Wellner, J. S., and Feakins, S. J.: Cenozoic Antarctic Peninsula Temperatures and Glacial Erosion Signals From a Multi-Proxy Biomarker Study, Paleoceanography and Paleoclimatology, 37, e2022PA004430, 2022.

1385 Tierney, J. E. and Tingley, M. P.: A TEX 86 surface sediment database and extended Bayesian calibration, Scientific data, 2, 1-10, 2015.

Toggweiler, J. R., Russell, J. L., and Carson, S. R.: Midlatitude westerlies, atmospheric CO2, and climate change during the ice ages, Paleoceanography, 21, 2006.

Torsvik, T. H., Van der Voo, R., Preeden, U., Mac Niocaill, C., Steinberger, B., Doubrovine, P. V., Van Hinsbergen, D. J.,
 Domeier, M., Gaina, C., and Tohver, E.: Phanerozoic polar wander, palaeogeography and dynamics, Earth-Science Reviews, 114, 325-368, 2012.

van de Lagemaat, S. H., Swart, M. L., Vaes, B., Kosters, M. E., Boschman, L. M., Burton-Johnson, A., Bijl, P. K., Spakman, W., and van Hinsbergen, D. J.: Subduction initiation in the Scotia Sea region and opening of the Drake Passage: When and why?, Earth-Science Reviews, 103551, 2021.

1395 van der Weijst, C. M., van der Laan, K. J., Peterse, F., Reichart, G.-J., Sangiorgi, F., Schouten, S., Veenstra, T. J., and Sluijs, A.: A 15-million-year surface-and subsurface-integrated TEX 86 temperature record from the eastern equatorial Atlantic, Climate of the Past, 18, 1947-1962, 2022.

Villa, G., Fioroni, C., Pea, L., Bohaty, S., and Persico, D.: Middle Eocene–late Oligocene climate variability: calcareous nannofossil response at Kerguelen Plateau, Site 748, Marine Micropaleontology, 69, 173-192, 2008.

1400 Weber, M., Raymo, M., Peck, V., Williams, T., Armbrecht, L., Bailey, I., Brachfeld, S., Cardillo, F., Du, Z., and Fauth, G.: Expedition 382 summary, in, IODP Publications, 2021a.

	Weber, M., Raymo, M., Peck, V., Williams, T., Armbrecht, L., Bailey, I., Brachfeld, S., Cardillo, F., Du, Z., and Fauth, G.: Site U1536, Weber, ME, Raymo, ME, Peck, VL, Williams, T., and the Expedition, 382, 2021b.	
405	Wei, W. and Wise Jr, S. W.: Biogeographic gradients of middle Eocene-Oligocene calcareous nannoplankton in the South Atlantic Ocean, Palaeogeography, Palaeoclimatology, Palaeoecology, 79, 29-61, 1990.	
	Wilson, D. S. and Luyendyk, B. P.: West Antarctic paleotopography estimated at the Eocene-Oligocene climate transition, Geophysical Research Letters, 36, 2009	Formatted: English (UK)
410	Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnet, J. S. K., Bohaty, S. M., De Vleeschouwer, D., Florindo, F., Frederichs, T., Hodell, D. A., Holbourn, A. E., Kroon, D., Lauretano, V., Littler, K., Lourens, L. J., Lyle, M., Palike, H., Rohl, U., Tian, J., Wilkens, R. H., Wilson, P. A., and Zachos, J. C.: An astronomically dated record of Earth's climate and its predictability over the last 66 million years, Science, 369, 1383-1387, 10.1126/science.aba6853, 2020.	
	Whittaker, J., Muller, R., Leitchenkov, G., Stagg, H., Sdrolias, M., Gaina, C., and Goncharov, A.: Major Australian- Antarctic plate reorganization at Hawaiian-Emperor bend time, Science, 318, 83-86, 2007.	
415	Zachos, J. C., Quinn, T. M., and Salamy, K. A.: High-resolution (104 years) deep-sea foraminiferal stable isotope records of the Eocene-Oligocene climate transition, Paleoceanography, 11, 251-266, 1996.	<b>Deleted:</b> Xing, Q., Munday, D., Klocker, A., Sauermilch, I., and Whittaker, J.: The sensitivity of the Eocene–Oligocene Southern Ocean to the strength and position of wind stress, Climate of the Past, 18, 2669-2693-2022.

32

Zhang, Y. G., Pagani, M., and Wang, Z.: Ring Index: A new strategy to evaluate the integrity of TEX86 paleothermometry, Paleoceanography, 31, 220-232, 2016.

Page 14: [1] Deleted	Hoem, F.S. (Frida)	29/04/2023 18:39:00	
v	· · ·		
<b>A</b>			
Page 14: [1] Deleted	Hoem, F.S. (Frida)	29/04/2023 18:39:00	
Υ			
<b>A</b>			
Page 14: [2] Deleted	Peterse, F. (Francien)	22/06/2023 14:54:00	
Υ			
<b>A</b>			
Page 14: [2] Deleted	Peterse, F. (Francien)	22/06/2023 14:54:00	
Υ			
A Dego 14: [2] Deleted	Dotoreo E (Eroneion)	22/06/2022 14:54:00	
Page 14: [2] Deleted	Peterse, F. (Francien)	22/06/2023 14:54:00	
<b>V</b>			
Page 14: [3] Deleted	Peterse, F. (Francien)	22/06/2023 14:55:00	
-			
V			
Page 14: [3] Deleted	Peterse, F. (Francien)	22/06/2023 14:55:00	
<u>ــــــــــــــــــــــــــــــــــــ</u>			
Page 14: [3] Deleted	Peterse, F. (Francien)	22/06/2023 14:55:00	
<b>V</b>			
<u> </u>			
Page 14: [4] Deleted	Hoem, F.S. (Frida)	11/04/2023 09:14:00	
۲			
<b>A</b>			
Page 14: [4] Deleted	Hoem, F.S. (Frida)	11/04/2023 09:14:00	
۲			
Page 14: [4] Deleted	Hoom E.S. (Erida)	11/04/2022 00:14:00	
Page 14: [4] Deleted	noem, r.S. (rnud)	11/04/2023 09:14:00	
<b>V</b>			
Page 14: [4] Deleted	Hoem, F.S. (Frida)	11/04/2023 09:14:00	
×			
Page 14: [5] Deleted	Sangiorgi, F. (Francesca)	29/06/2023 16:20:00	
T			
<b>A</b>			
Page 14: [5] Deleted	Sangiorgi, F. (Francesca)	29/06/2023 16:20:00	
▼			
<b>A</b>			
Page 14: [5] Deleted	Sangiorgi, F. (Francesca)	29/06/2023 16:20:00	
▼			
Page 14: [6] Deleted	Bijl, P.K. (Peter) 23/04/	2023 12:50:00	

٧.

Page 14: [8] Deleted	Peterse, F. (Francien)	22/06/2023 15:07:00	
۷			
Page 14: [8] Deleted	Peterse F (Francien)	22/06/2023 15:07:00	
	receise, it (italicien)	22/00/2023 13:07:00	
×			
Page 14: [8] Deleted	Peterse, F. (Francien)	22/06/2023 15:07:00	
Υ			
A Dago 14 [9] Delotod	Dotorso E (Eronsion)	22/06/2022 15:07:00	
Page 14: [8] Deleted	Peterse, r. (Francien)	22/06/2023 15:07:00	
▼			
Page 14: [8] Deleted	Peterse, F. (Francien)	22/06/2023 15:07:00	
▼			
	<b></b>		
Page 14: [8] Deleted	Peterse, F. (Francien)	22/06/2023 15:07:00	
×			
Page 14: [8] Deleted	Peterse, F. (Francien)	22/06/2023 15:07:00	
▼			
A			
Page 14: [9] Deleted	Peterse, F. (Francien)	22/06/2023 15:09:00	
¥			
Page 14: [9] Deleted	Peterse, F. (Francien)	22/06/2023 15:09:00	
v			
<b>A</b>			
Page 14: [9] Deleted	Peterse, F. (Francien)	22/06/2023 15:09:00	
<b>V</b>			
Page 17: [10] Deleted	Hoem, F.S. (Frida)	09/04/2023 10:06:00	
Page 17: [10] Deleted	Hoom ES (Erida)	00/04/2023 10:06:00	
		09/04/2023 10:06:00	
Page 17: [10] Deleted	Hoem, F.S. (Frida)	09/04/2023 10:06:00	
Page 17: [10] Deleted	Hoem, F.S. (Frida)	09/04/2023 10:06:00	
Page 17: [11] Deleted	Hoem, F.S. (Frida)	08/04/2023 15:22:00	
v			
<u> </u>			
Page 17: [11] Deleted	Hoem, F.S. (Frida)	08/04/2023 15:22:00	

Bijl, P.K. (Peter) 28/06/2023 09:24:00

 Page 17: [11] Deleted
 Hoem, F.S. (Frida)
 08/04/2023 15:22:00

**v**....

Page 14: [7] Deleted

Page 17: [12] Deleted	Hoem, F.S. (Frida)	09/04/2023 09:50:00	
<b>v</b>			 
Page 17: [12] Deleted	Hoem, F.S. (Frida)	09/04/2023 09:50:00	
▼			_
<b>_</b>			 
Page 17: [12] Deleted	Hoem, F.S. (Frida)	09/04/2023 09:50:00	
T			_
<b>A</b>			 
Page 17: [12] Deleted	Hoem, F.S. (Frida)	09/04/2023 09:50:00	
•			-
<b>A</b>			
Page 17: [12] Deleted	Hoem, F.S. (Frida)	09/04/2023 09:50:00	
•			-
▲			
Page 17: [13] Deleted	Peterse, F. (Francien)	22/06/2023 15:29:00	1
			-
¥			
Page 17: [13] Deleted	Peterse, F. (Francien)	22/06/2023 15:29:00	1
-		· ·	-
V			 
Page 17: [13] Deleted	Peterse, F. (Francien)	22/06/2023 15:29:00	1
-			
V			
Page 17: [14] Deleted	Biil, P.K. (Peter) 28/06/	2023 09:43:00	1
-			
A			 
Page 17: [14] Deleted	Bijl, P.K. (Peter) 28/06/	2023 09:43:00	1
			-
A			
Page 17: [14] Deleted	Biil, P.K. (Peter) 28/06/	2023 09:43:00	1
-			-
A			
Page 17: [15] Formatted	Hoem, F.S. (Frida)	04/05/2023 16:00:00	
Font colour: Text 1			
Page 17: [15] Formatted	Hoem, F.S. (Frida)	04/05/2023 16:00:00	]
Font colour: Text 1			
Page 17: [16] Deleted	Biil, P.K. (Peter) 28/06/	2023 09:25:00	1
-	,,		4
X			 
Page 17: [16] Deleted	Biil, P.K. (Peter) 28/06/	2023 09:25:00	 1
. age 17. [10] Deleten			3
X			 
Page 17: [16] Delated	Rijl DK (Deter) 28/06/	2023 00.25.00	1
rage 17. [10] Deleted	Diji, F.K. (Felei) 20/00/	2023 09.23.00	

**x**.....

Page 17: [18] Formatted	Hoem, F.S. (Frida)	04/05/2023 16:00:00	
Font colour: Text 1			
Page 17: [18] Formatted	Hoem, F.S. (Frida)	04/05/2023 16:00:00	
Font colour: Text 1			
Page 17: [19] Deleted	Peterse, F. (Francien)	22/06/2023 15:36:00	
X			
Page 17: [19] Deleted	Peterse, F. (Francien)	22/06/2023 15:36:00	
<b>X</b>			
Page 17: [20] Deleted	Hoem, F.S. (Frida)	30/06/2023 09:55:00	
٧			
<b>A</b>			
Page 17: [21] Deleted	Hoem, F.S. (Frida)	30/06/2023 10:28:00	
V			
Page 17: [21] Deleted	Hoem, F.S. (Frida)	30/06/2023 10:28:00	
▼			
Page 17: [21] Deleted	Hoem, F.S. (Frida)	30/06/2023 10:28:00	
<b>V</b>			
Page 17: [21] Deleted	Hoem, F.S. (Frida)	30/06/2023 10:28:00	
<b>v</b>			
Page 17: [21] Deleted	Hoem, F.S. (Frida)	30/06/2023 10:28:00	
▼			
Page 17: [21] Deleted	Hoem, F.S. (Frida)	30/06/2023 10:28:00	
<b>v</b>			
Page 17: [21] Deleted	Hoom E.S. (Erida)	20/06/2022 10:29:00	
Page 17: [21] Deleteu	noem, r.s. (riua)	50/06/2025 10:28:00	
▼			
Page 17: [22] Deleted	Peterse, F. (Francien)	22/06/2023 15:57:00	
<b>v</b>			
A Dage 17: [22] Deleted	Potorso E (Francian)	22/06/2023 15:57:00	
▼			
Page 17: [23] Deleted	Peterse, F. (Francien)	22/06/2023 15:58:00	
<b>v</b>			
Page 17: [23] Delated	Deterse E (Francian)	22/06/2023 15:58:00	
raye 17: [23] Deleted	receise, r. (rrancien)	22/00/2023 13:30:00	

Page 21: [24] Deleted	Peterse, F. (Francien)	22/06/2023 16:32:00	
<b>v</b>			
Page 21: [25] Deleted	Deterse E (Francien)	22/06/2023 16:32:00	
Page 21. [25] Deleteu	Peterse, F. (Francien)	22/00/2023 10.52.00	
×			
Page 21: [25] Deleted	Peterse, F. (Francien)	22/06/2023 16:32:00	
▼			
Page 21: [25] Deleted	Peterse, F. (Francien)	22/06/2023 16:32:00	
V	, , , ,		
<b>A</b>			
Page 21: [25] Deleted	Peterse, F. (Francien)	22/06/2023 16:32:00	
▼			
Page 21: [25] Deleted	Peterse, F. (Francien)	22/06/2023 16:32:00	
▼			
		22/06/2022 46.22.00	
Page 21: [25] Deleted	Peterse, F. (Francien)	22/06/2023 16:32:00	
×			
Page 21: [26] Deleted	Peterse, F. (Francien)	22/06/2023 16:44:00	
Υ			
Page 21: [26] Deleted	Peterse, F. (Francien)	22/06/2023 16:44:00	
· · · · · · · · · · · · · · · · · · ·			
<u> </u>			
Page 21: [26] Deleted	Peterse, F. (Francien)	22/06/2023 16:44:00	
▼			
Page 21: [27] Deleted	Hoem, F.S. (Frida)	09/04/2023 12:06:00	
٧			
Page 21: [27] Deleted	Hoem, F.S. (Frida)	09/04/2023 12:06:00	
▼			
Page 21: [27] Deleted	Hoem, F.S. (Frida)	09/04/2023 12:06:00	
Ψ			
Page 21: [27] Deleted	Hoem F.S. (Frida)	09/04/2023 12:06:00	
• • • • • • • • • • • • • • • • • • •			
<u> </u>			
Page 21: [27] Deleted	Hoem, F.S. (Frida)	09/04/2023 12:06:00	
<b>v</b>			
Page 21: [28] Deleted	Hoem, F.S. (Frida)	26/06/2023 17:08:00	
	, , , , , , , , , , , , , , , , , , , ,		

Υ.,

Page 21: [29] Deleted	Peterse, F. (Francien)	22/06/2023 16:49:00	
<b>v</b>			
<b>A</b>			
Page 21: [30] Deleted	Peterse, F. (Francien)	22/06/2023 16:53:00	
▼			
Page 21: [30] Deleted	Peterse, F. (Francien)	22/06/2023 16:53:00	
-			
▲			
Page 21: [31] Deleted	Hoem, F.S. (Frida)	04/05/2023 16:14:00	
▼			
<b>A</b>		1	
Page 21: [31] Deleted	Hoem, F.S. (Frida)	04/05/2023 16:14:00	
▼			
Page 21: [31] Deleted	Hoem ES (Erida)	04/05/2023 16:14:00	
rage 21. [31] Deleted		07/03/2023 10:14:00	
<b>V</b>			
Page 21: [31] Deleted	Hoem, F.S. (Frida)	04/05/2023 16:14:00	
V			
<u> </u>			
Page 21: [31] Deleted	Hoem, F.S. (Frida)	04/05/2023 16:14:00	
▼			
A Dago 21, [21] Deloted	Hoom E.S. (Erida)	04/05/2022 16:14:00	
Page 21: [51] Deleteu	noem, r.s. (rnua)	04/05/2025 10:14:00	
▼			
Page 21: [32] Deleted	Peterse, F. (Francien)	22/06/2023 16:55:00	
V			
<b>A</b>			
Page 21: [32] Deleted	Peterse, F. (Francien)	22/06/2023 16:55:00	
▼			
A Desco 21, [22] Delet		22/06/2022 16/FF-00	
Page 21: [32] Deleted	Peterse, F. (Francien)	22/00/2023 10:55:00	
▼			
Page 21: [32] Deleted	Peterse, F. (Francien)	22/06/2023 16:55:00	
×			
Page 21: [32] Deleted	Peterse, F. (Francien)	22/06/2023 16:55:00	
▼			
<b>A</b>			