



Storm characteristics influence nitrogen removal in an urban estuarine environment

Anne Margaret H. Smiley¹, Suzanne P. Thompson², Nathan Hall², Michael F. Piehler^{1,2,3}

¹Department of Earth, Marine, and Environmental Sciences, University of North Carolina, Chapel Hill, 27514, USA

5 ²UNC Institute of Marine Sciences, Morehead City, 28557, USA

³UNC Institute for the Environment, Chapel Hill, 27517, USA

Correspondence to: Anne Margaret H. Smiley (ahsmiley@live.unc.edu)

Abstract. Sustaining water quality is an important component of coastal resilience. Floodwaters deliver reactive nitrogen (NO_x) to sensitive aquatic systems and can diminish water quality. Coastal habitats in flooded areas can be effective at removing reactive nitrogen through denitrification (DNF). However, less is known about this biogeochemical process in urbanized environments. This study assessed the nitrogen removal capabilities of flooded habitats along an urban estuarine coastline in the upper Neuse River Estuary (NRE), NC, USA under two nitrate concentrations (16.8 μM and 52.3 μM NO_x, respectively). We also determined how storm characteristics (e.g., precipitation and wind) affect water column NO_x concentrations and consequently DNF by flooded habitats. Continuous flow-through sediment core incubation experiments quantified gas and nutrient fluxes across the sediment-water interface in marsh, swamp forest, undeveloped open space, stormwater pond, and shallow subtidal sediments. All habitats exhibited net DNF. Additionally, all habitats increased DNF rates under elevated nitrate conditions compared to low nitrate. Structured habitats with high sediment organic matter had higher nitrogen removal capacity than unstructured, low sediment organic matter habitats. High precipitation-high wind storm events produced concentrations significantly lower than other types of storms (e.g., low precipitation-high wind, high wind-low precipitation, low wind-low precipitation), which likely results in relatively low DNF rates by flooded habitats and low removal percentages of total dissolved nitrogen loads. These results demonstrate the importance of natural systems to water quality in urbanized coastal areas subject to flooding.

1 Introduction

Tropical cyclones often cause extensive flooding that can harm ecosystems, damage infrastructure, and disrupt the lives of coastal residents. There is evidence to suggest that anthropogenic climate change has produced conditions (e.g., warmer sea surface temperatures, increased atmospheric moisture) that make these high magnitude events more likely (Knutson et al., 2013; Min et al., 2011). Since the mid-1990's there has been an observed shift in storm activity in the United States where tropical cyclones have become slower and rainier, and result in catastrophic flooding at higher frequencies than historical averages (Easterling et al., 2017; Kossin, 2018; Kunkel et al., 2010). As climate change continues, some models predict an increase in the most intense storms and up to a 20 % increase in precipitation rates by 2100 (Knutson et al., 2010).



Floodwaters introduce allochthonous materials, including nutrients, to downstream receiving waters. Storm-related upstream discharge typically contains high concentrations of inorganic nitrogen and organic carbon, that constitute up to 80 % of annual loads into receiving waterbodies (Paerl et al., 2020). Estuaries are often nitrogen limited and sensitive to sudden influxes of reactive nitrogen (Howarth & Marino, 2006), therefore floodwaters can trigger water quality degradation by fueling algal blooms (Nixon, 1995) that can disrupt aquatic ecosystems by outcompeting other vegetated habitats for sunlight and nutrients (Wasson et al., 2017). One of the largest cyanobacterial blooms in the Okeechobee region has been attributed to Hurricane Irma in 2017 (Hampel et al., 2019) and models showed a strong biological response in Apalachicola Bay following Hurricane Michael in 2018 (D'Sa et al., 2019). Remineralization of algal biomass and terrigenous organic matter by heterotrophic bacteria depletes oxygen in the water column, which can affect health of aquatic organisms and the ecosystem overall (Diaz & Roseberg, 1995).

Watershed urbanization has been shown to exacerbate water quality degradation in tropical, subtropical, and temperate coastal regions by interfering with hydrologic, geomorphic, and biogeochemical processes (Bowen & Valiela, 2001; Gold et al., 2019, 2021; Lee et al., 2006; Ortiz-Zayas et al., 2006). Population growth has led to increased point source nutrient loading via wastewater effluent into receiving waterways (Carey & Migliaccio, 2009; Naden et al., 2016). Furthermore, impervious surfaces and stormwater pipes can streamline flow paths and enhance the export of non-point source anthropogenic nitrogen (Bernhardt et al., 2008). Agricultural landscapes can also deliver nutrients to receiving waterways. In some regions, high densities of agricultural operations substantially increase nutrient concentrations from nitrogen-based fertilizer and animal waste (Duda, 1982; Dupas et al., 2015).

Some natural habitats have been shown to be effective at removing terrigenous and anthropogenic nitrogen through a series of biogeochemical reactions (Groffman & Crawford, 2003; Pérez-Villalona et al., 2015; Piehler & Smyth, 2011; Reisinger et al., 2016; Rosenzweig et al., 2018). Denitrification (DNF) is a process by which sediment microbes convert bioavailable forms of nitrogen (nitrate and nitrite) to dinitrogen gas (N_2) under anaerobic conditions using carbon as an energy source. DNF is an important process by which reactive nitrogen is naturally and permanently removed from a system. It can be an effective strategy for maintaining water quality during flood conditions that favor DNF, namely, elevated dissolved nitrate and carbon, and anoxia (Adame et al., 2019; Velinsky et al., 2017). Much work has been done to understand DNF by natural habitats, such as emergent wetlands and oyster reefs (Ensign et al., 2008; Grabowski et al., 2012; Onorevole et al., 2018; Piehler & Smyth, 2011; Velinsky et al., 2017), but much less is known about nitrogen processing by urban landscapes, such as stormwater ponds and lawns/undeveloped open space (UOS), despite their being prolific in developed settings. The primary objective of this study is to quantify nitrogen removal by denitrification in flood-prone habitats, both natural and human influenced, including marsh, forested wetland, stormwater pond, undeveloped open space, and shallow subtidal sediments under varied nutrient conditions in Neuse River Estuary (NRE), North Carolina (NC).



65


Storms exhibit unique characteristics which can affect water chemistry differently (Davis et al., 2004; Mallin et al., 2002; Wetz & Paerl, 2008). Some storms produce elevated nutrient concentrations. Studies have shown that sustained winds at high speeds can increase nutrient concentrations by mixing stratified waters and resuspending sediments (Goodrich et al., 1987; Miller et al., 2006; Wengrove et al., 2015). Storms characterized by high precipitation can dilute the nutrients in the water column (Minaudo et al., 2019). Paerl et al. (2020) described the Neuse River Estuary (NRE), in eastern North Carolina, as either a “processor” under relatively lower discharge periods where nutrients are able to be partially processed, or a “pipeline” during high discharge periods where nutrients are delivered to the Albemarle-Pamlico Sound with little processing in the NRE. Therefore, the nitrogen removal capacity of flooded landscapes via DNF is likely influenced by water quality produced during varied storm conditions as well as contact time of floodwaters prior to export from the system. A secondary objective of this work is to compare nitrogen loads during multiple storm types to projected nitrogen removal rates by predominant landscape habitats including inundated marshes, forested wetland, UOS, and subtidal sediments.

As urban landscapes expand, resulting in losses of natural habitats and wetlands (Aguilera et al., 2020) concomitant with increased anthropogenic nutrient loads, it is essential that we understand the role that both natural and human influenced landscapes play in removing reactive nitrogen. Additionally, assessing how these habitats perform under a range of nutrient conditions will enable us to estimate landscape scale nitrogen removal capacities during different types of storms. These data will improve our understanding of estuarine nutrient budgets along urbanized coastlines in a new regime of tropical cyclone activity and inform strategic coastal development that conserves ecosystem services to maximize benefits for coastal residents.

2 Methods

2.1 Approach

This study combines laboratory, computational modeling, and geographic information systems (GIS) methods to understand landscape-scale DNF capacity during different types of storms. Storm types were defined based on precipitation and wind characteristics. NO_x concentrations, NO_x loads, and TDN loads during those storm events were modeled using weighted regressions. Habitat-specific DNF rates under ambient and elevated nitrate conditions were determined through laboratory experiments. These nitrate treatments represented low and high water column NO_x concentrations that are likely associated with different types of storms. Nitrogen removal during these storms was estimated based on experimental DNF values and inundated area of each habitat treatment at maximum inundation. These results were used to draw conclusions about the influence of storm characteristics on water column NO_x concentrations, and consequently, biogeochemical processes in flooded landscapes.



95

Figure 1. Site map of New Bern, NC in the upper Neuse River Estuary and sampling locations for sediment core and water collection.

2.2 Site description

The upper reaches of the Neuse River Estuary (NRE), North Carolina, USA are influenced by both riverine and coastal hydrologic processes, making them prone to multiple forms of flooding (e.g., fluvial, pluvial, and coastal storm surge). This region also includes both highly urbanized areas and natural ecosystems, affording the opportunity to assess anthropogenic impacts on ecosystem functioning in the context of a built environment. NRE is nutrient sensitive (Boyer et al., 1994; Pinckney et al.; Rudek et al., 1991); primary production is primarily nitrogen limited and episodic loading events can result in water quality degradation. With headwaters at the urban center of Raleigh and several smaller cities distributed along the river and throughout the watershed, the NRE receives inputs from a 16,000 km² drainage basin (Christian et al., 1991). Extensive agricultural use paired with rapid urbanization within the watershed makes NRE and similar locations susceptible to water quality degradation during major flood events.



2.3 Storm classification and water quality characteristics

2.3.1 Storm types based on wind and precipitation

Paerl et al. (2018) categorized tropical cyclones that made landfall in North Carolina between 1996 and 2016, based on river discharge at Fort Barnwell and wind speeds at Cape Lookout, NC (NOAA National Data Buoy Center Station CLKN7). The same criteria were used to categorize storms between 2017 and 2019. Storms that resulted in a 7-day mean Neuse River discharge above the 90th percentile of weekly averages ($191 \text{ m}^3 \text{ s}^{-1}$) were designated “high precipitation” (HP) events. Those that exhibited a maximum hourly average wind speed above the 90th percentile (14.1 m s^{-1}) between the 12 hours prior to and 24 hours after landfall, were considered “high wind” (HW) events. Storms that produced riverine discharges or wind speeds below these thresholds were considered “low precipitation” (LP) and “low wind” (LW) events, respectively. Storm types were assigned based on both precipitation and wind classifications (Table 1). For example, Hurricane Florence produced both high precipitation and high wind conditions and is therefore labelled as a HP-HW storm. Furthermore, storms that were considered both LP-LW were thought of as “baseline storm” conditions.

120 **Table 1. Summary matrix of named storms that made landfall on North Carolina’s coast between 1996 and 2019 categorized by storm type derived from Neuse River discharge and average wind speeds. Red text indicates storms with available floodplain footprints that were assessed for nitrogen removal.**

	High precipitation (HP)	Low precipitation (LP)
High wind (HW)	Fran (1996) Josephine (1996) Dennis (1999) Floyd (1999) Irene (1999) Gordon (2000)	Ernesto (2006) Irene (2011) Joaquin (2015) Matthew (2016) Florence (2018)
Low wind (LW)	Charley (2004) Nicole (2010) Ana (2015) Dorian (2019)	Arthur (1996) Bertha (1996) Bonnie (1998) Earl (1998) Helene (2000) Gustav (2002) Isabel (2003)



125 2.3.2 Water quality characteristics during multiple storms

This study compared water quality characteristics during multiple tropical cyclones that affected the North Carolina coast. Average NO_x concentrations, NO_x loads, and TDN loads on the day of maximum riverine discharge were estimated using Weighted Regressions on Time Discharge and Season (WRTDS) (Hirsch et al., 2010) as described in Paerl et al. (2018) with the following exception. The half window width for the flow term in the model was reduced from the default of 2 $\ln(\text{flow})$ increments down to 1 $\ln(\text{flow})$ increment to provide greater resolution of flow impacts on concentration and fluxes. This was necessary to capture the observed strong dilution effect of nitrate during extreme, storm-induced flood events. Average concentrations and loads were compared across storm types.

2.4 Nitrogen flux experiment

2.4.1 Sample collection

135 Sampling for nitrogen flux experiments occurred in October of 2020, timed to capture typical environmental conditions during hurricane season. Sediment cores (6.4 cm diameter with a height of approximately 17 cm) were collected in triplicate from habitats subject to storm flooding including subtidal sediments, stormwater pond, UOS, swamp forest and two marshes—one upstream and one downstream of the outfall from the City of New Bern’s wastewater treatment facility (Figure 1). The two marshes did not exhibit significant differences between mean flux rates and were, therefore, treated as a single habitat treatment.

2.4.2 Dissolved gas and nutrient fluxes across the sediment-water interface

The sediment cores and water were taken to UNC-CH’s Institute of Marine Sciences in Morehead City, NC to conduct dissolved gas and soluble nitrogen flux experiments using methods described by Piehler and Smyth (2011). Sediments and site water were incubated in a temperature-controlled chamber (Bally Inc.) set to *in situ* water temperature (19 °C) in a continuous flow-through system of water collected from the Neuse River between the two marshes (feedwater). The ambient NO_x concentration of the feedwater was 16.8 μM . A peristaltic pump was used to pull feedwater through the sediment cores at a rate of 0.6 L h^{-1} , equating to a turnover time of 5-6 hours for water over the sediment in each core. After an overnight equilibration period, water samples were collected from each sediment core outflow for three timepoints, each 5 hours apart. Water pumped directly from feedwater bins was collected at each timepoint to assess inflow concentrations of dissolved gases and nutrients. After collecting the third timepoint, the feedwater was enriched with sodium nitrate to a concentration of 52.3 μM to simulate a relatively high nitrogen scenario. Following a second overnight equilibration period under nitrate enriched conditions, three more timepoints were collected 5 hours apart. These two nitrogen treatments are referred to as “low nitrate” (16.8 μM ambient NO_x concentration) and “high nitrate” (52.3 μM enriched NO_x concentration).



155 Directly following each water collection (6 total), a membrane inlet mass spectrometer (MIMS; Bay Instruments, Easton, MD)
was used to analyze ratios of concentrations of dissolved gases, including $N_2:Ar$ and $O_2:Ar$, within water samples pulled from
cores as well as those collected directly from the feedwater. These measurements were used to calculate net DNF rates and
sediment oxygen demand (SOD). At timepoints 2 and 5, additional 50 mL water samples were collected to measure nutrient
fluxes based on core inflow and outflow concentrations. Samples were filtered through 0.7 μm Whatman GF/F filters and
160 stored at $-18^\circ C$ prior to analysis with a Lachat Quick-chem 8000 (Lachat Instruments, Milwaukee, WI, USA). Nutrient analytes
included dissolved inorganic nitrogen species (DIN): nitrate + nitrite (NO_x) and ammonium (NH_4^+) as well as total dissolved
nitrogen (TDN) allowing calculation of dissolved organic nitrogen (DON) by difference. At the end of the experiment, water
was drained from the cores and sediment samples were collected from the top 2 cm to determine percent sediment organic
material (SOM) based on loss on ignition (Byers et al., 1978; Smyth et al., 2015).

165 **2.5 Spatial data acquisition**

2.5.1 Habitat treatments

Distributions and total surface area of sampled habitats were determined using a variety of spatial datasets. Marshes, swamp
forests, and UOS were delineated using the National Land Cover Dataset (NLCD), a 30m raster dataset obtained from remotely
sensed Landsat imagery. Land cover classified as emergent and forested wetlands were considered marsh and swamp forest,
170 respectively. Area of UOS was calculated by combining the herbaceous classification and weighted estimates from the various
development categories (e.g., open space, low-high intensity). Pixels considered “Developed, open space” in the NLCD are
defined as those comprised of less than 20 percent constructed surfaces. The remaining 80 % of the pixel area was considered
UOS, colloquially referred to as lawns and grasses. Low, medium, and high intensity developed pixels were considered 51 %,
21 %, and 0 % UOS, respectively. NLCD datasets have been updated roughly in 2 to 3-year intervals. This work references
175 datasets from multiple years, including 2004, 2016, and 2019.

Shallow subtidal sediments were identified using NOAA’s Continuously Updated Digital Elevation Model (CUDEM) and
were defined as those within 1 meter of the surface. Stormwater infrastructure data were obtained from the City of New Bern
and include managed stormwater ponds. ArcGIS Pro 2.8.7 was used to extract NLCD and CUDEM data from 2 HUC12
180 watersheds in the upper NRE.

2.5.2 Inundation extents for multiple storms

Flood footprints that delineated inundated landscapes for seven selected storms were generated from the Advanced Circulation
(ADCIRC) model and acquired from the Coastal Emergency Risks Assessment website (<https://cera.coastalrisk.live/>). This
analysis includes Hurricanes Charley (8/23/2004), Arthur (7/4/2014), Joaquin (10/12/2015), Hermine (9/2/2016), Matthew



185 (10/8/2016), Florence (9/26/2018), and Dorian (9/6/2019). Flood footprints for LP-LW storms were not available, so these types of baseline storm events were not considered.

2.6 Calculations and statistical analysis

2.6.1 Dissolved gas and nutrient fluxes

Fluxes of nutrient and dissolved gases across the sediment water interface were calculated by multiplying the difference
190 between inflow and outflow concentrations by the peristaltic pump/flow rate and dividing by the surface area of the sediment core as in Eq. (1).

$$Flux = \frac{(C_{outflow} - C_{inflow}) \times F}{A} \quad (1)$$

Denitrification efficiency was calculated by dividing N_2 -N fluxes by the total inorganic nitrogen flux out of the sediments,
195 then multiplying 100, following Eq. (2).

$$Denitrification\ efficiency = \frac{Flux_{(N_2-N)}}{Flux_{(N_2-N)} + Flux_{(NO_x-N)} + Flux_{(NH_4-N)}} \times 100 \quad (2)$$

Mean flux rates for dissolved gases and nutrients were compared using Kruskal Wallis and post hoc Dunn tests to identify
200 differences across landscape treatments and between nutrient treatments. Linear regressions were performed to compare variations in N_2 -N fluxes to variations in SOD under ambient and nitrate enriched conditions. Additional linear regressions were used to compare variability in DNF to SOM under both low nitrate and high nitrate conditions. All statistical tests were done using R version 4.1.1 (R Core Team, 2011) and were considered significant when $p < 0.05$.

2.6.2 Nitrogen concentrations and loads during storms

Mean NO_x concentrations, NO_x loads, and TDN loads on the day of maximum riverine discharge were compared across storm
205 types were compared using Kruskal Wallis and post hoc Dunn tests. Differences between mean NO_x concentrations were used to draw comparisons between experimental nitrate treatments to environmental NO_x concentrations during different types of storms. Mean load values were compared to estimate nitrogen removal by flooded landscapes during multiple storms.

2.6.3 Nitrogen removal by the flooded landscape

Nitrogen removal was estimated for seven selected storms with available flood footprints. Tools in ArcGIS Pro were used to
210 extract land cover data from each storm's flood footprint within the two HUC12 watersheds in the upper NRE. The 2004 NLCD dataset was used to estimate inundated area during Hurricane Charley, the 2016 dataset was used to estimate inundated area for Hurricanes Arthur, Joaquin, Hermine, and Matthew, and the 2019 dataset was used for Hurricanes Florence and Dorian. For each habitat type, inundated surface areas and mean DNF rates obtained from the nitrogen flux experiments were




multiplied to estimate habitat-specific N removal rates, as in Eq. (3). Areas of shallow subtidal sediments were assumed to
 215 have remained constant over this time range. Nitrogen removal by stormwater ponds was not considered in this analysis.
 Removal rates under both high and low nitrate conditions were calculated.

$$N \text{ removal rate} = DNF \text{ rate} \times \text{surface area} \quad (3)$$

3 Results

220 3.1 Storm characteristics

HP-HW storms yielded a mean NO_x concentration of $11.7 \pm 2.50 \mu\text{M}$ on the day of maximum riverine discharge, which was
 lower than mean concentrations for the other three storm types (HP-LW: 25.2 ± 6.00 ; LP-HW: 29.3 ± 2.23 ; LP-LW: $24.5 \pm$
 225 2.16 ; Figure 2). The low nitrate experimental treatment ($16.8 \mu\text{M}$) was considered more representative of the HP-HW events,
 while the high nitrate treatment ($52.3 \mu\text{M}$) was considered more analogous to the other three storm types. Mean NO_x loads on
 the day of maximum discharge during low precipitation storms were significantly lower than loads during high precipitation
 storms (Figure 2). TDN loads during HP-HW events were higher than low precipitation storms, and LP-LW events produced
 loads lower than high precipitation events. There were no significant differences in mean TDN load between HP-LW and LP-
 HW storms (Figure 2).



230

Figure 2. Average NO_x concentrations, NO_x loads, and average TDN loads on the day of maximum river discharge for each storm type.

3.2 Nitrogen fluxes across the sediment-water interface

Under the low nitrate conditions, all habitats exhibited net DNF (Figure 3a). $\text{N}_2\text{-N}$ fluxes in marsh sediments were significantly
 235 higher than shallow subtidal, swamp forest, and stormwater pond sediments (Figure 3a). Following nitrate enrichment, all



landscapes experienced a significant increase in DNF rates compared to respective flux rates under low nitrate conditions (Figure 3a). Under high nitrate conditions, marsh and stormwater pond cores produced significantly higher DNF rates than both UOS and subtidal sediments. Swamp forest cores also exhibited higher rates than the subtidal cores (Figure 3a).

240 NO_x flux hovered near $0 \mu\text{mol m}^{-2} \text{h}^{-1}$ for each habitat under low nitrate conditions (Figure 3b), with no significant differences evident between means. Following the nitrate addition, each habitat exhibited a significant decrease in flux rates (Figure 3b). High nitrate NO_x flux rates were negative for all habitat treatments, indicating NO_x moving from the water column into the sediments; thus, each habitat acted as a NO_x sink post-enrichment. NO_x flux rates were not different between habitats. NH_4^+ fluxes were an order of magnitude lower than N_2 and NO_x fluxes (Figure 3c). Some NH_4^+ fluxes were positive while
 245 others were negative, although no significant differences across habitat or nitrate treatments were evident.

Average DNF efficiencies for all habitats and nitrate treatments were above 70 % (Figure 3d). Under low nitrate conditions, UOS was the most efficient habitat, significantly more efficient than marsh, stormwater pond, and shallow subtidal sediments. Following nutrient enrichment, marsh, stormwater pond, and shallow subtidal sediments showed a significant increase, and all
 250 habitat treatments nearly reached 100 % DNF efficiency.




Figure 3. Flux rates ($\mu\text{mol m}^{-2} \text{h}^{-1}$) across the sediment water interface for multiple nitrogen species, including: (A) $\text{N}_2\text{-N}$, (B) NO_x , and (C) NH_4 , as well as (D) DNF efficiencies (%). Positive fluxes indicate movement from the sediments to the overlying water column.



255

A linear regression analysis revealed that under low nitrate conditions, variability in SOD explains approximately 70 % of the variability in DNF (Figure 4). No significant relationship was evident between DNF and SOD under high nitrate conditions. There was a significant relationship between SOM and high nitrate DNF rates, where variability in SOM accounted for roughly 62 % of the variability in N₂-N flux.

260




Figure 4. Scatterplot and linear regression for the relationship between SOD and DNF under low (top left) and high (bottom left) nitrate conditions as well as SOM and DNF under low (top right) and high (bottom right) nitrate conditions.


3.3 Nitrogen removal during storms

265 Nitrogen removal was calculated for seven named storms (of 3 precipitation/wind types, Table 1) with available flood footprints. This was done by multiplying habitat specific DNF rates across inundated surface areas for each habitat treatment using rates produced during both low and high nutrient treatments. Flood footprints were not available for any LP-LW storms, so these baseline storm events were not included in this analysis. Removal rates calculated using low nitrate DNF rates ranged between 15.3 and 65.5 kg N h⁻¹. High nitrate removal rates ranged between 58.4 and 257 kg N h⁻¹. Low nitrate removal rates were considered more representative for HP-HW events and high nitrate removal rates were considered more representative for HP-LW and LP-HW events.

270



275 These removal rates were used to calculate the percent of TDN and NO_x loads that were removed by habitats within the floodplains of the seven selected storms. Under low nitrate conditions, the percentage of TDN load removed ranged from 1.15 to 5.95; under high nitrate conditions, they ranged from 4.81 to 24.6. Regarding NO_x loads, under low nitrate conditions, percent removed ranged from 5.71 to 21.6. Under high nitrate conditions, they ranged from 21.8 to 84.6.



280 **Figure 5. NLCD land cover classifications within HUC12 watershed boundaries and floodplain footprints for multiple storms affecting the Neuse River Estuary. Habitat treatments are derived from these land cover classifications.**

285 Floodplain footprints varied in size, with each storm inundating different proportions of each habitat (Figure 5). In each case, swamp forests were the most abundant inundated habitat in the floodplain, comprising between 44.9 and 66.2 % of the flooded habitat area. Their abundance paired with their relatively high DNF rates is reflected in their high contribution to nitrogen removal overall (Figure 6). Swamp forests were likely responsible for removing between 51.6 and 70.1 % of



nitrogen removed




Figure 6. Projected nitrogen removal rates (top) and percent contribution (bottom) by each habitat under low and high nitrate conditions during multiple storms. Higher color intensity indicates applicable nitrogen level based on storm type.

290

via DNF by all habitats under low nitrate conditions and between 64.0 and 79.4 % under high nitrate conditions. Shallow subtidal sediments also consistently comprised a large proportion of flooded habitat area, between 12.4 and 52.2 %. Although DNF rates in this habitat are relatively low, their abundance led to large contributions to nitrogen removal during storms, between 10.1 and 43.3 % under low nitrate conditions and between 7.27 and 31.6 % under high nitrate conditions. Marshes are relatively sparse in this region of the NRE that reliably contributed a small percentage of nitrogen removal. UOS also consistently made up a small portion of inundated landscapes, though the contribution of UOS seems to have increased in the most recent storms: Matthew, Florence, and Dorian (2.33, 13.0, and 6.57 % of the flooded habitat area, respectively).

295

300

Another consideration is the effect that NO_x concentrations have on the relative contribution to overall nitrogen removal by each habitat. Not all habitats respond the same to elevated nitrate conditions. For example, UOS sediments do not increase DNF rates in response to elevated NO_x concentrations to the same degree as other habitats, like swamp forest, do (Figure 2). Therefore, under high nitrate conditions, UOS contribute a smaller proportion to nitrogen removal than under low nitrate conditions (Figure 6). The same is observed with subtidal sediments. Swamp forests, on the other hand, reliably increase the proportion of their contribution under high nitrate conditions.

305



Table 2. Summary of NO_x concentrations (μM) and average nitrogen loads (NO_x and TDN; kg h⁻¹) on the day of maximum discharge, and percent of load removed by habitats under low and high nitrate conditions. Asterisks indicate the more representative percentage based on water column nitrate concentrations during different storm types.

Storm	Type	[NO _x]	NO _x load	TDN load	% TDN load removed- Low nitrate	% TDN load removed- High nitrate	% NO _x load removed- Low nitrate	% NO _x load removed- High nitrate
Joaquin	HP-HW	21.1	295	898	3.05*	12.8	9.27*	38.8
Matthew	HP-HW	8.50	597	3060	1.15*	4.81	5.90*	24.6
Florence	HP-HW	5.33	303	2730	2.40*	9.40	21.6*	84.6
Charley	HP-LW	17.3	212	659	3.31	13.3*	10.3	41.3*
Dorian	HP-LW	22.7	350	1010	4.01	16.5*	11.6	47.7*
Arthur	LP-HW	36.0	218	431	5.95	24.6*	11.8	48.7*
Hermine	LP-HW	24.8	268	726	2.11	8.05*	5.71	21.8*

310

4 Discussion

The results of this study shed light on nitrogen removal capacities of multiple flood-prone natural and human-influenced habitats. Few studies have investigated nitrogen removal by habitats in the context of a built environment (Denman et al., 2016; Groffman & Crawford, 2003; Reisinger et al., 2016; Rosenzweig et al., 2018), and even fewer studies have quantified nitrogen processing by urban landscapes, such as stormwater ponds and lawns (Gold et al., 2017; Hohman et al., 2021; Raciti et al., 2011). As coastlines continue urbanizing, these features will become increasingly abundant and comprise an important piece of coastal nutrient budgets. Additionally, this study explores how characteristics of storms can influence nitrogen removal capacity by coastal landscapes. This information is important in the context of climate change and the projected increase in the rainiest storms.

4.1 Nitrogen removal by habitat treatments

Positive N₂-N flux rates indicate that all habitats are capable of permanently removing nitrogen under high and low nitrate conditions, although, DNF rates varied across habitats and between nitrate treatments for some habitats. Under low nitrate conditions, marsh sediments produced the highest DNF rates and shallow subtidal sediments produced the lowest. Differences in DNF rates between structured habitats, like marshes, and unstructured habitats, like subtidal sediments have been documented in previous studies (Piehler & Smyth, 2011). These differences have been attributed predominantly to the availability of organic carbon. Carbon availability may explain differences between marshes and other habitats as well. Swamp forests are structured habitats, like marshes; however, the forest sediments produced DNF rates that were lower than the marsh

325



330 sediments under low nitrate conditions. It is possible that the quality of carbon affects DNF (Hill & Cardaci, 2004; Seitzinger, 1994). The organic carbon supplied to the sediments by forests, could be more recalcitrant than that of marshes. In subtropical systems, it has been shown that the molecular structure of sediment organic carbon in marshes are simpler and more readily decomposed than their woody counterparts, mangroves, which were dominated by recalcitrant forms of carbon (Sun et al., 2019).

335 There was not a significant difference between DNF rates in marshes and UOS under low nitrate conditions. The few studies that have examined nitrogen processing in urban UOS, have shown that grasses can exhibit high DNF activity (Groffman et al., 1991), but is spatially and temporally heterogenous (McPhillips et al., 2016; Raciti et al., 2011). Multiple factors can influence nitrogen processing, including fertilizer application, soil saturation, species of grass, and temperature (Mancino et al., 1988; Wang et al., 2013). A 1998 study (Mancino et al.) examined nitrogen dynamics in turfgrasses under multiple temperature and soil saturation conditions, and found that when soils are saturated and water temperatures are high, grasses
340 can remove up to 93 % of applied nitrogen via DNF. Experimental conditions in this NRE study are similar to those in the 1998 study—saturated soils and warm temperatures—and results suggest that OUS makes up an important piece of this system’s nitrogen budget under low nitrate conditions.

345 UOS sediments were unique in that they were the only habitat that exhibited 100 % DNF efficiency under low nitrate conditions; all nitrogen that fluxed out of the sediments was in the form of N_2 -N. This could be explained, in part, by organic carbon availability. Eyre and Ferguson (2009) describe critical carbon loading range to maximize DNF efficiency. It is possible that the high DNF efficiency exhibited by UOS sediments under low nitrate conditions is due to the presence of labile organic carbon in the soils that falls within a critical range; extremely high organic carbon may create a thick anoxic layer that is unsuitable for aerobic nitrifying bacteria that produce nitrate used in DNF. In contrast, too low organic carbon may promote
350 an aerobic layer unsuitable for the anaerobic denitrifying bacteria. Additionally, grasses have been shown to be extremely efficient at nutrient assimilation (Petrovic, 1990). It is plausible that much of the inorganic nitrogen that would have otherwise fluxed out of the sediments was integrated into biomass.

355 Like grasses, stormwater ponds are prolific features of developed landscapes, and yet few studies have examined their nitrogen processing capabilities. Low nitrate-DNF rates exhibited by the stormwater pond sampled in our study were low relative to the marsh but were higher than those from other studies (Gold et al., 2017; Gold et al., 2021). The pond sampled in this study is part of a restored wetland that feeds a tributary creek of the Neuse River. High DNF rates in this pond relative to others suggest that hydrological design as well as ecological connectivity could increase nitrogen removal by stormwater ponds. Flooding from the Neuse River could increase circulation to reduce stratification and prolonged anoxia that appears to favor alternate
360 nitrogen pathways. Additionally, the natural vegetation that surrounds the stormwater pond could provide a source of organic carbon to the sediments, much higher than a typical urban stormwater pond (Blaszczak et al., 2018; Hohman et al., 2021).



Following nitrate enrichment, all habitats exhibited significant increases in DNF. This type of biogeochemical response has been observed in other studies (Gold et al., 2021; Seitzinger, 1994; Smyth et al., 2015); however, DNF rates under the high nitrate condition in this study were exceptionally high, with marsh, swamp forest, and stormwater pond sediments producing the highest rates. These habitats also showcased the highest percentages of SOM. It is possible that these high SOM environments were nitrate limited and organic carbon was in excess; therefore, under low nitrate conditions there is a portion of SOM that was not used in the DNF process. This is supported by the weak linear relationship between SOM and DNF under low nitrate conditions. The significant positive linear relationship between SOM and DNF under high nitrate conditions supports that an external source of nitrate may have alleviated this limitation with abundant SOM readily available. A similar phenomenon was observed in Arango et al.'s (2007) study examining denitrification in streams in the Midwest of the US. To summarize, habitats showed increased DNF capacity in response to elevated NO_x concentrations, with high SOM environments playing the most important roles in nitrogen removal when water column NO_x concentrations are high.

The significant positive linear relationship between DNF and SOD at low nitrate concentrations is consistent with results obtained by Piehler and Smyth (2011) and Seitzinger et al. (2006). This suggests that DNF is tightly coupled with nitrification; the nitrate fueling DNF is produced *in situ*. Under high nitrate conditions, the relationship between SOD and DNF is no longer significant. This weak relationship paired with negative NO_x fluxes post-nitrate enrichment, could point to an increased importance of direct DNF, where overlying water supplies nitrate for DNF in the sediments.

4.2 Nitrogen removal during different types of storms

Comparing multiple storms that have affected North Carolina's coast revealed that HP-HW storms result in water column NO_x concentrations that are significantly lower than HP-LW, LP-HW, and LP-LW storms. It is possible that during HP-HW events, there is a compound dilution effect from both riverine discharge and wind-driven storm surge. NO_x concentrations can significantly alter biogeochemical processes, namely DNF. Therefore, it is likely that the effectiveness of nitrogen removal by the coastal landscape is dependent on the type of storm impacting the region. Results from this work suggest that flooded landscapes are permanently removing water column nitrogen through direct DNF at higher rates during HP-LW, LP-HW, and LP-LW storms compared to HP-HW storms when NO_x concentrations are relatively low and coupled nitrification-DNF is likely the dominant process.

These results largely reinforce the idea put forth by Paerl et al. (2018) where the Neuse River acts as a 'pipeline', delivering nitrogen to Pamlico Sound during these rainier events, as opposed to a 'processor' during drier events. Though NO_x concentrations during HP-HW storms were relatively low, the high volume of water during wetter storms delivers larger loads of TDN and nitrate to the estuary compared to drier storms. Reduced nitrogen removal capacity of the coastal landscape during these flood events paired with increases in nitrogen loads has implications for greater nitrogen export into Pamlico Sound,



395 especially as climate changes increase the magnitude and frequency of these rainier storms (Easterling et al., 2017; Knutson et al., 2010; Paerl et al., 2019).

Potentially exacerbating this threat to water quality is development within the Neuse River watershed. This study sheds light on how human impacts on the landscape influence distributions of nitrogen sinks as anthropogenic nitrogen sources increase. 400 As urban and suburban landscapes expand, UOS will become more abundant and their role in regulating water quality will grow. These results suggest that under low nitrate conditions, DNF rates in UOS sediments are comparable to marshes and they will play an important role during flooding from HP-HW storms, and other low nitrate scenarios. They likely will not play as large a role during other types of storms when water column nitrate concentrations are relatively high.

405 Just as developed landscapes are expanding within watersheds and along coastlines, it seems as though floodplains are infringing on these built environments (Sebastian et al., 2019; Wang et al., 2017; Wobus et al., 2017). Some experts cite a regime shift in tropical cyclone activity where annual exceedance percentages historically used to delineate floodplains (e.g., 100-year and 500-year floodplains) are no longer representative of the current conditions (Paerl et al., 2019). When assessing landcover within each storm's floodplain and the nitrogen removal by each habitat, UOS played an increasingly important role 410 during the most recent storms (Matthew, Florence, and Dorian). It is possible that their growing abundance within storm floodplains and their increased contribution to nitrogen removal informs a trend where floodplain boundaries are encroaching further inland (Knutson et al., 2013; Min et al., 2011).

In the upper NRE, when NO_x concentrations are high, more natural landscapes—including the hydrologically and ecologically 415 connected stormwater pond sampled in this study—produced the highest DNF rates. However, the limited areal extent of marshes and stormwater ponds within each storm's floodplain rendered them incapable of removing substantial nitrogen in this region. Swamp forests, on the other hand, appear to play an important role in maintaining water quality during storms. They were consistently the most extensive habitat within the storms' floodplains and, as such, made the largest contribution to nitrogen removal under both low and high nitrate conditions. Therefore, swamp forests appear to be essential for regulating 420 water quality in the NRE during storms of varying characteristics.

5 Conclusions

The results of this study provide information about nitrogen removal capacities by multiple natural habitats and urban landscape features in a flood-prone, developed, estuarine environment. All habitats showcased the ability to permanently remove nitrogen from the system under low nitrate conditions and increase nitrogen removal capacity in response to additional 425 nutrients delivered to the system. Flooded UOS can play an important role in regulating nitrogen when water column concentrations are relatively low (e.g., during rainier & windier storms). When water column nitrate concentrations are high,



more “natural” structured habitats, including marshes and swamp forest along with a somewhat unique stormwater/wetland pond, were more effective at removing nitrogen than shallow subtidal sediments and UOS. These differences in processing suggest that abundance and spatial distributions of these habitats within a floodplain can influence overall nitrogen removal capacity at the watershed scale.

Water column nutrient concentrations produced by different types of storms likely influence biogeochemical processing by flooded habitats and subsequent nitrogen export downstream and into Pamlico Sound. Results of this study suggest that flooded landscapes may be less effective at removing reactive nitrogen during HP-HW storms compared to other storm types. Low water column NO_x concentrations produced during HP-HW events, likely result in relatively low DNF rates. DNF rates are likely higher during storm events that produce relatively high water column NO_x concentrations, such as HP-LW, LP-HW, and LP-LW storms. Swamp forests are the most extensive of the habitats in this study area and play an important role in removing nitrogen and regulating water quality, regardless of storm characteristics. Management strategies should continue prioritizing swamp forest conservation in this region, as in North Carolina’s Riparian Buffer Protection Program.

As coastlines and watersheds become more developed and climate change increases the frequency and magnitude of storms and major flooding events, the export of both anthropogenic and terrigenous nitrogen will likely increase. Understanding nitrogen removal capabilities and limitations of flooded natural coastal habitats as well as those urban landscapes that will become more and more prevalent, will enable us to make informed management decisions to benefit the integrity of our coastal waters.

Data availability

Land cover data, watershed boundaries, and flood footprint data are open source and accessible online (<https://www.mrlc.gov/data>; <https://www.usgs.gov/national-hydrography/watershed-boundary-dataset>; <https://cera.coastalrisk.live/>). Wind and riverine discharge datasets are also open source and accessible online (https://www.ndbc.noaa.gov/station_page.php?station=clkn7; <https://waterdata.usgs.gov/monitoring-location/02091814/>). Data collected during the sediment core incubation experiment are available upon request.

Acknowledgements

This research was supported by funding from The National Science Foundation Growing Convergent Research Award #2021086 and the North Carolina Collaboratory award. We would like to thank Piehler lab members at the UNC Institute of Marine Sciences, Mollie Yacano and Chelsea Brown, for their assistance with sample collection and laboratory experiments. We also thank the anonymous reviewers for their helpful comments.



References

- Adame, M. F., Roberts, M. E., Hamilton, D. P., Ndehedehe, C. E., Reis, V., Lu, J., Griffiths, M., Curwen, G., & Ronan, M. (2019). Tropical Coastal Wetlands Ameliorate Nitrogen Export During Floods. *Frontiers in Marine Science*, 6. <https://doi.org/10.3389/fmars.2019.00671>
- 460 Aguilera, M. A., Tapia, J., Gallardo, C., Nunez, P., & Varas-Belemmi, K. (2020). Loss of coastal ecosystem spatial connectivity and services by urbanization: Natural-to-urban integration for bay management. *J Environ Manage*, 276, 111297. <https://doi.org/10.1016/j.jenvman.2020.111297>
- 465 Blaszczyk, J. R., Steele, M. K., Badgley, B. D., Heffernan, J. B., Hobbie, S. E., Morse, J. L., Rivers, E. N., Hall, S. J., Neill, C., Pataki, D. E., Groffman, P. M., & Bernhardt, E. S. (2018). Sediment chemistry of urban stormwater ponds and controls on denitrification. *Ecosphere*, 9(6). <https://doi.org/10.1002/ecs2.2318>
- Bowen, J. L., & Valiela, I. (2001). The ecological effects of urbanization of coastal watersheds: historical increases in nitrogen loads and eutrophication of Waquoit Bay estuaries. *Canadian Journal of Fisheries and Aquatic Sciences*, 58(8), 1489-1500. <https://doi.org/10.1139/cjfas-58-8-1489>
- 470 Boyer, J. N., Stanley, D. W., & Christian, R. R. (1994). Dynamics of NH₄⁺ and NO₃⁻ uptake in the water column of the Neuse River Estuary, North Carolina. *Estuaries*, 17(2), 361-371.
- Byers, S. C., Mills, E. L., & Stewart, P. L. (1978). A comparison of methods of determining organic carbon in marine sediments, with suggestions for a standard method. *Hydrobiologia*, 58(1), 43-47.
- Carey, R. O., & Migliaccio, K. W. (2009). Contribution of wastewater treatment plant effluents to nutrient dynamics in aquatic systems: a review. *Environ Manage*, 44(2), 205-217. <https://doi.org/10.1007/s00267-009-9309-5>
- 475 Christian, R. R., Boyer, J. N., & Stanley, D. W. (1991). Multi-year distribution patterns of nutrients within the Neuse River Estuary, North Carolina. *Marine Ecology Progress Series*, 71, 259-274.
- D'Sa, E. J., Joshi, I. D., Liu, B., Ko, D. S., Osburn, C. L., & Bianchi, T. S. (2019). Biogeochemical Response of Apalachicola Bay and the Shelf Waters to Hurricane Michael Using Ocean Color Semi-Analytic/Inversion and Hydrodynamic Models. *Frontiers in Marine Science*, 6. <https://doi.org/10.3389/fmars.2019.00523>
- 480 Davis, S. E., Cable, J. E., Childers, D. L., Coronado-Molina, C., Day, J. W., Hittle, C. D., Madden, C. J., Reyes, E., Rudnick, D., & Sklar, F. (2004). Importance of Storm Events in Controlling Ecosystem Structure and Function in a Florida Gulf Coast Estuary. *Journal of Coastal Research*, 20(4), 1198-1208. <https://doi.org/10.2112/03-0072r.1>
- Denman, E. C., May, P. B., & Moore, G. M. (2016). The Potential Role of Urban Forests in Removing Nutrients from Stormwater. *J Environ Qual*, 45(1), 207-214. <https://doi.org/10.2134/jeq2015.01.0047>
- 485 Diaz, R. J., & Rosenberg, R. (1995). Marine benthic hypoxia: A review of its ecological effects and the behavioural responses of benthic macrofauna. *Oceanography and Marine Biology*, 33, 245-303.
- Duda, A. M. (1982). Municipal point source and agricultural nonpoint source contributions to coastal eutrophication. *Journal of American Water Resources Association*, 18(3), 397-407.
- 490 Dupas, R., Delmas, M., Dorioz, J.-M., Garnier, J., Moatar, F., & Gascuel-Oudou, C. (2015). Assessing the impact of agricultural pressures on N and P loads and eutrophication risk. *Ecological Indicators*, 48, 396-407. <https://doi.org/10.1016/j.ecolind.2014.08.007>
- Easterling, D. R., Arnold, J., Knutson, T., Kunkel, K., LeGrande, A., Leung, L. R., Vose, R., Waliser, D., & Wehner, M. (2017). Precipitation change in the United States. *Climate Science Special Report: Fourth National Climate Assessment*, 1, 207-230.
- Ensign, S. H., Piehler, M. F., & Doyle, M. W. (2008). Riparian zone denitrification affects nitrogen flux through a tidal freshwater river. *Biogeochemistry*, 91(2-3), 133-150. <https://doi.org/10.1007/s10533-008-9265-9>
- 495 Eyre, B. D., & Ferguson, A. J. P. (2009). Denitrification efficiency for defining critical loads of carbon in shallow coastal ecosystems. *Hydrobiologia*, 629(1), 137-146. <https://doi.org/10.1007/s10750-009-9765-1>
- Gold, A. C., Thompson, S. P., & Piehler, M. F. (2017). Coastal stormwater wet pond sediment nitrogen dynamics. *Sci Total Environ*, 609, 672-681. <https://doi.org/10.1016/j.scitotenv.2017.07.213>
- 500 Gold, A. C., Thompson, S. P., & Piehler, M. F. (2019). The Effects of Urbanization and Retention-Based Stormwater Management on Coastal Plain Stream Nutrient Export. *Water Resources Research*, 55(8), 7027-7046. <https://doi.org/10.1029/2019wr024769>
- Gold, A. C., Thompson, S. P., & Piehler, M. F. (2021). Seasonal Variation in Nitrate Removal Mechanisms in Coastal Stormwater Ponds. *Water Resources Research*, 57(10). <https://doi.org/10.1029/2021wr029718>
- Goodrich, D. M., Boicourt, W. C., Hamilton, P., & Pritchard, D. W. (1987). Wind-induced destratification in Chesapeake Bay. *Journal of Physical Oceanography*, 17(12), 2232-2240.
- 505 Grabowski, J. H., Brumbaugh, R. D., Conrad, R. F., Keeler, A. G., Opaluch, J. J., Peterson, C. H., Piehler, M. F., Powers, S. P., & Smyth, A. R. (2012). Economic Valuation of Ecosystem Services Provided by Oyster Reefs. *BioScience*, 62(10), 900-909. <https://doi.org/10.1525/bio.2012.62.10.10>
- Groffman, P. M., Axelrod, E. A., Lemunyon, J. L., & Sullivan, W. M. (1991). Denitrification in Grass and Forest Vegetated Filter Strips. *Journal of Environmental Quality*, 20(3), 671-674. <https://doi.org/10.2134/jeq1991.00472425002000030027x>
- 510



- Groffman, P. M., & Crawford, M. K. (2003). Denitrification Potential in Urban Riparian Zones. *Journal of Environmental Quality*, 32(3), 1144-1149.
- Hampel, J. J., McCarthy, M. J., Reed, M. H., & Newell, S. E. (2019). Short Term Effects of Hurricane Irma and Cyanobacterial Blooms on Ammonium Cycling Along a Freshwater–Estuarine Continuum in South Florida. *Frontiers in Marine Science*, 6. <https://doi.org/10.3389/fmars.2019.00640>
- 515 Hill, A. R., & Cardaci, M. (2004). Denitrification and Organic Carbon Availability in Riparian Wetland Soils and Subsurface Sediments. *Soil Science Society of America Journal*, 68(1), 320-325. <https://doi.org/10.2136/sssaj2004.3200a>
- Hirsch, R. M., Moyer, D. L., & Archfield, S. A. (2010). Weighted regressions on time, discharge, and season (WRTDS), with an application to Chesapeake Bay River Inputs. *Journal of American Water Resources Association*, 46(5), 857-880. <https://doi.org/10.1111/j.1752-1688.2010.00482.x>
- 520 Hohman, S. P., Smyth, A. R., Bean, E. Z., & Reisinger, A. J. (2021). Internal nitrogen dynamics in stormwater pond sediments are influenced by pond age and inorganic nitrogen availability. *Biogeochemistry*, 156(2), 255-278. <https://doi.org/10.1007/s10533-021-00843-2>
- Howarth, R. W., & Marino, R. (2006). Nitrogen as the limiting nutrient for eutrophication in coastal marine. *Limnology and Oceanography*, 51(1 part 2), 364-376.
- 525 Knutson, T. R., McBride, J. L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J. P., Srivastava, A. K., & Sugi, M. (2010). Tropical cyclones and climate change. *Nature Geoscience*, 3(3), 157-163. <https://doi.org/10.1038/ngeo779>
- Knutson, T. R., Sirutis, J. J., Vecchi, G. A., Garner, S., Zhao, M., Kim, H.-S., Bender, M., Tuleya, R. E., Held, I. M., & Villarini, G. (2013). Dynamical Downscaling Projections of Twenty-First-Century Atlantic Hurricane Activity: CMIP3 and CMIP5 Model-Based Scenarios. *Journal of Climate*, 26(17), 6591-6617. <https://doi.org/10.1175/jcli-d-12-00539.1>
- 530 Kossin, J. P. (2018). A global slowdown of tropical-cyclone translation speed. *Nature*, 558(7708), 104-107. <https://doi.org/10.1038/s41586-018-0158-3>
- Kunkel, K. E., Easterling, D. R., Kristovich, D. A. R., Gleason, B., Stoecker, L., & Smith, R. (2010). Recent increases in U.S. heavy precipitation associated with tropical cyclones. *Geophysical Research Letters*, 37(24), n/a-n/a. <https://doi.org/10.1029/2010gl045164>
- 535 Lee, S. Y., Dunn, R. J. K., Young, R. A., Connolly, R. M., Dale, P. E. R., Dehayr, R., Lemckert, C. J., McKinnon, S., Powell, B., Teasdale, P. R., & Welsh, D. T. (2006). Impact of urbanization on coastal wetland structure and function. *Austral Ecology*, 31(2), 149-163. <https://doi.org/10.1111/j.1442-9993.2006.01581.x>
- Mallin, M. A., Posesy, M. H., McIver, M. R., Parsons, D. C., Ensign, S. H., & Alphin, T. D. (2002). Impacts and recovery from multiple hurricanes in a piedmont-coastal plain river system: human development of floodplains greatly compounds the impacts of hurricanes on water quality and aquatic life. *BioScience*, 52(11), 999-1010.
- 540 Mancino, C. F., Torello, W. A., & Wehner, D. J. (1988). Denitrification losses from Kentucky Bluegrass sod. *Agronomy Journal*, 80(1), 148-153.
- McPhillips, L. E., Groffman, P. M., Schneider, R. L., & Walter, M. T. (2016). Nutrient Cycling in Grassed Roadside Ditches and Lawns in a Suburban Watershed. *J Environ Qual*, 45(6), 1901-1909. <https://doi.org/10.2134/jeq2016.05.0178>
- 545 Miller, W. D., Harding, L. W., & Adolf, J. E. (2006). Hurricane Isabel generated an unusual fall bloom in Chesapeake Bay. *Geophysical Research Letters*, 33(6). <https://doi.org/10.1029/2005gl025658>
- Min, S. K., Zhang, X., Zwiers, F. W., & Hegerl, G. C. (2011). Human contribution to more-intense precipitation extremes. *Nature*, 470(7334), 378-381. <https://doi.org/10.1038/nature09763>
- 550 Minaudo, C., Dupas, R., Gascuel-Oudou, C., Roubeix, V., Danis, P.-A., & Moatar, F. (2019). Seasonal and event-based concentration-discharge relationships to identify catchment controls on nutrient export regimes. *Advances in Water Resources*, 131. <https://doi.org/10.1016/j.advwatres.2019.103379>
- Naden, P., Bell, V., Carnell, E., Tomlinson, S., Dragosits, U., Chaplow, J., May, L., & Tipping, E. (2016). Nutrient fluxes from domestic wastewater: A national-scale historical perspective for the UK 1800-2010. *Sci Total Environ*, 572, 1471-1484. <https://doi.org/10.1016/j.scitotenv.2016.02.037>
- 555 Nixon, S. W. (1995). Coastal marine eutrophication: A definition, social causes, and future concerns. *Ophelia*, 41(1), 199-219. <https://doi.org/10.1080/00785236.1995.10422044>
- Onorevole, K. M., Thompson, S. P., & Piehler, M. F. (2018). Living shorelines enhance nitrogen removal capacity over time. *Ecological Engineering*, 120, 238-248. <https://doi.org/10.1016/j.ecoleng.2018.05.017>
- 560 Ortiz-Zayas, J. R., Cuevas, E., Mayol-Bracero, O. L., Donoso, L., Trebs, I., Figueroa-Nieves, D., & McDowell, W. H. (2006). Urban influences on the nitrogen cycle in Puerto Rico. *Biogeochemistry*, 79(1-2), 109-133. <https://doi.org/10.1007/s10533-006-9005-y>
- Paerl, H. W., Crosswell, J. R., Van Dam, B., Hall, N. S., Rossignol, K. L., Osburn, C. L., Hounshell, A. G., Sloup, R. S., & Harding, L. W. (2018). Two decades of tropical cyclone impacts on North Carolina's estuarine carbon, nutrient and phytoplankton dynamics: implications for biogeochemical cycling and water quality in a stormier world. *Biogeochemistry*, 141(3), 307-332. <https://doi.org/10.1007/s10533-018-0438-x>



- 565 Paerl, H. W., Hall, N. S., Hounshell, A. G., Luettich, R. A., Jr., Rossignol, K. L., Osburn, C. L., & Bales, J. (2019). Recent increase in catastrophic tropical cyclone flooding in coastal North Carolina, USA: Long-term observations suggest a regime shift. *Sci Rep*, 9(1), 10620. <https://doi.org/10.1038/s41598-019-46928-9>
- Paerl, H. W., Hall, N. S., Hounshell, A. G., Rossignol, K. L., Barnard, M. A., Luettich, R. A., Rudolph, J. C., Osburn, C. L., Bales, J., & Harding, L. W. (2020). Recent increases of rainfall and flooding from tropical cyclones (TCs) in North Carolina (USA): implications for organic matter and nutrient cycling in coastal watersheds. *Biogeochemistry*, 150(2), 197-216. <https://doi.org/10.1007/s10533-020-00693-4>
- 570 Pérez-Villalona, H., Cornwell, J. C., Ortiz-Zayas, J. R., & Cuevas, E. (2015). Sediment Denitrification and Nutrient Fluxes in the San José Lagoon, a Tropical Lagoon in the Highly Urbanized San Juan Bay Estuary, Puerto Rico. *Estuaries and Coasts*, 38(6), 2259-2278. <https://doi.org/10.1007/s12237-015-9953-3>
- 575 Petrovic, A. M. (1990). The fate of nitrogenous fertilizers applied to turfgrass. *Frontiers of Environmental Quality*, 19(1), 1-14.
- Piehl, M. F., & Smyth, A. R. (2011). Habitat-specific distinctions in estuarine denitrification affect both ecosystem function and services. *Ecosphere*, 2(1). <https://doi.org/10.1890/es10-00082.1>
- Pinckney, J. L., Paerl, H. W., Harrington, M. B., & Howe, K. E. Annual cycles of phytoplankton community-structure and bloom dynamics in the Neuse River Estuary, North Carolina. *Marine Biology*, 131(2), 371-381.
- 580 Raciti, S. M., Burgin, A. J., Groffman, P. M., Lewis, D. N., & Fahey, T. J. (2011). Denitrification in suburban lawn soils. *J Environ Qual*, 40(6), 1932-1940. <https://doi.org/10.2134/jeq2011.0107>
- Reisinger, A. J., Groffman, P. M., & Rosi-Marshall, E. J. (2016). Nitrogen cycling process rates across urban ecosystems. *FEMS Microbiol Ecol*, 92(12). <https://doi.org/10.1093/femsec/fiw198>
- Rosenzweig, B. R., Groffman, P. M., Zarnoch, C. B., Branco, B. F., Hartig, E. K., Fitzpatrick, J., Forgiione, H. M., & Parris, A. (2018). Nitrogen regulation by natural systems in “unnatural” landscapes: denitrification in ultra-urban coastal ecosystems. *Ecosystem Health and Sustainability*, 4(9), 205-224. <https://doi.org/10.1080/20964129.2018.1527188>
- Rudek, J., Paerl, H. W., Mallin, M. A., & Bates, P. W. (1991). Seasonal and hydrological control of phytoplankton nutrient limitation in the lower Neuse River Estuary, North Carolina. *Marine Ecology Progress Series*, 75(2), 133-142.
- Sebastian, A., Gori, A., Blessing, R. B., van der Wiel, K., & Bass, B. (2019). Disentangling the impacts of human and environmental change on catchment response during Hurricane Harvey. *Environmental Research Letters*, 14(12). <https://doi.org/10.1088/1748-9326/ab5234>
- 590 Seitzinger, S., Harrison, J. A., Böhlke, J. K., Bouwman, A. F., Lowrance, R., Peterson, B., Tobias, C., & Drecht, G. V. (2006). Denitrification across landscapes and waterscapes: A synthesis. *Ecological Applications*, 16(6), 2064-2090.
- Seitzinger, S. P. (1994). Linkages between organic matter mineralization and denitrification in eight riparian wetlands. *Biogeochemistry*, 25(1), 19-39.
- 595 Smyth, A. R., Piehl, M. F., Grabowski, J. H., & Frid, C. (2015). Habitat context influences nitrogen removal by restored oyster reefs. *Journal of Applied Ecology*, 52(3), 716-725. <https://doi.org/10.1111/1365-2664.12435>
- Sun, H., Jiang, J., Cui, L., Feng, W., Wang, Y., & Zhang, J. (2019). Soil organic carbon stabilization mechanisms in a subtropical mangrove and salt marsh ecosystems. *Sci Total Environ*, 673, 502-510. <https://doi.org/10.1016/j.scitotenv.2019.04.122>
- 600 Velinsky, D. J., Paudel, B., Quirk, T., Piehl, M., & Smyth, A. (2017). Salt Marsh Denitrification Provides a Significant Nitrogen Sink in Barnegat Bay, New Jersey. *Journal of Coastal Research*, 78, 70-78. <https://doi.org/10.2112/si78-007.1>
- Wang, H., Marshall, C. W., Cheng, M., Xu, H., Li, H., Yang, X., & Zheng, T. (2017). Changes in land use driven by urbanization impact nitrogen cycling and the microbial community composition in soils. *Sci Rep*, 7, 44049. <https://doi.org/10.1038/srep44049>
- Wang, W., Haver, D., & Pataki, D. E. (2013). Nitrogen budgets of urban lawns under three different management regimes in southern California. *Biogeochemistry*, 121(1), 127-148. <https://doi.org/10.1007/s10533-013-9942-1>
- 605 Wasson, K., Jeppesen, R., Endris, C., Perry, D. C., Woolfolk, A., Beheshti, K., Rodriguez, M., Eby, R., Watson, E. B., Rahman, F., Haskins, J., & Hughes, B. B. (2017). Eutrophication decreases salt marsh resilience through proliferation of algal mats. *Biological Conservation*, 212, 1-11. <https://doi.org/10.1016/j.biocon.2017.05.019>
- Wengrove, M. E., Foster, D. L., Kalnejais, L. H., Percuoco, V., & Lippmann, T. C. (2015). Field and laboratory observations of bed stress and associated nutrient release in a tidal estuary. *Estuarine, Coastal and Shelf Science*, 161, 11-24. <https://doi.org/10.1016/j.ecss.2015.04.005>
- Wetz, M. S., & Paerl, H. W. (2008). Estuarine Phytoplankton Responses to Hurricanes and Tropical Storms with Different Characteristics (Trajectory, Rainfall, Winds). *Estuaries and Coasts*, 31(2), 419-429. <https://doi.org/10.1007/s12237-008-9034-y>
- Wobus, C., Gutmann, E., Jones, R., Rissing, M., Mizukami, N., Lorie, M., Mahoney, H., Wood, A. W., Mills, D., & Martinich, J. (2017). Climate change impacts on flood risk and asset damages within mapped 100-year floodplains of the contiguous United States. *Natural Hazards and Earth System Sciences*, 17(12), 2199-2211. <https://doi.org/10.5194/nhess-17-2199-2017>
- 615