



A revised parameterization for aerosol, cloud and precipitation pH for use in chemical forecasting systems (EQSAM4Clim-v12)

Swen Metzger¹, Samuel Rémy², Jason E Williams³, Vincent Huijnen³, and Johannes Flemming⁴

¹ResearchConcepts Io GmbH, Freiburg, Germany

²HYGEOS, Lille, France

³R and D Weather and Climate modeling, Royal Netherlands Meteorological Institute, De Bilt, Netherlands

⁴European Centre for Medium Range Weather Forecasts, Reading, UK and Bonn, Germany

Correspondence: Swen Metzger (sm@researchconcepts.io)

Abstract. The Equilibrium Simplified Aerosol Model for Climate version 12 (EQSAM4Clim-v12) has recently been revised to provide an accurate and efficient method for calculating the acidity of atmospheric particles. EQSAM4Clim is based on an analytical concept that is not only sufficiently fast for numerical weather prediction (NWP) applications, but also free of numerical noise, which makes it attractive also for air quality forecasting. EQSAM4Clim allows the calculation of aerosol composition based on the gas-liquid-solid and the reduced gas-liquid partitioning with the associated water uptake for both cases, and can therefore provide important information about the acidity of the aerosols. Here we provide a comprehensive description of the recent changes made to the aerosol acidity parameterization (referred to a version 12) which builds on the original EQSAM4Clim. We evaluate the pH improvements using a detailed box-model and compare against previous model calculations and both ground-based and aircraft observations from US and China covering different seasons and scenarios. We show that, in most cases, the simulated pH is within reasonable agreement with the results of the E-AIM reference model and of satisfactory accuracy.

1 Introduction

In order to address the relevance of gas-aerosol partitioning and aerosol water for climate and air quality studies, the Equilibrium Simplified Aerosol Model (EQSAM) was developed as a compromise between numerical speed and accuracy (Metzger et al., 2002). EQSAM has been widely used in many air quality and climate modelling systems worldwide (Metzger et al., 2018), including the IFS (Flemming et al., 2015) and the OpenIFS (Huijnen et al., 2022). Recently, the EQSAM version for Climate Applications (EQSAM4Clim) (Metzger et al., 2016a) has been implemented in IFS-COMPO (see the accompanying paper). In contrast to EQSAM, EQSAM4Clim is entirely based on a compound specific single-solute coefficient (ν_i), which was introduced in Metzger et al. (2012) to accurately parameterise the single solution hygroscopic growth, considering the Kelvin effect. This ν_i -approach accounts for the water uptake of concentrated nanometre-sized particles up to dilute solutions, i.e. from the compounds relative humidity of deliquescence (RHD) up to supersaturation (Köhler theory). EQSAM4Clim extends the ν_i -approach to multicomponent mixtures, including semi-volatile ammonium compounds and major crustal elements.



The advantage of EQSAM4Clim is that the entire gas–liquid–solid aerosol phase partitioning and water uptake, including major mineral cations, is solved analytically without iterations and thus computationally very efficient. This makes EQSAM4Clim suited not only for climate simulations, but also applicable to high resolution Numerical Weather Predictions (NWP) coupled with comprehensive atmospheric chemistry providing global values of particulate matter, as done in the Copernicus Atmosphere Monitoring Service (CAMS, Peuch et al. (2022); Rémy et al. (2022)) for example, using the ECMWF Integrated Forecasting System (IFS).

Previously, the use of EQSAM4Clim has undergone a rigorous assessment across different time scales, through a comparison with various observations and reference simulations on climate time scales using more than a decade of independent observations (e.g. Metzger et al. (2018)). Moreover, a comparison of simulated Aerosol Optical Depth (AOD) has been made against various satellite data at NWP time scales to validate the Polar Multi-sensor Aerosol properties (PMAp) AOD product version 2 AOD at an 1 hourly time resolution (Metzger et al., 2016b). EQSAM4Clim has been also used as part of air quality assessments through, e.g., the 2019 European Monitoring and Evaluation Programme (EMEP) report on transboundary particulate matter, photo-oxidants, acidifying and eutrophying components (Fagerli et al., 2019), and evaluated in the air quality modeling system CAMx over the continental US with 12 km grid resolution for winter and summer months (Koo et al., 2020). On all time scales, it was found that EQSAM4Clim accurately parameterises the gas/liquid/solid aerosol partitioning and associated aerosol water uptake sufficiently fast and free of numerical noise. This is due to its unique analytical structure, which makes it particularly also attractive for air quality assessments such as those provided by CAMS. Most recently, the latest version of EQSAM4Clim has been implemented in IFS-COMPO as presented in Metzger et al. (2022) and Metzger et al. (2023). A more comprehensive evaluation of the performance on global pH values and resulting effects on Particulate matter in IFS-COMPO will be presented in two companion papers.

This Technical Note provides a description of the improved aerosol acidity parameterization applied in EQSAM4Clim-v12. We show an extensive validation against reference model calculations using Extended Aerosol Inorganics Model (E-AIM) as described in Wexler and Clegg (2002) and Friese and Ebel (2010), using the detailed case study on aerosol acidity provided by Pye et al. (2020).

2 Description of EQSAM4Clim-v12

The overall gas/liquid/solid partitioning and aerosol water uptake parameterization is the same as described and evaluated in Metzger et al. (2012) and Metzger et al. (2016a), with further evaluation being provided in Metzger et al. (2018). Here we limit the description to those new features added to previous versions.

2.1 General Features

A schematic of the various input parameters needed for use in EQSAM4Clim is shown in Figure 1, where chemical species from each phase type is given. EQSAM4Clim is based on a compound specific single-solute coefficient (ν_i), which was introduced in Metzger et al. (2012) for single solute solutions and extended to multi-component mixtures by Metzger et al. (2016a) to




Figure 1. A schematic of the components included in EQSAM4Clim.

55 include semi-volatile ammonium (NH_4^+) compounds and major crustal elements. A feature of the ν_1 -approach is that the entire gas–liquid–solid aerosol phase partitioning and water uptake can be solved analytically without iterations, and hence without numerical noise.

EQSAM4Clim takes as input (i) the meteorological parameters air temperature (T) and relative humidity (RH), (ii) the aerosol precursor gases, i.e., major oxidation products of natural sources and anthropogenic air pollution represented by ammonia (NH_3), hydrochloric acid (HCl), nitric acid (HNO_3), sulphuric acid (H_2SO_4), and (iii) the ionic aerosol concentrations, i.e., lumped (both liquid and solid) anions, sulphate (SO_4^{2-}), bi-sulphate (HSO_4^-), nitrate (NO_3^-), chloride (Cl^-), and lumped (liquid+solid) cations, NH_4^+ , sodium (Na^+), potassium (K^+), magnesium (Mg^{2+}) and calcium (Ca^{2+}).

The equilibrium aerosol composition and aerosol Associated Water mass (AW) is calculated by EQSAM4Clim through the neutralization of anions by cations, which yields numerous salt compounds, i.e., the sodium salts Na_2SO_4 , NaHSO_4 , NaNO_3 , NaCl , the potassium salts K_2SO_4 , KHSO_4 , KNO_3 , KCl , the ammonium salts $(\text{NH}_4)_2\text{SO}_4$, NH_4HSO_4 , NH_4NO_3 , NH_4Cl , the magnesium salts MgSO_4 , $\text{Mg}(\text{NO}_3)_2$, MgCl_2 , and the calcium salts CaSO_4 , $\text{Ca}(\text{NO}_3)_2$, CaCl_2 . All salt compounds (except CaSO_4) can partition between the liquid and solid aerosol phase, depending on T, RH, AW and the temperature-dependent Relative Humidities of Deliquescence of (a) single solute compound solutions (RHD) and (b) of mixed salt solutions (Metzger et al., 2016a).



70 Based on the RHD of the single solutes, the (mixed) solution liquid/solid partitioning is calculated, whereby all compounds for which the RH is below the RHD are assumed to be precipitated, such that a solid and liquid phase can coexist. The liquid-solid partitioning is strongly influenced by mineral cations and in turn largely determines the aerosol pH (Sect. 3).

EQSAM4Clim estimates the concentration of the hydronium ion (H^+) [$\text{mol}/\text{m}^3(\text{air})$] and, subsequently, the pH of the solution from electroneutrality (Z^0 [$\text{mol}/\text{m}^3(\text{air})$]) after neutralization of all anions by all cations in the system (following
75 the neutralization reaction order given by Table 3 of Metzger et al. (2016a)). Note that the auto dissociation of H_2O is taken into account, but currently no dissolution and dissociation of aerosol precursor gases such as sulphur dioxide (SO_2), nitric acid (HNO_3), hydrogen chloride (HCl), or ammonia (NH_3) is taken into account, as this is typically considered in the aqueous phase chemistry module of any global chemistry forecast model. The initial $H^{+,0}$ concentration [$\text{mol}/\text{m}^3(\text{air})$] after cation-anion neutralization is obtained from:

$$80 \quad Z^0 = tAnions - tCations = \sum_i [Z^-]_i - \sum_j [Z^+]_j \quad (1)$$

$$[H^{+,0}] = Z^0 = 2[SO_4^{2-}] + [HSO_4^-] + [NO_3^-] + [Cl^-] - [K^+] - 2[Ca^{2+}] - 2[Mg^{2+}] - [Na^+] - [NH_4^+] \quad (2)$$

with $tAnions$ and $tCations$ (hereafter referred to as $tCAT$) representing the total of all anions and cations [$\text{mol}/\text{m}^3(\text{air})$], respectively and where $[H^{+,0}]$ denotes the initial hydronium ion concentration per volume air and which also depends on the auto-dissociation of H_2O (K_w) [$\text{mol}^2/\text{kg}^2(\text{water})$]. This is derived from Eq. (3) considering the temperature dependency as
85 widely assumed in equilibrium models.

$$K_w = 1.010 \times 10^{-14} \cdot \exp\left(-22.52 \cdot \left(\frac{T_0}{T} - 1\right) + 26.920 \cdot A_T\right) \quad \text{where} \quad A_T = \left(1 + \log\left(\frac{T_0}{T}\right) - \frac{T_0}{T}\right), \quad (3)$$

with $T_0 = 298K$.

2.2 Updates to the acidity component

2.2.1 Dependency of H^+ on the Chemical Domain

90 The neutralization equation does not correct for non-ideal solutions, such as described in Pye et al. (2020) and the references therein. For that purpose, with v12, we introduce for EQSAM4Clim a new factor XN , which is dependant on the aerosol composition, to correct the initial $[H^{+,0}]$ (Eq. 2). XN is obtained from:

$$XN = [X]/[Y] \quad (4)$$

with X denoting the sum of all anions noted above, while $Y = tNH_4$, i.e., the sum of tNH_3 and tNH_4^+ . XN is applied without
95 further scaling factors for ranges of $XN < 0.9$ with ambient temperatures below 293K.

For cases outside this range, XN needs to be scaled by 10 and multiplied by the factor N given in Table 1, in order to account for chemical processes which are not resolved by the parameterizations (particularly concerning HSO_4^- and free H_2SO_4).



Table 1. H^+ correction factors introduced with EQSAM4Clim-v12 for the chemical domains introduced in Metzger et al. (2016a).

Domain	Characterization	Regime	Correction factor N	Relation
D1	CATION RICH	$tCAT - tNH_4 \geq tSO_4$	N=1	XN
D2	SO_4^{2-} NEUTRAL	$tCAT \geq tSO_4$	N=1	XN
D3	SO_4^{2-} RICH	$tCAT \geq tHSO_4$ AND $tCAT < tSO_4$	N=1e1	XN
D4	SO_4^{2-} VERY RICH	$tCAT \geq MIN$ AND $tCAT < tHSO_4$	N=1e3	–

Following Table 2 of Metzger et al. (2016a), four chemical domains are considered to correct $[H^{+,0}]$ obtained with Eq. (2). No additional correction ($N = 1$) is needed for the neutral cases (D1-D2), i.e. where cations are in excess of total SO_4^{2-} , thus preventing the formation of all HSO_4^- salts (see Table 1 of Metzger et al. (2016a)). For the SO_4^{2-} rich case (D3), XN and N from Table 1 are multiplied, while for the SO_4^{2-} very rich case (D4), only a constant correction factor (N) is applied to correct Eq. (2), where tCAT denotes the sum of cations given in Eq. (2), t SO_4 is the sum of all SO_4^{2-} , including HSO_4^- and H_2SO_4 .

Additionally, we consider three cases for estimating the H^+ concentration, according to the possible solutions of Eq. (1), i.e.:

$$Z^* < 0 \quad | \quad [H^{+,*}] = \frac{LWC_{tot}}{10^{(7.0 + \log(-Z^* \cdot \frac{10^4}{LWC_o \cdot \mu_s^o}))}} \cdot \mu_s^o \quad (5a)$$

$$Z^* = 0 \quad | \quad [H^{+,*}] = [H^{+,neutral}] \times 10^{-6} \quad (5b)$$

$$Z^* > 0 \quad | \quad [H^{+,*}] = Z^* \times 10^{-6} \quad (5c)$$

with LWC_{tot} being the total Liquid Water Content [$kg(H_2O)/m^3(air)$] as defined below in Eqs. (9a-9d). $LWC_o = 1$ [$kg/m^3(air)$] and $\mu_s^o = 1$ [$mol/kg(H_2O)$], a reference solution and reference molality, respectively, to match units (Metzger et al., 2012, 2016a; Pye et al., 2020). Z^* is given by Eq. (6) and denotes the sum of our initial hydrogen concentration $[H^{+,0}]$ and $[H^{+,neutral}]$, an effective hydrogen concentration in a neutral solution (pH=7), which is given by Eq. (7), but empirically derived for our parameterization:

$$Z^* = [H^{+,neutral}] + [H^{+,0}] \quad (6)$$

$$[H^{+,neutral}] = \frac{B \cdot LWC_o \cdot K_w^{0.5}}{(1.0 - RH^2)} \quad (7)$$

with K_w from Eq. (3), a constant $B = 1/(\mu_s^o \cdot m_w) = 55.51$ [–], the molar mass of water, m_w [kg/mol], and the reference molality μ_s^o [$mol/kg(H_2O)$] and a reference solution $LWC_o = 1$ [$kg/m^3(air)$] to match units; RH denotes the fractional relative humidity [0-1].

Finally, the H^+ concentration of a given solution is obtained from:

$$[H^+] = [H^{+,*}] \cdot XN \quad (8)$$



2.2.2 Dependency of pH on the Liquid Water Content

125 For EQSAM4Clim-v12, five different pH values can be computed from the revised H^+ [$\text{mol}/\text{m}^3(\text{air})$] computation (Sect. 2.2.1) for diagnostic output. Therefore, EQSAM4Clim-v12 allows the differentiation of the various LWC [$\text{kg}(\text{H}_2\text{O})/\text{m}^3(\text{air})$] values associated with different type of atmospheric aerosols, haze/fog, or cloud droplets contained in the troposphere as defined in Eqs. (9a-9e):

$$pH_{equil} = -\log_{10} \left(\frac{[H^+]}{LWC_{equil}} \cdot \frac{1}{\mu_s^o} \right) \quad (9a)$$

130

$$pH_{noneq} = -\log_{10} \left(\frac{[H^+]}{LWC_{noneq}} \cdot \frac{1}{\mu_s^o} \right) \quad (9b)$$

$$pH_{cloud} = -\log_{10} \left(\frac{[H^+]}{LWC_{cloud}} \cdot \frac{1}{\mu_s^o} \right) \quad (9c)$$

135

$$pH_{precip} = -\log_{10} \left(\frac{[H^+]}{LWC_{precip}} \cdot \frac{1}{\mu_s^o} \right) \quad (9d)$$

$$pH_{total} = -\log_{10} \left(\frac{[H^+]}{LWC_{total}} \cdot \frac{1}{\mu_s^o} \right) \quad (9e)$$

Here, (i) LWC_{equil} [$\text{kg}(\text{H}_2\text{O})/\text{m}^3(\text{air})$] denotes the equilibrium water content calculated within EQSAM4Clim (from Eq.(22) in Metzger et al. (2016a)), (ii) LWC_{noneq} is the aerosol liquid water content associated with aerosol species not considered in the equilibrium computations of EQSAM4Clim (e.g., from chemical aging of pre-existing organic or black carbon particles as used e.g. in Metzger et al. (2016b) and Metzger et al. (2018)), (iii) LWC_{cloud} denotes the cloud liquid water content, (iv) LWC_{precip} denotes the liquid water content of a given precipitation flux and finally (v) LWC_{total} denotes the sum of LWC of Eqs. (9a-9d), where $\mu_s^o = 1$ [$\text{mol}/\text{kg}(\text{H}_2\text{O})$] is the reference molality to match units. The pH values of Eqs. (9b-9e) are an optional output feature and requires the corresponding input to EQSAM4Clim-v12 (e.g., in any 3-D application these are provided by the forecasting model). It is important to note that all pH and H^+ values are only for diagnostic output, as these values are not used within EQSAM4Clim.

In contrast to other aerosol equilibrium models such as E-AIM, EQSAM4Clim has a fully analytical structure which does not depend on the pH. However, accounting for the different pH values is important for air quality and climate applications, because of the influence of solution pH on aqueous phase chemistry in terms of SO_4^{2-} production and the subsequent deposition processes. Also note that in the case where an accurate pH calculation is needed, reference calculations from E-AIM should be considered instead. With EQSAM4Clim-v12 we try to find a compromise between computational speed and accuracy, so not always the pH parameterization might be applicable, although we show EQSAM4Clim-v12 performs well over a wide range of atmospheric conditions (see Sect. 3).



155 3 Results and Evaluation

In this section, we present comparisons of the revised pH parameterization of EQSAM4Clim-v12 against a wide range of data from different measurement sites and field campaigns spanning different locations for different years, where this data has been used in the comprehensive pH review study of Pye et al. (2020). In Pye et al. (2020) five distinct cases were defined and used to evaluate the simulated pH of various thermodynamic models applied in large-scale models. We therefore select the same
160 observational data for input to the box model calculations and also use the E-AIM model output as a reference to evaluate the revised pH parameterization. It should be noted that E-AIM is much too computationally expensive to be applied in a large-scale atmospheric model. Details of the five cases used are provided below (see also Table 5 of Pye et al. (2020) and the references therein), i.e., here sorted by complexity of the chemical system with respect to the aerosol composition:


– Europe: Cabauw Experimental Site for Atmospheric Research (CESAR), Cabauw (51.970° N, 4.926° E), The Nether-
165 lands, 2 May 2012–4 Jun 2013, 2646 data points, composition including:



– Asia: Measurement site Tianjin (39.7° N, 117.1° E), China, 9–22 Aug 2015, 241 data points, composition including:



– SE US: Southern Oxidant and Aerosol Study (SOAS) campaign, Centreville (32,9° N, 87,1° W), US, 6 Jun–14 Jul 2013,
170 787 data points, reduced composition (no mineral cations):



– SW US: California Nexus (CalNex) campaign, Pasadena (34,15° N, 118,1° W), CA, USA, 17 May–15 Jun 2010, 493
data points, reduced composition (no mineral cations, and no sodium):



– NE US: Wintertime Investigation of Transport, Emissions, and Reactivity (WINTER) campaign, Eastern US aloft, 3 Feb
175 2015, 3613 data points, reduced composition (no mineral cations, and no sodium):



A total of more than 7700 data points are available for evaluation from these campaigns covering a wide range of RH (above 20 %RH) and T (\approx 250 to 310 K). Moreover, relevant input is provided for assessing the performance over a complete year
180 (Cabauw), summertime (e.g. SOAS) and wintertime (WINTER). Note that only the correction factors needed for the revised H^{+} computations (shown in Table 1) have been iteratively derived by comparing the diagnostic pH output of EQSAM4Clim-v12 with the reference pH computations of E-AIM for these five cases (using error minimizing on the log-scale), while the water uptake calculation is identical to that described in Metzger et al. (2016a).



3.1 Aerosol pH

185 To evaluate the revised pH parameterization of EQSAM4Clim-v12, we compare the resulting values against the output of the E-AIM reference model for the five field campaign cases as discussed in Pye et al. (2020). In Figure 2 the results of EQSAM4Clim-v12 and E-AIM are shown and compared to measurements at the Cabauw Experimental Site for Atmospheric Research (CESAR, Guo et al. (2018b)), for the period of May 2012 - June 2013. The AW and aerosol pH are shown together with the corresponding T and the RH data, which are used as meteorological input to this box modelling study together, with
190 the ion concentrations taken from the reference study. There is a seasonal cycle in T throughout the year, with a typical range in RH of between 60-90% with the measurement station being representative of a polluted rural site. For pH, the spread simulated by EQSAM4Clim-v12 ranges from pH 1.8-4.5 which is close to that from E-AIM, whose minimum pH is around 2.0 in a range between 2.0-5.0 i.e. less acidic. Also the aerosol water predictions of both models compare well throughout all seasons, with only a few noticeable exceptions in spring 2013 (around step 2000). Note the input data here is unfiltered and may include a
195 few outliers that are not valid. Also note that the pH_F refers to the free- H^+ approximation of pH which is only included for completeness, but not further used and discussed here (we refer the interested reader to Pye et al. (2020)).

Figures 3–5 show similar comparisons for summertime for the Tianjin (Shi et al. (2019)), SOAS (Alabama Forest, US; Guo et al. (2015)) and CalNex (Pasadena, US; Guo et al. (2018a)) campaigns, representing both urban and forest scenarios between the years 2010 and 2015. The range in T for these campaigns is typically limited to between 290-310K, with distinct signatures
200 of diurnal variability in the RH. This results in similar variability in the AW content, with the pH range in SOAS (-1.0 to 2.0) being order of magnitude more acidic than either CalNex (1.0-5.0) or Tianjin (2.0-5.0). Again the spread in the pH values from EQSAM4Clim-v12 is here similar to that of the E-AIM reference model and only a bit wider for the CalNex case compared to the Cabauw case, due to limitations of the bi-sulfate/sulfate partitioning of the EQSAM4Clim version.

Finally for wintertime under polluted conditions we use the data from one flight taken as part of the WINTER flight campaign
205 (US East Coast; Guo et al. (2016)). Here EQSAM4Clim-v12 is tested for lower temperatures across a wide range of RH values at various altitudes with high values of sea-salt. Figure 6 shows comparisons using data from the flight taken on 3rd February 2015. Both EQSAM4Clim-v12 pH and E-AIM simulate low pH values estimates, with similar variability and correlated well with respect to pH values < 0.0 .

Figure 7 shows a comparison of the EQSAM4Clim-v12 pH results of the previous version 10 (left) (used e.g. in (Fagerli et al., 2019) and (Koo et al., 2020)) and the current version 12 (right) versus the pH results of E-AIM for all five cases. Clearly,
210 the pH results of EQSAM4Clim-v12 pH are closer to E-AIM compared to the v10, now more closely following the one-by-one line for a wide range of atmospheric conditions, although some scatter still remains. Note that this scatter is acceptable for the EQSAM4Clim parameterization concept. A more explicit treatment of the phase partitioning will be subject of a follow-up study.




Figure 2. First case study: CESAR site, Cabauw, The Netherlands, 2 May 2012–4 Jun 2013, 2646 data points. The results of EQSAM4Clim-v12 (green, squares) in comparison with E-AIM (pink, cross) using the data provided by Pye et al. (2020). Both models use the T [K] (panel a) and RH [0-1] (panel b) together with the lumped ion concentrations [$\mu\text{g}/\text{m}^3$ (air)] of Mg^{2+} , Ca^{2+} , K^+ , Na^+ , $\text{HCl}+\text{Cl}^-$, $\text{NH}_3+\text{NH}_4^+$, $\text{HNO}_3+\text{NO}_3^-$, $\text{H}_2\text{SO}_4+\text{SO}_4^{2-}+\text{HSO}_4^-$ as input to calculate the aerosol water mass, H_2O [$\mu\text{g}/\text{m}^3$ (air)] (panel c) and aerosol pH [-] (panel d), assuming the metastable aerosol phase (no solid/liquid, only gas/liquid partitioning aerosol partitioning). Additionally, the E-AIM output for the free pH (pH_F , orange X) is included (see Pye et al. (2020)).




Figure 3. Second case study: Tianjin, China, 9–22 Aug 2015, 241 data points, using the aerosol system: Mg^{2+} , Ca^{2+} , K^+ , Na^+ , $HCl+Cl^-$, $NH_3+NH_4^+$, $HNO_3+NO_3^-$, $H_2SO_4+SO_4^{2-}+HSO_4^-$.




Figure 4. Third case study: SOAS campaign, Centreville, US, 6 Jun–14 Jul 2013, 787 data points, using the aerosol system: Na^+ , $HCl+Cl^-$, $NH_3+NH_4^+$, $HNO_3+NO_3^-$, $H_2SO_4+SO_4^{2-}+HSO_4^-$.




Figure 5. Fourth case study: CalNex campaign, Pasadena, CA, USA, 17 May–15 Jun 2010, 493 data points, using the aerosol system: $\text{HCl}+\text{Cl}^-$, $\text{NH}_3+\text{NH}_4^+$, $\text{HNO}_3+\text{NO}_3^-$, $\text{H}_2\text{SO}_4+\text{SO}_4^{2-}+\text{HSO}_4^-$.




Figure 6. Fifth case study: WINTER campaign, Eastern US aloft, 3 Feb 2015, 3613 data points, using the aerosol system: $\text{HCl}+\text{Cl}^-$, $\text{NH}_3+\text{NH}_4^+$, $\text{HNO}_3+\text{NO}_3^-$, $\text{H}_2\text{SO}_4+\text{SO}_4^{2-}+\text{HSO}_4^-$.




Figure 7. Comparison of the EQSAM4Clim pH results of v10 (panel a) and v12 (panel b) versus the pH results of E-AIM for all five cases. The CPU consumption per step is included for each case. Chip: Apple M1 Ultra; Memory: 128 GB; llvm-11/flang compiler with O3.

Table 2. Statistical metrics for the pH results of EQSAM4Clim pH results of v12 (left) and E-AIM (right column).

Campaign	Data Min	Data Max	Data Mean	Std.-Dev.	Bias	Corr.	Count				
Cabauw	1.308	2.000	4.906	5.617	3.575	3.448	0.493	0.521	0.127	0.829	2646
Tianjin	2.171	1.921	3.270	4.565	2.838	2.743	0.250	0.389	0.095	0.595	241
SOAS	-0.719	-0.908	2.622	1.909	0.988	0.763	0.640	0.534	0.225	0.564	787
CalNex	0.428	0.844	3.418	2.836	1.907	1.957	0.649	0.288	-0.05	0.731	493
WINTER	-1.000	-0.996	3.609	3.472	0.934	1.019	0.936	0.831	-0.085	0.874	3613

215 To scrutinize the sensitivity and computational costs of these results, the results of two EQSAM4Clim versions including
 the CPU consumption per step are given in the panels of Figure 7 for each case. Comparing the two EQSAM4Clim versions
 (left and right panel) shows (a) that the pH results differ mostly for the Cabauw, Tianjin and SOAS campaigns, which represent
 different aerosol compositions and neutralization levels as defined by where the measurement campaign took place. While the
 Cabauw and Tianjin campaigns represent the most complex aerosol system with SO_4^{2-} being fully neutralized (Sect. 3), where
 220 both locations are affected by anthropogenic precursors which undergo gas/aerosol partitioning. Conversely, data from the
 SOAS, Calnex and the WINTER campaigns represent cases where SO_4^{2-} is not fully neutralized. Especially, the measurements
 from CalNex and the flight during the WINTER campaign represent often highly acidic cases.

225 Additionally, comparing cases shows that (b) the variability in the observed pH ranges across campaigns exceeds the vari-
 ability in pH simulated by the different modeling code versions. For instance, the pH values are for the WINTER campaign
 generally much lower compared to e.g., the Cabauw campaign, which shows throughout all results the highest pH values,
 reflecting the predominance of cations in the aerosol system for the Cabauw case.

Table 2 summarizes the key metrics and shows for each campaign the minimum and maximum pH value, together with
 the data mean and standard deviation for EQSAM4Clim (v12) and E-AIM, as well as the correlation of both. While the data



mean is with a variation of less than 0.25 pH units generally satisfactorily close for all campaigns, the correlation coefficient is
230 only above 0.7 for the Cabauw, CalNex and WINTER campaign. Tianjin, which represents besides Cabauw the most complex
aerosol system, shows a slightly lower correlation coefficient of 0.6, while SOAS is with a value of 0.56 at the lower end, due
to the influence of sulfate/bi-sulfate partitioning. Bi-sulfates are not always captured in the gas/liquid partitioning compared
to cases which include semi-volatile compounds (Cabauw, Tianjin, WINTER). Also note that the correlation coefficient is
strongly influenced by the number of data points, such that the WINTER and Cabauw cases are statistically more significant.
235 This complexity of the Cabauw data is also reflected in the highest computing consumption per step (where CPU/step values
are given in the legend within each panel of Figure 7), while the WINTER campaign represents the least complex system (no
cations and low temperatures) and, therefore, requires also the least CPU time. Note that there is some uncertainty in these
numbers due to the load imbalance of the system ($\leq 1\%$), while the CPU consumption for EQSAM4Clim-v10 is higher due to
the fact that double precision is used. For EQSAM4Clim-v12, the choice of precision is optional and single precision is used
240 throughout this work, since this alone can speed up the computations of up to 50% for these run-time optimized cases.

3.2 Application to IFS

Figure 8 extends Figure 2 by showing an example of application and implementation of EQSAM4Clim-v12 into a compre-
hensive high resolution atmospheric chemistry NWP forecasting system, with the chosen model being IFS-COMPO (Peuch et al.,
2022). We use a version similar to that described in Rémy et al. (2022), where the gaseous precursors such as HNO_3 are derived
245 using the chemistry scheme given in Williams et al. (2022). The implementation provides global pH values, whose impact on
PM_{2.5} will be evaluated in a companion paper. Here we only compare the new optional IFS pH results using a 3 hourly output
frequency with those from the EQSAM4Clim-v12 and E-AIM box models at an hourly frequency for the Cabauw case as an
example. Note that only a qualitative comparison is possible due to the cumulative effects of the different time averages, the
difference in resolution (with IFS-COMPO being ran at $\approx 25\text{km}$ scale) and in that the IFS-COMPO pH results are representa-
250 tive for 2019, while the box models show the results for the year 2012/2013. Nevertheless, overall the seasonal trend of the pH
values is captured rather well. During summertime (June - September) the IFS-COMPO pH values are on average less acidic
than those calculated in the box models, while for the autumn and winter months the agreement is on average very good. This
evaluation furthermore illustrates that uncertainties to the input quantities (aerosol composition, relative humidity, temperature)
dominate the overall uncertainty, indicating that EQSAM4Clim is fit for purpose.




Figure 8. Figure 2 extended to include IFS (3 hourly averages) in comparison with EQSAM4Clim-v12 and E-AIM box model results.



255 4 Conclusions

In this technical note we have provided a description of the revised EQSAM4Clim-v12 pH parameterization developed for use in regional and global chemical forecasting systems. Using a range of diverse case studies we have performed box model calculations with the results being compared against those from the E-AIM reference model calculations, covering a range of seasons and scenarios ranging from forest measurements to maritime seaboard measurements. Generally, the pH values are mostly within the range given by E-AIM and now more closely following the one-by-one line for a wide range of atmospheric conditions compared to the previous EQSAM4Clim version (v10). Although some deviations of the EQSAM4Clim-v12 pH estimates are noted, the scatter is acceptable for the EQSAM4Clim parameterization concept. The case studies reveal that the pH results of the revised parameterization provide satisfactory representation for the most complex aerosol cases, i.e., the sulfate neutral conditions which are characterized by the gas/aerosol partitioning of semi-volatile ammonium compounds. Additionally, EQSAM4Clim has a low cost with respect to the overall CPU consumption across aerosols with a range of composition complexity (as seen across different campaigns). This provides confidence that the revised pH parameterization is suitable for applications to large-scale chemical forecasting systems where near-real time results are mandatory.

Code availability. The current version of model is available from GitHub: <https://github.com/rc-io/eqsam> under the licence CC-BY-SA-4.0. The exact version of the model used to produce the results used in this paper is archived on Zenodo (Metzger, 2023), as are input data and scripts to run the model and produce the plots for all the simulations presented in this paper (Metzger, 2023).

Author contributions. SM drafted the paper and developed and implemented the revised pH formulation of EQSAM4Clim-v12 into IFS; SR, VH, SM, JW and JF maintain and carry out general aerosol and trace gas developments on IFS-COMPO; all contributed to drafting and revising this article.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Geoscientific Model Development.

Acknowledgements. This work is supported by the Copernicus Atmospheric Monitoring Services (CAMS) program managed by ECMWF on behalf of the European Commission. We acknowledge the Pye et al. (2020) study for providing the data of the box modelling study.



References

- Fagerli, H., Svetlana Tsyro, Jan Eiof Jonson, Ágnes Nyíri, Michael Gauss, David Simpson, Peter Wind, Anna Benedictow, Heiko Klein, Augustin Mortier, Wenche Aas, Anne-Gunn Hjellbrekke, Sverre Solberg, Stephen Matthew Platt, Karl Espen Yttri, Kjetil Tørseth, Silke Gaisbauer, Katarina Mareckova, Bradley Matthews, Sabine Schindlbacher, Carlos Sosa, Melanie Tista, Bernhard Ullrich, Robert Wankmüller, Thomas Scheuschner, Robert Bergström (on leave from SMHI), Lasse Johanson, Jukka-Pekka Jalkanen, Swen Metzger, Hugo A.C. Denier van der Gon, Jeroen J.P. Kuenen, Antoon J.H. Visschedijk, and Lars Barregård, Peter Molnár, Leo Stockfelt: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components, EMEP Status Report 1 “, Norwegian Meteorological Institute, https://www.emep.int/publ/common_publications.html, https://emep.int/publ/reports/2019/EMEP_Status_Report_1_2019.pdf, 2019.
- 280 Flemming, J., Huijnen, V., Arteta, J., Bechtold, P., Beljaars, A., Blechschmidt, A.-M., Diamantakis, M., Engelen, R. J., Gaudel, A., Inness, A., Jones, L., Josse, B., Katragkou, E., Marecal, V., Peuch, V.-H., Richter, A., Schultz, M. G., Stein, O., and Tsikerdekis, A.: Tropospheric chemistry in the Integrated Forecasting System of ECMWF, *Geoscientific Model Development*, 8, 975–1003, <https://doi.org/10.5194/gmd-8-975-2015>, 2015.
- 285 Friese, E. and Ebel, A.: Temperature Dependent Thermodynamic Model of the System $\text{H}^+ \text{-NH}_4^+ \text{-Na}^+ \text{-SO}_4^{2-} \text{-NO}_3^- \text{-Cl}^- \text{-H}_2\text{O}$, *The Journal of Physical Chemistry A*, 114, 11 595–11 631, <https://doi.org/10.1021/jp101041j>, 2010.
- 290 Guo, H., Xu, L., Bougiatioti, A., Cerully, K. M., Capps, S. L., Jr., J. R. H., and S.-H. Lee, A. G. C., Bergin, M. H., Ng, N. L., Nenes, A., , and Weber, R. J.: Fine-particle water and pH in the southeastern United States, *Atmospheric Chemistry and Physics*, 15, 5211–5228, <https://doi.org/10.5194/acp-15-5211-2015>, 2015.
- 295 Guo, H., Sullivan, A. P., Campuzano-Jost, P., Schroder, J. C., Lopez-Hilfiker, F. D., Dibb, J. E., Jimenez, J. L., Thornton, J. A., Brown, S. S., Nenes, A., and Weber, R. J.: Fine particle pH and the partitioning of nitric acid during winter in the northeastern United States, *Journal of Geophysical Research: Atmospheres*, 121, 10 355–10 376, <https://doi.org/10.1002/2016JD025311>, 2016.
- 300 Guo, H., Liu, J., Froyd, K. D., Roberts, J. M., Veres, P. R., Hayes, P. L., Jimenez, J. L., Nenes, A., , and Weber, R. J.: Fine particle pH and gas–particle phase partitioning of inorganic species in Pasadena, California, during the 2010 CalNex campaign, *Atmospheric Chemistry and Physics*, 17, 5703–5719, <https://doi.org/10.5194/acp-17-5703-2017>, 2018a.
- 305 Guo, H., Otjes, R., Schlag, P., Kiendler-Scharr, A., Nenes, A., and Weber, R. J.: Effectiveness of ammonia reduction on control of fine particle nitrate, *Atmospheric Chemistry and Physics*, 18, 12 241–12 256, <https://doi.org/10.5194/acp-18-12241-2018>, 2018b.
- Huijnen, V., Le Sager, P., Köhler, M. O., Carver, G., Rémy, S., Flemming, J., Chabrillat, S., Errera, Q., and Van Noije, T.: OpenIFS/AC: atmospheric chemistry and aerosol in OpenIFS 43r3, *Geoscientific Model Development*, 15, 6221–6241, <https://doi.org/10.5194/gmd-15-6221-2022>, 2022.
- 310 Koo, B., Metzger, S., Vennam, P., Emery, C., Wilson, G., and Yarwood, G.: Comparing the ISORROPIA and EQSAM Aerosol Thermodynamic Options in CAMx, in: *Air Pollution Modeling and its Application XXVI*, edited by Mensink, C., Gong, W., and Hakami, A., pp. 93–98, Springer International Publishing, Cham, https://doi.org/10.1007/978-3-030-22055-6_16, series Title: Springer Proceedings in Complexity, 2020.
- Metzger, S.: The EQSAM Box Model (for eqsam4clim-v12), <https://doi.org/10.5281/zenodo.10276178>, 2023.
- 315 Metzger, S., Dentener, F., Pandis, S., and Lelieveld, J.: Gas/aerosol partitioning: 1. A computationally efficient model, *Journal of Geophysical Research: Atmospheres*, 107, <https://doi.org/10.1029/2001JD001102>, 2002.



- Metzger, S., Steil, B., Xu, L., Penner, J. E., and Lelieveld, J.: New representation of water activity based on a single solute specific constant to parameterize the hygroscopic growth of aerosols in atmospheric models, *Atmospheric Chemistry and Physics*, 12, 5429–5446, <https://doi.org/10.5194/acp-12-5429-2012>, 2012.
- 315 Metzger, S., Steil, B., Abdelkader, M., Klingmüller, K., Xu, L., Penner, J. E., Fountoukis, C., Nenes, A., and Lelieveld, J.: Aerosol water parameterisation: a single parameter framework, *Atmospheric Chemistry and Physics*, 16, 7213–7237, <https://doi.org/10.5194/acp-16-7213-2016>, 2016a.
- Metzger, S., Abdelkader, M., Klingmueller, K., Steil, B., and Lelieveld, J.: Comparison of Metop PMAp Version 2 AOD Products using Model Data, <https://www.eumetsat.int/PMAp>, final Report ITT 15/210839, EUMETSAT, 2016b.
- 320 Metzger, S., Abdelkader, M., Steil, B., and Klingmüller, K.: Aerosol water parameterization: long-term evaluation and importance for climate studies, *Atmospheric Chemistry and Physics*, 18, 16 747–16 774, <https://doi.org/10.5194/acp-18-16747-2018>, 2018.
- Metzger, S., Remy, S., Williams, J. E., Huijnen, V., Meziane, M., Kipling, Z., Flemming, J., and Engelen, R.: Representing acidity in the IFS using a coupled IFS-EQSAM4Clim approach, other, display, <https://doi.org/10.5194/egusphere-egu22-11122>, 2022.
- Metzger, S., Rémy, S., Huijnen, V., Williams, J., Chabrilat, S., Bingen, C., and Flemming, J.: Impact of pH computation from EQSAM4Clim on inorganic aerosols in the CAMS system, other, pico, <https://doi.org/10.5194/egusphere-egu23-16085>, 2023.
- 325 Peuch, V.-H., Engelen, R., Rixen, M., Dee, D., Flemming, J., Suttie, M., Ades, M., Agusti-Panareda, A., Ananasso, C., Andersson, E., Armstrong, D., Barre, J., Bousserez, N., Dominguez, J. J., Garrigues, S., Inness, A., Jones, L., Kipling, Z., Letertre-Danczak, J., Parrington, M., Razinger, M., Ribas, R., Vermoote, S., Yang, X., Simmons, A., de Marcilla, J. G., and Thepaut, J.-N.: The Copernicus Atmosphere Monitoring Service: From Research to Operations, *Bulletin of the American Meteorological Society*, 103, E2650 – E2668, <https://doi.org/https://doi.org/10.1175/BAMS-D-21-0314.1>, 2022.
- 330 Pye, H. O. T., Nenes, A., Alexander, B., Ault, A. P., Barth, M. C., Clegg, S. L., Collett Jr, J. L., Fahey, K. M., Hennigan, C. J., Herrmann, H., Kanakidou, M., Kelly, J. T., Ku, I.-T., McNeill, V. F., Riemer, N., Schaefer, T., Shi, G., Tilgner, A., Walker, J. T., Wang, T., Weber, R., Xing, J., Zaveri, R. A., and Zuend, A.: The acidity of atmospheric particles and clouds, *Atmospheric Chemistry and Physics*, 20, 4809–4888, <https://doi.org/10.5194/acp-20-4809-2020>, 2020.
- 335 Rémy, S., Kipling, Z., Huijnen, V., Flemming, J., Nabat, P., Michou, M., Ades, M., Engelen, R., and Peuch, V.-H.: Description and evaluation of the tropospheric aerosol scheme in the Integrated Forecasting System (IFS-AER, cycle 47R1) of ECMWF, *Geoscientific Model Development*, 15, 4881–4912, <https://doi.org/10.5194/gmd-15-4881-2022>, 2022.
- Shi, X., Nenes, A., Xiao, Z., Song, S., Yu, H., Shi, G., Zhao, Q., Chen, K., Feng, Y., and Russell, A. G.: High-Resolution Data Sets Unravel the Effects of Sources and Meteorological Conditions on Nitrate and Its Gas-Particle Partitioning, *Environmental Science and Technology*, 53, 3048–3057, <https://doi.org/10.1021/acs.est.8b06524>, 2019.
- 340 Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- , and H_2O , *Journal of Geophysical Research: Atmospheres*, 107, <https://doi.org/10.1029/2001JD000451>, 2002.
- Williams, J. E., Huijnen, V., Bouarar, I., Meziane, M., Schreurs, T., Pelletier, S., Marécal, V., Josse, B., and Flemming, J.: Regional evaluation of the performance of the global CAMS chemical modeling system over the United States (IFS cycle 47r1), *Geoscientific Model Development*, 15, 4657–4687, <https://doi.org/10.5194/gmd-15-4657-2022>, 2022.
- 345