

1 **What is the stability of additional organic carbon stored thanks to alternative** 2 **cropping systems and organic wastes products application? A multi-methods** 3 **evaluation**

4
5 Tchodjowiè P. I. Kpemoua^{1,2,3}, Pierre Barré², Sabine Houot¹, François Baudin⁴, Cédric
6 Plessis¹, Claire Chenu¹

7 ¹ UMR Ecosys, Université Paris-Saclay, INRAE, AgroParisTech, Palaiseau, 91120,
8 France

9 ² Laboratoire de Géologie, UMR 8538, Ecole Normale Supérieure, PSL Research University, CNRS,
10 Paris 75005, France

11 ³ Agence de la transition écologique, ADEME, 49004 Angers, France

12 ⁴ UMR IStEP 7193, Sorbonne Université, CNRS, France

13 **Corresponding author:**

14 E-mail address: claire.chenu@inrae.fr (C. Chenu)

16 **Abstract**

17 The implementation of agroecological practices often leads to additional soil organic carbon storage,
18 and we have sought to assess the biogeochemical stability of this additional carbon. To achieve this, we
19 implemented a multi-method approach using particle size and density fractionation, Rock-Eval®
20 thermal analyses and long-term incubation (484 days), that we applied to topsoil samples (0-30 cm)
21 from temperate luvisols that had been subjected, in > 20 years long-term experiments in France, to
22 conservation agriculture (CA), organic agriculture (ORG) and conventional agriculture (CON-LC) in
23 La Cage experiment, and to organic wastes products (OWPs) applications in QualiAgro experiment,
24 including biowaste composts (BIOW), residual municipal solid waste composts (MSW), farmyard
25 manure (FYM) and conventional agriculture without organic inputs (CON-QA). The additional carbon
26 resulting from agroecological practices is the difference between the carbon stock of the bulk soil,
27 physical fractions or carbon pools in a soil under the agroecological practices and that of the same soil
28 under a conventional practice as control. The incubations provided information on the additional carbon
29 stability in the short term (i.e., MRT <2 years) and showed that the additional soil organic carbon
30 mineralized faster than the carbon in the conventional control at La Cage but slower at QualiAgro. In
31 OWPs-treated plots at QualiAgro, 60-66% of the additional carbon was stored as mineral-associated
32 organic matter (MAOM-C), and 34-40% as particulate organic matter (POM-C). In CA and ORG
33 systems at La Cage, 77-84% of the additional carbon was stored in MAOM-C, versus 16-23% as POM-
34 C. Management practices hence influenced the distribution of additional carbon in physical fractions.
35 Utilizing the PARTYSOC model with Rock-Eval® thermal analysis parameters, we found that most, if
36 not all, of the additional carbon belonged to the active carbon pool (MRT ~ 30-40 years). In summary,

37 our comprehensive multi-methods evaluation indicates that the additional soil organic carbon is less
38 stable over decadal and pluri-decadal time-scales compared to soil carbon under conventional controls.
39 Our results show that particle size and density fractions can be heterogenous in their biogeochemical
40 stability. On the other hand, although the additional carbon is mainly associated with MAOM, the high
41 proportion of this carbon in the active pool suggests that it has a mean residence time not exceeding ~50
42 years. On the other hand, agroecological practices with equivalent additional carbon stocks (MSW,
43 FYM vs CA) exhibited a higher proportion of additional carbon in POM-C under MSW (40%) and FYM
44 (34%) compared to CA (16%), which suggests a high chemical recalcitrance of POM-C under OWPs
45 management relative to conservation agriculture. Additional soil organic carbon deriving from organic
46 wastes, i.e., biomass that has been partially decomposed and transformed through its processing prior to
47 its incorporation in soil, would be more biogeochemically stable in soil than that deriving directly from
48 plant biomass. The apparent contradictions observed between method can be explained by the fact that
49 they address different kinetic pools of organic carbon. Care must be taken to specify which range of
50 residence times is considered when using any method intending to evaluate the biogeochemical stability
51 of soil organic matter, as well as when using the terms stable or labile. In conclusion, the contrasting
52 biogeochemical stabilities observed in the different management options highlight the need to maintain
53 agroecological practices to keep these carbon stocks at a high level over time, given that the additional
54 carbon is stable on a pluri-decadal scale.

55
56 *Keywords:* soil organic carbon, additional carbon, agroecological practices, Rock-Eval®,
57 biogeochemical stability, incubation, physical fractions.

58

59 **1 Introduction**

60 Soil organic matter (SOM) plays a crucial role in the functioning of terrestrial ecosystems and
61 can contribute to mitigate climate change. A minor change of soil organic carbon (SOC) content can
62 make a significant difference to global climate because soil contains more carbon than vegetation and
63 atmosphere combined (Lal, 2004). The 4p1000 initiative encourages the implementation of agricultural
64 practices that increase and/or maintain soil carbon stocks (www.4p1000.org, Rumpel et al., 2020). At
65 the field scale, changes in SOC stocks result from an imbalance between C inputs (crop residues, litter,
66 root exudates, exogenous organic matter such as organic wastes products (OWPs)) and C outputs from
67 the system due to crop residue export, SOC mineralization, leaching, or erosion (Lal, 2018). Although
68 some agricultural practices can reduce mineralization rates (e.g., reduced tillage, see review by
69 Haddaway et al., 2017), it is generally accepted that the most effective way to increase SOC stocks is to
70 increase carbon inputs (e.g., Virto et al., 2012; Autret et al., 2016; Fujisaki et al., 2018; Chenu et al.
71 2019). This can be achieved by increasing biomass production in the field and residue return (e.g., cover

72 crops, Poeplau and Don, 2015; Autret et al., 2016), or by mobilizing external carbon resources such as
73 OWPs (Peltre et al., 2012; Paetsch et al., 2016).

74

75 The implementation of selected agroecological practices and systems such as conservation
76 agriculture, agroforestry, OWPs application allows for additional carbon storage in soils (Peltre et al.,
77 2012; Autret et al., 2016; Paetsch et al., 2016; Pellerin et al., 2019; Bohoussou et al., 2022). The
78 additional carbon storage linked to agricultural practice B is the difference between the carbon stock in
79 a soil under practice B and that of the same soil under a reference practice (Pellerin et al., 2019). This
80 additional carbon storage is not necessarily the result of recent carbon inputs, but can also include the
81 legacy carbon. However, knowledge on the biogeochemical stability of this additional carbon is lacking,
82 questioning the reversibility of this storage. The carbon sink effect will indeed be more effective if the
83 additional carbon storage is realized in the form of persistent organic carbon (OC) and not in the form
84 of labile OC. We propose to evaluate and compare the biogeochemical stability of additional organic
85 carbon following implementation of various agroecological practices.

86

87 Several methods have been reported in the literature to assess the organic carbon temporal
88 stability in soils. These methods isolate kinetic pools or carbon fractions with contrasting mean residence
89 times (MRT), e.g., particle size fractionation, Balesdent, 1996; density fractionation, Sollins et al., 2006;
90 sequential extraction, Heckman et al., 2018; thermal analysis, Barré et al., 2016 and incubation, Schädel
91 et al., 2020. Physical fractionation is probably the most used method so far to evaluate SOM stability.
92 Physical fractionation methods isolate fractions based on size, density, or a mixture of both (Chenu et
93 al., 2015). In a study comparing several fractionation methods, Poeplau et al. (2018) found that particle
94 size fractionation was well suited to isolate particulate organic matter (POM) fractions from mineral
95 associated organic matter (MAOM) with contrasting MRT. Fractionation of SOM into POM and
96 MAOM components can reveal insights about the sources and stability of SOC (Kim et al., 2022).
97 However, some studies have shown that SOC fractionation methods fail to accurately separate stable
98 SOC from active SOC, and in particular that the isolated MAOM fractions are mixtures of labile SOC
99 (MRT of months to year) and stable centennial SOC (Balesdent et al. 1987; Jastrow et al., 1996;
100 Sanderman et al., 2013; Torn et al., 2013; Balesdent, 1996; Hsieh, 1992; von Lützow et al., 2007;
101 Sanderman and Grandy, 2020). This may be due to methodological challenges as much as the fact that
102 there are multiple pathways for SOM formation and stabilization (Cotrufo et al., 2013; Sokol et al.,
103 2019).

104

105 Thermal analysis techniques, long used in petroleum exploration and clay mineralogy, offer a
106 promising alternative or complement to physical and chemical fractionation methods, and are
107 increasingly applied to studies of SOC stability (Peltre et al., 2013; Plante et al., 2009). Indeed, several
108 parameters obtained using thermal analysis are strongly related to SOM biogeochemical stability (Barré

109 et al., 2016; Poeplau et al., 2019). However, these parameters do not allow us to separate the kinetic
110 carbon pools (Schiedung et al., 2017). And so, recently, Cécillon et al. (2018, 2021) developed a
111 machine-learning model, called PARTY_{SOC}, showing that Rock-Eval® parameters can be used to
112 predict the fraction of SOC that is stable at a centennial timescale. Kanari et al. (2022) evidenced that
113 SOC fractions calculated using PARTY_{SOC} matched the stable (MRT >100 years) and active (MRT ~
114 30-40 years) OC pools of the AMG model, a model widely validated to simulate SOC stock evolution
115 in French and European croplands (Clivot et al., 2019; Bruni et al., 2022). As a result, one can consider
116 that a Rock-Eval® analysis associated to the PARTY_{SOC} model allows for the quantification of carbon
117 fractions that are stable at a centennial timescale and active *sensu* AMG model.

118

119 The incubation method is, however, the only direct test for the biological stability of SOC, that
120 results from chemical resistance to decomposition and/or organo-mineral associations and/or
121 inaccessibility of organic substrates to microbial decomposition. Long-term incubations (months to
122 years) may diverge from the conditions prevailing in the soil profile but provide insights into the
123 potential decomposability of slower-cycling SOC (e.g., Schädel et al., 2014). In early laboratory
124 incubations, fast-cycling C respiration dominates total respired SOC, but rapidly declines, while slow-
125 cycling SOC accounts for most of the respired SOC after the fast SOC pool is depleted.

126

127 These different methods do not separate similar carbon kinetic pools. Indeed, the incubation
128 method isolates carbon with MRT ranging from days to years (Schädel et al., 2014) while others isolate
129 carbon with longer MRT (decades to centuries) (Cécillon et al., 2018, 2021, Balesdent, 1996). Thus, a
130 multi-method approach will further improve our knowledge of the biogeochemical stability of SOC in
131 the short (<2 years), medium (2-50 years) or long term (>50 years). The objective of this study was to
132 evaluate the biogeochemical stability of additional SOC stored upon the implementation of carbon
133 storing agroecological practices using a multi-method approach. To do so, we characterized SOM using
134 particle size and density fractionation, Rock-Eval® (RE) thermal analysis and incubation in soil from
135 plots managed using various agroecological practices such as addition of OWPs (composts and farmyard
136 manure) and alternative cropping systems including no tillage, permanent cover crop and the
137 introduction of legumes in the rotation. The application of OWPs is likely to provide organic matter
138 (OM) that has been pre-stabilized by the storage (manure) or composting process and is hence less
139 decomposable than the fresh matter provided by plant biomass in alternative cropping systems. Then,
140 we hypothesized (i) that the biogeochemical stability of additional SOC depends on the management
141 practices implemented and (ii) that the additional SOC originating from OWPs would be more stable
142 than that directly originating from plant biomass, but, (iii) overall, that the additional SOC would be less
143 stable than the SOC stored in the conventional controls.

144 2 Materials and methods

145 2.1 Field Site and soil sampling

146 This study focuses on two French long-term experiments (LTEs) developed on Luvisols in the
147 same region, where agroecological practices including conservation agriculture, organic agriculture and
148 OWPs application (composts and manure) were implemented.

149 **La Cage experiment** is conducted in Versailles (48°48'N, 2°08'E, alt 120 m). During the 21 years
150 of experimentation, the mean annual temperature and precipitation were 11.6 °C and 633 respectively.
151 The soil is a well-drained deep Luvisol (IUSS Working Group WRB, 2015). The field experiment is
152 arranged in a randomized complete block design, divided into two blocks, themselves divided into four
153 plots for each cropping system. Each plot is divided into two subplots of 0.56 ha, so that two different
154 crops of the crop rotation are present each year, wheat being grown every year in one of the two subplots
155 (Autret et al., 2020). A detailed presentation of crop rotations, soil management and fertilization were
156 given by Autret et al. (2016). The 4 year's crop rotation mainly consisted of rapeseed (*Brassica napus*
157 *L.*), winter wheat (*Triticum aestivum L.*), spring pea (*Pisum sativum L.*) and winter wheat.



- 158 - Conventional agriculture (CON-LC) is characterized by a soil and crop management
159 representative of the Paris Basin cereal production, with annual soil ploughing, the absence of
160 organic amendment, a mineral N fertilization (average rate = 143 kg N ha⁻¹ yr⁻¹) and a
161 systematic use of pesticides.
- 162 - Conservation agriculture (CA) includes a permanent soil cover, initially fescue (*Festuca rubra*)
163 and since 2008 alfalfa, grown under the main crops, except pea. In the rotation, rapeseed is
164 replaced by maize (*Zea mays L.*) in CA and direct seeding is performed.
- 165 - Organic agriculture (ORG) is characterized by alfalfa-alfalfa-wheat-wheat rotation with annual
166 soil ploughing and no synthetic fertilizers nor pesticides.

167 **The QualiAgro experiment** is located at Feucherolles, 20 km west of Versailles (48°52'N, 1°57'E,
168 alt 150 m). During the 21 years of experimentation, the mean annual temperature and precipitation were
169 11.0 °C and 614 respectively. The soil is a Luvisol (IUSS Working Group WRB, 2015). The crop
170 rotation mainly consisted of wheat and maize (Peltre et al., 2012). It is a field experiment conducted in
171 collaboration with INRAE and Veolia Environment Research and Innovation since 1998, on which
172 composts of OWPs are applied every 2 years for a dose equivalent to ~4 t C. ha⁻¹ from 1998 to 2013 and
173 ~2 t C. ha⁻¹ from 2015 to 2020. The unit plots are 10 x 45 m². Each treatment has 4 replicates and OWPs
174 are applied every two years on wheat stubble. Soils are ploughed every year on this experimental site.
175 Since 2015, wheat and maize residues are buried in the soil. Four treatments are considered in this study.

- 176 – Conventional agriculture (CON-QA): is characterized by a soil and crop management
 177 representative of the Paris Basin cereal, the absence of organic amendment, a mineral N
 178 fertilization (average rate = 167 kg N ha⁻¹ yr⁻¹).
- 179 – Biowaste compost (BIOW): composting of the fermentable fraction of selectively collected
 180 household waste, mixed with green waste;
- 181 – Municipal solid waste compost (MSW): composting of the residual fraction of household waste
 182 after selective collection of packaging;
- 183 – Farmyard manure (FYM) which represents a reference amendment.

184 At both sites, four replicate plots were available per treatment. From each plot, 3 sub-samples were
 185 taken from the topsoil 30 ± 1 cm (in September 2019 at QualiAgro and in November 2020 at La Cage),
 186 thoroughly mixed and combined into one sample. The samples were sieved to 4 mm, homogenized, the
 187 plant material was removed and the soil was oven dried at 35°C for 72h before particle size and density
 188 fractionation and RE thermal analysis.

189
 190 **Table 1** Soil organic carbon (SOC), soil organic nitrogen (SON) and C/N measured in topsoil. Values
 191 in brackets are standard deviations. CON-QA: conventional agriculture without organic inputs, MSW:
 192 municipal solid waste compost, FYM: farmyard manure, BIOW: biowaste compost, CON-LC:
 193 conventional agriculture, ORG: organic agriculture and CA: conservation agriculture.

Site	Soil textured	Agricultural Practices	SOC content g.kg ⁻¹	SOC stocks t C. ha ⁻¹	SOC gain (%)	SON g.kg ⁻¹	C/N
La Cage 	Luvisol 17% Clay 58% Silt 25% Sand	CON-LC	9.82 ± 0.48	42.22 ± 2.08	-	1.01 ± 0.07	10.58 ± 1.58
		ORG	10.39 ± 0.42	44.66 ± 1.80	6	1.09 ± 0.03	9.52 ± 0.12
		CA	13.30 ± 1.05	57.17 ± 4.53	35	1.29 ± 0.10	10.29 ± 0.28
QualiAgro 	Luvisol 15% Clay 78% Silt 7% Sand	CON-QA	9.92 ± 0.63	39.31 ± 2.49	-	0.97 ± 0.08	10.35 ± 1.61
		MSW	13.84 ± 0.16	54.03 ± 0.59	33	1.35 ± 0.04	10.26 ± 0.42
		FYM	13.91 ± 0.37	54.77 ± 1.40	42	1.36 ± 0.02	10.21 ± 0.37
		BIOW	16.04 ± 0.68	63.17 ± 2.56	64	1.62 ± 0.01	9.87 ± 0.41

2.2 Calculation of SOC stocks and additional carbon stocks

SOC stocks were calculated at equivalent soil mass in both long-term experiments. Thus, at QualiAgro the SOC stock was calculated by multiplying the SOC content by bulk density (data provided by QualiAgro) and was normalized to a depth of 10 cm (factor 10^{-3}) (reference soil mass of 3963 kg. ha^{-1}). Bulk densities between 1998 and 2019 increased significantly in all plots. We calculated the additional soil thickness required to achieve this equivalent soil mass in treatments with lighter tilled layers as described by Ellert and Bettany (1995):

$$T_{\text{add}} = \frac{(M_{\text{soil equiv}} - M_{\text{soil topsoil}}) * 10^{-4}}{\rho_{\text{b subsoil}}}, \quad (1)$$

where T_{add} is the additional thickness of the sub-soil layer expressed in cm needed to reach the equivalent soil mass, $M_{\text{soil equiv}}$ is the equivalent soil mass of the denser horizon in kg. ha^{-1} . In our study, the dense 0-29 cm layer was the reference treatment in 2019 with a bulk density of 1.37 g.cm^{-3} giving an equivalent soil mass ($M_{\text{soil equiv}}$) of 3963 kg. ha^{-1} . $M_{\text{soil topsoil}}$ is the soil mass in the surface (tilled) layer and $\rho_{\text{b subsoil}}$ is the bulk density of the underlying 29-35 cm layer (in g.cm^{-3}). Carbon stocks per hectare in equivalent soil masses ($\text{Stock}_{\text{C equiv}}$) were calculated by adding the carbon stock in the surface layers ($\text{Stock}_{\text{C topsoil}}$) and in the additional underlying layers ($\text{Stock}_{\text{C, Tadd}}$) with the following formula:

$$\text{Stock}_{\text{C equiv}} = \text{Stock}_{\text{C topsoil}} + \text{Stock}_{\text{C Tadd}}, \quad (2)$$

At La Cage experiment, the soil sampling strategy was designed to calculate SOC stocks on an equivalent soil mass (ESM) basis Ellert and Bettany (1995) over a depth at least equal to the deepest tillage event. The ploughing depth was *ca.* 30 cm before 1998 and shallower afterwards, about 25 cm (Autret et al., 2016). The sample was taken at the depth equivalent to a soil mass of 4300 kg. ha^{-1} . The carbon stocks were calculated by multiplying the SOC contents with this equivalent soil mass.

We then calculated the additional carbon storage ($\Delta\text{SOC stock}$) considering each time the conventional control at La Cage (CON-LC) and the conventional control at QualiAgro (CON-QA). The following formula was used:

$$\Delta\text{SOC stock} = \text{Stock}_{\text{C Practice}} - \text{Stock}_{\text{C conventional}}, \quad (3)$$


With $\text{Stock}_{\text{C Practice}}$: the carbon stock of the agroecological practice and $\text{Stock}_{\text{C conventional}}$: the carbon stock of the conventional control. The standard deviation used for the additional carbon stock was calculated based on the equation described by Kuzyakov and Bol, (2004) as follows:

$$SD_{\Delta\text{SOC stock}} = \sqrt{(SD_{\text{stock C Practice}})^2 + (SD_{\text{stock C conventional}})^2}, \quad (4)$$

2.3 Particle size and density fractionation

The method uses a preliminary disaggregation aiming at the best compromise between maximum destruction of micro-aggregates of size $< 50 \mu\text{m}$, and respect of the integrity of organic debris (Balesdent et al., 1991) and combines fractionation by particle size to separate POM from OM associated with clays and silts minerals with water flotation to separate POM from sands. For this purpose, approximately 50 g of soil was suspended in 180 mL of 0.5% sodium hexametaphosphate (SHMP) saline solution in a 250

230 mL polyethylene bottle; 10 glass beads were added and the whole set to agitation by inversion (REAX
 231 2 type inversion mixers) for 16 hours, at a speed of approximately 50 rpm to destroy the aggregates. The
 232 SHMP solution and the glass beads allows to completely disperse soil aggregates > 50 μm diameter in
 233 these soils (Balesdent et al., 1991). After agitation, the suspension was first sieved on a 200 μm sieve
 234 from which the refusal, the coarse fraction, was recovered in a 250mL glass beaker. We separated the
 235 coarse POM (cPOM) from the coarse sands (cSand) by flotation in water. The suspension <200 μm in
 236 a second time was submitted to a second sieving at 50 μm and the same operations were performed to
 237 separate the fine POM (fPOM) from the fine sands (fSand) using water flotation. The suspension <50
 238 μm is submitted to ultrasounds by imposing an energy of 300 J.mL^{-1} necessary to disperse the micro-
 239 aggregates (Balesdent et al., 1998). After this step, we sieved the suspension <50 μm to 20 μm to recover
 240 the coarse silts of size between 20 and 50 μm (cSilt) remaining on the sieve. The suspensions containing
 241 particles <20 μm were pooled in a 2L beaker. The separation of the fine silts between 2 and 20 μm (fSilt)
 242 from the clays was performed by centrifugation of the <20 μm filtrate at 64 g (circa 500 rpm) for 10min.
 243 The supernatant containing the clays was collected in a 5L beaker. The same process was repeated 4 to
 244 5 times by resuspending the pellet for an optimal recovery of fine silts by decantation. The supernatant
 245 collected in the 5L beaker constitutes the clay fraction (<2 μm) and the pellet after repeated
 246 centrifugation constitutes the fine silt fraction. To reduce the volume of the clay suspension to be freeze-
 247 dried, we added CaCl_2 to flocculate the clay particles and by centrifugation for 20 min at 16000 g (circa
 248 8000 rpm) we recovered the pellet which constitutes the clay fraction. An aliquot of the supernatant was
 249 taken to determine the dissolved organic carbon (DOC). The particle-size and density fractionation
 250 resulted in recovery rates of $95 \pm 2\%$ of the initial sample mass and $98 \pm 6\%$ recovery of carbon.



251

252 **Fig. 1** Particles size and density fractionation protocol (adapted from Balesdent et al., 1998).
253 The POM fraction is the sum of the cPOM, fPOM, cSand and fSand fractions, while the
254 MAOM fraction is the sum of the cSilt, fSilt and Clay fractions.

255 **2.3.1 Fractions preparation and elemental analysis (C, N)**

256 The fractions obtained during this fractionation were dried or freeze-dried. The organic (cPOM,
257 fPOM) and mineral (cSand, fSand, cSilt) fractions were oven dried at 50°C for 3 days, while the fine
258 silt (fSilt) and clay (Clay) fractions were freeze dried. Each fraction was weighed and C, N were
259 determined using dry combustion (elemental analyzer, Elemantar Vario ISOTOP).

260 **2.4 Rock-Eval® (RE) thermal analysis**

261 We analyzed 28 samples of bulk soil, using a RE6 Turbo apparatus (Vinci Technologies). A small
262 amount of soil (about 60 mg) was required for the analysis, which was performed in two consecutive
263 steps, during which carbon-containing effluents were directly detected. First, the sample underwent
264 pyrolysis in an inert atmosphere (N₂), followed by oxidation in the presence of O₂ (ambient air). The
265 heating routine applied during pyrolysis was that proposed by Disnar et al. (2003) and Baudin et al.
266 (2015), including a three-minute isotherm at 200 °C, followed by a 30 °C·min⁻¹ heating ramp to 650 °C.
267 Oxidation began with a one-minute isotherm at 300 °C, followed by a 20 °C·min⁻¹ heating ramp to 850
268 °C and a final five-minute isotherm at 850 °C (oxidation routine presented in Baudin et al. (2015) as the
269 "bulk rock/basic" method). Simultaneous detection of effluents during both analytical steps generated a
270 total of five thermograms per sample describing the evolution of hydrocarbons during pyrolysis
271 (HC_PYR), and CO and CO₂ during both pyrolysis and oxidation steps (CO_PYR, CO₂_PYR, CO_OX,
272 CO₂_OX).
273
274

275 **2.4.1 Rock-Eval® parameters**

276 The classical Rock-Eval® parameters were acquired using the RockSix software (Vinci
277 Technologies) with a good reproducibility (Pacini et al., 2023). They include: six automatically
278 generated "peaks" defined as specific areas of the three pyrolysis thermograms (S1, S2, S3, S3', S3CO
279 and S3'CO; Lafargue et al. (2018)), the amount of pyrolyzed carbon (PC corresponding to the sum of
280 organic C released as HC, CO, and CO₂ during pyrolysis), total organic carbon (TOC corresponding to
281 the amount of organic C released during analysis), inorganic carbon (MinC corresponding to the amount
282 of C released from carbonate cracking), hydrogen index (HI corresponding to the ratio of hydrocarbons
283 released to TOC), and oxygen index (OI_{RE6} corresponding to the ratio of organic oxygen released to
284 TOC). In addition, other parameters used as predictors by the PARTYSOCv2.0EU model were

285 calculated based on thermograms obtained using R scripts available on Zenodo
286 (<https://zenodo.org/record/4446138#.YDe84Xlw2SQ>) (Cécillon et al., 2021, Kanari et al., 2021). These
287 include: PseudoS1 (the sum of carbon released during the first 200 s of isothermal 200°C pyrolysis as
288 HC, CO, and CO₂), the S2/PC ratio (the ratio of the amount of hydrocarbons released excluding the first
289 200 s of pyrolysis to the pyrolyzed carbon), the PC/TOC ratio, the HI/OIRE6 ratio, and ten temperature
290 parameters (e.g., T30, T50, T70, T90) that describe the evolutionary steps, i.e., at what temperature 30,
291 50, 70, and 90% of a given gas was released. A detailed description of the definition, units, and equations
292 used to calculate all parameters can be found in the study of Kanari et al. (2021). The HI and OIRE6 are
293 commonly reported indices that represent proxies of the SOM H/C and O/C ratios respectively.
294

295 **2.4.2 PARTYSOC model based on Rock-Eval® (RE)**

296 In this study, we used the random forest model based on RE results PARTY_{SOCv2.0EU}
297 (<https://zenodo.org/record/4446138#.YDe84Xlw2SQ>) proposed by Cécillon et al. (2021). This model
298 was calibrated on data from 6 long-term agricultural experiments including a bare fallow treatment in
299 northwestern Europe and can predict the proportion of persistent SOC at a centennial timescale in topsoil
300 samples (0-30 cm). The model requires a set of 18 RE parameters (e.g., Kanari et al., 2021) characteristic
301 of a sample and provides a prediction of the proportion of stable SOC for soils from the La Cage and
302 the QualiAgro long-term experiments. The 18 RE parameters retained were the RE temperature
303 parameters T70HC_PYR, T90HC_PYR, T30CO₂_PYR, T50CO₂_PYR, T70CO₂_PYR, T90CO₂_PYR,
304 T70CO_ox, T50CO₂_ox, T70CO₂_ox, and T90CO₂_ox and the RE parameters PseudoS1, S2, S2 /
305 PC, HI, HI / OI_{RE6}, PC, PC / TOC_{RE6}, and TOC_{RE6} (Cécillon et al., 2021).

306 **2.5 Long-term incubation**

307 Polyvinyl chloride (PVC) cylinders 5.7 cm in diameter and 4 cm in height with 2 mm perforations
308 were used to build soil microcosms. A 50 µm mesh fabric at the bottom of the cylinder supported the
309 soil while promoting gas exchange. Each cylinder was weighed empty and then with fresh soil
310 equivalent to 100 g of dry soil. The soil samples were then brought to a bulk density of 1.3 g.cm⁻³ with
311 a hand press and mold. Knowing the initial water content, the samples were gradually brought to pF 2.5
312 by adding water with a Pasteur pipette. Then, the microcosms were mounted in 0.5 L jars. The soil
313 cylinders were placed on PVC racks and 15 mL of water was added to the bottom of the jars to stabilize
314 the humidity. The jars were sealed and the whole set was placed in the incubator at 20°C for one week
315 pre-incubation. Four replicates per agricultural practice were prepared.
316

317 After the pre-incubation period, we readjusted the water content of the soil cylinders to pF 2.5
318 when necessary. A total of 28 soil cylinders were incubated for 484 days under the same temperature
319 (20°C) and moisture (pF2.5) conditions.

320 ▪ **Mineralization measurement**

321 Soil organic carbon (SOC) mineralization in samples from both long-term experiments (LTEs)
322 was measured nondestructively using a micro gas chromatograph (μ GC 490; Agilent Technologie;
323 USA). Measurements were performed 1, 3, 7, 14, 28, 35 days, then one measurement every 2 weeks
324 until the sixth month and finally, one measurement every month until the end of incubation. The CO₂
325 emitted is measured in parts per million (ppm). It is then converted to $\mu\text{g C-CO}_2 \text{ g}^{-1}$ of dry soil using the
326 following formula: $\mu\text{g C - CO}_2 \cdot \text{g}^{-1} \text{ dry soil} = \frac{\text{CO}_2(\text{ppm}) * M_c * V_b}{V_M * M_{\text{soil}}}$ (5), (Védère et al., 2020).

327 With CO₂ (ppm): amount of CO₂ emitted measured by gas phase microchromatograph; M_c: molar mass
328 of carbon in $\text{g}\cdot\text{mol}^{-1}$; V_b: volume of the jar in L; V_M: molar volume of the gas in $\text{L}\cdot\text{mol}^{-1}$ and M_{soil}: mass
329 of the incubated dry soil in g. The absolute amount of carbon mineralized was expressed per unit of
330 SOC to obtain the specific SOC mineralization in $\mu\text{g C-CO}_2/100 \mu\text{g SOC}$, i.e., % SOC mineralized
331 (Kpemoua et al., 2023). To calculate the amount of additional carbon mineralized over the 484 days, we
332 first calculated the difference in absolute carbon mineralization between the agroecological practice and
333 the conventional control. We assume that the extra absolute carbon mineralized in the agroecological
334 practice relative to the conventional control comes from the additional carbon. Given the amount of
335 additional carbon (ΔSOC), we then expressed this extra absolute carbon mineralization in terms of
336 additional carbon (% ΔSOC).

337 **2.6 statistical analysis**

338 All data were tested for normality and homogeneity of variance. Log-transformation was applied
339 to the data for the cSand and fSand fractions, as the transformation improved the normality and variance
340 substantially. A one-way ANOVA was used to detect significant differences at the 5% threshold in bulk
341 soil carbon stocks, fractions, carbon pools and amount of carbon mineralized (C_{min}). Once a significant
342 difference was detected, Tukey's multiple comparison test was used to compare carbon stocks, additional
343 carbon stocks, percentage of total carbon storage and percentage of additional carbon storage in either
344 bulk soil, fractions and carbon pools according to agricultural practices. All statistical analyses were
345 completed in R (version 4.0.2).

346

347 **3 Results**

348 **3.1 SOC stocks**

349 The application of organic wastes products (OWPs), increased soil organic carbon (SOC) contents
350 in soils by 64% in biowaste composts treatment (BIOW), 40% in farmyard manure treatment (FYM)
351 and 39% in residual solid waste compost treatment (MSW) compared to the conventional control (CON-

352 QA); while, at La Cage, the implementation of organic agriculture (ORG) and conservation agriculture
353 (CA) increased SOC contents by 6% and 35% respectively, relative to conventional control (CON-LC)
354 (Table 1). The OWPs application significantly increased carbon stocks at QualiAgro. The SOC stocks
355 were in the order: BLOW > FYM \geq MSW > CON-QA (Table. 1). At La Cage, SOC stocks were in this
356 order: CA > ORG \geq CON-LC (Table 1).


357

358 **3.2 SOC distribution in fractions**

359 The mass proportion, carbon content and % carbon distribution of the physical fractions after
360 particle size and density fractionation are presented in the supplementary data Table. S1 and Table. S2.
361 The distribution of SOC stocks over the fractions obtained, expressed in t C. ha⁻¹, is given in Figs. 2a
362 and 2b. Carbon distribution in conventional controls (CON-LC and CON-QA) showed that 19-22% of
363 carbon was found in POM fractions, versus 78-81% in MAOM fractions (Figs. 4a and 4b). Overall, most
364 of the organic carbon was located in the clay fraction (64 -72% SOC, see Table. S1 and S2) regardless
365 of site and the agricultural practice implemented. The carbon distribution in QualiAgro indicated a
366 significant increase of SOC stocks in the cSand, fPOM, cSilt and Clay fractions after OWPs application
367 ($p < 0.05$), while no significant difference was observed in the cPOM, fSand and fSilt fractions ($p > 0.05$).
368 In La Cage, the implementation of conservation agriculture significantly increased SOC stocks as fPOM
369 and Clay fractions compared to organic and conventional agriculture which remained statistically equal.

370

371 We calculated the distribution of additional carbon (Δ SOC) in the fractions by difference of the
372 carbon stock of the bulk soil or physical fractions under agroecological practices with the carbon stock
373 of the bulk soil or physical fractions under conventional control (CON-QA or CON-LC respectively for
374 the QualiAgro and the La Cage experiments). The additional carbon stock at QualiAgro was $23.86 \pm$
375 1.79 t C. ha⁻¹ in BLOW compared to 15.46 ± 1.43 t C. ha⁻¹ in FYM and 14.72 ± 1.28 t C. ha⁻¹ in MSW
376 (Fig. 2c). At La Cage, the additional SOC stock was 14.95 ± 2.49 t C. ha⁻¹ in CA compared to $2.44 \pm$
377 1.38 t C. ha⁻¹ in ORG (Fig. 2d). In terms of percentage, we observed that the coarse mineral fractions
378 (cSand and fsand) have a negligible proportion of additional carbon at La Cage, representing 1% in CA
379 and 0% in ORG, while this proportion was raised to 2% in FYM, 5% in MSW and 7% in BLOW at
380 QualiAgro. This non-negligible proportion of carbon in the sand at QualiAgro suggests that not all
381 particulate organic matter (POM-C) has probably been isolated from the sand. Hence, in the following,
382 to define the POM-C fraction class, we combine the fractions of cPOM, fPOM, cSand and fSand
383 together. Thus, we observed in the QualiAgro experiment that, 60-66% of the additional carbon was
384 localized in mineral-associated organic matter fractions (MAOM-C), which included the cSilt, fSilt, and
385 Clay fractions, versus 34-40% in POM-C; whereas, in La Cage experiment, 77-84% of the additional
386 SOC stock was located in the MAOM-C versus 16-23% in the POM-C. Furthermore, among practices
387 with equivalent additional carbon stocks (MSW, FYM, CA), OWPs application resulted in a higher
388 proportion of additional carbon in POM-C (MSW: 34%; FYM: 40%) compared to CA (16%).



390


391 **Fig. 2** Soil organic carbon stock and additional carbon (Δ SOC) stock of bulk soils and physical fractions
 392 ($n = 4$) at QualiAgro and La Cage experiments. The error bars represent the standard deviations. Grouped
 393 bars with different letters are significantly different between agricultural practices (Tukey-HSD, $p <$
 394 0.05). CON-QA: conventional agriculture without organic inputs, BIOW: biowaste compost, MSW:
 395 municipal solid waste compost, FYM: farmyard manure, CON-LC: conventional agriculture, CA:
 396 conservation agriculture and ORG: organic agriculture. The POM fraction is the sum of the cPOM,
 397 fPOM, cSand and fSand fractions, while the MAOM fraction is the sum of the cSilt, fSilt and
 398 Clay fractions.

399

400 3.3 Estimating stable and active SOC pools with the PARTY_{SOC} model

401 The PARTY_{SOC} machine learning model was used to estimate the proportion of stable SOC under
 402 the different managements. The distribution of organic carbon stocks in the active and stable pools are
 403 shown in Fig. 3. In conventional controls, 38-43% of the soil carbon is found in the active pool, versus
 404 57-62% in the stable pool. The organic wastes products (OWPs) application significantly increased the
 405 size of the active pool relative to conventional control (Fig. 3a, ANOVA, $p < 0.05$). It was of $31.87 \pm$
 406 $2.23 \text{ t C. ha}^{-1}$ in BIOW compared to $29.62 \pm 1.97 \text{ t C. ha}^{-1}$ in FYM, $26.97 \pm 1.07 \text{ t C. ha}^{-1}$ in MSW and
 407 $16.76 \pm 1.69 \text{ t C. ha}^{-1}$ in CON-QA. The OWPs application significantly increased the size of the stable
 408 SOC pool in the BIOW ($31.29 \pm 0.91 \text{ t C. ha}^{-1}$) and MSW ($25.15 \pm 1.36 \text{ t C. ha}^{-1}$) treatments compared
 409 to the FYM ($25.15 \pm 1.44 \text{ t C. ha}^{-1}$) and CON-QA ($22.55 \pm 1.31 \text{ t C. ha}^{-1}$) which were statistically similar.
 410 Contrastingly, at La Cage experiment, 20 years of contrasted management had no significant effect on

411 the size of the stable SOC pool ($28.02 \pm 2.95 \text{ t C. ha}^{-1}$, $26.31 \pm 0.93 \text{ t C. ha}^{-1}$, and $26.08 \pm 1.89 \text{ t C. ha}^{-1}$
 412 for CA, ORG, and CON-LC respectively). However, CA significantly increased the size of the active
 413 pool ($29.15 \pm 5.79 \text{ t C. ha}^{-1}$) compared to ORG and CON-LC in which it was similar ($18.35 \pm 3.47 \text{ t C. ha}^{-1}$
 414 ha^{-1} and $16.14 \pm 0.97 \text{ t C. ha}^{-1}$ respectively) (Fig. 3b).
 415
 416
 417



418
 419 **Fig. 3** Soil organic carbon stock and additional carbon (Δ SOC) stock of bulk soils, active carbon (Ca)
 420 and stable carbon (Cs) ($n = 4$) at QualiAgro and La Cage experiments. The error bars represent the
 421 standard deviations. Grouped bars with different letters are significantly different between agricultural
 422 practices (Tukey-HSD, $p < 0.05$). CON-QA: conventional agriculture without organic inputs, BIOW:
 423 biowaste compost, MSW: municipal solid waste compost, FYM: farmyard manure, CON-LC:
 424 conventional agriculture, CA: conservation agriculture and ORG: organic agriculture.
 425

426 The results of additional carbon (i.e., the difference between the active or stable carbon pool of
 427 agroecological practices and the active or stable carbon pool of conventional control) distribution in the
 428 active and stable carbon pools are shown in Figs. 3c and 3d. In the QualiAgro experiment, BIOW (15.85
 429 $\pm 1.48 \text{ t C. ha}^{-1}$) and FYM ($13.36 \pm 1.16 \text{ t C. ha}^{-1}$) had similar active carbon pool size, higher than in the
 430 MSW ($12.34 \pm 0.75 \text{ t C. ha}^{-1}$). This active pool, represented 63-83 % of the additional carbon storage
 431 (Fig. 4c). Additional stable carbon pools ordered as follows: BIOW ($8.74 \pm 0.79 \text{ t C. ha}^{-1}$) > MSW (4.51
 432 $\pm 0.94 \text{ t C. ha}^{-1}$) > FYM ($2.60 \pm 0.97 \text{ t C. ha}^{-1}$) and represented between 17% (FYM) to 37% (BIOW) of

433 the additional carbon. At La Cage, 87% (CA) to 91% (ORG) of the additional carbon was in the active
 434 pool versus 9% (ORG) to 13% (CA) in the stable pool (Fig. 4d).

435


436 3.4 Carbon mineralization kinetics

437 At the end of soil incubation (day 484), the cumulative amounts of mineralized carbon expressed
 438 as percent soil organic carbon (%SOC) at La Cage experiment, differed significantly between the 3
 439 cropping systems, i.e., 12.60 ± 0.29 %SOC in ORG versus 11.52 ± 1.19 %SOC in CA and 10.21 ± 1.36
 440 %SOC in CON-LC (Fig. 4b). In the QualiAgro experiment, the specific carbon mineralization kinetics
 441 were significantly higher on the conventional control (CON-QA) without organic inputs compared to
 442 the soils receiving OWPs, where the mineralization carbon of the MSW and FYM plots were statistically
 443 identical but higher than the BIOW plot (Fig. 4a).

444

445 Overall, these two experiments show opposite trends. On the one hand, higher carbon
 446 mineralization under agroecological practices (ORG and CA) in La Cage experiment relative to the
 447 conventional control (CON-LC), and on the other hand, lower carbon mineralization under
 448 agroecological practices (MSW, FYM and BIOW) in QualiAgro experiment relative to the conventional
 449 control (CON-QA). Moreover, the percentage of additional carbon mineralized (% Δ SOC) at La Cage in
 450 CA (15% of Δ SOC) and ORG (57% of Δ SOC) was higher than at QualiAgro (4-5% of Δ SOC) (Figs. 4c
 451 and 4d). It must be noted however that the additional carbon stock was very small in the ORG treatment,
 452 which numerically explains the high % Δ SOC calculated value.

453



454

455 **Fig. 4.** Distribution of total carbon and additional carbon in carbon kinetic pools [C_{min} (carbon
456 mineralized), Active and Stable carbon] or fractions [POM and MAOM] under agricultural practices.
457 The error bars represent the standard errors. Grouped bars with different letters are significantly different
458 between agricultural practices (Tukey-HSD, $p < 0.05$).

459 **4 Discussion**

460 **4.1 Both POM-C and MAOM-C are sensitive to management**

461 The observed distribution of SOC in the fractions, i.e., most of the SOC (70-80%) being located
462 in the fine fraction ($<50 \mu\text{m}$), regardless of the agricultural practice implemented (Figs. 4a and 4b), is in
463 agreement with the literature (Christensen, 1985;1987; Balesdent, 1996; Baldock and Skjemstad, 2000;
464 Jolivet et al., 2003; Carter et al., 2003; Beare et al., 2014, Curtin et al., 2016, Poeplau et al., 2018).

465
466 Many studies have indicated that the POM-C fraction is more sensitive to land use and
467 management changes than the MAOM-C fraction (Cambardella and Elliot, 1992, Elliot et al., 1994,
468 Bayer et al., 2001, Salvo et al., 2010). However, our study showed that both POM-C and MAOM-C
469 fractions were both highly sensitive to the implementation of agroecological practices ca. 20 years. The
470 application of OWP resulted in additional soil organic carbon both as POM-C (34-40% of ΔSOC) and
471 MAOM-C (60-66% of ΔSOC), while conservation and organic agriculture resulted in additional organic
472 carbon mainly as MAOM-C (77-84% of ΔSOC) and less as POM-C (16-23% of ΔSOC). A significant
473 proportion of the additional carbon is associated with soil minerals, particularly in the clay fraction (See
474 Table. S1 and Table. S2).

475
476 Studies comparing no-tillage versus conventional tillage management showed in the surface layer
477 an increase in the POM-C fraction with no-tillage and no difference in MAOM-C (Wander et al., 1998,
478 Hussain et al., 1999; Carbonell-Bojollo et al., 2015; Samson et al., 2020). A recent meta-analysis by
479 Prairie et al. (2023) indicated that no-tillage increased both POM-C and MAOM-C fractions in soils
480 when this practice was maintained up to 6 years. However, the increase in the MAOM-C fraction was
481 less important than that of the POM-C fraction. We therefore ascribe the observed increase of MAOM-
482 C at La Cage to the no-tillage and the introduction of cover crops and the diversification of species (e.g.,
483 legumes) in the rotation. Interestingly, an earlier analysis of SOC distribution at La Cage, after 5 years
484 of differentiation, showed a significant increase of POM-C in the conservation agriculture system, while
485 no change of POM-C in the organic system and no significant change of the MAOM-C (Balabane et al.,
486 2005), suggesting either that it took more than 5 years for the additional POM-C to be broken down and
487 biodegraded as MAOM-C, or that the introduction of alfalfa as the cover crop instead of fescue since
488 2008 (i.e., 12 years later) resulted in more direct rhizodeposits inputs to MAOM-C. Indeed, according
489 to Autret et al. (2016), the estimated inputs from fescue were lower ($0.88 \text{ t C. ha}^{-1} \cdot \text{yr}^{-1}$) than those
490 coming from alfalfa as a cover crop ($1.12 \text{ t C. ha}^{-1} \cdot \text{yr}^{-1}$), about half of these amounts deriving from root

491 material. The cover crops and legume rotation in conservation agriculture and the legume rotations in
492 organic agriculture would likely have affected carbon input via the root system as dead roots (POM)
493 and rhizodeposits (MAOM). This would explain the high proportion of carbon associated with MAOM-
494 C. Typically, the cover crops characterized by low litter quality (e.g., grass) resulted in higher
495 accumulation of POM that was abundant in plant carbon, while cover crops with high litter quality (e.g.,
496 legumes) contributed to higher accumulation of SOC (Cotrufo et al., 2013), and of microbial necromass
497 carbon (Zhang et al., 2022) in MAOM-C. Thus, the high proportion of carbon in the MAOM-C at La
498 Cage (77-84% of Δ SOC) compared to QualiAgro (60-66% of Δ SOC) could be explained by the type
499 and quality of the carbon input. Because, the cover crops increase the time period in which plant roots
500 interact with the soil environment (Tiemann et al., 2015), they deliver an additional source of root litter
501 and exudates, providing greater diversity in belowground inputs (Austin et al., 2017). This promotes the
502 microbial growth and turnover in rhizosphere hotspots, processes that can enhance the formation of
503 MAOM (Kallenbach et al., 2016).

504
505 The recent meta-analysis, conducted by Zhang et al. (2022), indicated that the application of OWPs
506 significantly increases both MAOM and POM fractions in the soil relative to the control treatment,
507 which is consistent with our results. Peltre (2010) observed that the short-term application (4 times) of
508 the OWPs at QualiAgro increased the additional carbon only in the POM-C fraction, the MAOM-C
509 fraction $<50\mu\text{m}$ being unchanged. Paesch et al. (2016) later found that 7 successive applications of the
510 OWPs led to additional carbon in occluded small POM ($<20\mu\text{m}$) and in the fine silt + clay fraction ($<$
511 $6.3\mu\text{m}$). After 11 applications of OWPs we observed an increase in fine POM (50-200 μm), coarse silt
512 (20-50 μm) and MAOM-C ($<50\mu\text{m}$). This series of results indicate that the application of the OWPs
513 increase in the short term the POM-C fraction and that in a longer term (>10 years) the organic carbon
514 in the POM-C is transferred to the MAOM-C through biological activity in the soil. The transfer of
515 additional carbon from POM to MAOM is however slower at QualiAgro compared to La Cage. Cotrufo
516 et al. (2015), shows that POM and MAOM likely form under different biochemical and physical
517 pathways. While there is certainly some transfer from POM to MAOM, much of MAOM formation
518 occurs from dissolve organic matter (DOM) early during decomposition. There might just be a great
519 abundance of DOM and labile inputs int La Cage to explain greater MAOM formation.

520

521 **4.2 POM heterogeneity can hamper SOC stability assessments**

522 The POM in this study consists of crop residues and/or added manure or composts and microbial
523 residues. The agroecological practices with equivalent additional carbon stocks (MSW, FYM, CA)
524 showed after 20 years a higher proportion of additional carbon in POM-C under MSW (40% of Δ SOC)
525 and FYM (34% of Δ SOC) compared to CA (16% of Δ SOC). These results show that it is likely that
526 different management (e.g., OWPs application, no-tillage, cover crops and legume) alter the way gross

527 organic carbon inputs were distributed among the different organic carbon fractions. These results can
528 be explained by the fact that the decomposition rate of organic amendments and the SOC formed and
529 remaining in the long term vary according to the intrinsic quality of the amendment (Lashermes et al.,
530 2009). For example, Paustian et al. (1992) observed that high lignin content of FYM, which was more
531 recalcitrant to decomposition, resulted in greater accumulation of C than lower lignin amendments, such
532 as straw. Previous studies demonstrated that the OWPs generally are partially stabilized by the
533 composting and storage processes (Benbi and Khosa, 2014), unlike plant biomass, which is fresh OM.
534

535 The incubations revealed that in La Cage experiment, a higher percentage of the additional carbon
536 was mineralized in conservation agriculture (15% of Δ SOC) over 484 days than additional carbon at
537 QualiAgro (4-5% of Δ SOC) (Figs. 4c and 4d). The low additional carbon mineralization at QualiAgro
538 raises questions about the degradability of POM derived from OWPs, which were in higher proportion
539 (34-40% of Δ SOC) than at La Cage experiment (16-23% of Δ SOC). It is therefore likely that the OWPs-
540 derived POM were more recalcitrant with higher mean residence times compared to plant-derived POM.
541 The mean residence time of < 20 years given to POM in the study by Balesdent, (1996) may not be
542 applicable to systems where pre-processed exogenous OM are applied, because in the formed is based
543 on situations where organic input were crop residues. Thus, we assume a greater chemical recalcitrance
544 of POM-C in plots receiving OWPs, thereby reducing decomposers activity and carbon transfer to the
545 fine soil fraction (<50 μ m).
546

547 **4.3 Different methods provide a contrasted evaluation of biogeochemical** 548 **stability**

549 We used different methods to assess the biogeochemical stability of the additional carbon stored
550 in soil thanks to specific management options. The incubation method isolates carbon with mean
551 residence time (MRT) ranging from days to years (MRT < 2y in this case), while particle size and density
552 fractionation isolate carbon fractions ranging from years to decades (POM with MRT < 20y and MAOM
553 with MRT > 50y) and PARTY_{SOC} model based to RE thermal analysis that isolate carbon pools ranging
554 from decades to centuries (Active pool with MRT ~30-40y and Stable pool with MRT > 100y).
555

556 In the QualiAgro experiment, the incubations results indicate greater stability of additional carbon
557 compared to bulk SOC in the conventional control (i.e., lower specific carbon mineralization for soils
558 receiving OWPs relative to CON-QA). However, the results of particle size and density fractionation
559 and PARTY_{SOC} based to RE thermal analysis indicate that the additional carbon stored by OWPs
560 application is on average less stable than the soil carbon in the conventional control (CON-QA). This is
561 because, in these plots the additional carbon has a higher proportion of POM (MRT < 20y) and Active
562 carbon (MRT ~ 30-40y) than the conventional control (Figs. 4a and 4c). As the incubations target carbon

563 with MRT of the order of incubation length (i.e., $MRT < 2y$ in this study), we posit that this difference
564 is due to the fact that the different methods do not target the carbon pools with the same MRT. Put
565 together, these results suggest that, on the scale of a few decades, soil additional carbon in QualiAgro
566 experiment is less stable than soil carbon in conventional control, but in a shorter term (i.e., $MRT < 2y$),
567 the additional carbon is quite resistant.

568

569 In La Cage experiment, the results of the incubations and the PARTYSOC model based to RE
570 thermal analysis are consistent and indicate that the additional carbon stored by conservation agriculture
571 and organic agriculture is less stable than the soil carbon in the conventional control (CON-LC), whereas
572 the particle size and density fractionation indicates a more stable additional carbon, i.e., a higher
573 proportion of MAOM ($MRT > 50$ years) than the conventional control (Fig. 4). However, the study by
574 von Lützow et al. (2006) showed that MAOM does not have a unique mean residence time. For example,
575 land-use change (native and cropped lands) studies have indicated a decrease in carbon content in
576 MAOM-C over time (Balesdent et al., 1998, Yeasmin et al., 2019). Lutfalla et al. (2019), using samples
577 from 42 plots in Versailles, observed a decrease in carbon content in the clay fraction ($< 2 \mu m$) after 52
578 years of bare fallow conditions, thus questioning the long-term persistence of carbon associated with
579 clays and MAOM-C. Our results provide evidence that at least part of the carbon contained in MAOM
580 may not persist in soils over the long term as shown by others previously (e.g., Balesdent, 1987,
581 Keiluweit et al., 2015, Lutfalla et al., 2019, Chassé et al., 2021). We therefore hypothesize that the
582 additional carbon stored in the form of MAOM has a lower MRT than the MAOM in conventional
583 control.

584

585 Based on these results, our hypothesis that the biogeochemical stability of additional carbon is less
586 stable than the carbon in the conventional control is not always verified. However, considering that
587 MAOM is kinetically heterogeneous, then the results of these methods can be reconciled. So, the
588 additional carbon is overall less stable at a decadal or pluri-decadal timescale than the carbon stored in
589 the conventional control in both long-term experiments. Furthermore, taking all these elements and the
590 complementary nature of the methods into consideration, it emerges that the additional carbon stored
591 thanks to OWPs application is more stable in the short ($MRT < 2y$) and long term ($MRT > 100y$) than
592 the additional carbon enabled by alternative cropping systems, but less in the decadal and pluridecadal
593 time scale. The large time scale of SOM persistence shows that qualifying SOC simply as stable or labile
594 is not sufficient. It is essential to always associate a temporality with the biogeochemical stability that
595 is described in order to better assess the persistence of carbon in soils.

596

597

5 Conclusion

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

Acknowledgements

622

623

624

625

626

627

628

629

630

631

This study provided detailed information on the biogeochemical stability of additional carbon via a multi-methods evaluation. Soils from the same experimental sites but under widely contrasting management have resulted in contrasting carbon contents and stocks ca. 20 years of management. The results of particle size and density fractionation and PARTY_{SOC} model suggests that the additional carbon contained in MAOM may not persist in soils over the long term (> 50 years). Incubation, on the other hand, provided information on the short-term stability of additional carbon (i.e., MRT <2 years). Overall, the multi-methods evaluation showed that additional carbon was less stable at the decadal and pluri-decadal time-scales than carbon under conventional controls. However, incubations and PARTY_{SOC} model based to RE thermal analysis revealed that additional SOC in the QualiAgro experiment was more stable in short- term (MRT < 2y) and long- term (MRT >100y) than that in La Cage experiment. Additional SOC deriving from organic wastes, i.e., biomass that has been partially decomposed and transformed through its processing (digestion by cattle, storage and composting) prior to its incorporation in soil, would have a different biogeochemical stability than that deriving directly from plant biomass. Widely used (incubation, particle size fractionation) and increasingly used methods (RE) provide seemingly inconsistent assessments of the biogeochemical stability of SOC. These apparent contradictions can be explained by the fact that they address different kinetic pools of organic carbon. Care must be taken to specify which range of residence times are considered when using any method intending to evaluate the biogeochemical stability of SOM, as well as when using the terms stable or labile. As we found that the additional SOC stored thanks to the implementation of different management options had contrasted biogeochemical stabilities, there is a need to evaluate the biogeochemical stability of the additional SOC stored via other management options (e.g., agroforestry, lengthening temporary leys, no tillage...).

This research has been supported by the French Agence Nationale de la Recherche (StoreSoilC project, grant [ANR- 17-CE32-0005](#)). C. Chenu also thanks the CLand programme (ANR-16-CONV-0003). T. P. I. Kpemoua acknowledges the support of ADEME. The QualiAgro field experiment forms part of the SOERE-PRO (network of long-term experiments dedicated to the study of impacts of organic waste product recycling) integrated as a service of the ‘‘Investment for future’’ infrastructure AnaEE-France, overseen by the French National Research Agency (ANR-11-INBS-0001)’. The QualiAgro site is conducted within a collaboration between INRAE and Veolia. The authors also thank the personnel from Unité expérimentale Versailles Grignon INRAE, Michel Bertrand for the maintenance of and access to the La Cage long term experiment and Fabien Ferchaud for the collaboration upon La Cage SOC stocks assessment.

632

Author contribution

633 TPIK, CC, PB and SH designed the study. TPIK performed soil fractionation and long-term incubation.
634 FB and CP performed the RE6 thermal analyses and elementary analyses respectively. PB performed
635 the R codes to PARTY_{SOC} machine learning model. TPIK wrote the R codes and performed all
636 statistical analyses. TPIK, CC, PB, SH and FB contributed to the interpretation of the results. TPIK
637 prepared the paper with contributions from all coauthors.

638

639

640

641

642

References

643 Austin, E.E., Wickings, K., McDaniel, M.D., Robertson, G.P., Grandy, A.S.: Cover crop root
644 contributions to soil carbon in a no-till corn bioenergy cropping system. *GCB Bioenergy* 9,
645 1252–1263. <https://doi.org/10.1111/gcbb.12428>, 2017.

646 Autret, B., Guillier, H., Pouteau, V., Mary, B., Chenu, C.: Similar specific mineralization rates of
647 organic carbon and nitrogen in incubated soils under contrasted arable cropping systems. *Soil
648 and Tillage Research* 204, 104712. <https://doi.org/10.1016/j.still.2020.104712>, 2020.

649 Autret, B., Mary, B., Chenu, C., Balabane, M., Girardin, C., Bertrand, M., Grandeau, G., Beaudoin, N.:
650 Alternative arable cropping systems: A key to increase soil organic carbon storage? Results
651 from a 16 year field experiment. *Agriculture, Ecosystems & Environment* 232, 150–164.
652 <https://doi.org/10.1016/j.agee.2016.07.008>, 2016.

653 Balabane, M., Bureau, F., Decaens, T., Akpa, M., Hedde, M., Laval, K., Puget, P., Pawlak, B., Barray,
654 S., Cluzeau, D., Labreuche, J., Bodet, J.M., Le Bissonnais, Y., Saulas, P., Bertrand, M.,
655 Guichard, L., Picard, D., Houot, S., Arrouays, D., Brygoo, Y., Chenu, C.: Restauration de
656 fonctions et propriétés des sols de grande culture intensive. Effets de systèmes de culture
657 alternatifs sur les matières organiques et la structure des sols limoneux et approche du rôle
658 fonctionnel de la diversité biologique des sols (Dmostra), Rapport final de contrat MEDD
659 01105. *Gestion durable des Sols*. INRA, p. 119, 2005.

660 Baldock, J.A., Skjemstad, J.O.: Role of the soil matrix and minerals in protecting natural organic
661 materials against biological attack. *Organic Geochemistry* 31, 697–710.
662 [https://doi.org/10.1016/S0146-6380\(00\)00049-8](https://doi.org/10.1016/S0146-6380(00)00049-8), 2000.

663 Balesdent, J., Besnard, E., Arrouays, D., Chenu, C.: The dynamics of carbon in particle-size fractions
664 of soil in a forest-cultivation sequence. *Plant and Soil* 201, 49–57.
665 <https://doi.org/10.1023/A:1004337314970>, 1998.

666 Balesdent, J.: The significance of organic separates to carbon dynamics and its modelling in some
667 cultivated soils. *European Journal of Soil Science* 47, 485–493. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2389.1996.tb01848.x)
668 [2389.1996.tb01848.x](https://doi.org/10.1111/j.1365-2389.1996.tb01848.x), 1996.

669 Balesdent, J., Pétraud, J.P., Feller, C.: Effet des ultrasons sur la distribution granulométrique des
670 matières organiques des sols. *Science du Sol* 29, 95-106, 1991.

671 Balesdent, J.: The turnover of soil organic fractions estimated by radiocarbon dating. *Science of The*
672 *Total Environment* 62, 405–408. [https://doi.org/10.1016/0048-9697\(87\)90528-6](https://doi.org/10.1016/0048-9697(87)90528-6), 1987.

673 Barré, P., Plante, A.F., Cécillon, L., Lutfalla, S., Baudin, F., Bernard, S., Christensen, B.T., Eglin, T.,
674 Fernandez, J.M., Houot, S., Kätterer, T., Le Guillou, C., Macdonald, A., van Oort, F., Chenu,
675 C. :The energetic and chemical signatures of persistent soil organic matter. *Biogeochemistry*
676 130, 1–12. <https://doi.org/10.1007/s10533-016-0246-0>, 2016.

677 Baudin, F., Disnar, J.-R., Aboussou, A., Savignac, F.: Guidelines for Rock–Eval analysis of recent
678 marine sediments. *Organic Geochemistry* 86, 71–80.
679 <https://doi.org/10.1016/j.orggeochem.2015.06.009>, 2015.

680 Bayer, C., Martin-Neto, L., Mielniczuk, J., Pillon, C.N., Sangoi, L. : Changes in Soil Organic Matter
681 Fractions under Subtropical No-Till Cropping Systems. *Soil Sci. Soc. Am. J.* 65, 1473–1478.
682 <https://doi.org/10.2136/sssaj2001.6551473x>, 2001.

683 Beare, M.H., McNeill, S.J., Curtin, D., Parfitt, R.L., Jones, H.S., Dodd, M.B., Sharp, J.: Estimating the
684 organic carbon stabilisation capacity and saturation deficit of soils: a New Zealand case study.
685 *Biogeochemistry* 120, 71–87. <https://doi.org/10.1007/s10533-014-9982-1>, 2014.

686 Benbi, D.K., Khosa, M.K.: Effects of Temperature, Moisture, and Chemical Composition of Organic
687 Substrates on C Mineralization in Soils. *Communications in Soil Science and Plant Analysis*
688 45, 2734–2753. <https://doi.org/10.1080/00103624.2014.950423>, 2014.

689 Bohoussou, Y.N., Kou, Y.-H., Yu, W.-B., Lin, B., Virk, A.L., Zhao, X., Dang, Y.P., Zhang, H.-L.:
690 Impacts of the components of conservation agriculture on soil organic carbon and total nitrogen
691 storage: A global meta-analysis. *Science of The Total Environment* 842, 156822.
692 <https://doi.org/10.1016/j.scitotenv.2022.156822>, 2022.

693 Bruni, E., Chenu, C., Abramoff, R.Z., Baldoni, G., Barkusky, D., Clivot, H., Huang, Y., Kätterer, T.,
694 Pikula, D., Spiegel, H., Virto, I., Guenet, B.: Multi-modelling predictions show high uncertainty
695 of required carbon input changes to reach a 4‰ target. *European J Soil Science*.
696 <https://doi.org/10.1111/ejss.13330>, 2022.

697 Cambardella, C.A., Elliott, E.T.: Particulate Soil Organic-Matter Changes across a Grassland
698 Cultivation Sequence. *Soil Sci. Soc. Am. J.* 56, 777–783.
699 <https://doi.org/10.2136/sssaj1992.03615995005600030017x>, 1992.

700 Carbonell-Bojollo, R., González-Sánchez, E.J., Ruibérriz de Torres, M.R., Ordóñez-Fernández, R.,
701 Domínguez-Gimenez, J., Basch, G.: Soil organic carbon fractions under conventional and no-

702 till management in a long-term study in southern Spain. *Soil Res.* 53, 113.
703 <https://doi.org/10.1071/SR13369>, 2015.

704 Carter, M.R., Angers, D.A., Gregorich, E.G., Bolinder, M.A.: Characterizing organic matter retention
705 for surface soils in eastern Canada using density and particle size fractions. *Can. J. Soil. Sci.* 83,
706 11–23. <https://doi.org/10.4141/S01-087>, 2003.

707 Cécillon, L., Baudin, F., Chenu, C., Christensen, B.T., Franko, U., Houot, S., Kanari, E., Kätterer, T.,
708 Merbach, I., van Oort, F., Poeplau, C., Quezada, J.C., Savignac, F., Soucémariadin, L.N.,
709 Barré, P.: Partitioning soil organic carbon into its centennially stable and active fractions with
710 machine-learning models based on Rock-Eval® thermal analysis
711 (PARTY<sub></sub>SOC<sub></sub>v2.0 and
712 PARTY<sub></sub>SOC<sub></sub>v2.0<sub></sub>EU<sub></sub>). *Geosci. Model Dev.*
713 14, 3879–3898. <https://doi.org/10.5194/gmd-14-3879-2021>, 2021.

714 Cécillon, L., Baudin, F., Chenu, C., Houot, S., Jolivet, R., Kätterer, T., Lutfalla, S., Macdonald, A., van
715 Oort, F., Plante, A.F., Savignac, F., Soucémariadin, L.N., Barré, P.: A model based on Rock-
716 Eval thermal analysis to quantify the size of the centennially persistent organic carbon pool in
717 temperate soils. *Biogeosciences* 15, 2835–2849. <https://doi.org/10.5194/bg-15-2835-2018>,
718 2018.

719 Chassé, M., Lutfalla, S., Cécillon, L., Baudin, F., Abiven, S., Chenu, C., Barré, P.: Long-term bare-
720 fallow soil fractions reveal thermo-chemical properties controlling soil organic carbon
721 dynamics. *Biogeosciences* 18, 1703–1718. <https://doi.org/10.5194/bg-18-1703-2021>, 2021.

722 Chenu, C., Rumpel, C., Lehmann, J.: Methods for Studying Soil Organic Matter, in: *Soil Microbiology,*
723 *Ecology and Biochemistry.* Elsevier, pp. 383–419. [https://doi.org/10.1016/B978-0-12-415955-
724 6.00013-X](https://doi.org/10.1016/B978-0-12-415955-6.00013-X), 2015.

725 Chenu, C., Angers, D.A., Barré, P., Derrien, D., Arrouays, D., Balesdent, J.: Increasing organic stocks
726 in agricultural soils: Knowledge gaps and potential innovations. *Soil and Tillage Research* 188,
727 41–52. <https://doi.org/10.1016/j.still.2018.04.011>, 2019.

728 Christensen, B.T.: Decomposability of organic matter in particle size fractions from field soils with
729 straw incorporation. *Soil Biology and Biochemistry* 19, 429–435.
730 [https://doi.org/10.1016/0038-0717\(87\)90034-4](https://doi.org/10.1016/0038-0717(87)90034-4), 1987.

731 Christensen, B.T.: Carbon and Nitrogen in Particle Size Fractions Isolated from Danish Arable Soils by
732 Ultrasonic Dispersion and Gravity-Sedimentation. *Acta Agriculturae Scandinavica* 35, 175–
733 187. <https://doi.org/10.1080/00015128509435773>, 1985.

734 Clivot, H., Mouny, J.-C., Duparque, A., Dinh, J.-L., Denoroy, P., Houot, S., Vertès, F., Trochard, R.,
735 Bouthier, A., Sagot, S., Mary, B.: Modeling soil organic carbon evolution in long-term arable
736 experiments with AMG model. *Environmental Modelling & Software* 118, 99–113.
737 <https://doi.org/10.1016/j.envsoft.2019.04.004>, 2019.

738 Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Deneff, K., Paul, E.: The Microbial Efficiency-Matrix
739 Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter
740 stabilization: do labile plant inputs form stable soil organic matter? *Glob Change Biol* 19, 988–
741 995. <https://doi.org/10.1111/gcb.12113>, 2013.

742 Curtin, D., Beare, M.H., Qiu, W.: Texture effects on carbon stabilisation and storage in New Zealand
743 soils containing predominantly 2 : 1 clays. *Soil Res.* 54, 30.
744 <https://doi.org/10.1071/SR14292>, 2016.

745 Disnar, J.R., Guillet, B., Keravis, D., Di-Giovanni, C., Sebag, D.: Soil organic matter (SOM)
746 characterization by Rock-Eval pyrolysis: scope and limitations. *Organic Geochemistry* 34, 327–
747 343. [https://doi.org/10.1016/S0146-6380\(02\)00239-5](https://doi.org/10.1016/S0146-6380(02)00239-5), 2003.

748 Ellert, B.H., Bettany, J.R.: Calculation of organic matter and nutrients stored in soils under contrasting
749 management regimes. *Can. J. Soil. Sci.* 75, 529–538. <https://doi.org/10.4141/cjss95-075>, 1995.

750 Fujisaki, K., Chapuis-Lardy, L., Albrecht, A., Razafimbelo, T., Chotte, J.-L., Chevallier, T.: Data
751 synthesis of carbon distribution in particle size fractions of tropical soils: Implications for soil
752 carbon storage potential in croplands. *Geoderma* 313, 41–51.
753 <https://doi.org/10.1016/j.geoderma.2017.10.010>, 2018.

754 Haddaway, N.R., Hedlund, K., Jackson, L.E., Kätterer, T., Lugato, E., Thomsen, I.K., Jørgensen, H.B.,
755 Isberg, P.-E.: How does tillage intensity affect soil organic carbon? A systematic review.
756 *Environ Evid* 6, 30. <https://doi.org/10.1186/s13750-017-0108-9>, 2017.

757 Hussain, I., Olson, K.R., Ebelhar, S.A.: Long-Term Tillage Effects on Soil Chemical Properties and
758 Organic Matter Fractions. *Soil Sci. Soc. Am. J.* 63, 1335–1341.
759 <https://doi.org/10.2136/sssaj1999.6351335x>, 1999.

760 Hsieh, Y.-P.: Pool Size and Mean Age of Stable Soil Organic Carbon in Croplands. *Soil Science Society*
761 *of America Journal* 56, NP-NP.
762 <https://doi.org/10.2136/sssaj1992.03615995005600020049x>, 1992.

763 IUSS Working Group WRB.: World Reference Base for Soil Resources 2014, Update 2015.
764 International Soil Classification System for Naming Soils and Creating Legends for Soil Maps.
765 World Soil Resources Reports No. 106, Rome: FAO, 2015.

766 Jastrow, J.D., Miller, R.M., Boutton, T.W.: Carbon Dynamics of Aggregate-Associated Organic Matter
767 Estimated by Carbon-13 Natural Abundance. *Soil Science Society of America Journal* 60, 801–
768 807. <https://doi.org/10.2136/sssaj1996.03615995006000030017x>, 1996.

769 Jolivet, C., Arrouays, D., Lévêque, J., Andreux, F., Chenu, C.: Organic carbon dynamics in soil particle-
770 size separates of sandy Spodosols when forest is cleared for maize cropping: Organic carbon
771 dynamics in soil fractions. *European Journal of Soil Science* 54, 257–268.
772 <https://doi.org/10.1046/j.1365-2389.2003.00541.x>, 2003.

773 Kallenbach, C.M., Frey, S.D., Grandy, A.S.: Direct evidence for microbial-derived soil organic matter
774 formation and its ecophysiological controls. *Nat Commun* 7, 13630.
775 <https://doi.org/10.1038/ncomms13630>, 2016.

776 Kanari, E., Cécillon, L., Baudin, F., Clivot, H., Ferchaud, F., Houot, S., Levavasseur, F., Mary, B.,
777 Soucémariadin, L., Chenu, C., and Barré, P.: A robust initialization method for accurate soil
778 organic carbon simulations, *Biogeosciences*, 19, 375–387, [https://doi.org/10.5194/bg-19-375-](https://doi.org/10.5194/bg-19-375-2022)
779 2022, 2022.

780 Kanari, E., Barré, P., Baudin, F., Berthelot, A., Bouton, N., Gosselin, F., Soucémariadin, L., Savignac,
781 F., Cécillon, L.: Predicting Rock-Eval® thermal analysis parameters of a soil layer based on
782 samples from its sublayers; an experimental study on forest soils. *Organic Geochemistry* 160,
783 104289. <https://doi.org/10.1016/j.orggeochem.2021.104289>, 2021.

784 Kim, K., Daly, E. J., Gorzelak, M., and Hernandez-Ramirez, G.: Soil organic matter pools response to
785 perennial grain cropping and nitrogen fertilizer, *Soil Till. Res.*, 220, 105376,
786 <https://doi.org/10.1016/j.still.2022.105376>, 2022.

787 Kpemoua, T.P.I., Leclerc, S., Barré, P., Houot, S., Pouteau, V., Plessis, C., Chenu, C. : Are carbon-storing soils
788 more sensitive to climate change? A laboratory evaluation for agricultural temperate soils, *Soil Biology*
789 *and Biochemistry*, Volume 183, 2023, 109043.
790 <https://doi.org/10.1016/j.soilbio.2023.109043>, 2023.

791 Kuzyakov, Y., Bol, R.: Using natural ¹³C abundances to differentiate between three CO₂ sources
792 during incubation of a grassland soil amended with slurry and sugar. *Z. Pflanzenernähr. Bodenk.*
793 167, 669–677. <https://doi.org/10.1002/jpln.200421412>, 2004.

794 Lafargue E., Marquis F., Pillot D.: Rock-Eval 6 applications in hydrocarbon exploration, production and
795 soil contamination studies, *Oil Gas Sci. Technol. – Rev. IFP* 1998; 53, 421–437.
796 doi.org/10.2516/ogst:1998036, 2018.

797 Lal, R.: Digging deeper: A holistic perspective of factors affecting soil organic carbon sequestration in
798 agroecosystems. *Global Change Biology* 24, 3285–3301.
799 <https://doi.org/10.1111/gcb.14054>, 2018.

800 Lal, R.: Soil Carbon Sequestration Impacts on Global Climate Change and Food Security. *Science* 304,
801 1623–1627. <https://doi.org/10.1126/science.1097396>, 2004.

802 Lashermes, G., Nicolardot, B., Parnaudeau, V., Thuriès, L., Chaussod, R., Guillotin, M.L., Linères, M.,
803 Mary, B., Metzger, L., Morvan, T., Tricaud, A., Villette, C., Houot, S.: Indicator of potential
804 residual carbon in soils after exogenous organic matter application. *European Journal of Soil*
805 *Science* 60, 297–310. <https://doi.org/10.1111/j.1365-2389.2008.01110.x>, 2009.

806 Lutfalla, S., Barré, P., Bernard, S., Le Guillou, C., Alléon, J., Chenu, C.: Multidecadal persistence of
807 organic matter in soils: multiscale investigations down to the submicron scale. *Biogeosciences*
808 16, 1401–1410. <https://doi.org/10.5194/bg-16-1401-2019>, 2019.

809 Pacini, L., Adatte, T., Barré, P., Boussafir, M., Bouton, N., Cécillon, L., Lamoureux-Var, V., Sebag, D.,
810 Verrecchia, E., Wattripont, A., Baudin, F.: Reproducibility of Rock-Eval® thermal analysis for
811 soil organic matter characterization. *Organic Geochemistry* 186, 104687.
812 <https://doi.org/10.1016/j.orggeochem.2023.104687>, 2023.

813 Paetsch, L., Mueller, C.W., Rumpel, C., Houot, S., Kögel-Knabner, I.: Urban waste composts enhance
814 OC and N stocks after long-term amendment but do not alter organic matter composition.
815 *Agriculture, Ecosystems & Environment* 223, 211–222,
816 <https://doi.org/10.1016/j.agee.2016.03.008>, 2016.

817 Paustian, K., Parton, W.J., Persson, J.: Modeling Soil Organic Matter in Organic-Amended and
818 Nitrogen-Fertilized Long-Term Plots. *Soil Science Society of America Journal* 56, 476–488.
819 <https://doi.org/10.2136/sssaj1992.03615995005600020023x>, 1992.

820 Pellerin, S., Bamière, L., Launay, C., Martin, R., Angers, D., Balesdent, J., Basile-Doelsch, I.,
821 Bellassen, V., Cardinael, R., Cécillon, L., Ceschia, E., Chenu, C., Constantin, J., Darroussin, J.,
822 Delacote, P., Delame, N., Gastal, F., Gilbert, D., Schiavo, M.: Stocker du carbone dans les sols
823 français, Quel potentiel au regard de l'objectif de 4 pour 1000 et à quel coût? Synthèse du
824 rapport d'étude, INRA (France), 2019.

825 Peltre, C.: Potential carbon storage in soil after exogenous organic matter applications. *Soil*
826 *study*. AgroParisTech. English. NNT: 2010AGPT0076. pastel-00602825, 2010.

827 Peltre, C., Christensen, B.T., Dragon, S., Icard, C., Kätterer, T., Houot, S.: RothC simulation of carbon
828 accumulation in soil after repeated application of widely different organic amendments. *Soil*
829 *Biology and Biochemistry* 52, 49–60. <https://doi.org/10.1016/j.soilbio.2012.03.023>, 2012.

830 Peltre, C., Fernández, J.M., Craine, J.M., Plante, A.F.: Relationships between Biological and Thermal
831 Indices of Soil Organic Matter Stability Differ with Soil Organic Carbon Level. *Soil Sci. Soc.*
832 *Am. j.* 77, 2020–2028. <https://doi.org/10.2136/sssaj2013.02.0081>, 2013.

833 Plante, A.F., Fernández, J.M., Leifeld, J.: Application of thermal analysis techniques in soil science.
834 *Geoderma* 153, 1–10. <https://doi.org/10.1016/j.geoderma.2009.08.016>, 2009.

835 Poeplau, C., Barré, P., Cécillon, L., Baudin, F., Sigurdsson, B.D.: Changes in the Rock-Eval signature
836 of soil organic carbon upon extreme soil warming and chemical oxidation - A comparison.
837 *Geoderma* 337, 181–190. <https://doi.org/10.1016/j.geoderma.2018.09.025>, 2019.

838 Poeplau, C., Don, A.: Carbon sequestration in agricultural soils via cultivation of cover crops – A meta-
839 analysis. *Agriculture, Ecosystems & Environment* 200, 33–41.
840 <https://doi.org/10.1016/j.agee.2014.10.024>, 2015.

841 Poeplau, C., Don, A., Six, J., Kaiser, M., Benbi, D., Chenu, C., Cotrufo, M.F., Derrien, D., Gioacchini,
842 P., Grand, S., Gregorich, E., Griepentrog, M., Gunina, A., Haddix, M., Kuzyakov, Y., Kühnel,
843 A., Macdonald, L.M., Soong, J., Trigalet, S., Vermeire, M.-L., Rovira, P., van Wesemael, B.,
844 Wiesmeier, M., Yeasmin, S., Yevdokimov, I., Nieder, R.: Isolating organic carbon fractions

845 with varying turnover rates in temperate agricultural soils – A comprehensive method
846 comparison. *Soil Biology and Biochemistry* 125, 10–26.
847 <https://doi.org/10.1016/j.soilbio.2018.06.025>, 2018.

848 Prairie, A.M., King, A.E., Cotrufo, M.F.: Restoring particulate and mineral-associated organic carbon
849 through regenerative agriculture. *Proc. Natl. Acad. Sci. U.S.A.* 120, e2217481120.
850 <https://doi.org/10.1073/pnas.2217481120>, 2023.

851 Rumpel, C., Amiraslani, F., Chenu, C., Cardenas, M.G., Kaonga, M., Koutika, L.S., Ladha, J., Madari,
852 B., Shirato, Y., Smith, P., Soudi, B., Soussana, J.F., Whitehead, D., Wollenberg, E.: The 4p1000
853 initiative: Opportunities, limitations and challenges for implementing soil organic carbon
854 sequestration as a sustainable development strategy. *Ambio* 49, 350-360. [https://doi-
855 org.ezproxy.universite-paris-saclay.fr/10.1007/s13280-019-01165-2](https://doi-org.ezproxy.universite-paris-saclay.fr/10.1007/s13280-019-01165-2), 2020.

856 Salvo, L., Hernández, J., Ernst, O.: Distribution of soil organic carbon in different size fractions, under
857 pasture and crop rotations with conventional tillage and no-till systems. *Soil and Tillage
858 Research* 109, 116–122. <https://doi.org/10.1016/j.still.2010.05.008>, 2010.

859 Samson, M.-E., Chantigny, M.H., Vanasse, A., Menasseri-Aubry, S., Royer, I., Angers, D.A.:
860 Management practices differently affect particulate and mineral-associated organic matter and
861 their precursors in arable soils. *Soil Biology and Biochemistry* 148, 107867.
862 <https://doi.org/10.1016/j.soilbio.2020.107867>, 2020.

863 Sanderman, J., Fillery, I. R. P., Jongepier, R., Massalsky, A., Roper, M. M., MacDonald, L. M.,
864 Maddern, T., Murphy, D. V., Baldock, J. A.: Carbon sequestration under subtropical perennial
865 pastures II: Carbon dynamics, *Soil Res.*, 51, 771–780, <https://doi.org/10.1071/SR12351>, 2013.

866 Sanderman, J., Grandy, A.S.: Ramped thermal analysis for isolating biologically meaningful soil organic
867 matter fractions with distinct residence times. *SOIL* 6, 131–144.
868 <https://doi.org/10.5194/soil-6-131-2020>, 2020.

869 Sokol, N.W., Sanderman, J., Bradford, M.A.: Pathways of mineral-associated soil organic matter
870 formation: Integrating the role of plant carbon source, chemistry, and point of entry. *Glob
871 Change Biol* 25, 12–24. <https://doi.org/10.1111/gcb.14482>, 2019.

872 Sollins, P., Swanston, C., Kleber, M., Filley, T., Kramer, M., Crow, S., Caldwell, B.A., Lajtha, K.,
873 Bowden, R.: Organic C and N stabilization in a forest soil: evidence from sequential density
874 fractionation. *Soil Biology and Biochemistry* 38, 3313-3324, [10.1016/j.soilbio.2006.04.014](https://doi.org/10.1016/j.soilbio.2006.04.014),
875 2006.

876 Tiemann, L.K., Grandy, A.S., Atkinson, E.E., Marin-Spiotta, E., McDaniel, M.D.: Crop rotational
877 diversity enhances belowground communities and functions in an agroecosystem. *Ecol Lett* 18,
878 761–771. <https://doi.org/10.1111/ele.12453>, 2015.

879 Torn, M.S., Kleber, M., Zavaleta, E.S., Zhu, B., Field, C.B., Trumbore, S.E.: A dual isotope approach
880 to isolate soil carbon pools of different turnover times. *Biogeosciences* 10, 8067–8081.
881 <https://doi.org/10.5194/bg-10-8067-2013>, 2013.

882 Virto, I., Barré, P., Burlot, A., Chenu, C.: Carbon input differences as the main factor explaining the
883 variability in soil organic C storage in no-tilled compared to inversion tilled agrosystems.
884 *Biogeochemistry* 108, 17–26. <https://doi.org/10.1007/s10533-011-9600-4>, 2012.

885 Von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E.,
886 Marschner, B.: SOM fractionation methods : Relevance to functional pools and to stabilization
887 mechanisms. *Soil Biology and Biochemistry* 39, 2183–2207.
888 <https://doi.org/10.1016/j.soilbio.2007.03.007>, 2007.

889 Von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B.,
890 Flessa, H.: Stabilization of organic matter in temperate soils: mechanisms and their relevance
891 under different soil conditions - a review: Mechanisms for organic matter stabilization in soils.
892 *Eur. J. Soil Sci.* 57 (4), 426–445. <https://doi.org/10.1111/j.1365-2389.2006.00809.x>, 2006.

893 Wander, M.M., Bidart, M.G., Aref, S.: Tillage Impacts on Depth Distribution of Total and Particulate
894 Organic Matter in Three Illinois Soils. *Soil Science Society of America Journal* 62, 1704–1711.
895 <https://doi.org/10.2136/sssaj1998.03615995006200060031x>, 1998.

896 Yeasmin, S., Singh, B., Johnston, C.T., Sparks, D.L., Hua, Q.: Changes in Particulate and Mineral
897 Associated Organic Carbon with Land Use in Contrasting Soils (preprint). *Biogeochemistry:*
898 *Soils.* <https://doi.org/10.5194/bg-2019-416>, 2019.

899 Zhang, F., Chen, X., Yao, S., Ye, Y., Zhang, B.: Responses of soil mineral-associated and particulate
900 organic carbon to carbon input: A meta-analysis. *Science of The Total Environment* 829,
901 154626. <https://doi.org/10.1016/j.scitotenv.2022.154626>, 2022.

902
903