



Nitrogen oxides emissions from selected cities in North America, Europe, and East Asia observed by TROPOMI before and after the COVID-19 pandemic

Chantelle R. Lonsdale¹ and Kang Sun^{1,2}

¹Department of Civil, Structural and Environmental Engineering, University at Buffalo, Buffalo, NY, USA

²Research and Education in Energy, Environment and Water Institute, University at Buffalo, Buffalo, NY, USA

Correspondence: Kang Sun (kangsun@buffalo.edu)

Abstract. Nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$) emissions are estimated in three regions in the Northern hemisphere, generally located in North America, Europe, and East Asia, by calculating the directional derivatives of NO_2 column amounts observed by the TROPOMI instrument with respect to the horizontal wind vectors. We present monthly averaged emissions from 1 May 2018 to 31 January 2023 to capture variations before and after the COVID-19 pandemic. We focus on a diverse collection of 5 54 cities, 18 in each region. A spatial resolution of 0.04° resolves intracity emission variations and reveals NO_x emission hot spots at city cores, industrial areas, and sea ports. For each selected city, COVID-19-induced changes in NO_x emissions are estimated by comparing monthly and annually averaged values to the pre-COVID-19 year of 2019. While emission reductions are initially found during the first outbreak of COVID-19 in early 2020 in most cities, the cities' paths diverge afterwards. We group the selected cities into 4 clusters according to their normalized annual NO_x emissions in 2019–2022 using an 10 unsupervised learning algorithm. All but one selected North American cities fall into cluster 1 characterized by weak emission reduction in 2020 (-7% relative to 2019) and increase in 2022 by $+5\%$. Cluster 2 contains mostly European cities and is characterized by the largest reduction in 2020 (-31%), whereas the selected East Asian cities generally fall into clusters 3 and 4 with the largest impacts in 2022 (-25% and -37%). This directional derivative approach has been implemented in object-oriented, open-source Python and is available publicly for high-resolution and low-latency emission estimation for different 15 regions, atmospheric species, and satellite instruments.

1 Introduction

The COVID-19 pandemic, which was caused by the SARS-CoV-2 virus emerged in 2019 and its still evolving variants as of writing in 2023, has resulted in unprecedented shifts in human activities and anthropogenic emissions to the Earth's atmosphere.

20 One of the most effective and important indicators of the COVID-19-induced emission perturbations is the emission of nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$) (Levelt et al., 2022, and references therein). The dominant NO_x emission source is anthropogenic



fossil fuel combustion. Because of its relatively short chemical lifetime, hot spots of NO_x abundance can be readily identified near the emission sources. Due to its adverse health effects, NO_x is a regulated primary pollutant with significant implications for secondary ozone and $\text{PM}_{2.5}$ formation and reactive nitrogen deposition (Seinfeld and Pandis, 2016; Zhang et al., 2012).

25 Accurate and timely quantification of NO_x emission is thus essential for environmental regulation, air quality forecasting, and improved understanding of atmospheric chemistry processes.

Bottom-up NO_x emission inventories have been extensively used in atmospheric composition, climate change, and human health studies from regional to global scales (Streets et al., 2003; Crippa et al., 2018; McDuffie et al., 2020; Zheng et al., 2021). However, the bottom-up emission estimates are subject to significant and often under-characterized uncertainties that originate
30 from the lack of knowledge of emission factors, chemical processes, and spatiotemporal proxies as well as the inconsistencies among different geographical datasets. Moreover, bottom-up emission inventories require significant effort and time to compile, leading to often years of lag time before producing results. It is especially challenging for the bottom-up approaches to represent COVID-19-induced emission changes, as both the spread of variants and the policy responses of governments worldwide have been rapidly changing.

35 Alternatively, satellite observations can assess NO_x emissions from a top-down perspective and in a more timely manner. Substantial efforts among the research community have been devoted to characterizing the NO_x emission responses in the early phase of the pandemic (Gkatzelis et al., 2021, and references therein). Satellite-observed NO_2 tropospheric column amounts have been used to infer COVID-induced NO_x emission perturbations through chemical transport models (CTMs) (Miyazaki et al., 2020; Ding et al., 2020; Riess et al., 2022; Kang et al., 2022), fitting of plume dispersion or box models (Sun et al.,
40 2021; Lange et al., 2022; Dammers et al., 2022; Xue et al., 2022; Godłowska et al., 2023; Zhang et al., 2023), and calculation of the divergence of horizontal NO_2 flux (the flux divergence approach hereafter, de Foy and Schauer, 2022; Dix et al., 2022; Rey-Pommier et al., 2022; Chen et al., 2023). Each approach comes with its own strengths and limitations. The CTM-based approach usually resolves emissions spatiotemporally and incorporates meteorological and chemical processes, but requires significant computation and auxiliary datasets, which hinders its agility. Analytical plume or box models are generally applied
45 to a single source region and do not resolve the spatial distribution of emissions. The flux divergence approach has the potential to rapidly map emissions over extensive areas, whereas only annual averaged emissions have been reported in specific regions.

Inspired by the flux divergence approach, Sun (2022a) proposed a unified framework capable of rapidly imaging NO_x emissions using only TROPOMI level 2 products and the ERA5 global reanalysis, both of which are available within a few days of lag time. Here we coin this framework as the directional derivative approach, as the flux divergence is not explicitly
50 calculated. Instead, the emission signal originates from the directional derivative of satellite observed column amount with respect to the horizontal wind vector. The impact of topography on emission estimation, which was neglected in the flux divergence literature, is accounted for through a similar directional derivative of surface altitude. In this work, we apply the directional derivative approach to map NO_x emissions at 0.04° grid size over extensive regions in North America, Europe, and East Asia. We focus on 18 selected cities in each region (54 cities in total) and quantify monthly NO_x emissions from 1 May
55 2018 to 31 January 2023. We systematically compare the emissions in 2019 as the pre-COVID-19 year with those in 2020–2022 as post-COVID-19 years. The large spatiotemporal variations of NO_x emissions after 2020 in comparison with 2019 highlight



the complexity of post-COVID-19 emission changes and the importance of timely and persistent observation-based constraints. The normalized annual NO_x emissions from all selected cities are grouped into 4 clusters using an unsupervised learning algorithm. While the initial emission reductions during the onset of the pandemic in 2020 are ubiquitous in all clusters, the 2021 and 2022 emissions diverge significantly. The directional derivative approach has been implemented in object-oriented, open-source Python (Sun, 2022b) and is available publicly for future applications in different regions, time periods, and other satellite instruments beyond TROPOMI.

2 Data

2.1 Data for emission calculation

Following Sun (2022a), we use the TROPOMI Products Algorithm Laboratory (PAL) level 2 NO_2 product from 1 May 2018 to 14 November 2021. The operational offline product is then merged, resulting in a seamless and consistent product generated by a single retrieval processor (version 2.3.1) (van Geffen et al., 2022a). The nadir TROPOMI level 2 pixel size was $3.5 \times 7 \text{ km}^2$ before 6 August 2019 and updated to $3.5 \times 5.5 \text{ km}^2$ thereafter. The equator crossing of TROPOMI is at around 13:30 local time, but due to its ground swath width of 2600 km, the measurement's local time at the swath edges may differ by ± 1 hour from the nadir. We use only the level 2 pixels with quality assurance values above 0.75 according to the recommendation from the product Algorithm Theoretical Basis Document (ATBD) (van Geffen et al., 2022b).

Besides the NO_2 tropospheric vertical column density, the TROPOMI product also provides surface altitude at each level 2 pixel sampled from the GMTED2010 digital elevation model and horizontal wind at 10 m above the surface sampled from ECMWF meteorology (Eskes et al., 2022). In addition, we sample horizontal winds at 100 m and 10 m above the surface from the ERA5 reanalysis (Hersbach et al., 2020) spatiotemporally at TROPOMI level 2 observations.

2.2 Data for urban area coverage

Although the NO_x emissions derived from TROPOMI-observed NO_2 column amounts cover all the regions, it is the urban areas that dominate the NO_x emission budget and respond most to the COVID-19-induced perturbations. Cities in different countries and continents underwent drastically different scenarios after the onset of the pandemic. The definition of each city boundary is often ambiguous and inconsistent among geographical regions and urban datasets. To consistently identify cities globally, we use the Global Human Built-up And Settlement Extent (HBASE) Dataset from Landsat as the indicator of urban area coverage (Wang et al., 2017). The HBASE dataset has a native resolution of 30 m for the target year 2010, whereas we use the aggregated version at 1 km resolution.

Cities in this study are selected from a world city list with population and city center coordinate information (Hernández, 2022). In each region of North America, Europe, and East Asia, we focus on two subregions in the north and south, based on latitude, climate, and proximity of city clusters. Within each subregion, we select 9 cities with the consideration of population and location. The bounds of each region, subregion, and the names of all selected cities are shown in Figures 2, 7, and 12.



For each city, we consider the urban area coverage given by the HBASE dataset within ± 50 km in the zonal and meridional directions from the city center coordinate as the extent of the city. This 100×100 km window covers most select cities sufficiently with frequent inclusion of surrounding satellite cities (see Figures 5, 6, 10, 11, 15, and 16 for the extents of individual cities). Large cities that are close together may be enveloped by the same window. For example, the Washington, DC window includes most of the area covered by Baltimore (Figure 6e), and the Wuxi window includes two similar sized cities, Changzhou and Suzhou (Figure 15g). Without attempting to separate them, we treat these cities in the same window as a single metropolitan area. However, we separate the adjacent cities at the USA-Mexico border, namely San Diego and Tijuana as well as El Paso and Juarez, because of significantly different NO_x emission patterns across the country border. Additionally, the windows for Los Angeles and Dallas are extended to cover the entire Los Angeles basin and the Dallas–Fort Worth–Arlington metropolitan area, and the windows for Wuxi, Tianjin, and Busan are slightly nudged to avoid cutting off significant emission sources near the defined city edge.

3 Methods

3.1 $\text{NO}_x:\text{NO}_2$ ratio

As TROPOMI only observes NO_2 column amounts, a molar ratio between NO_x and NO_2 is needed to derive NO_x column amounts and NO_x emissions. Here we use a constant $\text{NO}_x:\text{NO}_2$ ratio of 1.32 as suggested in many NO_x emission estimation studies (Beirle et al., 2011; Goldberg et al., 2019; Beirle et al., 2019; Dix et al., 2022; Goldberg et al., 2022; Sun, 2022a; Dammers et al., 2022) for all cities. More sophisticated considerations exist, which are based on the photostationary steady state assumption and model simulated ozone concentration (Beirle et al., 2021; Lange et al., 2022) or directly from model simulated NO and NO_2 (Lorente et al., 2019; Zhang et al., 2023). As the main focus of this work is the relative emission changes in the pre- and post-COVID-19 years, the impact of variable $\text{NO}_x:\text{NO}_2$ ratio will largely cancel out. Additionally, the tropospheric mean $\text{NO}_x:\text{NO}_2$ ratio estimated by Beirle et al. (2021) also does not show excessive variations over the three regions included in this study. Moreover, the $\text{NO}_x:\text{NO}_2$ ratio can be readily updated by dividing the emissions from this study by 1.32 and then multiplying any city- and/or season-specific value.

3.2 NO_x emission estimation

The derivation of emissions (E) from satellite-observed column amounts (Ω) is based on the principle of mass conservation as in the following:

$$\langle E \rangle = \langle \mathbf{u} \cdot (\nabla \Omega) \rangle + X \langle \Omega \mathbf{u}_0 \cdot (\nabla z_0) \rangle + \frac{\langle \Omega \rangle}{\tau}. \quad (1)$$

Here $\langle \rangle$ is the spatiotemporal averaging operator already implemented in the physical oversampling framework (Sun et al., 2018; Sun, 2022b), z_0 is the surface altitude from level 2 files, and \mathbf{u} and \mathbf{u}_0 are horizontal wind vectors in the planetary boundary layer (PBL) and near the surface, respectively, represented by 100-m and 10-m winds sampled from ERA5. X and τ represent the inverse scale height and vertically integrated chemical lifetime and can be inferred as linear regression



coefficients using monthly or further aggregated images. The full derivation of Eq. 1 can be found in Sun (2022a). Despite
 120 the similarity between Eq. 1 and its counterpart in the flux divergence literature (Beirle et al., 2019, 2021; Liu et al., 2021;
 Dix et al., 2022; de Foy and Schauer, 2022; Rey-Pommier et al., 2022; Veeffkind et al., 2023), Eq. 1 accounts for the impacts
 from the horizontal divergence of wind and topography to the estimated emission, both of which are not included in the flux
 divergence equation and scale linearly with column amounts, Ω . The first and second terms on the right-hand side of Eq. 1
 are based on the directional derivatives of the column amount (Ω) and surface altitude (z_0) with respect to the horizontal wind
 125 vectors (\mathbf{u} and \mathbf{u}_0). Therefore, we refer to emission estimation using this equation as the directional derivative approach.

The inverse scale height (X) and chemical lifetime (τ) remain as key unknowns and can be estimated from observational
 data. At locations where emission $\langle E \rangle$ is small, Eq. 1 can be rewritten as a multi-linear regression model by neglecting the
 emission term:

$$\langle \mathbf{u} \cdot (\nabla \Omega) \rangle = \beta_0 + \beta_1 \langle \Omega \mathbf{u}_0 \cdot (\nabla z_0) \rangle + \beta_2 \langle \Omega \rangle + \varepsilon. \quad (2)$$

130 Here β_0 and ε represent the offset and random error in the predicted variable (i.e., $\langle \mathbf{u} \cdot (\nabla \Omega) \rangle$) that cannot be explained by the
 linear combination of predictors (i.e., $\langle \Omega \mathbf{u}_0 \cdot (\nabla z_0) \rangle$ and $\langle \Omega \rangle$). β_1 is an estimate of the negative inverse of scale height, and β_2
 is an estimate of the negative of the first order rate constant or equivalently the inverse of chemical lifetime.

For each region, terms $\langle \mathbf{u} \cdot (\nabla \Omega) \rangle$, $\langle \Omega \mathbf{u}_0 \cdot (\nabla z_0) \rangle$, and $\langle \Omega \rangle$ are calculated and saved at 0.04° grid size and monthly res-
 olution. Then regressions as in Eq. 2 are conducted at a subset of grid cells for each subregion. We first focus on fitting
 135 β_1 over relatively rough terrains where NO_x emissions are generally much smaller than the observational error and hence
 negligible. This fit can be done at a relatively high time resolution (monthly) given the high signal-to-noise ratio. β_2
 is also included in this fitting, although the results are usually very noisy. This first round of fitting is limited to grid cells
 with $0.001 \text{ m s}^{-1} < \langle \mathbf{u}_0 \cdot (\nabla z_0) \rangle < 0.1 \text{ m s}^{-1}$, which represent moderately rough terrains that are abundant in all regions and
 subregions. In the second round, the monthly fitted β_1 in the previous round is fixed, and only β_2 is fitted in flat terrains
 140 ($\langle \mathbf{u}_0 \cdot (\nabla z_0) \rangle < 0.001 \text{ m s}^{-1}$) that are free of strong NO_x emission sources ($\langle \mathbf{u} \cdot (\nabla \Omega) \rangle < 1 \text{ nmol m}^{-2} \text{ s}^{-1}$) and meanwhile
 characterized by moderate NO_2 column amount ($\langle \Omega \rangle > 2.5 \times 10^{-5} \text{ mol m}^{-2}$). To address the issue of low signal-to-noise ratio,
 this second round fitting is conducted over climatological months. Namely, the same months for all years are aggregated before
 fitting. This two-round fitting procedure is generally consistent with the study over the contiguous USA by Sun (2022a). The
 main improvements here are that the fittings are conducted in smaller subregions and that the seasonal variations of lifetime
 145 are resolved.

Figure 1 shows the fitted NO_x scale heights (top) and chemical lifetimes (bottom) for each month, although the monthly
 lifetime is from climatology and hence the same for different years. We caution here that the resultant scale heights and lifetimes
 are fundamentally fitting parameters in a multi-linear regression model (Eq. 2) that minimize the impacts of topography and
 chemical loss on the estimated emission. Qualitatively, the seasonality of NO_x scale heights is consistent with higher PBL
 150 height in the summer than winter, and the fact that the southern subregion in North America shows significantly higher scale
 height than other subregions is consistent with the high PBL height over southwest USA and northern Mexico (Ding et al.,
 2021; Ayazpour et al., 2023). The low scale heights in East Asia may be explained by higher levels of pollution and thus



more NO_x distributed near the surface. The chemical lifetimes in all subregions span a broad range and are generally longer in winter than summer. The seasonal variation of lifetime in the southern subregion in East Asia is comparable to previous lifetime estimates (Mijling and Van Der A, 2012; Shah et al., 2020), whereas the lifetimes in other subregions are significantly higher. The most plausible explanation is that the lifetime as in Eqs 1 and 2 is integrated through the vertical column, so the free tropospheric NO_x contributes more in relatively clean regions. The lifetime results in the northern subregion in Europe become unreliable in winter due to low data coverage as shown by occasional negative values. We keep using these results without modification as they do not have significant impacts on the estimated emissions.




Figure 1. (Top) Monthly scale heights fitted using Eq. 2. NA, EU, and AS represent the regions of North America, Europe, and East Asia. S and N after the underscore denote the southern and northern subregions in each region. (Bottom) Chemical lifetimes fitted for these six subregions. Each monthly data point is from the climatology so all years have the same seasonality.

160 Once the monthly NO_x emissions E are obtained using the monthly fitted X and τ , NO_x emissions from each selected city are calculated by averaging NO_x emission grid cells under the geographical coverage of the city (see Sect. 2.2 for the determination of city coverage). The NO_x emission grid cells are weighted by the fraction of urban area coverage during the averaging.

3.3 Algorithms for city clustering based on annual emissions

165 The monthly NO_x emissions from each city (9 cities per subregion and 54 cities in total) are aggregated annually for cluster analysis in Sect. 4.4. The normalized emissions in 2019–2022 by the 4-year mean are considered the feature for each city and clustered using the k-means clustering algorithm (Likas et al., 2003). K-means algorithm partitions a set of data points in n -dimensional space ($n = 4$ here, corresponding to annual emissions in 2019–2022) into k clusters, where each data point



belongs to the cluster with the nearest mean. The mean or centroid of each cluster is representative of the general pattern of
170 data points in the cluster. A total number of 4 clusters are used by locating the elbow point of the sum of squared errors as a
function of the number of clusters. Additionally, we reduce the feature dimension of 4 to 2 using principal component analysis,
such that each city can be projected to a 2-dimensional scatter plot as shown by Figure 17.

4 Results

This section dives into the regions in North America, Europe, and East Asia each with two subregions in the north and south
175 and 9 selected cities in each subregion. Section 4.4 synthesizes the annual NO_x emissions from all selected cities through the
cluster analysis.

4.1 North America

Figure 2 overviews the region in North America, its two subregions delineated by red rectangles, and selected cities with loca-
tions indicated by black arrows. The spatial distribution of NO_x emissions shown by the central map are estimated following
180 Sect. 3.2 in 2019–2022, except that the scale height and chemical lifetime are fitted using the entire region instead of a specific
subregion. The southern subregion covers the southwest USA and northern Mexico, and the northern subregion covers the mid-
west and northeast USA and part of Canada. The annual NO_x emissions in 2019–2022 averaged spatially over each selected
city are illustrated as pie charts around the edges of the plot. The sizes of the pies scale with the average city emissions over
the 4 years. One may compare the size of slices for 2020–2022 to 2019 as an indication of post-COVID-19 emission changes.
185 Note that the emissions in 2021 and 2022 are generally higher than 2019 for the selected cities in the southern subregion. The
slices of 2019 are not larger than a quarter of the pie in all the cities in the southern subregion. Namely, despite the impacts
of COVID-19, the post-COVID annual emissions in these cities are not lower than the pre-COVID year of 2019. The cities in
Mexico (Tijuana, Juarez, and Chihuahua) show faster growth of NO_x emissions year-to-year and stronger emissions compared
with neighboring cities in the USA. The northern subregion is quite different, in that the NO_x emissions in 2019 are all higher
190 than the 4-year average, i.e., the 2019 slices are larger than a quarter of the pies. This indicates decreased emissions after
COVID-19 that may be attributed to the direct and indirect impacts of pandemic-control measures.

Figures 3 and 4 show the monthly NO_x emissions that are averaged to obtain the annual emissions for cities in the southern
and northern subregions in North America (see Figure 2). In these plots, each city corresponds to one panel, and the panels
are ordered by population as provided by the city list (same for all the following 9-panel, 1-panel-per-city figures). For each
195 panel, the top subpanel shows the absolute monthly NO_x emissions, and the bottom subpanel shows the relative emissions to
the same months in 2019. For both subpanels, the values in 2019 are also repeated in the same months for the other years (2018
and 2020–2022) as a baseline for comparison. Higher monthly emissions relative to the same months in 2019 are indicated
by red shade, and blue shade is used otherwise. We remove the monthly data with city-wide average level 2 data coverage
lower than 2, i.e., the entire city area has to be on average covered at least twice in the month by TROPOMI observations. This
200 threshold is determined using all selected cities in this study, and monthly emissions with coverage lower than this value tend




Figure 2. Geographical locations of the 18 cities, 9 in each of the two subregions, in the region of North America. The subregions are outlined by dashed red rectangles. The annual NO_x emissions in 2019–2022 for each city are displayed as pie charts. The emission values for each year are labeled on or near the corresponding slice of the pie in $\text{nmol m}^{-2} \text{s}^{-1}$. The 4-year average emissions are labeled beside the city names. The central map shows 4-year average NO_x emissions throughout the region. The grid is coarsened from the native size of 0.04° to 0.12° for visualization purposes.

to be unreliable. To make consistent interannual comparison, if one month is removed for a particular year, the same months for all other years are also removed for a given city. This results in loss of some winter months in high latitude cities due to high solar zenith angle and snow coverage and wet-season months in some cities due to frequent cloud coverage. Although the monthly emissions are subject to significant variability, NO_x emissions for most cities dropped in early 2020 relative to the same months in 2019, coincident with the initial wave of COVID-19. The relative decrease is more significant in the northern subregion than in the southern subregion. The emissions, relative to 2019, diverge more in 2021 and 2022 among these cities. Some cities in the northeast USA show comparable or even more emission reduction in 2022 than in 2020 (e.g., Chicago and Philadelphia), whereas strong growth can be identified in south and southwest USA (e.g., Dallas, Houston, San Diego, and Phoenix) and in Mexico.




Figure 3. The red dots and lines show monthly NO_x emissions from cities in the southern subregion in the region of North America. For each panel, the top subpanel shows the absolute emissions, and the bottom subpanel shows the relative emissions to the corresponding months in 2019. The blue dashed lines show the 2019 values repeated in the same months for the other years (2018 and 2020–2022). Red and blue shades indicate higher and lower monthly emissions relative to the months in 2019.

210 Strong seasonal variations with higher emissions in winter months are observed in some cities (e.g., all cities in the northern subregion, Dallas, Houston, San Diego, and Juarez). One may hypothesize that these observed seasonal variations might be caused by seasonally varying artifacts, such as retrieval biases, vertical sensitivity of the retrieval at the surface, and the uncertainties in the wind vectors. However, we note that no clear seasonality can be identified in Tijuana, whereas the adjacent San Diego shows a much more prominent seasonal pattern. This is inconsistent with the aforementioned hypotheses, because



215 these artifacts should have impacted the NO_2 retrieval and NO_x emission estimation algorithms similarly at such a close distance. Moreover, similar seasonalities are not so common in the regions of Europe and East Asia to be shown in the following sections. Further validation of the emission values and seasonality will be the subject of future studies.




Figure 4. Absolute (top subpanels) and relative (bottom subpanels) monthly NO_x emissions from cities in the northern subregion in the region of North America. This figure is similar to Figure 3.

The geographical urban area coverage and spatial distribution of NO_x emissions for each city are shown by Figure 5 (southern subregion) and 6 (northern subregion). The averaged NO_x emissions in 2019–2022 are displayed in native grid size of 0.04° as a colored map. The city extent is illustrated as a mask where non-urban area is in black with 95% transparency, resulting in a gray hue. The city-covered area is fully transparent. The urban sprawl is significant in the USA (Barrington-Leigh and

220



Millard-Ball, 2015), as the city areas in the USA are generally larger than similar sized cities in other countries. Emission hot spots are often collocated with the downtown areas, but industrial areas and sea ports show higher emissions, e.g., the Port of Long Beach in Los Angeles (Figure 5a) and the Houston ship channel (Figure 5c). The emissions within the city window of Washington, DC are actually dominated by Baltimore to the northeast (Figure 6e). Emissions in Tijuana and Juarez are clearly higher than the adjacent American cities San Diego and El Paso, presumably due to different emission regulations.




Figure 5. Maps of NO_x emissions and urban area coverage for the 9 selected cities in the southern subregion in North America.




Figure 6. Maps of NO_x emissions and urban area coverage for the 9 selected cities in the northern subregion in North America.

4.2 Europe

Figure 7 is a similar overview for the region of Europe, where the southern and northern subregions are delineated by red dashed rectangles. The NO_x emissions from shipping lanes over the Atlantic ocean and the Mediterranean sea are prominent. The annual NO_x emissions in 2019–2022 are shown similarly as pie charts for each city. Note that the southern subregion includes an African city, Algiers in Algeria, due to the rectangular shape of the subregion. Algiers also stands out among other cities in this region in that its 2019 emission is lower than the 4-year average; the 2019 slice is smaller than a quarter of the pie, and its 2020 emission is higher than 2019. All other cities show emission reductions in 2020 relative to 2019, which often extend to the following years. Two large cities in developing countries, Algiers and Istanbul, are characterized by larger emissions overall and stronger rebound of emissions after 2020.




Figure 7. Geographical locations of the 18 cities in the region of Europe. The subregions are outlined by dashed red rectangles. The annual NO_x emissions in 2019–2022 in each city are displayed as pie charts. The emission values each year are labeled on or near the corresponding slice of the pie in $\text{nmol m}^{-2} \text{s}^{-1}$. The 4-year average emissions are labeled beside the city names. The central map shows 4-year average NO_x emissions throughout the region. The grid is coarsened from the native size of 0.04° to 0.12° for visualization purposes.

Figures 8 and 9 show the monthly NO_x emissions that are averaged to obtain the annual emissions for cities in the southern and northern subregions in the region of Europe (see Figure 7). Compared with cities in North America, cities in the European region generally had much larger emission decreases during the initial COVID-19 wave as indicated by larger blue shaded areas. In some cases, most noticeably Madrid, Lisbon, and London, the emission reductions extend almost throughout 2020–2022. In contrast, the emissions quickly rebounded after the initial impact in some central and eastern Europe cities, such as Bucharest, Warsaw, and Prague, as well as in Istanbul and Algiers as mentioned earlier. Some cities in the northern subregion are also subject to significant loss of winter month coverage due to high solar zenith angle and cloud coverage.

The geographical urban area coverage and spatial distribution of NO_x emissions for each city are shown by Figure 10 (southern subregion) and 11 (northern subregion), similar to city maps in the North American region (Figures 5 and 6). Unlike




Figure 8. Absolute (top subpanels) and relative (bottom subpanels) monthly NO_x emissions from cities in the southern subregion in the region of Europe. This figure is similar to Figure 3.




Figure 9. Absolute (top subpanels) and relative (bottom subpanels) monthly NO_x emissions from cities in the northern subregion in the region of Europe. This figure is similar to Figure 3.



245 the American cities that tend to sprawl into a large continuum, the selected European cities tend to be more concentrated with smaller satellite cities and towns scattered in the surrounding area. The most prominent emission features are generally located at the city centers, with Rotterdam as an exception where most observed emissions concentrate along the port of Rotterdam, the world's largest seaport outside of East Asia (Figure 11i).




Figure 10. Maps of NO_x emissions and urban area coverage for the 9 selected cities in the southern subregion in Europe.




Figure 11. Maps of NO_x emissions and urban area coverage for the 9 selected cities in the northern subregion in Europe.

4.3 East Asia

250 Figure 12 is a similar overview for the region of East Asia, where the southern and northern subregions are delineated by red dashed rectangles. Note that the overall emission background and emissions from individual cities are significantly higher than the regions of North America and Europe, as indicated by the enhanced scale of the color map and pie chart sizes. Selected cities in this region ubiquitously show lower mean annual emissions after COVID-19, i.e., the 2019 pie slices are all larger than a quarter. Unlike the regions of North America and Europe, the selected cities here all show lower emissions in 2022 than the
255 4-year average, and in many cases, the 2022 emissions are lower than 2021 and 2020. Out of 18 selected in this region, 16 are in China, which underwent widespread and stringent measures in 2022 to control the spread of the Omicron variant.




Figure 12. Geographical locations of the 18 cities, 9 in each of the two subregions, in the region of East Asia. The subregions are outlined by dashed red rectangles. The annual NO_x emissions in 2019–2022 in each city are displayed as pie charts. The emission values each year are labeled on or near the corresponding pie slice in $\text{nmol m}^{-2} \text{s}^{-1}$. The 4-year average emissions are labeled beside the city names. The central map shows 4-year average NO_x emissions throughout the region. The grid is coarsened from the native size of 0.04° to 0.08° for visualization purposes.




Figure 13. Absolute (top subpanels) and relative (bottom subpanels) monthly NO_x emissions from cities in the southern subregion in the region of East Asia. This figure is similar to Figure 3.




Figure 14. Absolute (top subpanels) and relative (bottom subpanels) monthly NO_x emissions from cities in the northern subregion in the region of East Asia. This figure is similar to Figure 3.



The temporal evolution of NO_x emissions and their relative changes to 2019 are shown in more detail in the monthly plotted Figures 13 and 14, with one of the strongest examples of this shown in the megacity of Shanghai in Figure 13a. Shanghai is one of the largest cities studied here with higher NO_x emissions than all other selected cities except Seoul. The data coverage for Shanghai is also complete without any missing months. The well-documented NO_x emission reductions during the spring festival in January and/or February are evident in 2019–2021, although the 2020 spring festival coincided with the initial control measures at the beginning of the COVID-19 outbreak (Liu et al., 2020; Huang and Sun, 2020). Emissions were back to 2019 levels during the second half of 2021. March 2022 marked the start of an unprecedented lockdown in Shanghai in response to the spread of the Omicron variant, and the resultant NO_x emission reductions overshadow those of early 2020. Emission levels largely recovered in August–October 2022 before plunging again due to a nation-wide spread of the Omicron variant, which ultimately led to a termination of most control measures in China in December 2022. The full effect of this policy change on NO_x emissions is not yet clear from the current data.

Figure 13b shows the monthly NO_x emissions in Wuhan, where COVID-19 first drew public attention in January 2020. Data in January and February of all years are not included due to insufficient TROPOMI coverage, resulting in missing peak lockdown months. Therefore, the relative reduction in 2020 is likely underestimated. Another noteworthy feature in Wuhan is that the October emissions in 2018 and 2020–2022 are all higher than October 2019. One likely cause is the emission reduction measures conducted to ensure good air quality during the 7th Military World Games held in Wuhan in October 2019 (Zhang et al., 2022). As a result, October 2019 emission in Wuhan was likely lower than the business-as-usual condition, leading to spurious enhancements in October of all other years. The other selected cities similarly show emission reductions in early 2020 during the onset of the pandemic and more extensive reductions in 2022 due to direct and indirect impacts of the Omicron variant spread. Some cities show consecutive months of recovery or increase of emissions in between, e.g., Hangzhou, Ningbo, and Jinan, for the summer and fall of 2020.

Similar to the regions of North America and Europe, the geographical urban area coverage and spatial distribution of NO_x emissions for each city in the region of East Asia are shown by Figure 15 (southern subregion) and 16 (northern subregion). Unlike most cities in North America and Europe, the strongest emissions in the selected Asian cities are often not located at the city centers, but in industrial areas or sea ports. Here we try to identify the most prominent emission hot spots on the city maps. The strongest emissions in Shanghai occur in the highly industrialized Baoshan and Pudong districts along the Yangtze River shore (Figure 15a). For Wuhan, the strongest emissions occur at Qingshan district where Wuhan Iron & Steel is located (Figure 15b). The emission hot spot in the southwest of Nanjing (Figure 15d) is the city of Maanshan, home of Maanshan Iron & Steel. The emission hot spot in the northeast of Wuxi (Figure 15g) is part of Zhanjiangang, a county-level city (in contrast to prefecture-level cities) under the administration of Suzhou. The emission hot spot to the east of Ningbo city (Figure 15i) appears to be part of the port of Ningbo-Zhoushan, the world's largest cargo handling port.

The city window around Seoul (Figure 16a) includes most of the Seoul metropolitan area, which was mapped for high-resolution NO_2 column amounts during the Korea–United States Air Quality (KORUS-AQ) campaign (Choo et al., 2023). The west-east extended hot spot in the west of the city window is associated with the Incheon industrial complex, while the more south-north extended hot spot at the center of the city window is located over the city of Seoul. The emission hot spot in the



south is located near Suwon, which is also industrialized. The emission hot spot to the southwest of the Tianjin city (Figure 16c) collocates with the port of Tianjin, the largest port in Northern China and the main maritime gateway to Beijing, and the adjacent industrial area in the Binhai New Area. The emission hot spot to the northeast of Tangshan city (Figure 16g) appears to be Qian'an, a county-level city under the administration of Tangshan, and is the location of the Yanshan Iron & Steel. The emission hot spot to the southwest of Qingdao city (Figure 16h) appears to be the Port of Qindao. The emission hot spot to the northeast of the Busan city (Figure 16i) appears to be the port of Ulsan, the largest industrial port in South Korea.




Figure 15. Maps of NO_x emissions and urban area coverage for the 9 selected cities in the southern subregion in East Asia.




Figure 16. Maps of NO_x emissions and urban area coverage for the 9 selected cities in the northern subregion in East Asia.

4.4 Clustering of city emissions

The monthly NO_x emissions calculated for each of the 54 selected cities contain a large amount of information that is challenging to digest. These monthly values are often subject to low signal-to-noise ratio, especially for the cold season months in high-latitude cities. As such, we aggregate to annual emissions for the years of 2019–2022 to obtain more insights into how different cities' emissions responded after the onset of COVID-19. For each city, the same months are included for all these years to enhance interannual consistency. The resultant annual emissions are already shown in Figures 2, 7, and 12. These annual emissions (4 values for each city) are normalized by the 4-year mean and then grouped into 4 clusters using the k-means algorithm. The normalized annual emission for each city corresponds to a point in the 4-dimensional space. To visualize the



clustering results, we reduce the dimension of the normalized annual emissions by calculating 2 principal components, which effectively projects the data to a 2-dimensional subspace.




Figure 17. (a) Clustering results using 54 cities selected in this study. The normalized annual emissions in 2019–2022 for each city are projected to 2-dimensional space using principal component analysis. The locations of Tianjin and Milan, defined as the two principal components, are nudged downward and right, respectively, to enhance visualization quality. (b–e) The bars indicate the cluster center, and error bars indicate the interquartile range within the cluster. The values are all relative to 2019.

Figure 17a shows the distributions of cities in the projected 2-dimensional subspace, where each city is a point marked differently according to the cluster it belongs to. The annual emissions for cities in the same cluster are shown as relative values to the 2019 emission in Figure 17b–e, where the bars indicate the cluster average, and error bars indicate the interquartile range within the cluster. All cities in the North American region, with Chicago as the only exception, are included in cluster 1 (Figure 17b), which is characterized by the least emission reduction in 2020 (−7% relative to 2019), with emissions recovering to the 2019 level in 2021, and increased emissions in 2022 (+5% relative to 2019). Bucharest, Warsaw, Prague, Algiers, and Istanbul



in the European region are also included in this cluster. The overall characteristic of cluster 1 is a minor reduction in 2020
315 relative to 2019 and steady increase afterwards.

Cluster 2 (Figure 17c) features the most significant emission reduction in 2020 (−31% relative to 2019), with a moderate
rebound in 2021 (−15%), and drop again in 2022 (−21%). Cities in this cluster are all in the European region, except Shanghai.
Cluster 3 (Figure 17d) differs from cluster 2 in that the emission reduction in 2020 is not as large (−8%), while the reduction
in 2022 is very significant (−25%), especially in comparison with its value in 2021 (−8.5%). Cluster 3 mainly includes Cities
320 in China with a few exceptions in Europe (Lisbon, Hamburg, and Berlin) and North America (Chicago). This is consistent with
the general evolution pattern of COVID-19 in China: quick recovery of emissions in later 2020 due to effective COVID-19
control measures, sporadic lockdowns in 2021, and much larger scale lockdowns in 2022.

Cluster 4 (Figure 17e) is characterized by the largest and sustained decrease of emissions from 2019 to 2022. The average
emission reduction in 2022 in this cluster is −37% relative to 2019, the lowest for all years in all clusters. The reduction in
325 2021 is −35%, also substantially more than all other clusters. Cities in this cluster are located in northern China and South
Korea with London as the only exception. Additionally, Tangshan, Seoul, and Busan in cluster 4 have large contributions
from industrial NO_x sources (see Figure 16), suggesting influences from economic factors besides direct COVID-19-control
impacts.

5 Conclusions and discussion

330 We apply the directional derivative approach developed by Sun (2022a) to estimate NO_x emissions in three northern hemisphere
regions; North America, Europe, and East Asia. For each region, the NO_x scale heights and chemical lifetimes, which are
necessary in the emission calculation, are estimated separately in two subregions. We focus on emissions from 9 selected cities
per subregion and present monthly averages and 4-year averaged emission maps at 0.04° grid size for a total number of 54
cities. The NO_x emissions maps reveal unprecedented levels of detail for a large and diverse collection of cities. NO_x emission
335 hot spots are consistently found at large city cores, while some cities feature significantly higher emissions than others, most
notably at the USA-Mexico border (San Diego vs. Tijuana and El Paso vs. Juarez). The spatial windows of some cities envelops
much more prominent emission hot spots than the city cores, which generally correspond to large seaports and industrial areas.
The average emissions in 2019–2022 are generally larger in East Asian cities, as 13 out of 18 cities in this region are higher
than 10 nmol m^{−2} s^{−1}. In contrast, only one city in the North American region (Tijuana) and two cities in the European region
340 (Algiers and Istanbul) are higher than this value.

With respect to the temporal variation of NO_x emissions, we choose the year of 2019 as the pre-COVID-19 baseline year, so
the relative emission changes in 2020–2022 to 2019 indicate the post-COVID-19 perturbations to each city. The initial impact
during the first outbreak of COVID-19 in early 2020 can be found in most cities, but their paths diverge afterwards. We average
the monthly emissions for each city to annual mean emissions in 2019–2022 and group the normalized annual mean emissions
345 into 4 clusters. All but one city in the North American region are grouped in cluster 1, which is characterized by the smallest
emission reduction in 2020 (−7%) and steady increase afterwards, resulting in a +5% increase in 2022. Limited representations



of Latin America (Tijuana, Juarez, and Chihuahua), Africa (Algiers), and Middle East (Istanbul) are all located in cluster 1. Future studies might be meaningful to test if the emission changing pattern of cluster 1 is common in these regions. The other clusters (2–4) feature much larger emission reductions than cluster 1 and differ by how these reductions are distributed in
350 2020–2022. The European cities are generally in cluster 2 with the largest impact in 2020 (–31%), whereas the East Asian cities are generally in cluster 3 and 4 with the largest impacts in 2022 (–25% and –37%).

In this study we fit scale heights at monthly resolution and fit chemical lifetimes for each climatological month to strike a balance between the quality of the fitting results and temporal resolution. However, we assume spatially homogeneous scale heights and chemical lifetimes within each subregion. Future investigations might be helpful to achieve higher spatial
355 granularity, especially considering the nonlinear effects of NO_x chemistry (Valin et al., 2013; Laughner and Cohen, 2019). We choose a constant NO_x : NO_2 ratio, given the emphasis of this study on relative emission changes and timeliness of emission estimation. An improved understanding of global NO_x : NO_2 ratio over the atmospheric columns through which satellite sensors integrate will likely enhance the quality of estimated NO_x emissions.

This work presents observation-based NO_x emission estimations over large areas (covering three major continents), with
360 fine spatial resolution (0.04° , resolving intracity emission variations), high temporal resolution (monthly), and timely results (until 31 January 2023). The current work flow requires only TROPOMI level 2 data and the ERA5 reanalysis, both publicly available with global coverage, and the open-source Python algorithm (Sun, 2022b). It is our hope that this tool will benefit future studies over more regions in the world and using additional remote sensing instruments.

Code availability. Code relevant to this paper can be found in Sun (2022b).

365 *Data availability.* TROPOMI NO_2 PAL data are available at <https://data-portal.s5p-pal.com/products/no2.html>. TROPOMI NO_2 offline data are available at <https://doi.org/10.5270/S5P-s4ljg54>. The ERA5 data are available at <https://doi.org/10.24381/cds.adbb2d47>.

Author contributions. CRL and KS edited the manuscript. KS curated the algorithm and datasets.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This research has been supported by the NASA Earth Science Division Rapid Response and Novel Research in Earth
370 Science program (RRNES, Award 80NSSC20K1295) and Atmospheric Composition: Modeling and Analysis (ACMAP, Award 80NSSC19K0988). We thank Guanyu Huang, Hyeong-Ahn Kwon, Caroline Nowlan, and Heesung Chong for helping identifying emission hot spots.



References

- Ayazpour, Z., Tao, S., Li, D., Scarino, A. J., Kuehn, R. E., and Sun, K.: Estimates of the spatially complete, observational-data-driven planetary boundary layer height over the contiguous United States, *Atmospheric Measurement Techniques*, 16, 563–580, <https://doi.org/10.5194/amt-16-563-2023>, 2023.
- 375 Barrington-Leigh, C. and Millard-Ball, A.: A century of sprawl in the United States, *Proceedings of the National Academy of Sciences*, 112, 8244–8249, <https://doi.org/10.1073/pnas.1504033112>, 2015.
- Beirle, S., Boersma, K. F., Platt, U., Lawrence, M. G., and Wagner, T.: Megacity emissions and lifetimes of nitrogen oxides probed from space, *Science*, 333, 1737–1739, 2011.
- 380 Beirle, S., Borger, C., Dörner, S., Li, A., Hu, Z., Liu, F., Wang, Y., and Wagner, T.: Pinpointing nitrogen oxide emissions from space, *Science Advances*, 5, eaax9800, <https://doi.org/10.1126/sciadv.aax9800>, 2019.
- Beirle, S., Borger, C., Dörner, S., Eskes, H., Kumar, V., de Laat, A., and Wagner, T.: Catalog of NO_x emissions from point sources as derived from the divergence of the NO₂ flux for TROPOMI, *Earth System Science Data*, 13, 2995–3012, <https://doi.org/10.5194/essd-13-2995-2021>, 2021.
- 385 Chen, Y.-C., Chou, C. C. K., Liu, C.-Y., Chi, S.-Y., and Chuang, M.-T.: Evaluation of the nitrogen oxide emission inventory with TROPOMI observations, *Atmospheric Environment*, p. 119639, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2023.119639>, 2023.
- Choo, G.-H., Lee, K., Hong, H., Jeong, U., Choi, W., and Janz, S. J.: Highly resolved mapping of NO₂ vertical column densities from GeoTASO measurements over a megacity and industrial area during the KORUS-AQ campaign, *Atmospheric Measurement Techniques*, 16, 625–644, <https://doi.org/10.5194/amt-16-625-2023>, 2023.
- 390 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth System Science Data*, 10, 1987–2013, <https://doi.org/10.5194/essd-10-1987-2018>, 2018.
- Dammers, E., Tokaya, J., Mielke, C., Hausmann, K., Griffin, D., McLinden, C., Eskes, H., and Timmermans, R.: Can TROPOMI-NO₂ satellite data be used to track the drop and resurgence of NO_x emissions between 2019–2021 using the multi-source plume method (MSPM)?, *Geoscientific Model Development Discussions*, 2022, 1–33, <https://doi.org/10.5194/gmd-2022-292>, 2022.
- 395 de Foy, B. and Schauer, J. J.: An improved understanding of NO_x emissions in South Asian megacities using TROPOMI NO₂ retrievals, *Environmental Research Letters*, 17, 24006, <https://doi.org/10.1088/1748-9326/ac48b4>, 2022.
- Ding, F., Iredell, L., Theobald, M., Wei, J., and Meyer, D.: PBL Height From AIRS, GPS RO, and MERRA-2 Products in NASA GES DISC and Their 10-Year Seasonal Mean Intercomparison, *Earth and Space Science*, 8, e2021EA001859, <https://doi.org/https://doi.org/10.1029/2021EA001859>, 2021.
- 400 Ding, J., van der A, R. J., Eskes, H. J., Mijling, B., Stavrou, T., van Geffen, J. H. G. M., and Veefkind, J. P.: NO_x Emissions Reduction and Rebound in China Due to the COVID-19 Crisis, *Geophysical Research Letters*, 47, e2020GL089912, <https://doi.org/https://doi.org/10.1029/2020GL089912>, 2020.
- Dix, B., Francoeur, C., Li, M., Serrano-Calvo, R., Levelt, P. F., Veefkind, J. P., McDonald, B. C., and de Gouw, J.: Quantifying NO_x Emissions from U.S. Oil and Gas Production Regions Using TROPOMI NO₂, *ACS Earth and Space Chemistry*, 6, 403–414, <https://doi.org/10.1021/acsearthspacechem.1c00387>, 2022.



- Eskes, H., Van Geffen, J., Boersma, F., Eichmann, K.-U., Apituley, A., Pedernana, M., Sneep, M., Veefkind, J., and Loyola, D.: Sentinel-5 precursor/TROPOMI Level 2 Product User Manual Nitrogen dioxide, <https://sentinel.esa.int/documents/247904/2474726/Sentinel-5P-Level-2-Product-User-Manual-Nitrogen-Dioxide.pdf>, 2022.
- 410 Gkatzelis, G. I., Gilman, J. B., Brown, S. S., Eskes, H., Gomes, A. R., Lange, A. C., McDonald, B. C., Peischl, J., Petzold, A., Thompson, C. R., and Kiendler-Scharr, A.: The global impacts of COVID-19 lockdowns on urban air pollution: A critical review and recommendations, *Elementa: Science of the Anthropocene*, 9, 176, <https://doi.org/10.1525/elementa.2021.00176>, 2021.
- Godłowska, J., Hajto, M. J., Lapeta, B., and Kaszowski, K.: The attempt to estimate annual variability of NO_x emission in Poland using Sentinel-5P/TROPOMI data, *Atmospheric Environment*, 294, 119 482, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2022.119482>,
415 2023.
- Goldberg, D. L., Lu, Z., Streets, D. G., de Foy, B., Griffin, D., McLinden, C. A., Lamsal, L. N., Krotkov, N. A., and Eskes, H.: Enhanced Capabilities of TROPOMI NO₂: Estimating NO_x from North American Cities and Power Plants, *Environmental Science & Technology*, 53, 12 594–12 601, <https://doi.org/10.1021/acs.est.9b04488>, 2019.
- Goldberg, D. L., Harkey, M., de Foy, B., Judd, L., Johnson, J., Yarwood, G., and Holloway, T.: Evaluating NO_x emissions and their effect
420 on O₃ production in Texas using TROPOMI NO₂ and HCHO, *Atmos. Chem. Phys.*, 22, 10 875–10 900, <https://doi.org/10.5194/acp-22-10875-2022>, 2022.
- Hernández, J.: World cities database, <https://doi.org/10.34740/KAGGLE/DSV/3360046>, 2022.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee,
425 D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, n/a, <https://doi.org/10.1002/qj.3803>, 2020.
- Huang, G. and Sun, K.: Non-negligible impacts of clean air regulations on the reduction of tropospheric NO₂ over East China during the COVID-19 pandemic observed by OMI and TROPOMI, *Science of The Total Environment*, 745, 141 023,
430 <https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.141023>, 2020.
- Kang, M., Zhang, J., Cheng, Z., Guo, S., Su, F., Hu, J., Zhang, H., and Ying, Q.: Assessment of Sectoral NO_x Emission Reductions During COVID-19 Lockdown Using Combined Satellite and Surface Observations and Source-Oriented Model Simulations, *Geophysical Research Letters*, 49, e2021GL095 339, <https://doi.org/https://doi.org/10.1029/2021GL095339>, 2022.
- Lange, K., Richter, A., and Burrows, J. P.: Variability of nitrogen oxide emission fluxes and lifetimes estimated from Sentinel-5P TROPOMI
435 observations, *Atmos. Chem. Phys.*, 22, 2745–2767, <https://doi.org/10.5194/acp-22-2745-2022>, 2022.
- Laughner, J. L. and Cohen, R. C.: Direct observation of changing NO_x lifetime in North American cities, *Science*, 366, 723 LP – 727, <https://doi.org/10.1126/science.aax6832>, 2019.
- Levelt, P. F., Stein Zweers, D. C., Aben, I., Bauwens, M., Borsdorff, T., De Smedt, I., Eskes, H. J., Lerot, C., Loyola, D. G., Romahn, F., Stavrou, T., Theys, N., Van Roozendaal, M., Veefkind, J. P., and Verhoelst, T.: Air quality impacts of COVID-19 lockdown measures detected from space using high spatial resolution observations of multiple trace gases from Sentinel-5P/TROPOMI, *Atmospheric Chemistry and Physics*, 22, 10 319–10 351, <https://doi.org/10.5194/acp-22-10319-2022>, 2022.
- 440 Likas, A., Vlassis, N., and Verbeek, J. J.: The global k-means clustering algorithm, *Pattern recognition*, 36, 451–461, 2003.



- 445 Liu, F., Page, A., Strode, S. A., Yoshida, Y., Choi, S., Zheng, B., Lamsal, L. N., Li, C., Krotkov, N. A., Eskes, H., van der A, R., Veefkind, P., Levelt, P. F., Hauser, O. P., and Joiner, J.: Abrupt decline in tropospheric nitrogen dioxide over China after the outbreak of COVID-19, *Science Advances*, 6, eabc2992, <https://doi.org/10.1126/sciadv.abc2992>, 2020.
- Liu, M., van der A, R., van Weele, M., Eskes, H., Lu, X., Veefkind, P., de Laat, J., Kong, H., Wang, J., Sun, J., Ding, J., Zhao, Y., and Weng, H.: A New Divergence Method to Quantify Methane Emissions Using Observations of Sentinel-5P TROPOMI, *Geophysical Research Letters*, 48, e2021GL094151, <https://doi.org/10.1029/2021GL094151>, 2021.
- 450 Lorente, A., Boersma, K. F., Eskes, H. J., Veefkind, J. P., van Geffen, J. H. G. M., de Zeeuw, M. B., Denier van der Gon, H. A. C., Beirle, S., and Krol, M. C.: Quantification of nitrogen oxides emissions from build-up of pollution over Paris with TROPOMI, *Scientific Reports*, 9, 20033, <https://doi.org/10.1038/s41598-019-56428-5>, 2019.
- McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., Zheng, B., Crippa, M., Brauer, M., and Martin, R. V.: A global anthropogenic emission inventory of atmospheric pollutants from sector- and fuel-specific sources (1970–2017): an application of the Community Emissions Data System (CEDS), *Earth System Science Data*, 12, 3413–3442, <https://doi.org/10.5194/essd-12-3413-2020>, 2020.
- 455 Mijling, B. and Van Der A, R. J.: Using daily satellite observations to estimate emissions of short-lived air pollutants on a mesoscopic scale, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2012JD017817>, 2012.
- Miyazaki, K., Bowman, K., Sekiya, T., Jiang, Z., Chen, X., Eskes, H., Ru, M., Zhang, Y., and Shindell, D.: Air Quality Response in China Linked to the 2019 Novel Coronavirus (COVID-19) Lockdown, *Geophysical Research Letters*, 47, e2020GL089252, <https://doi.org/10.1029/2020GL089252>, 2020.
- 460 Rey-Pommier, A., Chevallier, F., Ciais, P., Broquet, G., Christoudias, T., Kushta, J., Hauglustaine, D., and Sciare, J.: Quantifying NO_x emissions in Egypt using TROPOMI observations, *Atmospheric Chemistry and Physics*, 22, 11 505–11 527, <https://doi.org/10.5194/acp-22-11505-2022>, 2022.
- Riess, T. C. V. W., Boersma, K. F., van Vliet, J., Peters, W., Sneep, M., Eskes, H., and van Geffen, J.: Improved monitoring of shipping NO₂ with TROPOMI: decreasing NO_x emissions in European seas during the COVID-19 pandemic, *Atmospheric Measurement Techniques*, 15, 1415–1438, <https://doi.org/10.5194/amt-15-1415-2022>, 2022.
- 465 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2016.
- Shah, V., Jacob, D. J., Li, K., Silvern, R. F., Zhai, S., Liu, M., Lin, J., and Zhang, Q.: Effect of changing NO_x lifetime on the seasonality and long-term trends of satellite-observed tropospheric NO₂ columns over China, *Atmospheric Chemistry and Physics*, 20, 1483–1495, <https://doi.org/10.5194/acp-20-1483-2020>, 2020.
- 470 Streets, D. G., Bond, T. C., Carmichael, G. R., Fernandes, S. D., Fu, Q., He, D., Klimont, Z., Nelson, S. M., Tsai, N. Y., Wang, M. Q., Woo, J.-H., and Yarber, K. F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD003093>, 2003.
- Sun, K.: Derivation of Emissions from Satellite-Observed Column Amounts and Its Application to TROPOMI NO₂ and CO Observations, *Geophysical Research Letters*, 49, e2022GL101102, <https://doi.org/10.1029/2022GL101102>, 2022a.
- Sun, K.: Physical oversampling in python (POPY), <https://doi.org/10.5281/zenodo.7331896>, 2022b.
- Sun, K., Zhu, L., Cady-Pereira, K., Chan Miller, C., Chance, K., Clarisse, L., Coheur, P.-F., González Abad, G., Huang, G., Liu, X., Van Damme, M., Yang, K., and Zondlo, M.: A physics-based approach to oversample multi-satellite, multispecies observations to a common grid, *Atmospheric Measurement Techniques*, 11, 6679–6701, <https://doi.org/10.5194/amt-11-6679-2018>, 2018.



- 480 Sun, K., Li, L., Jagini, S., and Li, D.: A satellite-data-driven framework to rapidly quantify air-basin-scale NO_x emissions and its application to the Po Valley during the COVID-19 pandemic, *Atmospheric Chemistry and Physics*, 21, 13 311–13 332, <https://doi.org/10.5194/acp-21-13311-2021>, 2021.
- Valin, L. C., Russell, A. R., and Cohen, R. C.: Variations of OH radical in an urban plume inferred from NO₂ column measurements, *Geophysical Research Letters*, 40, 1856–1860, <https://doi.org/10.1002/grl.50267>, 2013.
- 485 van Geffen, J., Eskes, H., Compernelle, S., Pinardi, G., Verhoelst, T., Lambert, J.-C., Sneep, M., ter Linden, M., Ludewig, A., Boersma, K. F., and Veefkind, J. P.: Sentinel-5P TROPOMI NO₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data, *Atmospheric Measurement Techniques*, 15, 2037–2060, <https://doi.org/10.5194/amt-15-2037-2022>, 2022a.
- van Geffen, J., Eskes, H., Veefkind, J., and Boersma, K.: TROPOMI ATBD of the total and tropospheric NO₂ data products, <https://sentinel.esa.int/documents/247904/2476257/Sentinel-5P-TROPOMI-ATBD-NO2-data-products>, 2022b.
- 490 Veefkind, J. P., Serrano-Calvo, R., de Gouw, J., Dix, B., Schneising, O., Buchwitz, M., Barré, J., van der A, R. J., Liu, M., and Levelt, P. F.: Widespread Frequent Methane Emissions From the Oil and Gas Industry in the Permian Basin, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037 479, <https://doi.org/https://doi.org/10.1029/2022JD037479>, 2023.
- Wang, P., Huang, C., Brown de Colstoun, E. C., Tilton, J. C., and Tan, B.: Global Human Built-up And Settlement Extent (HBASE) Dataset From Landsat, <https://doi.org/10.7927/H4DN434S>, 2017.
- 495 Xue, R., Wang, S., Zhang, S., He, S., Liu, J., Tanvir, A., and Zhou, B.: Estimating city NO_x emissions from TROPOMI high spatial resolution observations – A case study on Yangtze River Delta, China, *Urban Climate*, 43, 101 150, <https://doi.org/https://doi.org/10.1016/j.uclim.2022.101150>, 2022.
- Zhang, L., Jacob, D. J., Knipping, E. M., Kumar, N., Munger, J. W., Carouge, C. C., van Donkelaar, A., Wang, Y. X., and Chen, D.: Nitrogen deposition to the United States: distribution, sources, and processes, *Atmospheric Chemistry and Physics*, 12, 4539–4554, <https://doi.org/10.5194/acp-12-4539-2012>, 2012.
- 500 Zhang, L., Wang, L., Wang, R., Chen, N., Yang, Y., Li, K., Sun, J., Yao, D., Wang, Y., Tao, M., and Sun, Y.: Exploring formation mechanism and source attribution of ozone during the 2019 Wuhan Military World Games: Implications for ozone control strategies, *Journal of Environmental Sciences*, <https://doi.org/https://doi.org/10.1016/j.jes.2022.12.009>, 2022.
- Zhang, Q., Boersma, K. F., Zhao, B., Eskes, H., Chen, C., Zheng, H., and Zhang, X.: Quantifying daily NO_x and CO₂ emissions from Wuhan using satellite observations from TROPOMI and OCO-2, *Atmospheric Chemistry and Physics*, 23, 551–563, <https://doi.org/10.5194/acp-23-551-2023>, 2023.
- 505 Zheng, B., Zhang, Q., Geng, G., Chen, C., Shi, Q., Cui, M., Lei, Y., and He, K.: Changes in China’s anthropogenic emissions and air quality during the COVID-19 pandemic in 2020, *Earth System Science Data*, 13, 2895–2907, <https://doi.org/10.5194/essd-13-2895-2021>, 2021.