

Correlation between marine aerosol optical properties and wind fields over remote oceans with use of spaceborne lidar observations

Kangwen Sun¹, Guangyao Dai¹, Songhua Wu^{1,2,3}, Oliver Reitebuch⁴, Holger Baars⁵, Jiqiao Liu⁶, Suping Zhang⁷

5 ¹College of Marine Technology, Faculty of Information Science and Engineering, Ocean University of China, 266100 Qingdao, China

²Laoshan Laboratory, 266237 Qingdao, China

³Institute for Advanced Ocean Study, Ocean University of China, 266100 Qingdao, China

10 ⁴Institut für Physik der Atmosphäre, Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), 82234 Oberpfaffenhofen, Germany

⁵Leibniz Institute for Tropospheric Research (TROPOS), 04318 Leipzig, Germany

⁶Laboratory of Space Laser Engineering, Shanghai Institute of Optics and Fine Mechanics, Chinese Academy of Sciences, 201800 Shanghai, China

15 ⁷Physical Oceanography Laboratory, Ocean University of China, 266100 Qingdao, China

Correspondence to: Guangyao Dai (daiguangyao@ouc.edu.cn)

Abstract. Marine aerosol is mainly produced by wind, which is also a vital element impacting the transport, evolution and dissipation of marine aerosol. The understanding of the accurate relationships between marine aerosol optical properties and wind speed will improve the global aerosol transport models, the satellite-retrieved AODs, the atmospheric correction of ocean color and the study of biogeochemical cycles. Aeolus, the worldwide first ever wind detection lidar satellite, had the ability to measure wind information and particulate optical properties simultaneously, which provide the opportunity to explore the absolutely synchronous relationships between marine aerosol optical properties and wind speeds. Furthermore, thanks to the Aeolus measurement of vertical profiles, the relationships can be discussed in different vertical layers. In this paper, utilizing Aeolus data, the relationships between the optical properties at 355 nm of marine aerosol and the corresponding instantaneous co-located wind speeds of three remote ocean areas are explored and discussed at two separate vertical atmospheric layers (0-1 km and 1-2 km, correspond to the heights within and above marine atmospheric boundary layer (MABL)), revealing the marine aerosol related atmospheric background states. The marine aerosol extinction/backscatter coefficients and the background wind speeds show positive relationships and they were fitted by power law functions, of which the corresponding R^2 are all higher than 0.9. Both the MABL and the higher layer above the MABL will receive the marine aerosol produced and transported by the wind from the air-sea interface. The marine aerosol load at the lower layer (MABL) is stronger than at the higher layer. The marine aerosol enhancements caused by the background wind are more intensive at the MABL. The gradient change points of marine aerosol extinction/backscatter coefficients appear during the growth of them with wind speed, above which the growth rate becomes lower. It might illustrate that the enhancement of marine aerosol driven by wind includes two phases, among which one is rapid growth phase with high dependency of wind, and another is slower growth phase after the

35 gradient change points. As derived data from Aeolus, the averaged marine aerosol optical depth along with wind speed is acquired and utilized to verify the results by the comparison with CALIPSO retrieved results reported in previous work, and besides, the averaged marine aerosol lidar ratio at 355 nm along with wind speed is discussed for the relationship between marine aerosol particle size and wind speed.

1 Introduction

40 According to the Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report, the total emission of marine aerosol (including marine primary organic aerosol) produced from ocean is 1400 to 6800 $\text{Tg} \cdot \text{yr}^{-1}$, which is considered the largest natural aerosol input to the atmosphere globally (Boucher et al., 2013). Accurate estimation of marine aerosol production, evolution and dissipation, and the knowledge of marine aerosol spatial and temporal distribution are significant for studying the global energy budget, aerosol-cloud interactions and visibility changes (Latham and Smith, 1990; Murphy et al., 1998; O'Dowd et al., 1999; Haywood et al., 1999; de Leeuw et al., 2000; Kaufman et al., 2002; Smirnov et al., 2012). Radiative forcing caused by marine aerosol is an important component of the global energy budget. It was reported that the average marine aerosol optical depth (AOD_{mar}) is approximately 0.15 while the volume concentration of cloud condensation nuclei from marine aerosol is around 60 cm^{-3} (Kaufman et al., 2002; Lewis and Schwartz, 2004). Therefore, marine aerosol has the direct impact and the indirect impact on radiative forcing, which are scattering and absorbing solar radiation, and converting cloud microphysical properties, respectively (Murphy et al., 1998; Pierce and Adams, 2006). The knowledge of the impact of the magnitude and changes of marine aerosol emissions on the shifts in climate and marine ecosystem processes is limited (IPCC, 2021).

Marine aerosols mainly include primary sea spray particles and secondary aerosols produced by the oxidation of emitted precursors. Sea spray particles, composed of sea-salt and primary organic aerosols, are produced by wind induced wave breaking as well as the wind driving direct mechanical disruption of waves crests (O'Dowd and de Leeuw, 2007; IPCC, 2021). Moreover, as a dynamical meteorological factor, wind speed also has vital influence on the transport, evolution and dissipation of aerosols. Consequently, the wind speed is a crucial factor which governs the production and life cycle of marine aerosol (Lewis and Schwartz, 2004). Exploring the accurate relationships between marine aerosol optical properties (aerosol optical depth (AOD), extinction coefficient (α), backscatter coefficient (β), etc.) and wind speed are significant for improving global aerosol transport models (Jaegle et al., 2011; Madry et al., 2011; Fan and Toon, 2011), for enhancing satellite-retrieved AODs (Kahn et al., 2010; Kleidman et al., 2012), for atmospheric correction of ocean color (Zibordi et al., 2011) and for the study of biogeochemical cycles (Meskhidze and Nenes, 2010). Several efforts have been reported to explore the relationship between the AOD or aerosol extinction coefficient over the ocean and wind speed. Utilizing either satellite-retrieved AODs (Glantz et al., 2009; Huang et al., 2010; Lehahn et al., 2010; O'Dowd et al., 2010; Grandey et al., 2011) or surface (coast, island or ship)-based measurement AODs (Platt and Patterson, 1986; Villevalde et al., 1994; Smirnov et al., 1995; Wilson and

Forgan, 2002; Smirnov et al., 2003; Shinozuka et al., 2004; Mulcahy et al., 2008; Lehahn et al., 2010; Adames et al., 2011; Sayer et al., 2012; Smirnov et al., 2012), major previous researches focused on the AOD measured by passive instruments (mainly sun-photometer). From these studies, various power-law or linear relationships revealing positive correlation between AODs over the ocean and surface wind speed were established. The passive instruments lack the abilities of distinguishing marine aerosol from other aerosols, acquiring vertical profiles of aerosols, and retrieving aerosol optical properties without sunlight (except for lunar-photometer) and under cloudy conditions (Kiliyanpilakkil and Meskhidze, 2011; Winker and Pelon, 2003). Active optical instruments for aerosol measurements, mainly like lidar, were also used in revealing the relationship between AOD/extinction coefficient of marine aerosol and wind speed. A shipborne depolarization lidar was occupied to acquire aerosol extinction coefficients over the East Sea of Korea near Busan and Pohang, associated with the measurement of an anemometer mounted on a mast, finding a positive linear relationship ($R^2=0.57$) between extinction (532 nm) at 300 ± 50 m and wind speed at 20 m (Shin et al., 2014). However, this relationship was established with data offshore thus it can not be representative for the global ocean. Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) onboard Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) mission is capable of measuring the global aerosol optical properties vertical distributions and recognizing aerosol types (include “clean marine”). Kiliyanpilakkil and Meskhidze (2011) selected CALIOP-retrieved pure AOD_{mar} below 2 km over ocean by utilizing the CALIOP aerosol subtype products and combined them with the surface wind speed provided by the Advanced Microwave Scanning Radiometer (AMSR-E) on board the Aqua satellite, acquiring a relatively complex increasing regression function, which will be presented and compared in Section 4.4.2 of this paper. Besides, Prijith et al. (2014) also made use of CALIOP-retrieved AODs below 0.5 km over ocean and the surface wind speed, obtaining nearly positive correlation linear relationships. Nevertheless, the assumed marine aerosol lidar ratio (LR_{mar}) (20 sr at 532nm) is used in the AOD_{mar} retrieval process of CALIOP (Kiliyanpilakkil and Meskhidze, 2011), but the LR_{mar} can vary from 10 sr to around 40 sr at 532 nm (Groß et al., 2013; Groß et al., 2015; Bohlmann et al., 2018; Floutsi et al., 2023), which could generate deviations during the retrieval of AOD_{mar} . In summary, to explore the accurate relationship between the marine aerosol optical properties and the wind speed, it is essential to conduct global continuous observations and obtain the information of aerosol type identification, while vertical profiles of aerosols can provide extra spatial information for further analysis. Moreover, previous studies mostly focused on the layer AOD_{mar} and ocean surface wind speed, exploring the probable production of marine aerosol driven by surface wind. The relationship between the vertical marine aerosol optical properties and the corresponding spatiotemporally synchronous wind speed is still to be investigated, which represents the marine-atmospheric background state and may reveal the transport and evolution of the marine aerosol vertically.

Atmospheric Laser Doppler Instrument (ALADIN) was a first ever spaceborne direct detection wind lidar, as the single payload installed on the Aeolus mission from the European Space Agency (ESA), which was launched into space in August 2018 (Stoffelen et al., 2005; Reitebuch et al., 2012; Kanitz et al., 2019). As a direct detection high-spectral-resolution lidar, ALADIN has the capability in providing the global aerosol optical properties (e.g., α and β) profiles at 355 nm (Level 2A product),

the horizontal-line-of-sight (HLOS) wind speed profiles (Level 2B product), and the wind vector profile from the European
100 Centre for Medium-Range Weather Forecasts (ECMWF) model along the Aeolus track (Level 2C product) (Rennie et al.,
2020). It should be emphasized that the aerosol and wind product are retrieved from the backscattered signal of the same laser
light pulse emitted from ALADIN to the atmosphere, hence the geolocation and time information of these products is
completely consistent for every profile. The detection altitude range of these products is from the earth surface to around 20
km and the vertical resolutions varies from 0.25 km to 2 km (from bottom to top). Though regarded as a by-product, the particle
105 optical property products are still demonstrated to provide valuable information about particles, especially on the detection
and characterization of aerosol and cloud layers and on the lidar ratios (LRs) (Baars et al., 2021; Flament et al., 2021; Abril-
Gago et al., 2022). It should be emphasized that the lowest altitude bins of Aeolus Level 2A and Level 2B products could be
contaminated by reflections from the land or ocean surface, and are thus not representative for the atmospheric wind speed and
the aerosol optical properties (Wu et al., 2022). Dai et al. (2022) conducted the first attempt on the combined application of
110 the aerosol products (Level 2A products) and the wind vector products (Level 2C products) of ALADIN, observing an
enormous dust transport event occurred in June 2020 from the Sahara to the Americas, describing the transport quantitatively
by calculating dust advection.

As mentioned above, Aeolus can provide global high spatial and temporal resolution aerosol optical properties profiles and
wind speed profiles despite the lack of the lowest bins close to the ground. Additionally, CALIOP can provide global aerosol
115 types information. Hence, the combination of Aeolus-CALIOP products is capable of analysing the relationship between the
marine aerosol optical properties (e.g., α , β , LR) at 355 nm and wind speed globally and vertically. In this paper, by utilizing
Aeolus Level 2A, Level 2C products and CALIOP aerosol subtype products, we firstly 1) select ocean areas far from land and
examine the domination of marine aerosol over these areas with the CALIOP aerosol classification products, and then 2) try
to acquire the pure marine aerosol optical properties (α , β , LR) at 355 nm and the corresponding wind speeds from Aeolus
120 products, and to analyse the spatial distributions of those atmospheric state parameters at two separate vertical layers (ocean
surface to 1 km, 1 km to 2 km, corresponding to the layers within and above the marine atmospheric boundary layer (MABL),
respectively), and finally 3) explore the relationship between the marine aerosol optical properties and the wind speeds
vertically above ocean. Generally, the highlights of this work mainly include 1) acquiring the spatiotemporally synchronous
relationship between the aerosol optical properties (α , β , LR) and the instantaneous wind speeds, which could indicate the
125 background atmosphere states within and above the MABL over remote ocean, 2) conducting analysis at two separate height
layers above ocean surface to explore the vertical differences in aspect of the wind-drive marine aerosol evolution.

The paper is organized as follows: section 2 introduces the spaceborne lidars and their specific products used in this study;
section 3 provides the methodology of study areas selection, data pre-processing and data analyses for relationship exploration
between marine aerosol optical properties and wind speed; section 4 presents the procedure of study areas selection, then
130 analyses and discusses the marine aerosol optical properties, the wind speed, and their relationship above three selected areas.

2 Spaceborne lidars and products

2.1 ALADIN/Aeolus

Since its launch in August 2018, ALADIN, has been globally observing the profiles of the component of the wind vector along the laser's line of sight (LOS), and the profiles of aerosol optical properties, for more than four years. Aeolus flies at a mean altitude of about 320 km in a sun-synchronous orbit with the local equatorial crossing time of about 06:00 and 18:00, a daily quasi-global coverage (about 16 orbits per day) with an orbit repeat cycle of 1 week (111 orbits) (Reitebuch, 2012). Designed as a high-spectral-resolution lidar with a laser wavelength of 354.8 nm, ALADIN has the ability to acquire wind profiles and particle optical properties simultaneously with its two separate optical frequency discrimination channels named as Rayleigh channel and Mie channel. The detailed descriptions of the instrument design and the measurement concept are introduced in, e.g., Ansmann et al. (2007), Dabas et al. (2008), Flamant et al. (2008), Reitebuch (2012), Lux et al. (2020) and Flament et al. (2021).

Processed in different phases, Aeolus data products are classified at several levels: Level 0 (instrument housekeeping data), Level 1B (engineering-corrected HLOS winds), Level 2A (aerosol and cloud layer optical properties), Level 2B (meteorologically representative HLOS winds) and Level 2C (Aeolus-assisted wind vectors) (Flamant et al., 2008; Tan et al., 2008; Rennie et al., 2020). It should be emphasized that Level 2C wind vectors are the outputs from the assimilation of the Aeolus Level 2B products in the ECMWF numerical weather prediction (NWP) operational model after 9 January 2020 (Rennie et al., 2021). In addition, the products of Aeolus are available into different Baselines which correspond to different processor versions used to derive the products. The products were firstly released as Baseline 07 at the beginning and updated to Baseline 14 until this study is conducted. As mentioned above, we use Level 2A and Level 2C products of Aeolus for the study of the relationship between marine aerosol optical properties and wind speeds. As the because Level 2C products can provide both components of the wind vector that we use Level 2C instead of Level 2B products of Aeolus. The time coverage of Aeolus products used in this study is from 20 April 2020 to 4 July 2022. Thus, in the aspect of the utilized Level 2A products, the data processors are Baseline 11 (20 April 2020 to 26 May 2021), Baseline 12 (26 May 2021 to 6 December 2021), Baseline 13 (6 December 2021 to 29 March 2022) and Baseline 14 (29 March 2022 to 4 July 2022), while as for the Level 2C products, the data processors are Baseline 09 (20 April 2020 to 9 July 2020), Baseline 10 (9 July 2020 to 8 October 2020), Baseline 11 (8 October 2020 to 26 May 2021), Baseline 12 (26 May 2021 to 6 December 2021), Baseline 13 (6 December 2021 to 29 March 2022) and Baseline 14 (29 March 2022 to 4 July 2022), respectively (<https://aeolus-ds.eo.esa.int/oads/access/>, last access: 16 February 2023). The Level 2C NWP wind vector products from ECMWF used in this study are obtained after assimilation of the Level 2B observed HLOS wind products.

2.2 CALIOP/CALIPSO

CALIOP, one of the payloads installed on CALIPSO, has been measuring global vertical aerosol and cloud optical properties profiles for more than 16 years since 2006. It can provide α at 532 nm and 1064 nm, β at 532 nm and 1064 nm,

depolarization ratio at 532 nm, vertical feature mask (VFM) products and so on (Winker et al., 2009). The VFM products comprise the vertical information along every profile on the identification of clouds and aerosols, and further, on the subtype classification of clouds and aerosols. For cloud and aerosols identification, the cloud-aerosol discrimination (CAD) algorithm is applied based on layer-integrated volume depolarization ratio, layer-integrated total attenuated color ratio, layer-mean attenuated backscatter at 532 nm, latitude and altitude (Liu et al., 2019). The aerosol sub-types are distinguished as “marine”, “dusty marine”, “dust”, “polluted dust”, “continental”, “polluted continental”, “elevated smoke” and “others” via the joint analysis of particulate depolarization ratio, integrated attenuated backscatter coefficient at 532 nm, layer top altitude, layer base altitude and surface type (Kim et al., 2018). In this study, CALIOP Level (L2) VFM products are applied to confirm the domination of the marine aerosol over the selected ocean areas. Different versions of the CALIOP L2 VFM product are used, respectively, the versions are 4.10 (20 April 2020 to 1 July 2020), 4.20 (1 July 2020 to 19 January 2022) and 3.41 (19 January 2022 to 4 July 2022).

3 Methodology

In general, the data processing and analysis procedure of this study can be concluded briefly as three parts including selection of the study areas, data pre-processing and data analyses, respectively. The flowchart of the study methodology is shown as Fig. 2.

Firstly, this work mainly focuses on the marine aerosol, hence the ocean areas for the study are supposed to be far away from land to reduce the influence of terrestrial aerosols, e.g. anthropogenic, dust, biomass burning. In this work, we selected three ocean areas located in the North Pacific ocean, South Pacific ocean, South Indian ocean, with the latitude and longitude range of 0° to 30° N and 150° E to 180° to 150° W, 20° S to 60° S and 100° W to 150° W, 20° S to 60° S and 60° E to 90° E, respectively, as shown in Fig. 1. Hence, we call these three remote ocean areas “the NP area”, “the SP area” and “the SI area” in this paper, respectively.




Figure 1: The selected study ocean areas.

The aerosol classification information from CALIOP VFM products are utilized to statistically analyse the aerosol types of the selected areas. It is found that the marine aerosols are mostly distributed at the altitude range of 0 km to 2 km during the VFM processing. Therefore, the statistical analysis of aerosol types is conducted at the same altitude range. It is considered that the

marine aerosol dominates in the selected area if the percentage of aerosol subtype “marine” is larger than 75% meanwhile the
190 percentage sum of “marine” and “dusty marine” is above 90%, then the study can be continued for this area.

α at 355 nm and β at 355 nm retrieved by the standard correction algorithm (SCA) from Aeolus Level 2A product are used
in this study, as the SCA processing is capable to produce more stable α and β than the Mie channel algorithm (Flament et
al., 2021). Furthermore, the mid-bin product (sca_optical_properties_mid_bins) of the SCA product are chosen as a result from
that the product retrieved as the mid-bin algorithm is more robust (Baars et al., 2021; Flament et al., 2021). To ensure a high
195 data quality and hence to acquire the relationship between the optical properties and wind speed, a rigorous quality control has
to be applied. In the aspect of quality control, negative α and β are excluded, and then the quality flags (“bin_1_clear” and
“processing_qc_flag”) provided in the Level 2A product are applied to filter out invalid data (Traçon et al., 2022). Additionally,
the outliers are eliminated by the method of Tukey’s test. By using the lower quartiles Q_L (25% positions of the data) and
upper quartiles Q_U (75% positions of the data), this method classifies the data below $Q_L - 3 \cdot (Q_U - Q_L)$ or above
200 $Q_L + 3 \cdot (Q_U - Q_L)$ as outliers (Hoaglin et al., 1986). As the Aeolus products do not distinguish the aerosol layers and cloud
layers, the marine aerosol optical properties may be contaminated by the cloud layers. Aeolus measured particulate β ,
combined with relative humidity (RH) and molecular β from the ECMWF NWP model provided in the Level 2A product are
utilized to screen the cloud layers. It is considered that a cloud is quite likely to exist if the backscatter ratio (BR) (total
backscatter coefficient/molecular backscatter coefficient) at 355 nm is larger than 2.5 or the RH is larger than 94% (Flamant
205 et al., 2020). Therefore, in this study, when the BR is larger than 2.5 or the RH is higher than 94%, the corresponding data bin
is regarded as cloud contaminated and is eliminated. With this cloud screening approach, in this study, 9%, 35%, 40% data in
the altitude range of 0-2 km was eliminated for the NP area, the SP area and the SI area, respectively. Due to the instrument
design of ALADIN, it can only detect the co-polar backscatter light, leading to the lack of the depolarized portion of the β
(Flamant et al., 2020). According to Groß et al. (2015), the depolarization ratio at 355 nm of marine aerosol ($\delta_{mar,355nm}$) is
210 approximately 0.02 when the RH is larger than 50%. Nevertheless, dried marine aerosol layers can significantly depolarize
and the depolarization ratio will vary from 0.02 to around 0.1, so the typical $\delta_{mar,355nm}$ of humid marine aerosol (RH>50%) is
not suitable for dried aerosol (Haarig et al. 2017; Bohlmann et al. 2018). Consequently, to correct the marine aerosol
backscatter coefficient with the typical $\delta_{mar,355nm}$ of humid marine aerosol, the data with RH>50% is retained (around 95%
data is retained), and thus with the typical $\delta_{mar,355nm}$ the total marine aerosol backscatter coefficient β_{mar} can be calculated by
215 the following Eq. (1):

$$\beta_{mar} = (1 + \delta_{mar,355nm}) \cdot \beta_{mar,Aeolus-co}, \quad (1)$$

where $\beta_{mar,Aeolus-co}$ is the original marine aerosol backscatter coefficient measured by ALADIN. It should be illustrated that
all the aerosol β s from Aeolus identified as $\beta_{mar,Aeolus-co}$ s and then utilized to calculate β_{mar} s by formula (1) is under the ideal

assumption that marine aerosol is the only aerosol type in the study areas. Though the study areas are all located in the remote
 220 ocean far away from land and are evaluated as “marine aerosol dominate” by CALIOP, there are a few terrestrial aerosols like
 dust, polluted dust, polluted continental and smoke, with the total proportion of no more than 10% (see Section 4.1 for the
 detail). For the part of terrestrial aerosols, the depolarization ratios at 355 nm of them are 0.22-0.24 for dust, 0.16 for polluted
 dust, 0.01 for polluted continental and 0.03 for smoke, among which the dust’s and the polluted dust’s are much larger than
 $\delta_{mar,355nm}$ (Floutsi et al., 2023). Consequently, regarding all the aerosols as marine aerosol and correcting β_{mar} by formula (1)
 225 leads to the obvious underestimation of the β for dust and polluted dust. Nevertheless, in view of the small proportions of
 dust (no more than 3.15%) and polluted dust (no more than 0.79%) above the study areas and thanks to the statistical analyses
 of data for a long term, the assumption that regarding all the aerosols as marine aerosol is considered not to critically impact
 the β_{mar} - wind speed relationship, while it should be noticed that the actual β is a little bit larger than the β_{mar} .

As for the wind vector data, Aeolus Level 2C product provides the u component (zonal components of wind vector) and v
 230 component (meridional components of wind vector) from the ECMWF model after assimilation of Level 2B observational
 wind product, at the same data bins of the Level 2A optical properties product. Hence the wind speed ws can be calculated
 with these two components by the following Eq. (2):

$$ws = \sqrt{u^2 + v^2} . \quad (2)$$

With the re-processed marine aerosol optical properties extinction coefficient α_{mar} and β_{mar} , and the corresponding ws , it is
 235 possible to explore the relationship between these parameters. At the beginning of data analyses, α_{mar} , β_{mar} and ws within
 the altitude range of 0 km to 2 km are selected, where the marine aerosol dominates according to the analysis of CALIOP
 VFM. Further, the whole study height range is divided into two individual layers. Referring the results of Luo et al. (2014),
 Luo et al. (2016) and Alexander et al. (2019), the MABL height of the remote ocean is summarized as around 1 km. Moreover,
 calculated with ECMWF provided boundary layer heights at the three study areas for the time period of 20 April 2020 to 26
 240 May 2021, the mean values and the standard deviations are 787.47 ± 231.77 m at the NP area, 939.39 ± 360.20 m at the SP
 area and 1005.29 ± 366.60 m at the SI area. Hence, the boundary height of the two vertical layers is set as 1 km, approximately
 corresponding to the mean MABL height of remote ocean. Though the MABL heights are variable and thus set as 1 km will
 lead to the potential inaccuracies, restricted by the relatively low height resolution of Aeolus (0.25 km below 0.5 km, 0.5 km
 in the range of 0.5 km to 2 km), utilizing more precise height boundaries won’t make more sense. It is considered that the
 245 statistical results of the 0-1 km layers and the 1-2 km layers are capable to generally represent the atmospheric conditions
 within the MABL and above the MABL. The lower layer with the altitude range of 0 km to 1 km is called $Layer_L$ in this paper
 and the higher layer with the altitude range of 1 km to 2 km is called $Layer_H$. It should be emphasized that in $Layer_L$, the
 lowest Aeolus Level 2A products (particle optical properties) data bins with the altitude of lower than about 0.25 km, are
 absent to avoid the ground return signals’ contamination. This leads to that the actual marine aerosol optical properties altitude
 250 range of $Layer_L$ is around 0.25 km to 1 km. Over the selected ocean areas, the spatial distribution of the α_{mar} , β_{mar} and ws

are acquired with the longitude-latitude grid of $5^\circ \times 5^\circ$ at two separate layers. Then the relationship analyses between the optical properties (α_{mar} , β_{mar}) and ws of these two layers are conducted by the average calculations of the optical properties along ws grids ($1 \text{ m} \cdot \text{s}^{-1}$) and the parametric curve fitting. For the average calculations, specifically, a grid with resolution of $1 \text{ m} \cdot \text{s}^{-1}$ from $0 \text{ m} \cdot \text{s}^{-1}$ to $30 \text{ m} \cdot \text{s}^{-1}$ is defined and the mean values and the standard deviations are calculated along the grid at both layers above the study areas, respectively. It should be emphasized primarily that before the calculation of mean of each wind speed grid, the outliers larger/less than the mean plus/minus one standard deviation are eliminated. About 70% to 80% α_{mar} and β_{mar} are retained after the elimination. The quite strict outlier removal is conducted here to reject the data unrepresentative for marine aerosol (may be contaminated by cloud thus become higher than the typical range). Hence, it can guarantee the data quality and validity of the pure marine aerosol optical properties in the statistical analysis process. Moreover, the wind speed grid of which the data counts are less than 100 is considered unrepresentative and the statistical result of this grid is abandoned. The derived data of α_{mar} , β_{mar} , averaged AOD_{mar} and LR_{mar} are obtained and discussed, as well. The AOD_{mar} is acquired by integrating Aeolus retrieved α_{mar} within 2 km of every single profile. The AOD_{mar} is calculated within the height of 2 km in order to compare with the previous result of CALIOP, where the integration height is the same as that in this study. The averaged AOD_{mar} along the ws grid are obtained and then are compared with the AOD_{mar} - ws relationships from a previous study in Section 4.4.1. The LR_{mar} are derived via dividing α_{mar} by β_{mar} for each corresponding data bin. The spatial distribution of the LR_{mar} are presented in Section 4.2, meanwhile the relationship between the variations of the LR_{mar} along ws grids and the marine aerosol particle size are discussed in Section 4.4.2.




Figure 2: Flowchart of the study methodology.

4.1 Aerosol types analysis

To verify the domination of marine aerosol, as introduced in Section 3 of this paper, the CALIOP VFM aerosol classification products are applied. The proportions of eight aerosol types (marine, dusty marine, dust, polluted dust, continental, polluted continental, smoke and others) are counted at two vertical layers defined in Section 3 over the NP area, the SP area and the SI area, respectively, as shown the histograms in Fig. 3. The proportions of marine aerosol at Layer_L in these three separate areas are 87%, 84% and 84% while the proportions at Layer_H are 84%, 79% and 79% respectively, which are all larger than 75%. Moreover, the percentage sums of marine aerosol and dusty marine aerosol are all above 90%, at both layers and over all study areas. Consequently, the selected areas NP, SP and SI can be regarded as the marine aerosol dominating areas. It should be illustrated that “dusty marine” was a new aerosol subtype raised for the first time in the version 4.10 of the CALIOP VFM product and was absent in the version 3.41, which was identified from part of version 3.41’s “polluted dust” with the criteria of “surface type” and “layer base altitude”. Using the version 3.41 of the CALIOP VFM data for the period of 19 January 2022 to 4 July 2022 led to the underestimate of “dusty marine” portion and the total marine aerosol portion. Even though under the condition of underestimate, the percentage of total marine aerosol are larger than 90%, which means the real proportion of total marine aerosol is higher, and hence the conclusion that the marine aerosol dominates in the altitude range of 0-2 km above these three areas is still valid.





Figure 3: Aerosol types statistical analyses over (a)(d) the NP area, (b)(e) the SP area and (c)(f) the SI area at two separate layers.

In this section, the study areas in this paper are introduced. With the aerosol types statistical analyses, the dominations of marine aerosol are confirmed over these three areas. It should be illustrated that among the areas, the NP area is mainly located in low latitudes or tropics, while the SP area and the SI area are in mid-latitude region.

4.2 Spatial distribution of wind speed and aerosol optical properties

With Aeolus L2A products (particle optical properties) and L2C products (ECMWF model winds) from April 2020 to July 2022, calculated for every $5^\circ \times 5^\circ$ grid, the averaged ws , α_{mar} , β_{mar} and LR_{mar} spatial distributions of $Layer_H$ and $Layer_L$ are acquired. The averaged atmospheric parameters spatial distributions of the NP area, the SP area and the SI area are presented in Fig. 4, Fig. 5 and Fig. 6, respectively. These figures describe the atmospheric background state of optical properties and wind speed within ($Layer_L$) and above ($Layer_H$) the MABL over the study areas. Additionally, the mean values and the standard deviations of these atmospheric parameters at $Layer_H$ and $Layer_L$ are calculated for each study area by averaging the spatial distributions of $5^\circ \times 5^\circ$ grid, and are presented in Fig. 7.



300 **Figure 4: Wind speed (ws), marine aerosol extinction coefficient (α_{mar}), marine aerosol backscatter coefficient (β_{mar}), and marine aerosol lidar ratio (LR_{mar}) spatial distributions above the North Pacific (NP) area at $Layer_H$ and $Layer_L$.**




Figure 5: Wind speed (ws), marine aerosol extinction coefficient (α_{mar}), marine aerosol backscatter coefficient (β_{mar}), and marine aerosol lidar ratio (LR_{mar}) spatial distributions above the South Pacific (SP) area at $Layer_H$ and $Layer_L$.

305





Figure 6: Wind speed (ws), marine aerosol extinction coefficient (α_{mar}), marine aerosol backscatter coefficient (β_{mar}), and lidar ratio (LR_{mar}) spatial distributions above the South Indian (SI) area at $Layer_H$ and $Layer_L$.



310

Figure 7: Mean values at $Layer_H$ and $Layer_L$ of (a) marine aerosol extinction coefficient (α_{mar}), (b) marine aerosol backscatter coefficient (β_{mar}), (c) marine aerosol lidar ratio (LR_{mar}) and (d) wind speed (ws) above the North Pacific (NP) area (blue standard deviation bars), the South Pacific (SP) area (red standard deviation bars), and the South Indian (SI) area (yellow standard deviation bars).

315

In Fig. 7, the mean values and the deviations of the parameters represent the background atmospheric states within the MABL ($Layer_L$) and over the MABL ($Layer_H$) above each study areas. The averaged ws are $8.1 \pm 1.0 \text{ m}\cdot\text{s}^{-1}$, $11.1 \pm 3.5 \text{ m}\cdot\text{s}^{-1}$, $12.0 \pm 3.5 \text{ m}\cdot\text{s}^{-1}$ at $Layer_L$, while $7.9 \pm 1.1 \text{ m}\cdot\text{s}^{-1}$, $11.5 \pm 4.2 \text{ m}\cdot\text{s}^{-1}$, $12.5 \pm 4.5 \text{ m}\cdot\text{s}^{-1}$ at $Layer_H$, above the NP area, the SP area, and the SI area, respectively. The averaged α_{mar} are $76 \pm 7 \text{ Mm}^{-1}$, $107 \pm 22 \text{ Mm}^{-1}$, $113 \pm 22 \text{ Mm}^{-1}$ at $Layer_L$, while $47 \pm 7 \text{ Mm}^{-1}$, $78 \pm 16 \text{ Mm}^{-1}$, $84 \pm 18 \text{ Mm}^{-1}$ at $Layer_H$, above the NP area, the SP area, and the SI area, respectively. The averaged β_{mar} are $4.1 \pm 0.5 \text{ Mm}^{-1}\cdot\text{sr}^{-1}$, $5.3 \pm 1.7 \text{ Mm}^{-1}\cdot\text{sr}^{-1}$, $5.6 \pm 0.7 \text{ Mm}^{-1}\cdot\text{sr}^{-1}$ at $Layer_L$, while $2.2 \pm 0.4 \text{ Mm}^{-1}\cdot\text{sr}^{-1}$,

$4.0 \pm 0.9 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$, $4.4 \pm 0.7 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$ at Layer_H, above the NP area, the SP area, and the SI area, respectively. The
 320 averaged LR_{mar} are $22.3 \pm 1.6 \text{ sr}$, $25.9 \pm 1.8 \text{ sr}$, $25.0 \pm 1.5 \text{ sr}$ at Layer_L, while $24.5 \pm 1.6 \text{ sr}$, $27.3 \pm 2.3 \text{ sr}$, $26.3 \pm 2.7 \text{ sr}$ at
 Layer_H, above the NP area, the SP area, and the SI area, respectively. It is reported that the typical ranges of α_{mar} and β_{mar}
 at 532 nm over remote ocean areas are around 60 Mm^{-1} to 80 Mm^{-1} and around $1 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$ to $5 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$, respectively,
 observed and retrieved by CALIOP (Prijith et al., 2014; Kiliyanpilakkil and Meskhidze, 2011). Applying the typical α_{mar}
 Ångström exponent from 532 nm to 355 nm of 0.7 ± 1.3 and the typical β_{mar} Ångström exponent from 532 nm to 355 nm of
 325 0.8 ± 0.1 (Floutsi et al., 2023), the converted typical ranges of α_{mar} and β_{mar} at 355 nm can be calculated, which are around
 47 Mm^{-1} to 180 Mm^{-1} and around $1.3 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$ to $7.2 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$. Compared with the typical ranges of α_{mar} and β_{mar} at
 355 nm, calculated from CALIOP retrieved typical ranges of marine aerosol optical properties and the typical conversion
 coefficients, it is considered that the Aeolus retrieved α_{mar} and β_{mar} are reasonable. The mean values of ws , α_{mar} and β_{mar}
 above the NP area are the lowest among the three areas, both at Layer_H and Layer_L, which may be because that this area is
 330 located in low latitudes region of the Northern Hemisphere. The highest mean wind speed of the SI area corresponds to the
 highest α_{mar} and β_{mar} . The mean wind speeds of Layer_H are both larger than those of Layer_L in the SP area and in the SI
 area, while the phenomenon is on the contrary in the NP area. It is worth noting that in all the study areas, the averaged α_{mar}
 and β_{mar} at Layer_L are larger than those at Layer_H, illustrating that the majority of the aerosol from ocean is trapped in the
 MABL while a fraction of marine aerosol can be elevated above the MABL. In the aspect of mean LR_{mar}, the values at Layer_H
 335 are all higher than at Layer_L, and all the values are in a reasonable range referring to Bohlmann et al. (2018), Groß et al. (2011),
 Groß et al. (2015) and Floutsi et al. (2023).

Figure 4, Fig. 5, and Fig. 6 presents the parameters distributions at two layers above the NP area, the SP area and the SI area.
 Primarily, the spatial variations of ws , α_{mar} , β_{mar} are more apparent along the meridian than zonally, both at Layer_H and at
 Layer_L. In the aspect of Layer_L, there are separate distinct high wind speed regions or belts along latitude in the three areas,
 340 which are 5° N to 20° N region of the NP area with the wind speed bins of approximately $8 \text{ m} \cdot \text{s}^{-1}$ to more than $10 \text{ m} \cdot \text{s}^{-1}$, 40° S to 60° S
 region of the SP area with the wind speed bins of more than $10 \text{ m} \cdot \text{s}^{-1}$ to approximately $17 \text{ m} \cdot \text{s}^{-1}$, and 35° S to 60° S
 region of the SI area with the wind speed bins of more than $10 \text{ m} \cdot \text{s}^{-1}$ to approximately $17 \text{ m} \cdot \text{s}^{-1}$ as well. Inspection of
 marine aerosol optical properties, α_{mar} and β_{mar} in the high wind speed regions are obviously larger than in other regions.
 Hence, it can be inferred that, in the MABL, the wind speed and the marine aerosol optical properties tend to be positively
 345 correlated. Referring to Layer_H, shown in the upper four panels of Fig. 4, Fig. 5 and Fig. 6, it can be found that the spatial
 variation trends of ws , α_{mar} , β_{mar} in the three areas are alike with those at Layer_L. The evident high wind speed regions,
 where the wind speeds are up to around $8\text{-}10 \text{ m} \cdot \text{s}^{-1}$ in 5° N to 20° N of the NP area, $15\text{-}18 \text{ m} \cdot \text{s}^{-1}$ in 40° S to 60° S of the SP

area and $13\text{-}19 \text{ m}\cdot\text{s}^{-1}$ in 35° S to 60° of the SI area, also exist at Layer_H while α_{mar} and β_{mar} are slightly enhanced in these regions, which indicates that the wind speed may still have weak positive influence on the marine aerosol optical properties at the higher atmosphere layer above the MABL. Some spatial distribution differences of ws , α_{mar} , β_{mar} between the three areas can be discovered as well. As for the SP area and the SI area, ws , α_{mar} , β_{mar} all mainly present increasing tendencies from north to south. In term of the NP area, besides the obvious enhancements of ws , α_{mar} , β_{mar} in the high wind speed belt, the gradual enhancements of these atmospheric parameters are presented from west to east in this area.

At both layers of the NP area and at Layer_L of the SP area, the LR_{mar} turn out lower in the relatively high wind speed regions, which illustrates a possible negative correlation between LR_{mar} and wind speed. The relationship between these two parameters is analysed and discussed in detail in Section 4.4.2 of this paper.

To conclude, this section presents the atmospheric background state of optical properties and wind speed, and analyses the spatial distributions of ws , α_{mar} , β_{mar} jointly at Layer_H and Layer_L above the NP area, the SP area and the SI area, respectively. The α_{mar} , β_{mar} retrieved from Aeolus Level 2A products are in reasonable agreement with CALIOP and the Aeolus-derived LR_{mar} are also reasonable. It is found that, both at Layer_H and at Layer_L , spatially, the wind speed and α_{mar} , β_{mar} show positive correlation though the optical properties at Layer_L are greater than those at Layer_H , indicating that both layers receive the input of the aerosol produced from ocean by the wind but the majority of the marine aerosol are trapped in the MABL while only a small fraction can be elevated into the higher layer. In addition, as the three study areas are located in different regions, the spatial distributions of ws , α_{mar} , β_{mar} are different.

4.3 Relationship between marine aerosol optical properties and wind speed

In order to determine the relationship between marine aerosol optical properties and corresponding wind speed, utilizing the method introduced in Section 3, the mean values and standard deviations (after the outlier removal) of α_{mar} and β_{mar} along with wind speed grid at two layers above the NP area, the SP area and the SI area are shown in the panel (a) and (b) of Fig. 8, Fig. 9, and Fig. 10, respectively. The regression curves of the optical properties are presented in those figures as well. Power law function is utilized for the curve fitting to describe the trend of marine aerosol optical properties with wind speed. Besides, the data counts in every wind speed grid are shown as the histograms in the panel (a) and (b) of Fig. 8, Fig. 9, and Fig. 10. In order to illustrate the variation tendencies α_{mar} and β_{mar} , the gradients of α_{mar} and β_{mar} with wind speed are also provided in the panel (c) and (d) of Fig. 8, Fig. 9, and Fig. 10. Table 1 summarizes the regression functions together with the corresponding R^2 , and the proportions of different wind speed bins together with the count sums, grouped by areas, layers and optical properties.


From the statistical results with wind speed grids and wind speed ranges, it can be found that most of the wind speeds are below $15 \text{ m}\cdot\text{s}^{-1}$ above the NP area, both at Layer_H and Layer_L , meanwhile the proportion of low wind speed ($0 < ws \leq 8$) is

slightly higher at Layer_H than at Layer_L. As for the SP area and the SI area, the high wind speed ($ws > 15$) accounts for around one fifth and a quarter respectively, and the low wind speed proportion over the SP area is higher than that over the SI area. The wind speed distribution is more concentrated at Layer_L than at Layer_H above these two areas, in view of the lower proportion of low/high wind speed and the higher proportion of middle wind speed ($8 < ws \leq 15$) at Layer_L.

Table 1: Regression functions of the averaged optical properties and the wind speed grids, together with the corresponding wind speed distributions, grouped by areas and layers.

Area	Layer	Optical property	Regression function	R^2	Proportion of wind speed bins ($m \cdot s^{-1}$)			Number of counts
					$0 < ws \leq 8$	$8 < ws \leq 15$	$ws > 15$	
NP	H	α_{mar}	$\alpha_{mar} = 0.084 \cdot ws^{1.8} + 42$	0.95	0.53	0.46	0.01	55758
		β_{mar}	$\beta_{mar} = 0.050 \cdot ws^{1.3} + 1.4$	0.97	0.54	0.45	0.01	64191
	L	α_{mar}	$\alpha_{mar} = 0.070 \cdot ws^{2.3} + 64$	0.99	0.49	0.50	0.01	44857
		β_{mar}	$\beta_{mar} = 0.15 \cdot ws^{1.2} + 2.1$	0.99	0.51	0.48	0.01	55117
SP	H	α_{mar}	$\alpha_{mar} = 5.0 \cdot ws^{0.84} + 54$	0.95	0.46	0.34	0.20	34088
		β_{mar}	$\beta_{mar} = 1.3 \cdot ws^{0.48} + 1.1$	0.96	0.49	0.31	0.20	30348
	L	α_{mar}	$\alpha_{mar} = 6.1 \cdot ws^{0.83} + 73$	0.98	0.45	0.38	0.17	25783
		β_{mar}	$\beta_{mar} = 1.8 \cdot ws^{0.47} + 1.1$	0.97	0.47	0.36	0.17	23854
SI	H	α_{mar}	$\alpha_{mar} = 0.65 \cdot ws^{1.4} + 56$	0.95	0.40	0.35	0.25	19552
		β_{mar}	$\beta_{mar} = 0.60 \cdot ws^{0.59} + 1.7$	0.96	0.42	0.33	0.25	16473
	L	α_{mar}	$\alpha_{mar} = 1.1 \cdot ws^{1.2} + 81$	0.92	0.38	0.41	0.21	15953
		β_{mar}	$\beta_{mar} = 1.2 \cdot ws^{0.47} + 1.7$	0.97	0.46	0.36	0.18	13923


Generally, in all cases shown in Fig. 8, Fig. 9 and Fig. 10, the optical properties at Layer_L are all larger than those at Layer_H in the same wind speed grid, while the variations of marine aerosol optical properties along with wind speed grid can be clearly observed that the tendency is increasing with the wind speed. Moreover, the regression curves are fitted pretty well as the R^2 are all above 0.90.



390 **Figure 8: Relationship between marine aerosol optical properties ((a) for α_{mar} , (b) for β_{mar}) and wind speed above the NP area. The blue circles and error bars represent the means and standard deviations of the optical properties along wind speed grids at Layer_L, while the reds represent the same items at Layer_H. The blue and red dotted-dashed lines are the optical property averages regression curves fitted along the wind speed grid at Layer_L and Layer_H, respectively. The blue and red histograms indicate the data counts of every wind speed grid at Layer_L and Layer_H, respectively. (c) and (d) represent the gradients of α_{mar} and β_{mar} with wind speed at Layer_L (blue lines) and Layer_H (red lines), respectively, while the blue dashed lines and the red dashed lines show the mean values of the gradients at two layers.**
 395


It can be found from the panel (a) and (b) of Fig. 8, in the NP area, α_{mar} at Layer_L increases from 64 Mm^{-1} at 0-1 $\text{m}\cdot\text{s}^{-1}$ wind speed interval to 113 Mm^{-1} at 16-17 $\text{m}\cdot\text{s}^{-1}$ wind speed interval, while it at Layer_H increases from 42 Mm^{-1} at 0-1 $\text{m}\cdot\text{s}^{-1}$ wind speed interval to 57 Mm^{-1} at 17-18 $\text{m}\cdot\text{s}^{-1}$ wind speed interval; β_{mar} at Layer_L increases from 2.2 $\text{Mm}^{-1}\cdot\text{sr}^{-1}$ at 0-1 $\text{m}\cdot\text{s}^{-1}$ wind speed interval to 6.6 $\text{Mm}^{-1}\cdot\text{sr}^{-1}$ at 16-17 m/s wind speed interval, while it at Layer_H increases from 1.6 $\text{Mm}^{-1}\cdot\text{sr}^{-1}$ at 0-1 $\text{m}\cdot\text{s}^{-1}$ wind speed interval to 3.3 $\text{Mm}^{-1}\cdot\text{sr}^{-1}$ at 17-18 m/s wind speed interval. The increments of these two parameters at Layer_L are much larger than at Layer_H. Moreover, the exponents of the regression functions are all larger than 1, indicating the growth rates of the optical properties along the wind grid increases. Referring to the panel (c) and (d) of Fig. 8, the gradients' mean values of the α_{mar} and β_{mar} at Layer_L are higher than those at Layer_H. Besides, the gradients at Layer_L

are mostly larger than that at Layer_H within the same wind speed interval, i.e., the optical properties at Layer_L will increase more rapidly with wind speed. It is worth to notice that for the case that the wind speed is above 10 m·s⁻¹, the gradients of β_{mar} show decreasing tendencies, whereas under the condition when the wind speed is lower than 10 m·s⁻¹, the values of the optical properties' gradients present increasing tendencies, indicating the better fitting by power law functions at lower wind speed. This phenomenon may imply that there might be two distinct variation trends of β_{mar} above or below the wind speed of 10 m·s⁻¹.



410

Figure 9: Relationship between marine aerosol optical properties and wind speed above the SP area. The items represent the same as those of Fig. 8.



415 **Figure 10: Relationship between marine aerosol optical properties and wind speed above the SI area. The items represent the same as those of Fig. 8.**

For the SP area and the SI area, the maximal wind speed can reach up to $28 \text{ m} \cdot \text{s}^{-1}$, while the variations of the optical properties along with wind speed are more complicated. In Fig. 9 (a), the α_{mar} over the SP area show approximate linear growth tendencies with wind speed both at Layer_L and at Layer_H, with the fitting functions' exponents of 0.93 and 0.82. The α_{mar} increase from 72 Mm^{-1} and 52 Mm^{-1} to 130 Mm^{-1} and 111 Mm^{-1} for Layer_L and Layer_H, respectively. Figure 9 (b) shows that the β_{mar} above the SP area increase from $2.7 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$ and $2.1 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$ to $7.0 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$ and $5.3 \text{ Mm}^{-1} \cdot \text{sr}^{-1}$, with the fitting functions' exponents of 0.51 and 0.38 for Layer_L and Layer_H. From Fig. 10 (a) and (b), it can be found that the variations of α_{mar} and β_{mar} with wind speed in the SI area are similar to those in the SP area, except for that the fitting functions' exponents of β_{mar} are larger than 1, which are 1.2 and 1.4 for Layer_L and Layer_H, respectively. In the Layer_H of SI area, α_{mar} at above $25 \text{ m} \cdot \text{s}^{-1}$ can reach up to 137 Mm^{-1} , much larger than that of around 110 Mm^{-1} in the SP area. The panel (c) and (d) of Fig. 9 and Fig. 10 show the gradients of α_{mar} and β_{mar} with the wind speed above the SP area and the SI area. In these four panels, the blue dashed lines (gradients' mean values at Layer_L) are all higher than the red lines (gradients' mean values at Layer_H), illustrating that the increments of α_{mar} and β_{mar} per unit wind speed at Layer_L are larger than those

420

425

at Layer_H, which implies that the input of marine aerosol driven by wind at Layer_L is stronger than at Layer_H. Focusing on the panel (c) of Fig. 9 and Fig. 10, it can be seen that, for both layers of the SP area and the SI area, the gradients of α_{mar} below 15 $\text{m}\cdot\text{s}^{-1}$ are almost all larger than 0 fluctuating slightly around the mean values, while the gradients of α_{mar} above 15 $\text{m}\cdot\text{s}^{-1}$ fluctuate drastically around 0. This phenomenon may indicate that below 15 $\text{m}\cdot\text{s}^{-1}$, both layers continually receive the input of marine aerosol driven by wind, nevertheless when the wind speed is higher than 15 $\text{m}\cdot\text{s}^{-1}$, the dependency of marine aerosol on wind become lower. As for the gradients of β_{mar} above the SP area and the SI area, from the panel (d) of Fig. 9 and Fig. 10, it can be found obviously that for both layers, the gradients of β_{mar} decrease above around 10 $\text{m}\cdot\text{s}^{-1}$. The corresponding variations of β_{mar} above the SP area and the SI area are shown in the panel (b) of Fig. 9 and Fig. 10, of which the β_{mar} increase with higher slopes at the wind speed range of 0-10 $\text{m}\cdot\text{s}^{-1}$ while the slopes of the increasing become lower when the wind speed is above 10 $\text{m}\cdot\text{s}^{-1}$. This phenomenon might indicate that the increasing of β_{mar} with wind speed includes two separate trends regarding 10 $\text{m}\cdot\text{s}^{-1}$ as the change point, consist with the surmise raised in the analysis of the NP area. Consequently, for all of the measurement cases, the marine aerosol optical properties at Layer_L are larger than those at Layer_H in any identical wind speed interval, indicating that the MABL possibly receive more marine aerosol produced and transport from the air-sea interface, while the higher layer above the MABL with the upper boundary of 2 km can also be influenced by the marine aerosol but less. The gradients' mean values of α_{mar} and β_{mar} at Layer_L are all larger than at Layer_H, which implies that the marine aerosol enhancements caused by the background wind are more intensive at the MABL. It should be emphasized that the gradients change during the growth of α_{mar} and β_{mar} along with wind speed. The gradient change point of α_{mar} (15 $\text{m}\cdot\text{s}^{-1}$) is greater than that of β_{mar} (10 $\text{m}\cdot\text{s}^{-1}$), and above them the enhancement rate becomes lower. It might illustrate that the enhancement of marine aerosol driven by wind includes two phases, among them one is growing rapidly phase with high dependency of wind, and another is after the gradient change points, the marine aerosol enhancement by wind is likely to diminish.

4.4 Dependency of aerosol optical depth and lidar ratio with wind speed

4.4.1 Marine aerosol optical depth vs. wind speed

As introduced in Section 1 of this paper, almost all the previous researches on the relationship between marine aerosol's optical properties and wind speed focused on the AOD of marine aerosol. In this study, as well, the attempt on the averaged 0-2 km AOD_{mar} of individual wind speed grid calculation has been conducted to compare the AOD_{mar} - *ws* relationship from previous study. The AOD_{mar} of every single profile is acquired by integrating Aeolus retrieved α_{mar} within 2 km. The wind speed profiles are also averaged for 2 km to correspond to the AOD_{mar} data. Then the relationship between the AOD_{mar} and the

wind speeds is obtained by averaging the AOD_{mar} in each wind speed interval ($0 \text{ m}\cdot\text{s}^{-1}$ - $30 \text{ m}\cdot\text{s}^{-1}$, stepped by $1 \text{ m}\cdot\text{s}^{-1}$). The AOD_{mar} - ws relationship is also explored utilizing the products from the A-Train satellites (Kiliyanpilakkil and Meskhidze, 2011). “clean marine” aerosol AOD at 532 nm above ocean surface (up to 2 km) provided by CALIOP and 10 m daily wind speed provided by AMSR-E were used. It should be noticed that the wind speed used in Kiliyanpilakkil and Meskhidze (2011) is daily ocean surface wind speed, different from that in this study, which is instantaneous layer-averaged wind speed. Collecting the data for the time period from 2006 to 2011 over 15 remote ocean regions globally, the regression curve of is acquired with the averaged AOD_{mar} at 532 nm for each wind speed grid and the surface wind speed which is up to $29 \text{ m}\cdot\text{s}^{-1}$, and the regression function is shown as the following Eq. 3:

$$AOD_{mar,532} = \frac{0.15}{1 + 6.7 \cdot e^{-0.17 \cdot U_{10}}}, \quad (3)$$


where the U_{10} represents daily 10 m ocean surface wind speed.

As described above, the AOD_{mar} data source (from spaceborne lidar observation), the study areas (remote ocean regions globally), and the wind speed range ($0 \text{ m}\cdot\text{s}^{-1}$ - $29 \text{ m}\cdot\text{s}^{-1}$) of the AOD_{mar} - ws relationship exploration in Kiliyanpilakkil and Meskhidze (2011) match well with those of this study. Hence, we select the AOD_{mar} - ws relationship established by Kiliyanpilakkil and Meskhidze (2011) for the comparison. Additionally, due to the different wavelengths of AOD_{mar} used in this study (355 nm) and in Kiliyanpilakkil and Meskhidze (2011) (532 nm), the effort on conversion of the AOD_{mar} at 532 nm to the AOD_{mar} at 355 nm is performed by applying the typical Ångström exponent of marine aerosol. It is reported that the marine aerosol Ångström exponent is surface wind speed related, and a linear relationship was established as the following Eq. 4 (Sayer et al., 2012):

$$A(ws) = 0.69 - 0.030 \cdot ws, \quad (4)$$

where A represents the Ångström exponent and ws represents the wind speed. Then the AOD_{mar} at 532 nm can be converted to the AOD_{mar} at 355 nm by the following Eq. 5:

$$AOD_{mar,355nm}(ws) = \exp\left[A(ws) \cdot \ln \frac{532}{355}\right] \cdot AOD_{mar,532nm}(ws). \quad (5)$$



480 **Figure 11: AOD_{mar} at 355 nm versus wind speed. The blue squares and the corresponding error bars represent the AOD_{mar} means and standard deviations along the ws grid of all the three study areas in this study; the red squares and line represent the AOD_{mar} at 355 nm along the ws grid converted from the regressive relationship between the AOD_{mar} at 532 nm and the ocean surface wind speed reported by Kiliyanpilakkil and Meskhidze (2011).**

In Fig. 11, the averaged AOD_{mar} and the corresponding standard deviations at 355 nm of all three study areas along the wind speed grid are represented as the blue squares and the error bars, while the regression curve of AOD_{mar} at 355 nm versus wind speed converted from Eq. 3 are represented as the red squares and line. Although instantaneous layer-averaged wind speed and daily ocean surface wind speed are used in this study and in Kiliyanpilakkil and Meskhidze (2011) individually, similar tendency of AOD_{mar} at 355 nm versus wind speed are obtained. It can be found that AOD_{mar} is increasing with wind speed, and the gradient of AOD_{mar} turns out becoming higher along the wind speeds when the wind speed is below $15 \text{ m}\cdot\text{s}^{-1}$ while the variation of AOD_{mar} becomes slower above $15 \text{ m}\cdot\text{s}^{-1}$. The converted CALIOP AOD_{mar} are lower than the Aeolus retrieved AOD_{mar} at $0\text{-}10 \text{ m}\cdot\text{s}^{-1}$, nevertheless the formers are all in the standard deviations range of the latter, thus it is considered that the Aeolus retrieved AOD_{mar} and their variation along the wind speed are reasonable. The lower AOD_{mar} from CALIOP after wavelength conversion at low wind speed may arise from using a fixed LR_{mar} of 20 sr at 532 nm for CALIOP AOD_{mar} retrievals while the LR_{mar} can vary with a quite large range of 10 sr to 90 sr (Masonis et al., 2003). Possible uncertainties of the CALIOP retrieved AOD_{mar} at 532 nm is discussed in detail in Kiliyanpilakkil and Meskhidze (2011). Besides, as discussed in Section 4.4.2 of this paper, the particle size of marine aerosol and the LR_{mar} will vary with wind speed, so using the CALIOP AOD_{mar} retrieved with the fixed LR_{mar} may generate additional error in the exploration of the relationship between the AOD_{mar} and the wind speed. Therefore, using Aeolus retrieved AOD_{mar} , which is integrated by independently retrieved extinction coefficient without the assumption of LR_{mar} , could make the AOD_{mar} - ws relationship more reliable.

485
490
495

500 4.4.2 Marine aerosol lidar ratio vs. wind speed

As one of the intensive optical properties, LR_{mar} independent of the aerosol concentration. It is reported that the LR_{mar} depends on the particle size, and specifically, with the reduction of the coarse mode, the total LR turns out to increase (Masonis et al., 2003). The possible reason for this phenomenon is that as the particles become smaller, the extinction is enhanced by the increasing sideward scattering and the backscatter gets weaker due to the decrease of the scattering cross section (Haarig et al., 2017). Aeolus L2A product provide particle extinction-to-backscatter ratio calculated with the raw β , which lacks the depolarized portion, as introduced in Section 3 of this paper. In this work, the corrected LR_{mar} is acquired by dividing the marine aerosol extinction to the marine aerosol depolarization-corrected backscatter. The calculation of the averaged LR_{mar} along wind speed grid has been conducted by averaging the LR_{mar} s of each $1 \text{ m} \cdot \text{s}^{-1}$ wind speed bin, meanwhile the standard deviations are acquired as well. It should be noted that before the statistical calculation, the outliers are eliminated by the method of Tukey's test, which is introduced in Section 3 of this paper.




Figure 12: LR_{mar} versus the wind speed. The dark blue curve, red curve, yellow curve and the corresponding error bars represent the averaged LR_{mar} and their standard deviations above the NP area, the SP area and the SI area, respectively. The purple curve and the corresponding error bars represent the CALIOP-retrieved LR_{mar} at 532 nm (Dawson et al., 2015). The green curve and the light blue curve represent the modelled LR_{mar} at 355 nm and at 532 nm, respectively (Sayer et al., 2012).

In Fig. 12, 0-2 km averaged LR_{mar} variations along with the wind speed above the NP area, the SP area and the SI area are represented as the blue curve, the red curve and the yellow curve, respectively. Generally, the distinct downward trend of the LR_{mar} at relatively low wind speeds ($0-14 \text{ m} \cdot \text{s}^{-1}$ of the NP area, $0-9 \text{ m} \cdot \text{s}^{-1}$ of the SP area and $0-10 \text{ m} \cdot \text{s}^{-1}$ of the SI area) can be observed in all cases. The results reported in this paper are similar to those in the previous studies, of which Dawson et al. (2015) and Sayer et al. (2012) investigated the relationship between LR_{mar} and wind speed utilizing measured LR_{mar} and

modelled LR_{mar} respectively. Combining the corrected CALIOP-retrieved LR_{mar} at 532 nm and 10 m ocean surface wind speed from the AMSR-E, the negative correlation between the LR_{mar} and wind speed is acquired with the wind speed bins of $0 \text{ m}\cdot\text{s}^{-1}$ up to $>15 \text{ m}\cdot\text{s}^{-1}$, shown as the purple curve in Fig. 12 (Dawson et al., 2015). The modelled LR_{mar} at 355 nm and at 532 nm also presents decreasing trends with the wind speed increases, presented as the green curve and the light blue curve in Fig. 12 (Sayer et al., 2012). These results seem to imply that as the wind speed increases for a low wind speed range, the particle size of marine aerosol get larger. The phenomenon is explained by the shift in marine aerosol volume size distribution with wind speed as wind speed increases, the fine mode volume size distribution of marine aerosol turns out decline while the coarse mode distribution becomes larger (Dawson et al., 2015; Smirnov et al., 2003; Sayer et al., 2012). The CALIOP LR_{mar} and the modelled LR_{mar} are all larger than the LR_{mar} of this study but are all in the standard deviation ranges. According to Groß et al. (2011), Groß et al. (2015), Bohlmann et al. (2018) and Floutsi et al. (2023), the pure LR_{mar} at 355 nm can vary from 10 sr to 40 sr, with the average of around 20 sr, thus it is considered that the averaged LR_{mar} in this study are reasonable. At the middle wind speed range ($14 \text{ m}\cdot\text{s}^{-1}$ - $18 \text{ m}\cdot\text{s}^{-1}$ of the NP area, $9 \text{ m}\cdot\text{s}^{-1}$ - $16 \text{ m}\cdot\text{s}^{-1}$ of the SP area, $10 \text{ m}\cdot\text{s}^{-1}$ - $20 \text{ m}\cdot\text{s}^{-1}$ of the SI area), the LR_{mar} show upward tendencies, implying that the marine aerosol particles might be broken up into smaller ones with a wind speed increase. At the very high wind speed above the SP area ($>16 \text{ m}\cdot\text{s}^{-1}$) and the SI area ($>20 \text{ m}\cdot\text{s}^{-1}$), the LR_{mar} turns out decreasing with the wind speed again, which indicates that the particle size of marine aerosol becomes larger at this wind speed condition.




Figure 13: Averaged LR_{mar} versus wind speed at Layer_L and Layer_H, in (a) the NP area, (b) the SP area and (c) the SI area, respectively.

Figure 13 shows the LR_{mar} variations at Layer_L and Layer_H along with the wind speed grid in three study areas. Some divergences of the LR_{mar} variations between the layers can be discovered. As for the NP area, the variation of LR_{mar} at Layer_L is from 29 sr at $1\text{-}2 \text{ m}\cdot\text{s}^{-1}$ to 19 sr at $12\text{-}13 \text{ m}\cdot\text{s}^{-1}$, larger than that at Layer_H, which is from 28 sr at $1\text{-}2 \text{ m}\cdot\text{s}^{-1}$ to 21 sr at $15\text{-}16 \text{ m}\cdot\text{s}^{-1}$. Regarding to the SP area and the SI area, the downward trend of LR_{mar} at high wind speed condition as

mentioned above is not apparent at Layer_L . Moreover, at Layer_H , the LR_{mar} can reach up to 27-28 sr at $15\text{-}25 \text{ m}\cdot\text{s}^{-1}$, close
545 to that at $0\text{-}5 \text{ m}\cdot\text{s}^{-1}$, implying that the marine aerosol particle sizes at low and high wind speed are similar.
Generally, the LR_{mar} dependency along with the wind speed shows a downward trend at relatively low wind speed, then an
upward trend at the middle wind speed, finally again a downward at the very high wind speed (if exist), which implies that the
marine aerosol particle size is increasing along with the wind speed at first, then might be broken up into smaller one by the
enhanced wind speed and ultimately turns out to a larger size again. Several differences of the LR_{mar} variations along with
550 the wind speed appear between the three study areas and the two vertical layers, which may be due to the differences in
meteorological and environmental conditions of the areas and the layers.

5 Summary and conclusion

By utilizing Level 2A products (particle optical properties and numerical weather prediction data) and Level 2C products
(numerical weather prediction wind vector assimilated with observed wind component) provided by ALADIN, and L2 vertical
555 feature mask (VFM) products provided by CALIOP, the optical properties at 355 nm of pure marine aerosol are derived. Then
the combined analysis of marine aerosol optical properties at 355 nm and instantaneous co-located wind speed above remote
ocean areas is conducted. Finally, their relationships are explored and discussed at two separate vertical atmospheric layers
(Layer_L with the height of 0-1 km and Layer_H with the height of 1-2 km, correspond to the heights within and above marine
atmospheric boundary layer (MABL)), revealing the marine aerosol related atmospheric background states within and above
560 the MABL over the remote oceans.

Three study areas located in remote ocean were selected, which were named the North Pacific (NP) area, the South Pacific
(SP) area and the South Indian (SI) area, respectively. Then we examined the domination of marine aerosol with the aerosol
classification data provided by CALIOP VFM products. The proportions of marine aerosol in these three areas are all larger
than 79% respectively while the percentage sums of marine aerosol and dusty marine aerosol are all above 90%. After quality
565 control, cloud screening was conducted with the criteria (relative humidity and backscatter ratio), and 9%, 35%, 40% data was
identified cloud contaminated in the altitude range of 0-2 km then was eliminated for the NP area, the SP area and the SI area,
respectively. Finally, backscatter correction is applied to the Aeolus L2A products. These procedures allow us to obtain reliable,
cloud-free marine aerosol optical properties and the corresponding wind speed.

The statistical results of the marine aerosol optical properties (extinction coefficient (α_{mar}) and backscatter coefficient (β_{mar}))
570 at 355 nm with the wind speed (ws) are analysed at Layer_L and Layer_H , for three study areas respectively. It is found that
both the MABL and the higher layer above the MABL can receive the marine aerosol produced and transported from air-sea
interface. Moreover, the marine aerosol load at the lower layer (MABL) is stronger than at the higher layer. The marine aerosol
enhancements caused by the background wind are more intensive at the MABL. Besides, the gradient change points of α_{mar}

($15 \text{ m} \cdot \text{s}^{-1}$) and of β_{mar} ($10 \text{ m} \cdot \text{s}^{-1}$) were found during the growth of α_{mar} and β_{mar} with wind speed, above which the growth rate of α_{mar} and β_{mar} becomes lower. It might illustrate that the enhancement of marine aerosol driven by wind includes two phases, among which one is growing rapidly phase with high dependency of wind, and another is after the gradient change points, the marine aerosol enhancement by wind is likely to diminish. The α_{mar} - ws curves and the β_{mar} - ws curves were fitted by power law functions and the corresponding R^2 are all higher than 0.9 for both layers above all three study areas. Regarding to the derived data, compared with the regression function between wavelength-converted CALIOP-retrieved AOD_{mar} and 10 m surface wind speed, the AOD_{mar} at 355 nm versus the wind speed in this work shows quite consistent tendency with CALIOP's though the wind speeds used are different. The LR_{mar} and marine aerosol particle size have negative relationship (Masonis et al., 2003). From the relationship between the LR_{mar} and the wind speed, it indicates that as the wind speed is increasing, the particle size of marine aerosol obviously becomes larger at relative low wind speed range, then could be broken up into smaller by wind at higher wind speed, and ultimately turns out a larger state again at very high wind speed. The α_{mar} - ws , β_{mar} - ws models within and above MABL at remote ocean areas were established with Aeolus provided data. Nevertheless, the regression curves of α_{mar} - ws and β_{mar} - ws above three study areas (the NP area, located in the Pacific Ocean, the low latitudes of the Northern Hemisphere; the SP area, located in the Pacific Ocean, the middle latitudes of the Southern Hemisphere; the SI area, located in the Indian Ocean, the middle latitudes of the Southern Hemisphere) are not totally consistent, while the meteorological and environmental conditions apart from wind are also distinct at different regions. It implies that in order to obtain more precise α_{mar} and β_{mar} models, besides wind speed, other meteorological and environmental factors, e.g., atmospheric stability, sea and air temperature, RH, etc. should participate in the establishment of the models, because the production, entrainment, transport and removal of the marine aerosol above the ocean are not only dominated by the wind, but also be impacted by these factors (Lewis and Schwartz, 2004). If future study is capable to obtain more other meteorological parameters above ocean, jointly analysing the aerosol optical properties and the wind together with them, more detailed information of marine aerosol production, entrainment, transport and removal will be acquired.

Data availability

The Aeolus data are downloaded via the website <https://aeolus-ds.eo.esa.int/oads/access/collection> (last access: 9 March 2023). Part of the Aeolus L2A and L2C data we used in this paper are not available publicly at the time the article was submitted. We are allowed to access the data through our participation as a Calibration and Validation team. The CALIOP data can be downloaded from <https://eosweb.larc.nasa.gov/project/CALIPSO> (last access: 9 March 2023).

Author contributions

G. Dai conceived of the idea for correlation between marine aerosol optical properties and wind fields over remote oceans with spaceborne lidars ALADIN, CALIOP; K. Sun wrote the manuscript; K. Sun, G. Dai, S. Wu, O. Reitebuch and H. Baars contributed to the data analyses; J. Liu and S. Zhang contributed to the scientific discussion. All the co-authors reviewed and edited the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Special issue statement

This article is part of the special issue “Aeolus data and their application”. It is not associated with a conference.

Acknowledgments

This study has been jointly supported by the Laoshan Laboratory Science and Technology Innovation Projects under grant LSKJ202201406, the National Natural Science Foundation of China (NSFC) under grant 61975191, 41905022 and U2106210. This work was also supported by Dragon 5 program which conducted by European Space Agency (ESA) and the National Remote Sensing Center of China (NRSCC) under grant 59089.

References

- Abril-Gago, J., Guerrero-Rascado, J. L., Costa, M. J., Bravo-Aranda, J. A., Sicard, M., Bermejo-Pantaleón, D., Bortoli, D., Granados-Muñoz, M. J., Rodríguez-Gómez, A., Muñoz-Porcar, C., Comerón, A., Ortiz-Amezcu, P., Salgueiro, V., Jiménez-Martín, M. M., and Alados-Arboledas, L.: Statistical validation of Aeolus L2A particle backscatter coefficient retrievals over ACTRIS/EARLINET stations on the Iberian Peninsula, *Atmos. Chem. Phys.*, 22, 1425–1451, <https://doi.org/10.5194/acp-22-1425-2022>, 2022.
- Adames, A. F., Reynolds, M., Smirnov, A., Covert, D. S., and Ackerman, T. P.: Comparison of MODIS ocean aerosol retrievals with ship-based sun photometer measurements from the “Around the America’s” expedition, *J. Geophys. Res.*, 116, D16303, <https://doi.org/10.1029/2010JD015440>, 2011.
- Alexander, S. P. and Protat, A.: Vertical profiling of aerosols with a combined Raman-elastic backscatter lidar in the remote Southern Ocean marine boundary layer (43–66°S, 132–150°E), *J. Geophys. Res.-Atmos.*, 124, 12107–12125, <https://doi.org/10.1029/2019JD030628>, 2019.

- Ansmann, A., Wandinger, U., Le Rille, O., Lajas, D., and Straume, A. G.: Particle backscatter and extinction profiling with the spaceborne high-spectral-resolution Doppler lidar ALADIN: methodology and simulations, *Appl. Optics*, 46, 6606, <https://doi.org/10.1364/AO.46.006606>, 2007.
- 630 Baars, H., Radenz, M., Floutsi, A. A., Engelmann, R., Althausen, D., Heese, B., Ansmann, A., Flament, T., Dabas, A., Trapon, D., Reitebuch, O., Bley, S., and Wandinger, U.: Californian wildfire smoke over Europe: A first example of the aerosol observing capabilities of Aeolus compared to ground-based lidar, *Geophysical Research Letters*, 48, e2020GL092194. <https://doi.org/10.1029/2020GL092194>, 2021.
- Bohlmann, S., Baars, H., Radenz, M., Engelmann, R., and Macke, A.: Ship-borne aerosol profiling with lidar over the Atlantic Ocean: from pure marine conditions to complex dust–smoke mixtures, *Atmos. Chem. Phys.*, 18, 9661–9679, <https://doi.org/10.5194/acp-18-9661-2018>, 2018.
- 635 Boucher, O., D. Randall, P. Artaxo, C. Bretherton, G. Feingold, P. Forster, V.-M. Kerminen, Y. Kondo, H. Liao, U. Lohmann, P. Rasch, S.K. Satheesh, S. Sherwood, B. Stevens and X.Y. Zhang, 2013: Clouds and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- 640 Dabas, A., Denneulin, M. L., Flamant, P., Loth, C., Garnier, A., and Dolfi-Bouteyre, A.: Correcting winds measured with a Rayleigh Doppler lidar from pressure and temperature effects, *Tellus A*, 60, 206–215, <https://doi.org/10.1111/j.1600-0870.2007.00284.x>, 2008.
- 645 Dai, G., Sun, K., Wang, X., Wu, S., E, X., Liu, Q., and Liu, B.: Dust transport and advection measurement with spaceborne lidars ALADIN and CALIOP and model reanalysis data, *Atmos. Chem. Phys.*, 22, 7975–7993, <https://doi.org/10.5194/acp-22-7975-2022>, 2022.
- de Leeuw, G., Neele, F. P., Hill, M., Smith, M. H., and Vignati, E.: Production of sea spray aerosol in the surf zone, *J. Geophys. Res.*, 105, 29397–29409, <https://doi.org/10.1029/2000JD900549>, 2000.
- 650 Dawson, K. W., Meskhidze, N., Josset, D., and Gassó, S.: Spaceborne observations of the lidar ratio of marine aerosols, *Atmos. Chem. Phys.*, 15, 3241–3255, <https://doi.org/10.5194/acp-15-3241-2015>, 2015.
- Fan, T. and Toon, O. B.: Modeling sea-salt aerosol in a coupled climate and sectional microphysical model: mass, optical depth and number concentration, *Atmos. Chem. Phys.*, 11, 4587–4610, <https://doi.org/10.5194/acp-11-4587-2011>, 2011.
- Flamant, P. H., Cuesta, J., Denneulin, M.-L., Dabas, A., and Huber, D.: ADM-Aeolus retrieval algorithms for aerosol and cloud products, *Tellus A*, 60, 273–286, <https://doi.org/10.1111/j.1600-0870.2007.00287.x>, 2008.
- 655 Flamant, P. H., Lever, V., Martinet, P., Flament, T., Cuesta, J., Dabas, A., Olivier, M., Huber, D., Trapon, D., and Lacour, A.: Aeolus Level-2A Algorithm Theoretical Basis Document, version 5.7, <https://earth.esa.int/eogateway/documents/20142/37627/Aeolus-L2A-Algorithm-Theoretical-Baseline-Document> (last access: 9 November 2022), 2020.

- 660 Flament, T., Traпон, D., Lacour, A., Dabas, A., Ehlers, F., and Huber, D.: Aeolus L2A aerosol optical properties product: standard correct algorithm and Mie correct algorithm, *Atmos. Meas. Tech.*, 14, 7851–7871, <https://doi.org/10.5194/amt-14-7851-2021>, 2021.
- Floutsi, A. A., Baars, H., Engelmann, R., Althausen, D., Ansmann, A., Bohlmann, S., Heese, B., Hofer, J., Kanitz, T., Haarig, M., Ohneiser, K., Radenz, M., Seifert, P., Skupin, A., Yin, Z., Abdullaev, S. F., Komppula, M., Filioglou, M., Giannakaki, E.,
665 Stachlewska, I. S., Janicka, L., Bortoli, D., Marinou, E., Amiridis, V., Gialitaki, A., Mamouri, R.-E., Barja, B., and Wandinger, U.: DeLiAn – a growing collection of depolarization ratio, lidar ratio and Ångström exponent for different aerosol types and mixtures from ground-based lidar observations, *Atmos. Meas. Tech.*, 16, 2353–2379, <https://doi.org/10.5194/amt-16-2353-2023>, 2023.
- Glantz, P., Nilsson, E. D., and von Hoyningen-Huene, W.: Estimating a relationship between aerosol optical thickness and
670 surface wind speed over the ocean, *Atmos. Res.*, 92, 58–68, <https://doi.org/10.1016/j.atmosres.2008.08.010>, 2009.
- Grandey, B. S., Stier, P., Wagner, T. M., Grainger, R. G., and Hodges, K. I.: The effect of extratropical cyclones on satellite-retrieved aerosol properties over ocean, *Geophys. Res. Lett.*, 38, L13805, <https://doi.org/10.1029/2011GL047703>, 2011.
- Groß, S., Tesche, M., Freudenthaler, V., Toledano, C., Wiegner, M., Ansmann, A., Althausen, D. and Seefeldner, M.:
675 Characterization of Saharan dust, marine aerosols and mixtures of biomass-burning aerosols and dust by means of multi-wavelength depolarization and Raman lidar measurements during SAMUM 2, *Tellus B: Chemical and Physical Meteorology*, 63(4), 706–724, <http://doi.org/10.1111/j.1600-0889.2011.00556.x>, 2011
- Groß, S., Esselborn, M., Weinzierl, B., Wirth, M., Fix, A., and Petzold, A.: Aerosol classification by airborne high spectral resolution lidar observations, *Atmos. Chem. Phys.*, 13, 2487–2505, <https://doi.org/10.5194/acp-13-2487-2013>, 2013.
- Groß, S., Freudenthaler, V., Wirth, M., and Weinzierl, B.: Towards an aerosol classification scheme for future EarthCARE
680 lidar observations and implications for research needs, *Atmos. Sci. Lett.*, 16: 77–82, <https://doi.org/10.1002/asl2.524>, 2015.
- Haarig, M., Ansmann, A., Gasteiger, J., Kandler, K., Althausen, D., Baars, H., Radenz, M., and Farrell, D. A.: Dry versus wet marine particle optical properties: RH dependence of depolarization ratio, backscatter, and extinction from multiwavelength lidar measurements during SALTRACE, *Atmos. Chem. Phys.*, 17, 14199–14217, <https://doi.org/10.5194/acp-17-14199-2017>, 2017.
- 685 Haywood, J. M., V. Ramaswamy, and B. J. Soden: Tropospheric aerosol climate forcing in clear-sky satellite observations over the oceans, *Science*, 283, 1299–1303, <https://doi.org/10.1126/science.283.5406.1299>, 1999.
- Hoaglin, D. C., Iglewicz, B., and Tukey, J. W.: Performance of some resistant rules for outlier labelling, *Journal of the American Statistical Association*, 81(396), 991–999, <https://doi.org/10.1080/01621459.1986.10478363>, 1986.
- Huang, H., Thomas, G. E., and Grainger, R. G.: Relationship between wind speed and aerosol optical depth over remote ocean,
690 *Atmos. Chem. Phys.*, 10, 5943–5950, <https://doi.org/10.5194/acp-10-5943-2010>, 2010.
- IPCC, 2021: Summary for Policymakers. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy,

- J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 3-32, doi:10.1017/9781009157896.001.
- 695 Jaegle, L., Quinn, P. K., Bates, T., Alexander, B., and Lin, J.-T.: Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, *Atmos. Chem. Phys.*, 11, 3137-3157, <https://doi.org/10.5194/acp-11-3137-2011>, 2011.
- Kahn, R. A., Gaitley, B. J., Garay, M. J., Diner, D. J., Eck, T. F., Smirnov, A., and Holben, B. N.: Multiangle Imaging SpectroRadiometer global aerosol product assessment by comparison with the Aerosol Robotic Network, *J. Geophys. Res.*, 700 115, D23209, <https://doi.org/10.1029/2010JD014601>, 2010.
- Kanitz, T., Lochard, J., Marshall, J., McGoldrick, P., Lecrenier, O., Bravetti, P., Reitebuch, O., Rennie, M., Wernham, D., and Elfving, A.: Aeolus first light: first glimpse, International Conference on Space Optics–ICSO 2018, 111801R, Chania, Greece, 9–12 October 2018, <https://doi.org/10.1117/12.2535982>, 2019.
- Kaufman, Y. J., D. Tanre, and O. Boucher: A satellite view of aerosols in the climate system, *Nature*, 419, 215-223, 705 <https://doi.org/10.1038/nature01091>, 2002.
- Kiliyanpilakkil, V. P. and Meskhidze, N.: Deriving the effect of wind speed on clean marine aerosol optical properties using the A-Train satellites, *Atmos. Chem. Phys.*, 11, 11401–11413, <https://doi.org/10.5194/acp-11-11401-2011>, 2011.
- Kim, M.-H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Trepte, C. R., Hu, Y., Liu, Z., Poole, L. R., Pitts, M. C., Kar, J., and Magill, B. E.: The CALIPSO version 4 automated aerosol classification and lidar ratio selection algorithm, 710 *Atmos. Meas. Tech.*, 11, 6107–6135, <https://doi.org/10.5194/amt-11-6107-2018>, 2018.
- Kleidman, R. G., Smirnov, A., Levy, R. C., Mattoo, S., and Tanre, D.: Evaluation and wind speed dependence of MODIS aerosol retrievals over open ocean, *IEEE T. Geosci. Remote*, 50, 429-435, <https://doi.org/10.1109/TGRS.2011.2162073>, 2012.
- Latham, J., and M. H. Smith: Effect on global warming of wind-dependent aerosol generation at the ocean surface, *Nature*, 347, 372-373, <https://doi.org/10.1038/347372a0>, 1990.
- 715 Lehahn, Y., Koren, I., Boss, E., Ben-Ami, Y., and Altaratz, O.: Estimating the maritime component of aerosol optical depth and its dependency on surface wind speed using satellite data, *Atmos. Chem. Phys.*, 10, 6711-6720, <https://doi.org/10.5194/acp-10-6711-2010>, 2010.
- Lewis, R. and Schwartz, E.: Sea salt aerosol production: mechanisms, methods, measurements and models – a critical review, American Geophysical Union, <https://doi.org/10.1029/GM152>, 2004.
- 720 Liu, Z., Kar, J., Zeng, S., Tackett, J., Vaughan, M., Avery, M., Pelon, J., Getzewich, B., Lee, K.-P., Magill, B., Omar, A., Lucker, P., Trepte, C., and Winker, D.: Discriminating between clouds and aerosols in the CALIOP version 4.1 data products, *Atmos. Meas. Tech.*, 12, 703–734, <https://doi.org/10.5194/amt-12-703-2019>, 2019.
- Luo, T., Yuan, R., and Wang, Z.: Lidar-based remote sensing of atmospheric boundary layer height over land and ocean, *Atmos. Meas. Tech.*, 7, 173–182, <https://doi.org/10.5194/amt-7-173-2014>, 2014.
- 725 Luo, T., Wang, Z., Zhang, D., and Chen, B.: Marine boundary layer structure as observed by A-train satellites, *Atmos. Chem. Phys.*, 16, 5891–5903, <https://doi.org/10.5194/acp-16-5891-2016>, 2016.

- Madry, W. L., Toon, O. B., and O'Dowd, C. D.: Modeled optical thickness of sea salt aerosol, *J. Geophys. Res.*, 116, D08211, <https://10.1029/2010JD014691>, 2011.
- 730 Masonis, S. J., Anderson, T. L., Covert, D. S., Kapustin, V., Clarke, A. D., Howell, S., and Moore, K.: A study of the extinction-to-backscatter ratio of marine aerosol during the Shoreline Environment Aerosol Study, *J. Atmos. Ocean. Tech.*, 20, 1388–1402, [https://10.1175/1520-0426\(2003\)020<1388:ASOTER>2.0.CO;2](https://10.1175/1520-0426(2003)020<1388:ASOTER>2.0.CO;2), 2003.
- Meskhidze, N. and Nenes, A.: Effects of ocean ecosystem on marine aerosol-cloud interaction, *Adv. Meteorol.*, 2010, 239808, <https://10.1155/2010/239808>, 2010.
- 735 Mulcahy, J. P., O'Dowd, C. D., Jennings, S. G., and Ceburnis, D.: Significant enhancement of aerosol optical depth in marine air under wind conditions, *Geophys. Res. Lett.*, 35, L16810, <https://10.1029/2008GL034303>, 2008.
- Murphy, D. M., Anderson, J. R., Quinn, P. K., McInnes, L. M., Brechtel, F. J., Kreidenweis, S. M., Middlebrook, A. M., Pósfai, M., Thomson, D. S., and Buseck, P. R.: Influence of sea-salt on aerosol radiative properties in the Southern Ocean marine boundary layer, *Nature*, 392, 62-65, <https://doi.org/10.1038/32138>, 1998.
- O'Dowd, C. D. and de Leeuw, G.: Marine aerosol production: a review of the current knowledge, *Phil. Trans. R. Soc.*, 365, 740 1753-1774, <https://doi.org/10.1098/rsta.2007.2043>, 2007.
- O'Dowd, C. D., Lowe, J. A., and Smith, M. H.: Coupling sea-salt and sulphate interactions and its impact on cloud droplet concentration predications, *Geophys. Res. Lett.*, 26, 1311-1314, <https://10.1029/1999GL900231>, 1999.
- O'Dowd, C. D., Scannell, C., Mulcahy, J., and Jennings, S. G.: Wind speed influences on marine aerosol optical depth, *Adv. Meteorol.*, 2010, 830846, <https://10.1155/2010/830846>, 2010.
- 745 Pierce, J. R. and Adams, P. J.: Global evaluation of CCN formation by direct emission of sea salt and growth of ultrafine sea salt, *J. Geophys. Res.*, 111, D06203, <https://10.1029/2005JD006186>, 2006.
- Platt, C. M. R. and G. R. Patterson: The interpretation of baseline atmospheric turbidity measurements at Cape Grim, Tasmania, *Journal of Atmospheric Chemistry*, 4, 187-197, <https://10.1007/BF00053778>, 1986.
- Prijith, S. S., Aloysius, M., and Mohan, M.: Relationship between wind speed and sea salt aerosol production: A new approach, 750 *Journal of Atmospheric and Solar-Terrestrial Physics*, 108, 34-40, <https://10.1016/j.jastp.2013.12.009>, 2014.
- Reitebuch, O.: The spaceborne wind lidar mission ADM-Aeolus, in: *Atmospheric Physics*, edited by: Schumann, U., Springer, 815-827, ISBN 978-3-642-30182-7, https://doi.org/10.1007/978-3-642-30183-4_49, 2012.
- Rennie, M., Tan, D., Andersson, E., Poli, P., Dabas, A., de Kloe, J., Marseille, G., and Stoffelen, A.: Aeolus Level-2B Algorithm Theoretical Basis Document, version 3.40, <https://earth.esa.int/eogateway/documents/20142/37627/Aeolus-L2B-Algorithm-ATBD.pdf> (last access: 7 November 2022), 2020.
- 755 Rennie, M. P., Isaksen, L., Weiler, F., de Kloe, J., Kanitz, T., and Reitebuch, O.: The impact of Aeolus wind retrievals on ECMWF global weather forecasts, *Q. J. Roy. Meteor. Soc.*, 147, 3555–3586, <https://doi.org/10.1002/qj.4142>, 2021.
- Sayer, A. M., Smirnov, A., Hsu, N. C., and Holben, B. N.: A pure marine aerosol model, for use in remote sensing applications, *J. Geophys. Res.*, <https://10.1029/2011JD016689>, 2012.

- 760 Shin, D. H., Müller, D., Choi, T., Noh, Y. M., Yoon, Y. J., Lee, K. H., Shin, S. K., Chae, N., Kim, K., and Kim, Y. J.: Influence of wind speed on optical properties of aerosols in the marine boundary layer measured by ship-borne DePolarization Lidar in the coastal area of Korea, *Atmospheric Environment*, 47, 282-290, <https://doi.org/10.1016/j.atmosenv.2013.10.027>, 2014.
- Shinozuka, Y., Clarke, A. D., Howell, S. G., Kapustin, V. N., and Huebert, B. J.: Sea-salt vertical profiles over the Southern and tropical Pacific oceans: Microphysics, optical properties, spatial variability, and variations with wind speed, *J. Geophys. Res.*, 109, D24201, <https://doi.org/10.1029/2004JD004975>, 2004.
- 765 Smirnov, A., Villevalde, Y., O'Neill, N. T., Royer, A., and Tarussov, A.: Aerosol optical depth over the oceans: Analysis in terms of synoptic air mass types, *J. Geophys. Res.*, 100, 16639-16650, <https://doi.org/10.1029/95JD01265>, 1995.
- Smirnov, A., Holben, B. N., Eck, T. F., Dubovik, O., and Slutsker, I.: Effect of wind speed on columnar aerosol optical properties at Midway Island, *J. Geophys. Res.*, 108, 4802, <https://doi.org/10.1029/2003JD003879>, 2003.
- 770 Smirnov, A., Sayer, A. M., Holben, B. N., Hsu, N. C., Sakerin, S. M., Macke, A., Nelson, N. B., Courcoux, Y., Smyth, T. J., Croot, P., Quinn, P. K., Sciare, J., Gulev, S. K., Piketh, S., Losno, R., Kinne, S., and Radionov, V. F.: Effect of wind speed on aerosol optical depth over remote oceans, based on data from the Maritime Aerosol Network, *Atmos. Meas. Tech.*, 5, 377-388, <https://doi.org/10.5194/amt-5-377-2012>, 2012.
- Stoffelen, A., Pailleux, J., Källén, E., Vaughan, J. M., Isaksen, L., Flamant, P., Wergen, W., Andersson, E., Schyberg, H., and
775 Culoma, A.: The atmospheric dynamics mission for global wind field measurement, *B. Am. Meteorol. Soc.*, 86, 73-88, <https://doi.org/10.1175/BAMS-86-1-73>, 2005.
- Tan, D. G., Andersson, E., Kloe, J. D., Marseille, G.-J., Stoffelen, A., Poli, P., Denneulin, M.-L., Dabas, A., Huber, D., and Reitebuch, O.: The ADM-Aeolus wind retrieval algorithms, *Tellus A*, 60, 191-205, <https://doi.org/10.1111/j.1600-0870.2007.00285.x>, 2008.
- 780 Reitebuch, O.: The Spaceborne Wind Lidar Mission ADM-Aeolus, in: *Atmospheric Physics, Research Topics in Aerospace*, edited by: Schumann, U., ISBN 978-3-642-30182-7, Springer-Verlag Berlin Heidelberg, 815-827, https://doi.org/10.1007/978-3-642-30183-4_49, 2012.
- Traou, D., Flament, T., Lacour, A. and Stieglitz, H.: L2A user guide, issue V 2.1, [https://earth.esa.int/eogateway/documents/20142/37627/Aeolus-Data-Innovation-Science-Cluster-DISC-Level-2A-user-](https://earth.esa.int/eogateway/documents/20142/37627/Aeolus-Data-Innovation-Science-Cluster-DISC-Level-2A-user-guide.pdf)
785 [guide.pdf](https://earth.esa.int/eogateway/documents/20142/37627/Aeolus-Data-Innovation-Science-Cluster-DISC-Level-2A-user-guide.pdf) (last access: 27 February 2023), 2022.
- Villevalde, Yu. V., Smirnov, A. V., O'Neill, N. T., Smyshlyayev, S. P., and Yakovlev, V. V.: Measurement of aerosol optical depth in the Pacific Ocean and the North Atlantic, *J. Geophys. Res.*, 99, 20983-20988, <https://doi.org/10.1029/94JD01618>, 1994.
- Wilson, S. R., and Forgan, B. W.: Aerosol optical depth at Cape Grim, Tasmania, 1986-1999, *J. Geophys. Res.*, 107, D8, <https://doi.org/10.1029/2001JD000398>, 2002.
- 790 Winker, D. M. and Pelon, J.: The CALIPSO mission, *Geoscience and Remote Sensing Symposium, IGARSS '03, Proceedings, IEEE International*, 2, 1329-1331, <https://doi.org/10.1109/IGARSS.2003.1294098>, 2003.

Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the CALIPSO mission and CALIOP data processing algorithms, *J. Atmos. Ocean. Tech.*, 26, 2310–2323, <https://doi.org/10.1175/2009JTECHA1281.1>, 2009.

795 Wu, S., Sun, K., Dai, G., Wang, X., Liu, X., Liu, B., Song, X., Reitebuch, O., Li, R., Yin, J., and Wang, X.: Inter-comparison of wind measurements in the atmospheric boundary layer and the lower troposphere with Aeolus and a ground-based coherent Doppler lidar network over China, *Atmos. Meas. Tech.*, 15, 131–148, <https://doi.org/10.5194/amt-15-131-2022>, 2022.

Zibordi, G., Berthon, J.-F., Melin, F., and D’Alimonte, D.: Cross-site consistent in situ measurements for satellite ocean color applications: The BiOMaP radiometric dataset, *Remote Sens. Env.*, 115, 2104–2115, <https://doi.org/10.1016/j.rse.2011.04.013>,

800 2011.