

Microwave radiometer observations of the ozone diurnal cycle and its short-term variability over Switzerland

Eric Sauvageat^{1,2}, Klemens Hocke^{1,2}, Eliane Maillard Barras³, Shengyi Hou¹, Quentin Errera⁴, Alexander Haefele³, and Axel Murk^{1,2}

¹Institute of Applied Physics, University of Bern, Bern, Switzerland

²Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

³Federal Office of Meteorology and Climatology MeteoSwiss, Payerne, Switzerland

⁴Royal Belgian Institute for Space Astronomy, BIRA-IASB, Brussels, Belgium

Correspondence: Eric Sauvageat (eric.sauvageat@unibe.ch)

Abstract. In Switzerland, two ground-based ozone microwave radiometers are operated in the vicinity of each other (ca. 40 km): GROMOS in Bern (Institute of Applied Physics) and SOMORA in Payerne (MeteoSwiss). Recently, their calibration and retrieval algorithms have been fully harmonized and updated time series are now available since 2009. Using these harmonized ozone time series, we investigate and cross-validate the strato-mesospheric ozone diurnal cycle derived from the two instruments and compare it with various model-based datasets: a dedicated GDOC ozone diurnal cycle climatology based on the Goddard Earth Observing System (GEOS-5) general circulation model, the Belgian Assimilation System for Chemical Observations (BASCOE) a chemical transport model driven by ERA5 dynamics, and a set of free-running simulations from the Whole Atmosphere Community Climate Model (WACCM). Overall, the two instruments show very similar ozone diurnal cycles at all seasons and pressure levels and the models compare well with each other. There is a good agreement between the models and the measurements at most seasons and pressure levels and the largest discrepancies can be explained by the limited vertical resolution of the microwave radiometers. However, as in a similar study over Mauna Loa, some discrepancies remain near the stratopause, at the transition region between ozone daytime accumulation and depletion. We report similar delays in the onset of the modelled ozone diurnal depletion in the lower mesosphere.

Using the newly harmonized time series of GROMOS and SOMORA radiometers, we present the first observations of short-term (sub-monthly) ozone diurnal cycle variability at mid-latitudes. The short-term variability is observed in the upper stratosphere during wintertime, when the mean monthly cycle has a small amplitude and when the dynamics is more important. It is shown in the form of strong enhancements of the diurnal cycle, reaching up to 4–5 times the amplitude of the mean monthly cycle. We show that BASCOE is able to capture some of these events and present a case study of one such event following the minor sudden stratospheric warming of January 2015. Our analysis of this event supports the conclusions of a previous modelling study, attributing regional variability of the ozone diurnal cycle to regional anomalies in nitrogen oxide (NO_x) concentrations. However, we also find period with enhanced diurnal cycle that do not show much change in NO_x and where other processes might be dominant (e.g. atmospheric tides). Given its importance, we believe that the short-term variability of the ozone diurnal cycle should be further investigated over the globe, for instance using the BASCOE model.

1 Introduction

25 Beyond its role in the earth protection from harmful ultraviolet radiation, ozone is a key species in the energy balance of the middle atmosphere, strongly influencing the radiation budget and thermal state of the stratosphere and mesosphere. Following the success of the Montreal Protocol in 1987, full recovery of the ozone layer is expected for the 21st century with significant regional variability and uncertainties. In particular, there is a high degree of uncertainty about the lower stratospheric ozone recovery and there are increasing observational evidence that ozone is still declining at some locations in the lower stratosphere
30 (Ball et al., 2018; Maillard Barras et al., 2022a; Godin-Beekmann et al., 2022), without satisfactory explanation to date.

Ozone is a very reactive molecule involved in many (photo-)chemical reactions in the middle atmosphere. The set of pure oxygen photochemical reactions leading to the production and destruction of ozone were first described by Chapman (1930) and are known as the Chapman cycle. Together with catalytic depletion cycles involving many different species (NO_y , Cl_y , HO_y , ...), they mostly drive the ozone amount in the middle atmosphere at multiple time scales. In particular, ozone **concentration**
35 concentrations are subject to complex diurnal cycle patterns depending on the geographic location, altitude, season and other factors (Schanz et al., 2014), which makes it both important and difficult to fully take into account in models or observations.

The importance of the ozone diurnal cycle in the mesosphere has been recognized early through the use of photochemical models and from early measurements (e.g. Prather, 1981; Vaughan, 1982; Pallister and Tuck, 1983; Zommerfelds et al., 1989). Its main patterns are well known and have been successfully observed and modelled (Connor et al., 1994; Ricaud et al., 1991;
40 Huang et al., 1997). In the stratosphere though, the ozone diurnal cycle is much weaker which makes accurate observations challenging. However, it needs to be taken into account when comparing different observing systems or to compute accurate trends (Bhartia et al., 2013; Maillard Barras et al., 2020). In recent years, there has been a renewed interest to improve the consideration of ozone and nitrogen oxides diurnal variations in satellite measurements (Frith et al., 2020; Schanz et al., 2021; Strode et al., 2022), partly because of the remaining uncertainties on stratospheric ozone post-2000 trends. For instance, Frith
45 et al. (2020) used a modified version of the GEOS-5 model to produce a global, zonally averaged ozone diurnal climatology: the GEOS-GMI Diurnal Ozone Climatology (GDOC). The idea was to publish an easy-to-use climatology of scaling factors to account for ozone diurnal variability in intercomparison studies or for the creation of merged ozone dataset. More recently, Strode et al. (2022) develops similar year-specific scaling factors for comparisons with the SAGE III/ISS measurements.

To validate such diurnal scaling factors, accurate observations are needed at different altitudes and locations. However, such
50 observations remain relatively sparse and challenging, especially in the stratosphere where the ozone diurnal cycle amplitude is small. Also, many satellites are sun-synchronous or use the sun as source which limits their ability to derive full diurnal cycle. Some satellite-based ozone diurnal cycle have been successfully derived however, from SAGE/ISS (Sakazaki et al., 2013, 2015) and from SABER on the Thermosphere Ionosphere Mesosphere Energetics and Dynamics (TIMED) satellite (Huang et al., 2010a). Whereas the satellite-based observations offer a global view on the ozone diurnal cycle, they do need to
55 aggregate data in space or time to derive diurnal cycle, therefore blurring any short-term or regional fluctuations in the cycle amplitude.

Passive microwave ground-based radiometers (MWRs) are well suited for ozone diurnal cycle observations because they operate continuously and do not use the sun as a source. These instruments have been used successfully by different groups to monitor the diurnal cycle, not only in the mesosphere but also in the stratosphere (Connor et al., 1994; Schneider et al., 2005; Haeferle et al., 2008; Parrish et al., 2014; Studer et al., 2014; Schranz et al., 2018). In the tropics, Parrish et al. (2014) derived detailed stratospheric ozone diurnal cycle over Mauna Loa from MWR measurements and compared it with satellite measurements and the Goddard Earth Observing System Chemistry Climate Model (GEOSCCM). They found a good agreement between the MWR and satellite observations and remaining discrepancies with the model in the upper stratosphere (3.2 to 1.8 hPa). In the polar region, Schranz et al. (2018) found larger discrepancies between MWR measurements and SD-WACCM simulations over Ny-Ålesund but only focused on a single year of measurements ~~though~~.

In this contribution, we derive updated ozone diurnal cycles over Switzerland from two collocated (ca. 40 km) ground-based MWRs between 2010 and 2022. The time series have recently been fully reprocessed with a harmonized algorithm (Sauvageat et al., 2022b) and provide a unique set of measurements to study the ozone diurnal cycle and validate model simulations over the mid-latitudes. Compared to the three previous studies on ozone diurnal cycle over Switzerland (Zommerfelds et al., 1989; Haeferle et al., 2008; Studer et al., 2014), this study combines the two MWRs over an extended time period (> 10 years) and focuses on the time where the two instruments used the same digital spectrometer (after 2009). The combination of the spectrometer update and of the recent harmonization extended the altitude range and improved the sensitivity of the ozone retrievals. We obtain significant improvements in the updated stratospheric ozone diurnal cycle ~~of GROMOS, which was strongly overestimated~~ measured by the GROUND-based Millimeter-wave Ozone Spectrometer (GROMOS) in Bern. In fact, the study from (Studer et al., 2014) showed overestimated diurnal cycle amplitude compared to the model simulations ~~in the last study (Studer et al., 2014). Although it was focused on the time period where GROMOS used a filter bank spectrometer (before 2009), we believe that part of the discrepancies were also due to the retrieval algorithm~~. Given that this instrument provided one of the main reference for ozone diurnal cycle comparison studies over the mid-latitudes in the last decade, we believe that it is highly valuable to present in details these updated results.

In fact, the objective of the present study is multiple. First we use the harmonized time series to derive updated diurnal cycle above Switzerland and provide a comparative basis for different models at mid-latitude. Especially, we aim at providing an additional validation for the dedicated GEOS-GMI Diurnal Ozone Climatology (GDOC), which has been published for use as data analysis tool. In addition, we compare our measurements with two other types of model-based datasets: the Belgian Assimilation System for Chemical Observations from Envisat (BASCOE) ~~chemistry-transport~~ chemistry transport model (CTM) and the Whole Atmosphere Community Climate Model (WACCM) ~~chemistry-climate~~ chemistry climate model (CCM). Finally, we present the first observations of short-term (sub-monthly) ozone diurnal variability and investigate the causes for such variations. We use the global, high-resolution simulations of BASCOE coupled with reanalysis data from ERA5 (Hersbach et al., 2020) to cross-validate our observations. We discuss a case study of the winter 2014-2015 and provide other examples where short-term fluctuations of the ozone diurnal cycle was observed.

The publication is organised as follow: Sect. 2 introduces the datasets and the methods used to compute the ozone diurnal cycle. Section 3 presents the results including the monthly ozone profile comparisons (Sect. 3.1), the intercomparisons of the

monthly averaged ozone diurnal cycle over Bern (Sect. 3.2), and an example of observed short-term variability during the boreal winter 2014-2015 (Sect. 3.3). Finally, Sect. 4 presents a summary of the main results and some conclusions.

2 Materials and Methods

95 In the following, we present succinctly the datasets and the methods used in our study. Regarding the datasets, we focus mostly on the microwave radiometer time series and on the main model characteristics, and provide references to the reader for additional details. Also, we summarise the most important features and relevant publications for each dataset in Table 1.

2.1 Microwave ground-based radiometers

Microwave ground-based radiometers (MWRs) are passive remote sensing instruments that can be used to derive trace gas or
100 temperature profiles in the atmosphere. MWRs measure the emission of atmospheric molecules in the microwave frequency range. Therefore, they do not rely on the sun for their observations while providing quite high temporal resolution and continuous sampling, which makes them excellent candidates for diurnal cycle studies.

In Switzerland, two ozone microwave radiometers are operated close to each other (ca. 40 km) on the Swiss Plateau. The
GROund-based Millimeter-wave Ozone Spectrometer (GROMOS) is operated by the Institute of Applied Physics (IAP) at the
105 University of Bern (46.95° N, 7.44° E, 560 masl) since 1994 and the Stratospheric Ozone MONitoring RAdiometer (SOMORA)
is operated by the Federal Office of Meteorology and Climatology MeteoSwiss in Payerne (46.82° N, 6.94° E, 491 masl) since
2000. The two instruments have been designed at the IAP, have similar design and use the rotational ozone emission line at
142.175 GHz to derive strato-mesospheric ozone profiles. Also, they have similar viewing geometries, both observe the sky at
~ 40° elevation angle and experience similar atmospheric opacity conditions. Following discrepancies identified between the
110 two instruments (Bernet et al., 2019; SPARC/IO3C/GAW, 2019), a complete harmonization of the data processing has recently
been performed for GROMOS and SOMORA. It resulted in harmonized, continuous, hourly time series of strato-mesospheric
ozone starting in 2009 and which are now freely available (Sauvageat et al., 2022b).

The vertical resolution of the MWRs is quite coarse (~ 10 km up to 3 hPa and ~ 15 km above) and the vertical extent of
the ozone profile is from 60 to 0.02 hPa (~ 20 to 75 km), corresponding to the range where the a-priori contribution to the
115 retrieved profile is lower than 20 % (Fig. 1). The MWRs coarser vertical resolution needs to be taken into account for inter-
comparison with higher resolution datasets (e.g. models), also for the diurnal cycle comparisons. The usual way is to apply a
smoothing procedure to the higher resolution dataset for the comparisons. In our study, we use the classical “averaging kernel
smoothing” which essentially convolve the high resolution dataset with the averaging kernels (AVKs) of the MWR retrieval
using Equation 1 (Rodgers and Connor, 2003). Equation 1 also applies the effect of the a-priori contribution of the MWR
120 retrievals onto the higher resolved profile and is usually expressed as:

$$\mathbf{x}_c = \mathbf{x}_a + \mathbf{A}(\mathbf{x} - \mathbf{x}_a) \quad (1)$$




Figure 1. Mean averaging kernels and measurement contribution for **(a)** GROMOS and **(b)** SOMORA. The black lines are the averaging kernel at individual pressure levels whereas the color lines are the respective measurement contribution (see upper x-axis). The shaded color area shows the standard deviation of the measurement contribution.

with x_a the a-priori profile (derived from monthly WACCM profiles in our case), A the averaging kernel matrix, x and x_c and x and x_c respectively the original and convolved high resolution profile.

GROMOS and SOMORA essentially have the same sensitivity, allowing to compare their observations directly. It can be seen by looking at the mean AVKs and the measurement contribution of the retrievals shown in Fig. 1. It also means that there is only little difference whether we use GROMOS or SOMORA AVKs for the smoothing procedure. In the following, all convolutions on the higher resolution profiles are performed using the GROMOS AVKs. Figure 1 shows the mean AVKs of the full GROMOS and SOMORA time series, however, for all the convolutions we use the appropriate monthly daytime or nighttime AVKs.

130 2.1.1 Satellite measurements

For validation purpose, we use measurements from the Microwave Limb Sounder (MLS) mounted on the Aura spacecraft (Waters et al., 2006). Since its launch in July 2004, the MLS instrument has been used extensively for trace gas observations

and is one of the main measurement reference for global ozone monitoring studies. More specifically, we use the latest ozone retrieval product (v5), following the screening guidelines provided in Livesey et al. (2022). The MLS ozone vertical resolution ranges from ~ 2.5 to ~ 5 km in the stratosphere and mesosphere whereas its horizontal resolution ranges between 300 and 500 km. As spatial collocation criteria, we keep only measurements around Switzerland ($\pm 1.8^\circ$ in latitude and $\pm 5^\circ$ longitude). Aura overpasses Switzerland twice a day, at 02:20 LST and 13:10 LST, thus providing the day-to-night ozone ratios but not the full diurnal cycle. In Sect. 3.3, we also show some measurements of temperature and nitrous oxide from MLS, however, as higher temporal resolution was needed for the short-term analysis, these were obtained with more relaxed collocation criteria ($\pm 3.6^\circ$ latitude and $\pm 10^\circ$ longitude) but following the same screening guidelines (Livesey et al., 2022).

2.2 Model-based datasets

2.2.1 GDOC

The GEOS-GMI Diurnal Ozone Climatology (GDOC) is a model-based climatology of ozone diurnal cycle derived from the NASA Goddard Earth Observing System general circulation model, version 5 (GEOS-5). The goal of this climatology is to provide a simple data analysis tool to account for ozone diurnal variability, e.g. when comparing different satellite profiles.

For the production, GEOS-5 was run in replay mode constrained to 3-hourly MERRA-2 assimilated meteorological fields from January 2017 to December 2018 (see Orbe et al. (2017); Frith et al. (2020) and references therein for model details). As final product, the GDOC provides zonally averaged (on 5° latitude bands) ozone diurnal cycle from 90 to 0.3 hPa (~ 20 to 55 km) with equivalent vertical resolution of ~ 1 km and a time resolution of 30 minutes. The climatology is also available (on request) on original model levels but has not been evaluated below 30 and above 0.3 hPa, which is the reason why we chose not to use it outside of this pressure range.

It provides monthly climatological ozone values as a function of local solar time (LST) normalized by-to midnight ozone values. The GDOC does not contain the original ozone profiles which prevents the application of the averaging kernel smoothing procedure on this dataset. Consequently, we only show the original high resolution profile from the GDOC dataset.

2.2.2 BASCOE

~~The Belgian Assimilation System for Chemical Observations (BASCOE) is a~~ This study uses chemistry transport model (CTM) ~~developed at the Royal Belgian Institute for Space Aeronomy (BIRA-IASB) (Errera and Fonteyn, 2001; Errera et al., 2008)~~ simulation performed by the Belgian Assimilation System for Chemical Observations (BASCOE, Errera and Fonteyn (2001); Errera et al. (2008)). The simulation covers the 2010-2020 period and is driven by 6-hourly wind and temperature fields taken from ERA5 (Hersbach et al., 2020) preprocessed in a similar setup as described in Chabrilat et al. (2018) ~~. BASCOE dynamics relies on 6-hourly winds and temperature taken from ERA5, the latest reanalysis dataset from the European Center for Meteorological and Weather Forecast (ECMWF). The model has~~ and Minganti et al. (2022). The simulation is performed with a time resolution of 30 minutes, ~~and~~

165 ~~a spatial~~ a horizontal resolution of 2° in latitude and 2.5° in longitude. ~~It runs on~~, and 42 hybrid pressure levels from the
surface to ~~the mesosphere (~ 0.01)~~ 0.01 hPa with a vertical resolution ~~of $\sim 1-4$ in the stratosphere and~~ between 1 km in
the lower stratosphere and 4 km in the lower mesosphere. The BASCOE model focuses on the calculation of the chemical
composition of the stratosphere. It includes around 60 chemical species interacting through around 200 reactions (gas phase,
photolysis and heterogeneous) and a parametrization to account for the effect of sulfate aerosols and polar stratospheric
170 clouds on the stratospheric composition (Huijnen et al., 2016). Background sulfate aerosols are taken from the Climate Model
Intercomparison Project Phase 6 (CMIP6) recommendations whereas surface emissions of long-lived species are taken from the
‘Historical Greenhouse Gas Concentrations’ (HGGC) recommendations also produced for CMIP6 experiments (Meinshausen et al., 2017)
. The model provides a realistic composition of the stratosphere when compared to independent observations (see e.g., Prignon et al. (2021);
). For this study, the model outputs have been interpolated at the location of Bern so that they can be compared with ~~our~~ the
175 two MWRs.

2.2.3 WACCM

We also use results from the Whole Atmosphere Community Climate Model (WACCM), version 4 in the configuration de-
scribed by Schanz et al. (2021). WACCM is a fully coupled global ~~chemistry-climate~~ chemistry climate model developed at
the National Center for Atmospheric Research (NCAR) with a stratospheric chemistry module based on the Model of Ozone
180 and Related Chemical Tracers (MOZART) (Kinnison et al., 2007). It simulates the atmosphere from the surface to ~ 150 km
altitude, with ~~vertical resolution ranging a~~ vertical resolution between 1.1 and 2 km in the middle-atmosphere. WACCM was
run with a horizontal resolution of 4° latitude by 5° longitude and for our study, we use the closest grid point to both Bern and
Payerne, which corresponds to 46° N and 5° E. The model was run with the pre-defined free-running “F 2000” scenario,
simulating a perpetual year 2000 but without data nudging.

185

Table 1. Summary of the datasets used in this study.

Dataset	Type	Coverage (horizontal, vertical)	Reference
GROMOS	MWR measurement	local, 60 – 0.02 hPa	Sauvageat et al. (2022b)
SOMORA	MWR measurement	local, 60 – 0.02 hPa	Sauvageat et al. (2022b)
MLS	Limb sounding measurement	global, 261 – 0.001 hPa	Waters et al. (2006); Froidevaux et al. (2008)
GDOC	Model based climatology	zonal, 30 – 0.3 hPa	Frith et al. (2020)
BASCOE	CTM	global, surface–0.05 hPa	Errera et al. (2008)
WACCM	CCM, free-running	global, surface– $5.1 \cdot 10^{-6}$ hPa	Garcia et al. (2007); Marsh et al. (2013)
ERA5	Reanalysis	global, surface–0.05 hPa	Hersbach et al. (2020)

2.3 Ozone profiles and day-to-night ratios

Before computing the ozone diurnal cycle from our different datasets, we first compare their monthly averaged ozone profiles during daytime and nighttime and compute their day-to-night ratios. We compare our two MWRs with MLS and ~~the BASCOE simulation~~ BASCOE, all averaged on the time period 2010-2020, therefore removing most of the year-to-year variability. WACCM is not included in these comparisons for two reasons. First, we use the free-running WACCM simulations for a single year, and it would make no sense to compare it with the other datasets which are multi-year averages, especially in the wintertime when the dynamics is an important modulator of the ozone amount. Second, the monthly averaged WACCM ozone profiles are actually used as a-priori for our MWR retrievals and therefore, they can not be used for validation against the retrieved MWR profiles.

For the ozone profile comparisons, we ~~use the approximate~~ choose to use profiles whose time corresponds approximately to the MLS overpass times ~~, keeping all (02:20 LST and 13:10 LST).~~ Therefore, we keep timestamps between 12:00 and 15:00 LST for the daytime profiles and between 01:00 and 04:00 LST for the nighttime profiles, regardless of the dataset. As explained in Sect. 2.1, we convolve the BASCOE simulations and the MLS measurements with monthly averaged daytime or nighttime GROMOS AVKs.

2.4 Ozone diurnal cycle

Similarly to the GDOC climatology, we choose to express the ozone diurnal cycle as ratio of ozone ~~VMR~~ volume mixing ratios (VMRs) relative to the midnight value. To compute the reference midnight value ($O_{3,NT}$ in Eq. 1), we have used different time periods to ~~reflects~~ reflect the different time resolution of the datasets. For the MWRs (~ 1 h time resolution), we compute the midnight reference value by taking an average of 2 nighttime measurements between 23:00 LST and 01:00 LST. For WACCM

205 and BASCOE (30 minutes time resolution), we use measurement between 00:00 and 01:00 LST, whereas the GDOC was normalized to the values between 23:45 to 00:15 LST [Frith et al. \(2020\)](#) ([Frith et al., 2020](#)).

For each hour and at each pressure level, we then compute the ratios to ozone at midnight using Equation 2. To simplify the notation, we do not explicitly write the pressure dependence of all the terms.

$$\Delta O_3(h) = \frac{O_3 - O_{3,NT}}{O_{3,NT}} \quad (2)$$

210 To compare with the monthly GDOC climatology, we compute monthly averaged $\Delta O_3(h)$ from GROMOS, SOMORA, WACCM and BASCOE. For each dataset, we use the available time series, i.e. 12 years of data for the two MWRs (2010-2022), 10 years for BASCOE (2010-2020), and the 1 year of the WACCM free-running model run. For the MWRs, we additionally filtered the time series to remove the measurements done at very high tropospheric opacity ($\tau > 1.5$) as they result in lower quality retrievals and can potentially contaminate the monthly averages. To some extent, it also helps to limit any seasonal bias
215 arising due to the summertime higher opacity although it is difficult to rule out this effect completely (e.g. see discussion on the effect of the opacity on GROMOS and SOMORA in Sauvageat et al. (2022b)). For the diurnal cycle, the effect of this filtering is not very large and for the interested reader, the unfiltered version is provided in the supplementary material (Figures S5 to S16).

We compute errors on the MWRs and BASCOE ozone diurnal cycles as standard error of the mean (SEM). For each month
220 and LST hour, we compute the standard deviation of $\Delta O_3(h)$ and divide it by the square root of the number of ozone profiles available for each hour.

2.5 Short-term variability of the ozone diurnal cycle

In addition to monthly averaged ozone diurnal cycle, we also show observations of short-term (sub-monthly) variability of
the ozone diurnal cycle. GROMOS and SOMORA provide a unique setup for short-term ozone diurnal cycle **observation**
225 **observations** because they have continuous, hourly, collocated measurements. Therefore, we can use them to compute the ozone diurnal cycle on sub-monthly period and cross-validate their measurements. Also, we use BASCOE simulations to compute the short-term variability of the ozone diurnal cycle over Switzerland and to investigate the cause of such variability.

In order to detect sub-monthly variations in the ozone diurnal cycle, we computed the day-to-night differences in ozone VMR for GROMOS, SOMORA and BASCOE. More specifically, we compute the anomalies of the day-to-night differences
230 (D_{O_3}) to a monthly climatology. As daily anomalies are too noisy, we average these differences on 5 days.

$$D_{O_3} = O_{3,DT} - O_{3,NT} \text{ [ppmv]} \quad (3)$$

In this contribution, we focus on the winter 2014-2015 and use BASCOE and MLS data to investigate and discuss potential reasons explaining a specific event during this winter. For further studies, we provide along with this publication the full time series of D_{O_3} daily anomalies for GROMOS, SOMORA and BASCOE.

3.1 Monthly ozone profiles and day-to-night ratio

The comparisons of the monthly averaged ozone daytime and nighttime profiles and day-to-night ratios are shown in Fig. 2 for December and Fig. 3 for June as proxies for the winter and summer season. Figure A1 and Fig. A2 in Appendix A show similar comparisons for March and September respectively. Similar comparisons but with respect to SOMORA MWR can be
 240 seen in the supplementary material (Figures S1 to S4).

~~(a) Monthly averaged profiles, (b) relative differences $-(X-GRO)/GRO$, and (c) day-to-night ratios of ozone VMR above Switzerland in December. In (a) and (b), the solid lines are the daytime profiles whereas the dashed lines are the nighttime profiles. Same as Fig. 2 but for June.~~

Overall we find a good agreement between measured ozone profiles (GROMOS, SOMORA and MLS), with relative differ-
 245 ences between the measured ozone profiles lower than 10 % up to 0.1 hPa. The differences between BASCOE and the MWR are within 15 % between 30 and 0.2 hPa, with slightly bigger bias during the winter months. BASCOE notably underestimates ozone amounts in the upper stratosphere to lower mesosphere (up to ~ 0.5 hPa) and overestimates ozone in the lower stratosphere, regardless of the month or the time of day. Above 0.2 hPa, the differences depend on the month considered but the model tends to underestimate the daytime ozone profiles, leading to a small overestimation of the day-to-night depletion
 250 ratio over 0.2 hPa. The ozone deficit of BASCOE in the middle atmosphere is consistent with previous studies (Skachko et al., 2016) and is a bias found in other models as well (see e.g. Fig. 3.1 in SPARC, 2010). One reason for this ozone deficit might be an overestimation of NO_2 in the model simulations, enhancing catalytic ozone destruction through the so-called "~~Crutzen~~" "Crutzen" cycle (Crutzen, 1970).

All datasets show the ozone daytime accumulation in the stratosphere and its transition to ozone daytime depletion around
 255 the stratopause ($\sim 2 - 1$ hPa). All datasets show stratospheric ozone accumulation during daytime (up to ~ 1 hPa) with stronger amplitude during the summertime, and they all show the strong mesospheric ozone depletion during daytime, growing in amplitude with altitude. They agree quite well on the pressure at which peak accumulation and peak depletion occur. During summer, there is an excellent agreement between day-to-night ratios up to ~ 0.6 hPa. Above this pressure level, the GDOC climatology systematically underestimates the daytime ozone loss, leading to less negative day-to-night ratio than the other
 260 datasets. In the mesosphere, BASCOE and MLS compare well with the MWRs up to 0.05 hPa, with day-to-night ratios in agreement within 15 %.

3.2 Comparison of monthly ozone diurnal cycle

Going beyond the day-to-night ratio, full monthly diurnal cycles over Switzerland are shown for summer and winter in Figures 4 and 5 for the 2 MWRs and the 3 model-based datasets (Figures B1 and B2 in Appendix B show similar results for spring and
 265 autumn respectively). These figures show the ratios to ozone midnight values as a function of the LST between 60 and 0.02 hPa (respectively 30 and 0.3 hPa for GDOC). As mentioned previously, the monthly averages correspond to different time periods for each dataset: 2010-2022 for GROMOS and SOMORA, 2010-2020 for BASCOE, 2017-2018 for the GDOC and 2000




Figure 2. (a) Monthly averaged profiles, (b) relative differences $((X-GRO)/GRO)$, and (c) day-to-night ratios of ozone VMR above Switzerland in December. In (a) and (b), the solid lines are the daytime profiles whereas the dashed lines are the nighttime profiles.




Figure 3. Same as Fig. 2 but for June.




Figure 4. Monthly averaged ozone diurnal cycle over Switzerland in December as seen in GROMOS, SOMORA, GDOC, BASCOE and WACCM datasets. Note that only the BASCOE and WACCM datasets have been convolved with the AVKs of GROMOS as explained in Sect. 2.1.

perpetual year for WACCM. Also, AVK smoothing procedures have been applied to the WACCM and BASCOE dataset but not to the GDOC. The original version without any AVK smoothing can be seen for all datasets in the supplementary material.

270 For better visualization, we also show these diurnal cycles averaged over 9 selected pressure ranges. This is shown in Figures 6 and 7 for winter and summer (in Figures B3 and B4 in Appendix B for spring and autumn). These figures show the original cycle from each dataset together with the AVK smoothed cycle, which enables to clearly see the effect of the AVK smoothing procedure on the high-resolution datasets. In order to better compare the different months and datasets, Fig. 8 shows vertical profiles of the diurnal cycle amplitude for all months and datasets. Here, the amplitude is defined as the percentage change

275 between the maximum and the minimum normalized ozone ratios (ΔO_3 as defined in Eq. 2) during the course of a day.

The new harmonized ozone time series from GROMOS and SOMORA have excellent agreement in ozone diurnal cycle. They agree well in patterns and amplitudes at all seasons and most altitudes. Some small discrepancies can be seen in summertime in the transition region (see e.g. July or August in Fig. 8), however, as shown in Sauvageat et al. (2022b), it is also the season where GROMOS and SOMORA experienced the larger discrepancies between their respective measurements. The two

280 regimes of the ozone diurnal cycle are clearly visible in all datasets. Namely, the accumulation of ozone during daytime in the stratosphere and the depletion of ozone during daytime above ~ 1 hPa are well captured by all datasets.

Among the model datasets, we observe most discrepancies of the diurnal cycle amplitude by the GDOC during wintertime in the upper stratosphere (see also the month of January and February, shown in the supplementary material January or February




Figure 5. Same as Fig. 4 but for June.

in Figure 8). To some extent, these discrepancies could be due to the temporal (different averaging periods) and longitudinal (zonal mean in GDOC) variability, which are both smoothed out in the GDOC. As mentioned by Frith et al. (2020), this is also the season where the ozone diurnal cycle is smaller and where the model uncertainties are higher. Below, we will present a summary of the differences between the MWRs and the models focusing on different altitude regions and discuss in more details the reasons for the observed discrepancies.

3.2.1 Mesosphere ($p < 0.3$ hPa)

Overall, we observe a tendency of the models to overestimate the diurnal ozone depletion in the mesosphere. It is mostly noticeable above ~ 0.1 hPa where the sensitivity of the MWRs are decreasing and the measurement error is growing fast. Therefore, even if the effect of the lower sensitivity should be included through the AVK smoothing, biases above this altitude should be considered with care. Note that at this altitude, BASCOE also has a limited vertical resolution as it only uses 2 pressure levels above 0.1 hPa. Considering the above limitations, we still observe quite a good agreement of the upper mesospheric diurnal cycle at all seasons. In agreement with Parrish et al. (2014) but in contradiction with the conclusions from Studer et al. (2014), we do not observe a significant seasonal variation of the mesospheric diurnal cycle amplitude. This is in better agreement with the model results, which show similar amplitude throughout the year.

Monthly-averaged ozone diurnal cycle over Switzerland in December as seen in GROMOS, SOMORA, GDOC, BASCOE and WACCM datasets. Note that only the BASCOE and WACCM datasets have been convolved with the AVKs of GROMOS as explained in Sect. 2.1.

Same as Fig. 4 but for June:




Figure 6. Monthly averaged ozone diurnal cycle over Switzerland at 4 selected pressure ranges in December. For BASCOE and WACCM we show both the diurnal cycle before (solid lines) and after convolution (dash lines) of the dataset with the AVKs of GROMOS. To account for the large changes of the diurnal cycle amplitude with altitude, the scale of the y-axis is adapted for each sub-plot.




Figure 7. Same as Fig. 6 but for June.




Figure 8. Monthly averaged vertical profiles of the ozone diurnal cycle amplitude. It shows the percentage change between the maximum and the minimum ozone ratio values. The dashed lines show the model results after convolution with the AVKs of GROMOS.

3.2.2 Lower mesosphere (1 – 0.3 hPa)

In the lower mesosphere, we note a consistent bias between the models and the observations around sunrise: the diurnal ozone
 300 depletion observed by the MWRs consistently starts earlier than the models. It is true for most months and could be partly explained by differences in the vertical resolution (e.g. see Fig. 7 at 51 and 56 km). Interestingly, it does not seem to impact the sunset period which rules out potential errors arising from the time conversion between the different datasets. This feature was also observed by Parrish et al. (2014) over Mauna Loa, and it seems to persist even after application of the AVKs (not for all months though), which gives us confidence that it does not result from the a-priori.

305 3.2.3 Stratopause region (3 – 1 hPa)

Around the stratopause, we can clearly see the complex transition region between the mesospheric diurnal depletion and the stratospheric accumulation. This is where we notice the largest biases between our different datasets. In fact, we observe discrepancies among the three model-based datasets, and between the observations and the ~~model-based-datasets.~~ The bias models. The biases around the stratopause (1 – 3 hPa) are similar to the ones reported by Parrish et al. (2014) and Haeefe
 310 et al. (2008), i.e. differing behaviour in the pre-dawn hours and after sunrise. They are seen at all seasons during daytime and reach values up to 2 % differences among the models themselves (e.g. between 1 and 2 hPa in Fig. 7). Between the models

and the MWRs, the biases are significantly reduced by the application of the AVK smoothing procedure but we still note biases up to 2 % in this region.

As will be shown in Sect. 3.3, the upper stratosphere and lower mesosphere are also experiencing short-term variability of the ozone diurnal cycle, which can influence the monthly averaged cycle. In particular, the datasets produced using only a few specific years (i.e. WACCM or GDOC in our case), will be influenced by the short-term variability of these years whereas it will be smoothed out in the MWR or the BASCOE datasets which are averaged on 10 years or more. As shown in Figure S11 from the supplementary material of Frith et al. (2020), whereas the inter-annual variability is generally limited below 5 hPa and during summer, inter-annual variations up to 5 % around 0.5 – 1 hPa can be seen during wintertime in the mid-latitudes. It supports the existence of short-term variability in the ozone diurnal cycle and might therefore explain some of the remaining discrepancies near the stratopause region.

3.2.4 Middle and lower stratosphere (30 – 3 hPa)

In the middle and lower stratosphere, we observe the typical behaviour of the stratospheric ozone diurnal cycle: a small dip after sunrise followed by a gentle accumulation reaching a maximum in the late afternoon. The stratospheric cycle shows a high seasonal variability, with a maximum diurnal cycle amplitude around the summer solstice and lower diurnal variations during winter. In summer, we observe a peak amplitude of the ozone diurnal cycle of 3 – 4 % in the afternoon around 5 hPa in July, reducing to less than 2 % in the wintertime. For this reason, the dip after sunrise, attributed to rapid dissociation of NO₂ at sunrise (Pallister and Tuck, 1983), is mostly visible during the summer months. Note that this is a significant improvement compared to the previous retrievals of [the](#) GROMOS time series, where the dip was not observed and where the amplitude of the stratospheric cycle was high compared to the models (Figure 6a and 6b in Studer et al. (2014)). With the new time series, the amplitude of stratospheric ozone cycle is well captured by GROMOS and SOMORA at most seasons. In fact, most of the discrepancies that we observe in the middle stratosphere are the consequences of the limited vertical resolution of the MWRs, whereas the differences of the lower stratosphere stay mostly within the error bars.

3.3 Short-term variability of the ozone diurnal cycle

In this section, we present the first measurements of short-term ozone diurnal cycle variability using the unique setup offered by the collocated, hourly resolved measurements from GROMOS and SOMORA. To our knowledge, it is the first time that short-term variability of the ozone diurnal cycle is observed and in the following, we try to identify some of the reasons leading to such events focusing on a case study from the boreal winter 2014-2015. The upper panel in Fig. 9 shows the ozone concentration in the upper stratosphere from GROMOS, SOMORA and BASCOE. From the ozone times series already, there are time periods where the ozone VMR shows some large fluctuations on a diurnal basis in a season where the mean ozone cycle is usually small (see Fig. 6, around 40 km). An example of such case can be seen at the beginning of January 2015 (arrow on Fig. 9) or, for instance, at the end of January 2015. From the BASCOE time series, we can even identify other [period-periods](#) with enhanced cycle which are not really seen as such in the MWRs measurements. These are some examples of what we refer as [“short-term ozone diurnal cycle variability”](#), generally lasting for a few days.




Figure 9. (a) Ozone VMR from GROMOS, SOMORA and BASCOE during the boreal winter 2014-2015. (b) Nitrogen oxides (NO and NO₂) simulated by BASCOE. All quantities are averaged between 3 and 0.81 hPa and the ozone time series over 2 h time periods. NO_x are shown as daily mean of nighttime (NO₂) respectively daytime (NO) values. The arrow highlights the period with an enhanced diurnal cycle.

345 These enhancements can be better seen in Fig. 10, in the form of day-to-night D_{O_3} anomalies in the middle atmosphere (60 – 0.1 hPa). It shows similar patterns in GROMOS and SOMORA time series, with large increase of D_{O_3} around 1 hPa at the beginning of January 2015, followed by a secondary peak in the second half of the month. To some extent, BASCOE is also able to reproduce these two peaks in the ozone diurnal cycle, somehow limited to below 1 hPa and with limited vertical resolution.

350 Focusing on the upper stratosphere (3 – 1 hPa) where the anomalies are highest, Fig. 11 shows the temporal evolution of different quantities during the winter 2014-2015. In particular, Fig. 11(b) shows the day and nighttime ozone values. Focusing on the early January event, it can be seen that there is a consistent increase in daytime ozone from the three datasets, somehow delayed slightly in time in BASCOE compared to the MWRs. Such increase is also visible during the second peak at the end of January for GROMOS and SOMORA and somehow less clearly in the BASCOE series. In terms of amplitude, this increase
 355 is substantial and corresponds approximately to 4 to 5 times the monthly averaged day-to-night difference in January which is ~ 0.1 ppmv at this pressure level.

Corresponding to these peaks in D_{O_3} anomalies, sharp decreases in nitrogen oxides (NO_x) are simulated from BASCOE. In fact, the decrease affects both nitric oxide (NO) and nitrogen dioxide (NO₂). The two species are in photochemical balance




Figure 10. Anomalies in day-night DO_3 from GROMOS, SOMORA and BASCOE during the boreal winter 2014-2015. For each dataset, we show the differences of DO_3 compared to a monthly climatology computed on the decade 2010-2020.

during the day as NO mainly originates from photolysis of NO_2 during daytime and can react with ozone to give back NO_2 ,
 360 forming a major catalytic ozone depletion process in the middle atmosphere (Crutzen, 1970). In fact, the effect of NO_x on ozone
 maximizes between 20 and 45 km altitude ($\sim 50 - 1$ hPa), corresponding well to the peaks of diurnal cycle enhancements
 seen on Fig. 10. To cross-validate these simulations, we also show some nitrous oxide (N_2O) measurements taken by the
 MLS instrument onboard the Aura satellite in Fig. 11(e). Also here, the early January peak is visible as a decrease in the
 N_2O measurement from MLS, which makes sense as N_2O is the main source of NO_x in this altitude region (McElroy and
 365 McConnell, 1971). Note that this is not the only event identified where such a behaviour can be seen. In fact, it seems that most
 winters seem to experience similar events (see e.g. similar plots for the boreal winter 2016-2017 shown in Appendix C)

In order to provide a more global picture and investigate the reasons for the N_2O decrease seen above Central Europe in the
 MLS measurements, we investigated the dynamical situation of the Northern Hemisphere by looking at the ERA5 reanalysis
 data during this period. In fact, this event follows closely a Sudden Stratospheric Warming (SSW) which took place early




Figure 11. Time series of different quantities during the winter 2014-2015, all averaged between 3 and 0.81 hPa. (a) is the DO_3 anomalies, (b) shows the ozone VMR of the three dataset during daytime and nighttime, (c) is temperature from MLS and ERA5, (d) are NO and NO_2 as simulated by BASCOE and (e) is N_2O measurements from MLS.

370 January. It was a minor warming but with significant disturbances on middle atmospheric chemistry and transport (Manney et al., 2015). In fact, Fig. 2 from Manney et al. shows how the polar vortex briefly splitted at the onset of the SSW, leading to a mixing of the air between the mid-latitude and the poles in the upper stratosphere. Following this event, some filaments of polar air containing little ozone and N_2O reached Central Europe as can be seen on the ozone map in Fig. 12. Such an irruption of polar air over Switzerland would explain the decrease in the N_2O MLS measurements seen early January and might well
 375 explain the subsequent changes NO_x and consequently in the ozone diurnal cycle.

Interestingly, we do not observe the strong ozone decrease associated with this filament of polar air reaching Bern, neither in the MWRs nor in the BASCOE time series. This could be a problem from the ERA5 reanalysis data as it does not feature any diurnal cycle at these altitude and therefore might also lack the reaction of ozone to greater sun illumination (resulting in more ozone production and therefore increasing ozone amount in the low-ozone polar air which would be missed by the model).
 380 Even though this event might be considered textbook example of such dynamical event, we find it interesting to find such a coherent picture of a short-term event from a combination of ground-based measurements, chemistry transport model, satellite measurements and reanalysis data.




Figure 12. Situation over Europe in the upper stratosphere ~ 5 hPa shortly after the minor SSW of early January 2015 as seen in the ERA5 reanalysis.

In this publication, we focused on a specific case of short-term diurnal cycle variability related to NO_x changes in the upper stratosphere. However, in the MWRs and in the BASCOE time series, the short-term variability is not always associated with changes in NO_x concentration. Among the non-chemical processes that can impact the diurnal cycle amplitude, solar tides are important, both through vertical transport of O_x and through their modulation of temperature which can have a significant impact on ozone photochemistry (Schanz et al., 2014). Solar tides have periods of 1 solar day (24 h) or of its harmonics (e.g. 12 or 8 h) and can therefore impact the diurnal cycle as well but generally their impact should be larger in the upper mesosphere and above (see e.g. Bjarnason et al. (1987); Huang et al. (2010b)). However, Sakazaki et al. (2013) reported significant influence from tidal vertical transport at ~ 35 km already, and therefore, short-term tidal variability could induce some short-term variations of the ozone diurnal cycle in the upper stratosphere. The reciprocal is also true as variability of the stratospheric ozone thermal forcing can influence the generation of tides (Goncharenko et al., 2012). There have been studies on the tidal variability in the middle atmosphere and it has been shown that solar tides are subject to a wide range of temporal variability, from intra-seasonal to short-term variability of a few days (Kopp et al., 2015; Baumgarten et al., 2018). It has also been shown that tides respond to SSW and can have non-linear interactions with planetary and gravity waves. Hence, it is likely

that a coupling exists between the tidal variability and the short-term variability observed in the ozone diurnal cycle, but it is difficult to conclude on any causal relation without additional data (e.g. high resolution temperature or wind measurements).

4 Conclusions

Using new harmonized ozone time series from two nearby located microwave radiometers enables us to study in great details
400 the ozone diurnal cycle over Switzerland. With more than 11 years of parallel, independent measurements, these instruments
provide a unique validation source for satellite and model-based datasets. We find that the recently published GDOC clima-
tology compares well with our MWRs above Switzerland and agrees well with the WACCM and BASCOE models in the
stratosphere and lower mesosphere. As reported by previous studies, we observe some remaining discrepancies between our
observations and the models near the stratopause, in the transition region between ozone daytime accumulation and depletion.
405 The discrepancies remain small and are significant only during summertime, where the diurnal cycle is stronger, providing
better signal-to-noise ratio for the observations. Some of our results contradict a previous study also based on the GROMOS
instrument (Studer et al., 2014), now providing a better agreement of the ozone diurnal cycle compared to model-based dataset
but also compared to other previous MWR diurnal cycle study (Parrish et al., 2014). These updated results motivated the present
study, and are a consequence of the spectrometer change and of the recent harmonization of the calibration and retrieval routines
410 of GROMOS and SOMORA (Sauvageat et al., 2022b).

For the first time, short-term variations of the ozone diurnal cycle could be detected in two collocated MWR time series,
highlighting the value of ground-based radiometric measurements to monitor the short-term dynamics and photochemistry in
the middle atmosphere. The quantification of these variations is limited by the rapidly increasing measurement noise, however
some enhancements of the diurnal cycle are clearly visible in the upper stratosphere during wintertime, where the diurnal cycle
415 is otherwise very small. Compared to the averaged monthly diurnal cycle, we find enhancement of 4 – 5 times the monthly
mean diurnal cycle amplitude lasting for a few days. In fact, the observed short-term variability of the ozone diurnal cycle
seems much higher than its intra-seasonal (month-to-month) or inter-annual variability during wintertime.

Regional (longitudinal) variability of the stratospheric ozone diurnal cycle has previously been identified by Schanz et al.
(2014) in a model-based study from WACCM. They attributed the regional variability to changes in temperature (atmospheric
420 tides), O_x and NO_y . Our study supports that, in some cases, short-term variability in the ozone diurnal cycle can be attributed
to changes in NO_x concentrations through dynamical transport. In other cases, other processes might be acting to modify the
amplitude of the ozone diurnal cycle, for instance changes in atmospheric tides. Our study also shows that a CTM like BASCOE
is able to simulate the changes of the ozone diurnal cycle amplitude due to changes in NO_x . In view of its significance, we
believe that the reasons and importance of the short-term variability of the diurnal cycle should be further investigated with
425 BASCOE over the globe.

It is beyond the scope of this publication to provide comprehensive analysis of this phenomenon but we aim at bringing
some new data to better understand stratospheric ozone diurnal cycle variability. It seems to be of particular interest in views
of the recent studies aiming at better accounting for the stratospheric ozone diurnal variability in satellite datasets (e.g. Frith

et al., 2020; Strode et al., 2022; Natarajan et al., 2023). Note that we focused our analysis on the upper stratosphere, where the
 430 short-term variability was most visible in our observations, but short-term variations are not limited to this region. In fact, our
 observations indicate that the variability is also present in the mesosphere and the lower stratosphere, where the role of NO_x is
 less important and where other processes likely dominate. To conclude, more work is definitely needed to assess the importance
 of the short-term variability of the ozone diurnal cycle and confirm the potential role of other mechanisms influencing it.

Code and data availability. The GROMOS and SOMORA level 2 data are available from the Bern Open Repository and Information System
 435 in the form of yearly netCDF files (Sauvageat et al., 2022a; Maillard Barras et al., 2022b). The recently harmonized calibration and retrieval
 routines are freely available at <https://doi.org/10.5281/zenodo.6799356>. The data and analysis code reproducing the results presented in this
 manuscript are freely available. MLS v5 data (Schwartz et al., 2020) are available from the NASA Goddard Space Flight Center Earth
 Sciences Data and Information Services Center (GES DISC): <https://disc.gsfc.nasa.gov/>. The ERA5 dataset (Hersbach et al., 2020) was
 downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store. The CMIP6 recommendations for sulfate aerosols used
 440 in BASCOE are available at ftp://iacftp.ethz.ch/pub_read/luo/CMIP6/

Appendix A: Monthly ozone profile comparisons

A1 Additional plots for spring and autumn




Figure A1. Same as Fig. 2 but for March.




Figure A2. Same as Fig. 2 but for September.

Appendix B: Monthly diurnal cycle

B1 Additional plots for spring and autumn




Figure B1. Same as Fig. 4 but for March.




Figure B2. Same as Fig. 4 but for September.




Figure B3. Same as Fig. 6 but for March.




Figure B4. Same as Fig. 6 but for September.

C1 Short-term

In this section, we show measurements and simulations of the short-term variability during the boreal winter 2016-2017. Whereas more work is needed to unravel the complete picture of this winter, it shows another example of diurnal cycle enhancement associated with a sharp decrease of nitrogen oxides in the upper stratosphere.




Figure C1. Anomalies in day-night DO_3 from GROMOS, SOMORA and BASCOE during the boreal winter 2016-2017. For each dataset, we show the differences of DO_3 compared to a monthly climatology computed on the decade 2010-2020.




Figure C2. Time series of different quantities during the winter 2016-2017, all averaged between 3 and 0.81 hPa. (a) is the DO_3 anomalies, (b) shows the ozone VMR of the three dataset during daytime and nighttime, (c) is temperature from MLS and ERA5, (d) are NO and NO_2 as simulated by BASCOE and (e) is N_2O measurements from MLS.

450 *Author contributions.* ES carried out the data analysis, analysed the results, and prepared the manuscript. KH conceived the project and contributed to the interpretation of the results. EM provided the SOMORA data and contributed to the interpretation of the results. SH performed a preliminary study within the frame of his master thesis. QE performed the BASCOE simulation and provided valuable help on the interpretation of all model results. AH and AM contributed to the interpretation of the results. All of the authors discussed the scientific findings and provided valuable feedback for the manuscript editing.

455 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This work has been partly funded by MeteoSwiss and the Swiss Global Atmospheric Watch program. The authors would like to acknowledge all the people involved in the design and operation of GROMOS and SOMORA. Also, they would like to thank the developers of the Atmospheric Radiative Transfer Simulator (Buehler et al., 2018), version 2.4 and their precious support to setup the ozone retrievals. In addition, we thank the numerous contributors to the free and open source software packages used for the data analysis,
 460 in particular xarray, matplotlib, Typhon and pyretrieval.

References

- Ball, W. T., Alsing, J., Mortlock, D. J., Staehelin, J., Haigh, J. D., Peter, T., Tummon, F., Stübi, R., Stenke, A., Anderson, J., Bourassa, A., Davis, S. M., Degenstein, D., Frith, S., Froidevaux, L., Roth, C., Sofieva, V., Wang, R., Wild, J., Yu, P., Ziemke, J. R., and Rozanov, E. V.: Evidence for a continuous decline in lower stratospheric ozone offsetting ozone layer recovery, *Atmospheric Chemistry and Physics*, 18, 1379–1394, <https://doi.org/10.5194/acp-18-1379-2018>, 2018.
- 465 Baumgarten, K., Gerding, M., Baumgarten, G., and Lübken, F.-J.: Temporal variability of tidal and gravity waves during a record long 10-day continuous lidar sounding, *Atmospheric Chemistry and Physics*, 18, 371–384, <https://doi.org/10.5194/acp-18-371-2018>, 2018.
- Bernet, L., Clarmann, T. v., Godin-Beekmann, S., Ancellet, G., Maillard Barras, E., Stübi, R., Steinbrecht, W., Kämpfer, N., and Hocke, K.: Ground-based ozone profiles over central Europe: incorporating anomalous observations into the analysis of stratospheric ozone trends, *Atmospheric Chemistry and Physics*, 19, 4289–4309, <https://doi.org/10.5194/acp-19-4289-2019>, 2019.
- 470 Bhartia, P. K., McPeters, R. D., Flynn, L. E., Taylor, S., Kramarova, N. A., Frith, S., Fisher, B., and DeLand, M.: Solar Backscatter UV (SBUV) total ozone and profile algorithm, *Atmospheric Measurement Techniques*, 6, 2533–2548, <https://doi.org/10.5194/amt-6-2533-2013>, 2013.
- Bjarnason, G. G., Solomon, S., and Garcia, R. R.: Tidal influences on vertical diffusion and diurnal variability of ozone in the mesosphere, *Journal of Geophysical Research: Atmospheres*, 92, 5609–5620, <https://doi.org/10.1029/JD092iD05p05609>, 1987.
- 475 Buehler, S. A., Mendrok, J., Eriksson, P., Perrin, A., Larsson, R., and Lemke, O.: ARTS, the Atmospheric Radiative Transfer Simulator - Version 2.2, the planetary toolbox edition, *Geoscientific Model Development*, 11, 1537–1556, <https://doi.org/10.5194/gmd-11-1537-2018>, 2018.
- Chabrilat, S., Vigouroux, C., Christophe, Y., Engel, A., Errera, Q., Minganti, D., Monge-Sanz, B. M., Segers, A., and Mahieu, E.: Comparison of mean age of air in five reanalyses using the BASCOE transport model, *Atmospheric Chemistry and Physics*, 18, 14 715–14 735, <https://doi.org/10.5194/acp-18-14715-2018>, 2018.
- 480 Chapman, S.: A theory of the upper-atmospheric ozone, *Memoirs of the Royal Meteorological Society*, 3, <https://www.rmets.org/sites/default/files/papers/chapman-memoirs.pdf>, 1930.
- Connor, B. J., Siskind, D. E., Tsou, J., Parrish, A., and Remsberg, E. E.: Ground-based microwave observations of ozone in the upper stratosphere and mesosphere, *Journal of Geophysical Research: Atmospheres*, 99, 16 757–16 770, <https://doi.org/10.1029/94JD01153>, 1994.
- 485 Crutzen, P. J.: The influence of nitrogen oxides on the atmospheric ozone content, *Quarterly Journal of the Royal Meteorological Society*, 96, 320–325, <https://doi.org/10.1002/qj.49709640815>, 1970.
- Errera, Q. and Fonteyn, D.: Four-dimensional variational chemical assimilation of CRISTA stratospheric measurements, *Journal of Geophysical Research: Atmospheres*, 106, 12 253–12 265, <https://doi.org/10.1029/2001JD900010>, 2001.
- 490 Errera, Q., Daerden, F., Chabrilat, S., Lambert, J. C., Lahoz, W. A., Viscardy, S., Bonjean, S., and Fonteyn, D.: 4D-Var assimilation of MIPAS chemical observations: ozone and nitrogen dioxide analyses, *Atmospheric Chemistry and Physics*, 8, 6169–6187, <https://doi.org/10.5194/acp-8-6169-2008>, 2008.
- Frith, S. M., Bhartia, P. K., Oman, L. D., Kramarova, N. A., McPeters, R. D., and Labow, G. J.: Model-based climatology of diurnal variability in stratospheric ozone as a data analysis tool, *Atmospheric Measurement Techniques*, 13, 2733–2749, <https://doi.org/10.5194/amt-13-2733-2020>, 2020.
- 495

- Froidevaux, L., Jiang, Y. B., Lambert, A., Livesey, N. J., Read, W. G., Waters, J. W., Browell, E. V., Hair, J. W., Avery, M. A., McGee, T. J., Twigg, L. W., Sumnicht, G. K., Jucks, K. W., Margitan, J. J., Sen, B., Stachnik, R. A., Toon, G. C., Bernath, P. F., Boone, C. D., Walker, K. A., Filipiak, M. J., Harwood, R. S., Fuller, R. A., Manney, G. L., Schwartz, M. J., Daffer, W. H., Drouin, B. J., Cofield, R. E., Cuddy, D. T., Jarnot, R. F., Knosp, B. W., Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., and Wagner, P. A.: Validation of Aura Microwave Limb Sounder stratospheric ozone measurements, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008771>, 2008.
- Garcia, R. R., Marsh, D. R., Kinnison, D. E., Boville, B. A., and Sassi, F.: Simulation of secular trends in the middle atmosphere, 1950–2003, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD007485>, 2007.
- Godin-Beekmann, S., Azouz, N., Sofieva, V. F., Hubert, D., Petropavlovskikh, I., Effertz, P., Ancellet, G., Degenstein, D. A., Zawada, D., Froidevaux, L., Frith, S., Wild, J., Davis, S., Steinbrecht, W., Leblanc, T., Querel, R., Tourpali, K., Damadeo, R., Maillard Barras, E., Stübi, R., Vigouroux, C., Arosio, C., Nedoluha, G., Boyd, I., Van Malderen, R., Mahieu, E., Smale, D., and Sussmann, R.: Updated trends of the stratospheric ozone vertical distribution in the 60°S–60°N latitude range based on the LOTUS regression model, *Atmospheric Chemistry and Physics*, 22, 11 657–11 673, <https://doi.org/10.5194/acp-22-11657-2022>, 2022.
- Goncharenko, L. P., Coster, A. J., Plumb, R. A., and Domeisen, D. I. V.: The potential role of stratospheric ozone in the stratosphere-ionosphere coupling during stratospheric warmings, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051261>, 2012.
- Haefele, A., Hocke, K., Kämpfer, N., Keckhut, P., Marchand, M., Bekki, S., Morel, B., Egorova, T., and Rozanov, E.: Diurnal changes in middle atmospheric H₂O and O₃: Observations in the Alpine region and climate models, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2008JD009892>, 2008.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Huang, F. T., Reber, C. A., and Austin, J.: Ozone diurnal variations observed by UARS and their model simulation, *Journal of Geophysical Research: Atmospheres*, 102, 12 971–12 985, <https://doi.org/10.1029/97JD00461>, 1997.
- Huang, F. T., Mayr, H. G., Russell III, J. M., and Mlynczak, M. G.: Ozone diurnal variations in the stratosphere and lower mesosphere, based on measurements from SABER on TIMED, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2010JD014484>, 2010a.
- Huang, F. T., McPeters, R. D., Bhartia, P. K., Mayr, H. G., Frith, S. M., Russell III, J. M., and Mlynczak, M. G.: Temperature diurnal variations (migrating tides) in the stratosphere and lower mesosphere based on measurements from SABER on TIMED, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD013698>, 2010b.
- Huijnen, V., Flemming, J., Chabrilat, S., Errera, Q., Christophe, Y., Blechschmidt, A.-M., Richter, A., and Eskes, H.: C-IFS-CB05-BASCOE: stratospheric chemistry in the Integrated Forecasting System of ECMWF, *Geosci. Model Dev.*, 9, 3071–3091, <https://doi.org/10.5194/gmd-9-3071-2016>, 2016.
- Kinnison, D., Brasseur, G. P., Walters, S., Garcia, R., Marsh, D., Sassi, F., Harvey, V., Randall, C., Emmons, L., Lamarque, J., et al.: Sensitivity of chemical tracers to meteorological parameters in the MOZART-3 chemical transport model, *Journal of Geophysical Research: Atmospheres*, 112, 2007.

- 535 Kopp, M., Gerding, M., Höffner, J., and Lübken, F.-J.: Tidal signatures in temperatures derived from daylight lidar soundings above Kühlungsborn (54°N, 12°E), *Journal of Atmospheric and Solar-Terrestrial Physics*, 127, 37–50, <https://doi.org/10.1016/j.jastp.2014.09.002>, layered Phenomena in the Mesopause Region, 2015.
- Livesey, N. J., Read, W. G., Wagner, P. A., Froidevaux, L., Santee, M. L., Schwartz, M. J., Lambert, A., Valle, L. F. M., Pumphrey, H. C., Manney, G. L., Fuller, R. A., Jarnot, R. F., Knosp, B. W., and Lay, R. R.: Earth Observing System (EOS) Aura Microwave Limb Sounder (MLS) Version 5.0x Level 2 and 3 data quality and description document., Tech. rep., <https://mls.jpl.nasa.gov/eos-aura-mls/data-documentation>, last access: 30 December 2022, 2022.
- 540 Maillard Barras, E., Haefele, A., Nguyen, L., Tummon, F., Ball, W. T., Rozanov, E. V., Rüfenacht, R., Hocke, K., Bernet, L., Kämpfer, N., Nedoluha, G., and Boyd, I.: Study of the dependence of long-term stratospheric ozone trends on local solar time, *Atmospheric Chemistry and Physics*, 20, 8453–8471, <https://doi.org/10.5194/acp-20-8453-2020>, 2020.
- 545 Maillard Barras, E., Haefele, A., Stübi, R., Jouberton, A., Schill, H., Petropavlovskikh, I., Miyagawa, K., Stanek, M., and Froidevaux, L.: Dynamical linear modeling estimates of long-term ozone trends from homogenized Dobson Umkehr profiles at Arosa/Davos, Switzerland, *Atmospheric Chemistry and Physics*, 22, 14 283–14 302, <https://doi.org/10.5194/acp-22-14283-2022>, 2022a.
- Maillard Barras, E., Sauvageat, E., Hocke, K., Murk, A., and Haefele, A.: Harmonized middle atmospheric ozone time series from SOMORA, <https://doi.org/10.48620/119>, [data set], last access: 3 March 2023, 2022b.
- 550 Manney, G. L., Lawrence, Z. D., Santee, M. L., Read, W. G., Livesey, N. J., Lambert, A., Froidevaux, L., Pumphrey, H. C., and Schwartz, M. J.: A minor sudden stratospheric warming with a major impact: Transport and polar processing in the 2014/2015 Arctic winter, *Geophysical Research Letters*, 42, 7808–7816, <https://doi.org/10.1002/2015GL065864>, 2015.
- Marsh, D. R., Mills, M. J., Kinnison, D. E., Lamarque, J.-F., Calvo, N., and Polvani, L. M.: Climate Change from 1850 to 2005 Simulated in CESM1(WACCM), *Journal of Climate*, 26, 7372 – 7391, <https://doi.org/10.1175/JCLI-D-12-00558.1>, 2013.
- 555 McElroy, M. B. and McConnell, J. C.: Nitrous Oxide: A Natural Source of Stratospheric NO, *Journal of Atmospheric Sciences*, 28, 1095 – 1098, [https://doi.org/10.1175/1520-0469\(1971\)028<1095:NOANSO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1971)028<1095:NOANSO>2.0.CO;2), 1971.
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J., Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I. G., Law, R. M., Lunder, C. R., O’Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J., and Weiss, R.: Historical greenhouse gas concentrations for climate modelling (CMIP6), *Geosci. Model Dev.*, 10, 2057–2116, <https://doi.org/10.5194/gmd-10-2057-2017>, 2017.
- 560 Minganti, D., Chabrillat, S., Christophe, Y., Errera, Q., Abalos, M., Prignon, M., Kinnison, D. E., and Mahieu, E.: Climatological impact of the Brewer–Dobson circulation on the N₂O budget in WACCM, a chemical reanalysis and a CTM driven by four dynamical reanalyses, *Atmos. Chem. Phys.*, 20, 12 609–12 631, <https://doi.org/10.5194/acp-20-12609-2020>, 2020.
- Minganti, D., Chabrillat, S., Errera, Q., Prignon, M., Kinnison, D. E., Garcia, R. R., Abalos, M., Alsing, J., Schneider, M., Smale, D., Jones, N., and Mahieu, E.: Evaluation of the N₂O Rate of Change to Understand the Stratospheric Brewer–Dobson Circulation in a Chemistry–Climate Model, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD036 390, <https://doi.org/10.1029/2021JD036390>, 2022.
- 565 Natarajan, M., Damadeo, R., and Flittner, D.: Solar occultation measurement of mesospheric ozone by SAGE III/ISS: impact of variations along the line of sight caused by photochemistry, *Atmospheric Measurement Techniques*, 16, 75–87, <https://doi.org/10.5194/amt-16-75-2023>, 2023.
- 570 Orbe, C., Oman, L. D., Strahan, S. E., Waugh, D. W., Pawson, S., Takacs, L. L., and Molod, A. M.: Large-Scale Atmospheric Transport in GEOS Replay Simulations, *Journal of Advances in Modeling Earth Systems*, 9, 2545–2560, <https://doi.org/10.1002/2017MS001053>, 2017.

- Pallister, R. C. and Tuck, A. F.: The diurnal variation of ozone in the upper stratosphere as a test of photochemical theory, *Quarterly Journal of the Royal Meteorological Society*, 109, 271–284, <https://doi.org/10.1002/qj.49710946002>, 1983.
- 575 Parrish, A., Boyd, I. S., Nedoluha, G. E., Bhartia, P. K., Frith, S. M., Kramarova, N. A., Connor, B. J., Bodeker, G. E., Froidevaux, L., Shiotani, M., and Sakazaki, T.: Diurnal variations of stratospheric ozone measured by ground-based microwave remote sensing at the Mauna Loa NDACC site: measurement validation and GEOSCCM model comparison, *Atmospheric Chemistry and Physics*, 14, 7255–7272, <https://doi.org/10.5194/acp-14-7255-2014>, publisher: Copernicus GmbH, 2014.
- Prather, M. J.: Ozone in the upper stratosphere and mesosphere, *Journal of Geophysical Research*, 86, 5325, <https://doi.org/10.1029/JC086iC06p05325>, 1981.
- 580 Prignon, M., Chabrilat, S., Friedrich, M., Smale, D., Strahan, S. E., Bernath, P. F., Chipperfield, M. P., Dhomse, S. S., Feng, W., Minganti, D., Servais, C., and Mahieu, E.: Stratospheric Fluorine as a Tracer of Circulation Changes: Comparison Between Infrared Remote-Sensing Observations and Simulations With Five Modern Reanalyses, *J. Geophys. Res.*, 126, e2021JD034995, <https://doi.org/10.1029/2021JD034995>, 2021.
- 585 Ricaud, P., Brillet, J., De La Noe, J., and Parisot, J. P.: Diurnal and seasonal variations of stratomesospheric ozone: Analysis of ground-based microwave measurements in Bordeaux, France, *Journal of Geophysical Research: Atmospheres*, 96, 18 617–18 629, <https://doi.org/10.1029/91JD01871>, 1991.
- Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002299>, 2003.
- 590 Sakazaki, T., Fujiwara, M., Mitsuda, C., Imai, K., Manago, N., Naito, Y., Nakamura, T., Akiyoshi, H., Kinnison, D., Sano, T., Suzuki, M., and Shiotani, M.: Diurnal ozone variations in the stratosphere revealed in observations from the Superconducting Submillimeter-Wave Limb-Emission Sounder (SMILES) on board the International Space Station (ISS), *Journal of Geophysical Research: Atmospheres*, 118, 2991–3006, <https://doi.org/10.1002/jgrd.50220>, 2013.
- Sakazaki, T., Shiotani, M., Suzuki, M., Kinnison, D., Zawodny, J. M., McHugh, M., and Walker, K. A.: Sunset–sunrise difference in solar 595 occultation ozone measurements (SAGE II, HALOE, and ACE–FTS) and its relationship to tidal vertical winds, *Atmospheric Chemistry and Physics*, 15, 829–843, <https://doi.org/10.5194/acp-15-829-2015>, 2015.
- Sauvageat, E., Hocke, K., Maillard Barras, E., Haefele, A., and Murk, A.: Harmonized middle atmospheric ozone time series from GROMOS, <https://doi.org/10.48620/65>, [data set], last access: 3 March 2023, 2022a.
- Sauvageat, E., Maillard Barras, E., Hocke, K., Haefele, A., and Murk, A.: Harmonized retrieval of middle atmospheric ozone from two mi- 600 crowave radiometers in Switzerland, *Atmospheric Measurement Techniques*, 15, 6395–6417, <https://doi.org/10.5194/amt-15-6395-2022>, 2022b.
- Schanz, A., Hocke, K., and Kämpfer, N.: Daily ozone cycle in the stratosphere: global, regional and seasonal behaviour modelled with the Whole Atmosphere Community Climate Model, *Atmospheric Chemistry and Physics*, 14, 7645–7663, <https://doi.org/10.5194/acp-14-7645-2014>, 2014.
- 605 Schanz, A., Hocke, K., Kämpfer, N., Chabrilat, S., Inness, A., Palm, M., Notholt, J., Boyd, I., Parrish, A., and Kasai, Y.: The Diurnal Variation in Stratospheric Ozone from MACC Reanalysis, ERA-Interim, WACCM, and Earth Observation Data: Characteristics and Intercomparison, *Atmosphere*, 12, 625, <https://doi.org/10.3390/atmos12050625>, 2021.
- Schneider, N., Selsis, F., Urban, J., Lezeaux, O., Noé, J. D. L., and Ricaud, P.: Seasonal and Diurnal Ozone Variations: Observations and Modeling, *Journal of Atmospheric Chemistry*, 50, 25–47, <https://doi.org/10.1007/s10874-005-1172-z>, 2005.

- 610 Schranz, F., Fernandez, S., Kämpfer, N., and Palm, M.: Diurnal variation in middle-atmospheric ozone observed by ground-based microwave radiometry at Ny-Ålesund over 1 year, *Atmospheric Chemistry and Physics*, 18, 4113–4130, <https://doi.org/10.5194/acp-18-4113-2018>, 2018.
- Schwartz, M., Froidevaux, L., Livesey, N., and Read, W.: MLS/Aura Level 2 Ozone (O₃) Mixing Ratio V005, <https://doi.org/10.5067/Aura/MLS/DATA2516>, [data set], last access: 3 March 2023, 2020.
- 615 Skachko, S., Ménard, R., Errera, Q., Christophe, Y., and Chabrilat, S.: EnKF and 4D-Var data assimilation with chemical transport model BASCOE (version 05.06), *Geoscientific Model Development*, 9, 2893–2908, <https://doi.org/10.5194/gmd-9-2893-2016>, 2016.
- SPARC: SPARC CCMVal Report on the Evaluation of Chemistry-Climate Models, in: SPARC Report, edited by Eyring, V., Shepherd, T. G., and Waugh, D. W., SPARC Report No. 5, WCRP-30/2010, WMO/TD – No. 40, available at www.sparc-climate.org/publications/sparc-reports/, 2010.
- 620 SPARC/IO3C/GAW: SPARC/IO3C/GAW Report on Long-term Ozone Trends and Uncertainties in the Stratosphere, in: SPARC Report, edited by Petropavlovskikh, I., Godin-Beekmann, S., Hubert, D., Damadeo, R., Hassler, B., and Sofieva, V., <https://doi.org/10.17874/f899e57a20b>, SPARC Report No. 9, GAW Report No. 241, WCRP-17/2018, available at www.sparc-climate.org/publications/sparc-reports/, 2019.
- Strode, S. A., Taha, G., Oman, L. D., Damadeo, R., Flittner, D., Schoeberl, M., Sioris, C. E., and Stauffer, R.: SAGE III/ISS ozone and NO₂ validation using diurnal scaling factors, *Atmospheric Measurement Techniques*, 15, 6145–6161, <https://doi.org/10.5194/amt-15-6145-2022>, 2022.
- 625 Studer, S., Hocke, K., Schanz, A., Schmidt, H., and Kämpfer, N.: A climatology of the diurnal variations in stratospheric and mesospheric ozone over Bern, Switzerland, *Atmospheric Chemistry and Physics*, 14, 5905–5919, <https://doi.org/10.5194/acp-14-5905-2014>, 2014.
- Vaughan, G.: Diurnal variation of mesospheric ozone, *Nature*, 296, 133–135, <https://doi.org/10.1038/296133a0>, number: 5853 Publisher: Nature Publishing Group, 1982.
- 630 Waters, J., Froidevaux, L., Harwood, R., Jarnot, R., Pickett, H., Read, W., Siegel, P., Cofield, R., Filipiak, M., Flower, D., Holden, J., Lau, G., Livesey, N., Manney, G., Pumphrey, H., Santee, M., Wu, D., Cuddy, D., Lay, R., Loo, M., Perun, V., Schwartz, M., Stek, P., Thurstans, R., Boyles, M., Chandra, K., Chavez, M., Chen, G.-S., Chudasama, B., Dodge, R., Fuller, R., Girard, M., Jiang, J., Jiang, Y., Knosp, B., LaBelle, R., Lam, J., Lee, K., Miller, D., Oswald, J., Patel, N., Pukala, D., Quintero, O., Scaff, D., Van Snyder, W., Tope, M., Wagner, P.,
- 635 and Walch, M.: The Earth observing system microwave limb sounder (EOS MLS) on the Aura Satellite, *IEEE Transactions on Geoscience and Remote Sensing*, 44, 1075–1092, <https://doi.org/10.1109/TGRS.2006.873771>, 2006.
- Zommerfelds, W. C., Kunzi, K. F., Summers, M. E., Bevilacqua, R. M., Strobel, D. F., Allen, M., and Sawchuck, W. J.: Diurnal variations of mesospheric ozone obtained by ground-based microwave radiometry, *Journal of Geophysical Research: Atmospheres*, 94, 12 819–12 832, <https://doi.org/10.1029/JD094iD10p12819>, 1989.