

Effect of different biopreparations on the soil physical properties and CO₂ emissions when are growing food type crops

Sidona Buragienė¹, Egidijus Šarauskis¹, Aida Adamavičienė², Kęstutis Romaneckas², Kristina Lekavičienė¹, Vilma Naujokienė¹

¹ Department of Agricultural Engineering and Safety, Vytautas Magnus University, Agriculture Academy, Lithuania
sidona.buragiene@vdu.lt, egidijus.sarauskis@vdu.lt, kristina.lekaviciene@vdu.lt, vilma.naujokiene@vdu.lt

² Department of Agroecosystems and Soil Science, Vytautas Magnus University, Agriculture Academy, Lithuania
aida.adamaviciene@vdu.lt, kęstutis.romaneckas@vdu.lt,

*Correspondence: vilma.naujokiene@vdu.lt; tel.: (+370 673 58 114)

Abstract

The introduction of innovative technologies in agriculture is key not only to improving the efficiency of agricultural production, crop yields, and quality but also to balancing energy use and preserving a cleaner environment. Biopreparations are environmentally friendly means of restoring the vitality of the soil on which plants can thrive. Biopreparations have an impact on soil health and alter greenhouse gas emissions. The aim of this study was to investigate the effects of different biopreparations on soil porosity, temperature, and CO₂ emission from the soil in North-East Europe (Lithuania) growing food type crops. The experimental studies were carried out over three years, and each spring, after the resumption of winter crops, the soil surface was sprayed with biopreparations of different properties or mixtures of biopreparations, under 7 scenarios, with one scenario left as a control. Soil porosity, temperature, and CO₂ emissions from the soil were measured regularly every month from April to August. The application of the biopreparations showed a cumulative effect on the soil properties. In the third year of the study, the total porosity of the soil was higher in all scenarios compared to the control, ranging between 51% and 74%. The aeration porosity of the soil was also higher in all years of the study than in the control, although no significant differences were obtained. The results of the studies on CO₂ emissions from the soil showed that in the first year, the application of the biopreparations increases emissions compared to the control. However, when assessing the cumulative effect of the biopreparations on soil respiration intensity, it was found that in the third year, most of the biopreparations led to a reduction in CO₂ emissions compared to the control. The lowest emissions were achieved with the biopreparations consisting of essential oils of plants, 40 species of various herbs extracts, marine algae extracts, *Azospirillum sp.*, *Frateruria aurentia*, *Bacillus megaterium*, mineral oils, *Azotobacter vinelandi*, humic acid, gibberellic acid, sodium molybdate, azotobacter chroococcum, azospirillum brasilense, etc.

Keywords: GHG emissions, carbon dioxide, bioproducts, soil porosity, soil temperature

Introduction

1.1 Importance of biopreparations

Decades of soil degradation have led to a search for ways to contribute to soil sustainability by preserving soil properties without harming the environment. Over the last decade, European agricultural policy has increasingly turned towards environmental sustainability, with the aim of reducing the use of chemicals and increasing the organic area (European Commission, 2020). An increasing number of agricultural operators and farmers have adopted environmentally friendly biotechnologies that use biopreparations, i.e., bioproducts designed to inhibit the growth of pathogenic fungi or bacteria, stimulate plant growth, improve plant nutrient uptake, and restore soil properties and fertility (Michalak et al., 2016; Trevisan et al., 2019; Szparaga et al., 2019). Consumers have started to increasingly value agricultural products with high nutritional and functional value and environmentally sustainable production (Caruso et al., 2019; Szparaga et al., 2018). Therefore, bioproducts used in agricultural practice aim to enhance the biological protection of plants by reducing the spread of pathogens and pests, increase crop productivity, improve soil microbiology, change the physical and chemical properties of soil, reduce environmental pollution, and weaken the properties of crop residues (Khattab et al., 2009; Vaitauskiene et al., 2015; Oskiera et al., 2017; Naujokienė et al., 2018). Blaszczyk et al. (2014) stated that *Trichoderma harzianum* and *Trichoderma atroviride* are common components of biopreparations used in agriculture. Fungi of the genus *Trichoderma* can effectively reduce phytopathogens in agricultural soils through various mechanisms (Oskiera et al., 2017). A combination of edaphic and dynamic factors, including crop rotation, residue management, soil type, tillage, and climate, affect the microorganism community (Bünemann et al., 2008; Gil et al., 2011; Zhang et al., 2014). A growing body of research demonstrates that plant-derived phytochemicals affect the soil microbiota through interactions between plant roots and soil (Bais et al., 2006; Kong et al., 2008; Lorenzo et al., 2013). Biopreparations have multiple effects, but scientists are placing more emphasis on their positive effects on plants and soil (Tarantino et al. 2018). Biopreparations are also used as seed diluents to increase germination and reduce seed contamination with pathogenic microorganisms (Selby et al., 2016; Rouphael et al., 2018). Kocira et al. (2020) report that the mixtures of seeds and biopreparations obtained from *Archangelica officinalis* L. significantly inhibit fungal development on the seed surface. Biopreparations have antimicrobial activity because they contain biologically active substances that can inhibit the development of microorganisms. The appropriate composition of the biopreparations to be used depends mainly on the plant species (Nostro et al., 2000; Sen and Batra, 2012; Shihabudeen et al., 2010). The use of biopreparations can reduce the cost of crop production and increase the efficiency of soil nutrient use by reducing the incidence of diseases caused by nutrient deficiencies. However, this effect is not easy to achieve, as it requires a lot of knowledge on the proper selection of biopreparations, their application method, and the correct adjustment of the amount and concentration (Ertani et al., 2018; Szparaga et al., 2019; Michałek et al., 2018).

1.2. Effects of biopreparations on soil

Soil microorganisms are an essential link in the nutrient cycle in the soil and maintain soil fertility. Their activity determines the physical and chemical properties of the soil, and these properties in turn determine how the microorganisms feel in the soil. Soil physical properties such as porosity and temperature are constantly changing under the influence of the environment. A research team from Poland investigating the influence of microorganisms on soil density and porosity found no significant changes over 5 years (Pranagal et al., 2020). Other researchers (Montemurro et al., 2010; Peltre et al., 2015; Juknevičius et al., 2020) have suggested that biopreparations increased the organic carbon content of the soil, which presumably led to a decrease in soil density and an increase in overall porosity. Researchers have pointed out that soil water content influences soil density (Lu et al., 2018; Tian et al., 2018; Tong et al., 2020). Naujokienė et al. (2018) reported that the use of differently prepared biopreparations reduced soil hardness by up to 28% and increased total porosity by up to

25% in the second year of the study, which resulted in lower diesel fuel consumption and reduced GHG emissions to the environment.

1.3. CO₂ emissions from soil

The agricultural sector is one of the most important GHG polluters of the environment, and cleaner production processes in this sector are of particular interest (Hamzei and Seyyedi, 2016; Wu et al., 2017). CO₂ emissions from soil are the second largest component of the carbon cycle and contribute to climate change (Mohammed et al., 2022). Agricultural producers are encouraged to increase agricultural production by developing alternative technologies that address climate change, i.e., reducing the carbon footprint of agriculture (Dias et al., 2016; Foley et al., 2011; Tilman et al., 2002). Soil bioactivity is the set of biological processes that determine soil respiration, enzyme activity, humification, and mineralization processes. A group of researchers (Ma et al., 2021) has observed that microorganism structure (community structure) and soil properties change together depending on environmental conditions and determine the dynamics of GHG emissions. After using the biological preparation, the amount of organic carbon in the soil increased from 1.8 to 2%, the difference in increase is 0.2% (Juknevičius et al., 2018). Stimulating soil microorganisms increases CO₂ release and improves nutrient mobilization (Klenz, 2015). Scientific results showed that the preparation of biocrusts biopreparation significantly improved soil physicochemical properties, respiration, and alkaline phosphatase, protease, and cellulose, and reduced CO₂ emissions in vegetation areas (Liu et al., 2017).

The dependence of soil respiration intensity, GHG emissions, and physical soil properties on tillage and other technological operations has already been studied quite extensively. However, the impact of environmentally friendly biopreparations on soil physical properties and the dynamics of CO₂ emissions during the growing season has not yet been sufficiently studied (Naujokienė et al., 2018). The limited number of scientific papers on this topic shows that research on the effects of biopreparations on soil under different meteorological conditions is new and relevant. The aim of this study was to investigate the effects of different biopreparation formulations on soil porosity, temperature, and CO₂ emission from the soil in Central Europe (Lithuania) growing food type crops.

2. Material and methods

2.1. Site description and experimental design

Experimental field research was carried out in 2014–2017 at the Experimental Station of Vytautas Magnus University Agriculture Academy (54°534'N, 23°50'E) in ugleytic satiated planosol (*Endohypogleyic-Eutric Planosol – PLe-gln-w*) (Buivydaite and Motuzas, 2001). Analysis of changes in soil physical properties and CO₂ emissions under the influence of biopreparations of different composition in North-East Europe (Lithuania) on the left bank of river Nemunas, in Kaunas district.

In the first year of the study, winter wheat (variety “Ada”) was grown, in the second year – winter wheat (“Famulus”) was grown, and in the third year – winter oilseed rape (“Cult”) was grown. Eight scenarios (SC) were selected to determine the effect of biopreparations on soil properties and CO₂ emissions from the soil, of which SC1 was the control with no biopreparations used. In the other seven SCs, biopreparations or mixtures of biopreparations were used. The components of the biopreparations are given in Table 1. The biopreparations were applied after the resumption of winter crops in the second half of April. The experimental plots were laid out in a linear pattern. The initial size of the plots was 600 m² and the reference size was 400 m² (Naujokienė et al., 2018, 2019). The layout of the experimental field scenarios is presented in Figure 1.




Fig. 1. Scheme of experimental field study scenarios

Table 1. Composition of the biopreparations used in different scenarios (Naujokienė et al., 2018, 2019)

The composition of biopreparations	Scenario							
	SC1	SC2	SC3	SC4	SC5	SC6	SC7	SC8
40 species of various herbs	-	+	+	-	+	-	+	-
Marine algae extracts	-	+	+	+	+	-	+	+
Essential oils of plants	-	+	+	-	+	-	+	-
Mineral oils	-	+	+	-	+	-	+	-
<i>Azospirillum spp.</i>	-	-	+	+	-	-	-	-
<i>Bacillus magetarium</i>	-	-	+	+	-	-	-	-
<i>Frateuria autentia</i>	-	-	+	+	-	-	-	-
<i>Azotobacter chroococcum</i>	-	-	-	-	+	-	-	+
<i>Azotospirillum brasilense</i>	-	-	-	-	+	-	-	+
4.5% of humic acids	-	-	-	-	-	+	+	-
0.5% gibberellic acid	-	-	-	-	-	+	+	-
0.01% copper (Cu)	-	-	-	-	-	+	+	-
0.01% zinc (Zn)	-	-	-	-	-	+	+	-
0.01% manganese (Mn)	-	-	-	-	-	+	+	-
0.01% iron (Fe)	-	-	-	-	-	+	+	-
0.01% calcium (Ca)	-	-	-	-	-	+	+	-
0.005% sodium molybdate (Na_2MoO_4)	-	-	-	-	-	+	+	-
Phosphorus P (P_2O_5)	-	-	-	-	+	-	-	+
Potassium K (K_2O)	-	-	-	-	+	-	-	+
<i>Azotobacter spp.</i>	-	-	-	-	-	+	+	-
Water (H_2O)	+	+	+	+	+	+	+	+

“+” – a compound is used; “-” – a compound is not used.

2.2. Measurements of soil physical properties

Soil properties were measured in April, May, June, July, and August over a three-year period. Depending on the meteorological conditions, a total of 14 tests were carried out (Table 2).

Table 2. Soil properties assessment plan (2015–2017)

2015	2016	2017
25.04.2015	29.04.2016	05.05.2017
11.05.2015	25.05.2016	30.05.2017
June was too dry	20.06.2016	28.06.2017
04.07.2015	20.07.2016 (after harvesting)	31.07.2017 (after harvesting)
07.08.2015 (after harvesting)	08.08.2016 (after soil tillage)	01.08.2017 (after soil tillage)

Soil porosity was sampled with a soil sampling drill from a depth of 0–10 cm. For each scenario, 5 measurements were taken. Soil porosity was determined with a vacuum air pycnometer after drying the samples to an air-dry mass. The total porosity P_b was calculated according to the formula (Maikštėnienė et al., 2007):

$$P_b = \left(1 - \frac{\rho_d}{\rho_{k.f.}} \right) \cdot 100, \quad (1)$$

where ρ_d – soil density, g cm⁻³;

$\rho_{k.f.}$ – soil solid phase density, g cm⁻³.

Aeration porosity $P_{aer.}$ was calculated according to the formula (Maikštėnienė et al., 2007):

$$P_{aer.} = P_b - (w \cdot \rho_d), \quad (2)$$

where w – soil water content, %.

Soil density was determined by weighing, taking samples with a Nekrasov drill and calculated according to the formula $\rho = m/v$ i.e., mass to volume ratio. The density of the solid phase was determined with a vacuum air pycnometer, after which the obtained results were inserted into the formulas presented in the article. Aeration porosity is a very important quantity for the soil, as it determines the amount of air spaces in the soil, and air is needed for plant roots to grow and develop normally.

Soil temperature at a depth of 0–5 cm in all treatments was determined with a hand-held portable device “HH2 Moisture Meter”, to which a “WET-2” type sensor was connected. The tests were carried out in 5 repetitions, and the depth of temperature measurement is indicated as 0-5 cm, as the rounding error is on the smaller side.

2.3. Measurement of CO₂ emissions from soil

CO₂ emissions from the soil were measured on the same dates as other physical soil properties. The measurements were carried out with the ADC BioScientific Lcpro+ System, a portable CO₂ gas analyzer consisting of a compact programming console, a soil respiration chamber, and a plastic ring to be inserted into

the soil. Carbon dioxide emissions were measured 5 times in each scenario. CO₂ gas emissions were measured in each repetition 5 times, the ring was placed in the soil at a depth of 20 mm, and all measurements were made in the first half of the day (from 10 a.m. to 2 p.m.). The soil temperature was measured in parallel with the measurement of CO₂ gas emissions.

The programming console is connected to the soil breathing chamber at the selected measurement location. A metal ring was inserted into the selected measurement site and the chamber attached to it. The ring is inserted perpendicular to the soil and left in place. The measurement site must be free of grass or other elements that could damage the sensors. The telescopic probe shall deliver CO₂ from the atmosphere at a height of 3 meters. This height was chosen to prevent the measurement from being influenced by the person taking the measurement. The measurement is carried out for 10 minutes, observing fluctuations in carbon dioxide. The data is automatically recorded on a memory stick.

2.4. Meteorological conditions

Meteorological data received from the Kaunas Meteorological Station (KMS). The distance between the KMS and the area where the experiments were conducted is approximately 500 m. The weather station provides multi-year data averages that are available calculated from 1974 until 2017 KMS provides multi-year data averages that are calculated since 1974 until 2017.

April 2015 was unusually warm. The average temperature for the month was 1 °C above the long-term average and precipitation was 7.6 mm above the long-term average (Fig. 2). May and June 2015 were 0.9 °C and 0.2 °C colder than the long-term average, with 10 mm of precipitation in May and 46.2 mm less than the long-term average in June. July 2015 was close to the long-term average, with 8.8 mm less precipitation than the long-term average. August was hot and dry, with an average air temperature of 20.3 °C and only 6.9 mm of precipitation. These data show that the 2015 growing season was very dry and deficient in moisture.




Fig. 2. Meteorological conditions during the study in 2015–2017

In April 2016, the average air temperature was 1 °C above the long-term average and in May it was 3.43 °C above the long-term average. April received 41.2 mm of precipitation, while May was a low-precipitation month, with only 36.4 mm, 17.4 mm below the long-term average. Warm and humid weather prevailed in summer. June was particularly warm, with an average air temperature of 17.21 °C, 1.61 °C above the long-term average. July and August were about 0.3 °C warmer than the long-term average. Compared to the long-

term average, precipitation was 21.1 mm higher in June, 81.7 mm higher in July, and 34.6 mm higher in August. The summer period of 2016 was humid.

Although the average temperature in April 2017 (5.61 °C) was close to the long-term average (6.1 °C), precipitation was 1.9 times higher than the long-term average. The weather in May was moderately warm and dry. The air temperature was 12.87 °C, 0.57 °C above the long-term average. Precipitation was very low, at just 10.5 mm, compared with the long-term average of 53.8 mm for May. The summer weather in Lithuania was humid and cool. The average temperature in June was no different from the long-term average, but precipitation was 1.28 times the long-term average. Meteorological conditions in July were close to the long-term average, with an air temperature of 16.77 °C and 79.6 mm of precipitation. The weather warmed up to 17.47 °C in August, with a long-term average of 16.6 °C. Precipitation in August was 25.3 mm lower than the long-term average. Precipitation in the summer of 2017 was in line with the long-term average.

2.5 Statistical analysis


One-way analysis of variance (ANOVA) was used to assess the statistical significance of the results. Dispersion analysis was performed on the LSD test for mathematical statistics (Raudonius, 2017; Olsson et al., 2007). We used the statistical software package SYSTAT, version 10. The probability level was indicated as follows: * – differences are significant at $P \leq 0.05 > 0.01$; ** – differences are significant at $P \leq 0.01 > 0.001$.

Non-parametric correlation analysis was applied to evaluate the causes of the studied traits. We used STAT and SIGMA PLOT software. The analysis matrix included data on normally distributed variables, such as: soil temperature and soil porosity forms. We calculated the correlations among all possible combinations.


3. Results and discussion

3.1. Soil total porosity

In the first year of the study (2015), the total porosity ranged from 50.4% to 55.4% before the application of the biopreparations (Fig. 3a). Two weeks after the spraying of the biopreparations (11 May 2015), the total porosity was measured and it was found that all treatments showed a decrease in total porosity ranging from 1.08% to 7.82%, except for treatments SC2 and SC8, which showed an increase in total porosity of 8.4% and 2.6% respectively. No studies were carried out in June due to drought. In July, total porosity varied from 47.5% to 52.2% for all treatments tested. Only one scenario, SC4, showed an increase in total porosity up to 3.9% compared to the total porosity found in May. Significant differences were obtained in scenarios SC3, SC4, SC7 and SC8. In August, the range of treatments in total porosity was between 47.9% and 51.7%. Significant differences were obtained in scenarios SC5 and SC6. Already in the first year of the study, a strong correlation between soil temperature and total porosity was found ($r_{2015} = -0.909$, $P \leq 0.01 > 0.001^{**}$).




LSD(A) _{.05} =2.16%	LSD(A) _{.01} =2.92%
LSD(B) _{.05} =4.17%	LSD(B) _{.01} =5.62%
LSD(D) _{.05} =2.62%	LSD(D) _{.01} =3.53%
LSD(E) _{.05} =2.77%	LSD(E) _{.01} =3.74%



LSD(A) _{.05} =3.03%	LSD(A) _{.01} =4.08%
LSD(B) _{.05} =3.89%	LSD(B) _{.01} =5.25%
LSD(C) _{.05} =3.22%	LSD(C) _{.01} =4.34%
LSD(D) _{.05} =3.11%	LSD(D) _{.01} =4.20%

LSD(E) _{.05} =5.04%	LSD(E) _{.01} =6.81%
------------------------------	------------------------------



LSD(A) _{.05} =3.03%	LSD(A) _{.01} =4.08%
LSD(B) _{.05} =6.15%	LSD(B) _{.01} =8.29%
LSD(C) _{.05} =4.28%	LSD(C) _{.01} =5.77%
LSD(D) _{.05} =5.00%	LSD(D) _{.01} =6.75%
LSD(E) _{.05} =4.07%	LSD(E) _{.01} =5.48%

Fig. 3. The effect of biopreparations on the dynamics of soil total porosity: a) 2015, b) 2016, c) 2017

Notes: SC1 – water (control); SC2 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, water; SC3 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, *Azospirillum spp.*, *Bacillus magetarium*, *Frateruria autentia*, water; SC4 – marine algae extracts, *Azospirillum spp.*, *Bacillus magetarium*, *Frateruria autentia*, water; SC5 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, *Azotobacter chroococcum*, *Azotospirillum brasilense*, Phosphorus, Potassium, water; SC6 – 4.5% of humic acids, 0.5% gibberellic acid, 0.01% copper (Cu), 0.01% zinc (Zn), 0.01% manganese (Mn), 0.01% iron (Fe), 0.01% calcium (Ca), 0.005% sodium molybdate (Na_2MoO_4), *Azotobacter spp.*, water; SC7 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, 4.5% of humic acids, 0.5% gibberellic acid, 0.01% copper (Cu), 0.01% zinc (Zn), 0.01% manganese (Mn), 0.01% iron (Fe), 0.01% calcium (Ca), 0.005% sodium molybdate (Na_2MoO_4), *Azotobacter spp.*, water; SC8 – marine algae extracts, *Azotobacter chroococcum*, *Azotospirillum brasilense*, Phosphorus, Potassium, water. * – significant differences from control treatment (SC1) at $P \leq 0.05$, ** – at $P \leq 0.01$ > 0.001. Intervals mean standard deviation.


In 2016, the post-winter soil total porosity ranged from 41.8% to 53.2% (Fig. 3b). Total porosity was measured on 23 May 2016 after the application of the biopreparations and showed an increase in total porosity in all SCs. In the control treatment SC1, an increase in total porosity was also found due to the meteorological conditions, as the warm and dry month of May prevailed. Carson et al. (2010) found that bacterial diversity increases with water potential ≤ 2.5 kPa in the sand and ≤ 4.0 kPa in silt + clay, which corresponds to a pore

space filled with $\leq 56\%$ water. The higher precipitation in June resulted in soil compaction, which reduced the total porosity in all scenarios except SC4, due to the presence of higher levels of microorganisms (*Azospirillum* sp., *Frateruria aurentia*, *Bacillus megaterium*) that prevented soil compaction. July was a high-precipitation month, which resulted in a decrease in total porosity of between 1.5% and 13% compared to June in all scenarios except SC1. In the control scenario, an increase of 3.4% was observed in July due to the filling of soil pores with water, which slightly increased the total porosity. In August, all scenarios showed an increase in total porosity compared to July, with the exception of scenarios SC1 and SC2, which consisted of non-bacterial components. These scenarios showed a decrease but not a significant one. Comparing the results obtained in April (before the application of the biopreparations) and August, it was found that the application of the biopreparations which were dominated by microorganisms, resulted in a more porous soil. The increase in total porosity ranged from 1.53% to 17.26% in most scenarios.


In 2017, total porosity at the beginning of May varied from 46.6% to 54.8% (Fig. 3c). Significant differences between the treatments compared to the control treatment were obtained in scenarios SC4, SC6, SC7, and SC8 at probability $P \leq 0.01 > 0.001$ and in scenario SC5 at $P \leq 0.05 > 0.01$. Biopreparations with higher bacterial content have a long-lasting effect, which is felt after overwintering with a higher total porosity index. The measurement of total porosity after spraying the biopreparations showed that in all SCs the total porosity increased from 18.54% to 26.54% because of the biopreparations and the environmental conditions. Scientists have found that biotreatments alter soil physicochemical properties (Banerjee, 2011; Cittenden et al., 2016). In June, when compared to the control, significant differences were obtained in scenarios SC6, SC7, and SC8 at $P \leq 0.01 > 0.001$, although almost all SCs showed a decrease in total porosity, except for scenario SC7, which used *Azotobacter* spp. bacteria in combination with mineral oils, seaweed, and various grass extracts, which affected total porosity. In 2017, the strong correlation was found between soil temperature and total porosity ($r_{2017} = 0.932^{**}$), with increasing temperature having a positive effect on total porosity. In 2017 in the first decade of May, very little precipitation fell (6.5 mm). This may have contributed to the positive correlation between soil temperature and total porosity. In 2015, almost 4 times more precipitation fell during the same period. Meteorological conditions are likely to have influenced the total porosity of the soil.

3.2 Soil aeration porosity


In the first year of the study, the aeration porosity before the application of the biopreparations ranged from 25.8% to 33.7% (Fig. 4a). Two weeks after the application of the biopreparations, the aeration porosity was measured as well. It was found that aeration porosity increased in all treatments, except for SC4, which showed a decrease of 10.13%. Scientists suggest that the application of biopreparations increases the organic carbon content of the soil, therefore decrease soil density and increase porosity (Montemurro et al., 2010; Peltre et al., 2015; Juknevičius et al., 2020). In July, aeration porosity varied from 23.8 % in SC1 to 30.0% in SC7 for the treatments studied. In August, the variation in aeration porosity ranged from 26.9% in SC2 to 32.4% in SC7 and SC8. After re-vegetation of plants, the SC1 option had the lowest aeration porosity, and after a month it increased 2.6 times, but in other options, where biological agents were used, the soil aeration porosity was found to be higher. It is due to meteorological conditions (soil moisture) and plant root system.



LSD(A) _{.05} =4.17%	LSD(A) _{.01} =5.62%
LSD(B) _{.05} =5.39%	LSD(B) _{.01} =7.28%
LSD(D) _{.05} =3.20%	LSD(D) _{.01} =4.31%
LSD(E) _{.05} =3.40%	LSD(E) _{.01} =4.59%



LSD(A) _{.05} =5.49%	LSD(A) _{.01} =7.41%
LSD(B) _{.05} =4.73%	LSD(B) _{.01} =6.38%
LSD(C) _{.05} =3.63%	LSD(C) _{.01} =4.89%
LSD(D) _{.05} =4.11%	LSD(D) _{.01} =5.54%
LSD(E) _{.05} =6.99%	LSD(E) _{.01} =9.43%



LSD(A) _{.05} =4.08%	LSD(A) _{.01} =5.50%
LSD(B) _{.05} =7.94%	LSD(B) _{.01} =10.72%
LSD(C) _{.05} =6.69%	LSD(C) _{.01} =9.02%
LSD(D) _{.05} =7.44%	LSD(D) _{.01} =10.04%
LSD(E) _{.05} =5.72%	LSD(E) _{.01} =7.71%

Fig. 4. The effect of biopreparations on the dynamics of soil aeration porosity: a) 2015, b) 2016, c) 2017
Notes: SC1 – water (control); SC2 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, water; SC3 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, *Azospirillum* spp., *Bacillus magetarium*, *Frateruria autentia*, water; SC4 – marine algae extracts, *Azospirillum* spp., *Bacillus magetarium*, *Frateruria autentia*, water; SC5 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, *Azotobacter chroococcum*, *Azotospirillum brasiliense*, Phosphorus, Potassium, water; SC6 – 4.5% of humic acids, 0.5% gibberellic acid, 0.01% copper (Cu), 0.01% zinc (Zn), 0.01% manganese (Mn), 0.01% iron (Fe), 0.01% calcium (Ca), 0.005% sodium molybdate (Na_2MoO_4), *Azotobacter* spp., water; SC7 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, 4.5% of humic acids, 0.5% gibberellic acid, 0.01% copper (Cu), 0.01% zinc (Zn), 0.01% manganese (Mn), 0.01% iron (Fe), 0.01% calcium (Ca), 0.005% sodium molybdate (Na_2MoO_4), *Azotobacter* spp., water; SC8 – marine algae extracts, *Azotobacter chroococcum*, *Azotospirillum brasiliense*, Phosphorus, Potassium, water. * – significant differences from control treatment (SC1) at $P \leq 0.05$, ** – at $P \leq 0.01$, *** – at $P \leq 0.001$. Intervals mean standard deviation.


In 2016, in the spring, at the resumption of vegetation, aeration porosity ranging from 12.1% in SC1 to 23.7% in SC5 (Fig. 4b). Other researchers (Yevtushenko et al., 2016) have found that aeration porosity was above 20% regardless of tillage technology. In our case, SC1 had the lowest aeration porosity of 12.08%, while in other SCs it was around 20%. Aeration porosity measurements taken two weeks after the application of the biopreparations showed an increase in aeration porosity in all the SCs compared to the April tests. The highest increase of 2.6 times in aeration porosity was found in the control scenario SC1. The increase in aeration porosity has been influenced not only by the sprayed biopreparations but also by favorable meteorological conditions. Many researchers suggest that porosity is particularly sensitive to tillage and environmental conditions (Cassaro et al., 2011, Lipiec et al., 2012, da Costa et al., 2014). The month of June was particularly warm with an average air temperature of 17.21 °C. The highest aeration porosity in June was found in SC4,

SC7, and SC8. The lowest aeration porosity of 23.7% was found in the control treatment. In July, the aeration porosity was similar to that in June. In August, all scenarios showed an increase in total porosity compared to July, except for scenarios SC1, SC2, and SC3.


In 2017, the aeration porosity at the beginning of May varied from 19.3% to 27.9% (Fig. 4c). Aeration porosity measurements after spraying biopreparations showed that in all SC2, aeration porosity increased compared to the measurements taken in May because of biopreparations and environmental conditions. The measurements carried out in the third decade of June showed a decrease in aeration porosity in all treatments compared to the measurements carried out in May. Researchers investigating effective microorganisms found no significant effect on porosity (Pranagal et al., 2020). At the end of July, aeration porosity ranged from 33.8% to 43.3%. After harvest, aeration porosity decreased in almost all scenarios except SC3 and SC8.

3.3 Soil temperature


Soil temperatures in April 2015 ranged from 10.5 °C to 11.8 °C (Fig. 5a). In April, compared to the control, soil temperature increased significantly in SC3, SC5, and SC7 scenarios ($P \leq 0.05 > 0.01$). Soil temperature is one of the most important variables influencing soil respiration and depends on environmental conditions (Moyano et al., 2013, Sierra et al., 2015). Our studies have also shown that environmental conditions have an effect on temperature changes, i.e. the use of a biopreparation influences the increase in temperature. As the soil gradually warmed in May, soil temperatures were found to be about 2–5 °C higher than in May. Compared to the control scenario SC1, significantly lower soil temperatures were observed in SC7 and SC8 at $P \leq 0.01 > 0.001$ and $P \leq 0.05 > 0.01$ in SC6. In July, the soil temperature ranged from 19.08 °C (SC7) to 22.04 °C (SC2). In July, a significant decrease in soil temperature was found between control SC1 and SC6 ($P \leq 0.05 > 0.01$) and between SC1 and SC7 ($P \leq 0.01 > 0.001$) predicting that there is a denser crop. In August, the soil temperature was the highest recorded, as the absence of vegetative cover resulted in a significant warming of the soil. Researchers (Dai et al., 2021) found that soil temperature was lowest in the non-arable soil with straw mulch. The uneven spread of crop residues after harvest decreased soil temperature. Significant increases were found between scenarios SC1 and SC6, SC1 and SC7 at the 95% probability level, and between SC1 and SC2, SC1 and SC8 at the 99% probability level.



LSD(A) _{.05} =1.10 °C	LSD(A) _{.01} =1.48 °C
LSD(B) _{.05} =0.48 °C	LSD(B) _{.01} =0.65 °C
LSD(D) _{.05} =1.20 °C	LSD(D) _{.01} =1.62 °C
LSD(E) _{.05} =3.12 °C	LSD(E) _{.01} =4.21 °C



LSD(A) _{.05} =0.71 °C	LSD(A) _{.01} =0.95 °C
LSD(B) _{.05} =1.41 °C	LSD(B) _{.01} =1.90 °C
LSD(C) _{.05} =0.91 °C	LSD(C) _{.01} =1.23 °C
LSD(D) _{.05} =1.91 °C	LSD(D) _{.01} =2.58 °C
LSD(E) _{.05} =1.34 °C	LSD(E) _{.01} =1.81 °C



LSD(A) _{.05} =0.67 °C	LSD(A) _{.01} =0.91 °C
LSD(B) _{.05} =2.37 °C	LSD(B) _{.01} =3.19 °C
LSD(C) _{.05} =1.85 °C	LSD(C) _{.01} =2.50 °C
LSD(D) _{.05} =1.42 °C	LSD(D) _{.01} =1.92 °C
LSD(E) _{.05} =2.50 °C	LSD(E) _{.01} =3.37 °C

Fig. 5. The effect of biopreparations on the dynamics of soil temperature: a) 2015, b) 2016, c) 2017 Notes: SC1 – water (control); SC2 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, water; SC3 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, Azospirillum spp., Bacillus magetarium, Frateuria autentia, water; SC4 – marine algae extracts, Azospirillum spp., Bacillus magetarium, Frateuria autentia, water; SC5 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, Azotobacter chroococcum, Azotospirillum brasilense, Phosphorus, Potassium, water; SC6 – 4.5% of humic acids, 0.5% gibberellic acid, 0.01% copper (Cu), 0.01% zinc (Zn), 0.01% manganese (Mn), 0.01% iron (Fe), 0.01% calcium (Ca), 0.005% sodium molybdate (Na₂MoO₄), Azotobacter spp., water; SC7 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, 4.5% of humic acids, 0.5% gibberellic acid, 0.01% copper (Cu), 0.01% zinc (Zn), 0.01% manganese (Mn), 0.01% iron (Fe), 0.01% calcium (Ca), 0.005% sodium molybdate (Na₂MoO₄), Azotobacter spp., water; SC8 – marine algae extracts, Azotobacter chroococcum, Azotospirillum brasilense, Phosphorus, Potassium, water. * – significant differences from control treatment (SC1) at $P \leq 0.05$, ** – at $P \leq 0.01$. Intervals mean standard deviation.


In 2016, soil temperatures varied from 13.2 °C to 14.1 °C after the resumption of plant growth (Fig. 5b). A significant increase was found between scenarios SC1 and SC5 ($P \leq 0.01 > 0.001$) and between SC1 and SC8 ($P \leq 0.05 > 0.01$). At the end of May, soil temperature increased on average by about 10 °C. In June, soil temperature ranged from 22 °C (SC6) to 23.3 °C (SC4). In the SC4 scenario, soil temperature was significantly higher than the control ($P \leq 0.05 > 0.01$). In July and August, soil temperatures were found to be similar due to the settled weather, and no significant differences were found between the scenarios and the control.

On 5 May 2017, the highest soil temperature was found in SC8 and the lowest in the control scenario (Fig. 5c). Soil temperatures were significantly higher in scenarios SC2 and SC5 ($P \leq 0.05 > 0.01$), SC6, SC7, and SC8 ($P \leq 0.01 > 0.001$) compared to the control scenario. Soil warming at the end of May resulted in a significant increase in soil temperature in all scenarios except SC2 at the 99% probability level. In SC8, a substantial increase was found at the 95% probability level compared to the control SC1. In June, the lowest soil temperature of 16.66 °C was found in SC1 and the highest of 21 °C in SC7. Significantly higher soil temperatures compared to the control were found in scenarios SC4, SC6, SC7, and SC8 at the 99% probability level. An increase in soil thermal conductivity increases temperature whereas an increase in soil heat capacity reduces temperature (Obia et al, 2020). At the end of July, soil temperatures ranged from 21.8 to 24.3 °C. A significant increase was found between control and SC4 ($P \leq 0.01 > 0.001$) and between control and SC5 and SC6 scenarios ($P \leq 0.05 > 0.01$). On 1 August, soil temperature increased in most of the scenarios studied compared to soil temperature at the end of July. However, a significant decrease in soil temperature ($P \leq 0.01 > 0.001$) was obtained after the harvest between control and SC5, and at the 95% probability level soil temperature was significantly lower in the control scenario than in SC4, SC6, and SC7. In all scenarios the higher soil surface temperatures were due to tillage, which allowed warm air to enter the soil. The data of other authors do not confirm these researches, because all the research of ours and other authors were in other spheres of soil composition and climate, so this was added as additional information, as it was obtained in other countries, but perfectly parallel studies were not found, only similar ones, due to the soil and the diversity of the area.


3.4 CO₂ emissions from soil

In April 2015, the highest CO₂ emissions were observed in scenario SC7 with 2.29 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 6a) and the lowest in SC2 with 0.73 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Soil moisture, temperature, and biopreparation composition were the main influences on soil respiration. Research groups suggest that soil moisture influences CO₂ emission, with continuous moisture conditions increasing the bacterial content of the soil, resulting in higher CO₂ emissions from the soil compared to reirrigation (Jiao et al. 2023; Gultekin et al., 2023; Barnard et al., 2015). Tillage technology also has an impact, as tilled soil emits up to 21% more CO₂, but this depends on the soil type, organic carbon, and microorganism content of the soil (Abdalla et al., 2016; Chaplot et al., 2015; Huang et al., 2013). Canarini et al. (2017) found that in soils with more than 2% organic carbon, CO₂ emissions increase after drought, in contrast to soils with low carbon content. In our case, substantial increases between the control SC1 and SC5 scenarios were found at the 95% probability level, and between SC1 and SC7 at the 99% probability level. In May, soil respiration increased or remained the same compared to April. A substantial increase was found between the control and scenarios SC4, SC5, SC6, SC7, and SC8 at the 99% probability level. It is likely that the bacteria present in the bioassay (*Azospirillum sp.*, *Frateruria aurentia*, *Bacillus megaterium*, *Azotobacter chroococum*, *Azospirillum brasilense*, *Azobacter vinelandii*) contributed to the substantial increases. Scientists suggest that soil microorganisms can increase CO₂ release (Klenz, 2015) under certain environmental conditions. According to the results of the May study, soil temperature had a significant influence on CO₂ emissions. The May results showed a strong correlation between soil temperature and CO₂ emissions ($r_{2015} = -0.903^{**}$), with rising temperature reducing emissions. Although in the following months of July and August, CO₂ emissions were increasing due to higher ambient temperatures in all scenarios. A group of researchers (Tóth et al., 2018) investigated that soil emissions may be higher during the growing season. In August, the highest CO₂ emissions were found in SC3 at 4.26 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and the lowest in SC2 at 2.15 $\mu\text{mol m}^{-2} \text{s}^{-1}$.


$\text{m}^{-2} \text{s}^{-1}$. Significant increases between control and SC8 were found at the 95% probability level, and at the 99% probability level – between scenarios SC1 and SC4.



LSD(A) ₀₅ =0.78 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(A) ₀₁ =1.05 $\mu\text{mol m}^{-2} \text{s}^{-1}$
LSD(B) ₀₅ =0.53 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(B) ₀₁ =0.71 $\mu\text{mol m}^{-2} \text{s}^{-1}$
LSD(D) ₀₅ =0.76 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(D) ₀₁ =1.03 $\mu\text{mol m}^{-2} \text{s}^{-1}$
LSD(E) ₀₅ =0.71 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(E) ₀₁ =0.96 $\mu\text{mol m}^{-2} \text{s}^{-1}$
LSD(A) ₀₅ =0.78 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(A) ₀₁ =1.05 $\mu\text{mol m}^{-2} \text{s}^{-1}$



LSD(A) ₀₅ =0.30 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(A) ₀₁ =0.40 $\mu\text{mol m}^{-2} \text{s}^{-1}$
LSD(B) ₀₅ =0.65 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(B) ₀₁ =0.88 $\mu\text{mol m}^{-2} \text{s}^{-1}$
LSD(C) ₀₅ =0.45 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(C) ₀₁ =0.60 $\mu\text{mol m}^{-2} \text{s}^{-1}$
LSD(D) ₀₅ =0.67 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(D) ₀₁ =0.90 $\mu\text{mol m}^{-2} \text{s}^{-1}$
LSD(E) ₀₅ =0.77 $\mu\text{mol m}^{-2} \text{s}^{-1}$	LSD(E) ₀₁ =1.04 $\mu\text{mol m}^{-2} \text{s}^{-1}$



LSD(A) ₀₅ =0.34 μmol m ⁻² s ⁻¹	LSD(A) ₀₁ =0.46 μmol m ⁻² s ⁻¹
LSD(B) ₀₅ =0.39 μmol m ⁻² s ⁻¹	LSD(B) ₀₁ =0.52 μmol m ⁻² s ⁻¹
LSD(C) ₀₅ =0.36 μmol m ⁻² s ⁻¹	LSD(C) ₀₁ =0.49 μmol m ⁻² s ⁻¹
LSD(D) ₀₅ =0.42 μmol m ⁻² s ⁻¹	LSD(D) ₀₁ =0.57 μmol m ⁻² s ⁻¹
LSD(E) ₀₅ =1.27 μmol m ⁻² s ⁻¹	LSD(E) ₀₁ =1.72 μmol m ⁻² s ⁻¹

Fig. 6. The effect of biopreparations on the dynamics of CO₂ emissions from soil a) 2015, b) 2016, c) 2017. Notes: SC1 – water (control); SC2 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, water; SC3 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, Azospirillum spp., Bacillus magetarium, Frateuria autentia, water; SC4 – marine algae extracts, Azospirillum spp., Bacillus magetarium, Frateuria autentia, water; SC5 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, Azotobacter chroococcum, Azotospirillum brasilense, Phosphorus, Potassium, water; SC6 – 4.5% of humic acids, 0.5% gibberellic acid, 0.01% copper (Cu), 0.01% zinc (Zn), 0.01% manganese (Mn), 0.01% iron (Fe), 0.01% calcium (Ca), 0.005% sodium molybdate (Na₂MoO₄), Azotobacter spp., water; SC7 – 40 species of various herbs, marine algae extracts, essential oils of plants, mineral oils, 4.5% of humic acids, 0.5% gibberellic acid, 0.01% copper (Cu), 0.01% zinc (Zn), 0.01% manganese (Mn), 0.01% iron (Fe), 0.01% calcium (Ca), 0.005% sodium molybdate (Na₂MoO₄), Azotobacter spp., water; SC8 – marine algae extracts, Azotobacter chroococcum, Azotospirillum brasilense, Phosphorus, Potassium, water. * – significant differences from control treatment (SC1) at $P \leq 0.05 > 0.01$, ** – at $P \leq 0.01 > 0.001$. Intervals mean standard deviation.

In 2016, CO₂ emissions from the resumption of vegetation ranged from 0.69 μmol m⁻² s⁻¹ to 1.06 μmol m⁻² s⁻¹ (Fig. 6b). Other researchers (Forte et al., 2017) who studied conventional tillage reported that it leads to higher CO₂ emissions due to higher decomposition rates of soil organic matter and higher temperature fluctuations. At the end of May, soil respiration was more intense in all scenarios, and a substantial increase compared to the control was found in SC7 and SC8 at $P \leq 0.05 > 0.01$. In June, with a decrease in air temperature of 1-2 °C, CO₂ release slowed down and varied between 1.31 μmol m⁻² s⁻¹ and 2.64 μmol m⁻² s⁻¹. A significant increase at $P \leq 0.01 > 0.001$ was found between SC1 and SC7. In July, CO₂ emissions increased, but at the 95% confidence level, a significant increase was found between SC1 and SC7. Soil respiration increased further after harvest in August, ranging from 2.79 to 5.78 μmol m⁻² s⁻¹. CO₂ emissions increased significantly compared to the control in SC6 and SC7 at the 99% probability level and in SC5 at the 95% probability level.

At the beginning of May 2017, CO₂ emissions were in the range of 0.55-1.37 μmol m⁻² s⁻¹ (Fig. 6c). In the SC1 scenario, soil respiration was found to peak at 1.37 μmol m⁻² s⁻¹. At the end of May, all scenarios had higher CO₂ emissions due to ambient conditions and were significantly different from the control, with all SCs at the 99% probability level except SC2. Many field experiments have shown that CO₂ is significantly and positively correlated with soil organic carbon (Liu et al., 2014) and soil temperature (Cartwright and Hui, 2014), but in our case, it is the opposite. In June, CO₂ emissions increased significantly in SC1 compared to SC2 ($P \leq 0.05 > 0.01$) and to the SC4 scenario ($P \leq 0.01 > 0.001$). Significant reductions were obtained in scenarios SC5 ($P \leq 0.05 > 0.01$) and SC3 ($P \leq 0.01 > 0.001$). The settled temperature in July, which was close

to the long-term average (around 10 °C), resulted in more intense soil respiration in all scenarios except SC7 and SC8. A significant decrease ($P \leq 0.01 > 0.001$) was found between the control SC1 and SC7, SC8 scenarios, which could be influenced by the different compositions of the biopreparations. Drulis et al. (2022) state that bioproducts are substances that can improve crop productivity and quality, increase nutrient availability in the soil, improve plant nutrient use efficiency, and promote organic matter decomposition and humification in the soil. In all scenarios, CO₂ emissions increased by a factor of 3 to 5 in the range of 4.89-11.07 $\mu\text{mol m}^{-2} \text{s}^{-1}$ after the harvest and tillage in August. These changes are due to the fact that the study was carried out immediately after tillage. The process of tillage greatly intensifies CO₂ emissions to the environment (Buragienė et., 2015). However, the differences between the scenarios were influenced by the different compositions of the biopreparations. Emissions were significantly higher only in scenario SC4 compared to the control ($P \leq 0.01 > 0.001$) and significantly lower in scenarios SC3, SC7, and SC8 compared to the control ($P \leq 0.01 > 0.001$).

Data analysis showed average and strong linear correlations between soil temperature, CO₂ emission, total and aeration porosity (Table 3). In 2015 and 2017, we strong negative correlations between soil temperature and CO₂ emission. In 2016, the opposite correlation was found. In 2016 were wetter compared to 2015 and 2017, especially the months of July and August. Amount of precipitation in 2016 during the vegetation period of the plants was evenly distributed, there were no periods of drought. Meanwhile, in 2015 and 2017, drier periods were identified when no more than 5 mm of precipitation fell per decade.

Soil temperature also correlated with soil porosity, however relations in 2015 were negative and in 2016-2017 – positive.

Table 3. Correlations between soil properties

Independent variables, <i>x</i>	Dependent variables, <i>Y</i>			
	Temperature, °C	CO ₂ emission, $\mu\text{mol m}^{-2} \text{s}^{-1}$	Total porosity, %	Aeration porosity, %
2015				
Temperature, °C	1.00	-0.914**	-0.752*	-0.856**
CO ₂ emission, $\mu\text{mol m}^{-2} \text{s}^{-1}$	-	1.00	0.712*	0.755*
Total porosity, %	-	-	1.00	0.986**
2016				
Temperature, °C	1.00	0.725*	0.804*	0.771*
CO ₂ emission, $\mu\text{mol m}^{-2} \text{s}^{-1}$	-	1.00	0.855**	0.824*
Total porosity, %	-	-	1.00	0.923**
2017				
Temperature, °C	1.00	-0.849**	0.822*	0.762*
CO ₂ emission, $\mu\text{mol m}^{-2} \text{s}^{-1}$	-	1.00	-0.728*	-0.842**
Total porosity, %	-	-	1.00	0.900**

Notes: * - significant at $P \leq 0.05 > 0.01$; ** - at $P \leq 0.01 > 0.001$.

Soil CO₂ emission correlated with soil porosity, however in 2017 this relation was negative. In addition, soil porosity forms closely correlated with each other.

Conclusions

-In the first and second years of the study, the total porosity of the soil varied between 41% and 62%, while in the third year, the total porosity of the soil increased in all scenarios and over the whole study period ranging from 51% to 74%. This increase was due to the interaction between the long-term use of biopreparations and

meteorological conditions.

-In the first year, soil temperature in August showed a significant increase compared to the control ($P \leq 0.05 > 0.01$) in scenarios SC6, SC7, SC2, and SC8. Similar trends were confirmed in the second and third years.

-The use of biopreparations had an impact on CO₂ emissions from soil. In the first year, it was found that, just scenario SC2 reduced CO₂ emissions from soil. The cumulative effect of biopreparation application was most pronounced in the third year.

-Tillage intensifies CO₂ emissions from the soil, these studies confirmed that biopreparations (SC3, SC7, SC8) can significantly reduce the CO₂ emission intensity from the soil after tillage, predictable due to the overlap of biocomposite components such as Marine algae extracts and bacteria.

-Data analysis showed average and strong linear correlations between soil temperature, CO₂ emission, total and aeration porosity. In 2015 and 2017, we strong negative correlations between soil temperature and CO₂ emission. In 2016, the opposite correlation was found due to higher precipitation.

Future research on the use of bacteria-based and environmentally friendly bioproducts should focus on increasing CO₂ storage in soil, simplifying agricultural operations, reducing inputs, and increasing the efficiency of crop production.

References

1. Abdalla K., Chivenge P., Ciais P., Chaplot V., 2016. No-tillage lessens soil CO₂ emissions the most under arid and sandy soil conditions: results from a meta-analysis. *Biogeosciences*. 13, 3619–3633. <https://doi.org/10.5194/bg-13-3619-2016>.
2. Bais H.P., Weir T.L., Perry L.G., Gilroy S., Vivanco J.M., 2006. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu. Rev. Plant Biol.* 57, 233–266. <https://doi.org/10.1146/annurev.arplant.57.032905.105159>.
3. Banerjee D., 2011. Endophytic Fungal Diversity in Tropical and Subtropical Plants. *Res. J. Microbiol.* 6, 54–62. <https://doi.org/10.3923/jm.2011.54.62>.
4. Barnard R.L., Osborne C.A., Firestone M.K., 2015. Changing precipitation pattern alters soil microbial community response to wet-up under a Mediterranean-type climate. *The ISME Journal*. 9, 946–957. <https://doi.org/10.1038/ismej.2014.192>.
5. Błaszczuk L., Siwulski M., Sobieralski K., Lisiecka J., Jędrzycka M., 2014. *Trichoderma* spp. – application and prospects for use in organic farming and industry. *Journal of Plant Protection Research*. 54 (4), 309–317. <https://doi.org/10.2478/jppr-2014-0047>.
6. Buivydaitė V., Juodis J., Motuzas A., Vaičys M. Lietuvos dirvožemių klasifikacijos // Lietuvos dirvožemiai: monografija / sudaryt. V.Vasiliauskienė. - Vilnius: PĮ „Lietuvos mokslas“, 2001, p. 244-408.
7. Bünemann E.K., Marscher P., Smernik R.J., Conyers M., McNeill A.M. 2008. Soil organic phosphorus and microbial community composition as affected by 26 years of different management strategies. *Biol. Fertil. Soils*. 44, 717–726. <https://doi.org/10.1007/s00374-007-0254-2>.
8. Buragienė S., Šarauskis E., Romaneckas K., Sasnauskienė J., Masilionytė L., Kriaučiūnienė Z., 2015. Experimental analysis of CO₂ emissions from agricultural soils subjected to five different tillage systems in Lithuania. *Science of the Total Environment*. 514, 1–9. <https://doi.org/10.1016/j.scitotenv.2015.01.090>.
9. Canarini A., Kiær L.P., Dijkstra F.A., 2017. Soil carbon loss regulated by drought intensity and available substrate: A meta-analysis. *Soil Biology and Biochemistry* 112, 90–99. <https://doi.org/10.1016/j.soilbio.2017.04.020>.
10. Carson J.K., Gonzalez-Quinones V., Murphy D.V., Hinz C., Shaw J.A., Gleeson D.B., 2010. Low pore connectivity increases bacterial diversity in soil. *Applied and Environmental Microbiology*. 76, 3936. <https://doi.org/10.1128/AEM.03085-09>.

11. Cartwright J., Hui D.F., 2015. Soil respiration patterns and controls in limestone cedar glades. *Plant Soil*. 389, 157–169. <https://doi.org/10.1007/s11104-014-2348-6>.
12. Caruso G., De Pascale S., Cozzolino E., Giordano M., El-Nakhel C., Cuciniello A., Cenvinzo V., Colla G., Roupahel Y., 2019. Protein Hydrolysate or Plant Extract-based Biostimulants Enhanced Yield and Quality Performances of Greenhouse Perennial Wall Rocket Grown in Different Seasons. *Plants*. 8, 208. <https://doi.org/10.3390/plants8070208>.
13. Cassaro F.A.M., Borkowski A.K., Pires L.F., Saab S.D.C., 2011. Characterization of a Brazilian clayey soil submitted to conventional and no-tillage management practices using pore size distribution analysis. *Soil Tillage Res.*, 111, 175–179. <https://doi.org/10.1016/j.still.2010.10.004>.
14. Chaplot V., Abdalla K., Alexis M., Bourennane H., Darboux F., Dlamini P., Everson C., Mchunu C., Muller-Nedebock D., Mutema M., Quenea K., Thenga H., Chivenge P., 2015. Surface organic carbon enrichment to explain greater CO₂ emissions from short-term no-tilled soils. *Agr. Ecosyst. Environ.*, 203, 110–118. <http://dx.doi.org/10.1016/j.agee.2015.02.001>.
15. Crittenden, S., Goede, R., 2016. Integrating soil physical and biological properties in contrasting tillage systems in organic and conventional farming. *Eur. J. Soil Biol.* 77, 26–33. <https://doi.org/10.1016/j.ejsobi.2016.09.003>.
16. da Costa P. A., Mota J. C. A., Romero R. E., Freire A. G., Ferreira T. O., 2014. Changes in soil pore network in response to twenty-three years of irrigation in a tropical semiarid pasture from northeast Brazil. *Soil Tillage Res.*, 137, 23–32. <https://doi.org/10.1016/j.still.2013.11.004>.
17. Dai Z., Hu J., Fan J., Fu W., Wang H., Hao M., 2021. No-tillage with mulching improves maize yield in dryland farming through regulating soil temperature, water and nitrate-N. *Agriculture, Ecosystems & Environment*, 309, 1–12. <https://doi.org/10.1016/j.agee.2020.107288>.
18. Dias G.A., Rocha R.H.C., Araújo J.L., De Lima J.F., Guedes W.A., 2016. Growth, yield, and postharvest quality in eggplant produced under different foliar fertiliser (*Spirulina platensis*) treatments. *Semin. Cienc. Agrar.* 37, 3893–3902. <https://doi.org/10.5433/1679-0359.2016v37n6p3893>.
19. Drulis S.P., Kriaučiūnienė Z., Liakas V., 2022. The Influence of Different Nitrogen Fertilizer Rates, Urease Inhibitors and Biological Preparations on Maize Grain Yield and Yield Structure Elements. *Agronomy*, 12(3). 1–15. <https://doi.org/10.3390/agronomy12030741>.
20. Ertani A., Francioso O., Tinti A., Schiavon M., Pizzeghello D., Nardi S., 2018. Evaluation of seaweed extracts from *Laminaria* and *Ascophyllum nodosum* spp. as biostimulants in *Zea mays* L. using a combination of chemical, biochemical and morphological approaches. *Front. Plant Sci.* 9, p. 428. <https://doi.org/10.3389/fpls.2018.00428>.
21. European Commission. European Green Deal. 2020. <https://ec.europa.eu/info/strategy/priorities-2019-2024/european-green-deal> (accessed 16 November 2022).
22. Foley J.A., Ramankutty N., Brauman K.A., Cassidy E.S., Gerber J.S., Johnston M., Mueller N.D., O'Connell C., Ray D.K., West P.C., et al. 2011. Solutions for a cultivated planet. *Nature*, 478, 337–342. <https://doi.org/10.1038/nature10452>.
23. Forte A., Fiorentino N., Fagnano M., Fierro A., 2017. Mitigation impact of minimum tillage on CO₂ and N₂O emissions from a Mediterranean maize cropped soil under low-water input management. *Soil Tillage Res.*, 166, 167–178. <https://doi.org/10.1016/j.still.2016.09.014>.
24. Gil S.V., Meriles J., Conforto C., Basanta M., Radl V., Hagn A., et al. 2011. Response of soil microbial communities to different management practices in surface soils of a soybean agroecosystem in Argentina. *Eur. J. Soil Biol.* 47, 55–60. <https://doi.org/10.1016/j.ejsobi.2010.11.006>.
25. Gultekin R., Avağ R., Görgişen C., Öztürk Ö., Yeter T., Bahçeci Alsan P., 2023. Effect of deficit irrigation practices on greenhouse gas emissions in drip irrigation. *Scientia Horticulturae*. 310, 111757. <https://doi.org/10.1016/j.scienta.2022.111757>.

26. Hamzei J., Seyyedi M., 2016. Energy use and input–output costs for sunflower production in sole and intercropping with soybean under different tillage systems. *Soil Tillage Res.* 157, 73–82. <https://doi.org/10.1016/j.still.2015.11.008>.
27. Huang M., Jiang L., Zou Y., Xu S., Deng G. 2013. Changes in soil microbial properties with no-tillage in Chinese cropping systems. *Biol Fertil. Soils*, 49, p. 373–377. <https://doi.org/10.1007/s00374-013-0778-6>.
28. Jiao P., Yang L., Nie X., Li Z., Liu L., Zheng P., 2023. Dependence of cumulative CO₂ emission and microbial diversity on the wetting intensity in drying-rewetting cycles in agriculture soil on the Loess Plateau. *Springer, Soil Ecology Letters*. 5, 220147. <https://doi.org/10.1007/s42832-022-0147-1>.
29. Juknevičius D., Kriauciūnienė Z., Jasinskas A., Šarauskis E., 2020. Analysis of changes in soil organic carbon, energy consumption and environmental impact using bio-products in the production of winter wheat and oilseed rape. *Sustainability*. 12(19), 8246. <https://doi.org/10.3390/su12198246>.
30. Juknevičius D., Šarauskis E., Rimkuvienė D., Rukaite J., Karayel D., 2018. Effect of biological preparation on changes in soil organic carbon and its environmental impact applying precision farming. *Engineering for Rural Development*. Jelgava. <https://doi.org/10.22616/ERDev2018.17.N398>.
31. Khattab R.Y., Arntfield S.D., Nyachoti C.M., 2009. Nutritional quality of legume seeds as affected by some physical treatments, Part 1: Protein quality evaluation. *LWT - Food Science and Technology*. 42. 6. 1107–1112. <https://doi.org/10.1016/j.lwt.2009.02.008>.
32. Klenz K., 2015. Promoting soil integrity: the use of cover crops and a microorganism stimulant on a striptill corn field in southeastern Minnesota. *Natural Lands Ecology Papers*. 1–35.
33. Kocira S., Hara P., Szparaga A., Czerwińska E., Beloev H., Findura P., Bajus P. 2020. Evaluation of the Effectiveness of the Use of Biopreparations as Seed Dressings. *Agriculture*. 10(4), 90. <https://doi.org/10.3390/agriculture10040090>.
34. Kong C.H., Wang P., Zhao H., Xu X.H., Zhu Y.D., 2008. Impact of allelochemical exuded from allelopathic rice on soil microbial community. *Soil Biol. Biochem.* 40, 1862–1869. <https://doi.org/10.1016/j.soilbio.2008.03.009>.
35. Lipiec J., Hajnos M., Swieboda R., 2012. Estimating effects of compaction on pore size distribution of soil aggregates by mercury porosimeter. *Geoderma*. 179–180, 20–27. <https://doi.org/10.1016/j.geoderma.2012.02.014>.
36. Liu X. P., Zhang W.J., Hu C.S., Tang X.G. 2014. Soil greenhouse gas fluxes from different tree species on Taihang Mountain, North China. *Biogeosciences*, 11, 1649–1666. <https://doi.org/10.5194/bg-11-1649-2014>.
37. Liu Y., Xing Z., Yang H., 2017. Effect of biological soil crusts on microbial activity in soils of the Tengger Desert (China). *Journal of Arid Environment*. 144, 201–211. <https://doi.org/10.1016/j.jaridenv.2017.04.003>.
38. Lorenzo P., Pereira C.S., Rodríguez-Echeverría S., 2013. Differential impact on soil microbes of allelopathic compounds released by the invasive *Acacia dealbata* link. *Soil Biol. Biochem.* 57, 156–163. <https://doi.org/10.1016/j.soilbio.2012.08.018>.
39. Lu Y., Horton R., Ren T., 2018. Simultaneous determination of soil bulk density and water content: A heat pulse-based method. *European Journal of Soil Science*. 69, 947–952. <https://doi.org/10.1111/ejss.12690>.
40. Ma S., Fan J., Chen Y., Lu X. 2021. Studying greenhouse gas emissions through interactions between phospholipid fatty acid content and soil properties of alpine grassland soil in Northern Tibet, China. *Global Ecology and Conservation*, 27, 1–16. <https://doi.org/10.1016/j.gecco.2021.e01558>.

41. Maikštėnienė S., Šlepetienė A., Masilionytė L., 2007. The effect of mouldboard nouldbo and on energetic efficiency of agrosystems. *Zemdirbyste=Agriculture*. 94, 3-23.
42. Michalak I., Chojnacka K., Dmytryk, A., Wilk R., Gramza M., Rój E., 2016. Evaluation of supercritical extracts of algae as biostimulants of plant growth in field trials. *Front. Plant Sci.* 7, 1591. <https://doi.org/10.3389/fpls.2016.01591>.
43. Michałek W., Kocira A., Findura P., Szparaga A., Kocira S., 2018. The Influence of Biostimulant Asahi SL on the Photosynthetic Activity of Selected Cultivars of *Phaseolus vulgaris* L. *Rocz. Ochr. Sr.* 20, 1286–1301.
44. Mohammed S., Mirzaei M., Toro A. P., Anari M. G., Moghiseh E., Asadi H., Szabo S., Kakuszi-Szeles A., Harsany E. 2022. Soil carbon dioxide emissions from maize (*Zea mays* L.) fields as influenced by tillage management and climate. *Irrig. and Drain.* 71, 228–240. <https://doi.org/10.1002/ird.2633>.
45. Montemurro F., Ferri D., Tittarelli F., Canali S., Vitti C., 2010. Anaerobic Digestate and On-Farm Compost Application: Effects on Lettuce (*Lactuca sativa* L.) Crop Production and Soil Properties. *Compost Science & Utilization*, 18:3, 184–193, <https://doi.org/10.1080/1065657X.2010.10736954>.
46. Moyano F.E., Manzoni S., Chenu C., 2013. Responses of soil heterotrophic respiration to moisture availability: an exploration of processes and models. *Soil Biology and Biochemistry*, 59, 72–85. <https://doi-org/10.1016/j.soilbio.2013.01.002>.
47. Naujokienė V., Šarauskis E., Lekavičienė K., Adamavičienė A., Buragienė S., Kriaučiūnienė Z., 2018. The influence of biopreparations on the reduction of energy consumption and CO₂ emissions in shallow and deep soil tillage. *Science of The Total Environment*. 626, 1402–1413. <https://doi.org/10.1016/j.scitotenv.2018.01.190>.
48. Naujokienė, V., Šarauskis, E., Bleizgys, R. and Sasnauskienė, J., 2019. Soil biotreatment effectiveness for reducing global warming potential from main polluting tillage operations in life cycle assessment phase. *Science of the Total Environment*, 671, pp.805-817.
49. Nostro A., Germano M.P., D'angelo V., Marino A., Cannatelli M.A., 2000. Extraction methods and bioautography for evaluation of medicinal plant antimicrobial activity. *Lett. Appl. Microbiol.* 30, 379–384. <https://doi.org/10.1046/j.1472-765x.2000.00731.x>.
50. Obia A., Cornelissen G., Martinsen V., Smebye A.B., Mulder J., 2020. Conservation tillage and biochar improve soil water content and moderate soil temperature in a tropical Acrisol. *Soil and Tillage Research*. 197, 1–12. <https://doi-org/10.1016/j.still.2019.104521>.
51. Olsson U., Engstrand U., Rupšys P., 2007. *Statistiniai metodai/Mokomoji knyga*. Akademija. p. 138.
52. Oskiera M., Szczech M., Bartoszewski G., 2017. Stębowska A. Smoliska U Monitoring of *Trichoderma* species in agricultural soil in response to application of biopreparations. *Biol. Control*. 113, 65–72. <https://doi.org/10.1016/j.biocontrol.2017.07.005>.
53. Peltre C., Nyord T., Bruun S., Jensen L.S., Magida J., 2015. Repeated soil application of organic waste amendments reduces draught force and fuel consumption for soil tillage. *Agric. Ecosyst. Environ.* 211, 94–101. <https://doi.org/10.1016/j.agee.2015.06.004>.
54. Pranagal J., Ligęza S., Smal H., 2020. Impact of Effective Microorganisms (EM) Application on the Physical Condition of Haplic Luvisol. *Agronomy*, 10(7), 1049. <https://doi.org/10.3390/agronomy10071049>.
55. Raudonius S. 2017. AppliScation, of statistics in plant and crop research: important issues. *Žemdirbystė–Agriculture*. 104 (4). 377–382. <https://doi.org/10.13080/z-a.2017.104.048>.
56. Roupheal Y., Giordano M., Cardarelli M., Cozzolino E., Mori M., Kyriacou M., Colla G., Bonini P., 2018. Plant-and seaweed-based extracts increase yield but differentially modulate nutritional quality of greenhouse spinach through biostimulant action. *Agronomy*. 8, 126. <https://doi.org/10.3390/agronomy8070126>.
57. Selby C., Carmichael E., Sharma H.S., 2016. Bio-refining of perennial ryegrass (*Lolium perenne*): Evaluation of aqueous extracts for plant defence elicitor activity using French bean

- cell suspension cultures. *Chem. Biol. Technol. Agric.* 3, 11. <https://doi.org/10.1186/s40538-016-0061-9>.
58. Sen A., Batra A., 2012. Evaluation of antimicrobial activity of different solvent extracts of medicinal plant: *Melia azedarach* L. *Int. J. Curr. Pharm Res.* 4(2), 67–73.
 59. Shihabudeen M.S., Priscilla D.H., Thirumurugan K., 2010. Antimicrobial activity and phytochemical analysis of selected Indian folk medicinal plants. *Int. J. Pharm Sci. Res.* 1(10), 430–434.
 60. Sierra C.A., Trumbore S.E., Davidson E.A., Vicca S., Janssens I., 2015. Sensitivity of decomposition rates of soil organic matter with respect to simultaneous changes in temperature and moisture. *Journal of Advances in Modeling Earth Systems*, 7, 335–356, <https://doi.org/10.1002/2014MS000358>.
 61. Szparaga A., Kocira S., Kocira A., Czerwińska E., Świeca M., Lorencowicz E., Kornas R., Koszel M., Oniszczyk T., 2018. Modification of growth, yield, and the nutraceutical and antioxidative potential of soybean through the use of synthetic biostimulants. *Front. Plant Sci.* 9, p. 1401. <https://doi.org/10.3389/fpls.2018.01401>.
 62. Szparaga A., Kocira S., Kocira A., Czerwińska E., Depo K., Erlichowska B., Deszcz E., 2019. Effect of applying a biostimulant containing seaweed and amino acids on the content of fiber fractions in three soybean cultivars. *Legume Res.* 42, 341–347. <https://doi.org/10.18805/LR-412>.
 63. Tarantino A., Lops F., Disciglio G., Lopriore G., 2018. Effects of Plant Biostimulants on Fruit Set, Growth, Yield and Fruit Quality Attributes of ‘Orange Rubis®’ Apricot (*Prunus armeniaca* L.) Cultivar in Two Consecutive Years. *Sci. Hortic.* 239, 26–34. <https://DOI:10.1016/j.scienta.2018.04.055>.
 64. Tian Z., Lu Y., Ren T., Horton R., Heitman J. H., 2018.. Improved thermo-time domain reflectometry method for continuous in-situ determination of soil bulk density. *Soil and Tillage Research.* 178, 118–129. <https://doi.org/10.1016/j.still.2017.12.021>.
 65. Tilman D., Cassman K.G., Matson P.A., Naylor R., Polasky S., 2002. Agricultural sustainability and intensive production practices. *Nature*, 418, 671–677. <https://doi.org/10.1038/nature01014>.
 66. Tong B., Kool D., Heitman J.L., Sauer T.J., Gao Z., Horton R., 2020. Thermal property values of a central Iowa soil as functions of soil water content and bulk density or of soil air content. *Soil Science.* 71(21), 169–178. <https://doi.org/10.1111/ejss.12856>.
 67. Tooth E., Gelybó G., Dencsó M., Kása I., Birkás M., Horel Á. 2018. Soil CO₂ Emissions in a Long-Term Tillage Treatment Experiment. *Soil Management and Climate Change*, p. 293–307. <https://doi-org/10.1016/B978-0-12-812128-3.00019-7>.
 68. Trevisan S., Manoli A., Quaggiotti S., 2019. A Novel Biostimulant, Belonging to Protein Hydrolysates, Mitigates Abiotic Stress Effects on Maize Seedlings Grown in Hydroponics. *Agronomy.* 9, p. 28. <https://doi.org/10.3390/agronomy9010028>.
 69. Vaitauskienė K., Šarauskis E., Naujokienė V., Liakas, V., 2015. The influence of free-living nitrogen-fixing bacteria on the mechanical characteristics of different plant residues under no-till and strip-till conditions. *Soil and Tillage Research.* 154, 91–102. <https://doi.org/10.1016/j.still.2015.06.007>.
 70. Wu H., Yuan Z., Geng Y., Ren J., Sheng H., Gao L., 2017. Temporal trends and spatial patterns of energy use efficiency and greenhouse gas emissions in crop production of Anhui Province, China. *Energy.* 15, 955–968. <https://doi.org/10.1016/j.energy.2017.05.173>.
 71. Yevtushenko T.V., Tonkha O.L., Pikovskaa O.V., 2016. Changes in bulk density and porosity of chernozem typical under different cultivation systems. *Annals of Agrarian Science.* 14, 299–302. <https://doi.org/10.1016/j.aasci.2016.09.005>.
 72. Zhang B., Li Y., Ren T., Tian Z., Wang G., He X., et al. 2014. Short-term effect of tillage and crop rotation on microbial community structure and enzyme activities of a clay loam soil. *Biol. Fertil. Soils.* 50, 1077–1085. <https://doi.org/10.1007/s00374-014-0929-4>.