

Technical Note: Improved synthetic routes to *cis*- and *trans*-(2-Methyloxirane-2,3-diyl)dimethanol (*cis*- and *trans*- β -isoprene epoxydiol).

Molly Frauenheim,¹ Jason D. Surratt,^{1,2} Zhenfa Zhang, Avram Gold¹

¹Department of Environmental Sciences and Engineering, Gillings School of Global Public Health, The University of North Carolina at Chapel Hill, NC, 27599-7431, USA

²Department of Chemistry, College of Arts and Sciences, The University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599-3290, USA




Figure S1. ¹H NMR (400 MHz, D₂O) mesaconic acid, dimethyl ester.




Figure S2. ^1H NMR (400 MHz, D_2O) *E*-2-methyl but-2-ene-1,4-diol.




Figure S3. ¹H NMR (400 MHz, D₂O) *trans*-(2-methyloxirane-2,3-diyl)dimethanol (*trans*-β-IEPOX).




Figure S4. ^1H NMR (400 MHz, D_2O) citraconic acid dimethyl ester.




Figure S5. ^1H NMR (400 MHz, D_2O) Z-2-methyl but-2-ene-1,4-diol.




Figure S6. ¹H NMR (400 MHz, D₂O) *cis*-(2-methyloxirane-2,3-diyl)dimethanol (*cis*-β-IEPOX).