



1



2



3

4 Figure S1. The J_{NO_2} in $50^\circ \text{ S} - 90^\circ \text{ S}$ from MIPAS and model considering different species at 38 km. (a)
 5 Only the reaction of NO with O_3 and the photolysis of NO_2 are considered. (b) In addition to the reactions
 6 in (a), the reaction of O with NO_2 is considered. (c) In addition to the reactions in (a), the reaction of O with
 7 NO_2 and the reaction of ClO with NO are also considered. Model data is for the same time and location as
 8 the satellite data. The color strip represents the latitude source of data points at the same solar zenith angle.
 9 Each point in J_{NO_2} -Model and J_{NO_2} -MIPAS represents the four-day running mean of the average J_{NO_2} of

10 multiple daily measurements at two latitude degree intervals. In the correlation plots, the abscissa is J_{NO_2} -
11 MIPAS and the ordinate is the J_{NO_2} -Model.

12 In order to better understand which species will have a significant impact on NOx chemistry, J_{NO_2}
13 calculated by considering different species at 38 km is shown in Figure S1. When only the reaction of NO
14 with O_3 and the photolysis of NO_2 are considered, the calculated J_{NO_2} value from satellite data is
15 significantly higher than the model values. After considering the reaction of O with NO_2 , the J_{NO_2} value
16 calculated by satellite data has changed substantially, which indicates that O has a large influence on NOx
17 chemistry. When the reaction of ClO and NO is also considered, the calculated J_{NO_2} value matches well
18 with the model data, which also indicates that ClO should be considered in NOx chemistry at 38 km. It is
19 worth noting that at other altitudes, due to the different profiles of each species, the importance to NOx
20 chemistry is also different. When the altitude is lower than 35 km, the concentrations of ClO and O are very
21 low, and hardly affect NOx chemistry. Moreover, the satellite data error of ClO is very large, so ClO should
22 be ignored in such calculations when the altitude is lower than 35km to avoid introducing unnecessary error.

23 When the altitude is higher than 40km, the concentration of HO_2 increases, so HO_2 can have a large impact
24 on NOx chemistry at higher altitudes than those considered here. HO_2 data can't be measured by MIPAS,
25 so the J_{NO_2} at an altitude higher than 40 km is not included in this work.

26