

Disentangling environmental drivers of Subarctic dinocyst assemblage compositional change during the Holocene

Sabrina Hohmann¹, Michal Kucera¹, Anne de Vernal²

5

¹ MARUM – Center for Marine Environmental Sciences, University of Bremen, Bremen, Leobener Straße 8, D-28359 Bremen, Germany

² Centre Geotop, Université du Québec à Montréal, Montréal, C.P. 8888, Succ. Centre-Ville, Montréal, QC, H3C 3P8, Canada

10 *Correspondence to:* Sabrina Hohmann (hohmannlls@gmail.de)

15 **Abstract.** Analysis of compositional changes in fossil organic-walled dinoflagellate cyst (dinocysts) assemblages is a widely used tool for the quantitative reconstruction of past environmental parameters. This approach assumes that the assemblage composition is significantly related to the reconstructed environmental parameter, which requires an independent correlation between the assemblage and the variable, meaning that the variable explains a dimension of the assemblage variance that is not also explained by other parameters. Theoretically, dinocyst assemblages can be used to reconstruct multiple environmental variables. However, there is evidence that primary and subordinate drivers for assemblage compositions regionally differ and it remains unclear whether a significant relationship to specific parameters in the present ocean always implies that this relationship is significant in fossil assemblages, questioning if past changes in these multiple parameters can be reconstructed simultaneously from fossil assemblages. Here, we analysed a local subset of the Northern Hemisphere dinocyst calibration dataset (n = 1968), including samples from the Baffin Bay area (n = 421), and evaluated the benefits of a local versus a more regional or global calibration for the environmental reconstruction of Baffin Bay oceanography during the Holocene. We determined the dimensionality of the dinocyst ecological response and identified environmental drivers in the Baffin Bay area for the modern dataset. We analysed four existing Holocene records along a north-south transect in the area and evaluated the statistical significance of downcore reconstructions by applying the local and global datasets with different techniques: the modern analogue technique (MAT), the weighted average partial least square (WA-PLS), and maximum likelihood (ML). We evaluated reconstructions tested as significant in the light of the existing state of knowledge about West Greenland's Holocene paleoceanography. Our analyses imply that present-day and Holocene dinocyst assemblages in the Baffin Bay area are primarily driven by salinity changes; other parameters were less important in driving assemblage compositions and their contribution differed among the studied records. We show that the objectively occurring and temporally coherent shifts in dinocyst assemblages in the Holocene of Baffin Bay can be robustly interpreted only by transfer functions that are locally calibrated. Transfer functions based on the broad North Hemisphere calibration yielded many insignificant environmental

20
25
30
35

reconstructions. At the same time, we show that even in the local calibration, not all parameters that appear to significantly affect dinocyst assemblages in the calibration dataset can be meaningfully reconstructed in the fossil record. A thorough evaluation of the calibration dataset and the significance of downcore applications is necessary to reveal the region-specific information contained in dinocyst assemblage composition.

40

1 Introduction

During the last decades, the growing interest in environmental responses to changing climate produced many studies documenting climate variations based on reconstructions making use of marine microfossils (e.g. Muller et al., 1983; Sarnthein et al., 1988; Berger et al., 1989; Meyers, 1997; Rühlemann et al., 1999; Fischer et al., 2000). Next to analyses of chemical and isotopic signals locked in microfossils skeletons, quantitative analyses of the changing taxonomic composition of microfossil assemblages, using ecological models, has been common in palaeoceanography since Imbrie & Kipp (1971). This approach assumes that the composition of marine plankton and benthic communities reflects the environment in which they lived and that this information is stored within the sediment by their fossils. The hypothesis that an assemblage of taxa can be expressed as a function of (a) certain environmental variable(s) (ter Braak, 1987), allows for quantitative modelling applying transfer functions and analogue methods (from now on called transfer functions). Multiple microfossils like planktonic foraminifera (e.g. Kucera et al., 2005), diatoms (e.g. Sha et al., 2014) and dinoflagellate cysts (dinocysts) (e.g. de Vernal et al., 2005) have been used for environmental reconstruction applying multiple transfer function methods.

However, the key assumption of this approach requires that the assemblage composition is significantly related to the given environmental parameter to reconstruct both at present for the purpose of calibration and in the past for the purpose of reconstruction. In the presence of multiple environmental parameters affecting assemblage composition, it is challenging to disambiguate the effect of each parameter and determine which one(s) acted on the analysed fossil assemblages in the past.

There is evidence that the assemblages of zooplankton microfossils, such as planktonic foraminifera and radiolarians being exclusively oceanic, are primarily controlled by temperature (Morey et al., 2005; Hernández-Almeida et al., 2017). Phytoplanktonic microfossils assemblages however, like diatoms and dinocysts, seem to be affected by multiple environmental factors (de Vernal et al., 2000, 2001; Dale et al., 2002; Holzwarth et al., 2007; Radi and de Vernal, 2008; Ribeiro and Amorim, 2008; Lopes et al., 2010; Hohmann et al., 2020). Apparently, phytoplankton communities have a complex relationship to the environment and reflect independently the attribute of hydrography and nutrient availability, rendering the possibility of multiple environmental factor reconstruction. It is tempting to explore whether such dependency on multiple factors can be used to reconstruct changes in all the driving factors in the past.

About 20 % of dinoflagellate species produce organic-walled cysts within their life cycles (Wall and Dale, 1968; Dale, 1983; Taylor and Pollinger, 1987; Head, 1996). As the walls of these cysts consist of a condensed predominantly aromatic macromolecular structure (Bogus et al., 2014), they are relatively well preserved in seafloor sediments (e.g. Zonneveld et al., 2008), not being prone to dissolution after death, unlike mineralised fossils as calcareous (e.g. foraminifera and nannoplankton)

70 or siliceous (e.g. diatoms and radiolarians) microfossils (Koç et al., 1993; Matthiessen et al., 2001; Seidenkrantz et al., 2007; Schröder-Adams and van Rooyen, 2011; Zamelczyk et al., 2012). However, during exposure on the seafloor, oxidation of the dinocyst may result in selective organic matter degradation which might affect the assemblage composition (Zonneveld et al., 2010). Nevertheless, if organic matter degradation is negligible, in settings with carbonate or opal dissolution, dinocyst-based transfer functions are often the best means to reconstruct past surface ocean properties.

75 Dinoflagellates form a major group of planktonic marine primary producers and include diverse autotrophic, mixotrophic, and heterotrophic forms that all thrive in the upper layers of the oceans (e.g. de Vernal and Marret, 2007). Dinocyst studies of marine surface sediment samples revealed that dinocysts are excellent tracers of hydrological and biological conditions (Mudie, 1992; Rochon and Vernal, 1994; Matthiessen, 1995; de Vernal et al., 1997, 2000, 2001, 2013; Radi et al., 2001; Radi and de Vernal, 2004, 2008; Rochon et al., 2008). While dinocyst assemblages can theoretically be applied to quantify multiple
80 environmental drivers, identification of the primary driving factor(s) for the dinocyst assemblages compositional changes is challenging. Relationships between the assemblages and environmental parameters are complex, pointing to regionally different relationships between assemblages and environmental conditions. For example, whereas temperature and salinity gradients are determinants in the Atlantic-Arctic region, primary productivity seems to explain a large portion of taxa variability in the Pacific (Radi and de Vernal, 2008; Hohmann et al., 2020). Studies suggest that the number and type of primary
85 factors driving dinocyst assemblages vary regionally (Zonneveld and Siccha, 2016; Hohmann et al., 2020). Additionally, the evaluation of the performance of transfer functions seems to be method-specific (Hohmann et al., 2020) and biased by spatial autocorrelation (Telford and Birks, 2005, 2009).

Hohmann et al. (2020) analysed the dimensionality of the dinocyst environmental response separately for assemblages from the Pacific and the Atlantic-Arctic regions, calibrated and evaluated multiple transfer-function methods for predicting
90 environmental variables and estimated their performances in the light of spatial autocorrelation. This study identified different primary drivers for two areas affecting dinocyst assemblage composition, highlighting the merit of local calibrations and the necessity to carry out variable selections of main driving factors for each study area separately. Defining a local calibration dataset has the advantage of reducing the complexity of the assemblage to environment composition. However, this comes at the cost of reducing the pool of compositional variability, leading to the risk of biasing the results. For example, in Holocene
95 cores off West Greenland, some of the best compositional analogues have been found in the Gulf of St. Lawrence (e.g. Allan et al., 2018).

However, because factor(s) driving microfossil assemblages differ between regions, it is possible that the same fossil assemblages may find equally good compositional analogues representing different combinations of environmental parameters. This is not to say that a specific taxon responds differently to a given environmental parameter in different regions
100 (the behaviour of a taxon within the environmental gradient is global), but as the importance of environmental parameters differ regionally due to a higher variance of certain parameters, the taxa assemblages react mostly to these variables. This may lead to situations where the assemblages are the same, although some parameters in the environment were different. Hence, while local calibrations limit the range of compositional variability available to derive the transfer function, they are more

likely to disambiguate the main driving variable in a certain region, which may render reconstructions in that region more
105 robust assuming the driving factors remain the same.

By calibrating locally, we can exclude areas where dinocysts are susceptible to lateral transport during sinking and
sedimentation processes. There is evidence that a dinocyst could be displaced up to thousands of kilometres from the location
at the surface where it was produced (Nooteboom et al., 2019). Long-distance cyst transport by bottom waters and sediment
flows was also suggested based on sediment trap studies in the central and North Atlantic Oceans (Dale, 1992; Dale and Dale,
110 1992). This implies that in areas with strong ocean currents, the surface above the sedimentation area does not necessarily
mirror the environmental conditions under which the plankton was produced, generating a nuisance in the calibration dataset.

The aim of this study is to investigate if a local dinocyst-based transfer function calibration on samples covering the Baffin
Bay and surrounding regions allows a robust disambiguation of the factors that affected the assemblage variation in this region
during the Holocene. Baffin Bay is a climate-sensitive region characterised by strong seasonal sea ice variability and meltwater
discharge from the Greenland Ice Sheet (GIS). In view of the current rapid warming in the Arctic, it is relevant to understand
115 how this marine basin reacted to climate changes during the Holocene. Indeed, to this end, many Holocene reconstructions
based on fossil dinocyst assemblages were made and could serve as a basis for a comprehensive analysis of factors affecting
the assemblage composition (e.g. Levac et al., 2001; Ouellet-Bernier et al., 2014; Allan et al., 2018; Caron et al., 2019).

We begin by carrying out an objective variable selection, guided by canonical ordination, and investigate the data structure of
120 the resulting designed local transfer functions. We proceed by testing their ability to interpret assemblage changes recorded in
Holocene sediments by their applications on four records covering a north-south transect throughout the Baffin Bay area. To
this end, we used data from three existing records and added new data from a core located in Southern Melville Bay (see Fig.
1 and Table 1). We specifically focus on Holocene sections, assuming that the magnitude of environmental change in the
Holocene was relatively small for the local calibration to be sufficient to capture the range of past environments recorded in
125 the cores. To estimate the significance of the reconstructions we apply a test designed by Telford and Birks (2011), evaluating
if reconstructions explain more of the variation in the fossil taxa data than random environmental variables assigned to the
same fossil data. To assess the merit of a local versus a larger regional calibration, we compared local transfer function results
to regional transfer functions based on a Northern Hemisphere calibration dataset.

130

135

Table 1: List of cores used in this study. Including core location and water depth (m), the modern sea surface conditions: sea surface temperature (SST) in summer, sea surface salinity (SSS) in summer, months per year of sea ice cover, and annual productivity of organic carbon for each location provided by the WOA and NSIDC.

Core ID	Core location	Latitude (° N)	Longitude (° W)	Water depth (m)	Modern sea surface conditions				Time interval covered (yrs BP)	References for dinocyst records
					SST summer (°C)	SSS summer (psu)	SeaIce (month yr ⁻¹)	PPannual (mgC m ⁻² day ⁻¹)		
HU91-039-008P	Nares Strait	77.16	74.19	663	1.42	31.93	7.64	4635.00	1549-6756	Levac et al. 2001
GeoB19927-3	Southern Melville Bay	73.35	58.05	932	3.89	32.74	7.22	4179.00	0-7677	This paper and Saini et al. 2020
MSM343310	Disco Bugt	68.38	53.49	855	5.45	32.54	3.84	6374.00	135-3578	Allan et al. 2018
HU2008-029-004	Northwest Labrador Sea	61.46	58.03	2674	7.31	34.16	0.72	4497.00	0-9936	Gibb et al. 2015







Figure 1: Map of the study area. (a) Schematic modern surface water circulation in the Baffin Bay area and around Greenland (BIC – Baffin Island Current, WGC – West Greenland Current, EGC – East Greenland Current, IC – Irminger Current); (b) Location of surface sediment samples (red dots) from the calibration dataset and cores (orange dots; see Table 1) used in this study. On the right, modern environmental parameter gradients gridded using DIVA gridding with Ocean Data View (Schlitzer 2018). Units are listed in Table 3. The six-month sea-ice edge in the sea-ice gradient map is indicated by the thick black line.

2 Material and methods

2.1 Data

The calibration dataset includes a local subset of the data from the Northern Hemisphere dataset (n=1968) in de Vernal et al. (2020). The local subset comprises census counts of recent organic-walled dinoflagellate cyst assemblages from core-top sediments at 421 sites from Baffin Bay with Kane Basin, Norwegian Bay, and Lancaster Sound (n=233), Davis Strait (n=103), Labrador Sea (n=30) and Hudson Bay with Hudson Strait and Foxe Basin (n=55) (Fig. 1b). In samples containing low dinocyst concentrations, minimum counts of 60 specimens were accepted. The core-top samples were collected in the uppermost 1 cm of box or gravity cores and represent a few tens of years to centuries, depending upon sedimentation rates and mixing due to bioturbation (Radi and de Vernal, 2008). Palynological sample processing followed the procedures for palynological preparation described in de Vernal et al. (2010), which exclude the use of oxidation techniques to avoid selective degradation of more sensitive cyst taxa such as those produced by Protoperidinales (Marret, 1993). The Northern Hemisphere dataset from (de Vernal et al., 2020) served also as the regional reference dataset which we use to compare the results of the local dataset to. With exception of the samples from Price et al. (2016, 2018), which were added at a later stage, the Northern Hemisphere dataset and therefore all samples included in the local dataset have been tested negative for a significant selective organic matter degradation (Zonneveld et al., 1997) in Hohmann et al. (2020). The taxonomy used corresponds to that presented in Rochon et al. (1999) and updated in van Nieuwenhove et al. (2020). The taxonomic resolution of the local dataset was reduced to 33 dinocyst taxa after grouping following Hohmann et al. (2020) (Table 2, List of taxa). The original dinocyst dataset holding taxon percentages is archived at PANGAEA (de Vernal et al., 2019); the local data matrices with reduced taxonomy used in this analysis are available as supplementary material (S1).

We chose to carry out a canonical-ordination-based parameter selection (Lopes et al., 2010) to characterise the relationship between environmental variables and dinocyst assemblage composition. We defined a set of potential environmental controlling variables for dinocyst distributions in the Baffin Bay region: surface layer (here represented by conditions at 0 m depth) temperature and salinity in summer, sea ice cover and productivity in spring and summer as well as for the whole year. We did not consider temperature, salinity, and productivity parameters for the cold seasons since most of the area is covered by sea ice and there is hardly any productivity and hence no dinocyst production. Apart from environmental factors, some life-cycle factors (e.g., Fensome et al., 1993) such as water depth and distance from shore seem to affect dinocyst assemblages. But as they represent a separate independent dimension of variance (Hohmann et al., 2020), we do not include them in the

185 local calibration dataset. The initial variable selection is listed in Table 3 and includes some of the variables considered in Hohmann et al. (2020) (c.f. for more detail). Their spatial distribution is shown in Fig. 1.

Table 2: List of dinocyst taxa in the local calibration dataset. Including abbreviations and highest relative abundances (minimum is always zero). Asterisks indicate heterotrophic taxa.

Taxa name	Abbreviation	Notes	Maximum %
<i>Ataxiodinium choane</i>	Atax		1.2
<i>Bitectatodinium tepikiense</i>	Btep		3.3
<i>Impagidinium aculeatum</i>	Iacu		2.1
<i>Impagidinium pallidum</i>	Ipal		7.5
<i>Impagidinium paradoxum</i>	Ipar		8.8
<i>Impagidinium patulum</i>	Ipat		1.8
<i>Impagidinium sphaericum</i>	Isph		6.2
<i>Nematosphaeropsis labyrinthus</i>	Nlab		71.6
<i>Operculodinium centrocarpum</i>	Ocen	Group including: <i>Operculodinium centrocarpum</i> , <i>O. centrocarpum</i> — short processes, <i>O. centrocarpum</i> — Arctic morphotype	93.5
<i>Pyxidinosopsis reticulata</i>	Pret		1.5
<i>Spiniferites elongatus</i>	Selo		27.3
<i>Spiniferites ramosus</i>	Sram		14.6
<i>Spiniferites lazus</i>	Slaz		0.3
<i>Spiniferites mirabilis-hyperacanthus</i>	Smir		0.6
<i>Spiniferites</i> spp.	Sspp		4.4
Cyst of <i>Pentapharsodinium dalei</i>	Pdal		95.3
Cyst of <i>Scrippsiella trifida</i>	Stri		4.2
<i>Islandinium minutum</i> *	Imin	Group including: <i>Islandinium minutum</i> *, <i>Islandinium ? cezare</i> *, <i>Islandinium brevispinosum</i> *	96.8
<i>Echinidinium karaense</i> spp.*	Ekar		19.6
<i>Brigantedinium</i> spp.*	Bspp	Group including: <i>Brigantedinium</i> spp., <i>Brigantedinium cariaeoense</i> , <i>Brigantedinium simplex</i> , Cyst of <i>Protoperidinium americanum</i> *	99.0

<i>Dubridinium</i> spp.*	Dubr		3.4
<i>Protoperidinioid</i> cyst*	Peri		1.3
<i>Lejeunecysta</i> spp.*	Lspp		0.7
<i>Selenopemphix nephroides</i> *	Snep		2.2
<i>Xandarodinium xanthum</i> *	Xxan		0.9
<i>Selenopemphix quanta</i> *	Squa	Group including: <i>Selenopemphix quanta</i> *, Cyst of <i>Protoperidinium nudum</i> *	20.8
<i>Trinovantedinium applanatum</i> *	Tapp		5.2
<i>Votadinium calvum</i> *	Vcal		0.5
<i>Votadinium spinosum</i> *	Vspi		0.6
<i>Quinquecuspis concreta</i> *	Qcon		0.7
Cyst of <i>Polykrikos kofoidii</i> *	Pkof		8.6
Cyst of <i>Polykrikos</i> sp. – Arctic*	Parc		23.1
<i>Echinidinium</i> spp.*	Espp		2.5

190

Table 3: Environmental variables used in the canonical-ordination based parameter selection and their ranges within the local calibration dataset.

195

Environmental variable	Minimum	Maximum	Range
SSTsummer [°C]	-0.81	13.24	14.05
SSSsummer [psu]	17.51	34.98	17.47
SeaIce [months yr ⁻¹]	0	12	12
PPAnnual [mgC m ⁻² day ⁻¹]	824	33047	32223
PPSpring [mgC m ⁻² day ⁻¹]	64	3626	3562
PPSummer [mgC m ⁻² day ⁻¹]	202	6021	5819

We used four sediment sequences to test the ability of the locally calibrated transfer functions to interpret assemblage changes during the Holocene. These sediment cores cover a north-south transect throughout the Baffin Bay region (Fig. 1(b), Table 1) including intervals within the last 10 ka.

200

We analysed dinocyst assemblages in core GeoB19927-3 (73°35,26' N, 58°05,66' W), which was taken at 932 meters of water depth by gravity coring during cruise MSM44 in 2015 (Dorschel et al., 2015). This core is from southern Melville Bay, an area where the relatively warm high-salinity West Greenland Current (WGC) interacts with the cold polar water of the Baffin Current (BC) originating from the Arctic Ocean and meltwater from glaciers of the North-West Greenland region. It is located optimally and offers a high resolution and continuous sedimentation (Dorschel et al., 2016). A high-resolution radiocarbon-based chronology and organic biomarker proxies for sea ice and primary productivity have already been presented by Saini et al. (2020). The core consists of 1147 cm of sediment. The working half of the core was sampled and divided into four sets of samples. One set was freeze-dried for dinocyst analysis. The other three sets were stored for other multi-proxy analyses and chronology (i.e., biomarker, foraminifera, and provenance studies). In reference to summer temperature anomalies during the Holocene for the Northern Hemisphere (Marcott et al., 2013; Kaufman et al., 2020), we expected a possible trend towards the last 2000 years and sampled at a higher resolution within the top 270 cm. We sub-sampled every centimetre within the top 25 cm, every 5 cm within the 25 cm to 280 cm interval, and every 10 cm within the 280 cm to 760 cm interval covering the last 7677 years, including 124 samples (see Table 1). Sample processing followed the procedure for palynological preparation described in de Vernal et al. (2010), which consists of repeated HCl and HF treatment. The taxonomy of dinocysts used here was based on Rochon et al. (1999) and de Vernal et al. (2020), considering all taxonomic categories that were resolved in the calibration dataset. To obtain statistically reliable assemblage counts, at least 300 dinocyst specimens were counted per sample when possible. For samples with low dinocyst abundance as many specimens as possible were enumerated. For the sample with the lowest dinocyst abundance, we counted 89 specimens. The raw assemblage dataset for GeoB19927-3 will be available on time via the PANGAEA database.

In addition, we re-evaluated three existing sediment records that have already been studied in detail and have been analysed for their palynological content and dinocyst assemblages. HU91-039-008P (Levac et al., 2001) was sampled every 20 cm from the top to 820 cm which corresponds to 1549-6756 years BP and includes 41 samples. MSM343310 (Allan et al., 2018) includes 194 samples taken around every 8 cm from top to 925 cm (135-3578 years BP), while HU2008-029-004 (Gibb et al., 2015) includes 30 samples from every 4 cm from top to 116,5 cm (0- 9936 years BP) (see Table 1).

225

2.2 Data analysis

To explore the local relationship between dinocyst assemblages and environmental variables, we followed the procedure schematically illustrated in Fig. 2 and used statistical ordination (McGarigal et al., 2000). We performed detrended correspondence analysis (DCA) (Birks, 1995) to evaluate the gradient length, expressed as standard deviation (SD) units in the first DCA axis to decide whether to apply linear ($SD < 3$) or unimodal ordination ($SD > 4$; $SD > 3$ to < 4 both methods possible but unimodal ordination is advised) (Lepš and Šmilauer, 2003). For DCA and further analyses, we used the vegan package (Oksanen et al., 2019) in R (R Core Team, 2017). The dataset was found to have an SD of 3.0 (S2 - supplementary

235 data) using log-transformed ($\log(x+1)$) dinocyst assemblage per mil data. As the SD of our dataset is biased towards the linear threshold, we proceeded with linear techniques.




Figure 2: Flowchart describing the approach and sequence of analyses in this study.

240

In the first step, we selected the main drivers affecting compositional changes in the dinocyst assemblages. We applied Redundancy Analysis (RDA), a constrained ordination (McGarigal et al., 2000) for linear data, to quantify the strength of the relationship between species variances and the tested forcing parameters. We applied Hellinger-transformation (Rao, 1995) to the dinocyst taxa ratios (33 taxa with abundances in percent). This transformation was proposed by Legendre & Gallagher (2001) for assemblage data that contains many double-zero abundances, as it allows using it without considering the common absence of species as a resemblance between assemblages. Hellinger-transformation also gives less weight to species with low counts and many zeros, focussing more on the composition of common species, as we assume that local datasets unlike global

ones do not include indicators or inter-regional endemic species. It also ensures the preservation of Euclidean distance in a linear ordination when unimodal data is used as our species data SD did not indicate strongly linear data. Environmental data were standardised as described above to ensure the compatibility of variables for comparisons.

RDA was performed after ter Braak (1986) using the platform R (R Core Team, 2017). For the variable selection, we followed the procedure described in Hohmann et al. (2020), which includes a variable selection considering Variance Inflation Factors (VIFs) ≤ 2 with the objective to extricate independent variables that show no collinearity between each other and explaining a significant amount of the variance in the species data. Following this procedure, we obtained a set of three independent forcing variables, meaning that each of these three parameters explains a separate dimension of variance in the taxa data (for more explanation see S3 – supplementary data). These three independent variables, that revealed to be summer sea-surface temperature (SSTsummer), summer sea-surface salinity (SSSummer), and spring productivity (PPSpring) (see section 3.2), were used for a final RDA and further analyses.

We determined the extent and type of autocorrelation in the independent data set, as an accurate evaluation of transfer function performances is only possible in case of spatial independence between sample sites in the data set (Telford and Birks, 2009). In the case of spatial autocorrelation, the transfer function performances can be overestimated, which could result in inappropriate choices of transfer function models and parameters, possibly including environmental variables with no ecological relevance as “reconstructable”. Autocorrelation could occur due to spatial interdependence of proxy variables, which would render the main requirement of spatial independence for adequate transfer functions violated. It could also be due to an environmental similarity of nearby sample sites. A spatial autocorrelation due to environmentally similar sites, as found in data sets from the North Pacific as well as the North Atlantic and Arctic (Hohmann et al., 2020), may meet the requirement of transfer functions. For autocorrelation tests and further analyses, we additionally used R packages fields (Nychka et al., 2017), palaeoSig (Telford, 2015), and rioja (Juggins, 2017) implementing the routines developed in Telford and Birks (2009).

For the transfer functions remaining in the independent calibration dataset, we did calibration and cross-validation based on the leave-one-out (LOO) (cf. Efron and Gong, 1983) and h-block technique (Burman et al., 1994; Telford and Birks, 2005, 2009) for three distinctly different transfer function approaches: the Maximum Likelihood (ML), the Weighted Averaging Partial Least Square (WA-PLS) and the Modern Analogue Technique (MAT) (for further details about the applied approaches see Hohmann et al., 2020). For calibration and cross-validation tests we additionally used R packages bioindic (Guiot and Gally, 2014), gstat (Pebesma, 2004), and sp (Pebesma and Bivand, 2005) and followed the R code developed by Trachsel and Telford (2016) for h-block cross-validation and the estimation of h.

We reconstructed the three independent variables (SSTsummer, SSSsummer, and PPSpring) within four sediment sequences collected along a north-south transect through the Baffin Bay area. To determine changes in the environment visible through assemblage shifts downcore, we performed principal component analysis (PCA) on percentages of downcore dinocyst assemblages.

We tested the ability of the calibrated transfer functions to interpret the assemblage changes recorded in the sediments by applying the Telford and Birks method (2011). To estimate the significance of the reconstructions, this method evaluates if reconstructions explain more of the variance in the fossil data than most reconstructions derived from transfer functions trained on random environmental data. It also determines the best reconstruction when reconstructing several environmental variables and if there is sufficient information in the proxy data to support multiple independent reconstructions. We use redundancy analysis (RDA) to estimate the proportion of variance in the fossil data explained by a single reconstruction. Then, using the same taxa data from the calibration dataset, reconstructions are inferred from transfer functions trained on random environmental variables drawn from a uniform distribution. The proportion of the variance explained by these random reconstructions is then estimated. If the tested variable explains more of the taxa variance than 95 % of the random reconstructions, the reconstruction is deemed statistically significant (cf. Telford and Birks, 2011). Reconstructions that fail this test should be interpreted with caution.

Reconstructions and significance tests within the four cores were performed by applying the calibrated local dataset and the regional Northern Hemisphere dataset, aiming to assess the advantages and disadvantages of using the local dataset.

3 Results

3.1 Redundancy analysis

The first step of the variable selection is based on RDA carried out for single variables. It revealed that, when considered individually, each of the six environmental variables appeared to explain a significant ($p \leq .01$) amount of variation in the assemblage composition in the dataset (Table 4). Together, they explain 32 % of the total inertia in the taxa dataset. We then evaluated the independence of the tested environmental variables through a selection, excluding variables with $VIF > 2$. This successive forward selection uncovered three separate dimensions in the taxa variance driven by SSTsummer, SSSsummer and PPSpring respectively. The remaining variables were excluded from further analyses due to their collinearity with SSTsummer, SSSsummer, or PPSpring.

Table 4: Explained inertia (as a single variable) and variance inflation factors of the tested environmental variables in the RDA model. VIF(a) when all variables are considered; VIF(b) after manual forward selection for variables without dependence.

Environmental variable	% inertia explained	VIF (a)	VIF (b)
SSTsummer	12.4	3.9	1.8
SSSsummer	6.9	2.0	1.0
SeaIce	9.7	2.9	

PPAnnual	4.2	394.7	
PPSpring	2.5	72.7	1.8
PPSummer	4.6	137.5	

310 When only the independent environmental variables are used, the constrained part of the RDA explains about 24 % of the taxa variance (Table 5) with environmental variables having significant correlations with RDA-axes (Table 6). The first axis explains 66 % of the constrained variance and is positively correlated with SSTsummer, while axes 2 and 3 explain 27 % and 7 % are positively correlated with SSSsummer and PPSpring, respectively (Fig. 3, Table 5). Regarding species-environment relationships, we find that the variance of *I. minutum* is strongly related to changes in SSTsummer, while the abundance of *N. labyrinthus* is related to SSSsummer. The cyst of *P. dalei* shows correlations with SSTsummer and PPSpring. Phototrophic cysts are generally associated with higher summer temperatures and heterotrophic cysts with lower temperatures (Figs. 3a, b). Sample-environment relationships (Figs. 3c, d) reveal that all samples contribute to the variance in SSTsummer. The variance for PPSpring is high for samples from Labrador Sea, Baffin Bay and Hudson Bay, while SSSsummer variance is highest in samples from Davis Strait and Hudson Bay.

320

325

Table 5: RDA results and results of permutation tests for the significance of RDA-axes after manual forward selection ($p \leq .01$).

		RDA-axis 1	RDA-axis 2	RDA-axis 3	Total inertia	Proportion of total variance explained
Eigenvalues		0.057	0.024	0.005	0.365	
Constrained	proportion	0.660	0.284	0.056		0.236
Constrained	cumulative	0.660	0.944	1.000		
F-value		85.2	36.6	7.3		

330

Table 6: Intraset correlation coefficients (r^2) between independent variables and RDA-axes and between the variables themselves.

	RDA-axis 1	RDA-axis 2	RDA-axis 3	SSTsummer	SSSsummer	PPSpring
SSTsummer	0.79	0.01	0.20	1	0.01	0.45
SSSsummer	0.02	0.96	0.01		1	0.01
PPSpring	0.08	0.01	0.91			1




Figure 3: RDA scores. (a and b): RDA ordination diagrams of dinocyst taxa (see Table 1 for abbreviations) vs environmental variables (arrows) showing the correlation with the axes (direction of arrows) and the strength of the dinocyst vs environmental variable relationship (length of arrows). Species in the centre region are without abbreviations. (c and d): Sample ordination for RDA constrained by independent environmental variables for axes 1, 2 and 3.

340

3.2 Spatial autocorrelation

We tested spatial and environmental autocorrelation, considering only the independent variables and following the procedure described by Telford and Birks (2009). Inevitably, the performance (r^2) deteriorates with increasing fraction of sites deleted (S4 - supplementary data). We find that the deterioration of performance is much worse with neighbourhood deletion than with random deletion, indicating the presence of spatial autocorrelation. When deleting environmentally similar sites, we find a pattern of performance loss that is slightly worse for neighbourhood deletion than for environmental deletion. Hence, removing geographically close sites causes a higher decline in performance than when the same fraction of environmentally similar sites is removed, indicating spatial autocorrelation over environmental autocorrelation. Variable performances drop below 0.55 when samples within a 200 km radius are deleted, implying that calibrated transfer functions need to be interpreted with caution.

350

3.3 Cross-validation

For transfer function development, LOO-cross-validation results indicate the best performances for MAT, with WA-PLS RMSEP values on average around 40 % higher and ML RMSEP values on average by around 100 % higher, depending upon the variable (Table 7). The differences in LOO-RMSEPs between the calibration and the verification subsets for MAT show differences of 25 % for SSSsummer and 8% for PPSpring, indicating an inadequate spatial coverage for some areas within the regional subset for these parameters. For SSTsummer reconstructions the spatial coverage in the dataset seems to be sufficient. All three environmental parameters show similar performances within each reconstruction method. The h-block cross-validation shows MAT RMSEP increasing by 60 %, 60 % and 95 % for SSTsummer, SSSsummer and PPSpring, respectively. RMSEPs of WA-PLS and ML remain largely unaffected by site deletion within the applied radius h for SSTsummer (1 %) and SSSsummer (7 %) (Table 7). For PPSpring RMSEPs increase by about 25 %.

360

An inspection of the distribution of residuals from cross-validation tests of the variables along their environmental gradient reveals uneven residual distributions for all independent variables. (S6 - supplementary material). For all transfer function methods, the spatial structure of the residuals is complex and large residual values emerge regionally. For ML large over- and underestimations suggest that the method is inaccurate for reconstructions. For MAT and WA-PLS, we observe a systematic underestimation in the Labrador Sea, especially for SSTsummer and PPSpring and an overestimation in parts of Baffin Bay and Hudson Bay with especially large overestimations for PPSpring in Hudson Bay. However, residuals for SSSsummer reconstructions regarding MAT and WA-PLS are quite small, suggesting solid transfer functions.

365

375 **Table 7: Cross-validation performances.** Performance for the local dataset applying LOO-cross-validation and h-block cross-validation
 for MAT (c - calibration dataset, v – verification dataset) with the number of analogues used (#ana.), for WA-PLS with the number of
 components used (#comp.) and for ML. The number of analogues and components that will perform best during reconstructions (#ana. and
 #comp.) have been determined calculating the RMSE of prediction as a function of the number of analogues used and a randomisation t-test
 for testing the significance of cross-validated components used after van der Voet (1994), respectively. h-values have been estimated by
 380 fitting a spherical variogram to detrended residuals of a WA model (S5 - supplementary material) following Telford and Birks (2009) as
 well as Trachsel and Telford (2016).

Cross-validation	LOO													
Method	MAT					WA-PLS					ML			
	RMSEP _c	r ² _c	RMSEP _v	r ² _v	#ana.	RMSE	r ²	RMSEP	r ²	#comp	RMSE	r ²	RMSEP	r ²
SSTsummer	1.38	0.76	1.35	0.80	5	1.90	0.58	2.03	0.52	4	2.73	0.41	2.75	0.40
SSSummer	1.35	0.68	1.01	0.78	5	1.55	0.56	1.60	0.53	2	2.34	0.49	2.37	0.47
PPSpring	324.16	0.61	298.14	0.75	7	446.20	0.30	466.17	0.24	3	748.44	0.04	750.99	0.04
Cross-validation	h-block									h [km]	Variogram model			
Method	MAT			WA-PLS			ML							
	RMSEP	r ²	#ana.	RMSEP	r ²	#comp.	RMSEP	r ²						
SSTsummer	2.25	0.42	5	1.93	0.59	4	2.74	0.43	256	Spherical				
SSSummer	2.14	0.22	5	1.68	0.64	4	2.48	0.57	623	Spherical				
PPSpring	636.67	0.01	7	536.96	0.36	5	958.37	0.02	745	Spherical				

3.4 Transfer function significances in sediment core reconstructions


Having established the transfer functions, we applied them to the selected four sediment cores covering the N-S transect. First, for each record, we identified the interval characterised by the largest changes in assemblage composition, which we assumed to be related to the most important environmental change. PC analysis of downcore dinocyst assemblages revealed one to two
390 assemblage changes within each of the four cores (Fig. 4). The first component (PC1) of the PCA identified three assemblage zones in the northernmost record HU91-039-008P, which account for 74.5 % of the total variance, while PC2 accounted for 17.0 %. In Southern Melville Bay (GeoB19927-3) and Disko Bugt (MSM343310) PC1 identified two assemblage shifts accounting for 76.6 % and 77.7 % of the total variance, while PC2 identified at least three shifts accounting for 17.0 % and 18.3 % respectively. Within the most southern record from the Labrador Sea (HU2008-029-004) PC1 revealed four shifts (69.3
395 %) and PC2 three (21.3 %).

The reconstructions based on the local calibration dataset for the northern core HU91-039-008P (Fig. 4a) show modest changes but SSTsummer resonates with the facies shifts. In core GeoB19927-3 (Fig. 4b), assemblage shifts correspond to variations in SSSsummer and PPSpring reconstructions based on ML and WA-PLS. In core MSM343310 (Fig. 4c), the facies shift is caught in SSSsummer and SSTsummer reconstructions based on ML and WA-PLS. In core HU2008-029-004 (Fig. 4d), we identify
400 shifts based on WA-PLS and ML reconstructions.

The different degree of correspondence between the environmental reconstructions inferred by the transfer functions and the principal community change trends in the dinocyst records is echoed in the significance tests made after the approach of Telford and Birks (2011) for downcore reconstructions. Here a significant reconstruction should explain more of the variance in the fossil dinocyst assemblage than a random transfer function calibrated using the same assemblages but random environmental
405 data. These tests reveal that different transfer function methods and environmental variables produce significant reconstructions for each core. Significance test results of transfer function applications with the local dataset and the regional Northern Hemisphere dataset reveal the same pattern (Table 8).

When the Northern Hemisphere dataset is used for reconstructions of the four downcore assemblages (Fig. 4), we find that 17 % of all the tested transfer functions produce significant estimates (SSSsummer – 25 %, SSTsummer – 17 %, PPSpring – 8
410 %) (Fig. 5, Table 8).

However, when applying the local calibration dataset, most transfer functions produce significant reconstructions for SSSsummer (75 %) and SSTsummer (50 %), and some for PPSpring (8 %) (Fig. 5, Table 8). For the northern core HU91-039-008P, summer salinity estimates are significant with all three transfer function methods (Fig. 5a). In the Southern Melville Bay core GeoB19927-3, the WA-PLS transfer function has been tested as significant for PPSpring, as well as SSSsummer and
415 SSTsummer (Fig. 5b). For records in Disko Bugt (MSM343310), both MAT and WA-PLAS produce significant estimates for SSSsummer, while WA-PLS shows additional significance for SSTsummer (Fig. 5c). Further south, the record from Labrador Sea (HU2008-029-004) yields significant estimates for SSSsummer and SSTsummer based on MAT and ML (Fig. 5d).



420 **Figure 4: Downcore reconstructions based on the local dataset (n = 421) and the Northern Hemisphere dataset (n = 1968) using three transfer function techniques.** MAT, WA-PLS and ML are represented by blue, green and purple curves respectively. The lighter blue curves correspond to maximum and minimum possible values calculated from a set of five modern analogues. The ecostratigraphic zones are determined from PC analyses (dashed horizontal lines). The “|” signs (red) on the upper horizontal axis correspond to modern conditions. (a) HU91-039-008P, (b) GeoB19927-3, (c) MSM343310 and (d) HU2008-029-004.

425

Table 8: Significances for downcore reconstructions after the approach of Telford and Birks (2011) using the local dataset and the regional Northern Hemisphere dataset. Significant reconstructions are marked by a green check mark, and the not significant ones by a red cross.

Transfer significances	function	Baffin Bay dataset			Northern Hemisphere dataset		
		SSTsummer	SSSsummer	PPspring	SSTsummer	SSSsummer	PPspring
HU91-039-008P	MAT	√	√	×	×	√	√
	WA-PLS	√	√	×	×	×	×
	ML	×	√	×	×	×	×
GeoB19927-3	MAT	×	×	×	×	×	×
	WA-PLS	√	√	√	×	√	×
	ML	×	√	×	×	×	×
MSM343310	MAT	×	√	×	×	√	×
	WA-PLS	√	√	×	×	×	×
	ML	×	×	×	√	×	×
HU2008-029	MAT	√	√	×	√	×	×
	WA-PLS	×	×	×	×	×	×
	ML	√	√	×	×	×	×



430

significances n = 421

significances n = 1968



(a)

HU91-039-008P





(b)

GeoB19927-3





(c)

MSM343310



(d)

HU2008-029-004



435 **Figure 5: Results of the significance tests for downcore reconstructions.** (a) HU91-039-008P, (b) GeoB19927-3, (c) MSM343310 and
(d) HU2008-029-004. For all significance tests, 999 random environmental variables were generated to produce the null distribution (grey
bars). Black lines mark the variances explained by observed variables. Red dotted lines mark the 95th percentile of the random distribution
($p \leq .05$) above which reconstructions are deemed as significant. The proportion of the variance explained by the first axis of a principal
component analysis (PCA) is also recorded (black dotted line), as this represents the maximum proportion of the variance in the fossil data
that a reconstruction could possibly explain. For significance test analysis we applied the number of analogues (MAT) and components
(WA-PLS) resulting in highest significances respecting all three variables.

440

4 Discussion

4.1 Methodology

The ecological models and transfer functions derived in this study rely on a few assumptions. Foremost is that the composition
of dinocyst assemblages was primarily affected by the selected environmental parameters and that the set of parameters is
445 comprehensive and includes all the relevant parameters. However, it is possible that a sedimentary assemblage reflects other
processes than the considered parameters. For example, non-oceanographic factors such as those related to the dinoflagellate
life cycle (Bravo and Figueroa, 2014) or post-depositional organic matter degradation (Zonneveld et al., 2019) could affect the
dinocyst assemblage composition in the calibration dataset. The production of cysts is a part of the dinoflagellate's sexual
reproduction. It protects the diploid cell, which may be dormant for a variable length of time. After dormancy the cell emerges
450 from the cyst and re-establishes its metabolic activities within the pelagic ecosystem (Taylor and Pollinger, 1987). During
encystment many cysts settle on the seafloor, requiring areas where the seafloor is not too deep to ensure the possibility of
excystment and ascendance of the cells into surface waters. Hohmann et al. (2020) showed that water depth and distance from
shore explain a significant amount of compositional variance in dinocyst assemblages ($p < .01$). However, these variables were
shown to describe a separate dimension of variability, orthogonal to the tested oceanographic and primary productivity factors.
455 Also, when applied to a sedimentary record, factors such as the distance from shore and water depth should be mostly constant
for the downcore record. The effect of organic matter degradation on the dinocyst assemblage composition can be detected
independently of the ecological model because the ratio between cysts of autotrophic and heterotrophic taxa should
systematically decrease with an increase in selective organic matter degradation, caused by their different wall polymer
composition (Zonneveld et al., 1997). In the Northern Hemisphere calibration dataset, there is no evidence of a pervasive effect
460 of degradation. Neither in the Northern Hemisphere (Hohmann et al., 2020), nor in the Baffin Bay local dataset (this study) a
systematic relationship between the residuals and the ratio of heterotrophic and autotrophic taxa has been observed.

Since selective degradation plays at most a subordinate role in biasing dinocyst assemblages in this study and life-cycle factors
explain an independent dimension of the taxa variance, we consider the dinocyst assemblage compositions in the calibration
datasets as being mostly driven by environmental factors. However, most of the taxa variance in the local calibration dataset
465 cannot be explained even when all tested environmental factors (including water depth and distance from shore) are combined.
First, the combination of taxa to achieve higher consistency may hide a finer ecologically relevant taxonomic level. Since

studies revealed the existence of cryptic species in modern dinoflagellates (Montresor et al., 2003; Parkinson et al., 2016; Wang et al., 2019) and since some taxa are grouped voluntarily during analysis, finer relevant taxonomic levels are very likely. It is, however, currently unavoidable because routine analyses (de Vernal et al., 2020) do not allow a refinement in regard to taxonomic levels.

Further, cyst-producing dinoflagellates lose their mobility at cyst formation, from which time onwards they are subject to ocean current transport until they reach the ocean floor. Although we excluded sample sides areas with strong oceanographic currents, small-size dinocyst particles might still be subject to transport by moderate currents when not incorporated in marine snow or faecal pellets (e.g. Turner 2002, 2015) or bottom water currents and sediment flows (Dale, 1992; Dale and Dale, 1992). This may result in a lateral relocation and sedimentation from a cysts formation environment and thus in an additional nuisance in the calibration dataset (Nooteboom et al., 2019). Biases in the reconstructions may also result from advected fossil dinocysts in the past when ocean currents were different. While this might not be eminent for stratigraphies in Baffin Bay during the Holocene, in general cores from such regions need to be treated with caution. Additional noise may also emerge from modern assemblages in the calibration dataset, which represent average fluxes over different time intervals. Assemblages may be to a different degree affected by anthropogenic effects, such as local pollution or global warming. Assemblages from sites with low sediment accumulation may represent averages over millennia and contain a mixture of microfossils deposited during distinct oceanographic events of the late Holocene (e.g. Wigley et al., 1981; Bradley and Jonest, 1993) or be affected by cysts deposited during conditions induced by modern anthropogenic global change.

Alternatively, it is possible that the tested parameters are not the most determinant ones or that the assemblage composition is to a certain degree driven by biotic interactions rather than by environmental factors. It is also possible that environmental factors, not yet considered, play an important role. These could include turbulence, micronutrients, stratification, or any other parameters that might be difficult to quantify for data treatment purposes.

Regardless of the limitations that can be due to missing or unidentified driving parameters, we find three mutually independent ecological gradients driving dinocyst assemblages in the local calibration dataset. This confirms previous findings that phytoplankton (Lopes et al., 2010) and hence dinocyst assemblages reflect multiple environmental drivers (e.g. Radi and de Vernal, 2008; Ribeiro and Amorim, 2008; Hohmann et al., 2020).

The existence of multiple environmental gradients affecting the dinocyst assemblage composition implies that it should be possible to extract information from the same fossil dinocyst assemblage on past variability for more than one environmental driver. Whilst this is correct theoretically, one must consider the (small) amount of variance constrained by the tested parameters against the (large) portion of the variance that remains unconstrained. For example, primary productivity, which represents the third gradient after SST and SSS, explains a maximum of 6 % of the constraint variability in the RDA (Table 5 and 6). This parameter does not contribute to the first two dimensions of the RDA (see also Radi and de Vernal, 2008; de Vernal et al., 2020) and although the extracted relationship appears statistically significant, it may well be overwhelmed by factors that are responsible for the unconstrained portion of the variance in assemblage composition.

500 Additionally, we note that the presence of multiple environmental gradients in the local calibration dataset is associated with spatial structuring of the environment (S6 – supplementary material). This may result in the fact that some environmental drivers deemed significant in the RDA, are only significant because they drive the taxa composition in a specific area with a strong gradient of the given factor. This is in line with the observed spatial autocorrelation of the tested variables (S4 – supplementary material) and explains the performance decrease of MAT in the h-block cross-validation (Table 7) and as well
505 as the strong spatial structure observed in the residuals of all methods (S6 – supplementary material). These observations are coherent with findings in Hohmann et al. (2020), where the observed multiple and mutually independent environmental factors driving the dinocyst assemblage composition in two larger regional datasets may be resulting from the agglomeration of information from regions with unlike environmental (primary) drivers and endemic taxon-environment relationships. Hence, our results suggest localised driving mechanisms for dinocyst assemblage compositions in very specific areas. This implies
510 that generating locally calibrated transfer functions will come at the cost of loss of generalisation and limited applicability to past oceanographic states.

Being based upon differing basic principles, the applied transfer function approaches seem to be affected differently by the detected local dataset's spatial structuring and the region-dependent primary drivers (Kucera et al., 2005). ML transfer functions appear to be unable to disclose and depict the taxon-environment relationship when multiple gradients affect the taxa
515 assemblages (highest RMSEPs, lowest r^2 , fewest significant transfer functions). However, ML and WA-PLS performances are less affected by the effect of spatial autocorrelation. During h-block cross-validation, their performances remain similar, while the performances of MAT transfer functions plummet, returning higher RMSEP and lower r^2 than ML and WA-PLS (Table 7). On the other hand, performances of MAT based on LOO cross-validation are more optimistic than for ML and WA-PLS. While MAT finds the local structure in the assemblage data, ML and WA-PLS, which are based on a unimodal species-
520 environment response model, find the general component of the variation in the assemblage data that is correlated with a specific environmental forcing. While all approaches have their advantages, ML and WA-PLS are more robust with regard to the detected spatial structure in the data than MAT (Telford and Birks, 2005). Cross-validation of the local dataset applying a calibration and verification dataset results in a 25 % lower RMSEP_v for the verification dataset compared to the calibration dataset for SSSsummer, while RMSEP_v are < 10 % lower for SSTsummer (2 %) and PPspring (8 %). This indicates that
525 additional sample sites might benefit performances for SSSsummer transfer functions as with an adequate, hence representative, spatial coverage RMSEP should be similar for both datasets (Table 7).

Regardless of the statistical approach, a theoretical high-performance transfer function for one of the independent driving variables for assemblage composition in the local calibration dataset does not necessarily produce a reliable reconstruction applied to fossil downcore dinocyst assemblages. This is because we can never be sure that the main drivers for past
530 assemblages were identical in the modern calibration dataset. To assess if the main past drivers resemble the main modern drivers, we compared the transfer function reconstructions to the major compositional trends in fossil dinocyst assemblages as revealed by multivariate analyses.

Across the four records throughout Baffin Bay, the main pattern of dinocyst assemblage change during the Holocene can be identified as two biofacies shifts occurring around 4.0 ka and 6.7 ka (Fig. 4 and 6) throughout the whole study area and one regime shift occurring in Baffin Bay and Disco Bay between 1.0 and 1.5 ka. Despite the presence of this clear pattern, most transfer functions based on the local and Northern Hemisphere data sets reconstruct modest changes close to uncertainty for SSTsummer, SSSsummer and PPSpring (Fig. 4), likely reflecting the difficulty in attributing the taxa assemblage change to specific factors. Alternatively, one may argue that we have not retained the correct determinant parameters during the manual forward selection. For example, the Holocene records could reflect changes that were due to large shifts in sea ice coverage. Indeed, during the selection process, the analysis suggested that SeaIce does not explain a separate dimension of variance in the taxa assemblages within the local Baffin Bay dataset (S3 – supplementary material), as it revealed high collinearity with SSTsummer. This means that regionally SeaIce and SSTsummer explain a similar taxa variance and while reconstructing one, we reconstruct a mixture of both parameters. It is also possible that the parameterisation of sea ice cover used here is not ideal to describe the relationship with dinocyst assemblages. In the context of Baffin Bay, SSSsummer, which is the parameter that appears to be best reconstructed, is likely a reflection of the regional hydrography closely related to the sea ice cover that matters for productivity. Therefore, we can only conclude that the environmental regime shifts around 1.3 ka, 4.0 ka and 6.7 ka appear to be driven by changes in local hydrography that were associated with salinity changes but considering the magnitude of the salinity changes implied by the transfer functions, it is unlikely that the dinocyst composition responded to salinity directly.

550




Figure 6: Dinocyst assemblage changes according to the first two PC-axes. Coloured dots represent PC transitions pointing to regime shifts around 1.3 ka (between 1.0 ka and 1.5 ka), 4.0 ka (between 3.1 ka and 4.3 ka) and 6.7 ka (orange shaded).

555

In contrast to the results of the local calibration, only a few of the reconstructions were identified as significant when applying the regional Northern Hemisphere dataset (SSTsummer = 16 %, SSSsummer = 25 %, PPspring = 8 %) (Table 8, Fig. 5). This suggests that transfer functions based on broad regional or even global calibration datasets are affected by nuisance variables, which makes it difficult to identify the driving mechanisms of dinocyst assemblages when the amplitude of environmental change is relatively small, as it is the case during the middle and late Holocene. For longer time intervals with large amplitude changes, global calibrations may work better and would be even necessary as larger datasets contain a wider range of analogues. Hence, we suggest a thorough evaluation of the calibration dataset and the significance of fossil applications ensuring that the driving mechanisms of dinocyst assemblages are caught through time and space.

565

4.2 Salinity reconstructions and regime shifts in the context of the existing regional paleoceanographic records

Results imply that in Baffin Bay most locally calibrated MAT and WA-PLS transfer functions produce robust Holocene reconstructions for summer salinity, being the main driver of dinocyst assemblage compositions while representing the regional hydrography closely related to sea ice cover and sea surface temperature driving the salinity changes. This poses the opportunity to interpret SSSsummer reconstructions within the context of the regional paleoceanography.

570

Until 7.6 ka Baffin Bay was still a region of extended sea ice and low primary productivity (e.g. Ouellet-Bernier et al., 2014; Jennings et al., 2014; Gibb et al., 2015; Caron et al., 2019; Limoges et al., 2020; Saini et al., 2020) (Fig. 7 d-g), while the Laurentide and the Greenland ice sheets continuously retreated (Fig. 7 a and b) (Dyke et al., 2003; Larsen et al., 2015).

With the beginning of the Holocene Climate Optimum, sea ice cover in Baffin Bay retreated and it became a region of marginal ice zones until 6.3 ka. During this time, the first observed regime shift occurred between 7.6 ka and 6.3 ka (Fig.6) which is in accordance to the shift towards mild postglacial conditions in Baffin Bay proposed by Gibb et al. (2015) and Caron et al. (2019). In the northwest Labrador Sea (core HU2008-029-004) however, this postglacial transition already occurred around 11.9 ka (Gibb et al., 2015) implying that the observed regime shift also visible in both PCA-axes of HU2008-029-004, cannot - like in Baffin Bay - be ascribed to the postglacial transition.

575

After the transition phase from 6.3 ka on, salinity reconstructions propose a continuous increase in the northwest Labrador Sea and a continuous decrease of salinity in southern Melville Bay (Fig. 7 i and k). For Labrador Sea this is in accordance to findings in Gibb et al. (2015), who ascribes the increase to a reduction of stratification in the upper waters caused by the reduction of meltwater input from the Laurentide ice sheet and the increased advection of North Atlantic water (de Vernal et al., 2001; Solignac et al., 2004). Hence, the observed regime shift in the northwest Labrador Sea between 7.6 ka and 6.3 ka is in line with the suggested regional shift towards modern water circulation south of Davis Strait (Gibb et al., 2015).

585

For southern Melville Bay, other records also propose a slow salinity decrease between 5.5 ka and 1.4 ka (Gibb et al., 2015; Caron et al., 2019), which might be attributed to the still active process of postglacial meltwater discharge north of Davis Strait. This can be supported by a continuous decrease in sea ice cover in this region (Fig. 7 d-f; Gibb et al., 2015, Caron et al., 2019; Saini et al., 2020) and the increase in primary productivity after the first observed regime shift (e.g. Caron et al., 2019; Limoges et al., 2020). At the Southwest Greenland margin dinoflagellate blooms in summer-fall have been associated with freshwater inputs due to meltwater runoff from the GIS, which triggers upwelling of nutrient-rich waters and high productivity (Juul-Pedersen et al., 2015; Krawczyk et al., 2015, 2018). Along most of the West Greenland margin, sea-ice melt and meltwater runoff from the GIS result in seasonal stratification of surface waters, which is amplified in summer by solar heat leading to relatively mild conditions (Juul-Pedersen et al., 2015; Tremblay et al., 2015) also enhancing primary productivity. Pollen data indicate a high influx of long-distance pollen over Northwest Greenland during this time, indicating a milder climate (Funder, 1978; Fredskild, 1985; Fredskild and Andrews, 1985).

In northernmost Baffin Bay, our reconstructions show no general trend but a continuously oscillating salinity (Fig. 7 h), which is supported by reconstructions in Levac et al. (2001). Like in Levac et al. (2001), our reconstructed values stay below 32, indicating lower values than today's modern instrumental values.

However, while a general trend towards lower salinity can be observed north of Davis Strait and south of the northernmost Baffin Bay after 5.5 ka, other studies (Gibb et al., 2015; Caron et al., 2019) also find a salinity peak around 3.7 ka, interrupting the general trend, which is not visible in our reconstruction, although, this salinity peak corresponds to the second regime shift indicated by the PCA of all four studied cores (Fig. 6). This change in dinocyst assemblage compositions is also mentioned in Caron et al. (2019) for southern Melville Bay, although other dinocyst records from this region did not observe a shift (Ouellet-Bernier et al., 2014; Gibb et al., 2015). In northernmost Baffin Bay, our reconstructions point to a period of slightly increased salinity during the regime shift (Fig. 7 h). For the Baffin Bay area this shift may mark the final adaptation of assemblage compositions to full interglacial conditions, supported by still continuously decreasing sea ice cover (Fig. 7) and other paleoclimate studies along the southwest Greenland margin suggesting that optimum conditions were attained between 6.0 and 3.5 ka (Moros et al., 2006; Lloyd et al., 2007; Seidenkrantz et al., 2007; Ren et al., 2009; Andresen et al., 2010; Perner et al., 2013).


In northernmost Baffin Bay and northwest Labrador Sea however, the changes in assemblage compositions are not stable, changing again within the supposed period of the regime shift (Fig. 6). For northern Baffin Bay Levac et al. (2001) find a general deterioration of sea surface conditions after 3.6 ka, which was accompanied by extensive sea ice cover. These findings are supported by the second regime shift, implying changes in the dinocyst assemblage composition.

The late Holocene is locally characterized by the Neoglacial cooling trend (Weidick et al., 2012), which along the West Greenland coast, is associated with subsurface cooling and glacier advances (e.g. Perner et al. 2013; Briner et al., 2016; Schweinsberg et al., 2017) over the last 4.0 ka (e.g. Knudsen et al., 2008; Seidenkrantz et al., 2008; Krawczyk et al., 2010; Andresen et al., 2010). This cooling trend is marked by climate oscillations, including the short warming phases of the Roman Warm Period (2250-1600 years BP) (Lamb, 2002) and the Medieval Climatic Optimum (1100-700 years BP) (Lamb, 1965;

620 Mann, 2002). In Baffin Bay records, the trend's impact becomes visible from around 2.2 ka (Fig. 7). Regional cooling
accompanied by decreasing summer solar insolation (Berger and Loutre, 1991), fostering regrowth of the GIS (Briner et al.,
2010; Weidick and Bennike, 2007; Young et al., 2011), as well as cooling over the GIS (Dahl-Jensen et al., 1998) led to more
extensive sea ice, which was accompanied by increasing sea-surface salinity and a decline in primary productivity. Around
625 1.3 ka the southern Melville Bay core (GeoB19927-3) and the Disco Bay core (MSM343330) suggest a regime shift supporting
the assumption of a transition towards colder climate. This is also in accordance with proxy records from eastern Baffin Bay
suggesting glacier advances (Ouellet-Bernier et al., 2014; Gajewski, 2015) and decreased terrigenous detrital supply (Caron et
al., 2020) in this time interval.

Overall, the reconstructions for summer salinity and the observed regime shifts fit into and support the current picture of West
Greenland's paleoceanography. Comparing salinity reconstructions for southern Melville Bay (GeoB19927-3) with
630 reconstructions of sea ice cover from the same core derived from the biomarker proxy $P_{IP_{25}}$ (Belt and Müller, 2013) for late-
spring sea ice cover (Saini et al., 2020), similar changes can be observed. Increasing sea ice cover correlates with increasing
salinity and vice versa. This additionally supports the significance of SSSsummer WA-PLS reconstructions in the Melville
Bay core and supports the assumption that salinity and sea ice cover are tightly coupled environmental drivers in this area.

635



640 **Figure 7: Time series for significant SSSsummer reconstructions (Table 8) (local MAT and WA-PLS) from the four studied cores compared to different environmental records from the Baffin Bay region.** (a) Greenland Ice Sheet area (Larsen et al., 2015), (b) Laurentide Ice Sheet area (Dyke et al., 2003) (plots after Briner et al. (2016)), (c) Holocene Greenland temperatures based on $\delta^{18}\text{O}$ from the

GISP ice core (Alley et al., 2010), (d) sea ice cover in Southern Melville Bay based on biomarkers (Saini et al., 2020) (MIZ – marginal ice zone), (e) sea ice cover in Upernavik based on dinocyst assemblages (3-point running mean) (Caron et al., 2019), (f) sea ice cover in Southern Baffin Bay based on dinocyst assemblages (5-point running mean) (Gibb et al., 2015; Jennings et al., 2014), (g) primary productivity in Upernavik based on dinocyst assemblages (3-point running mean) (Caron et al., 2019), (h) SSSsummer from core HU91-039-008P, (i) SSSsummer from core GeoB19927-3, (j) SSSsummer from core MSM343310 and (k) SSSsummer from core HU2008-029-004. Regime shifts (Figure 6) are shaded in grey.

650 **5 Conclusion**

Understanding past natural variability of ecosystem drivers in Arctic regions requires reliable reconstructions. We calibrated local transfer functions from Baffin Bay samples for three independent parameters driving dinocyst assemblages in this area. We estimated how well these local functions were able to estimate past environmental parameters in the Baffin Bay region during the Holocene, comparing them to estimates derived from regional transfer functions from Northern Hemisphere samples and afterwards comparing the significant local transfer functions with the existing regional paleoceanographic records. Statistical analyses of surface sediment samples and transfer function application on downcore samples from a Holocene transect along the Baffin Bay indicate that salinity is the main driver of present-day and Holocene dinocyst assemblage composition in the region, with temperature being an important but minor driver.

Analyses of four Holocene records in a north-south transect along the Baffin Bay show three regime shifts that occurred throughout the region. The main environmental parameter to which the dinoflagellate communities seem to have reacted during regime shifts is salinity, but further minor drivers may have differed regionally. The reconstructed salinity records and regime shifts support the current state of knowledge of West Greenland's Holocene paleoceanography.

We show that the relationship between dinocyst assemblage composition and multiple environmental drivers can be modelled by MAT and WA-PLS transfer functions. In our study, the identification of main driving variables and regime shifts was achieved through the application of a local calibration dataset, while through the application of the Northern Hemisphere dataset, we did not succeed in extracting this information from the taxa variability. While our results reinforce the assumption of counteracting nuisance in spatially large calibration datasets, suggesting the application of local datasets, regions with spatially or temporarily broader environmental gradients might supposedly benefit from larger calibration datasets regardless of a certain nuisance.

We suggest a thorough evaluation of transfer function performances and significances for downcore applications to disclose the specific drivers for present and fossil dinocyst assemblages in a studied core location. To this end, a local calibration dataset, which reduces several nuisances, is presumably beneficial when the amplitude of parameter change is relatively small, like during the middle and late Holocene.

Data availability

675 This study contains supplementary material. This material includes the matrices of taxa abundances with all tested
environmental (oceanographic and non-oceanographic) variables from the local $n = 421$ dataset (S1), DCA analysis plot (S2),
script of manual forward selection (S3), spatial autocorrelation analysis plots (S4), variogram plots of the tested variables for
the estimation of h in the h -block cross-validation (S5) and residual plots from cross-validation tests (S6). The R codes applied
for analyses and the script for manual forward selection (S3) can be accessed on ZENODO
680 (<https://zenodo.org/record/8304438>). Dinocyst assemblage data from the newly analysed core GeoB19927-3 are uploaded to
PANGAEA and will be available soon.

Author contribution

All three (co-) authors collectively contributed to the conceptualisation of this study. Michal Kucera and Anne de Vernal
685 acquired funding and provided resources. Sabrina Hohmann developed the methodology with the supervision of Michal
Kucera, compiled the necessary data and performed all formal analyses. Visualisation of research results and original draft
preparation was done by Sabrina Hohmann. Michal Kucera and Anne de Vernal reviewed and edited the original draft.

Competing interests

The authors declare that they have no conflict of interest.

690

Acknowledgements

This study is a contribution of the International Research Training Group “Processes and Impacts of Climate Change in the
North Atlantic Ocean and the Canadian Arctic” (ArcTrain), which was supported jointly by the German Research Foundation
(DFG) (IRTG 1904) and by the Natural Sciences and Engineering Research Council of Canada (NSERC). We also
695 acknowledge the support from the Fonds de recherche du Québec – Nature et Technologie (FRQNT). We are grateful to
Taoufik Radi and Sebastien Zaragosi for their help in the preparation of the reference oceanographical data set and we
especially thank Maryse Henry for her extraordinary support during sample analyses.

New sample material has been provided by the GeoB Core Repository at the MARUM – Center for Marine Environmental
Sciences, University of Bremen, Germany. The data reported in this paper are archived in Pangaea (www.pangaea.de).

- Allan, E., de Vernal, A., Knudsen, M.F., Hillaire-Marcel, C.: Late Holocene sea-surface instabilities in the Disko Bugt area, west Greenland, in phase with $\delta^{18}\text{O}$ -oscillations at Camp Century, *Paleogeography and Paleoclimatology*, 33, 227-224, <https://doi.org/10.1002/2017PA003289>, 2018.
- Alley, R.B., Andrews, J.T., Brigham-Grette, J., Clarke, G.K.C., Cuffey, K.M., Fitzpatrick, J.J., Funder, S., Marshall, S.J.,
705 Miller, G.H., Mitrovica, J.X., Muhs, D.R., Otto-Bliesner, B.L., Polyak, L., White, J.W.C.: History of the Greenland Ice Sheet: paleoclimatic insights, *Quat. Sci. Rev.*, 29, 1728-1756, <https://doi.org/10.1016/j.quascirev.2010.02.007>, 2010.
- Andresen, C.S., Mccarthy, D.J., Dylmer, C.V., Seidenkrantz, M., Kuijpers, A., Lloyd, J.M.: Interaction between subsurface ocean waters and calving of the Jakobshavn Isbræ during the late Holocene, *The Holocene*, 21, 211-224, <https://doi.org/10.1177/0959683610378877>, 2010.
- 710 Belt, S.T., Müller, J.: The Arctic sea ice biomarker IP25: a review of current understanding, recommendations for future research and applications in palaeo sea ice reconstructions, *Quat. Sci. Rev.*, 79, 9–25, <https://doi.org/10.1016/j.quascirev.2012.12.001>, 2013.
- Berger, A., Loutre, M.F.: Insolation values for the climate of the last 10 million years, *Quat. Sci. Rev.*, 10, 297–317, 1991.
- Berger, W.H., Smetacek, V., Wefer, G.: Ocean productivity and paleoproductivity - an overview, in: *Productivity of the Ocean: Present and Past*, John Wiley & Sons Limited, 1–34, 1989.
- 715 Birks, J.B.: Quantitative palaeoenvironmental reconstructions, in: *Statistical Modelling of Quaternary Science Data*, Maddy, D., Brew, I.S. (Eds.), Quaternary Research Association, Cambridge, 271, 1995.
- Bogus, K., Mertens, K.N., Lauwaert, J., Harding, I.C., Vrielinck, H., Zonneveld, K.A.F., Versteegh, G.J.M., 2014. Differences in the chemical composition of organic-walled dinoflagellate resting cysts from phototrophic and heterotrophic
720 dinoflagellates. *J. Phycol.* 20, 254–266. <https://doi.org/10.1111/jpy.12170>
- Bradley, R.S., Jonest, P.D.: ' Little Ice Age ' summer temperature variations: their nature and relevance to recent global warming trends, *Holocene*, 3, 367–376, <https://doi.org/10.1177/095968369300300>, 1993.
- Bravo, I., Figueroa, R.: Towards an ecological understanding of dinoflagellate cyst functions, *Microorganisms*, 2, 11–32, <https://doi.org/10.3390/microorganisms2010011>, 2014.
- 725 Briner, J.P., Stewart, H.A.M., Young, N.E., Philipps, W., Losee, S.: Using proglacial-threshold lakes to constrain fluctuations of the Jakobshavn Isbræ ice margin, western Greenland, during the Holocene, *Quat. Sci. Rev.*, 29, 3861–3874, <https://doi.org/10.1016/j.quascirev.2010.09.005>, 2010.
- Briner, J.P., McKay, N.P., Axford, Y., Bennike, O., Bradley, R.S., de Vernal, A., Fisher, D., Francus, P., Fréchette, B., Gajewski, K., Jennings, A., Kaufman, D.S., Miller, G., Rouston, C., Wagner, B.: Holocene climate change in Arctic
730 Canada and Greenland, *Quat. Sci. Rev.*, 147, 340–364, <https://doi.org/10.1016/j.quascirev.2016.02.010>, 2016
- Burman, P., Bhow, E., Nonal, D.: A cross-validatory method for dependent data, *Biometrika*, 81, 351–358, <https://doi.org/10.1093/biomet/81.2.351>, 1994.

- Caron, M., Rochon, A., Carlos, J., Serrano, M., Onge, G.S.T.: Evolution of sea-surface conditions on the northwestern Greenland margin during the Holocene, *J. Quat. Sci.*, 34, 569–580, <https://doi.org/10.1002/jqs.3146>, 2019.
- 735 Caron, M., Montero-Serrano, J.C., St-Onge, G., Rochon, A.: Quantifying Provenance and Transport Pathways of Holocene Sediments From the Northwestern Greenland Margin, *Paleoceanogr. Paleoclimatology*, 35, 1–23, <https://doi.org/10.1029/2019PA003809>, 2020.
- Dahl-Jensen, D., Mosegaard, K., Gundestrup, N., Clow, G.D., Johnsen, S.J., Hansen, A.W., Balling, N.: Past Temperatures Directly from the Greenland Ice Sheet, *Science*, 282, 268–271, <https://doi.org/10.1126/science.282.5387.268>, 1998.
- 740 Dale, B.: *Dinoflagellate resting cysts: “benthic plankton,” survival strategies of the algae*, Cambridge University Press, 1983.
- Dale, B.: Dinoflagellate contributions to the open ocean sediment flux, in: *Dinoflagellate Contributions to the Deep Sea*, Dale, B., Dale, A.L. (Eds.), *Ocean Biocoenosis Ser.*, 1–31, 1992.
- Dale, B., Dale, A.L.: Dinoflagellate contributions to the sediment flux of the Nordic Seas, in: *Dinoflagellate Contributions to the Deep Sea*, Dale, B., Dale, A.L. (Eds.), *Ocean Biocoenosis Ser.*, 45–75, 1992.
- 745 Dale, B., Dale, A.L., Jansen, J.H.F.: Dinoflagellate cysts as environmental indicators in surface sediments from the Congo deep-sea fan and adjacent regions, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 185, 309–338, [https://doi.org/10.1016/S0031-0182\(02\)00380-2](https://doi.org/10.1016/S0031-0182(02)00380-2), 2002.
- de Vernal, A., Marret, F.: Organic-walled dinoflagellate cysts: Tracers of sea-surface conditions, in: *Developments in Marine Geology*, Elsevier B.V., 371–408, [https://doi.org/10.1016/S1572-5480\(07\)01014-7](https://doi.org/10.1016/S1572-5480(07)01014-7), 2007.
- 750 de Vernal, A., Rochon, A., Turon, J.-L., Matthiessen, J.: Organic-walled dinoflagellate cysts: Palynological tracers of sea-surface conditions in middle to high latitude marine environments, *Geobios.*, 30, 905–920, [https://doi.org/10.1016/S0016-6995\(97\)80215-X](https://doi.org/10.1016/S0016-6995(97)80215-X), 1997.
- de Vernal, A., Hillaire-Marcel, C., Turon, J.-L., Matthiessen, J.: Reconstruction of sea-surface temperature, salinity, and sea-ice cover in the northern North Atlantic during the last glacial maximum based on dinocyst assemblages, *Can. J. Earth Sci.*, 37, 725–750, 2000.
- 755 de Vernal, A., Henry, M., Matthiessen, J., Mudie, P.J., Rochon, A., Boessenkool, K., Eynaud, F., Grøsfjeld, K., Guiot, J., Hamel, D., Harland, R., Head, M.J., Kunz-Pirrung, M., Levac, E., Loucheur, V., Peyron, O., Pospelova, V., Radi, T., Turon, J.-L., Voronina, E.: Dinoflagellate cyst assemblages as tracers of sea-surface conditions in the northern North Atlantic, Arctic and sub-Arctic seas: the new “n= 677” data base and its application for quantitative palaeoceanographic reconstruction, *J. Quat. Sci.*, 16, 681–698, <https://doi.org/10.1002/jqs.659>, 2001.
- 760 de Vernal, A., Eynaud, F., Henry, M., Hillaire-Marcel, C., Londeix, L., Mangin, S., Matthiessen, J., Marret, F., Radi, T., Rochon, A., Solignac, S., Turon, J.L.: Reconstruction of sea-surface conditions at middle to high latitudes of the Northern Hemisphere during the Last Glacial Maximum (LGM) based on dinoflagellate cyst assemblages, *Quat. Sci. Rev.*, 24, 897–924, <https://doi.org/10.1016/j.quascirev.2004.06.014>, 2005.
- 765 de Vernal, A., Henry, M., Bilodeau, G.: *Micropaleontological preparation techniques and analyses*, Cahiers du Geotop, 3, 2010.

- de Vernal, A., Rochon, A., Fréchette, B., Henry, M., Radi, T., Solignac, S.: Reconstructing past sea ice cover of the Northern Hemisphere from dinocyst assemblages: Status of the approach, *Quat. Sci. Rev.*, 79, 122–134, <https://doi.org/10.1016/j.quascirev.2013.06.022>, 2013.
- 770 de Vernal, A., Radi, T., Zaragosi, S., Van Nieuwenhove, N., Rochon, A., Allan, E., De Schepper, S., Eynaud, F., Head, M.J., Limoges, A., Londeix, L., Marret, F., Matthiessen, J., Penaud, A., Pospelova, V., Price, A., Richerol, T.: Percentages of common modern dinoflagellate cyst taxa in surface sediments of the Northern Hemisphere and corresponding environmental parameters, *PANGAEA*, <https://doi.org/10.1594/PANGAEA.908494>, 2019.
- 775 de Vernal, A., Radi, T., Zaragosi, S., Van Nieuwenhove, N., Rochon, A., Allan, E., De Schepper, S., Eynaud, F., Head, M.J., Limoges, A., Londeix, L., Marret, F., Matthiessen, J., Penaud, A., Pospelova, V., Price, A., Richerol, T.: Distribution of common modern dinoflagellate cyst taxa in surface sediments of the Northern Hemisphere in relation to environmental parameters: The new n=1968 database, *Mar. Micropaleontol.*, 159, 101796, <https://doi.org/10.1016/j.marmicro.2019.101796>, 2020.
- 780 Dorschel, B., Afanasyeva, V., Bender, M., Dreutter, S., Eisermann, H., Gebhardt, A.C., Hansen, K., Hebbeln, D., Jackson, R.: Past Greenland ice sheet dynamics, palaeoceanography and plankton ecology in the Northeast Baffin Bay - Cruise No. MSM44 “BAFFEAST” - June 30-July 30, 2015 - Nuuk (Greenland), *MARIA S. MERIAN-Berichte*, 51, https://doi.org/10.2312/cr_msm2344, 2015.
- Dyke, A.S., Moore, A., Robertson, L.: Deglaciation of North America, *Geol. Surv. Can, Open File*, 2003.
- 785 Efron, B., Gong, G.: A leisurely look at the bootstrap, the jackknife, and cross-validation, *Am. Stat.*, 37, 36–48, <https://doi.org/10.1080/00031305.1983.10483087>, 1983.
- Fensome, R.A., Taylor, F.J.R., Norris, G., Sarjeant, W.A.S., Wharton, D.I., Williams, G.L.: A classification of living and fossil dinoflagellates, *Micropaleontol. Spec. Pap.*, 7, 1–351, 1993.
- Fischer, G., Ratmeyer, V., Wefer, G.: Organic carbon fluxes in the Atlantic and the Southern Ocean: Relationship to primary production compiled from satellite radiometer data, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 47, 1961–1997, 790 [https://doi.org/10.1016/S0967-0645\(00\)00013-8](https://doi.org/10.1016/S0967-0645(00)00013-8), 2000.
- Fredskild, B.: The Holocene vegetational development of Tugtulligssuaq and Qeqertat, northwest Greenland, *Meddelelser om Grønland, Geoscience*, 14, 1-20, 1985.
- Fredskild, B., Andrews, J.T.: Holocene pollen records from West Greenland, in: Andrews, J.T. (Ed.), *Quaternary Environments: Eastern Canadian Arctic, Baffin Bay and Western Greenland*, Allen and Unwin, Boston, 643–681, 1985.
- 795 Funder, S.: Holocene (10,000-0 years BC) climates in Greenland, and North Atlantic atmospheric circulation, *Danish Meteorol. Inst. Climatol. Pap.*, 4, 175–181, 1978.
- Gajewski, K.: Quantitative reconstruction of Holocene temperatures across the Canadian Arctic and Greenland, *Glob. Planet. Change*, 128, 14–23, <https://doi.org/10.1016/j.gloplacha.2015.02.003>, 2015.
- 800 Gibb, O.T., Steinhauer, S., Frechette, B., de Vernal, A., Hillaire-Marcel, C.: Diachronous evolution of sea surface conditions in the Labrador Sea and Baffin Bay since the last deglaciation, *The Holocene*, 25, 1882–1897,

<https://doi.org/10.1177/0959683615591352>, 2015.

Guiot, J., Gally, Y.: R Package: bioindic, 2014.

Head, M.J.: Modern dinoflagellate cysts and their biological affinities, in: *Palynology: Principles and Applications*, Jansonius, J., McGregor, D.C. (Eds.), American Association of Stratigraphic Palynologists Foundation, 1197–1248, 1996.

805 Hernández-Almeida, I., Cortese, G., Chen, M.-T., Kucera, M.: Environmental determinants of radiolarian assemblages in the western Pacific since the last deglaciation, *Paleoceanography*, 32, 830–847, <https://doi.org/10.1002/2017PA003159>, 2017.

Hohmann, S., Kucera, M., de Vernal, A.: Identifying the signature of sea-surface properties in dinocyst assemblages: Implications for quantitative palaeoceanographical reconstructions by transfer functions and analogue techniques, *Mar. Micropaleontol.*, 159, 101816, <https://doi.org/https://doi.org/10.1016/j.marmicro.2019.101816>, 2020.

Holzwarth, U., Esper, O., Zonneveld, K.A.F.: Distribution of organic-walled dinoflagellate cysts in shelf surface sediments of the Benguela upwelling system in relationship to environmental conditions, *Mar. Micropaleontol.*, 64, 91–119, <https://doi.org/10.1016/j.marmicro.2007.04.001>, 2007.

815 Imbrie, J., Kipp, N.G.: A new micropaleontological method for quantitative paleoclimatology: Application to a late Pleistocene Caribbean core, in: *The Late Cenozoic Glacial Ages*, Turekian, K.K. (Ed.), Yale University Press, New Haven, 71–181, 1971.

Jennings, A.E., Walton, M.E., Ó Cofaigh, C., Kilfeather, A., Andrews, J.T., Ortiz, J.D., De Vernal, A., Dowdeswell, J.A.: Paleoenvironments during Younger Dryas-Early Holocene retreat of the Greenland Ice Sheet from outer Disko Trough, central west Greenland, *J. Quat. Sci.*, 29, 27–40, <https://doi.org/10.1002/jqs.2652>, 2014.

820 Juggins, S.: *Rioja: Analysis of Quaternary science data*, 2017.

Juul-Pedersen, T., Arendt, K.E., Mortensen, J., Blicher, M.E., Søgaard, D.H., Rysgaard, S.: Seasonal and interannual phytoplankton production in a sub-Arctic tidewater outlet glacier fjord, SW Greenland, *Mar. Ecol. Prog. Ser.*, 524, 27–38, <https://doi.org/10.3354/meps11174>, 2015.

825 Kaufman, D., McKay, N., Routson, C., Erb, M., Dätwyler, C., Sommer, P.S., Heiri, O., Davis, B.: Holocene global mean surface temperature, a multi-method reconstruction approach, *Sci. Data*, 7, 201, <https://doi.org/10.1038/s41597-020-0530-7>, 2020.

Knudsen, K.L., Stabell, B., Seidenkrantz, M.S., Eiríksson, J., Blake, W.: Deglacial and Holocene conditions in northernmost Baffin Bay: Sediments, foraminifera, diatoms and stable isotopes, *Boreas*, 37, 346–376, <https://doi.org/10.1111/j.1502-3885.2008.00035.x>, 2008.

830 Koc, N., Jansen, E., Hafliðason, H.: Paleoceanographic reconstruction of surface ocean conditions in the Greenland, Iceland, and Norwegian seas through the last 14,000 years based on diatoms, *Quat. Sci. Rev.*, 12, 115–140, [https://doi.org/10.1016/0277-3791\(93\)90012-B](https://doi.org/10.1016/0277-3791(93)90012-B), 1993.

Kokinos, J.P., Eglinton, T.I., Gon, M.A.: Characterization of a highly resistant biomacromolecular material in the cell wall of a marine dinoflagellate resting cyst, *Organic*, 28, 265–288, [https://doi.org/10.1016/S0146-6380\(97\)00134-4](https://doi.org/10.1016/S0146-6380(97)00134-4), 1998.

- 835 Krawczyk, D., Witkowski, A., Moros, M., Lloyd, J., Kuijpers, A., Kierzek, A.: Late-Holocene diatom-inferred reconstruction of temperature variations of the West Greenland Current from Disko Bugt, Central West Greenland, *The Holocene*, 20, 659–666, <https://doi.org/10.1177/0959683610371993>, 2010.
- Krawczyk, D.W., Witkowski, A., Juul-Pedersen, T., Arendt, K.E., Mortensen, J., Rysgaard, S.: Microplankton succession in a SW Greenland tidewater glacial fjord influenced by coastal inflows and run-off from the Greenland Ice Sheet, *Polar Biol.*, 38, 1515–1533, <https://doi.org/10.1007/s00300-015-1715-y>, 2015.
- 840 Krawczyk, D.W., Meire, L., Lopes, C., Juul-Pedersen, T., Mortensen, J., Li, C.L., Krogh, T.: Seasonal succession, distribution, and diversity of planktonic protists in relation to hydrography of the Godthåbsfjord system (SW Greenland), *Polar Biol.*, 41, 2033–2052, <https://doi.org/10.1007/s00300-018-2343-0>, 2018.
- Kucera, M., Weinelt, Mara, Kiefer, T., Pflaumann, U., Hayes, A., Weinelt, Martin, Chen, M., Mix, A.C., Barrows, T.T., Cortijo, E., Duprat, J., Juggins, S., Waelbroeck, C.: Reconstruction of sea-surface temperatures from assemblages of planktonic foraminifera: multi-technique approach based on geographically constrained calibration data sets and its application to glacial Atlantic and Pacific Oceans, *Quat. Sci. Rev.*, 24, 951–998, <https://doi.org/10.1016/j.quascirev.2004.07.014>, 2005.
- 845 Lamb, H.H.: The early medieval warm epoch and its sequel, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 1, 13–37, [https://doi.org/10.1016/0031-0182\(65\)90004-0](https://doi.org/10.1016/0031-0182(65)90004-0), 1965.
- Lamb, H.H.: *Climate, history and the modern world*, Routledge, 2002.
- Larsen, N.K., Kjær, K.H., Lecavalier, B., Bjørk, A.A., Colding, S., Huybrechts, P., Jakobsen, K.E., Kjeldsen, K.K., Knudsen, K.-L., Odgaard, B.V., Olsen, J.: The response of the southern Greenland ice sheet to the Holocene thermal maximum, *Geology*, 43, 291–294, 2015.
- 855 Legendre, P., Gallagher, E.D.: Ecologically meaningful transformations for ordination of species data, *Oecologia*, 129, 271–280, <https://doi.org/10.1007/s004420100716>, 2001.
- Lepš, J., Šmilauer, P. (Eds.): *Multivariate analysis of ecological data using CANOCO*, Cambridge university press, 2003.
- Levac, E., De Vernal, A., Blake, W.: Sea-surface conditions in northernmost Baffin Bay during the Holocene: Palynological evidence, *J. Quat. Sci.*, 16, 353–363, <https://doi.org/10.1002/jqs.614>, 2001.
- 860 Limoges, A., Weckström, K., Ribeiro, S., Georgiadis, E., Hansen, K.E., Martinez, P., Seidenkrantz, M.S., Giraudeau, J., Crosta, X., Massé, G.: Learning from the past: Impact of the Arctic Oscillation on sea ice and marine productivity off north-west Greenland over the last 9,000 years, *Glob. Chang. Biol.*, 26, 6767–6786, <https://doi.org/10.1111/gcb.15334>, 2020.
- Lloyd, J.M., Kuijpers, A., Long, A., Moros, M., Park, L.A.: Foraminiferal reconstruction of mid- to late-Holocene ocean circulation and climate variability in Disko Bugt, West Greenland, *The Holocene*, 17, 1079–1091, <https://doi.org/10.1177/0959683607082548>, 2007.
- 865 Lopes, C., Mix, A.C., Abrantes, F.: Environmental controls of diatom species in northeast Pacific sediments, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 297, 188–200, <https://doi.org/10.1016/j.palaeo.2010.07.029>, 2010.
- Mann, M.E.: Medieval Climatic Optimum. *Encycl. Glob. Environ. Chang.*, 1, 514–516, 2002.

- Marcott, S.A., Shakun, J.D., Clark, P.U., Mix, A.C.: A Reconstruction of regional and global temperature for the past 11,300 years, *Science*, 339, 1198–1201, <https://doi.org/10.1126/science.1228026>, 2013.
- 870
- Marret, F.: Les effets de l'acétytolysé sur les assemblages de kystes de dinoflagelle's, *Palynosciences*, 2, 267–272, 1993.
- Matthiessen, J.: Distribution patterns of dinoflagellate cysts and other organic-walled microfossils in recent Norwegian-Greenland Sea sediments, *Mar. Micropaleontol.*, 24, 307–334, [https://doi.org/10.1016/0377-8398\(94\)00016-G](https://doi.org/10.1016/0377-8398(94)00016-G), 1995.
- Matthiessen, J., Baumann, K.H., Schröder-Ritzrau, A., Hass, C., Andruleit, H., Baumann, A., Jensen, S., Kohly, A., Pflaumann, U., Samtleben, C., Schäfer, P., Thiede, J.: Distribution of calcareous, siliceous and organic-walled planktic microfossils in surface sediments of the Nordic Seas and their relation to surface-water masses, in: *The Northern North Atlantic*, Schäfer, P., Ritzrau, W., Schlüter, M., Thiede, J. (Eds.), Springer, Berlin Heidelberg, 105–127, https://doi.org/10.1007/978-3-642-56876-3_7, 2001.
- 875
- McGarigal, K., Stafford, S., Cushman, S.: Discriminant analysis, in: *Multivariate statistics for wildlife and ecology research*, Springer, New York, 129–187, https://doi.org/10.1007/978-1-4612-1288-1_4, 2000.
- 880
- Meyers, P.A.: Organic geochemical proxies of paleoceanographic, paleolimnologic, and paleoclimatic processes, *Org. Geochem.*, 27, 213–250, [https://doi.org/10.1016/S0146-6380\(97\)00049-1](https://doi.org/10.1016/S0146-6380(97)00049-1), 1997.
- Montresor, M., Sgroso, S., Procaccini, G., Wiebe, H.C.F.: Intraspecific diversity in *Scrippsiella trochoidea* (Dinophyceae): evidence for cryptic species, *Phycologia*, 42, 56–70, <https://doi.org/10.2216/i0031-8884-42-1-56.1>, 2003.
- 885
- Morey, A.E., Mix, A.C., Pisias, N.G.: Planktonic foraminiferal assemblages preserved in surface sediments correspond to multiple environment variables, *Quat. Sci. Rev.*, 24, 925–950, <https://doi.org/10.1016/j.quascirev.2003.09.011>, 2005.
- Moros, M., Jensen, K.G., Kuijpers, A.: Mid- to late-Holocene hydrological and climatic variability in Disko Bugt, central West Greenland, *The Holocene*, 16, 357–367, <https://doi.org/10.1191/0959683606hl933rp>, 2006.
- Mudie, P.J.: Circum-Arctic Quaternary and Neogene marine palynofloras: paleoecology and statistical analysis, *Neogene Quat. Dinoflag. cysts acritarchs*, 10, 347–390, 1992.
- 890
- Muller, P.J., Erlenkeuser, H., Von Grafenstein, R.: Glacial-interglacial cycles in oceanic productivity inferred from organic carbon contents in eastern north Atlantic sediment cores, in: *Coastal upwelling, its sediment record, part B: Sedimentary records of ancient coastal upwelling*, Thiede J., Suess, E. (Ed.), Plenum Press, New York, 65–398, 1983.
- Nooteboom, P.D., Bijl, P.K., Sebille, E. Van, Heydt, A.S. Von Der, Dijkstra, H.A.: Transport bias by ocean currents in sedimentary microplankton assemblages: Implications for paleoceanographic reconstructions, *Paleogeography and Paleoclimatology*, 34, 1178–1194, <https://doi.org/10.1029/2019PA003606>, 2019.
- 895
- Nychka, D., Furrer, R., Sain, S.: *fields: Tools for spatial data*, <https://doi.org/10.5065/D6W957CT>, 2017.
- Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H.: *Vegan: Community ecology package (version 2.5-6)*, The Comprehensive R Archive Network, 2019.
- 900
- Ouellet-Bernier, M.M., de Vernal, A., Hillaire-Marcel, C., Moros, M.: Paleoceanographic changes in the Disko Bugt area, West Greenland, during the Holocene, *Holocene*, 24, 1573–1583, <https://doi.org/10.1177/0959683614544060>, 2014.

- Parkinson, J.E., Baumgarten, S., Michell, C.T., Baums, I.B., Lajeunesse, T.C., Voolstra, C.R.: Gene expression variation resolves species and individual strains among coral-associated dinoflagellates within the genus *Symbiodinium*, *Genome Biol. Evol.*, 8, 665-680, <https://doi.org/10.1093/gbe/evw019>, 2016.
- 905
- Pebesma, E.J.: Multivariable geostatistics in S: the gstat package, *Comput. Geosci.*, 30, 683-691, <https://doi.org/10.1016/j.cageo.2004.03.012>, 2004.
- Pebesma, E.J., Bivand, R.S.: S classes and methods for spatial data: the sp package, *R news*, 5, 9-13, 2005.
- Perner, K., Moros, M., Jennings, A.E., Lloyd, J.M., Knudsen, K.L.: Holocene palaeoceanographic evolution off West Greenland, *The Holocene*, 23, 374-387, <https://doi.org/10.1177/0959683612460785>, 2013.
- 910
- Price, A.M., Pospelova, V., Coffin, M.R.S., Latimer, J.S., Chmura, G.L.: Biogeography of dinoflagellate cysts in northwest Atlantic estuaries, *Ecol. Evol.*, 6, 5648-5662, <https://doi.org/10.1002/ece3.2262>, 2016.
- Price, A.M., Baustian, M.M., Turner, R.E., Rabalais, N.N., Chmura, G.L.: Dinoflagellate cysts track eutrophication in the Northern Gulf of Mexico, *Estuaries and Coasts*, 41, 1322-1336, <https://doi.org/10.1007/s12237-017-0351-x>, 2018.
- 915
- R Core Team: R: A language and environment for statistical computing, 2017.
- Radi, T., de Vernal, A.: Dinocyst distribution in surface sediments from the northeastern Pacific margin (40-60°N) in relation to hydrographic conditions, productivity and upwelling, *Rev. Palaeobot. Palynol.*, 128, 169-193, [https://doi.org/10.1016/S0034-6667\(03\)00118-0](https://doi.org/10.1016/S0034-6667(03)00118-0), 2004.
- Radi, T., de Vernal, A.: Dinocysts as proxy of primary productivity in mid-high latitudes of the Northern Hemisphere, *Mar. Micropaleontol.*, 68, 84-114, <https://doi.org/10.1016/j.marmicro.2008.01.012>, 2008.
- 920
- Radi, T., de Vernal, A., Peyron, O.: Relationships between dinoflagellate cyst assemblages in surface sediment and hydrographic conditions in the Bering and Chukchi seas, *J. Quat. Sci.*, 16, 667-680, <https://doi.org/10.1002/jqs.652>, 2001.
- Rao, C.R.: A review of canonical coordinates and an alternative to correspondence analysis using Hellinger distance, *Quaestio*, 19, 23-63, 1995.
- 925
- Ren, J., Jiang, H., Seidenkrantz, M.S., Kuijpers, A.: A diatom-based reconstruction of Early Holocene hydrographic and climatic change in a southwest Greenland fjord, *Mar. Micropaleontol.*, 70, 166-176, <https://doi.org/10.1016/j.marmicro.2008.12.003>, 2009.
- Ribeiro, S., Amorim, A.: Environmental drivers of temporal succession in recent dinoflagellate cyst assemblages from a coastal site in the North-East Atlantic (Lisbon Bay, Portugal), *Mar. Micropaleontol.*, 68, 156-178, <https://doi.org/10.1016/j.marmicro.2008.01.013>, 2008.
- 930
- Rochon, A., de Vernal, A.: Palynomorph distribution in recent sediments from the Labrador Sea, *Can. J. Earth Sci.*, 31, 115-127, <https://doi.org/10.1139/e94-010>, 1994.
- Rochon, A., de Vernal, A., Turon, J.-L., Matthießen, J., Head, M.J.: Distribution of recent dinoflagellate cysts in surface sediments from the North Atlantic Ocean and adjacent seas in relation to sea-surface, *Am. Assoc. Stratigr. Palynol. Contrib. Ser.*, 35, 1-164, <https://doi.org/10013/epic.14283>, 1999.
- 935

- Rochon, A., Eynaud, F., de Vernal, A.: Dinocysts as tracers of hydrographical conditions and productivity along the ocean margins: Introduction, *Mar. Micropaleontol.*, 68, 1–5, <https://doi.org/10.1016/j.marmicro.2008.04.001>, 2008.
- 940 Rühlemann, C., Müller, P.J., Schneider, R.R.: Organic carbon and carbonate as paleoproductivity proxies: Examples from high and low productivity areas of the tropical Atlantic, in: *Use of proxies in paleoceanography: Examples from the South Atlantic*, Fischer, G., Wefer, G. (Eds.), Springer, Berlin Heidelberg, 315–344, https://doi.org/10.1007/978-3-642-58646-0_12, 1999.
- Saini, J., Stein, R., Fahl, K., Weiser, J., Hebbeln, D., Hillaire, C., de Vernal, A.: Holocene variability in sea ice and primary productivity in the northeastern Baffin Bay, *Arktos*, <https://doi.org/10.1007/s41063-020-00075-y>, 2020.
- 945 Sarnthein, M., Winn, K., Duplessy, J.-C., Fontugne, M.R.: Global variations of surface ocean productivity in low and mid latitudes: Influence on CO₂ reservoirs of the deep ocean and atmosphere during the last 21,000 years, *Paleoceanography*, 3, 361–399, <https://doi.org/10.1029/PA003i003p00361>, 1988.
- Schlitzer, R.: Ocean Data View, odv.awi.de, 2018.
- 950 Schröder-Adams, C.J., van Rooyen, D.: Response of recent benthic foraminiferal assemblages to contrasting environments in Baffin Bay and the northern Labrador Sea, *Northwest Atlantic, Arctic*, 64, 317–341, <https://doi.org/10.14430/arctic4122>, 2011.
- Schweinsberg, A.D., Briner, J.P., Miller, G.H., Bennike, O., Thomas, E.K.: Local glaciation in West Greenland linked to North Atlantic ocean circulation during the Holocene, *Geology*, 45, 195–198, <https://doi.org/10.1130/G38114.1>, 2017.
- 955 Seidenkrantz, M.S., Aagaard-Sørensen, S., Sulsbrück, H., Kuijpers, A., Jensen, K.G., Kunzendorf, H.: Hydrography and climate of the last 4400 years in a SW Greenland fjord: Implications for Labrador Sea palaeoceanography, *Holocene*, 17, 387–401, <https://doi.org/10.1177/0959683607075840>, 2007.
- Seidenkrantz, M.S., Roncaglia, L., Fischel, A., Heilmann-Clausen, C., Kuijpers, A., Moros, M.: Variable North Atlantic climate seesaw patterns documented by a late Holocene marine record from Disko Bugt, West Greenland, *Mar. Micropaleontol.*, 68, 66–83, <https://doi.org/10.1016/j.marmicro.2008.01.006>, 2008.
- 960 Sha, L., Jiang, H., Seidenkrantz, M.S., Knudsen, K.L., Olsen, J., Kuijpers, A., Liu, Y.: A diatom-based sea-ice reconstruction for the Vaigat Strait (Disko Bugt, West Greenland) over the last 5000 yr, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 403, 66–79, <https://doi.org/10.1016/j.palaeo.2014.03.028>, 2014.
- Solignac, S., de Vernal, A., Hillaire-Marcel, C.: Holocene sea-surface conditions in the North Atlantic - Contrasted trends and regimes in the western and eastern sectors (Labrador Sea vs. Iceland Basin), *Quat. Sci. Rev.*, 23, 319–334, <https://doi.org/10.1016/j.quascirev.2003.06.003>, 2004.
- 965 Taylor, F.J.R., Pollinger, U.: The ecology of dinoflagellate, in: *The biology of dinoflagellates*, Taylor, F.J.R. (Ed.), Blackwell Scientific Publications, Oxford, 398–529, 1987.
- Telford, R.J.: *palaeoSig*: Significance tests of quantitative palaeoenvironmental reconstructions, R package version, 1, 2015.
- 970 Telford, R.J., Birks, H.J.B.: The secret assumption of transfer functions: Problems with spatial autocorrelation in evaluating model performance, *Quat. Sci. Rev.*, 24, 2173–2179, <https://doi.org/10.1016/j.quascirev.2005.05.001>, 2005.

- Telford, R.J., Birks, H.J.B.: Evaluation of transfer functions in spatially structured environments, *Quat. Sci. Rev.*, 28, 1309–1316, <https://doi.org/10.1016/j.quascirev.2008.12.020>, 2009.
- Telford, R.J., Birks, H.J.B.: A novel method for assessing the statistical significance of quantitative reconstructions inferred from biotic assemblages, *Quat. Sci. Rev.*, 30, 1272–1278, <https://doi.org/10.1016/j.quascirev.2011.03.002>, 2011.
- 975 ter Braak, C.J.F.: Canonical correspondence analysis: A new eigenvector technique for multivariate direct gradient analysis, *Ecology*, 67, 1167–1179, <https://doi.org/10.2307/1938672>, 1986.
- ter Braak, C.J.F.: Ordination, in: *Data analysis in community and landscape ecology*, Jongman, R.H., ter Braak, C.J.F., van Tongeren, O.F.R. (Eds.), Center for Agricultural Publishing and Documentation, Wageningen, The Netherlands, 91–169, 1987.
- 980 Trachsel, M., Telford, R.J.: Technical note: Estimating unbiased transfer-function performances in spatially structured environments, *Clim. Past*, 12, 1215–1223, <https://doi.org/10.5194/cp-12-1215-2016>, 2016.
- Tremblay, J.É., Anderson, L.G., Matrai, P., Coupel, P., Bélanger, S., Michel, C., Reigstad, M.: Global and regional drivers of nutrient supply, primary production and CO₂ drawdown in the changing Arctic Ocean, *Prog. Oceanogr.*, 139, 171–196, <https://doi.org/10.1016/j.pocean.2015.08.009>, 2015.
- 985 Turner, J.T.: Zooplankton fecal pellets, marine snow, phytodetritus and the ocean’s biological pump, *Prog. Oceanogr.*, 130, 205–248, <https://doi.org/10.1016/j.pocean.2014.08.005>, 2015.
- Turner, J.T.: Zooplankton fecal pellets, marine snow and sinking phytoplankton blooms, *Aquat. Microb. Ecol.*, 27, 57–102, <https://doi.org/10.3354/ame027057>, 2002.
- van Nieuwenhove, N., Head, M.J., Limoges, A., Pospelova, V., Mertens, K.N., Matthiessen, J., De Schepper, S., de Vernal, A., Eynaud, F., Londeix, L., Marret, F., Penaud, A., Radi, T., Rochon, A.: An overview and brief description of common marine organic-walled dinoflagellate cyst taxa occurring in surface sediments of the Northern Hemisphere, *Mar. Micropaleontol.*, 159, 101814, <https://doi.org/10.1016/j.marmicro.2019.101814>, 2020.
- 990 Wall, D., Dale, B.: Modern dinoflagellate cysts and evolution of the Peridinales, *Micropaleontology*, 14, 265–304, 1968.
- Wang, N., Mertens, K.N., Krock, B., Luo, Z., Derrien, A., Pospelova, V., Liang, Y., Bilien, G., Smith, K.F., De Schepper, S., Wietkamp, S., Tillmann, U., Gu, H.: Cryptic speciation in *Protoceratium reticulatum* (Dinophyceae): Evidence from morphological, molecular and ecophysiological data, *Harmful Algae*, 88, 101610, <https://doi.org/10.1016/j.hal.2019.05.003>, 2019.
- 995 Weidick, A., Bennike, O.: Quaternary glaciation history and glaciology of Jakobshavn Isbræ and the Disko Bugt region, West Greenland: A review, *Geological Survey of Denmark and Greenland Bulletin*, <https://doi.org/10.34194/geusb.v14.4985>, 2007.
- 1000 Weidick, A., Bennike, O., Citterio, M., Nørgaard-Pedersen, N.: Neoglacial and historical glacier changes around Kangarsuneq fjord in southern West Greenland, *Geological Survey of Denmark and Greenland Bulletin*, <https://doi.org/10.34194/geusb.v27.4694>, 2012.
- Wigley, T.M.L., Ingram, M.J., Farmer, G.: Past climates and their impact on man: A review, in: *Climate and History*,

- 1005 Cambridge University Press, New York, 3–50, 1981.
- Young, N.E., Briner, J.P., Stewart, H.A.M., Axford, Y., Csatho, B., Rood, D.H., Finkel, R.C.: Response of Jakobshavn Isbræ, Greenland, to Holocene climate change, *Geology*, 39, 131–134, <https://doi.org/10.1130/G31399.1>, 2011.
- Zamelczyk, K., Rasmussen, T.L., Husum, K., Hafliðason, H., de Vernal, A., Ravna, E.K., Hald, M., Hillaire-Marcel, C.: Paleoceanographic changes and calcium carbonate dissolution in the central Fram Strait during the last 20 ka, *Quat. Res.*, 1010 78, 405–416, <https://doi.org/10.1016/j.yqres.2012.07.006>, 2012.
- Zonneveld, K.A.F., Siccha, M.: Dinoflagellate cyst based modern analogue technique at test - A 300 year record from the Gulf of Taranto (Eastern Mediterranean), *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 450, 17–37, <https://doi.org/10.1016/j.palaeo.2016.02.045>, 2016.
- Zonneveld, K.A.F., Versteegh, G.J.M., de Lange, G.J.: Preservation of organic-walled dinoflagellate cysts in different oxygen 1015 regimes: a 10,000 year natural experiment, *Mar. Micropaleontol.*, 29, 393–405, [https://doi.org/10.1016/S0377-8398\(96\)00032-1](https://doi.org/10.1016/S0377-8398(96)00032-1), 1997.
- Zonneveld, K.A.F., Versteegh, G., Kodrans-Nsiah, M.: Preservation and organic chemistry of Late Cenozoic organic-walled dinoflagellate cysts: A review, *Mar. Micropaleontol.*, 68, 179–197, <https://doi.org/10.1016/j.marmicro.2008.01.015>, 2008.
- 1020 Zonneveld, K.A.F., Versteegh, G.J.M., Kasten, S., Eglinton, T.I., Emeis, K., Huguet, C., Koch, B.P.: Selective preservation of organic matter in marine environments; processes and impact on the sedimentary record, *Biogeosciences*, 7, 483–511, <https://doi.org/10.5194/bg-7-483-2010>, 2010.
- Zonneveld, K.A.F., Gray, D., Kuhn, G., Versteegh, G.J.M.: Postdepositional aerobic and anaerobic particulate organic matter degradation succession reflected by dinoflagellate cysts: The Madeira Abyssal Plain revisited, *Mar. Geol.*, 408, 87–109, 1025 <https://doi.org/10.1016/j.margeo.2018.11.010>, 2019.