

Multi-scale variations of subglacial hydro-mechanical conditions at Kongsvegen glacier, Svalbard

Coline Bouchayer^{1,2}, Ugo Nanni², Pierre-Marie Lefeuvre³, John Hult², Louise Steffensen Schmidt², Jack Kohler³, François Renard^{1,4}, and Thomas V. Schuler²

¹The Njord Centre, Departments of Geosciences and Physics, University of Oslo, 0316 Oslo, Norway

²Department of Geosciences, University of Oslo, 0316 Oslo, Norway

³Norwegian Polar Institute, Tromsø, Norway

⁴ISTerre, Univ. Grenoble Alpes, Grenoble INP, Univ. Savoie Mont Blanc, CNRS, IRD, Univ. Gustave Eiffel, 38000 Grenoble, France

Correspondence: Coline Bouchayer (colili@uio.no)

Abstract. The flow of glaciers is largely controlled by changes at the ice-bed interface, where basal slip and sediment deformation drive basal glacier motion. Determining subglacial conditions and their responses to hydraulic forcing remains challenging due to the difficulty of accessing the glacier bed. Here, we monitor the interplay between surface runoff and hydro-mechanical conditions at the base of the Kongsvegen glacier in Svalbard. From ~~spring~~-July 2021 to ~~summer~~-August 2022, we measure both subglacial water pressure and till strength. Additionally, we derive subglacial hydraulic gradient and radius ~~over a kilometre scale~~ of channelized subglacial drainage system from seismic power, recorded at the glacier surface. To characterize the variations in the subglacial conditions caused by changes in surface runoff, we investigate the variations of the following hydro-mechanical properties: measured water pressure, measured sediment ploughing forces and derived hydraulic gradient and radius, over seasonal, multi-day and diurnal time-scales. We discuss our results in light of existing theories of subglacial hydrology and till mechanics to describe subglacial conditions. We find that during the short, low melt intensity season in 2021, the subglacial drainage system ~~evolves~~-evolved at equilibrium with runoff, increasing its capacity as the melt season ~~progresses~~progressed. In contrast, during the long and high melt intensity season in 2022, the subglacial drainage system ~~evolves~~-evolved transiently to respond to the abrupt and large water supply. We suggest that in ~~this~~-the latter configuration, the drainage capacity of the preferential drainage axis ~~is~~-was exceeded, promoting the expansion of hydraulically connected regions, and local weakening of ice-bed coupling and hence, enhanced sliding.

1 Introduction

Glacier ice loss represents one of the greatest global environmental risks in a changing climate due to its contribution to sea level rise (Moon et al., 2018; Rounce et al., 2023)(IPCC, 2021). Glacier flow is controlled by (i) viscous deformation of the ice, (ii) slip at the ice-bed interface, and (iii) subglacial sediment deformation. Basal motion (ii and iii) is responsible for most of the short-term velocity variations and rapid ice flow (e.g., Kamb and Engelhardt, 1991; Kamb, 2001)(e.g., Lliboutry, 1968; Weertman, 1957; but underlying processes are still poorly understood due to the difficulty in observing the subglacial environment -Unravelling

(Rada and Schoof, 2018; Gimbert et al., 2021a, e.g.,). Unraveling the response of subglacial conditions to surface water input and its consequences for glacier dynamics is key to reduce uncertainties in future projections of ice mass loss to the oceans (e.g., Stoeker, 2014; IPCC, 2019)(e.g., Maier et al., 2022; Rounce et al., 2023).

25 Rates of basal motion depend on the gravitational driving stress imposed by the glacier geometry, and on frictional properties at the base of the glacierbase, which, in turn, are governed by the thermal regime and hydro-mechanical conditions (e.g., Alley, 1989; Engelhardt and Kamb, 1997)(e.g., Lliboutry, 1968; Gilbert et al., 2022, 2014). The rate of water supply from the glacier surface to the glacier base and the state of the subglacial drainage system control the basal water pressure (Röthlisberger, 1972). Basal conditions therefore vary on several time scales (e.g., hourly, daily, seasonally), inherited from typical variations of runoffglacier surface runoff (hereafter referred to as runoff), and subglacial drainage system evolution(Weertman, 1957; Lliboutry, 1968; Schoof, 2005). Following the variations of surface energy balance and rainfall, glacier surface runoff (hereafter referred to as runoff) runoff typically displays characteristic diurnal variations, superimposed on multi-day weather cycles and a pronounced seasonality (Röthlisberger and Lang, 1987; Nienow et al., 2017) (Sugiyama and Gudmundsson, 2003; Nanni et al., 2023). Variations in water pressure are caused both by variations in runoff as well as by the state and evolution of bulk transmissivity the efficiency of the subglacial drainage system (Rada and Schoof, 2018) . Previous works have identified different components of the subglacial drainage system that convey and store water along the glacier bed. These include water sheets (Weertman, 1972; Walder, 1982; Creyts and Schoof, 2009), cavities in the lee side of bedrock obstacles (Lliboutry, 1968; Iken, 1981), linked cavities (Kamb, 1987; Walder, 1986) (Kamb, 1987; Walder, 1986; Nanni et al., 2023) and channels incised into the ice or subglacial substrate (Röthlisberger, 1972; Nye, 1976; Hooke et al., 1990; Walder and Fowler, 1994) 40 -Water sheets (Röthlisberger, 1972; Nye, 1976; Gulley, 2009). Water sheet and cavity systems are spatially distributed across the glacier base, flow bed. Flow pathways are typically tortuous and these systems are characterized as hydraulically inefficient and often operate, operating at high water pressures. In contrast, channelized drainage systems along preferential drainage axes is highly are localized and often hydraulically efficient, which permits fast flow of water, and operates operating typically at low water pressures. Glaciers resting on a tillbase exhibit a complex 45 For glaciers resting on deformable sediments, i.e. till, the subglacial drainage system -While some water drains through the pore space of the granular material, also drainage along the ice-till interface has been described and various drainage structures have been proposed -For glaciers lying on fine grain sediments, water is expected to flow through localized flow pathways termed canals. These canals is more complex. For instance, Flowers and Clarke (2002a, b) proposed a macro-porous horizon as a continuum concept to comprise inter-granular pore spaces, thin films, cavities, or larger gaps. Additionally, water may circulate through channel-like structures called canals that are incised into the sediment and/or ice by erosion and close through the creep of ice and sediments (Walder and Fowler, 1994; Ng, 2000). Flowers and Clarke (2002a, b) proposed a macro-porous horizon as a continuum concept to comprise inter-granular pore spaces, thin films, cavities, or larger gaps- 50 Water pressure at the glacier base directly influences ice-bed coupling, as well as it affects the shear strength of the till (Lliboutry, 1968; Clarke, 2005; Damsgaard et al., 2020; Zoet and Iverson, 2020; Hansen and Zoet, 2022; Tsai et al., 2022) 55 -This double-tracked influence gives rise and the ice-bed coupling (Damsgaard et al., 2020; Zoet and Iverson, 2020; Hansen and Zoet, 2022) . This influence leads to complex behavior in the relationship between water pressure and basal motion, with an exact formula-

tion still being debated (De Fleurian et al., 2014; Benn et al., 2019; Thøgersen et al., 2019; Gimbert et al., 2021a; Gilbert et al., 2022). Whereas high water pressure weakens the ice-bed coupling and therefore promotes sliding, low water pressure has the opposite effect. Similarly, water pressure influences the yield strength of the till, for which (De Fleurian et al., 2014; Zoet and Iverson, 2020). Currently, till deformation is best described by a Coulomb-plastic behavior is widely accepted. Motivation rheology. Motivations for this rheology comes come from granular and soil mechanics (e.g., Schofield and Wroth, 1968; Terzaghi et al., 1996; Mitchell et al., 2005), field measurements on subglacial till deformation (Hooke et al., 1997; Kavanaugh and Clarke, 2006), laboratory experiments on till (e.g., Kamb and Engelhardt, 1991; Iverson et al., 1998; Tulaczyk et al., 2000; Iverson et al., 2007; Zoet and Iverson et al., 1998, 2007; Zoet and Iverson, 2020), inversion of subglacial mechanics from ice-surface velocities (e.g., Tulaczyk et al., 2000; Walker et al., 2012; Goldberg et al., 2014; Minchew et al., 2016; Gillet-Chaulet et al., 2016), and numerical experiments (Iverson and Iverson, 2001; Kavanaugh and Clarke, 2006; Damsgaard et al., 2013, 2016). Pressure-gradient driven transport of water into or out of the till pore space affects rheological properties of the till and thus influence rates of sediment deformation (Iverson et al., 1995; Hooke et al., 1997; Tulaczyk et al., 2000; Hansen and Zoet, 2022; Warburton et al., 2023) (Iverson et al., 1995; Warburton et al., 2023). Alterations in subglacial hydrological and mechanical hydro-mechanical conditions have the capacity to change the overall glacier dynamics, sometimes leading to partial or complete destabilisation (Thøgersen et al., 2019).

destabilization (Thøgersen et al., 2019; Zhan, 2019). An example of glacier destabilization are glacier surge, characterized by significant ice flow accelerations often accompanied by sudden and rapid advances of glaciers (Truffer et al., 2021).

To investigate subglacial properties, borehole measurements have provided direct access to the glacier bed and have often been instrumented to monitor water pressure (e.g. Hubbard et al., 1995; Sugiyama et al., 2011; Andrews et al., 2014; Doyle et al., 2018; Sugiyama et al., 2019; Rada and Schoof, 2018; Rada Giacaman and Schoof, 2023). Studies based on numerous boreholes highlighted the simultaneous and different behaviour experienced by different areas of the bed (e.g., Andrews et al., 2014; Rada and Schoof, 2018). Such processes are yet poorly described by current drainage models (Flowers, 2015). For example, some areas of the glacier bed can be connected to the main subglacial drainage pathway, whereas other parts are not, and further areas even display alternating behavior (e.g., Gordon et al., 1998; Kavanaugh and Clarke, 2000). Additional studies pointed out that some areas of the bed show signs of hydraulic isolation (e.g., Murray and Clarke, 1995; Andrews et al., 2014; Hoffman et al., 2016; Rada and Schoof, 2018). Many long-term observations have a marked spatial show the spatial and temporal heterogeneity of water pressure variations, even at small spatial scales (Murray and Clarke, 1995; Iken and Truffer, 1997; Fudge et al., 2008; Andrews et al., 2014; Rada and Schoof, 2018), as well as sudden changes, indicative for reorganisation of indicative for reorganization of the drainage system (Gordon et al., 1998; Kavanaugh and Clarke, 2000; Schuler et al., 2002), and the presence of hydraulic isolation (e.g., Murray and Clarke, 1995; Andrews et al., 2014; Hoffman et al., 2016; Rada and Schoof, 2018). Adequately representing such heterogeneity remains a challenge for subglacial drainage models (Flowers, 2015), although some progress have been made recently (Downs et al., 2018; Gilbert et al., 2022; Sommers et al., 2023). Borehole instrumentation has also been used to collect information on till shear strength using ploughmeters, on sliding rates using drag spools, and on till deformation using inclinometers (e.g., Fischer and Clarke, 1994; Fischer et al., 1998, 1999, 2001; Porter et al., 1997). Borehole studies have therefore provided crucial information on the local hydro-mechanical adaptation of the subglacial environment to changes in

runoff and ~~the following its~~ impact on glacier dynamic. Due ~~to~~ the co-existence of different subglacial drainage system components and their temporal evolution, the interpretation of the hydro-mechanical conditions from borehole studies solely remains very local and challenging to extrapolate to the glacier-scale.

95 Recent studies have shown the potential of near-surface cryoseismology to bridge the gap between observations at different scales (Podolskiy and Walter, 2016), for instance to detect brittle fractures related to crevasse opening (e.g., Roux et al., 2008; Nanni et al., 2022), stick-slip motion at the glacier base (e.g., Wiens et al., 2008; Gräff et al., 2021; Köpfler et al., 2022; Hudson et al., 2023), iceberg calving (~~e.g., Köhler et al., 2015; Sergeant et al., 2018~~)(e.g., Köhler et al., 2015), or to infer subglacial hydraulic conditions across various temporal (sub-daily to multi-year) and spatial (decameter to kilometer) 100 scales (Bartholomäus et al., 2015; Nanni et al., 2020; Lindner et al., 2020; Nanni et al., 2021; Labedz et al., 2022a). The later case is based on the principle that turbulent-water-flow generates high-frequency seismic ~~tremor-vibrations~~ that can be used to quantify relative changes in the most active part of the subglacial drainage system (Gimbert et al., 2016; Nanni et al., 2021).

~~The Given the wide range of scales over which subglacial conditions can vary subglacial conditions and the multitude of involved processes, request processes at play, appropriate sampling and analysis strategies to better understand are essential~~ 105 ~~to understand how the drainage system evolves, how till behaves, and the interplay between drainage system evolution, till rheology and~~ consequences for basal motion. ~~In this study~~Here, we present the ~~results analysis~~ of a comprehensive, subglacial multi-sensor record ~~, complemented by a cryoseismological~~ dataset, all simultaneously acquired at Kongsvegen glacier, a surge-type glacier in Svalbard. ~~From June 2021 to August 2022, we measured both subglacial water pressure and till strength, and derived subglacial hydraulic gradient and radius of channelized subglacial drainage from cryoseismology.~~ The field instru- 110 mentation has been designed to optimize interpretation by co-locating the instruments in a single borehole and accompanying measurements of glacier surface velocity and energy-balance driven estimates of surface runoff. ~~The records span the period from June 2021 to August 2022 and cover two contrasting melt seasons.~~ Our results document runoff-induced variations in the subglacial hydro-mechanical conditions at different glaciologically significant time scales (diurnal, multi-day and seasonal).

2 Study area

115 Kongsvegen glacier (hereafter, Kongsvegen, $78^{\circ} 48' \text{N}$, $12^{\circ} 59' \text{E}$) is located near the Ny-Ålesund research station on the ~~northwest north-western~~ coast of Svalbard (Fig. 1a). The glacier covers ~~a surface area of~~ $\sim 108 \text{ km}^2$, is $\sim 25.5 \text{ km}$ long (in 2010, RGI, 2017) and its ice thickness at the drilling site ($78^{\circ} 18' \text{N}$, $17^{\circ} 13' \text{E}$) is $\sim 350 \text{ m}$. The surface slope ranges from 0.5° to 2.5° with a north-western orientation (Hagen et al., 1993). The glacier drains into Kongsfjorden fjord and its terminus is grounded below sea level. Typical for an Arctic glacier, its ice is polythermal with ~~a temperate base and~~ an 120 upper layer of 50-130 m thick cold ice ~~covering temperate ice until the bed~~. Ice in the accumulation zone is temperate while it is frozen to the lateral margins in the ablation zone (Björnsson et al., 1996). The glacier rests on fine-grained sandstone and sand/silt glacio-marine sediments (Hjelle, 1993; Murray and Booth, 2010).

Kongsvegen is a surge-type glacier, with surges reported around 1800, 1869 and 1948 (Liestøl, 1988; Woodward et al., 2002). ~~The glacier is in its quiescent phase since~~Since the last surge~~and~~, ~~the glacier~~ exhibits low velocities of about 3 m a^{-1} .

125 Melvold and Hagen (1998) showed that the mass transported down glacier is only about 3-20% of the annual mass gained in the accumulation area, symptomatic for a surge-type glacier in its quiescent stage. Surface velocities recorded nearby the equilibrium line indicate that the glacier is accelerating since 2014, indicating the possibility for an imminent **fast-flow-event surge** (Fig. 1 b).

We conducted field campaigns to install and maintain a set of instruments at Kongsvegen starting in September 2020. Borehole and surface instrumentation **was-were** completed in April 2021 and the data collected **cover-the-period-span** from June 26, 2021 to August 8, 2022. The following section details the instrumentation **and-its-usage**, the data collected and the associated processing.




Figure 1. Study site and field methods. (a) Location of Kongsvegen glacier in Svalbard. The green star indicates the instrument site where data were collected and the red circles indicate the position of the GNSS KNG6 and KNG7 (Credits: NPI/Copernicus Sentinel data). (b) Annual surface velocity near the equilibrium line of Kongsvegen glacier from 2005 to 2022, with an acceleration around 2014 witnessing that this glacier is closer to a surge event (personal communication from J. Kohler). (c) Main borehole with the smaller secondary borehole where the return pump is installed. (d) Drilling installation. (e) Sediment sample recovered with a sediment sampling tool at the bottom of the borehole.

3 Methods

3.1 Borehole

135 On April 25, 2021, we drilled a borehole near the long term equilibrium line ($78^{\circ}18'N$, $17^{\circ}13' E$, Fig. 1a) of Kongsvegen and placed instruments along the borehole and at its base. The borehole was drilled using a hot water drilling system, consisting of three high-pressure hot water machines (Kärcher HDS1000D), a 1/2 inch diameter high-pressure hose, a 2 m drill stem with a 2.3 mm diameter nozzle, a pulley, a tripod, and water tanks (three 1000-liter IBC tanks, Fig. 1d). Since the glacier was in winter conditions and liquid water was not available, the water flowing out of the borehole was captured in an auxiliary hole for recycling (Fig. 1c). During drilling, the ~~borehole water level started dropping~~ level of water in the borehole started to drop when the drilling reached a depth of 260 m, indicating a connection to an active part of the drainage system. A sediment sample collected at the bottom of the ~ 350 m borehole provides evidence for the existence of a sediment layer at the glacier bed (Fig. 1e). The borehole location has been chosen based on the work of Scholzen et al. (2021); Pramanik et al. (2020) who ~~suggest~~ suggested the existence of a preferential drainage ~~pathway~~ axis in close proximity to this site.

145 At the bottom of the borehole, we installed a ploughmeter, i.e. a 1.4 m long steel rod equipped with strain gauges, to monitor mechanical conditions of the subglacial till (e.g. Humphrey et al., 1993; Iverson et al., 1994; Fischer and Clarke, 1994; Porter et al., 1997; Fischer et al., 1998, 2001; Murray and Porter, 2001; Boulton et al., 2001). The tip of the instrument penetrates into the till whereas its upper part remains in the borehole and is trapped in the ice. As the glacier moves across its bed, the ploughmeter tip is dragged through the sediment and the device bends, which is sensed by the strain gauges. Strain on the ploughmeter is measured using a Wheatstone bridge for each pair of strain gauges in two perpendicular axes (Hoffmann, 1974). The exact insertion depth of the device into the till is ~~unknown~~ uncertain. However, based on previous experiences with identical devices, we estimate the penetration depth to be ~~around~~ in the range 10 to 40 cm, which is sufficient to ensure that all strain gauges are immersed in ~~subglacial material~~ the subglacial till. Just above the upper end of the ploughmeter, ~~(which is about 1 m above the glacier bed)~~, we installed a vibrating wire pressure sensor (Geokon 4500SH, <2 kPa accuracy and 0.5 kPa resolution) to monitor subglacial water pressure, p . Sensor readings and data recording ~~is~~ are performed using a Campbell Scientific CR1000X data-logger, recording data at one minute intervals. Ploughmeter ~~readings~~ measurements are converted to force F experienced by the instrument while being dragged through the sediment, ~~based on a calibration performed~~ F . To do the conversion, we performed a calibration in the laboratory prior to field deployment. ~~To perform the calibration, loads: masses of 10 kg and 50 kg (~ 100 and ~ 500 N)~~ have been applied to the free end of the horizontally fixed ploughmeter ~~by hanging a known mass on its free end. We repeated these measurements~~ in eight orientations (0 to 315° every 45°) ~~using masses of 10 kg and 50 kg corresponding to loads of ~ 100 and ~ 500 N, respectively~~). After applying the calibration component-wise, we derive F from the ~~X and Y~~ x and y components using Pythagoras' theorem.

3.2 Near-surface instrumentation

At ~~~ 100 m~~ ~ 100 m from the borehole, we installed a three-component geophone (DiGOS, 4.5 Hz) ~~was installed~~ ~ 1.5 m into the ice to ensure good coupling ~~and~~ with the ice and to prevent melt-out during summer. A DiGOS datacube, that comprises a

digitizer and a data-logger, controlled the sampling rate (100 Hz) and recorded the signals. In this study, we ~~analyse~~analyze seismicity in the 3-10 Hz frequency band as a proxy for hydraulic conditions (Bartholomaus et al., 2015; Gimbert et al., 2016; Nanni et al., 2020; Lindner et al., 2020; Labeledz et al., 2022a). In addition, we used data of two surface stations (KNG6 and KNG7) that recorded positions using Global Navigation Satellite System (GNSS), from which we ~~derive~~derived surface velocities. The stations are located at distances of 740 ± 10 m (KNG6, 78.78067°N , 13.15153°E) and 3100 ± 10 m (KNG7, 78.76770°N , 13.23962°E) upstream of the drill site. Data are ~~recorded at five-second intervals continuously~~recorded at five-second intervals between April 1 and September 1, when solar panels help ~~conserving~~conserve battery capacity, but only for one hour per day during the rest of the year. Gaps occurred in the time series when low battery voltage caused ~~failure of~~ the data logger to fail. The GNSS data are processed assuming that the rover station is static for one hour. Then the velocities are averaged daily to reduce the velocity uncertainties caused by the relatively low speed of the glacier. The Norwegian Mapping Authority's permanent network base station in Ny Ålesund is used as reference (baseline of ~ 30 km). The two velocity records are merged, i.e. we consider the velocity derived from KNG6 when available and the one from KNG7 for the rest of the period (the original records for KNG6 and KNG7 can be seen in the Appendix C). We apply a one-week moving median for KNG7 velocity to smooth the record especially during the winter period when the velocities are low. For the long term averaged velocity from 2005 to 2022 (Fig. 1b), annual (April to April) glacier surface velocity was derived from annual GNSS surveys of mass balance stakes Nuth et al. (2012).

3.3 Surface runoff and meteorological conditions

The surface runoff is ~~modelled~~modeled using the CryoGrid community model (Westermann et al., 2023), ~~coupling~~. This model couples a surface energy balance to model and a multi-layer snow pack model enabling simulation of glacier mass balance and freshwater runoff (Schmidt et al., 2023). The model is forced by 3-hourly fields of near surface conditions from the Copernicus Arctic Regional Reanalysis (CARRA, Schyberg et al., 2020; Yang et al., 2021) for 2021 and single-member forecasts by AROME-Arctic (Müller et al., 2017) for 2022. ~~Meteorological forcing includes 2 m air temperature and precipitation among other variables.~~ Differences in model results due to different forcing datasets are small in our study area (i.e., $< 2\%$ of the total runoff, Schmidt et al., 2023). Meteorological forcing includes 2 m air temperature and precipitation among other variables. The CryoGrid community model ~~then~~ calculates the surface energy balance to simulate the mass balance components as well as the build-up and decay of seasonal snow. The available surface water in a grid cell is either retained in snow or firn, or runs off under the influence of gravity. The retention is governed by the hydraulic conductivity of the snow, parameterized based on snow grain size, density, and effective water saturation. Depending on temperature conditions, retained water may refreeze, thereby releasing latent energy. Once the retention capacity of a layer is reached, excess water may run off with a time scale depending on surface slope. Schmidt et al. (2023) estimated a standard error of runoff of $0.12 \text{ m w.e. a}^{-1}$. The surface runoff is ~~modelled~~modeled on a 2.5 by 2.5 km grid ~~and we assume that all runoff produced within an area of 6.25 km^2 upstream of our borehole is conveyed at the base without any delay.~~ We determine the surface runoff routed through our study area by summing all surface runoff generated upstream from our study site within the glacierized catchment of Kongsvegen's glacier (Scholzen et al., 2021). Since our analysis considers relative changes, only the timing but not the absolute magnitude ~~are of~~

200 ~~interest. We calculate the surface runoff transiting through our study area by integrating all runoff produced in its upglacier hydrological basin, the latter computed using the TopoToolbox (Schwanghart and Scherler, 2014). The temporal resolution of the runoff output is three hours~~ of the runoff is of interest. A complete description of the workflow is given by Schmidt et al. (2023). Using simulated surface runoff to represent local discharge through a given cross-section implicitly assumes transfer of water between the surface and the base within short time, ~~which is supported.~~ The suitability of this hypothesis is reinforced
 205 by in-situ observations from other Svalbard glaciers similar to Kongsvegen (Benn et al., 2009; Gulley, 2009; Bælum and Benn, 2011; Irvine-Fynn et al., 2011) ~~and,~~ and by the good agreement between daily values of simulated runoff, and measured proglacial discharge at the catchment scale (Schmidt et al., 2023). Therefore, we consider relative variations in surface runoff to represent those of subglacial discharge, even though large uncertainties on the magnitude of the subglacial discharge remain.

3.4 Derived subglacial variables

210 3.4.1 Seismic power, hydraulic radius and hydraulic gradient

We calculate the seismic power, P , from the vertical component of the ground velocity using Welch's method over a ~~two-second~~ 2-second time window with a 50% overlap (Welch, 1967; Beyreuther et al., 2010) within the frequency band 3 to 7 Hz. Our choice of this frequency band is based on the dominance of turbulent-water flow-induced seismicity in this ~~band (Bartholomaus et al., 2015;~~ frequency band (Gimbert et al., 2016; Nanni et al., 2020), as opposed to bedload transport that generates seismicity at higher
 215 frequencies (Gimbert et al., 2016). This ~~has been previously observed in other glacial settings (e.g., Preiswerk and Walter, 2018; Lindner et al., 2018)~~ turbulent-water flow-induced seismicity has been observed on different glacial settings (e.g., Bartholomaus et al., 2015; Preiswerk and Walter et al., 2018). Variations in this frequency band are related to changes in hydraulic radius, R , i.e. the ratio of the cross-sectional area of a channelized flow to its wetted perimeter, and in hydraulic gradient, S , i.e. the water pressure gradient in the along flow direction. For open ~~flow;~~ channel flow, i.e. flow with a free surface, as opposed to pressurized flow, where the channel cross-section
 220 is filled with water in all directions, R scales with flow depth and S with slope ~~in the along~~ along the flow direction. For glaciers, Gimbert et al. (2016) expressed P as a function of R and S . The relative changes of these variables are derived from relative changes in P and Q (Gimbert et al., 2016):

$$\frac{S}{S_{ref}} = \left(\frac{P}{P_{ref}} \right)^{24/41} \left(\frac{Q}{Q_{ref}} \right)^{-30/41} \quad (1)$$

$$225 \quad \frac{R}{R_{ref}} = \left(\frac{P}{P_{ref}} \right)^{-9/82} \left(\frac{Q}{Q_{ref}} \right)^{-21/41} \quad (2)$$

where the subscript *ref* represents a reference state that must be defined over the same period for Q and P , but not necessarily for R and S . In our case, the reference state was taken on June 24, 2021 ($Q_{ref} = 0.08 \text{ m}^3 \text{ s}^{-1}$ and $P_{ref} = 180.59 \text{ dB}$) which corresponds to the pre-melt season 2021 value against which we want to evaluate relative changes happening during the melt season. The method is described in detail by Gimbert et al. (2016). Here, we neglect changes in conduit shape, degree of
 230 fullness and number as they have limited impact on the derivation of R and S (Gimbert et al., 2016; Nanni et al., 2020). This

approach allows to estimate the evolution of R and S of the dominating drainage system over an area of $e \sim 1 \text{ km}^2$ around the seismic station (Gimbert et al., 2016; Nanni et al., 2020; Lindner et al., 2020; Nanni et al., 2021; Labeledz et al., 2022a). To simplify notation, we hereafter refer to R/R_{ref} and S/S_{ref} as R and S respectively.

~~We evaluate our results against theoretical considerations relating P and Q (Gimbert et al., 2016), and hydraulic properties and Q (Röthlisberger, 1972; Nanni et al., 2020) to~~ To discuss the state of the subglacial drainage system during the period of our records, ~~we evaluate our results against scaling relationships linking P with Q (Gimbert et al., 2016), and R - S with Q (Röthlisberger, 1972; Nanni et al., 2020).~~ The scaling relationship proposed by Gimbert et al. (2016) defines cases where steady-state channels incised in the ice adapt to changes in Q by only adjusting R (constant S) or the opposite case (constant R). Nanni et al. (2020) adapted Röthlisberger’s (1972) theory to derive scaling relationships between R and Q , and S and Q , both for a steady-state channel evolution incised in the ice ~~as well as and~~ for a channel of static cross-sectional area evolving ~~thus~~ as a rigid pipe (Table 1).

Table 1. Scaling relationships between P and Q , S and Q , and R and Q , for special cases, derived by Gimbert et al. (2016) and Nanni et al. (2020) from Röthlisberger (1972) theory to assess variations in channelized drainage conditions.

Context	Relation	Reference
Change in runoff occurring at constant hydraulic radius	$P \propto Q^{14/3}$	Gimbert et al. (2016)
Change in runoff occurring at constant hydraulic gradient	$P \propto Q^{5/4}$	
Steady-state channel	$R \propto Q^{9/22}$	Röthlisberger (1972); Nanni et al. (2020)
	$S \propto Q^{-2/11}$	

3.5 Processing of time series, catalog of events and classification

To characterize ~~the responses of the subglacial environment to runoff, Q , we analyse~~ subglacial conditions, we analyze the responses of the ~~subglacial variables recorded in the present study, i.e. the measured variables~~ (force, F , ~~the and~~ water pressure, p , ~~and derived, i.e., the~~) and derived variables (turbulent-water-flow-induced seismic power, P , the hydraulic radius, R and the hydraulic gradient, S , ~~in terms of phase relationship to~~) to the runoff, Q . Here we use the term ~~phase~~ ‘phase’ to characterize the time ~~relationships relationship~~ between one of the subglacial variables and ~~events detected in the time series of Q the runoff~~. These phase relationships are ~~analysed~~ analyzed at three different, glaciologically relevant time scales: seasonal, multi-day, and diurnal. To extract the corresponding components at these three time scales, we filtered the time series using a low-pass filter with cutoff at 20 days, a band-pass filter between four and eight days, and a band-pass filter between six hours and 36 hours, respectively. We subdivide the multi-day and diurnal time series into individual events based on Q variations. We define an event by two subsequent minima of Q within the bandwidth investigated. We normalize both Q and the subglacial variables by their respective maxima and subdivide the time into 50 equidistant steps. ~~We choose the~~ The number of time steps was chosen empirically as the minimum number of data points that preserves the shape and characteristics of the event time series (see also Appendix B, Fig. B1). We synthesize the different responses of each subglacial variables, X , to changes in Q using a classification scheme (Fig. 2). Our workflow resembles that developed by Nanni et al. (2020) to understand sub-glacial

subglacial hydrology on hard-bed glaciers, and by Javed et al. (2021) to study storm-induced hydrological conditions variations. The period of records is subdivided ~~;~~ at a multi-day time scales into twelve events (melt season 2021, see Appendix D, Tab. D1) and eight events (melt season 2022, see Appendix D, Tab. D2); and at a diurnal time scale into 96 events (melt season 2021, see Appendix D, Tab. D3) and 85 events (melt season 2022, see Appendix D, Tab. D4).




Figure 2. Phase relationship classification for events. Below each class, the plots in the first row correspond to a representative event for this class with runoff (Q_{norm}) plotted in blue and one subglacial variable (X_{norm} , with X being F , P , p , R or S) plotted in pink against time. The magnitude of the variables is ~~normalised~~normalized between 0 and 1 and the time is re-sampled into 50 time steps. The plots in the second row show the shape of the relationship between the two variables after classification. The solid color points refer to the mean behavior in this class, all individual events from the filtered time series are shown in shaded colors. The black line is the linear regression fitted to the scatter plot. *Preceding class* and *Lagging class* correspond to clockwise or anti-clockwise hysteresis (or time-lag) between the runoff, Q , and the observed variable, while *In-phase class* and *Anti-phase class* correspond to linear relationships. The color scale indicates chronology.

Our classification scheme is based upon the following metrics: the slope, m , of the linear regression between the subglacial variable X (with X being F , P , p , R or S) and Q (black line, Fig. 2); the squared residuals, RSS , between the linear regression and the $X_{norm} - Q_{norm}$ hysteresis loop; θ whose sign indicates the direction of the hysteresis loop, θ . The spread of the data relative to the regression is quantified by the squared residuals (Appendix F, Fig. F1b):

$$RSS = \sum_{i=1}^n r_i^2 \quad (3)$$

where r_i are the residuals of the regression.

The parameter θ expresses the asymmetry of the response with respect to the forcing ~~and its~~. Its sign determines whether the hysteresis is clockwise ~~or counterclockwise~~ (positive) or counterclockwise (negative). We compute θ by comparing the mean of the subglacial variable ~~considered~~ during the rising limb of Q , $\bar{X}_{Q_{rising}}$, to its counterpart during decreasing Q , $\bar{X}_{Q_{falling}}$

270 (Appendix F, Fig. F1a):

$$\theta = \frac{\bar{X}_{Q_{rising}} - \bar{X}_{Q_{falling}}}{\bar{X}_{Q_{falling}}} \quad (4)$$

The sign of θ indicates whether the signal precedes ($\theta > 0$) or lags ($\theta < 0$) Q .

Events are classified according to the phase relationship between Q and the subglacial variable. We distinguish four classes, representing the following cases (Fig. 2):

- 275
- *Preceding class*: The subglacial variable considered precedes Q , $RSS > 2$ & $\theta > 0$,
 - *In-phase class*: The subglacial variable considered lags behind Q , $RSS > 2$ & $\theta < 0$,
 - *In-phase class*: The subglacial variable considered and Q vary in phase, $RSS \leq 2$ & $m > 0$,
 - *Anti-phase class*: The subglacial variable considered and Q vary in anti-phase, $RSS \leq 2$ & $m < 0$.

To discriminate between linear and hysteresis relations, we use a threshold of $RSS = 2$, corresponding to a phase difference
280 of about $\pi/10$ for two sinusoidal variations. ~~Our analysis does not target a two member classification (steady-state or not), but a four member classification of phase relationships which are subsequently interpreted~~ This threshold allows some deviation from a strictly linear behavior and accepts $RSS \leq 2$ still representing linear behavior, thus taking into account uncertainties symptomatic for observations of natural systems. The choice of this threshold is motivated from visual observation of clustering of phase relations (Fig. 2).

285 4 Results

4.1 Overview over the dataset

~~The~~ In Figure 3, we show the meteorological conditions extracted from CARRA/AROME-Arctic, the velocity, u_s , the runoff, Q , the turbulent-water-flow-induced seismic power, P , the water pressure, p and the force acting on the ploughmeter, F ; ~~are displayed in Figure 3.~~ We obtain time series covering two melt seasons (2021 and 2022, Fig. 3) and one winter period
290 (Appendix A, Fig. A1). The meteorological conditions (Figure 3a and b) control the timing and volume of Q resulting from meltwater production or rainfall (Figure 3c and d). We note that the dataset covers two very different melt seasons. While the 2021 melt season is short (67 days from July 1 ~~2021~~ to September 6 ~~2021~~), marked by low temperature oscillating around 0°C and continuous low runoff (lower than $20 \text{ m}^3 \text{ s}^{-1}$), the 2022 melt season is long (at least 83 days because we do not capture the end of this melt season, from May 25 ~~2022~~ to August 16 ~~2022~~) marked by high temperatures (up to 7°C) ~~and~~
295 $^\circ\text{C}$) accompanied by frequent and large excursions of runoff above $20 \text{ m}^3 \text{ s}^{-1}$.

In response to temperature and rainfall variations, Q displays variations on several time scales (Fig. 3c and d, blue line): (i) the seasonal time scale (>20 days) is marked by Q generally being limited to the melt season; (ii) the multi-day superimposed time scale (four to eight days), typically reflects weather variability (warm-spells, e.g., Fig. 3c②), or rainfall, e.g., Fig. 3c④);




Figure 3. Time series of physical quantities measured during the melt seasons 2021 (a, c, e, g) and 2022 (b, d, f, h): (a-b) Temperature (black line), snow fall rate (grey-gray bars) and rainfall rate (light blue) from CARRA/AROME-Artic (Schmidt et al., 2023). The three variables are extracted for the grid point closest to the borehole location. (c-d) Modelled runoff (blue) and measured glacier surface velocity (red). Circled numbers refer to episodes described in the main text. (e-f) seismic power recorded at the surface of the glacier in the 3-10 Hz frequency band (yellow). (g-h) Borehole water pressure (green) and force acting on the ploughmeter (dark red). Blue shaded areas represent the melt seasons. Grey shaded areas represent periods of missing data. The complete uninterrupted record spanning from spring 2021 to the end of summer 2022 is presented in Appendix A, Fig. A1.

(iii) and being melt-dominated, the pronounced diurnal variability of Q reflects the diurnal reflecting the variability of surface energy balance.

The turbulent-water-flow-induced seismic power, P_s (Fig. 3e and f, yellow) follows the variations in Q throughout the recorded period, increasing at the beginning of the two melt seasons and decreasing towards their end. Such behaviour has been previously observed in other settings and confirms the sensitivity of the selected frequency band to Q (Bartholomäus et al., 2015; Gimbert et al., 2016; Nanni et al., 2020, 2021; Lindner et al., 2020; Labedz et al., 2022a) (Bartholomäus et al., 2022). In contrast, variations in water pressure in the borehole, p , do not follow Q variations in a simple way (Fig. 3g and h, green). At the beginning of the 2021 melt season, p is high, close to the overburden pressure (3.2 MPa). As Q continues to increase, p decreases until it reaches its minimum values (~ 2 MPa) after peak-an abrupt increase in Q marked by (2). p increases again during the winter to reach a value (2.9 MPa) close to overburden pressure (3.2 MPa, (2.9 MPa, Appendix A, Fig. A1d). In the second year of the record, p remains high and increases to the overburden pressure in mid-July 2022 before decreasing again to levels close to its winter value (2.9 MPa) at the end of August 2022.

Similar to p , the force acting on the ploughmeter, F , shows different behaviors between the two melt seasons (Fig. 3g and h, dark red). During the melt season 2021, ~~it~~ F remains fairly stable until August 2021, when it suddenly undergoes variations of large amplitude (~ 150 N maximum amplitude) ~~and~~ high-frequency variations. As the instrument site becomes snow-free around August 15 ~~2021~~ (Fig. 3a), F gradually decreases towards the end of the melt season until it reaches its minimum value in early October 2021 (~ 70 N). At the end of September ~~2021~~ occurs a major precipitation episode (Fig. 3c④) ~~with heavy rainfall going over to snowfall as the temperature dropped~~. F does not react to this event. After this precipitation event, F gradually increases until it stabilizes at ~ 170 N in January 2022. Note that this value is almost twice the value observed before the melt season 2021 (~ 90 N).

The surface velocity of the glacier ~~is related to~~ co-varies with Q -induced changes ~~and~~. We observe several glacier speed-up episodes ~~are identified~~ (i) early in the melt season 2021 ~~when~~ Q increased (Fig. 3c①), (ii) during large rainfall episode (Fig. 3c④), (iii) and during sudden influx of meltwater during the 2022 melt season (Fig. 3d, ⑥, ⑦, ⑧).

To interpret the responses of the subglacial drainage system and glacier dynamics to variations in runoff, Q , we analyze the data using a phase relationship analysis on seasonal (above 20 days, Sec. 4.2), multi-day (four to eight days, Sec. 4.3), and diurnal (six hours to 36 hours, Sec. 4.4) time scales.

325 4.2 Analysis of seasonal variations

To examine the phase relationship between the subglacial variables and Q at the seasonal scale, all time series are low-pass filtered with a cut-off frequency of 20 days. We first describe the results obtained for the melt season 2021 and then for the melt season 2022.

We ~~divided~~ observe four regimes during the melt season ~~2021 into four regimes~~ 2021. At the beginning of the melt season, 330 the preferential drainage axis evolves predominantly by adjusting its capacity, R (constant hydraulic gradient, S , Fig. 4a, 1). As Q increases, ~~this is followed by~~ we observe an adjustment of S at constant R (Fig. 4a, 2). When Q decreases in August 2021, the preferential drainage axis evolves by adjusting R (constant S ; Fig. 4a, 3). At the end of the melt season 2021, ~~the trajectory of~~ P is indicative of we observe an adjustment of S (Fig. 4a, 4). ~~The~~ Simultaneously, $R - Q$ trajectory is parallel to the scaling ~~relationships~~ relationship for channels evolving at steady-state (Fig. 4b) and ~~we observe that the~~ $S - Q$ trajectory 335 follows the steady-state relationship, though not always strictly parallel (Fig. 4c). ~~We observe that the~~ At this time, $p - Q$ relationship is ~~characterised~~ characterized by a clockwise hysteresis (Fig. 4d) indicating that the peak in p precedes the peak in Q . The linear relationship between F and p during the first half of the melt season indicates that the two subglacial variables are anti-correlated (Fig. 4e) and, at the end of the melt season, $F - p$ trajectory shows a counter-clockwise hysteresis, indicating ~~the that~~ the peak in F lags after the peak in p (Fig. 4e).

340 Similar as in 2021, the melt season 2022 can be divided into four regimes, but these phases describe different ~~behaviours~~ behaviors. At the beginning of the melt season, the preferential drainage axis evolves predominantly by adjusting R (Fig. 4f, 1), as observed in 2021. Then, the preferential drainage axis briefly leaves this regime to follow an evolution that is not ~~described by the theoretical relations and that are neither governed by~~ indicative of either constant R ~~nor or~~ constant S (Fig. 4f, 2). ~~This is followed by a return to a regime indicative of predominant~~ After this period, the preferential drainage axis follows a regime




Figure 4. Relationships between two variables at the seasonal scale for the melt seasons 2021 and 2022. Color scales are scale-scaled to the number of days in each melt seasons. ~~Note that the variables R and S have been derived only from July 19 to August 31, 2021 and from May 25 to August 1, 2022 explaining that the color scale is not entirely represented (outside of these periods, runoff is too low to derive R and S , see Section 3.4.1. P , R and S are expressed in terms of relative changes to a reference stage, in our case on June 14, 2021, when $Q_{ref} = 0.2\text{m}^3\text{s}^{-1}$.~~ (a) and (f) Relationships between scaled runoff (Q/Q_{ref}) and scaled turbulent-water-flow-induced seismic power (P/P_{ref}). The x -axis is in logarithmic scale. The superimposed lines show the relations derived by Gimbert et al. (2016) for a constant hydraulic gradient (pink lines, $P \propto Q^{5/4}$) and for a constant hydraulic radius (purple curve, $P \propto Q^{14/3}$). (b) and (g) Relationship between scaled runoff (Q/Q_{ref}) and scaled hydraulic radius (R/R_{ref}). Both x and y -axes are in logarithmic scale. Superimposed lines show the relations of steady-state preferential drainage axis evolution (Nanni et al. (2020), $R \propto Q^{9/22}$). (c) and (h) Relationship between scaled runoff (Q/Q_{ref}) and scaled hydraulic gradient (S/S_{ref}). Both x and y -axes are in logarithmic scale. Superimposed lines show the relations of Nanni et al. (2020) for a preferential drainage axis at steady-state (black lines; $S \propto Q^{-2/11}$) and for a preferential drainage axis evolving as a fixed cross sectional area channel referred to as rigid pipe (red line; $S \propto Q^2$). ~~For the panels a to e and f to h, we interpret our observations as aligning with one of the scenarios detailed in Gimbert et al. (2016); Nanni et al. (2020), where the slope of the hysteresis curve is parallel to the theoretical scaling.~~ (d) and (i) Relationship between normalised-normalized water pressure and normalised-normalized runoff. (e) and (j) Relation between normalised-normalized force and normalised-normalized water pressure. Arrows indicate the direction of time and numbers refer to different periods described in the main text. The numbers do not correspond to the same periods between each panels and are unrelated to the periods identified by the circled numbers in Figure 3.

345 that is predominantly governed by adjustment of S (constant R) for the remaining increase in runoff (Fig. 4f, 3). The runoff decrease phase is not completely captured ~~in the record~~ because the records do not extend to the end of the melt season 2022. ~~However, for~~ For the period covered by data, we observe that the evolution of P is not indicative ~~for neither of either~~ constant R ~~nor or~~ constant S (Fig. 4f, 4). We identify these four phases in the relationship of S and Q . During phases 1 and 3, the behavior is indicative of a preferential drainage axis evolving in equilibrium with Q (Fig. 4g and h, 1 and 3) ~~and while~~ during
350 phases 2 and 4, the ~~observed behavior more behavior~~ behavior closely resembles that of a rigid-pipe (Fig. 4g and h, 2 and 4). ~~As Over the whole melt season and as~~ As Over the whole melt season and as opposed to the melt season 2021, p and Q evolve almost in parallel. At the beginning of the melt

season, p and Q are positively related even though the relationship displays some clockwise hysteresis (Fig. 4i). As for the melt season 2021, F and p are anti-correlated [during the melt season 2022](#) (Fig. 4j).

4.3 Analysis at multi-day scale




Figure 5. Phase relationships between the subglacial variables (S , R , p and F) and runoff (Q) at a multi-day time scale. [Panel \(a\)](#) shows the classification for the melt season 2021, and [panel \(b\)](#) for the melt season 2022. Grey fields refer to periods when data is missing and the vertical [grey lines](#) delineate the events. *Preceding class* and *Lagging class* correspond to clockwise or anti-clockwise hysteresis between the runoff and the observed variable while classes *In-phase class* and *Anti-phase class* correspond to linear relationships. In addition, the time series of velocity ([grey line](#)) and runoff (blue line) are super-imposed on each panel. Circled numbers refers to episodes described in Section 4.1.

355 To understand the relationships on [the](#) time-scales of weather variations, we filtered the time series with a band-pass filter, removing the variations with periods below four days and above eight days. Then, we applied our phase relationship classification scheme to each event (Fig. 5). We investigate the phase relationships between the subglacial variables (the force, F , the water pressure, p , the hydraulic radius, R , the hydraulic gradient, S) and the runoff, Q , for each event during the melt seasons in 2021 (eleven events) and 2022 (eight events, see Section 3.5). Similar phase relationships are observed during both melt

360 seasons between (i) R and Q , and (ii) S and Q . During both melt seasons, R evolves in phase with Q (*In-phase class*, Fig. 5 a and c) ~~and~~, S [evolves in phase with \$Q\$ only](#) at the beginning of the melt season 2021 (*In-phase class*, Fig. 5a). We cannot compare the relationship between S and Q at the beginning of the melt season 2022 ~~as we have removed the events due to the inconsistency due to inconsistencies between the filtered data and the raw data during this period and raw data, necessitating their exclusion from the analysis~~ (Appendix G, Fig. G1). For the remaining part of the season, the [behaviour](#) behavior of S in

365 response to Q is very similar in 2021 and 2022. During the first half of both melt seasons, S is first lagging after Q (*Lagging class*, Fig. 5a and c). During Q important excursions ~~occurring in of~~ August 2021 (Fig. 5b②) and July 2022 (Fig. 5d, ⑦), S precedes Q changes (*Preceding class*, Fig. 5a and b). In contrast to the phase relationships of R and S to Q , F do not show the same time-evolution across both melt seasons ~~but~~, $F - Q$ phase relationship is sensitive to glacier acceleration. During high velocity episodes (Fig. 5b and d, ①, ③, ⑦), F systematically lags behind Q (*Lagging class*, Fig. 5a and c). Conversely, when

370 velocity is low and stable during the melt season 2021 (Fig. 5b and d, from ① to ③), F is anti-correlated with Q (Fig. 5a

and c). We do not have GNSS data in 2022 at this period to compare with the observations in 2021. ~~$p-Q$ phase relationship is contrasted between~~ Between the melt season 2021 and ~~2022.~~ 2022, $p-Q$ relationship is contrasted. On one hand, $p-Q$ phase relationship cannot be easily linked to specific Q regimes or speed-up episodes and shows various responses across the melt season 2021 (*Lagging class*, *In-phase class* and *Anti-phase class*, Fig. 5a and c). On the other hand, p always precedes Q during the melt season 2022 (*Preceding class*, Fig. 5c).




Figure 6. Relationship between the subglacial variables (P/P_{ref} , R/R_{ref} , S/S_{ref}) and runoff (Q/Q_{ref}) during the two melt seasons at a multi-day time scale. The color scale indicates the timing during both melt seasons and is scaled according to the length of each season. (a) and (d) Relationship between scaled runoff (Q/Q_{ref}) and scaled turbulent-water-flow-induced seismic power (P/P_{ref}). The x -axis is in logarithmic scale. The superimposed lines show the relations derived by Gimbert et al. (2016) for a constant hydraulic gradient (pink lines, $P \propto Q^{5/4}$) and for a constant hydraulic radius (purple curve, $P \propto Q^{14/3}$). (b) and (e) Relationship between scaled runoff (Q/Q_{ref}) and scaled hydraulic radius (R/R_{ref}). Both, x and y -axis are in logarithmic scale. Superimposed lines show the relations for a steady-state channel evolution (Nanni et al. (2020) $R \propto Q^{9/22}$). (c) and (f) Relationship between scaled runoff (Q/Q_{ref}) and scaled hydraulic gradient (S/S_{ref}). x and y axes are in logarithmic scale. Superimposed lines show the relations of Nanni et al. (2020) for a channel evolution (black lines; $S \propto Q^{-2/11}$) and for a channel evolving as a rigid-pipe of static cross-section (red line; $S \propto Q^2$).

Figure 6 shows the comparison between the multi-day scale observations of P , R and S and the scaling relationships by Gimbert et al. (2016) and Nanni et al. (2020). At the beginning of the melt season 2021 (July 2021), the slope of the ~~$P-Q$ relation~~ relationship exceeds those of the ~~relations expected for a constant R or constant S~~ before aligning with a slope indicative of scaling relationships. Then, ~~$P-Q$ relationship indicates~~ a preferential drainage axis evolving its capacity at a constant S in the middle of the melt season (August 2021, Fig. 6a). At the end of the melt season, P switches back to the behavior seen at the beginning of the melt season 2021 (Fig. 6a). In general, R evolves with Q similar to what is expected for a steady-state channel (Fig. 6b) whereas S shows a more complex ~~behaviour~~ behavior that is difficult to disentangle (Fig. 6c).

During the melt season 2022, the evolution of P is neither clearly dominated by a constant R or S (Fig. 6d). In general, R increases with Q but at a slope that differs from the one expected for steady-state channels (Fig. 6e). As in 2021, the evolution of S exhibits complex ~~behaviour~~behavior at this scale during the melt season 2022 (Fig. 6f).

4.4 Analysis of diurnal variations




Figure 7. Phase relationships between the subglacial variables (S , R , p and F) and runoff (Q) at a diurnal time scale. (a) Classes from the phase relationship classification per event for the melt season 2021 and (b) for the melt season 2022. Grey fields refer to periods when data is missing and the vertical ~~grey~~gray line delineate the events. In addition, the time series of velocity (~~grey~~gray) and runoff (blue) are super-imposed on each panel. Circled numbers refer to episodes described in Section 4.1.

Symptomatic for glacier meltwater runoff, Q exhibits strong diurnal variations. To examine the glacier response to changes in Q at a diurnal scale, we filtered the time series using a ~~band-pass filter, cutting-off variations beyond the lower and upper limits of bandpass filter removing the variations with periods below~~bandpass filter removing the variations with periods below six hours and above 36 hours ~~, respectively~~ (Fig. 7) ~~respectively~~
 390 ~~to capture the primary diurnal frequency~~. By analyzing variations within this band, we primarily aim at capturing the diurnal variation (around 24 hours) and also ~~to account for some variations and at accounting for some~~ fluctuations around this period. The filtered time series are then subdivided into 95 events in 2021 and 84 events in 2022 (see Section 3.5) and we applied our phase relationship classification scheme.

The phase relationships between the subglacial variables (the force, F , the water pressure, p , the hydraulic radius, R , the
 395 hydraulic gradient, S) and Q on a diurnal time scale are displayed in Figure 7. ~~Results indicate~~We observe that R and S show consistent phase relationships with Q during ~~both melt seasons~~the melt seasons 2021 and 2022, alternating between *Lagging class* and *Preceding class*, and *Preceding class* and *Anti-phase class*, respectively (Fig. 2). However, $p - Q$ and $F - Q$ ~~phase~~
 400 relationships vary across all classes without an easily identifiable pattern (Fig. 7a-c). Except for during short episodes, p and F do not display pronounced diurnal variations (Appendix H, Fig. H1). Therefore, we focus on the analysis of diurnal variations on the responses of P , R and S .

During the 2021 melt season, we observe that R mostly varies in phase with Q (*In-phase class*, Fig. 7a). S is mostly anti-correlated with or precedes Q (*Preceding class* or *Anti-phase class*, Fig. 7a).

During the melt season 2022, we observe a shift from linear responses of R and S (*In-phase class* and *Anti-phase class*, Fig. 7c) towards more hysteretic responses (*Preceding class* and *Lagging class*, Fig. 7c) when Q shows the first significant increase

405 in June (Fig. 7c and d, ⑥). Before this event, R varies with changes in Q ~~before this event~~ (*In-phase class*, Fig. 7c) but after, R lags behind Q (*Lagging class*, Fig. 7c). Similarly, S shifts regimes from being anti-correlated with Q before the ~~episode event~~ ⑥ (*Anti-phase class*, Fig. 7c) to a regime where S precedes Q after this ~~period event~~ (*Preceding class*, Fig. 7c).

5 Discussion

5.1 Interpreting the evolution of subglacial conditions

410 In this study, we have analyzed variations in subglacial hydro-mechanical conditions, i.e., the response of subglacial variables to changes in runoff and we interpret now the observed behavior in terms of subglacial drainage system evolution (Sec. 5.2) and till rheology (Sec. 5.3). We divided the ~~phase~~ relationships observed between different subglacial variables (the force, F , the water pressure, p , the hydraulic gradient, S , and the hydraulic radius, R) and runoff, Q , into four classes (Fig. 2). Here, we first consider the expected responses of R , S (Eqs. 1 and 2) and p to changes in Q for different typical stages of subglacial channel
415 evolution. In addition, we discuss the expected responses of F for ~~different till rheologies~~ the Coulomb-plastic rheology and discuss why the observed relationships may not differ from current theory. Second, we apply this interpretation scheme to the observed behavior separately for each of the considered time scales, before we consolidate these interpretations into a coherent picture of subglacial conditions.

Due to the relatively low bed slope and the long distance between our borehole location and the glacier front, unpressurized
420 drainage is unlikely to persist ~~in these conditions and open flowpaths~~ and open flow paths are expected to close quickly (Nye, 1976). For a drainage axis with a fixed cross sectional area (rigid pipe), an increase in runoff, Q ~~results~~, is expected to result in increasing water pressure, p , that translates to a positive, linear $p - Q$ relationship (*In-phase class*, Fig. 2). In this situation, we expect constant R , unaffected by variations in Q (not classified) and concomitant acceleration events. Since we always measure p at the same location and the glacier terminus is fixed at sea-level, for a spatially homogeneous drainage system, we expect that
425 variations in S are closely related to those of p . However, spatio-temporal complexity in the drainage system downstream of our borehole may lead to incoherent relations between local p and spatially integrated S . According to Röthlisberger's theory for ice-walled channels (Röthlisberger, 1972), the channel cross-section is determined by the two counter-acting processes of ~~melt opening~~ 1) opening by melting due to dissipation of potential energy and 2) creep-closure of the surrounding ice. In steady-state, these two processes balance each other, and a ~~larger runoff would be~~ large runoff is associated with a ~~larger~~
430 large channel, thus ~~requiring lower~~ causing low S and p (Schoof, 2010; Werder et al., 2013). ~~This inverse $p - Q$ relationship is one of the best-known characteristics of ice-walled drainage. Hence, we~~ In this situation, the glacier velocity is expected to remain constant or decrease. We interpret a negative, linear relationship (*Anti-phase class*, Fig. 2) between Q and p (and similar for S and Q), as indicative for steady-state drainage of a preferential drainage axis. ~~This situation also entails that~~ In this configuration, R increases with Q (*In-phase class*, Fig. 2). The evolution of the drainage system in response to Q ~~typically~~
435 is is typically transient between the two end-members described above, ~~with a fixed cross sectional area referred to~~ evolving as rigid pipe on one hand, and a steady-state channel on the other onehand. For transient evolution between these two end-members, we expect a hysteretic behavior in the phase relationships between subglacial variables and Q . Evolution towards

steady-state occurs with some time delay; if variations of the forcing term Q occur faster than this delay, the variations of R lag the variation variations of Q , resulting in a counter-clockwise hysteresis (*Lagging class*, Fig. 2). For such a transient evolution, R is smaller during the rising limb of Q than during the decline, emanating as its decline, resulting in a counter-clockwise $R - Q$ hysteresis (*Lagging class*, Fig. 2). At the same time, the larger R during the decline of Q requires a lower p to drive the flow, resulting in a clockwise hysteresis in the $p - Q$ relationship (*Preceding class*, Fig. 2). As stated above, we expect S to vary in a similar fashion as similarly to p . The response of turbulent water flow induced seismic power P to changes in Q is complex because it integrates the responses of both R and S , hence direct interpretation of the $P - Q$ relations is difficult (Gimbert et al., 2016; Nanni et al., 2020).

We explore two different assumptions for the interpretation of F variations experienced by the ploughmeter ; following Fischer and Clarke (1994) and Kavanaugh and Clarke (2006). For assuming a Coulomb-plastic till (Iverson et al., 1998) ; the flow law for till (Iverson et al., 1998). The till shear strength depends directly on the pore-water pressure, which we assume to co-vary with p at the bottom of the borehole (Iken and Bindshadler, 1986; Hooke et al., 1997; Zoet and Iverson, 2020; Gilbert et al., 2020). This behavior results would result in a negative, linear response of relationship between F to variations in and p (*Anti-phase class*, Fig. 2). For a linear-viscous type till (Alley, 1990; Kamb and Engelhardt, 1991) F increases linearly with sliding velocity. Usually, glacier sliding laws acknowledge a water pressure control on sliding speed (Iken and Bindshadler, 1986; Hooke et al., 1997) which would result in a positive, linear relationship between F and p (*In-phase class*, Fig. 2). In that case, a high water pressure would weaken the till, facilitating its deformation (e.g., Weertman, 1957; Lliboutry, 1968). However, more complex $F - p$ relationships can emanate as well, accounting for that the relationship between occur, arising from the non-linearity of p and sliding speed may be non-linear (Alley, 1989; Boulton and Hindmarsh, 1987). In general, interpreting a direct relation between F and Q is complicated by the circumstance that variations in sliding speed relationship (Alley, 1989; Boulton and Hindmarsh, 1987) and from the degree of ice-till coupling (Iverson et al., 2007). p and F can also be influenced by potential distal forcing, e.g. longitudinal stress-transfer, complicating their interpretation. Additionally, as p (and S) also depend on depends both on Q and the efficiency of the drainage system to evolve to cope with runoff input, hence depend both on subglacial drainage system, $F - Q$ and P relationships are difficult to interpret.

Our interpretation scheme described here lets us expect two to three a limited number of options for the classification of phase relations between each subglacial variable and Q . For $p - Q$ and $S - Q$, we expect behaviors according to *Preceding class*, *In-phase class* or *Anti-phase class*; we expect $R - Q$ to display either *Lagging class* or *In-phase class* behavior; $F - p$ is expected to fall either in *In-phase class* or *Anti-phase class*, direct $F - Q$ relations are not easily comprehended interpretable. In practice, we observe that some events are classified outside the expected range (Figs. 5 and 7). The occurrence of such behavior may be attributed partly to artifacts introduced by the spectral filtering applied to the time series for the analysis. Although we manually checked the consistency between the unfiltered and filtered signals to remove the most apparent differences (see Section 3.5), some inconsistencies may still remain. Small shifts in timing of peaks may be amplified by normalizing the event time axis and hence lead to mis-classification of some events. In addition, the definition of the four classes is motivated by noticing that phase relations may be linearly positive or negative or exhibit some transitory stage (preceding or lagging). To account for uncertainties symptomatic for observations of natural systems, we allow some deviation from strictly linear behavior and

~~accept $RSS \leq 2$ still representing linear behavior. The choice of this threshold is motivated from visual impression of clustering of phase relations.~~

475 5.2 Subglacial drainage system evolution

The two observed melt seasons considerably differ in terms of duration, variability and intensity (Fig. 3). Whereas in 2021, melting occurs over a relatively short period and yields low levels of water supply, ~~in 2022,~~ the melt season 2022 lasts longer and is characterized by higher temperatures ~~and thus yields~~ yielding to higher water supply rates. This difference provides the opportunity to study the evolution of the subglacial drainage system in response to very different forcing.

480 5.2.1 The melt season 2021: short duration and low intensity

The melt season 2021 is short (67 days) and marked by runoff usually lower than $20 \text{ m}^3 \text{ s}^{-1}$ (Fig. 3a). Applying our interpretation scheme to the observed responses of the water pressure, p , the turbulent-water-flow-induced seismic power, P , the hydraulic radius, R , and the hydraulic gradient, S , yields a largely consistent picture of a subglacial drainage system that fully adapts to seasonal runoff variations: ~~$p - Q$ relationship~~ relationship exhibits clockwise hysteresis, indicative of system capacity growing with runoff (Fig. 4d); ~~$R - Q$ variations are positively-linearly related (Fig. 4b); the $S - Q$ relationship covers only the declining phase of Q and shows an increase of S during this decline, consistent with the above interpretation (Fig. 4c).~~ ~~As mentioned above, due to the composite nature of turbulent water flow induced seismic power, the $P - Q$ relationship is more difficult to interpret, but comparison to idealised behavior suggests that system adjustment is dominated by different mechanisms during different stages (Fig. 4a).~~ Although all records draw the picture of a system adjusting to Q variations, there is a noteworthy difference in the interpretations of borehole measurements and those of cryoseismology records. While long-term variations of R and S suggest that the system capacity ~~obtains reaches~~ reaches an equilibrium with Q ; ~~the~~ The variations of p ~~measured in the borehole,~~ indicate a transient evolution in response to changes in Q . ~~$p - Q$ exhibits clockwise hysteresis,~~ indicative of system capacity growing with runoff ~~also this in agreement with expected behavior (Fig. 4d).~~ We explain this apparent disagreement by the different spatial scales of sensing of the different instruments: while the geophone records are likely dominated by seismicity generated by turbulent flow in the channelized system (Nanni et al., 2021), the pressure record is representative of the local conditions at the borehole location, which is more likely to sense the local distributed system (Rada and Schoof, 2018)

The theoretical timescale of channel adjustment is usually longer (several days to weeks) than typical variations in Q (hours) (Röthlisberger, 1972). At short time scales, drainage pathways are ~~then thus~~ either overwhelmed when Q increases or partially filled when Q decreases, which results in a response similar to that of a rigid pipe (fixed cross-sectional area channel) rather than that of a steady-state channel (variable cross-section determined by the balance between melt opening and creep-closure to cope with runoff variations). We therefore expect a predominance of *In-phase class* for the $p - Q$ and $S - Q$ relationships over multi-day and diurnal time scales (Fig. 5a and 7a). The multi-day classifications of $p - Q$ and $S - Q$ relationships mainly support this view by displaying *Lagging class* and *In-phase class* behaviors (Fig. 5a), ~~however;~~ however, on diurnal time scales (Fig. 7a), the picture is less clear ~~. On these shorter time scales, we expect only minor variations of R , lagging those of Q~~

(Lagging-class). However, we observe mainly *In-phase-class* behavior at both multi-day and diurnal time scales (Fig. 5a and 7a), suggesting that geometrical adjustment of the drainage system takes place already over short time scales; however this implies that the observed behaviors of p and S are caused by likely due to changes in hydraulic roughness since R already has adjusted (Nanni et al., 2020).

510 ~~Towards the end~~ We observe that the melt-induced runoff increase at the beginning of the melt season 2021, and a major rainfall event at its end (Fig. 3, ① and ③) ~~left-leave~~ a clear impact on p at the multi-day time scale (Fig. 3d, Fig. 5a) ~~and coincided~~, which coincides with a considerable glacier acceleration. During ~~this event, the these events~~, $p - Q$ relationship indicates that the drainage system evolves in a transient manner (*Lagging class* and *In-phase class*, Fig. 5a). ~~Prior to this event~~ We suggest that at the beginning of the melt season, the drainage system is not yet developed when the
515 melt begins leading the immediate increase in p . At the end of the melt season, Q was at low levels for a period of about 10 days (Fig. 3d), and presumably, the capacity of the drainage system had decreased, when it suddenly became overwhelmed by the arrival of new water volumes. This results in a sharp increase of p , provoking a short-term acceleration of the glacier. Similar late season events have also been reported in other studies (~~Andrews et al., 2014; Rada and Schoof, 2018~~)
(Andrews et al., 2014; Rada and Schoof, 2018; Nanni et al., 2023). Even though p supports the interpretation of the velocity
520 data in this context, $p - Q$ relationship displays similar patterns throughout the 2021 melt season, which do not lead to acceleration events.

5.2.2 The melt season 2022: long duration and high intensity

In contrast to the melt season 2021, the melt season 2022 is long (at least, 83 days since our records end before the melt season ceases) and contains frequent and large excursions of runoff Q above $20 \text{ m}^3 \text{ s}^{-1}$ (Fig. 3a) leading to a different evolution of the
525 subglacial drainage system ~~compared to the preceding year. The~~. $R - Q$ relationship is linearly positive, indicating ~~drainage system size evolving that the drainage system capacity evolves~~ at equilibrium with Q (Fig. 4g). ~~The~~ $S - Q$ relationship follows a trajectory that first is typical of a steady-state channel (Fig. 4h, 1) before ~~bending over shifting~~ to a positive slope similar to that expected for a rigid pipe (Fig. 4h, 2, 3 and 4). Such behavior is typical for a system that is continuously overwhelmed since the subglacial preferential drainage axes cannot adapt fast enough to increasing Q . This period (Fig. 4h, 2, 3 and 4) coincides
530 with a noticeable glacier acceleration event aligning with a subglacial drainage system out of equilibrium. This acceleration indicates a direct link between glacier velocity and state of the subglacial drainage system as measured locally and possibly in isolated bed regions (Figs. 3h ⑦, 4i). $P - Q$ relationship in 2022 shows similar behavior as in 2021, if we account for that the 2022 record does not cover the falling limb of Q (Fig. 4f). The view of a continuously overwhelmed drainage system is further supported by the generally positive slope of the $p - Q$ relationship ~~which that~~ has a considerably smaller clockwise hysteresis
535 compared to the preceding year (Fig. 4i).

Over shorter time scales, the classifications of $R - Q$ and $S - Q$ exhibit similar behavior as in 2021 (Fig. 5b and 7b), indicative for system adjustment also on multi-day and diurnal times. However, the *Lagging class* behavior of $S - Q$ in the beginning of the melt season visible in Figure 5b, remains difficult to explain. On a diurnal time scale, the dominance of *Anti-phase class* in the $S - Q$ ~~relation-relationship~~ before July 2022 suggests that preferential drainage axis evolution is in equilibrium with Q

540 (Fig. 7b). ~~The data then displays a switch~~ We observe a switch in the data to mainly *Preceding class* behavior, which in turn is indicative of a transient evolution, typical for a drainage system that cannot adapt fast enough to the runoff variations. $R - Q$ relationship exhibits a similar switch (from *In-phase class* to class II 7b) with implications consistent ~~to~~ with the interpretation above. The *Preceding class* behavior of $p - Q$ on multi-day scales further supports the view of geometric adjustment at this scale. ~~The same analysis on the~~ However, as opposed to the acceleration events during the melt season 2021 (Fig. 3h ① and
545 ③), there is no straightforward explanation between $p - Q$ relationship and the acceleration event observed during the melt season 2022 (Fig. 3h ⑦). We suggest that here, the glacier acceleration is controlled by processes not happening locally and thus not recorded locally by the water pressure sensor. The diurnal scale (Fig. 7b) reveals that there is more diurnal variations in p during the melt season 2022, compared to 2021 (more events are classified). However, the diurnal analysis renders a blurry picture since all classes occur and no clear pattern can be depicted (Fig. 3d).

550 5.2.3 Ambiguous interpretation from borehole and cryoseismic records

In the previous section, we interpreted the multi-variable record in terms of drainage system evolution in response to runoff (~~summarised~~ summarized in Figure 9). We note that sometimes, interpretations derived from different records are ambiguous. For instance, during the melt season 2021 ~~,~~ on a seasonal scale, the relationships between the cryoseismic record (P) with Q , and derived variables (R and S) ~~and~~ with Q , yield a picture of a subglacial drainage system in equilibrium with Q . In contrast,
555 ~~the~~ $p - Q$ relationship based on the borehole record is symptomatic of a transient evolution where geometric adjustments lag variations in Q . Another example of the ambiguity is found in the analysis of diurnal variations in 2022 (Fig. 7b) where the cryoseismic records indicate a switch from an equilibrium to a transient evolution coinciding with a major increase in Q (Fig. 7d ⑥). The corresponding classification of ~~the~~ $p - Q$ ~~relation~~ relationship is less conclusive about a similar switch and exhibits variations over all classes with no clearly recognizable pattern. In this section, we discuss potential sources for these
560 inconsistencies and how these may be resolved.

~~Percieving the drainage system as a macroporous sheet goes along with the spatial heterogeneity of the subglacial drainage system, with parts of the glacier base being located in areas that are well connected to the main water drainage, whereas other parts only indirectly communicate with that system (Murray and Clarke, 1995; Rada and Schoof, 2018), for instance through a porous medium of lower permeability. Such a situation would apply when interpreting records from sensors that~~
565 ~~are physically installed at different locations along the base (e.g., in different boreholes), but also, when the view field of the individual sensors differs such that the recorded signals have different characteristics of noise and signal amplitude.~~ In our case, P integrates the seismicity in the 3-10 Hz frequency band in which seismic wavelengths are ~~on~~ of the order of 150-500m ~~(for typical surface wave velocity on the order of 1500 m sec⁻¹ Köhler et al., 2012; Gimbert et al., 2021b)~~ (for typical surface wave velocity
. Accordingly, the cryoseismologically derived variables P , R and S are sensitive to an area of ~ 1 km² around the geophone
570 location, ~~and therefore, they will always represent the hydraulically most active part within this area (Nanni et al., 2021).~~

. In addition, since the seismicity in this frequency range is mainly generated by turbulent water flow, we expect that the observed signal is mainly indicative of large channels (Nanni et al., 2021). In contrast, water pressure p is measured at the bottom of the borehole, ~~the precise dimensions of which at the base of the glacier are unknown, but~~ estimated to be ~ 20 cm in

diameter [at the base of the glacier](#). Depending on the hydraulic connection of the borehole and the ice-till coupling, p may be representative for about 1 m² in case of hydraulic isolation, or for a several orders of magnitude larger area in case of a direct connection to a preferential drainage axis (Murray and Clarke, 1995; Mair et al., 2001, 2003). [As a consequence, interpreting the cause of glacier acceleration through cryoseismology or local basal water pressure might not be suitable. On one hand, cryoseimology is sensitive to the most efficient part of the subglacial drainage system that are often not the primary controllers of glacier velocity. On the other hand, glacier velocity results from basal conditions over a larger area \(scale of ice thickness\) than what is locally sampled by the measured water pressure \$p\$.](#)

In addition, the hot-water drilling operation might have disturbed the subglacial environment (excavation of fines, volume of water pushed to the bed), influencing the water pressure observation. However, the volume of water injected through hydraulic connection of the borehole to the bed is limited (~ 0.5 m³ for the observed 30 m drop of water column at connection) and the borehole has been drilled beginning of May 2021. In the absence of surface melting before late June 2021, it seems unlikely that a potential initial connection could be maintained. Geometrically controlled patterns of channelization on a hard bed may be persistent, but a soft sediment bed provides less geometrical controls on the spatial patterns and year-to-year variability of channel location within a few meters seems plausible. In addition, several studies report that water pressure records displayed regime changes and suggest that these may reflect reorganization of the drainage system (Gordon et al., 1998; Kavanaugh and Clarke, 2000; Schuler et al., 2002; Andrews et al., 2014; Rada and Schoof, 2018). Resuming the idea of a spatially heterogeneous and discontinuous drainage system aids resolving the apparent discrepancies derived from the different records: if the active drainage system within 1 km² around our instrument site is in equilibrium with Q , the seismological record would reveal this effect, whereas the borehole record may reveal a different interpretation if the borehole itself is located in a less-well connected or even isolated part of the glacier bed. The apparent discrepancy derived from different records therefore supports the comprehension of the drainage system as a discontinuous, spatially heterogeneous sheet. Although the minor diurnal variability [and high values](#) of p during the melt season 2022 may be interpreted as symptomatic for hydraulic isolation ([Rada and Schoof, 2018; Rada Giacaman and Schoof, 2022](#)), p displays seasonal variability responding to Q . This observation suggests that differently connected parts of the glacier bed hydraulically communicate, at least when the capacity of the drainage system is overwhelmed in times of high water supply. Sufficiently high water pressure in connected bed areas can cause the expansion of the connected drainage system (Murray and Clarke, 1995). During such episodes, high water pressures would occur over a large part of the glacier bed, possibly promoting glacier sliding. Indeed, during the melt season 2022, several episodes of glacier acceleration are recorded during episodes of high Q that coincided with p close to overburden pressure (Fig. 3 b ⑥, ⑦), even though the borehole was not well connected to the main drainage system. Conversely, when the connected areas of the bed operate at low water pressure, areas of the bed adjacent to the preferential drainage axes are hydraulically isolated by stress-bridging (Lappégard et al., 2006), resulting in areas of the glacier bed switching back and forth between connected and isolated (Murray and Clarke, 1995).

5.3 Till changes and glacier dynamics

In the Section 5.1 above, we have proposed an interpretation scheme based on the phase relationship between the force experienced by the ploughmeter, F , and the pore water pressure of the till in the till layer, here taken as adequately represented by p . In this section, we explore subglacial processes that may explain the complex relationship observed between the subglacial hydrology (represented by the subglacial water pressure) and the mechanical properties of the till (represented by the force experienced by the ploughmeter) assuming a Coulomb-plastic rheology. We recall that for such a constitutive flow law for the till, a negative $p - F$ relationship is expected (*In-phase-Anti-phase class*).




Figure 8. Comparison between the variations of water pressure p and ploughmeter force F (a) at the seasonal, and (b) to (d) multi-day time scales. (a) Relationship between p and F at the seasonal time scale. F and p are normalised-normalized (min-max normalisationnormalization) to be comparable. Blue-yellow and yellow-purple color scales indicate time during melt seasons and winter, respectively. (b) Evolutions of F (red) and p (blue) for an event indicative of apparent viscous behavior (assuming that p is positively related to basal motion). Similar behaviour-behavior has been observed also during other periods (e.g., from July 13, 2021 to July 22, 2021, and August 8, 2021 to August 17, 2021). The time of the event is normalised-normalized over 50 time steps. (c) Evolutions of F (red) and p (blue) for an event indicative of Coulomb-plastic behavior. Similar behaviour-behavior has been observed also during other periods (e.g., from July 22, 2021 to July 31, 2021, from July 7, 2021 to August 8, 2021, and from July 11, 2022 to July 23, 2022). The time of the event is normalised-normalized over 50 time steps. (d) Evolutions of F (red) and p (blue) that remain unclear as the interpretation scheme does not provide a clear indication of till rheology. Similar behaviour-behavior has been observed also during other periods (e.g., from September 15, 2021 to September 27, 2021, and from May 28, 2022 to June 6, 2022). The time of the event is normalised-normalized over 50 time steps. Simultaneous multi-day variations of F and p can be assessed only for five and two events during the melt seasons 2021 and 2022, respectively. All panels show normalized variations of F and p .

Over-At a seasonal scale during the melt seasons, the, $p - F$ relation-relationship displays a generally negative slope (Figs. 4e and j and 8a). For a Coulomb-plastic material, an increase in pore water pressure results in a decrease in shear strength, which in turn would cause a decrease in F (Fig. 9). Furthermore, the observed anti-correlation between p and F is in good agreement with the modeling results of Kavanaugh and Clarke (2006) for a Coulomb-plastic material, an interpretation that is in line with the findings of Fischer and Clarke (1994); Fischer et al. (1998, 2001) at Trapridge glacier, Storglaciären and Unteraargletscher,

respectively. However, during winter 2021/22, the $p - F$ relationship exhibits a positive slope (Fig. 8a) which is unexpected for Coulomb-plastic rheology. Apparent-viscous behavior entails a velocity dependency of basal resistance, resulting in a positive
620 $p - F$ relationship. However, we do not observe glacier acceleration during the same period in winter 2021/22.

Over shorter time scales, time series of p and F exhibit complex behavior: correlation (Fig. 8b), anti-correlation (Fig. 8c), and lagging after each configuration ~~other~~-occur (Fig. 8d). Applying our interpretation scheme suggests that during episodes similar to those displayed in Figure 8c, the inverse $p - F$ correlation results from weakening of the sediment at times of high water pressure due to reduced effective pressure and vice versa, as expected for a near-Coulomb rheology (Fig. 9). However, a
625 positive relationship between p and F as pictured in ~~Figures 8b and d~~ Figure 8 does not agree with Coulomb-plastic rheology. ~~The illustrated episodes apparently do not coincide with periods of high surface velocity.~~ Similar $p - F$ correlations have been observed previously (e.g., Murray and Porter, 2001; Rousselot and Fischer, 2007; Thomason and Iverson, 2008) but not extensively discussed. A range of mechanisms have been proposed to explain such behavior, ~~such as e.g.~~, the sediments loaded towards their yield point (e.g., Murray and Porter, 2001), the state of the mechanical coupling between the ice and the till and
630 its influence on pore-pressure variations (Iverson et al., 1995; Fischer and Clarke, 1997; Boulton et al., 2001; Mair et al., 2003; Iverson, 2010), the varying ~~mobilisation~~ mobilization of the till at depth (e.g., Iverson et al., 1998; Tulaczyk, 1999; Tulaczyk et al., 2001; Truffer et al., 2000; Truffer, 2004). However, a direct explanation on how these mechanisms would explain the correlation between F and p is not straightforward. We further point out that the ~~attitude~~ vertical position of the ploughmeter relative to the till may have changed, ~~for instance e.g.~~ through changes in ~~tilt or vertical position~~ ploughmeter tilt, but these
635 effects cannot be disentangled from till behavior without further ~~accompanying~~ measurements. Such measurements will be subject for future ploughmeter deployments.

6 Conclusions

In this study, we adopt a multi-method, multi-scale analysis to examine the responses of the subglacial environment of the Kongsvegen glacier, Svalbard, to changes in runoff. ~~We Here, we~~ synthesize the broad spectrum of ~~sediment deformation and~~
640 ~~subglacial drainage behaviors~~ relationships between the observed data series in four classes and interpret them in terms of drainage system evolution and till rheology (Fig. 9).

Our data cover two contrasting melt seasons: during the short and less intensive melt season 2021, we ~~conjecture~~ suggest that our borehole intersected a well-connected part of the subglacial drainage system, whereas in the longer and intensive melt season 2022, the borehole recorded characteristics of a poorly connected subdomain of the glacier bed (Fig. 9). Seismologi-
645 cal records indicate the existence of an efficient drainage system in both periods. Considering the different footprints of our sensors (e.g., meter-scale sensitivity for the ploughmeter and the water pressure sensor and hundreds of meter-scale sensitivity for the seismic investigation due to the selected frequencies), complementary information can be obtained that allows us to propose a consistent picture of the subglacial environment. ~~We therefore suggest the co-existence of efficient and inefficient~~ drainage system components together with isolated patches, rendering the perception ~~The apparent disagreement between~~
650 seismic data and borehole-sensed pressure records can be explained by the spatial heterogeneity of the subglacial drainage

system, supporting the concept of a discontinuous, spatially heterogeneous drainage system (Fig. 9, Rada Giacaman and Schoof, 2023) (Fig. 9, Andrews et al., 2014; Hoffman et al., 2016; Nanni et al., 2021; Rada Giacaman and Schoof, 2023).

The relationship between the force experienced by the ploughmeter and the water pressure reveals complex till rheology behavior. As expected, the till behaves mainly mostly as a Coulomb-plastic material ~~but episodically, the $p-F$ relation is indicative of~~ deviating behavior. Such behavior may be caused by different mechanisms, e.g. till loading towards its yield strength, or ~~variations in the mobilization of the till at depth, but we cannot exclude possible effects of changes in instrument-till coupling that may cause similar behavior. In future, additional measurements of instrument attitude.~~ In the future, monitoring instrument vertical position could help disentangling instrument behavior from till behavior and assessments of changes in till properties using seismic noise interferometry (Zhan, 2019) over a large area could complement the local ploughmeter record.

~~In this study, we~~ By constraining the spatial and temporal extents of our investigation, this study investigated local subglacial hydro-mechanical conditions rather than explored the multi-annual and likely large scale processes responsible of the surge build-up. We show the importance of a multi-sensor multi-scale approach to study the complex variations of hydro-mechanical conditions in response to runoff changes. Such an approach may also be beneficial for studying the dynamics of other transient geological systems that are characterized by build-up and evolution of internal states. These states may reach critical thresholds ~~causing instabilities over different scales from short-lived events to large-scale by destabilization as exemplified in many~~ geohazards such as glacial surges or collapses, volcanic eruptions, landslides, or earthquakes.




Figure 9. Sketch of the adjustment of hydro-mechanical conditions below Kongsvegen glacier to variations in runoff over the period from June 2021 to August 2022. (a) Hydraulic quantities, i.e. water pressure (p), hydraulic gradient (S), and hydraulic radius (R), used to characterize the evolution of the subglacial drainage system during the short and low intensity melt season of 2021, and the long and high intensity melt season of 2022. Our findings indicate that during the 2021 season, the subglacial drainage system adapted to runoff changes in steady-state, leading to an increase in its capacity over time. However, during the 2022 season, we observed a transient evolution of the drainage system in response to the continued and high-intensity input of runoff. As a result, the drainage capacity of the main drainage system was exceeded, causing water to leak into poorly connected areas of the bed increasing the water pressure, thereby triggering speed-up events. (b) The mechanical quantity, i.e., force (F), was used to examine the rheological behavior of the till. The till rheology behaved mainly as a Coulomb-plastic material (anti-correlation between p and F), but episodically showed deviating behavior (correlation between p and F), the underlying mechanisms for which remain unclear.

Code and data availability. The data are uploaded in Zenodo (<https://doi.org/10.5281/zenodo.7648444>, Bouchayer (2023a)) and the code to reproduce the figures in the manuscript can be found on GitHub (https://github.com/Colinebouch/mammamia_alldata_processing, Bouchayer (2023b)).

670 *Author contributions.* CB, UN and TVS designed the concept of the study. CB developed the code, produced all tables and figures and wrote the initial draft. UN and TVS provided help and ideas in all phases of the study and wrote parts of the manuscript. FR provided ideas and feedbacks that lead to the finalization of the manuscript. TVS provided all the organizational and technical support. JK and TVS provided the logistical support to conduct the field campaigns. JH built the ploughmeter. CB, UN, JH, TVS, PML and JK conducted fieldwork. TVS and CB processed the ploughmeter and water pressure data. UN and CB processed the seismic data. PML processed the GNSS velocity data. JK
675 provided the long term velocity data at KNG6. LS contributed with CARRA and AROME-ARCTIC forcing, and simulated surface runoff. All authors contributed to the final manuscript with input and suggestions.

Competing interests. The authors declare no conflict of interest.

Acknowledgements. This project has received support from the Research Council of Norway through the projects MAMMAMIA (grant no. 301837) and SLIDE (no. 337228) and from the Faculty of Mathematics and Natural Sciences at the University of Oslo through the
680 strategic research initiative EarthFlows. TVS and FR acknowledge support from the project FricFrac funded by the Center for Advanced Study (CAS) at the Norwegian Academy of Science and Letters during academic year 2023-2024. We are grateful to the Governor of Svalbard for permitting fieldwork at Kongsvegen. The energetic help of Wenxue Cao, Ceslav Czyk, Basile De Fleurian, Jean-Charles Gallet, Adrien Gilbert, Stephen Hudson, Simon Filhol, Urs Fischer, Catherine Larose, and Ashley Morris during the field campaigns is greatly acknowledged. UN acknowledges Andreas Köhler for support in the seismic data analysis. We thank the [previous](#) editor, Prof. Elisa Mantelli,
685 [the current editor, Prof. Nanna B. Karslon](#) and two anonymous reviewers for thorough reviews that greatly improved an earlier version of the manuscript.

References

- Alley, R.: Water-pressure coupling of sliding and bed deformation: II. Velocity-depth profiles, *Journal of Glaciology*, 35, 119–129, 1989.
- Alley, R. B.: Multiple steady states in ice-water-till systems, *Annals of Glaciology*, 14, 1–5, 1990.
- 690 Andrews, L. C., Catania, G. A., Hoffman, M. J., Gulley, J. D., Lüthi, M. P., Ryser, C., Hawley, R. L., and Neumann, T. A.: Direct observations of evolving subglacial drainage beneath the Greenland Ice Sheet, *Nature*, 514, 80–83, 2014.
- Bælum, K. and Benn, D.: Thermal structure and drainage system of a small valley glacier (Tellbreen, Svalbard), investigated by ground penetrating radar, *The Cryosphere*, 5, 139–149, 2011.
- Bartholomaeus, T. C., Amundson, J. M., Walter, J. I., O’Neel, S., West, M. E., and Larsen, C. F.: Subglacial discharge at tidewater glaciers revealed by seismic tremor, *Geophysical research letters*, 42, 6391–6398, 2015.
- 695 Benn, D., Gulley, J., Luckman, A., Adamek, A., and Glowacki, P. S.: Englacial drainage systems formed by hydrologically driven crevasse propagation, *Journal of Glaciology*, 55, 513–523, 2009.
- Benn, D., Fowler, A. C., Hewitt, I., and Sevestre, H.: A general theory of glacier surges, *Journal of Glaciology*, 65, 701–716, 2019.
- Beyreuther, M., Barsch, R., Krischer, L., Megies, T., Behr, Y., and Wassermann, J.: ObsPy: A Python toolbox for seismology, *Seismological Research Letters*, 81, 530–533, 2010.
- 700 Björnsson, H., Gjessing, Y., Hamran, S.-E., Hagen, J. O., Liestøl, O., Pálsson, F., and Erlingsson, B.: The thermal regime of sub-polar glaciers mapped by multi-frequency radio-echo sounding, *Journal of Glaciology*, 42, 23–32, 1996.
- Bouchayer, C.: Dataset at a 3h resolution used in the paper ‘The MAMMAMIA project: A multi-scale multi- method approach to understand runoff-induced changes in the subglacial environment and consequences for surge dynamic in Kongsvegen glacier, Svalbard’, <https://doi.org/10.5281/zenodo.7648444>, 2023a.
- 705 Bouchayer, C.: GitHub repository to process the data, published in Zenodo, https://github.com/Colinebouch/mammamia_alldata_processing/commits/v1.0, <https://doi.org/10.5281/zenodo.7648470>, last update: 17/02/2022, 2023b.
- Boulton, G. and Hindmarsh, R.: Sediment deformation beneath glaciers: rheology and geological consequences, *Journal of Geophysical Research: Solid Earth*, 92, 9059–9082, 1987.
- 710 Boulton, G., Dobbie, K., and Zatsepin, S.: Sediment deformation beneath glaciers and its coupling to the subglacial hydraulic system, *Quaternary International*, 86, 3–28, 2001.
- Clarke, G. K.: Subglacial processes, *Annu. Rev. Earth Planet. Sci.*, 33, 247–276, 2005.
- Clyne, E., Alley, R. B., Vore, M., Gräff, D., Anandakrishnan, S., Walter, F., and Sergeant, A.: Glacial hydraulic tremor on Rhonegletscher, Switzerland, *Journal of Glaciology*, 69, 370–380, 2023.
- 715 Creyts, T. T. and Schoof, C. G.: Drainage through subglacial water sheets, *Journal of Geophysical Research: Earth Surface*, 114, 2009.
- Damsgaard, A., Egholm, D. L., Piotrowski, J. A., Tulaczyk, S., Larsen, N. K., and Tylmann, K.: Discrete element modeling of subglacial sediment deformation, *Journal of Geophysical Research: Earth Surface*, 118, 2230–2242, 2013.
- Damsgaard, A., Egholm, D. L., Beem, L. H., Tulaczyk, S., Larsen, N. K., Piotrowski, J. A., and Siegfried, M. R.: Ice flow dynamics forced by water pressure variations in subglacial granular beds, *Geophysical Research Letters*, 43, 12–165, 2016.
- 720 Damsgaard, A., Goren, L., and Suckale, J.: Water pressure fluctuations control variability in sediment flux and slip dynamics beneath glaciers and ice streams, *Communications Earth & Environment*, 1, 66, 2020.
- De Fleurian, B., Gagliardini, O., Zwinger, T., Durand, G., Le Meur, E., Mair, D., and Råback, P.: A double continuum hydrological model for glacier applications, *The Cryosphere*, 8, 137–153, 2014.

- Downs, J. Z., Johnson, J. V., Harper, J. T., Meierbachtol, T., and Werder, M. A.: Dynamic hydraulic conductivity reconciles mismatch between
725 modeled and observed winter subglacial water pressure, *Journal of Geophysical Research: Earth Surface*, 123, 818–836, 2018.
- Doyle, S. H., Hubbard, B., Christoffersen, P., Young, T. J., Hofstede, C., Bougamont, M., Box, J., and Hubbard, A.: Physical conditions of
fast glacier flow: 1. Measurements from boreholes drilled to the bed of Store Glacier, West Greenland, *Journal of Geophysical Research:
Earth Surface*, 123, 324–348, 2018.
- Engelhardt, H. and Kamb, B.: Basal hydraulic system of a West Antarctic ice stream: constraints from borehole observations, *Journal of
730 Glaciology*, 43, 207–230, 1997.
- Fischer, U. H. and Clarke, G. K.: Ploughing of subglacial sediment, *Journal of Glaciology*, 40, 97–106, 1994.
- Fischer, U. H. and Clarke, G. K.: Stick–slip sliding behaviour at the base of a glacier, *Annals of Glaciology*, 24, 390–396, 1997.
- Fischer, U. H., Iverson, N. R., Hanson, B., Hooke, R. L., and Jansson, P.: Estimation of hydraulic properties of subglacial till from plough-
meter measurements, *Journal of Glaciology*, 44, 517–522, 1998.
- 735 Fischer, U. H., Clarke, G. K., and Blatter, H.: Evidence for temporally varying “sticky spots” at the base of Trapridge Glacier, Yukon Territory,
Canada, *Journal of Glaciology*, 45, 352–360, 1999.
- Fischer, U. H., Porter, P. R., Schuler, T., Evans, A. J., and Gudmundsson, G. H.: Hydraulic and mechanical properties of glacial sediments
beneath Unteraargletscher, Switzerland: implications for glacier basal motion, *Hydrological Processes*, 15, 3525–3540, 2001.
- Flowers, G. E.: Modelling water flow under glaciers and ice sheets, *Proceedings of the Royal Society A: Mathematical, Physical and Engi-
740 neering Sciences*, 471, 20140907, 2015.
- Flowers, G. E. and Clarke, G. K.: A multicomponent coupled model of glacier hydrology 1. Theory and synthetic examples, *Journal of
Geophysical Research: Solid Earth*, 107, ECV–9, 2002a.
- Flowers, G. E. and Clarke, G. K.: A multicomponent coupled model of glacier hydrology 2. Application to Trapridge Glacier, Yukon, Canada,
Journal of Geophysical Research: Solid Earth, 107, ECV–10, 2002b.
- 745 Fudge, T., Humphrey, N. F., Harper, J. T., and Pfeffer, W. T.: Diurnal fluctuations in borehole water levels: configuration of the drainage
system beneath Bench Glacier, Alaska, USA, *Journal of Glaciology*, 54, 297–306, 2008.
- Gilbert, A., Gagliardini, O., Vincent, C., and Wagnon, P.: A 3-D thermal regime model suitable for cold accumulation zones of polythermal
mountain glaciers, *Journal of Geophysical Research: Earth Surface*, 119, 1876–1893, 2014.
- Gilbert, A., Gimbert, F., Thøgersen, K., Schuler, T. V., and Kääb, A.: A Consistent Framework for Coupling Basal Friction With Subglacial
750 Hydrology on Hard-Bedded Glaciers, *Geophysical Research Letters*, 49, e2021GL097507, 2022.
- Gillet-Chaulet, F., Durand, G., Gagliardini, O., Mosbeux, C., Mougnot, J., Rémy, F., and Ritz, C.: Assimilation of surface velocities acquired
between 1996 and 2010 to constrain the form of the basal friction law under Pine Island Glacier, *Geophysical Research Letters*, 43, 10–311,
2016.
- Gimbert, F., Tsai, V. C., Amundson, J. M., Bartholomäus, T. C., and Walter, J. I.: Subseasonal changes observed in subglacial channel
755 pressure, size, and sediment transport, *Geophysical Research Letters*, 43, 3786–3794, 2016.
- Gimbert, F., Gilbert, A., Gagliardini, O., Vincent, C., and Moreau, L.: Do existing theories explain seasonal to multi-decadal changes in
glacier basal sliding speed?, *Geophysical Research Letters*, 48, e2021GL092858, 2021a.
- Gimbert, F., Nanni, U., Roux, P., Helmstetter, A., Garambois, S., Lecointre, A., Walpersdorf, A., Jourdain, B., Langlais, M., Laarman, O.,
et al.: A multi-physics experiment with a temporary dense seismic array on the Argentière glacier, French Alps: The RESOLVE project,
760 *Seismological Society of America*, 92, 1185–1201, 2021b.

- Goldberg, D., Schoof, C., and Sergienko, O.: Supporting Material for Stick-slip motion of an Antarctic Ice Stream: the effects of viscoelasticity, 2014.
- Gordon, S., Sharp, M., Hubbard, B., Smart, C., Ketterling, B., and Willis, I.: Seasonal reorganization of subglacial drainage inferred from measurements in boreholes, *Hydrological Processes*, 12, 105–133, 1998.
- 765 Gräff, D., Köpfl, M., Lipovsky, B. P., Selvadurai, P. A., Farinotti, D., and Walter, F.: Fine structure of microseismic glacial stick-slip, *Geophysical Research Letters*, 48, e2021GL096043, 2021.
- Gulley, J.: Structural control of englacial conduits in the temperate Matanuska Glacier, Alaska, USA, *Journal of Glaciology*, 55, 681–690, 2009.
- Hagen, J. O., Liestøl, O., Roland, E., and Jørgensen, T.: *Glacier atlas of Svalbard and Jan Mayen*, vol. 129, Norsk Polarinstitutt Oslo, 1993.
- 770 Hansen, D. and Zoet, L.: Characterizing sediment flux of deforming glacier beds, *Journal of Geophysical Research: Earth Surface*, 127, e2021JF006544, 2022.
- Hjelle, A.: *Geology of Svalbard*, 1993.
- Hoffman, M. J., Andrews, L. C., Price, S. F., Catania, G. A., Neumann, T. A., Lüthi, M. P., Gulley, J., Ryser, C., Hawley, R. L., and Morriss, B.: Greenland subglacial drainage evolution regulated by weakly connected regions of the bed, *Nature communications*, 7, 13903, 2016.
- 775 Hoffmann, K.: *Applying the wheatstone bridge circuit*, HBM Germany, 1974.
- Hooke, R. L., Laumann, T., and Kohler, J.: Subglacial water pressures and the shape of subglacial conduits, *Journal of Glaciology*, 36, 67–71, 1990.
- Hooke, R. L., Hanson, B., Iverson, N. R., Jansson, P., and Fischer, U. H.: Rheology of till beneath Storglaciären, Sweden, *Journal of Glaciology*, 43, 172–179, 1997.
- 780 Hubbard, B., Sharp, M., Willis, I., Nielsen, M., and Smart, C.: Borehole water-level variations and the structure of the subglacial hydrological system of Haut Glacier d’Arolla, Valais, Switzerland, *Journal of Glaciology*, 41, 572–583, 1995.
- Hudson, T., Kufner, S., Brisbourne, A., Kendall, J., Smith, A., Alley, R., Arthern, R., and Murray, T.: Highly variable friction and slip observed at Antarctic ice stream bed, *Nature Geoscience*, pp. 1–7, 2023.
- Humphrey, N., Kamb, B., Fahnestock, M., and Engelhardt, H.: Characteristics of the bed of the lower Columbia Glacier, Alaska, *Journal of Geophysical Research: Solid Earth*, 98, 837–846, 1993.
- 785 Iken, A.: The effect of the subglacial water pressure on the sliding velocity of a glacier in an idealized numerical model, *Journal of Glaciology*, 27, 407–421, 1981.
- Iken, A. and Bindschadler, R. A.: Combined measurements of subglacial water pressure and surface velocity of Findelengletscher, Switzerland: conclusions about drainage system and sliding mechanism, *Journal of Glaciology*, 32, 101–119, 1986.
- 790 Iken, A. and Truffer, M.: The relationship between subglacial water pressure and velocity of Findelengletscher, Switzerland, during its advance and retreat, *Journal of Glaciology*, 43, 328–338, 1997.
- IPCC: *Special Report on the Ocean and Cryosphere in a Changing Climate* [H.-O. Portner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)], vol. In Press, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157964>, 2019.
- 795 IPCC: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, vol. In Press, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.

- Irvine-Fynn, T. D., Hodson, A. J., Moorman, B. J., Vatne, G., and Hubbard, A. L.: Polythermal glacier hydrology: A review, *Reviews of Geophysics*, 49, 2011.
- 800 Iverson, N. R.: Shear resistance and continuity of subglacial till: hydrology rules, *Journal of Glaciology*, 56, 1104–1114, 2010.
- Iverson, N. R. and Iverson, R. M.: Distributed shear of subglacial till due to Coulomb slip, *Journal of Glaciology*, 47, 481–488, 2001.
- Iverson, N. R., Jansson, P., and Hooke, R. L.: In-situ measurement of the strength of deforming subglacial till, *Journal of Glaciology*, 40, 497–503, 1994.
- Iverson, N. R., Hanson, B., Hooke, R. L., and Jansson, P.: Flow mechanism of glaciers on soft beds, *Science*, 267, 80–81, 1995.
- 805 Iverson, N. R., Hooyer, T. S., and Baker, R. W.: Ring-shear studies of till deformation: Coulomb-plastic behavior and distributed strain in glacier beds, *Journal of Glaciology*, 44, 634–642, 1998.
- Iverson, N. R., Hooyer, T. S., Fischer, U. H., Cohen, D., Moore, P. L., Jackson, M., Lappégard, G., and Kohler, J.: Soft-bed experiments beneath Engabreen, Norway: regelation infiltration, basal slip and bed deformation, *Journal of Glaciology*, 53, 323–340, 2007.
- Javed, A., Hamshaw, S. D., Lee, B. S., and Rizzo, D. M.: Multivariate event time series analysis using hydrological and suspended sediment
- 810 data, *Journal of Hydrology*, 593, 125–132, 2021.
- Kamb, B.: Glacier surge mechanism based on linked cavity configuration of the basal water conduit system, *Journal of Geophysical Research: Solid Earth*, 92, 9083–9100, 1987.
- Kamb, B.: Basal zone of the West Antarctic ice streams and its role in lubrication of their rapid motion, *The West Antarctic ice sheet: behavior and environment*, 77, 157–199, 2001.
- 815 Kamb, B. and Engelhardt, H.: Antarctic ice stream B: conditions controlling its motions and interactions with the climate system, *IAHS Publication*, 208, 145–154, 1991.
- Kavanaugh, J. L. and Clarke, G. K.: Evidence for extreme pressure pulses in the subglacial water system, *Journal of Glaciology*, 46, 206–212, 2000.
- Kavanaugh, J. L. and Clarke, G. K.: Discrimination of the flow law for subglacial sediment using in situ measurements and an interpretation
- 820 model, *Journal of Geophysical Research: Earth Surface*, 111, 2006.
- Köhler, A., Chapuis, A., Nuth, C., Kohler, J., and Weidle, C.: Autonomous detection of calving-related seismicity at Kronebreen, Svalbard, *The Cryosphere*, 6, 393–406, 2012.
- Köhler, A., Nuth, C., Schweitzer, J., Weidle, C., and Gibbons, S. J.: Regional passive seismic monitoring reveals dynamic glacier activity on Spitsbergen, Svalbard, *Polar Research*, 34, 26–33, 2015.
- 825 Köpfli, M., Gräff, D., Lipovsky, B. P., Selvadurai, P. A., Farinotti, D., and Walter, F.: Hydraulic Conditions for Stick-Slip Tremor Beneath an Alpine Glacier, *Geophysical Research Letters*, 49, e2022GL100286, 2022.
- Labeledz, C. R., Bartholomäus, T. C., Amundson, J. M., Gimbert, F., Karplus, M. S., Tsai, V. C., and Veitch, S. A.: Seismic mapping of subglacial hydrology reveals previously undetected pressurization event, *Journal of Geophysical Research: Earth Surface*, 127, e2021JF006406, 2022a.
- 830 Labeledz, C. R., Bartholomäus, T. C., Amundson, J. M., Karplus, M. S., Veitch, S. A., and Shugar, D. H.: Swarm-Like Behavior of Icequakes Associated with Surface Crevassing Activity on a Mountain Glacier, in: *AGU Fall Meeting Abstracts*, vol. 2022, pp. NS43A–04, 2022b.
- Lappégard, G., Kohler, J., Jackson, M., and Hagen, J. O.: Characteristics of subglacial drainage systems deduced from load-cell measurements, *Journal of Glaciology*, 52, 137–148, <https://doi.org/10.3189/172756506781828908>, 2006.
- Liestøl, O.: The glaciers in the Kongsfjorden area, Spitsbergen, *Norsk Geografisk Tidsskrift-Norwegian Journal of Geography*, 42, 231–238, 835 1988.

- Lindner, F., Walter, F., Laske, G., and Gimbert, F.: Glaciohydraulic seismic tremors on an Alpine glacier, *The Cryosphere*, 14, 287–308, 2020.
- Liboutry, L.: General theory of subglacial cavitation and sliding of temperate glaciers, *Journal of Glaciology*, 7, 21–58, 1968.
- Maier, N., Gimbert, F., and Gillet-Chaulet, F.: Threshold response to melt drives large-scale bed weakening in Greenland, *Nature*, 607, 714–720, 2022.
- Mair, D., Nienow, P., Willis, I., and Sharp, M.: Spatial patterns of glacier motion during a high-velocity event: Haut Glacier d’Arolla, Switzerland, *Journal of Glaciology*, 47, 9–20, 2001.
- Mair, D., Willis, I., Fischer, U. H., Hubbard, B., Nienow, P., and Hubbard, A.: Hydrological controls on patterns of surface, internal and basal motion during three “spring events”: Haut Glacier d’Arolla, Switzerland, *Journal of Glaciology*, 49, 555–567, 2003.
- Melvold, K. and Hagen, J. O.: Evolution of a surge-type glacier in its quiescent phase: Kongsvegen, Spitsbergen, 1964–95, *Journal of Glaciology*, 44, 394–404, 1998.
- Minchew, B., Simons, M., Björnsson, H., Pálsson, F., Morlighem, M., Seroussi, H., Larour, E., and Hensley, S.: Plastic bed beneath Hofsjökull Ice Cap, central Iceland, and the sensitivity of ice flow to surface meltwater flux, *Journal of Glaciology*, 62, 147–158, 2016.
- Mitchell, J. K., Soga, K., et al.: *Fundamentals of soil behavior*, vol. 3, John Wiley & Sons New York, 2005.
- Moon, T., Ahlstrøm, A., Goelzer, H., Lipscomb, W., and Nowicki, S.: Rising oceans guaranteed: Arctic land ice loss and sea level rise, *Current climate change reports*, 4, 211–222, 2018.
- Müller, M., Homleid, M., Ivarsson, K. I., Køltzow, M. A., Lindskog, M., Midtbø, K. H., Andrae, U., Aspelien, T., Berggren, L., Bjørge, D., Dahlgren, P., Kristiansen, J., Randriamampianina, R., Ridal, M., and Vignes, O.: AROME-MetCoOp: A nordic convective-scale operational weather prediction model, *Weather and Forecasting*, 32, 609–627, <https://doi.org/10.1175/WAF-D-16-0099.1>, 2017.
- Murray, T. and Booth, A. D.: Imaging glacial sediment inclusions in 3-D using ground-penetrating radar at Kongsvegen, Svalbard, *Journal of Quaternary science*, 25, 754–761, 2010.
- Murray, T. and Clarke, G. K.: Black-box modeling of the subglacial water system, *Journal of Geophysical Research: Solid Earth*, 100, 10 231–10 245, 1995.
- Murray, T. and Porter, P. R.: Basal conditions beneath a soft-bedded polythermal surge-type glacier: Bakaninbreen, Svalbard, *Quaternary International*, 86, 103–116, 2001.
- Nanni, U., Gimbert, F., Vincent, C., Gräff, D., Walter, F., Piard, L., and Moreau, L.: Quantification of seasonal and diurnal dynamics of subglacial channels using seismic observations on an Alpine glacier, *The Cryosphere*, 14, 1475–1496, 2020.
- Nanni, U., Gimbert, F., Roux, P., and Lecointre, A.: Observing the subglacial hydrology network and its dynamics with a dense seismic array, *Proceedings of the National Academy of Sciences*, 118, e2023757 118, 2021.
- Nanni, U., Roux, P., Gimbert, F., and Lecointre, A.: Dynamic Imaging of Glacier Structures at High-Resolution Using Source Localization With a Dense Seismic Array, *Geophysical Research Letters*, 49, e2021GL095 996, 2022.
- Nanni, U., Scherler, D., Ayoub, F., Millan, R., Herman, F., and Avouac, J.-P.: Climatic control on seasonal variations in mountain glacier surface velocity, *The Cryosphere*, 17, 1567–1583, 2023.
- Ng, F. S.: Canals under sediment-based ice sheets, *Annals of Glaciology*, 30, 146–152, 2000.
- Nienow, P., Sole, A., Slater, D. A., and Cowton, T.: Recent advances in our understanding of the role of meltwater in the Greenland Ice Sheet system, *Current Climate Change Reports*, 3, 330–344, 2017.
- Nuth, C., Schuler, T. V., Kohler, J., Altena, B., and Hagen, J. O.: Estimating the long-term calving flux of Kronebreen, Svalbard, from geodetic elevation changes and mass-balance modeling, *Journal of Glaciology*, 58, 119–133, 2012.

- Nye, J. F.: Water flow in glaciers: jökulhlaups, tunnels and veins, *Journal of Glaciology*, 17, 181–207, 1976.
- 875 Podolskiy, E. A. and Walter, F.: Cryoseismology, *Reviews of geophysics*, 54, 708–758, 2016.
- Porter, P. R., Murray, T., and Dowdeswell, J. A.: Sediment deformation and basal dynamics beneath a glacier surge front: Bakaninbreen, Svalbard, *Annals of Glaciology*, 24, 21–26, 1997.
- Pramanik, A., Kohler, J., Lindbäck, K., How, P., Van Pelt, W., Liston, G., and Schuler, T. V.: Hydrology and runoff routing of glacierized drainage basins in the Kongsfjord area, northwest Svalbard, *The Cryosphere Discussions*, pp. 1–33, 2020.
- 880 Preiswerk, L. E. and Walter, F.: High-Frequency (> 2 Hz) Ambient Seismic Noise on High-Melt Glaciers: Green’s Function Estimation and Source Characterization, *Journal of Geophysical Research: Earth Surface*, 123, 1667–1681, 2018.
- Rada, C. and Schoof, C.: Channelized, distributed, and disconnected: subglacial drainage under a valley glacier in the Yukon, *The Cryosphere*, 12, 2609–2636, 2018.
- Rada Giacaman, C. A. and Schoof, C.: Channelised, distributed, and disconnected: Spatial structure and temporal evolution of the subglacial drainage under a valley glacier in the Yukon, *The Cryosphere Discussions*, pp. 1–39, 2022.
- 885 Rada Giacaman, C. A. and Schoof, C.: Channelized, distributed, and disconnected: spatial structure and temporal evolution of the subglacial drainage under a valley glacier in the Yukon, *The Cryosphere*, 17, 761–787, <https://doi.org/10.5194/tc-17-761-2023>, 2023.
- RGI, C.: Randolph Glacier Inventory (RGI) - A dataset of Global Glacier Outlines: Version 6.0, <http://www.glims.org/RGI/randolph60.html>, 2017.
- 890 Röthlisberger, H.: Water pressure in intra-and subglacial channels, *Journal of Glaciology*, 11, 177–203, 1972.
- Röthlisberger, H. and Lang, H.: *Glacial hydrology*, John Wiley & Sons, 1987.
- Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D., Compagno, L., Copland, L., et al.: Global glacier change in the 21st century: Every increase in temperature matters, *Science*, 379, 78–83, 2023.
- Rousselot, M. and Fischer, U. H.: A laboratory study of ploughing, *Journal of Glaciology*, 53, 225–231, 2007.
- 895 Roux, P.-F., Marsan, D., Métaxian, J.-P., O’Brien, G., and Moreau, L.: Microseismic activity within a serac zone in an alpine glacier (Glacier d’Argentiere, Mont Blanc, France), *Journal of Glaciology*, 54, 157–168, 2008.
- Schmidt, L. S., Schuler, T. V., Thomas, E. E., and Westermann, S.: Meltwater runoff and glacier mass balance in the high Arctic: 1991–2022 simulations for Svalbard, *The Cryosphere*, 17, 2941–2963, 2023.
- Schofield, A. N. and Wroth, P.: *Critical state soil mechanics*, vol. 310, McGraw-hill London, 1968.
- 900 Scholzen, C., Schuler, T. V., and Gilbert, A.: Sensitivity of subglacial drainage to water supply distribution at the Kongsfjord basin, Svalbard, *The Cryosphere*, 15, 2719–2738, 2021.
- Schoof, C.: The effect of cavitation on glacier sliding, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 461, 609–627, 2005.
- Schoof, C.: Ice-sheet acceleration driven by melt supply variability, *Nature*, 468, 803–806, 2010.
- 905 Schuler, T., Fischer, U. H., Sterr, R., Hock, R., and Gudmundsson, G. H.: Comparison of Modeled Water Input and Measured Discharge Prior to a Release Event: Unteraargletscher, Bernese Alps, Switzerland: Selected paper from EGS General Assembly, Nice, April-2000 (Symposium OA36), *Hydrology Research*, 33, 27–46, 2002.
- Schwanghart, W. and Scherler, D.: TopoToolbox 2 – MATLAB-based software for topographic analysis and modeling in Earth surface sciences, *Earth Surface Dynamics*, 2, 1–7, <https://doi.org/10.5194/esurf-2-1-2014>, 2014.
- 910 Schyberg, H., Yang, X., Kjøltzow, M., Amstrup, B., Bakketun, Å., Bazile, E., Bojarova, J., Box, J., Dahlgren, P., Hagelin, S., Homleid, M., Horányi, A., Høyer, J., Johansson, Å., Killie, M., Körnich, H., Le Moigne, P., Lindskog, M., Manninen, T., Nielsen Englyst, P., and Wang,

- Z.: Arctic regional reanalysis on single levels from 1991 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS), <https://doi.org/10.24381/cds.713858f6>, 2020.
- 915 Sergeant, A., Yastrebov, V. A., Mangeney, A., Castelnaud, O., Montagner, J.-P., and Stutzmann, E.: Numerical modeling of iceberg capsizing responsible for glacial earthquakes, *Journal of Geophysical Research: Earth Surface*, 123, 3013–3033, 2018.
- Sommers, A., Meyer, C., Morlighem, M., Rajaram, H., Poinar, K., Chu, W., and Mejia, J.: Subglacial hydrology modeling predicts high winter water pressure and spatially variable transmissivity at Helheim Glacier, Greenland, *Journal of Glaciology*, pp. 1–13, 2023.
- Stocker, T.: *Climate change 2013: the physical science basis: Working Group I contribution to the Fifth assessment report of the Intergovernmental Panel on Climate Change*, Cambridge university press, 2014.
- 920 Sugiyama, S. and Gudmundsson, G. H.: Diurnal variations in vertical strain observed in a temperate valley glacier, *Geophysical Research Letters*, 30, 2003.
- Sugiyama, S., Skvarca, P., Naito, N., Enomoto, H., Tsutaki, S., Tone, K., Marinsek, S., and Aniya, M.: Ice speed of a calving glacier modulated by small fluctuations in basal water pressure, *Nature Geoscience*, 4, 597–600, 2011.
- Sugiyama, S., Navarro, F. J., Sawagaki, T., Minowa, M., Segawa, T., Onuma, Y., Otero, J., and Vasilenko, E. V.: Subglacial water pressure and ice-speed variations at Johnsons Glacier, Livingston Island, Antarctic Peninsula, *Journal of Glaciology*, 65, 689–699, 2019.
- 925 Terzaghi, K., Peck, R. B., and Mesri, G.: *Soil mechanics in engineering practice*, John wiley & sons, 1996.
- Thøgersen, K., Gilbert, A., Schuler, T. V., and Malthe-Sørensen, A.: Rate-and-state friction explains glacier surge propagation, *Nature communications*, 10, 1–8, <https://doi.org/10.1038/s41467-019-10506-4>, 2019.
- Thomason, J. F. and Iverson, N. R.: A laboratory study of particle ploughing and pore-pressure feedback: a velocity-weakening mechanism for soft glacier beds, *Journal of Glaciology*, 54, 169–181, 2008.
- 930 Truffer, M.: The basal speed of valley glaciers: an inverse approach, *Journal of Glaciology*, 50, 236–242, 2004.
- Truffer, M., Harrison, W. D., and Echelmeyer, K. A.: Glacier motion dominated by processes deep in underlying till, *Journal of Glaciology*, 46, 213–221, 2000.
- Truffer, M., Kääh, A., Harrison, W. D., Osipova, G. B., Nosenko, G. A., Espizua, L., Gilbert, A., Fischer, L., Huggel, C., and Burns, P. A. C. e. a.: Glacier surges, in: *Snow and Ice-Related Hazards, Risks, and Disasters*, pp. 417–466, Elsevier, 2021.
- 935 Tsai, V. C., Smith, L. C., Gardner, A. S., and Seroussi, H.: A unified model for transient subglacial water pressure and basal sliding, *Journal of Glaciology*, 68, 390–400, 2022.
- Tulaczyk, S.: Ice sliding over weak, fine-grained tills: dependence of ice-till interactions on till granulometry, *SPECIAL PAPERS-GEOLOGICAL SOCIETY OF AMERICA*, pp. 159–178, 1999.
- 940 Tulaczyk, S., Kamb, W. B., and Engelhardt, H. F.: Basal mechanics of ice stream B, West Antarctica: 1. Till mechanics, *Journal of Geophysical Research: Solid Earth*, 105, 463–481, 2000.
- Tulaczyk, S. M., Scherer, R. P., and Clark, C. D.: A ploughing model for the origin of weak tills beneath ice streams: a qualitative treatment, *Quaternary International*, 86, 59–70, 2001.
- Vincent, C. and Moreau, L.: Sliding velocity fluctuations and subglacial hydrology over the last two decades on Argentière glacier, Mont Blanc area, *Journal of Glaciology*, 62, 805–815, 2016.
- 945 Walder, J. S.: Stability of sheet flow of water beneath temperate glaciers and implications for glacier surging, *Journal of Glaciology*, 28, 273–293, 1982.
- Walder, J. S.: Hydraulics of subglacial cavities, *Journal of Glaciology*, 32, 439–445, 1986.
- Walder, J. S. and Fowler, A.: Channelized subglacial drainage over a deformable bed, *Journal of glaciology*, 40, 3–15, 1994.

- 950 Walker, R. T., Christianson, K., Parizek, B. R., Anandakrishnan, S., and Alley, R. B.: A viscoelastic flowline model applied to tidal forcing of Bindschadler Ice Stream, West Antarctica, *Earth and Planetary Science Letters*, 319, 128–132, 2012.
- Warburton, K., Hewitt, D., and Neufeld, J.: Shear dilation of subglacial till results in time-dependent sliding laws, *Proceedings of the Royal Society A*, 479, 20220536, 2023.
- Weertman, J.: On the sliding of glaciers, *Journal of glaciology*, 3, 33–38, 1957.
- 955 Weertman, J.: General theory of water flow at the base of a glacier or ice sheet, *Reviews of Geophysics*, 10, 287–333, 1972.
- Welch, P.: The use of fast Fourier transform for the estimation of power spectra: a method based on time averaging over short, modified periodograms, *IEEE Transactions on audio and electroacoustics*, 15, 70–73, 1967.
- Werder, M. A., Hewitt, I. J., Schoof, C. G., and Flowers, G. E.: Modeling channelized and distributed subglacial drainage in two dimensions, *Journal of Geophysical Research: Earth Surface*, 118, 2140–2158, 2013.
- 960 Westermann, S., Ingeman-Nielsen, T., Scheer, J., Aalstad, K., Aga, J., Chaudhary, N., Etzelmüller, B., Filhol, S., Käab, A., Renette, C., et al.: The CryoGrid community model (version 1.0)—a multi-physics toolbox for climate-driven simulations in the terrestrial cryosphere, *Geoscientific Model Development*, 16, 2607–2647, 2023.
- Wiens, D. A., Anandakrishnan, S., Winberry, J. P., and King, M. A.: Simultaneous teleseismic and geodetic observations of the stick–slip motion of an Antarctic ice stream, *Nature*, 453, 770–774, 2008.
- 965 Woodward, J., Murray, T., and McCaig, A.: Formation and reorientation of structure in the surge-type glacier Kongsvegen, Svalbard, *Journal of Quaternary Science: Published for the Quaternary Research Association*, 17, 201–209, 2002.
- Yang, X., Nielsen, K. P., Amstrup, B., Peralta, C., Høyer, J., Englyst, P. N., Schyberg, H., Homleid, M., Kpltzow, M. ø., Randriamampianina, R., Dahlgren, P., Stpylen, E., Valkonen, T., Palmason, B., Thorsteinnsson, S., Bojarova, J., Körnich, H., Lindskog, M., Box, J., and Mankoff, K.: C3S Arctic regional reanalysis – Full system documentation, Tech. rep., 2021.
- 970 Zhan, Z.: Seismic noise interferometry reveals transverse drainage configuration beneath the surging Bering Glacier, *Geophysical Research Letters*, 46, 4747–4756, 2019.
- Zoet, L. K. and Iverson, N. R.: A slip law for glaciers on deformable beds, *Science*, 368, 76–78, 2020.

Appendix A: Full time serie




Figure A1. Time series of physical quantities measured from spring 2021 to summer 2022. (a) Temperature (black line), precipitation (light blue bars) from CARRA/AROME-Artic, and relative glacier surface height (grey line) from Cryogrid simulations (Schmidt et al., 2023). The three variables are extracted for the closest grid point of the borehole. In 2022, the surface height is negative which corresponds to ice melt. (b) Modelled runoff (blue line) and glacier surface velocity measured (red line). Circled numbers refer to different episodes described in the main text. (c) turbulent-water-flow-induced seismic power recorded at the surface of the glacier in the 3-10 Hz frequency band (yellow line). (d) Borehole water pressure (green line) and force acting on the ploughmeter (dark red line). Blue shaded areas represent the melt seasons. Grey shaded areas represent periods of missing data.

Appendix B: Pre-processing of the time series

975 [Need some text here]

Appendix C: Surface velocity data

The velocity record presented in this study combines two different GNSS records. One of the GNSS stations is positioned at stake 6 (KNG-6: 13.15153°E 78.78067°N) and the other one at stake 7 (KNG-7: 13.23962°E 78.76770°N). In figure C1, we present the two datasets that have been combined. We apply a one-week moving median for KNG7 velocity to smooth the record especially during the winter period when the velocities are low and thus the daily velocity derivation is less accurate.

980




Figure B1. Pre-processing workflow applied to the time series. The original time-series (see Fig. 3 below) have been filtered at three time-scales. The multi-day and diurnal filtered data have been inspected against the unfiltered data to remove spurious artefacts that can be created by the filtering technique (see also Appendix G, G1). We then segmented the recorded data into multiple events and ~~normalised~~normalized the magnitude and duration of each.




Figure C1. Velocity data series from GNSS stations KNG6 (red curve) and KNG7 (blue curve).

Appendix D: Events

Table D1. Description of the twelve multi-day time scale events during the melt season 2021

Event	Start	Stop	Duration
0	2021-07-01 00:00:00	2021-07-03 03:00:00	2 days 03:00:00
1	2021-07-03 03:00:00	2021-07-13 06:00:00	10 days 03:00:00
2	2021-07-13 06:00:00	2021-07-22 03:00:00	8 days 21:00:00
3	2021-07-22 03:00:00	2021-07-31 03:00:00	9 days 00:00:00
4	2021-07-31 03:00:00	2021-08-08 12:00:00	8 days 09:00:00
5	2021-08-08 12:00:00	2021-08-17 21:00:00	9 days 09:00:00
6	2021-08-17 21:00:00	2021-08-28 00:00:00	10 days 03:00:00
7	2021-08-28 00:00:00	2021-09-04 15:00:00	7 days 15:00:00
8	2021-09-04 15:00:00	2021-09-15 21:00:00	11 days 06:00:00
9	2021-09-15 21:00:00	2021-09-27 03:00:00	11 days 06:00:00
10	2021-09-27 03:00:00	2021-10-04 12:00:00	7 days 09:00:00
11	2021-10-04 12:00:00	2021-10-04 21:00:00	0 days 09:00:00

Table D2. Description of the eight multi-day time scale events during the melt season 2022

Event	Start	Stop	Duration
0	2022-05-25 00:00:00	2022-05-28 21:00:00	3 days 21:00:00
1	2022-05-28 21:00:00	2022-06-10 12:00:00	12 days 15:00:00
2	2022-06-10 12:00:00	2022-06-26 06:00:00	15 days 18:00:00
3	2022-06-26 06:00:00	2022-07-11 06:00:00	15 days 00:00:00
4	2022-07-11 06:00:00	2022-07-23 18:00:00	12 days 12:00:00
5	2022-07-23 18:00:00	2022-08-07 18:00:00	15 days 00:00:00
6	2022-08-07 18:00:00	2022-08-16 00:00:00	8 days 06:00:00
7	2022-08-16 00:00:00	2022-08-16 15:00:00	0 days 15:00:00

Table D3. Description of the 96 diurnal time scale events during the melt season 2021

Event	Start	Stop	Duration	Event	Start	Stop	Duration
0	2021-07-01 00:00:00	2021-07-02 09:00:00	1 days 09:00:00	36	2021-08-06 12:00:00	2021-08-07 03:00:00	0 days 15:00:00
1	2021-07-02 09:00:00	2021-07-03 18:00:00	1 days 09:00:00	37	2021-08-07 03:00:00	2021-08-07 15:00:00	0 days 12:00:00
2	2021-07-03 18:00:00	2021-07-04 15:00:00	0 days 21:00:00	38	2021-08-07 15:00:00	2021-08-08 15:00:00	1 days 00:00:00
3	2021-07-04 15:00:00	2021-07-05 15:00:00	1 days 00:00:00	39	2021-08-08 15:00:00	2021-08-09 15:00:00	1 days 00:00:00
4	2021-07-05 15:00:00	2021-07-06 15:00:00	1 days 00:00:00	40	2021-08-09 15:00:00	2021-08-10 15:00:00	1 days 00:00:00
5	2021-07-06 15:00:00	2021-07-07 15:00:00	1 days 00:00:00	41	2021-08-10 15:00:00	2021-08-11 15:00:00	1 days 00:00:00
6	2021-07-07 15:00:00	2021-07-08 18:00:00	1 days 03:00:00	42	2021-08-11 15:00:00	2021-08-12 15:00:00	1 days 00:00:00
7	2021-07-08 18:00:00	2021-07-09 09:00:00	0 days 15:00:00	43	2021-08-12 15:00:00	2021-08-13 12:00:00	0 days 21:00:00
8	2021-07-09 09:00:00	2021-07-10 15:00:00	1 days 06:00:00	44	2021-08-13 12:00:00	2021-08-14 12:00:00	1 days 00:00:00
9	2021-07-10 15:00:00	2021-07-11 12:00:00	0 days 21:00:00	45	2021-08-14 12:00:00	2021-08-15 15:00:00	1 days 03:00:00
10	2021-07-11 12:00:00	2021-07-12 12:00:00	1 days 00:00:00	46	2021-08-15 15:00:00	2021-08-16 06:00:00	0 days 15:00:00
11	2021-07-12 12:00:00	2021-07-13 18:00:00	1 days 06:00:00	47	2021-08-16 06:00:00	2021-08-16 18:00:00	0 days 12:00:00
12	2021-07-13 18:00:00	2021-07-14 15:00:00	0 days 21:00:00	48	2021-08-16 18:00:00	2021-08-17 18:00:00	1 days 00:00:00
13	2021-07-14 15:00:00	2021-07-15 15:00:00	1 days 00:00:00	49	2021-08-17 18:00:00	2021-08-18 15:00:00	0 days 21:00:00
14	2021-07-15 15:00:00	2021-07-16 12:00:00	0 days 21:00:00	50	2021-08-18 15:00:00	2021-08-19 15:00:00	1 days 00:00:00
15	2021-07-16 12:00:00	2021-07-17 15:00:00	1 days 03:00:00	51	2021-08-19 15:00:00	2021-08-20 18:00:00	1 days 03:00:00
16	2021-07-17 15:00:00	2021-07-18 15:00:00	1 days 00:00:00	52	2021-08-20 18:00:00	2021-08-21 18:00:00	1 days 00:00:00
17	2021-07-18 15:00:00	2021-07-19 15:00:00	1 days 00:00:00	53	2021-08-21 18:00:00	2021-08-22 15:00:00	0 days 21:00:00
18	2021-07-19 15:00:00	2021-07-20 12:00:00	0 days 21:00:00	54	2021-08-22 15:00:00	2021-08-24 12:00:00	1 days 21:00:00
19	2021-07-20 12:00:00	2021-07-21 12:00:00	1 days 00:00:00	55	2021-08-24 12:00:00	2021-08-25 15:00:00	1 days 03:00:00
20	2021-07-21 12:00:00	2021-07-22 15:00:00	1 days 03:00:00	56	2021-08-25 15:00:00	2021-08-26 09:00:00	0 days 18:00:00
21	2021-07-22 15:00:00	2021-07-23 12:00:00	0 days 21:00:00	57	2021-08-26 09:00:00	2021-08-27 15:00:00	1 days 06:00:00
22	2021-07-23 12:00:00	2021-07-24 12:00:00	1 days 00:00:00	58	2021-08-27 15:00:00	2021-08-28 15:00:00	1 days 00:00:00
23	2021-07-24 12:00:00	2021-07-25 12:00:00	1 days 00:00:00	59	2021-08-28 15:00:00	2021-08-29 15:00:00	1 days 00:00:00
24	2021-07-25 12:00:00	2021-07-26 12:00:00	1 days 00:00:00	60	2021-08-29 15:00:00	2021-08-30 12:00:00	0 days 21:00:00
25	2021-07-26 12:00:00	2021-07-27 15:00:00	1 days 03:00:00	61	2021-08-30 12:00:00	2021-08-31 12:00:00	1 days 00:00:00
26	2021-07-27 15:00:00	2021-07-28 15:00:00	1 days 00:00:00	62	2021-08-31 12:00:00	2021-09-01 09:00:00	0 days 21:00:00
27	2021-07-28 15:00:00	2021-07-29 12:00:00	0 days 21:00:00	63	2021-09-01 09:00:00	2021-09-02 06:00:00	0 days 21:00:00
28	2021-07-29 12:00:00	2021-07-30 12:00:00	1 days 00:00:00	64	2021-09-02 06:00:00	2021-09-04 09:00:00	2 days 03:00:00
29	2021-07-30 12:00:00	2021-07-31 15:00:00	1 days 03:00:00	65	2021-09-04 09:00:00	2021-09-05 03:00:00	0 days 18:00:00
30	2021-07-31 15:00:00	2021-08-01 12:00:00	0 days 21:00:00	66	2021-09-05 03:00:00	2021-09-06 00:00:00	0 days 21:00:00
31	2021-08-01 12:00:00	2021-08-02 12:00:00	1 days 00:00:00	67	2021-09-06 00:00:00	2021-09-06 15:00:00	0 days 15:00:00
32	2021-08-02 12:00:00	2021-08-03 06:00:00	0 days 18:00:00	68	2021-09-06 15:00:00	2021-09-07 03:00:00	0 days 12:00:00
33	2021-08-03 06:00:00	2021-08-04 12:00:00	1 days 06:00:00	69	2021-09-07 03:00:00	2021-09-08 00:00:00	0 days 21:00:00
34	2021-08-04 12:00:00	2021-08-05 12:00:00	1 days 00:00:00	70	2021-09-08 00:00:00	2021-09-08 18:00:00	0 days 18:00:00
35	2021-08-05 12:00:00	2021-08-06 12:00:00	1 days 00:00:00	71	2021-09-08 18:00:00	2021-09-09 18:00:00	1 days 00:00:00

Table D3. (Following) Description of the 96 diurnal time scale events during the melt season 2021

Event	Start	Stop	Duration
72	2021-09-09 18:00:00	2021-09-10 18:00:00	1 days 00:00:00
73	2021-09-10 18:00:00	2021-09-11 09:00:00	0 days 15:00:00
74	2021-09-11 09:00:00	2021-09-12 06:00:00	0 days 21:00:00
75	2021-09-12 06:00:00	2021-09-13 03:00:00	0 days 21:00:00
76	2021-09-13 03:00:00	2021-09-14 09:00:00	1 days 06:00:00
77	2021-09-14 09:00:00	2021-09-15 06:00:00	0 days 21:00:00
78	2021-09-15 06:00:00	2021-09-19 00:00:00	3 days 18:00:00
79	2021-09-19 00:00:00	2021-09-20 12:00:00	1 days 12:00:00
80	2021-09-20 12:00:00	2021-09-22 15:00:00	2 days 03:00:00
81	2021-09-22 15:00:00	2021-09-23 06:00:00	0 days 15:00:00
82	2021-09-23 06:00:00	2021-09-24 09:00:00	1 days 03:00:00
83	2021-09-24 09:00:00	2021-09-25 18:00:00	1 days 09:00:00
84	2021-09-25 18:00:00	2021-09-27 00:00:00	1 days 06:00:00
85	2021-09-27 00:00:00	2021-09-27 12:00:00	0 days 12:00:00
86	2021-09-27 12:00:00	2021-09-28 03:00:00	0 days 15:00:00
87	2021-09-28 03:00:00	2021-09-29 03:00:00	1 days 00:00:00
88	2021-09-29 03:00:00	2021-09-29 18:00:00	0 days 15:00:00
89	2021-09-29 18:00:00	2021-09-30 09:00:00	0 days 15:00:00
90	2021-09-30 09:00:00	2021-10-01 03:00:00	0 days 18:00:00
91	2021-10-01 03:00:00	2021-10-02 18:00:00	1 days 15:00:00
92	2021-10-02 18:00:00	2021-10-03 12:00:00	0 days 18:00:00
93	2021-10-03 12:00:00	2021-10-04 00:00:00	0 days 12:00:00
94	2021-10-04 00:00:00	2021-10-04 15:00:00	0 days 15:00:00
95	2021-10-04 15:00:00	2021-10-04 21:00:00	0 days 06:00:00

Table D4. Description of the 85 diurnal time scale events during the melt season 2022

Event	Start	Stop	Duration	Event	Start	Stop	Duration
0	2022-05-25 00:00:00	2022-05-25 06:00:00	0 days 06:00:00	36	2022-06-29 12:00:00	2022-06-30 12:00:00	1 days 00:00:00
1	2022-05-25 06:00:00	2022-05-26 12:00:00	1 days 06:00:00	37	2022-06-30 12:00:00	2022-07-01 12:00:00	1 days 00:00:00
2	2022-05-26 12:00:00	2022-05-27 03:00:00	0 days 15:00:00	38	2022-07-01 12:00:00	2022-07-02 06:00:00	0 days 18:00:00
3	2022-05-27 03:00:00	2022-05-27 18:00:00	0 days 15:00:00	39	2022-07-02 06:00:00	2022-07-03 12:00:00	1 days 06:00:00
4	2022-05-27 18:00:00	2022-05-28 15:00:00	0 days 21:00:00	40	2022-07-03 12:00:00	2022-07-04 12:00:00	1 days 00:00:00
5	2022-05-28 15:00:00	2022-05-29 15:00:00	1 days 00:00:00	41	2022-07-04 12:00:00	2022-07-05 09:00:00	0 days 21:00:00
6	2022-05-29 15:00:00	2022-05-30 15:00:00	1 days 00:00:00	42	2022-07-05 09:00:00	2022-07-06 09:00:00	1 days 00:00:00
7	2022-05-30 15:00:00	2022-05-31 15:00:00	1 days 00:00:00	43	2022-07-06 09:00:00	2022-07-07 09:00:00	1 days 00:00:00
8	2022-05-31 15:00:00	2022-06-01 12:00:00	0 days 21:00:00	44	2022-07-07 09:00:00	2022-07-08 09:00:00	1 days 00:00:00
9	2022-06-01 12:00:00	2022-06-02 15:00:00	1 days 03:00:00	45	2022-07-08 09:00:00	2022-07-09 09:00:00	1 days 00:00:00
10	2022-06-02 15:00:00	2022-06-03 12:00:00	0 days 21:00:00	46	2022-07-09 09:00:00	2022-07-10 12:00:00	1 days 03:00:00
11	2022-06-03 12:00:00	2022-06-04 09:00:00	0 days 21:00:00	47	2022-07-10 12:00:00	2022-07-11 09:00:00	0 days 21:00:00
12	2022-06-04 09:00:00	2022-06-05 15:00:00	1 days 06:00:00	48	2022-07-11 09:00:00	2022-07-12 06:00:00	0 days 21:00:00
13	2022-06-05 15:00:00	2022-06-06 18:00:00	1 days 03:00:00	49	2022-07-12 06:00:00	2022-07-13 06:00:00	1 days 00:00:00
14	2022-06-06 18:00:00	2022-06-07 12:00:00	0 days 18:00:00	50	2022-07-13 06:00:00	2022-07-14 12:00:00	1 days 06:00:00
15	2022-06-07 12:00:00	2022-06-08 15:00:00	1 days 03:00:00	51	2022-07-14 12:00:00	2022-07-15 09:00:00	0 days 21:00:00
16	2022-06-08 15:00:00	2022-06-09 18:00:00	1 days 03:00:00	52	2022-07-15 09:00:00	2022-07-16 09:00:00	1 days 00:00:00
17	2022-06-09 18:00:00	2022-06-10 18:00:00	1 days 00:00:00	53	2022-07-16 09:00:00	2022-07-17 09:00:00	1 days 00:00:00
18	2022-06-10 18:00:00	2022-06-11 15:00:00	0 days 21:00:00	54	2022-07-17 09:00:00	2022-07-18 09:00:00	1 days 00:00:00
19	2022-06-11 15:00:00	2022-06-12 15:00:00	1 days 00:00:00	55	2022-07-18 09:00:00	2022-07-19 09:00:00	1 days 00:00:00
20	2022-06-12 15:00:00	2022-06-13 15:00:00	1 days 00:00:00	56	2022-07-19 09:00:00	2022-07-20 09:00:00	1 days 00:00:00
21	2022-06-13 15:00:00	2022-06-14 15:00:00	1 days 00:00:00	57	2022-07-20 09:00:00	2022-07-21 09:00:00	1 days 00:00:00
22	2022-06-14 15:00:00	2022-06-15 15:00:00	1 days 00:00:00	58	2022-07-21 09:00:00	2022-07-22 09:00:00	1 days 00:00:00
23	2022-06-15 15:00:00	2022-06-16 15:00:00	1 days 00:00:00	59	2022-07-22 09:00:00	2022-07-23 06:00:00	0 days 21:00:00
24	2022-06-16 15:00:00	2022-06-17 18:00:00	1 days 03:00:00	60	2022-07-23 06:00:00	2022-07-24 09:00:00	1 days 03:00:00
25	2022-06-17 18:00:00	2022-06-18 12:00:00	0 days 18:00:00	61	2022-07-24 09:00:00	2022-07-25 06:00:00	0 days 21:00:00
26	2022-06-18 12:00:00	2022-06-19 15:00:00	1 days 03:00:00	62	2022-07-25 06:00:00	2022-07-26 09:00:00	1 days 03:00:00
27	2022-06-19 15:00:00	2022-06-21 15:00:00	2 days 00:00:00	63	2022-07-26 09:00:00	2022-07-27 09:00:00	1 days 00:00:00
28	2022-06-21 15:00:00	2022-06-22 12:00:00	0 days 21:00:00	64	2022-07-27 09:00:00	2022-07-28 09:00:00	1 days 00:00:00
29	2022-06-22 12:00:00	2022-06-23 09:00:00	0 days 21:00:00	65	2022-07-28 09:00:00	2022-07-29 09:00:00	1 days 00:00:00
30	2022-06-23 09:00:00	2022-06-24 12:00:00	1 days 03:00:00	66	2022-07-29 09:00:00	2022-07-30 06:00:00	0 days 21:00:00
31	2022-06-24 12:00:00	2022-06-25 12:00:00	1 days 00:00:00	67	2022-07-30 06:00:00	2022-07-31 09:00:00	1 days 03:00:00
32	2022-06-25 12:00:00	2022-06-26 12:00:00	1 days 00:00:00	68	2022-07-31 09:00:00	2022-08-01 06:00:00	0 days 21:00:00
33	2022-06-26 12:00:00	2022-06-27 15:00:00	1 days 03:00:00	69	2022-08-01 06:00:00	2022-08-02 09:00:00	1 days 03:00:00
34	2022-06-27 15:00:00	2022-06-28 09:00:00	0 days 18:00:00	70	2022-08-02 09:00:00	2022-08-03 09:00:00	1 days 00:00:00
35	2022-06-28 09:00:00	2022-06-29 12:00:00	1 days 03:00:00	71	2022-08-03 09:00:00	2022-08-04 06:00:00	0 days 21:00:00

Table D4. (Following) Description of the 85 diurnal time scale events during the melt season 2022

Event	Start	Stop	Duration
72	2022-08-04 06:00:00	2022-08-05 09:00:00	1 days 03:00:00
73	2022-08-05 09:00:00	2022-08-06 09:00:00	1 days 00:00:00
74	2022-08-06 09:00:00	2022-08-07 09:00:00	1 days 00:00:00
75	2022-08-07 09:00:00	2022-08-08 09:00:00	1 days 00:00:00
76	2022-08-08 09:00:00	2022-08-09 09:00:00	1 days 00:00:00
77	2022-08-09 09:00:00	2022-08-10 09:00:00	1 days 00:00:00
78	2022-08-10 09:00:00	2022-08-11 09:00:00	1 days 00:00:00
79	2022-08-11 09:00:00	2022-08-12 09:00:00	1 days 00:00:00
80	2022-08-12 09:00:00	2022-08-13 09:00:00	1 days 00:00:00
81	2022-08-13 09:00:00	2022-08-14 09:00:00	1 days 00:00:00
82	2022-08-14 09:00:00	2022-08-15 12:00:00	1 days 03:00:00
83	2022-08-15 12:00:00	2022-08-16 09:00:00	0 days 21:00:00
84	2022-08-16 09:00:00	2022-08-16 15:00:00	0 days 06:00:00

Appendix E: Spectrogram

Large seasonal changes in turbulent-water-flow-induced seismic power, P are observed within the 3–10 Hz frequency range, in which P is higher by more than two orders of magnitude during the melt season (mid-May to September) compared to winter (Fig. E1). Changes in P are also observed within the 10–20 Hz frequency range, with P during the melt season being about an order of magnitude larger than in winter (Fig. E1).




Figure E1. Spectrogram of the observed seismic power P as a function of time on the x -axis and frequency on the y -axis. ~~Colours~~ Colors represent seismic power on a decimal logarithmic scale. White bands are data gaps. The two black lines delineate the frequency band between 3 and 10 Hz that we used in this study.

Appendix F: Metrics used in the classification

Figure F1 provides a visual representation of the metrics used to classify the events.




Figure F1. Description of the metrics used to classify the events. (a) Description of one event with the rising and falling part of the runoff to calculate the hysteresis θ . (b) The linear fit used to calculate the residual sum of squares RSS and its slope, m .

Appendix G: **Band-pass-filtered** Bandpass-filtered time series

990 We have filtered the original time series to determine the different frequency signatures separately. For the seasonal scale, we have applied a low-pass filter with a cutoff at 20 days (Fig. G1 a). For the multi-day scale, we have applied a band-pass filter between four and eight days (Fig. G1 b). For the daily time-scale, we have applied a band-pass filter between six hours and 36 hours (Fig. G1 b).




Figure G1. Band-pass filtered series for (a) seasonal (>20 days), multi-day (four to eight days) and diurnal (6h to 36h) time scales. For the last two series, we have computed the band-pass time series for the melt season time period only.

Appendix H: Frequency content of the time series between 6h and 10 days

995 Figure H1 shows the frequency content of all variables over the recording period between a period of 6h and a period of 10 days. While the force and the water pressure have only episodic frequency content on the scale of one day, the other variables exhibit longer periods with variations on that time scale, but still concentrated during the melting season.




Figure H1. Frequency content intensity for all the variables between 6h and 10 days over the year. The red lines delineate the diurnal frequency content (between 6h and 36h).