

1 **Brief Communication: An Ice-Debris Avalanche in the Nupchu Valley, Kanchenjunga Conservation**
2 **Area, Eastern Nepal**

3
4 **Alton C. Byers,¹ Marcelo Somos-Valenzuela,² Dan H. Shugar,³ Daniel McGrath,⁴ Mohan B. Chand,⁵**
5 **Ram Avtar⁶**

6
7 ¹Institute of Arctic and Alpine Research, University of Colorado at Boulder, Boulder, Colorado USA
8 80309

9 ²Department of Forest Sciences, Faculty of Agriculture and Environmental Sciences, University of La
10 Frontera, Temuco, Chile, 4780000

11 ³Water, Sediment, Hazards, and Earth-surface Dynamics (waterSHED) Lab, Department of Geoscience,
12 University of Calgary, 2500 University Drive NW, Calgary, Alberta, T2N 1N4
13 Canada

14 ⁴Department of Geosciences, Colorado State University, Fort Collins, CO, USA, 80523

15 ⁵Department of Environmental Sciences, Patan Multiple Campus, Tribhuvan University, Lalitpur, Nepal

16 ⁶Faculty of Environmental Earth Science, Hokkaido University, Sapporo Japan 060-0810

17
18 **Correspondence:** Alton C. Byers (alton.byers@colorado.edu)

19
20 **Abstract** Beginning in December 2020, a series of small-to-medium, torrent-like pulses commenced
21 upon a historic debris cone located within the Nupchu valley, Kanchenjunga Conservation Area (KCA),
22 Nepal. Sometime between 16 and 21 August 2022 a comparatively large ice-debris avalanche event
23 occurred, covering an area of 0.6 km² with a total estimated volume of order 10⁶ m³. The area of the
24 debris cone left by the August 2022 event increased the historic debris cone area by 0.2 km² (total
25 area: 0.6 km²). Although no human or livestock deaths occurred, the increase in torrent-like pulses of
26 debris upon this historic debris cone since 2020 exemplifies a style of mass movement that may
27 become increasingly common as air temperatures rise in the region. Although the magnitude of this
28 event was small compared to events like the 2021 Chamoli avalanche, the widespread distribution and
29 frequency of comparable events presents a substantial, and potentially increasing, hazard across High
30 Mountain Asia.

31
32
33 **1 Introduction**

34
35 Large magnitude but low frequency events in the high mountains can include a variety of familiar and
36 poorly understood cryospheric processes, including glacial lake outburst floods (GLOFs) (Lamsal et al.
37 2014), snow/ice/rock avalanches (Shugar et al. 2021), landslide-induced avalanches and floods (Byers
38 et al. 2019), englacial conduit floods (Rounce et al. 2017), and others (see: Byers et al. 2022). Today,
39 enhanced communications and remote sensing technologies enable rapid identification and location
40 of such events, often within hours of their occurrence. Many, however, remain unreported because of
41 their remoteness, inaccessibility, poor communications, and/or absence of people (see: Byers et al.
42 2020). In this *Brief Communication*, we report on a large ice-debris avalanche that occurred sometime
43 between 16 and 21 August 2022 in the Kanchenjunga Conservation Area (KCA), eastern Nepal. The
44 event is noteworthy not only because of its probable linkages to climate change impacts in the region,
45 but also because local residents were unaware of its occurrence, as were the Government of Nepal
46 and climate change research entities in Kathmandu. Here we briefly document the event and describe
47 its present and future implications for local communities, scientists, and governments.




Figure 1. Kanchenjunga Conservation Area and location of the Nupchu ice-debris avalanche.

2 Setting

The KCA is a 2,035 km² protected area established in 1997 by the Nepalese Department of National Parks and Wildlife Conservation, with management responsibility handed over to local communities in 2006 (WWF Nepal 2018) (Figure 1). It is home to a range of ethnic groups of primarily Tibeto-Burman origin that include Limbu, Rai, Tamang, Gurung, Magar, Chhetri, and Sherpa (Thapa 2009). Livelihoods were traditionally based upon agriculture, livestock raising, and trade with Tibet, but globalization, outmigration, and new road construction over the past 15 years has rapidly changed the character of both the social and environmental landscape. The South Asian monsoon dominates weather patterns, with most rainfall falling between June and September (Kandel et al. 2019).

Based upon an analysis of 1962–2000 satellite imagery, valley and mountain glaciers in the KCA cover approximately 488 ± 29 km² and exhibit an overall negative glacier surface area loss of 0.5 ± 0.2% yr⁻¹

65 (Racoviteanu et al. 2015). Valley glaciers are largely debris-covered and have been receding since the
66 most recent maximum during the Little Ice Age. Hooker (1854), for example, wrote in 1849 of
67 observing glacial moraines that provided proof “...of glaciers having once descended to from 8,000 to
68 10,000 feet in every Sikkim and east Nepal valley...” (Hooker 1854: 166). The British alpinist Freshfield
69 (1903: 236) writes of the “glacial shrinkage” he encountered in the Lhonak region in 1899, as well as
70 throughout both the Nepal and Sikkim sides of the Kanchenjunga massif. Although the Kanchenjunga
71 region received some of the earliest study, exploration, and mountaineering expeditions in Nepal by
72 outsiders (Thapa 2009), relatively little glacier and cryospheric hazards research has been conducted
73 to date. For example, until 2019 only one GLOF event was on record for the region (Watanabe et al.
74 1998; ICIMOD 2011), although subsequent research revealed that at least seven others had occurred
75 since 1921 (Byers et al. 2020). The Nupchu valley, where the ice-debris avalanche of concern occurred,
76 is used seasonally for yak herding, potato farming, and tourism, with four operational tourist lodges in
77 the village of Kampuchen as of the fall of 2022 (Figure 1).

78 3 Methods

79
80 Field-based observations and assessments of Nupchu Pokhari (glacial lake), other nearby lakes, and
81 the ice-debris avalanche were conducted between 1–20 September 2022. Methods included GPS-
82 based route mapping, photography of avalanche features, oral testimony, and literature reviews.
83 Historic (declassified KH-9 Hexagon satellite imagery; see: Maurer et al. 2019; Dehecq et al. 2020) and
84 recent (Planet Dove and SuperDove) satellite imagery revealed the sequence of avalanche/debris flow
85 events between 1975 and 2023 (Figure 2). Numerical simulations of the avalanche were conducted
86 using R.Avaflow version 3 (Mergili and Pudasaini 2014–2023; Mergili et al. 2017), a state of the art
87 software that has been used globally to study ice/rock avalanches events (Zhang 2022). Numerical
88 simulations were used to provide upper limit volume estimation of the avalanche, which were
89 constrained by field observations. We used the parameters from Zhang et al. (2022) to produce a
90 single-phase model scenario for three different volumes: 1, 2.5, and 5 million m³. For calibration, we
91 modified the internal friction angle of the mixture to match the extension of the debris left by event.
92 For the terrain elevation, ~~an~~ two DEMs were used: the ALOS PALSAR Radiometric Terrain Corrected
93 high resolution ALOS PALSAR 12.5 m DEM was used (AP_13152_FBD_F0540_RT1) (ASF DAAC 2014)
94 and the High Mountain Asia 8m resolution DEM (Shean, 2017), that was void filled.-

Formatted: Font: (Default) +Body (Calibri), Font color: Auto

Formatted: Font: (Default) +Body (Calibri), Font color: Auto

95 4 The Event

96 The investigation of the Nupchu valley was initiated by local concerns about Nupchu Pokhari
97 (27.790708° N, 87.934275° E) as being one of the most dangerous glacial lakes in terms of a potential
98 GLOF (Figure 1). Periodic, smaller floods from the upper Nupchu valley were reported, and assumed
99 locally to have originated in the Nupchu Pokhari, although no supporting evidence was available. Our
100 field reconnaissance results of Nupchu Pokhari on 12 September 2022, however, suggested that the
101 lake posed a moderate risk of flooding, largely based on the absence of overhanging ice and other
102 potential flood triggers. This assessment corroborates the findings of Rounce et al. (2017) which
103 concluded that the 0.129 km² Nupchu Pokhari presented only a moderate risk of flooding because of
104 (a) no apparent growth between 2000 and 2015, (b) absence of avalanche pathways into the lake (i.e.,
105 in line with the direction of the lake and its outflow), and (c) absence of landslide pathways entering
106 the lake.

107 The August 2022 ice/debris avalanche event was unexpected. Field staff had conducted a
108 reconnaissance of the valley below Nupchu Pokhari in early August 2022 to check out potential
109 camping sites, at which time the upper valley was primarily pastureland. When the field team and A.C.
110 Byers returned in early September, the original path was blocked by massive ice-debris avalanche
111 material (27.774328° N, 87.941064° E) that had clearly occurred at some point in the interim (Figure
112 2). Our team and *dzopkio* (yak-cattle crossbreeds used as pack animals) were nevertheless able to
113 climb up and over the avalanche debris to the upper Nupchu valley, but at the time the source and
114 triggers related to the event remained unknown.

115 The original historic debris cone was found to have covered an area 0.402 km² that had been relatively
116 stable for at least 45 years, based upon the oldest satellite imagery available (i.e., 1975) (Dehecq et al.
117 2020) (Panel A, Figure 2). Time series satellite images revealed the periodic occurrence of surficial
118 debris flows upon this original deposition. That is, beginning in 2020, a series of small-to-medium,
119 torrent-like pulses commenced (Panel C through G, Figure 2), culminating in the relatively large event
120 that occurred sometime between 16 and 21 August 2022 (Panel H, Figure 2). The area of the debris
121 cone left by the August 2022 event increased the original area covered by 0.2 km² (total area: 0.6
122 km²). Of the three different volume estimates tested (1, 2.5, and 5 million m³) using two DEMs and
123 R.Avaflow, an avalanche volume of 1×10^6 m³ using an Alos-Palsar/ALOS PALSAR RTC DEM most
124 consistently matched the extent (red line) and depth of the new debris cone deposited as determined
125 by our field observations (Figure 3).

126
127 Our team was unable to locate the event on any seismographs, most likely related to the absence of
128 instrumentation in this part of the eastern Himalayas. Based upon direct field observations as well as
129 satellite imagery, the avalanche had clearly blocked and temporarily dammed the water from the
130 Nupchu Khola (river) at its onset, which was nevertheless able to cut down through the ice and
131 sediment deposited to form a steep canyon estimated at >10 m depth. The presence of shrubs (e.g., *J.*
132 *indica*) fully stripped of their bark was testimony to the high velocities of the flood- and meltwater
133 produced by frictional forces during the event, a phenomenon reported for other rockfall-induced
134 landslides in Nepal (Byers et al. 2019).


135 136 **5 Discussion**

137
138 Interestingly, individuals in the community of Kampuchen, only 5 km downstream of the event, were
139 unaware of the avalanche. Yak herds had already returned from the high pastures to the village by
140 early August, the community was busy harvesting potatoes and preparing for the fall tourist season,
141 and no obvious changes in the Nupchu Khola had been observed (e.g., Kargel 2014 for a description of
142 changes in the Seti Kosi prior to the catastrophic flooding of 5 May 2012). Thus, authorities in
143 Taplejung and Kathmandu were also unaware of the event as of September 2022, which is typical of
144 many large-scale cryospheric events in remote regions of the Himalayas (e.g., Byers et al. 2022).

145
146 Still, the acceleration of torrent-like pulses of debris upon the historic debris cone since 2020 suggests
147 that these events may be linked to contemporary warming trends, similar to those that may have
148 triggered larger-scale mass wasting events elsewhere in the Himalaya (e.g., Shugar et al. 2021; Kääh et
149 al. 2021; Taylor et al. 2023). The frequency of such ice-debris flow events within the KCA region, and
150 more broadly across the Himalaya, is unknown. However, with projections of continued warming in
151 these regions (e.g., Lalande et al. 2021), a more systematic approach to determining their historic
152 frequency, as well as a better understanding of their triggers, is warranted. After further evaluation,


Formatted: Indent: Left: 0 px

153 vulnerable villages, such as Kampuchen, may wish to consider the installation of preventative
154 floodwater diversion mechanisms, such as the rock-filled gabion walls currently protecting tourist
155 lodges in the Mt. Everest region (e.g., Rounce et al. 2017; Byers et al. 2022) using participatory
156 processes as outlined in Watanabe et al. (2016).




157
158 **Figure 2.** Time series satellite images showing the periodic occurrence of surficial debris flows upon the
159 original deposition. These appear to have accelerated in both frequency in magnitude beginning several years
160 ago, leading up to the main event that occurred between 16 and 21 August 2022. Blue dashed outline in panel

161 H is the 1975 outline of the debris cone, while the dashed black circle identifies the failure zone. The
162 photograph at the bottom provides an oblique view of the ice/debris avalanche about three weeks after it
163 occurred (photograph by A. Byers). Panel H shows an image from early 2023, as imagery from immediately
164 after the ice-avalanche in mid-August 2022 were partially obscured by clouds (KH-9 imagery courtesy of USGS;
165 Planet Dove and SuperDove imagery courtesy of Planet Lab PBC).
166



Formatted: Indent: Hanging: 0.13"



168
 169 **Figure 3.** Base change modeled with R.Avaflow for three different avalanche volumes: a) $1 \times 10^6 \text{ m}^3$ (top) using an
 170 Alos-LOS PalsarALSAR RTC DEM, b) $1 \times 10^6 \text{ m}^3$ using the void filled HMA-DEM, c) $2.5 \times 10^6 \text{ m}^3$ (bottom left) using
 171 an Alos-LOS PalsarALSAR RTC DEM, and d) $5 \times 10^6 \text{ m}^3$ (bottom right) using an Alos PalsarALOS PALSAR RTC DEM.
 172 Of the three estimates and 2 DEMs, $1 \times 10^6 \text{ m}^3$ using an Alos PalsarALOS PALSAR RTC DEM most consistently
 173 matched the extent and depth of the new debris cone deposited in August 2022 (red line; the yellow line
 174 represents the extent of the historic debris cone).
 175

176 5 Conclusion

177
 178 Beginning in 2020, a series of small-to-medium, torrent-like pulses commenced upon a historic debris
 179 cone located approximately 2 km down valley from the lake, culminating in a relatively large
 180 avalanche event that occurred sometime between 16 and 21 August 2022. The August 2022 event
 181 deposited debris with an area of 0.6 km^2 and estimated volume in the order of 10^6 m^3 . No fatalities

Formatted: Indent: Left: 0 px

182 from the event occurred because of the absence of humans and livestock in the vicinity when the
183 event occurred. Likewise, no impoundment of the Nupchu Khola, and formation of a potentially
184 dangerous backwater lake, occurred as a result of debris blockage, although such scenarios happen
185 routinely in high mountain environments.

186
187 The improvement of remote area event reporting mechanisms, especially to authorities in the capital,
188 Kathmandu, could help with the development of hazard mitigation technologies and response.
189 Likewise, more systematic monitoring of cryospheric events by scientists, using remote sensing
190 platforms and hazard mapping tools, could help with the development of more effective early warning
191 systems for vulnerable communities, livestock, and adventure tourists. Ultimately, this could lead to a
192 minimization of losses and damage due to multi-hazard events.

193
194 *Data availability.* Declassified KH-9 Hexagon satellite imagery is available at
195 <https://earthexplorer.usgs.gov/>. Planet Dove and SuperDove satellite imagery is available
196 at <https://www.planet.com/explorer/>. The *ALOS PALSAR Radiometric Terrain Corrected high-res*
197 *DEM/ALOS PALSAR DEM* "AP_13152_FBD_F0540_RT1" used for the R.Avaflow is available
198 at <https://search.asf.alaska.edu/>, <https://asf.alaska.edu/>.

199
200 *Author contributions.* ACB conceived the study and wrote the original narrative, with contributions
201 from MS-V, DS, DM, MBC, and RA. MBC created Figure 1, DS and RA created Figure 2, and MS-V
202 created Figure 3. MS-V conducted the numerical simulations of avalanche volumes shown in Figure 3.
203 All authors revised and contributed to the final manuscript.

204
205 *Competing Interests.* The contact author has declared that none of the authors has any competing
206 interests.

207
208 *Acknowledgements.* The Fulbright Nepal Scholar Program is thanked for its support of A.C. Byers
209 during his six-month field study of contemporary impacts on alpine ecosystems in the Kanchenjunga
210 Conservation Area, eastern Nepal. The Department of National Parks and Wildlife Conservation,
211 Kanchenjunga Conservation Area, Department of Geography at Tribhuvan University are also thanked
212 for their interest in and support of the project. Support for M. Somos-Valenzuela during the
213 preparation of this paper was provided by the Chilean Science Council (ANID) through the Program of
214 International Cooperation (PII-180008). Support for D.H. Shugar was provided by the Natural Sciences
215 and Engineering Research Council of Canada (DG-2020-04207) and Alberta Innovates. Support for D.
216 McGrath was provided by NASA award 80NSSC20K1343.

217 218 219 **References**

220
221 ASF DAAC 2014. ALOS PALSAR Radiometric Terrain Corrected high res; Includes Material ©
222 JAXA/METI; 2008 Accessed through <https://search.asf.alaska.edu/> <https://asf.alaska.edu/> on 11
223 February 2023, 2008. <https://doi.org/10.5067/10.5067/Z97HFCNKR6VA>

224
225 Byers, A.C., Shugar, D., Chand, M., Portocarrero, C., Shrestha, M., Rounce, D., and Watanabe, T.: Three
226 recent and lesser-known glacier-related flood mechanisms in high mountain environments.
227 Mountain Research and Development, 42(2):A12-A22, 2022. [https://doi.org/10.1659/MRD-](https://doi.org/10.1659/MRD-JOURNAL-D-21-00045.1)
228 [JOURNAL-D-21-00045.1](https://doi.org/10.1659/MRD-JOURNAL-D-21-00045.1)

229

Formatted: Font: (Default) +Body (Calibri), 11 pt

Formatted: Font: (Default) +Body (Calibri), 11 pt

Formatted: English (United States)

Commented [M1]: This DOI doesn't appear to link to the dataset.

230 Byers, A.C., Chand, M.B., Lala, J., Shrestha, M., Byers, E.A., and Watanabe, T. 2020. Reconstructing the
231 history of glacial lake outburst floods (GLOF) in the Kanchenjunga Conservation Area, east Nepal: an
232 interdisciplinary approach. *Sustainability* 2020, 12, 5407. [https://www.mdpi.com/2071-](https://www.mdpi.com/2071-1050/12/13/5407)
233 [1050/12/13/5407](https://www.mdpi.com/2071-1050/12/13/5407)
234
235 Byers, A.C., Rounce, D.R., Shugar, D.H. and Regmi, D.: A rockfall-induced glacial lake outburst flood,
236 upper Barun valley, Nepal. *Landslides* (2019) 16: 533, 2019. [https://doi.org/10.1007/s10346-018-](https://doi.org/10.1007/s10346-018-1079-9)
237 [1079-9](https://doi.org/10.1007/s10346-018-1079-9)
238
239 Dehecq, A., Gardner, A.S., Alexandrov, O., McMichael, S., Hugonnet, R., Shean, D., and Marty, M.:
240 Automated processing of declassified KH-9 Hexagon satellite images for global elevation change
241 analysis since the 1970s, *Frontiers Earth Science*, Vol. 8, 09 November, 2020.
242 <https://doi.org/10.3389/feart.2020.566802>
243
244 Freshfield, D.W.: Round Kangchenjunga: A Narrative of Mountain Travel and Exploration (historic
245 reprint), 1903. www.forgottenbooks.com
246
247 Hooker, J.D.: Himalayan Journals Vol. 1, 1854, Orlando: The Perfect Library (historic reprint).
248
249 Kääh, A., Jacquemart, M. n., Gilbert, A., Leinss, S., Girod, L., Huggel, C., Falaschi, D., Ugalde, F.,
250 Petrakov, D., Chernomorets, S., Dokukin, M., Paul, F., Gascoin, S., Berthier, E. and Kargel, J.: Sudden
251 large-volume detachments of low-angle mountain glaciers: more frequent than thought?
252 *Cryosphere* 15(4), 1751–1785, 2021. <https://doi:10.5194/tc-15-1751-2021>
253
254 Kandel, P., Chettri, N., Chaudhary, R.P., Badola, H.M., Gaira, K.S., Wangchuk, S., Bidha, N., Uprety, Y.,
255 Sharma, E.: Plant Diversity of the Kangchenjunga Landscape, Eastern Himalayas. *Plant Diversity*,
256 Volume 41, Issue 3, 2019, pp. 153-165, 2019. <https://doi.org/10.1016/j.pld.2019.04.006>
257
258 Kargel J.: One scientist’s search for the causes of the deadly Seti River flash flood. *Earth Observatory*,
259 24 January, 2014.
260 <https://earthobservatory.nasa.gov/blogs/fromthefield/2014/01/24/setiriverclues/>; accessed on 15
261 December 2022.
262
263 Lalande, M., Ménégos, M., Krinner, G., Naegeli, K., and Wunderle, S.: Climate change in the High
264 Mountain Asia in CMIP6, *Earth Syst. Dynam.*, 12, 1061–1098, [https://doi.org/10.5194/esd-12-1061-](https://doi.org/10.5194/esd-12-1061-2021)
265 [2021](https://doi.org/10.5194/esd-12-1061-2021), 2021.
266
267 Maurer, J.M., Schaeffer, J.M., Rupper, S., and Corley, A.: Acceleration of ice loss across the Himalayas
268 over the past 40 years. *Science Advances*, Vol 5, Issue 6, 2019. [https://doi: 10.1126/sciadv.aav7266](https://doi:10.1126/sciadv.aav7266)
269
270 Mergili, M., Fischer, J. T., Krenn, J. y Pudasaini, S. P.: R.avafLOW v1, an advanced open-source
271 computational framework for the propagation and interaction of two-phase mass flows, *Geosci.*
272 *Model Dev.*, 10(2), 553–569, 2017. <https://doi:10.5194/gmd-10-553-2017>.
273
274 Mergili, M. and Pudasaini, S.P., 2014–2023. r.avafLOW - The Mass Flow Simulation Tool.
275 <https://www.avafLOW.org>
276

277 Racoviteanu, A.E., Arnaud, Y., Williams, M.W., and Manley, W.F.: Spatial patterns in glacier
 278 characteristics and area changes from 1962 to 2006 in the Kanchenjunga–Sikkim area, eastern
 279 Himalaya. *The Cryosphere*, 9, 505–523, 2015. www.the-cryosphere.net/9/505/2015/
 280 doi:10.5194/tc-9-505-2015.

281

282 Shean, D. (2017). High Mountain Asia 8-meter DEMs Derived from Cross-track Optical Imagery.
 283 Boulder, CO: NASA NSIDC DAAC: NASA National Snow and Ice Data Center Distributed Active Archive
 284 Center. <https://doi.org/10.5067/0MCWJJH5ABYO>

285

286 Taylor, C., Robinson, T.R., Dunning, S., Carr, R. and Westoby, M.: Glacial lake outburst floods threaten
 287 millions globally. *Nature Communications* 14, 487 (2023). [https://doi.org/10.1038/s41467-023-](https://doi.org/10.1038/s41467-023-36033-x)
 288 [36033-x](https://doi.org/10.1038/s41467-023-36033-x)

289

290 Thapa, R: *Kanchenjunga: The Unique Gift of Nature*. Kathmandu: Kanchan Printing Press, published by
 291 Phupu Chowang Sherpa, 2009.

292

293 Watanabe, T., Khanal, N.R., and Gautam, M.P.: The Nangama glacial lake outburst flood occurred on
 294 23 June 1980 in the Kanchanjunga area, eastern Nepal. *Ann. Hokkaido Geogr. Soc.*, 72, 13–20, 1998.
 295 DOI:10.14917/hgs1959.1998.13.

296

297 Watanabe, T., A. C. Byers, M, A. Somos-Valenzuela and D. C. McKinney: The need for community
 298 involvement in glacial lake field research: the case of Imja Glacial Lake, Khumbu, Nepal Himalaya.
 299 Chapter 13. In: Singh, R. B., Schickhoff, U., and Mal, S. (eds.): *Climate Change, Glacier Response, and*
 300 *Vegetation Dynamics in the Himalaya: Contributions Toward Future Earth Initiatives*, Springer
 301 International Publishing Switzerland, pp. 235-250, 2016, doi:10.1007/978-3-319-28977-9_13

302

303 Zhang, T., Wang, W., Shen, Z., Zhan, N., Wang, Z. and An, B.: Understanding the 2004 glacier
 304 detachment in the Amney Machen Mountains, northeastern Tibetan Plateau, via multi-phase
 305 modeling. *Landslides*, (October), 2022. <https://doi.org/10.1007/s10346-022-01989-2>.

Formatted: Font: (Default) +Body (Calibri), 11 pt, Font color: Custom Color(RGB(34,34,34)), Pattern: Clear (White)

Commented [M2]: I believe this needs date accessed at the end of the citation.

Formatted: Hyperlink, Font: (Default) +Body (Calibri), 11 pt, Pattern: Clear (White)

Field Code Changed

Formatted: Font color: Custom Color(RGB(34,34,34))

Formatted: List Paragraph