

Opinion: The strength of long-term comprehensive observations to meet multiple grand challenges at different environments and in the atmosphere

Markku Kulmala^{1,2,3}, Anna Lintunen^{1,4}, Hanna Lappalainen¹, Annele Virtanen⁵, Chao Yan^{1,2,3}, Ekaterina Ezhova¹, Tuomo Nieminen^{1,6}, Ilona Riipinen⁷, Risto Makkonen^{1,8}, Johanna Tamminen⁹, Anu-Maija Sundström⁹, Antti Arola⁸, Armin Hansel¹⁰, Kari Lehtinen⁵, Timo Vesala¹, Tuukka Petäjä¹, Jaana Bäck⁴, Tom Kokkonen¹ and Veli-Matti Kerminen¹

¹Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, Finland

²Joint International Research Laboratory of Atmospheric and Earth System Sciences, School of Atmospheric Sciences, Nanjing University, Nanjing, China

³Aerosol and Haze Laboratory, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing, China

⁴Institute for Atmospheric and Earth System Research / Forest Sciences, Faculty of Agriculture and Forestry, University of Helsinki, Finland

⁵Department of Technical Physics, University of Eastern Finland, Kuopio, Finland

⁶Department of Physics, University of Helsinki, Finland

⁷Bolin Centre for Climate Research, Department of Environmental Science (ACES), Stockholm University, Sweden

⁸Climate Research Programme, Finnish Meteorological Institute, Helsinki, Finland

⁹Space and Earth Observation Centre, Finnish Meteorological Institute, Helsinki, Finland

¹⁰Institute for Ion and Applied Physics, University of Innsbruck, Austria

Correspondence to: Markku Kulmala (markku.kulmala@helsinki.fi)

Abstract. To be able to meet global grand challenges (climate change; biodiversity loss; environmental pollution; scarcity of water, food and energy supplies; acidification; deforestation; chemicalization; pandemics), which all are closely interlinked with each other, we need comprehensive open data with proper metadata, along with open science. The large data sets from ground-base in situ observations, ground and satellite remote sensing and multiscale modelling need to be utilized seamlessly.

In this opinion paper, we demonstrate the power of the SMEAR (Station for Measuring Earth surface – Atmosphere Relations) concept via several examples, such as detection of new particle formation and their subsequent growth, quantifying atmosphere-ecosystem feedback loops, combining comprehensive observations with emergency science and services, as well as studying the effect of COVID restrictions on different air quality and climate variables. The future needs and the potential of comprehensive observations of the environment are summarized.

1 Background

The Earth is facing several environmental challenges on a global scale, often called “Grand Challenges” (<https://www.wcrp-climate.org/grand-challenges/>). The growing population (<https://pdp.unfpa.org>) needs fresh air, fresh water, food and energy,

while at the same time climate is changing, many cities have challenges with air quality, biodiversity is decreasing and supplies of fresh water, food and energy are diminishing. Since these Grand Challenges are highly connected and interlinked, not only with each other but also with e.g. pandemics, they cannot be solved separately since potential solutions are tightly coupled with each other (e.g. Kulmala et al., 2015; Lappalainen et al., 2016; Nolan et al., 2018; Laughner et al., 2021; Wang et al., 2023). However, the solutions may also include unexpected trade-offs (e.g. Fastre et al., 2020). Therefore, integrated, comprehensive, big open data are required (Kulmala et al., 2021), together with a research and innovation framework in which 35 a multidisciplinary research with critical mass of scientists utilising proper resources is connected to fast-tracked policy making and wide stakeholder community. This allows aiming for practical solutions based on deep scientific understanding.

The global challenges are intimately linked to interactions and feedbacks between the different compartments of the planet Earth at different spatial and temporal scales. Fundamentally, the atmosphere is closely interconnected with various other parts 40 of the Earth, including biosphere, hydrosphere, cryosphere and lithosphere as well as urban surfaces over a range of time and spatial scales varying from seconds to millennia (Wanner et al., 2008). The sources, sinks and atmospheric concentrations of reactive trace gases, greenhouse gases and aerosol particles depend strongly on each other via physical, chemical and biological processes (e.g. Arneth et al., 2010, Stocker et al., 2013; Kulmala et al., 2014a; Unger, 2014, Green et al., 2017; Smith et al., 2023). Furthermore, both human actions and natural feedback mechanisms between the biosphere and atmosphere have 45 substantial impacts on interactions between these atmospheric constituents and their influences on air quality and climate (Raes et al., 2010; Stocker et al., 2013; Kulmala et al., 2015; Kulmala, 2015; Nolan et al., 2018; Doherty et al., 2022; Wang et al., 2023). The importance of atmospheric aerosol particles on climate and human health in both regional and global scales has 50 attracted a plenty of research interest during the recent years (Butt et al., 2017; Boy et al., 2019; Bellouin et al., 2020; Lappalainen et al., 2022; Lintunen et al., 2023). Despite these efforts, atmospheric aerosol particles remain perhaps the least 55 known factor influencing radiative forcing, causing thereby large uncertainties in predicting the future behavior of the climate system (IPCC 2013, 2021).

The global annual cost of climate change impacts are estimated to reach hundreds of billions of euros by 2030 (UNEP, 2016; Köberle et al., 2021) and, with an increasing global warming, this cost is expected to increase strongly in the future. According 60 to the World Economic Forum, climate change has cost the EU €145 billion in a decade (<https://www.weforum.org/agenda/2022/12/climate-europe-gdp-emissions/>). There is an urgent need for improving climate projections, reducing greenhouse gas emissions and developing options to sequester terrestrial carbon by simultaneously taking 65 into account the other climate forcers, such as atmospheric aerosol particles and trace gases. In addition, in order to have better understandings of natural and anthropogenic sources and sinks of carbon and of atmospheric processes influencing air quality, we need to develop existing monitoring and forecasting systems of the terrestrial carbon cycle, and to both enhance and improve measurements from process levels to a global scale. These practical needs provide emerging business opportunities for various industries. For example, European Green Deal Investment will mobilize at least one trillion euros of sustainable

investments over the next decade (COM 2020). The information produced by using new verification systems is essential for society to design economically and socially optimal sustainability strategies and climate-neutrality pathways and to be able to meet Paris Agreement targets (Kriegler et al., 2018). The importance of comprehensive and standardized measurements is also underlined by rapid development of the environmental data analysis based on artificial intelligence (AI). The quality of the AI based environmental analysis is as good as the measured source data. The novel Earth Virtualization Engines (EVE) concept (<https://eve4climate.org/>) in under development and the foreseen digital infrastructure of multi-tiered climate information for various type of users would build on optimal earth-system data integration and monitoring by using AI methods like machine learning (Stevens et al. 2023).

Open science and open data are essential. Open science, the sharing of knowledge and data as early as possible in the research process (Vicente-Saez and Martinez-Fuentes, 2018) are essential as addressed by bodies like European Commission (European Commission, 2021). Such *in-situ* data can be obtained from global networks, such as GAW (Global Atmospheric Watch) and FluxNet (Smith et al. 2012; Baldocci, 2019), and continental-scale infrastructures such as TERN (Terrestrial Ecosystem Research Network in Australia), NEON (National Ecological Observatory Network in US), ASCENT (Atmospheric Science and Chemistry mEasurement NeTwork in US), AmeriFlux (Boden et al. 2016), ChinaFlux (Yu et al. 2006) AsiaFlux (Mizoguchi et al. 2008), AfriFlux (Ciais et al. 2011), and the European RI's ICOS (Integrated Carbon Observation System), ACTRIS (Aerosols, Clouds, and Trace gases Research Infrastructure), and eLTER (Integrated European Long-Term Ecosystem, Critical Zone & Socio-Ecological Research Infrastructure) (Loescher et al., 2022). However, often the *in -situ* data is covering only some components of the Earth system and the proper integration of different datasets is lacking. This can be overcome by co-locating different measurements, which enables new knowledge of the interactions and feedbacks between the Earth components (biosphere, hydrosphere, atmosphere, and geosphere), and allows for science-based solutions related to the interlinked Grand Challenges. An example of co-located measurements at a single station enabling new knowledge of the interactions and feedbacks between the Earth components, spearheading science-based solutions related to the interlinked Grand Challenges, is Station(s) for Measuring Earth surface – Atmosphere Relations (SMEAR).

The primary objective of this paper is, by using a few examples based mainly on data from the SMEAR II station located in Hyytiälä, Finland, to demonstrate the power of comprehensive, continuous and integrated long-term observations in a way toward addressing some of the Grand Challenge discussed above. This kind of an approach is closely tied with the SMEAR concept introduced earlier and summarized briefly in the next section. Besides research associated with the Grand Challenging, we demonstrate the strength of the SMEAR concept when rapid and unexpected changes occur in the environment and atmosphere.

100 **2 SMEAR concept**

The SMEAR concept is based on comprehensive, continuous, and integrated long-term observations (Hari and Kulmala, 2005; Hari et al., 2016). It has been developed to meet Environmental Grand Challenges and to collect big open data sets in order to test theories and develop models at the interfaces of different Earth components. Such unique environmental open data can contribute to solving burning questions of society – even questions that are currently unforeseen.

105

Current observations (see IPCC 2013, 2021) are fragmented, which means that typically different infrastructures are measuring greenhouse gases, aerosols, air quality, ecosystems, climate and biodiversity. These measurements are conducted in different locations and environments, and often during relatively short campaigns. However, to meet the ongoing environmental challenges, an integrated approach with long-term measurements is needed. In practice this means co-location of various
110 infrastructures, which enables simultaneous measurements of different Earth components (Guo, 2018; Kulmala et al., 2021; Lintunen et al., 2023). Changes in one of these components are directly or indirectly communicated to the others via intricately linked processes and feedbacks occurring at their interfaces (e.g. Stocker et al., 2013; Nolan et al., 2018; Smith et al., 2023).

115

The SMEAR stations, together with the SMEAR concept, were established before the various international environmental research infrastructures were established (Hari and Kulmala, 2005). Today the four Finnish SMEAR stations and international
120 SMEAR-like stations (Kulmala et al., 2021) are contributing to the European ESFRIs focused on standardized measurements on atmospheric composition and ecosystem processes like ICOS, ACTRIS, eLTER and AnaEE. SMEAR II is participating to several global Earth Observation systems and networks such as WMO GAW and/or EMEP, GEO-GEOSS, FluxNet, AERONET and SolRad-Net. The key aspect is the co-location of these different thematic networks into a single site. This allows multi- and interdisciplinarity open science based on the collected data.

125

Crucial in the SMEAR concept is that it measures a set of variables needed for the atmosphere-Earth surface interactions and feedback analysis, providing fundamental knowledge for the climate change mitigation and adaptation plans. The Earth surface can be, for example forest, lake, ocean, peatland, urban area, glacier, agricultural land. The most established station, SMEAR II, is located in Hyytiälä, Finland, and it includes measurements of over 1200 different variables (Kulmala et al., 2021). The measurements are conducted at different scales from small chamber enclosures to a regional scale, which is made possible by the 128-m-high measurement tower at the SMEAR II station. The measurements include meteorological variables,
130 atmospheric compositions and fluxes (aerosols, clouds, atmospheric chemistry, greenhouse gases etc.), as well as variables describing ecosystem functioning and soil dynamics. Long-term *in situ* measurements are accompanied by remote sensing, experiments (both lab and field) and multi-scale modelling. Such an approach enables us to track the regional and long-range

transboundary pollution transport and to evaluate e.g. trends in measured concentrations and fluxes, process dynamics, and feedbacks between processes and Earth components, such as soil-forest-atmosphere and forest-soil-streams-lake.

135

An important part of the SMEAR concept is open access to the research infrastructure and open data (<https://smear.avaa.csc.fi/>). These data are massive and heterogeneous, and thus challenging to manage, but the easy access to these data and both harmonised and standardised ways to analyse it are important (Junninen et al., 2009). As a summary, we can state that we need to work towards an open, integrated approach that can be accomplished with a global SMEAR network,
140 a global Earth observatory (Hari et al., 2016; Kulmala 2018).

In the following section, we give examples of capabilities of the SMEAR concept addressing different kinds of research questions: atmospheric new particle formation, COBACC feedback loop that combines terrestrial carbon sink to aerosol source, COVID impacts on air quality, and long-term trends in some of the quantities essential for air quality and climate
145 research. We also discuss the future role of comprehensive measurements to detect unexpected changes in the environment and atmosphere.

3 The utilization of comprehensive data sets

Here we provide examples on how we have used comprehensive data sets to meet several scientific and societal challenges and discuss briefly what has been learnt from these investigations.

150 3.1 Atmospheric new particle formation (NPF)

A well-known example of the usefulness of comprehensive, long-term observations are investigations related to atmospheric NPF (e.g., Mäkelä et al., 1997; Kulmala et al., 2013). Before such observations, more emphasis was placed on binary nucleation in the stratospheric conditions (e.g, Hamill et al., 1982). When looking at the scientific literature published prior to mid-1990's, the common thought appears to have been that in the troposphere, NPF is a relatively rare and local phenomenon with minor
155 contributions to regional or global aerosol particle budgets. However, the first long-term observations of particle number concentrations revealed a frequent occurrence of NPF and its regional character in a boreal forest environment (Mäkelä et al., 1997), and later observations confirmed the same to be the case in many other types of atmospheric environments (e.g. Kerminen et al., 2018; Nieminen et al., 2018; Chu et al., 2019; Brean et al., 2023). Motivated by long-term observations, explicit description of NPF was then included in several large-scale modeling frameworks. Simulations using such models
160 demonstrated that NPF is the dominant source of the particle number concentration in the global atmosphere, and an important contributor to concentrations of cloud condensation nuclei (e.g. Merikanto et al., 2009; Gordon et al., 2017).

In order to understand how atmospheric NPF is connected with climate and air pollution, or emissions to the atmosphere, one needs to quantify the mechanistic pathways of atmospheric NPF and its basic characteristics, such as its frequency and 165 associated particle formation and growth rates. Our mechanistic understanding on the initial steps of atmospheric NPF, clustering, relied for a long time on theories and laboratory experiments, and with a belief that the only important clustering mechanisms in the atmosphere is the binary nucleation between sulfuric acid and water vapors (Malila, 2018). Atmospheric observations provided increasing evidences on the existence of multiple and possibly more complex clustering pathways (e.g. Kulmala et al., 2014b). Such findings inspired comprehensive and dedicated laboratory experiments in the CLOUD chamber 170 at CERN (e.g. Kirkby et al., 2011; Almeida et al., 2013; Lehtipalo et al., 2018, He et al., 2021). Many of the clustering pathways quantified in these CLOUD experiments have recently been found in atmospheric observations (e.g. Sipilä et al., 2016; Jokinen et al., 2018; Lehtipalo et al., 2018; Beck et al., 2021; Yan et al., 2021), confirming the diversity of NPF in various atmospheric environments.

175 Atmospheric observations have revealed large differences in the NPF characteristics between different sites, as well as between different seasons at individual sites (e.g. Nieminen et al., 2018; Chu et al., 2019; Deng et al., 2020, Brean et al., 2023). At sites with multi-year observations, there appears to be a notable inter-annual variability in both frequency and intensity of NPF, and in some cases also a long-term trend has been reported (Asmi et al., 2011; Nieminen et al., 2014; Saha et al., 2018; Kalivitis et al., 2019; Neefjes et al., 2022). Figure 1 shows the monthly medians of nucleation mode particle concentrations measured 180 at the SMEAR II station. Over the 27-year observation period from 1996 until 2022, the monthly median concentrations decreased at a rate of $-0.9\%/\text{year}$. The temporal variability in NPF characteristics has been ascribed to changes in meteorological conditions and aerosol precursor sources, including clear influences of reduced sulfur emissions and other air pollution control actions in Europe and North America (Hamed et al., 2010; Kyrö et al., 2014; Wang et al., 2017; Saha et al., 2018) and more recently also in China (Zhao et al., 2021; Zhu et al., 2021).

185




Figure 1: Monthly medians of the nucleation mode particle (3–25 nm in the mobility diameter) concentrations at the SMEAR II station in Hyytiälä, Finland. The line is a linear lin-log fit to the data and shows clearly the decreasing trend.

190

Long-term atmospheric observations have played a central role in atmospheric model development. The first semi-empirical parameterizations of new particle formation rates for modeling purposes were based on simultaneous and continuous measurements of gas-phase sulfuric acid concentrations and particle number size distributions (Sihto et al., 2006; Paasonen et al., 2010; Semeniuk and Dastoor, 2018). Later, long-term observations have been essential in testing the performance of large-scale models in simulating NPF and subsequent growth of newly formed particles to CCN (e.g. Spracklen et al., 2010; Fountoukis et al., 2012; Yu et al., 2015; Qi et al., 2018).

While models are likely to be the main tool for estimating the future impacts of atmospheric NPF on climate and air quality, they regularly need observations to verify their performance. In addition, many related scientific issues remain that cannot be solved without comprehensive and continuous observations. One of them is the relative importance of different clustering pathways in different environments and due to continually changing atmospheric composition in these environments. The second issue is the quantification of factors dictating the frequency and intensity of NPF, including the role of “quiet NPF”, i.e. relatively weak NPF not captured by traditional NPF event analysis methods (Kulmala et al., 2022a). The third issue is to

205 understand the growth of newly formed particles into sizes where they may act as CCN or contribute to haze formation (e.g. Ren et al., 2021; Kulmala et. 2022b; Stolzenburg et al., 2023). Related to this issue, we need long-term observations to better understand how small clusters survive while growing larger sizes, especially in polluted environments (Kulmala et al., 2017; Tuovinen et al., 2022), in order to find out the relative importance of condensation and heterogeneous reactions in growth, and to quantify the most important precursor vapors causing this growth.

210 **3.2 COBACC feedback loop**

To understand better the complex feedbacks between the atmosphere and ecosystems, we have developed a concept called the Continental Biosphere-Atmosphere-Clouds-Climate (COBACC) feedback loop (Kulmala et al., 2013). It utilises a multidisciplinary-and integrated approach to quantify the feedbacks. The loop consists of several interrelated processes (for detailed description see Kulmala et al., 2014a; Artaxo et al., 2022, Kulmala et al., 2023): 1) the atmospheric temperature and
215 CO₂ influences on biogenic volatile organic compound (BVOC) emissions, 2) the influence of BVOCs on the formation and growth of aerosol particles, 3) the effect of aerosol particles on clouds, 4) the effect of aerosol particles and clouds on solar radiation, in particular, on its diffuse fraction, and 5) the link between diffuse radiation and photosynthesis, and carbon sink in general.

220 The various processes involved in the interactions within the COBACC feedback loop occur over different time scales. Increases in atmospheric temperatures and carbon dioxide concentrations occur at inter-annual time scales, carbon cycling including photosynthesis and emission of BVOCs vary on scales from sub-hourly to seasonal, while the cloud variability and its effect on radiation that drive photosynthesis operate at sub-hourly time scales. Therefore, continuous and comprehensive observations, together with modelling, are key to solving such complex questions. In what follows, we summarize our current
225 understanding of the feedback loop in the boreal zone (for a more comprehensive review see Artaxo et al., 2022), and indicate future directions. Continuous observations serve as a base for most of the studies cited below, and the SMEAR II data set, used in most of them, remains the most comprehensive one up to date.

Paasonen et al. (2013) considered the effect of warming climate on BVOC emissions and associated increases in >100 nm
230 aerosol particle number concentrations (a proxy for CCN), quantifying the potential cooling effect due to this feedback. The pilot study of Kulmala et al. (2014a) made the first estimate of the COBACC feedback loop using SMEAR II data, focusing on the direct aerosol effect and excluding clouds from the consideration. After that, Ezhova et al. (2018) refined and extended the analysis of the aerosol-diffuse radiation-photosynthesis part of the feedback loop using data from five sites in the boreal zone, also excluding clouds. Clouds were included in the Earth System Model (ESM) studies on the feedback loop (Rap et al.,
235 2018; Sporre et al., 2019). However, the link between BVOC, aerosol particles and clouds in various ESMs is a source of substantial discrepancies, even of different sign, in the radiation – the main driver of photosynthesis (Sporre et al., 2020). Therefore, observations remain an extremely relevant source of data for this complex question.

Based on the COBACC feedback loop, Kulmala et al. (2020) developed the CarbonSink+ concept, which, beside aerosols, takes into account the effect of forest on clouds and surface albedo. The next step is to include all radiative forcers. Current COBACC feedback loop studies are directed towards quantifying the role of clouds (Fig. 2), including their interaction with the surface-based parameters and their effects on radiation and photosynthesis, based on observations. The combination of on-site and satellite observations was employed to show that clouds become optically thicker in a warmer climate with larger amounts of organic aerosol particles (Yli-Juuti et al., 2021). Furthermore, Petäjä et al. (2022) showed that continuous interaction of an air mass with emissions from the boreal forest changes the properties of this air mass over a time period of several days, including both aerosol physical and chemical characteristics and humidity. Both factors are important for the formation and evolution of clouds. Räty et al. (2023) extended this approach to a data set covering more than a decade and confirmed the main conclusions showing an increase in, e.g., cloud condensation nuclei, specific humidity as well as cloud optical thickness and precipitation frequency (Fig. 2).

250




Figure 2: Illustration of forest-boundary layer clouds link (Räty et al., 2023): fraction of air masses with the parameter value above its median after 10-30 h and 50-75 h interaction with boreal forest. Parameters are cloud condensation nuclei at 0.2% supersaturation (CCN, median value 180 cm^{-3}), specific humidity (q , median value 5 g kg^{-1}), cloud optical thickness (COT, median value 11). Note also an increase in precipitation frequency from 7% to 12% (P_{freq}). Results are obtained from 11-years data set featuring growing seasons, SMEAR II/MODIS.

However, the outcome from this study regarding the effect of forest on cloud properties remains somewhat obscure: cloud properties were taken from the satellite data sets, which drastically decreases the number of data available for analysis. To overcome this problem, e.g. the cloud classification algorithm by Ylivinkka et al. (2020) can be used. The algorithm allows quantifying some cloud properties, e.g. optical thickness for some types of clouds, whereas cloud fraction is linked to patchiness. The radiation measurements, an input parameter for this algorithm, have been measured at SMEAR II for more

than two decades, and therefore the cloud-related data set can potentially be extended significantly. Overall, continuous comprehensive observations play a key role in tackling multidisciplinary problems with multiple time scales.

265 While the COBACC feedback has, until now, been studied primarily in the boreal ecosystem, more data to constrain similar feedbacks within other ecosystems – particularly tropical and (semi)arid as well as urban – are urgently needed.

BVOC and semi volatile organic compounds (SVOC) are closely linked to SOA formation as a function of ecosystems. Therefore, quantification and research of the fluxes of these VOCs is crucial. Field measurements of fluxes of BVOC and their
270 oxidation products exhibiting reduced volatility, such as SVOC are challenging, and they can only be measured with rather short inlets to avoid wall losses during sampling. Recently a PTR3 instrument was used on top of the SMEAR II tower in Hyytiälä (Fischer et al., 2021) at 36 m above ground level and 15 m above the canopy of a forested ecosystem dominated by terpenoid emitters. The PTR3 instrument was installed approximately 4 m away from the tower structure and the virtually wall-less inlet was successfully tested allowing undisturbed gas sampling from this distance. For the first time emission fluxes
275 of sesquiterpene ozonolysis products and diterpenes were recorded. With the low flux signal-to-noise ratio achieved with the new instrumentation, we can now track and study clear diurnal patterns, even for the smallest emissions rates virtually in real time. Such intensive campaigns demonstrate the feasibility of new technology to be integrated in Flagship stations providing an extended parameter set in the future.


280 3.3 COVID restrictions

By the end of 2021, the global spread of COVID-19 caused by the SARS-CoV-2 virus has resulted in the loss of over 10 million lives (Adam, 2022; Msemburi et al., 2023). In China, national interventions were implemented starting from January 2020 to prevent the spread of the virus (China NHC, 2020). The strict lockdown measures associated with the COVID-19
285 pandemic provide a unique opportunity to investigate, in a real-world atmospheric laboratory, the direct and indirect effects of reduced emissions, as well as atmospheric chemistry and interacting processes associated with these emission changes, on air quality (e.g. Kroll et al., 2020; Jiang et al., 2021; Wang et al., 2021; Sokhi et al., 2021; Amouei Torkmahalleh et al., 2021).

This unique opportunity is a good demonstration of the strength of the SMEAR concept applied to atmospheric observations. First, in such an unplanned situation where new research activities have also been restricted, it is impossible to organize and
290 carry out targeted intensive observations. Second, although there are several functioning observing stations, the relatively poor measurement capacity in most of them is unable to support the in-depth analyses needed for new scientific insights. To date, there are hundreds of atmospheric science studies relevant to the COVID-19 lockdown (https://docs.google.com/document/d/1UTQvW_OytC37IatMNR5qJK7qKfSylNpI2fT3pdteVZA/edit). However, a large proportion of these studies only report variations of a few atmospheric parameters and are far from providing a mechanistic

295 understanding of changes in atmospheric processes. There are also studies that use regional models to understand the atmospheric processes during the lockdown, but these modeling results have limited verification due to the lack of comprehensive observations.

The Aerosol and Haze Laboratory of Beijing University of Chemical Technology (AHL/BUCT; Liu et al., 2020) is one of the 300 stations fully implementing the SMEAR concept. This station was established in January 2018, and since then it has been operating uninterruptedly with a full measurement capacity. Our comprehensive observations showed that the lockdown caused changes of different magnitudes in various atmospheric parameters (Fig. 3). In general, most of the primary pollutants, such as NOx, SO₂, BC, and VOCs, showed a reduction in their abundance, but at different levels. For example, NOx was reduced by more than 50%, SO₂ by ~25% and VOCs only by ~15%. This suggests that emissions from different source sectors 305 were affected differently by the lockdown. In contrast to the primary pollutants, most of the secondary pollutants showed increased concentrations. Particulate nitrate, sulfate, ammonia, organics, and gas-phase highly oxygenated organic molecules (HOMs) increased by ~50–150%. This indicates that secondary pollution, i.e. the conversion of primary pollutants into secondary ones, became more efficient. This phenomenon is closely related to the increased oxidation capacity of the atmosphere, as indicated by the increased concentrations of OH, NO₃ and O₃.



310

Figure 3: Variations of primary and secondary pollutants caused by the lockdown. Relative changes of atmospheric variables between the COVID-19 lockdown period (24th Jan – 5th Mar 2020) and pre-lockdown period (1st Jan – 23rd Jan 2020). The relative changes are defined as $\frac{[X]_{lock} - [X]_{pre}}{[X]_{pre}} \times 100\%$, where [X] is the average of each variable. Variables associated with new particle formation (NPF) are shown only for NPF days during the daytime.

Our comprehensive data sets allow us to obtain cutting-edge knowledge in several research directions. Here, we provide two examples that provide direct observational evidence, showing the substantial influence of anthropogenic emissions on the
320 atmospheric oxidative capacity in both daytime and nighttime.

3.3.1 How did the atmospheric new particle formation respond to COVID-19 lockdown

Yan et al. (2022) explored how NPF responded to emissions reductions in Beijing during the COVID-19 lockdown. Clustering between SA and base molecules drove the initial NPF in both the pre-lockdown and lockdown periods. Our results show that
325 this clustering was insensitive to emission reductions. Through direct observation, this study provided evidence that traffic emissions do not appear to be a significant source of NPF in Beijing, in contrast to conclusions drawn from some recent urban studies (Rönkkö et al., 2017; Guo et al., 2020).

During the lockdown period, we hypothesized that the reduction in nitrogen oxides (NO_x) concentrations would promote
330 particle growth. This is because NO can suppress particle growth by changing the composition of oxidized organic molecules (OOMs) and making them more volatile on average (Yan et al., 2020). However, our study found otherwise. Although we noted changes in the composition of OOMs, especially in molecules arising from the oxidation of aromatic volatile organic compounds, there were only negligible changes in the volatility of OOMs. These results indicate that the reaction between
335 RO_2 and NO still plays a vital role in OOM formation even after a dramatic reduction in NO_x levels. It has been suggested that the autoxidation of RO_2 will become more important in atmospheric chemistry as NO_x concentrations continue to decrease in North America (Praske et al., 2018), leading to increased toxicity of peroxide-driven particles and the formation of secondary organic aerosols (Zhao et al., 2017). However, our findings suggest that these harmful effects on human health and air quality in Beijing are less likely to arise in the immediate future.

340 3.3.2 Enhanced formation of secondary organic carbon associated with NO_3 radical


Carbonaceous aerosols are acknowledged to have significant impacts on climate change, Earth's radiation balance, visibility, and human health (Donahue et al., 2009; Bond et al., 2013; IPCC, 2021). We examined carbonaceous aerosols measured with an OC/EC analyzer between 1 December 2019 and 15 March 2020 in Beijing, encompassing the COVID-19 pandemic period (Feng et al., 2022). Our findings showed that anthropogenic gas-phase pollutants and primary organic
345 compounds were greatly reduced during the lockdown period. However, we also observed the emergence of enhanced nighttime secondary organic carbon, which we attributed to nocturnal chemistry associated with the oxidation by NO_3 radical. Our results indicate that this nocturnal chemistry phenomenon warrants greater attention in efforts to reduce PM concentration in China.

350 3.4 Long-term trends in comprehensive observations of atmospheric variables

The long-term observations at the SMEAR II station in Hyytiälä, Finland, cover measurements of trace gas concentrations (SO₂, O₃, NO_x, CO, CO₂) as well as volatile organic compounds (VOCs, such as monoterpenes), which are measured on multiple heights above the ground at the 128-m-high mast. The continuous time-series of measurements starting from 1996 allow us to quantify long-term trends in these variables (Fig. 4). In addition, mass spectrometer measurements of sulphuric acid (H₂SO₄, the main oxidation product of SO₂) started in 2016, and proxy calculations based on the measured H₂SO₄ concentrations enable extending this time series (Petäjä et al., 2009; Dada et al., 2020).

The monthly-median concentration of the H₂SO₄ proxy has a decreasing trend of $-2.6\%/\text{year}$ (Fig. 4a). The H₂SO₄ proxy is calculated based on the production of H₂SO₄ due to the oxidation of SO₂ by OH radicals and via stabilized Criegee intermediates which are produced in ozonolysis of monoterpenes (Dada et al., 2020), and on the loss of H₂SO₄ due to its condensation into pre-existing particles. Both the source and sink terms of H₂SO₄ have decreasing trends in Hyytiälä during 1996–2022, being $-2.7\%/\text{year}$ for the SO₂ concentration $-1.0\%/\text{year}$ for the condensation sink (Fig. 4c-d). The stronger decrease in the H₂SO₄ precursor vapor concentration compared with the H₂SO₄ sink seems to determine the long-term trend observed in the sulphuric acid proxy concentrations.

The monoterpene concentrations are characterized by a large year-to-year variability and do not show a statistically significant trend (Fig. 4b) (see also e.g., Tarvainen et al., 2005; Taipale et al., 2011; Rantala et al.; 2015, Hellen et al., 2018). In the summertime, the monoterpene concentrations are highest during the year and have stayed relatively constant, whereas the annually lowest concentrations during winter and spring show also the largest variability between years. The long-term data show strong seasonal and diurnal patterns in emission rates, which mostly relate to changes in temperature and partly also light availability. Furthermore, vegetation phenological events and biotic and abiotic stresses produce high emission peaks (e.g., Aalto et al., 2014). The amplitude of the daily and seasonal variations in monoterpene emission rates is high and masks the potential climate change effect over the years. This emphasizes the importance of versatile, comprehensive measurements for quantifying atmospheric processes.



375 **Figure 4: Trends in (a) sulphuric acid concentration, (b) monoterpene concentration, (c) sulphur dioxide concentration, and (d)** condensation sink. Note the different time periods: in panels (a) and (b) 2010–2022, and in panels (c) and (d) 1996–2022. The dotted black lines show monthly medians of observations and the grey solid lines are trends fitted to the logarithmic values of the monthly data. Monoterpene concentrations do not have a statistically significant trend and therefore the trend line is not shown in panel (b).

380 **4 Integration of the data from satellites, models and in situ observations**

4.1 Satellite and Airborne observation

Satellites provide data on a global scale of atmospheric composition, radiation, surface properties and meteorology. Passive satellite measurements of atmospheric gases and aerosols are representative over an entire atmospheric column, and hence they are not directly comparable to in situ measurements. Although satellites cannot provide as detailed and wide range of

385 different atmospheric parameters as comprehensively equipped in-situ measurement stations, such as SMEAR II, with their spatial coverage they can provide very valuable and complementary information. After understanding and analyzing in-situ and satellite measurements together on a station-by-station basis, satellite data can enable the transition from point-like measurements to the interpretation of regional variability in atmospheric processes (e.g. Viatte et al., 2021; Pseftogkas et al., 2022; Hakala et al., 2019).

One example of utilizing satellite data, when moving from pointwise to global (or regional) analysis, has been to better understand the new particle formation (NPF) phenomena on a larger spatial scale (Kulmala et al., 2011; Sundström et al., 2015). As satellites cannot essentially detect aerosol particles smaller than 100 nm in diameter, these observations as such are not directly applicable for NPF studies. However, satellites provide information on many atmospheric parameters tightly linked to NPF, such as UV radiation, trace gas concentrations and estimations on ambient aerosol loads. By analyzing these data with detailed in situ measurements, it is possible to develop merged satellite variables that can be used to study the regional variation of NPF. Other examples are various climate-related feedback loops, e.g. between the atmosphere and the biosphere, where satellite observations could have the potential of increasing the understanding of the spatial scale variation. Such development work would not be possible without SMEAR-type observations that have the capability of providing process-level understanding of the phenomena. It is also essential that such extensive in situ observations exist in various environments, so that the sensitivity of satellite observations in these kinds of applications could be properly tested.

Passive satellite instruments provide typically so-called columnar measures, and for instance aerosol optical depth (AOD) is the vertically-integrated aerosol extinction. Similarly, gas concentrations measured by satellite represent column or partial column concentrations, typically over a tropospheric column. Therefore, comparison with surface in-situ measurements cannot offer a direct validation for the measurements made by satellite instruments. However, all possible columnar measurements (e.g. AOD by ground-based sun-photometers) at the same ground station, on the other hand, would facilitate satellite data validation and provide information on the accuracy of the satellite observations. Currently, the SMEAR II station in Hyytiälä is accompanied with an Aerosol Robotic Network station (AERONET, Holben et al., 1998), which allows a direct validation of satellite-based aerosol observations. Moreover, satellite validation would strongly benefit from reliable gas and aerosol vertical profile observations from the surface level up to the stratosphere.

One possible future pathway in better bridging the spatial-scale gap between pointwise in-situ measurements and large-scale satellite measurements would be to utilize unmanned aerial vehicle (UAV) measurements (Motlagh et al., 2023). Satellite data have spatial resolutions limited to a few hundred meters at best, and in atmospheric observations more typically to kilometers. Drone measurements (e.g., Kezoudi et al., 2021) could bring more insight into the sub-kilometer scale variations if carried out in the vicinity of SMEAR-type stations at the time of the satellite overpass.

4.2 Model frameworks

Model frameworks for the global climate and Earth systems have been constructed to replicate real-world processes and interactions as closely as necessary to understand the current state of these systems, reasons for past changes, and ultimately to simulate future climate pathways in order to support adaptation and mitigation efforts (Bauer et al., 2021). Modern Earth System Models (ESMs) combine an increasing number of individual components, including not only the physical ocean and


atmosphere models but also detailed descriptions of chemistry, aerosols and the biosphere (e.g. Döscher et al., 2022). This increase in model complexity has established groundbreaking research of Earth system feedbacks and quantification of their
425 strength in current and future climate (Sporre et al., 2019; Thornhill et al., 2021).

Despite rigorous validation of its individual process components, evaluation and constraining of highly-coupled ESMs remains difficult due to the large number of interactions and feedbacks within the Earth system (e.g. Sporre et al., 2020). In the temporal and spatial scales of ESMs even the observational record as a whole remains brief and irregular, and therefore the observations
430 must be extended by proxies of changes in the historical period (Wandji Nyamsi et al., 2020) and towards the Earth's deep past over millions of years (Wong et al., 2021). Only integrated long-term observations can provide multidisciplinary data to support evaluation of simulated Earth system feedbacks and their components.

With increasing complexity and process details, ESMs can arrive at correct results via wrong reasons and counteracting biases.
435 The advancement of spatial resolution and process descriptions within ESMs already allows evaluation at the process-scale. For example, NPF events can be co-analyzed from ESMs and long-term datasets (Bergman et al., 2022). Such process-oriented analysis is essential for validating the reasons for biases or systematic errors in simulated properties (e.g. CCN), but this requires dedicated long-term observations to constrain the models throughout distinct climate states and changing environment (Fanourgakis et al., 2019). Recent advances in trajectory-based analysis of ESMs provide a novel way to investigate simulated
440 air-masses and station footprints co-located with observations. To complete a global 4D evaluation of ESM performance, the long-term stationary datasets should be complemented with surface or airborne transects, vertical profiles and satellite retrievals (e.g. van Noije et al., 2021).

With the competition between increasing spatial resolution and more detailed process descriptions in ESMs, data-driven
445 approaches have been suggested to replace computationally intensive modules (Ahola et al., 2022). Whether through emulation, neural networks or other machine learning techniques, the teaching and learning process requires comprehensive understanding of the model realm complemented with integrated observational suite (e.g. Schreck et al., 2022).

4.3 Integration of different approaches



450

Figure 5: Combination of methods to integrate “bottom-up” and “top-down” insights on atmospheric aerosol and its interactions with clouds, as outlined within the FORCeS project (see forces-project.eu. Figure courtesy of Tinja Olenius.

Combination of emerging long-term in-situ measurements, satellite data and process understanding bear a great potential for finding new ways to evaluate and constrain ESMs, and to reduce uncertainties in their projections (see Fig. 5, adopted from the FORCeS project). The inevitable spatial limitation related to in-situ observations can, to some degree, be overcome by ensuring long enough temporal coverages and enhancing the number of representative data points (to be compared with satellites and models) this way (e.g. Isokäntä et al., 2022; Khadir et al., 2023).

455

Long-term, global in-situ observations are specifically useful in pin-pointing the model weaknesses and strengths, as well as in providing detailed observations with various techniques at well-defined altitudes, as opposed to sampling the entire atmospheric column. Long-term in-situ observations offer great opportunities to compare detailed process-level observations with satellite observations and large-scale models. Detailed measurements enable e.g. investigations of size-segregated trends in aerosol loadings in a regional context using both ESM and observational data (e.g. Leinonen et al., 2022). While the number

460

of relevant data is steadily increasing, more long-term observations from under-sampled parts of the atmosphere (global south, 465 highly remote areas) are needed. In addition, combining long-term in-situ measurements, satellite data, ESM model outputs and process understanding offers a great potential for finding new ways to evaluate and constrain the biosphere-climate feedbacks in ESMs, the magnitude of which is still highly variable between models (Thornhill et al., 2021). By separating the different processes (e.g. those that relate biogenic emissions and resulting aerosol concentrations to air temperatures and cloud properties), and by combining long-term observations with satellite data and multiscale modelling, one can facilitate the 470 evaluation of the predictive abilities of the models (Blichner et al., to be submitted). This approach can help to isolate the impact of individual factors and improve our understanding of the underlying processes.

For example, the uncertainty in the effective radiative forcing due to aerosol-cloud interactions is governed by the cloud susceptibility to aerosol perturbations (Bellouin et al., 2020). This is split into two components which are (i) the response of 475 the cloud droplet number concentration to aerosol perturbations – relevant for the radiative forcing due to aerosol-cloud interactions, also known as the Twomey effect – and (ii) the rapid adjustments in particular of cloud liquid water path and cloud fraction (Bellouin et al., 2020). In-situ long term aerosol and cloud observations enable investigations of cloud-susceptibility to aerosol perturbations. In-situ observations (both long-term and campaign-wise) and process understanding combined with ESM model outputs (or with satellites), facilitate pin-pointing specific processes or factors that 480 should be improved in order to be able to describe the cloud activation and aerosol indirect forcing correctly in the models.

5 Future perspectives and possibilities

Currently, the speed of climate change along with its unpredictable consequences are challenging the capacities of existing observation systems. In addition, the ability to analyze the yet unknown questions and challenges, the “black swans” (Taleb, 485 2010), calls for comprehensive continuous observations. For example, COVID19 gave an unexpected opportunity to demonstrate the effect of exceptional reductions in anthropogenic emissions on air quality and climate (e.g. Gettelman et al., 2020; Wijnands et al., 2022). In this case the already running SMEAR-type, comprehensive measurements at the BUCT/AHL station in Beijing enabled us to investigate the atmospheric processes in detail. Other examples of this type of unusual and extraordinary events could be volcanic eruptions, gas pipe attacks, extreme weather events, forest fires, exceptionally dry periods, economic collapses or chemical weapons. All these events have both short- and long-term dynamic effects on air 490 quality and the climate system as well as on the functioning of societies. Also, the possibility of realized global tipping points, such as permafrost loss or boreal forest shift towards tundra, may lead to unexpected environmental episodes, events and feedbacks (Rockström et al., 2009, Kulmala et al., 2015, Lenton et al., 2019). The key questions are whether the majority of current observation systems contain sufficiently comprehensive set of variables to capture these events, and whether we have the preparedness to detect, analyze and quantify these events.

Recently proposed geoengineering approaches for mitigating global warming include a clear potential for large magnitude feedbacks which can have significant, yet unpredictable consequences to other processes. Risks related to uncontrolled geoengineering without international laws and the manipulation of the atmosphere highlight the value of continuous, comprehensive measurements detecting the changes. For example, operational Solar Radiation Modification (SRM) deployment would introduce new environmental and socio-economic threats like damaging the ozone layer and overcompensating climate change at regional scales (UNEP, 2023).

We need open big data to meet the present grand challenges, and we need to collect comprehensive data to be able to answer questions which do not exist today. The questions can be societal, economic or scientific, or any combination of these. To effectively collect, distribute and utilize big data, there are several key actions that needs to be considered:

Firstly, it is important to promote open data flows and storages globally via open access data platforms and structures. This can be achieved through optimizing data flows by also considering how to access and analyze the data at the storage site instead of transferring huge amounts of data. Advanced AI and data mining techniques should be employed to explore and utilize the data effectively. As importantly the in situ community should actively take part of the ongoing development of AI based Earth Virtualization Engines (EVE) which aims to animate the Earth observations, in situ and satellite data together, at 1 km scale for different users (Stevens et al., 2023). It is also important to transfer knowledge to make the data more accessible and to develop and provide examples and roadmaps for local data owners that highlights how to get merits via open data. The more local the data and data needs are, the more challenging it is to have them fully open. Therefore, it is crucial to demonstrate the benefits of offering open access to the data.

Secondly, global collaboration is needed to develop measurement protocols and data standards to reliably observe concentrations, fluxes and changes in the atmosphere and the environment. When using low-cost sensors, it is crucial to establish a proper calibration system that ensures data quality and traceability. Also, existing observation station types require calibrated sensors and enhanced harmonization. It is important to connect to existing harmonization actions by international organizations, such as the European Committee for Standardization (CEN), European environmental Research infrastructures and Network of Air Quality Reference Laboratories (AQUILA) etc. Several World Meteorological Organization working groups are already active towards these goals. However, we need to go to the next level to make in-house processes more effective.

525

Finally, it is necessary to establish a hierarchy of stations ranging from cost-effective sensors (low cost) to comprehensive flagship stations, such as SMEAR – Stations to Measure Earth surface Atmosphere Relationships, by utilizing the knowledge and experience from the European Strategy Forum on Research Infrastructures (ESFRI) as well as from operational observation networks pertinent to different domains in the atmosphere – environment continuum. Within the next 20 years, we should have

530 a station network utilizing the hierarchy of stations with three steps, namely Flagship stations, Median stations and low cost sensors with enough Flagship stations scattered globally spatially and in ecosystem level to have enough representativeness of varying conditions. The comprehensive Flagship stations should be preferably part of GAW network with 500–1000 stations like SORPES, SIOS and SMEAR (Kulmala, 2018). The Median stations are high-end stations, but typically focusing on a specific topic (e.g. flux stations, AQ networks). Low cost sensors need calibration from Flagship and Median stations and
535 utilizing of AI and 5G/6G/7G networks (Rebeiro-Hargrave et al., 2021).

Taking these actions will improve our understanding of the environment and our ability to respond to environmental challenges. An important question is who is willing to take the lead? Probably large international organizations are needed like WMO. In practice, we need to develop steps towards a GAW+ and maybe even to establish International climate /atmosphere
540 institute. The institute should be multinational and multi-institutional research center following e.g. the model of CERN, but focusing on the atmospheric and Earth system research.

This would be based on combining the experiences from WMO Global Atmospheric Watch (GAW) program (WMO 2017), COPERNICUS, and international *in situ* research infrastructures like the ESFRIs ICOS, ACTRIS and eLTER (see Section 1).
545 The observation systems and research infrastructures, present standards, protocols and recommendations are consensus-based. For example, essential atmospheric variables and data products management alone have been developed by several different actors, such as the Global Climate Observing System (GCOS) (WMO 2022) and GAW program (WMO 2017). These standardized systems have taken years to develop and are still in progress but need to be continued.

550 Under the WMO leadership, we should aim to the establishment of the global observatory for comprehensive data set(s) on *weather, climate, water and environment*. This framework would provide a wide range of benefits, such as creating a real-world component and comparison for digital twin(s). It will also allow a proper WMO contribution to share integrated big datasets. The global observatory will provide a seamless connection between in-situ observations, remote sensing and multiscale model data. This will enable easy access and utilization of remote sensing products, such as inland water altimetry
555 for rivers, lakes and reservoirs as well as arctic snow and ice cover. Furthermore, it would provide observational support for global food forecasting through real-time dissemination of river level and discharge data, air quality forecasting and food and water supply forecasting. It would also enable us to predict future climate and find out feedbacks and interactions between various environmental factors.

560 Once we have collected all these data, it is crucial that it is utilized effectively. To be able to use the big data, open access is typically needed. However, there are several barriers before the data can be used. The barriers include the lack of documentation, unknown data, misunderstood user needs, discipline specific jargon, bad and unusable interfaces, authorization problems, wrong terminology, training problems, unknown formats, difficulties in licensing and documentation, etc. To

overcome the barriers of information, we need to have mutual trust and understanding of the needs, in addition to which we
565 need to have access to the data to make new discoveries. This can be achieved by implementing the FAIR principles (Findable,
Accessible, Interoperable, Reusable) (Wilkinson et al., 2016), open data policies, proper knowledge transfer and conducting
impact investigations e.g. IIASA.

It is worth noting that the most important reason to investigate multiple variables with continuous measurements is that we
570 never know beforehand, when we will meet a ‘black swan’. When we have comprehensive, continuous, open data, we can
analyze the data to study unexpected phenomena and answer to the upcoming challenges.

6 Conclusions

The need for comprehensive open data sets is obvious. The climate emergency and fast development of AI-based climate
analysis force us to develop a new generation observation system. Within the next 20 years, we should have a well-established
575 station network, a “Global Earth Observatory”, providing standardized data from different environments and scales.

Traditionally, and even in many cases today, there exist distinct infrastructures and their designated users, with experiences
rather far from each other. Different research groups have typically their own instruments, raw data, data analysis methods and
publications. This is not an efficient way to meet grand challenges that can only be tackled with interdisciplinary approach. In
580 the future, we need to utilize more joint efforts, including co-location of research infrastructures. Already to store raw data
jointly, and to analyze it together in systematic ways, provides a big surplus. To have common data repositories for storing
analyzed and published data is a big step forward.

One example on how comprehensive observations can be utilized is to solve air quality issues. In order to be able to understand
585 the chemistry of air pollution, we need to observe multiple pollutants in existing air pollution cocktails (Kulmala, 2015). We
should also remember that we spend 90% of our time indoors, and therefore also indoor air quality need to be understood. Air
quality is important e.g. for health effects, visibility and from the acidification point of view. The multiple pollutants include
PM_{2.5}, size-resolved particle number, black carbon, O₃, NO_x, SO₂, CO, acids, various organic compounds etc., and their
interactions and feedbacks. Our typical framework is a seamless chain from deep understanding to solutions, starting from
590 observations and then continuing to understanding processes, feedbacks and interactions. These steps are needed to control air
pollution and to improve air quality.

In basic and applied research, we make new discoveries and new knowledge from the resources we have with the money we
have at our disposal. New discoveries and knowledge often lead to innovations and with innovations people make money from

595 that knowledge. Innovations and new knowledge are a key for society to maintain and enhance wellbeing and this way the circle closes.

It is crucial to utilize multidimensional, multidisciplinary, multiscale approach to be able to answer questions related to grand challenges. It is also important to have a clear and ambitious vision from deep understanding to practical solutions. Also, we
600 need seamless chain to connect measurements, modelling and theory as well as from research to innovations, economic growth and human wellbeing.

The main benefits that research community can gain from using integrated research infrastructure approach includes higher-quality science and higher visibility and recognition from the society and its various stakeholders and higher amount of
605 scientific users both nationally and internationally. Moreover, collaboration possibilities will be enhanced, including feedbacks between domains, landscape analysis, up- and downscaling. Detailed experiments and observations can support each other, so that new observational and numerical methods can be developed. The improved utilization of data flows and synergies in data use are foreseen.

610 In order to meet grand challenges and answer open scientific, societal and economic questions, we need to build upon a network of domain-specific research infrastructures, such as ACTRIS, eLTER and ICOS. We need to acknowledge that for example ACTRIS is already integrating several subfields, namely aerosol in-situ, trace gas and cloud in-situ observations and ground-based remote sensing of aerosols, clouds and trace gases. The human-environment relationship, i.e., the social-ecology research, is part of the transdisciplinary approach in eLTER, enabling policy-relevant research and interaction with society at
615 different scales. The SMEAR concept in essence includes co-location and integration of the observations performed in the domain specific environmental RIs. A further connection to and integration with e.g., health and societal data are needed. Furthermore, we need excellent science, with high quality, critical mass and interdisciplinary research as well as education and training, i.e. knowledge exchange. We need to contribute to innovation ecosystem and have continuous, long-term dialogue with policy makers. Internationally, this enables clear contributions to science diplomacy based on integrated
620 scientific viewpoint.

Acknowledgements

We acknowledge the following projects: ACCC Flagship funded by the Academy of Finland grant number 337549, Academy professorship funded by the Academy of Finland (grant no. 302958), Academy of Finland projects no. 1325656, 311932, 334792, 316114, 325647, 325681, 347782, “Quantifying carbon sink, CarbonSink+ and their interaction with air quality”
625 INAR project funded by Jane and Aatos Erkko Foundation, “Gigacity” project funded by Wihuri foundation, European Research Council (ERC) project ATM-GTP Contract No. 742206, and European Union’s Horizon 2020 research and innovation programme under Grant agreement No 101003826 via project CRiceS (Climate Relevant interactions and

feedbacks: the key role of sea ice and Snow in the polar and global climate system) and Horizon Europe research and innovation programme under Grant agreement No 101056783 via project FOCI (Non-CO₂ Forcers and their Climate, Weather, Air Quality and Health Impacts). University of Helsinki support via ACTRIS-HY is acknowledged. Support of the technical and scientific staff in Hyttiälä and BUCT/AHL are acknowledged.

635

References

Aalto J., Kolari P., Hari, P. Kerminen V. -M., Schiestl-Aalto P., Aaltonen H., Levula J., Siivola E., Kulmala M., and Bäck J.: New foliage growth is a significant, unaccounted source for volatiles in boreal evergreen forests, *Biogeosci.*, 11, 1331-1344, 2014.

640

Adam, D.: True COVID death toll could be more than double official count, *Nature*, 605, 206, 2022.

Ahola, J., Raatikainen, T., Alper, M. E., Keskinen, J.-P., Kokkola, H., Kukkurainen, A., Lipponen, A., Liu, J., Nordling, K., Partanen, A.-I., Romakkaniemi, S., Räisänen, P., Tonttila, J., and Korhonen, H.: Technical note: Parameterising cloud base updraft velocity of marine stratocumuli, *Atmos. Chem. Phys.*, 22, 4523–4537, 10.5194/acp-22-4523-2022, 2022.

645

Almeida, J., Schobesberger, S., Kurten, A., Ortega, I. K., Kupiainen-Maatta, O., Praplan, A. P., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurten, T., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppa, J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petaja, T., Riccobono, F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld, J. H., Simon, M., Sipila, M., Stozhkov, Y., Stratmann, F., Tome, A., Trostl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P. L., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkamaki, H., and Kirkby, J.: Molecular understanding of sulphuric acid-amine particle nucleation in the atmosphere, *Nature*, 502, 359-363, 10.1038/nature12663, 2013.

650

Amouei Torkmahalleh, M., Akhmetvaliyeva, Z., Omran, A. D., Faezeh Darvish Omran, F., Kazemitabar, M., Naseri, M., Naseri, M., Sharifi, H., Malekipirbazari, M., Kwasi Adotey, E., Gorjinezad, S., Eghtesadi, N., Sabanov, S., Alastuey, A., de Fátima Andrade, M., Buonanno, G., Carbone, S., Cárdenas-Fuentes, D. E., Cassee, F. R., Dai, Q., Henríquez, A., Hopke, P.

K., Keronen, P., Khwaja, H. A., Kim, J., Kulmala, M., Kumar, P., Kushta, J., Kuula, J., Massagué, J., Mitchell, T., Mooibroek, D., Morawska, L., Niemi, J. V., Ngagine, S. H., Norman, M., Oyama, B., Oyola, P., Öztürk, F., Petäjä, T., Querol, X., Rashidi, Y., Reyes, F., Ross-Jones, M., Salthammer, T., Savvides, C., Stabile, L., Sjöberg, K., Söderlund, K., Sunder Raman, R.,
665 Timonen, H., Umezawa, M., Viana, M., and Xie, S.: Global Air Quality and COVID-19 Pandemic: Do We Breathe Cleaner Air?, *Aerosol and Air Quality Research*, 21, 200567, 10.4209/aaqr.200567, 2021.

Arneth, A., Harrison, S. P., Zaehle, S., Tsigaridis, K., Menon, S., Bartlein, P., J., Feichter, J., Korhola, A., Kulmala, M., O'Donnell, D., Schurgers, G., Sorvari, S., and Vesala, T.: Terrestrial biogeochemical feedbacks in the climate system, *Nature Geosci.*, 3, 525-532, 2010.
670

Artaxo, P., Hansson, H.-C., Andreae, M. O., Bäck, J., Gomes Alves, E., Barbosa, H. M. J., Bende,r F., Bourtsoukidis, E., Carbone, S., Chi, J., Decesari, S., Despres, B. V. R., Ditas, F., Ezhova, E., Fuzzi, S., Hasselquist, N. J., Heitzenberg, J., Holanda, B. A., Guenther, A., Hakola, H., Heikkinen, L., Kerminen, V.-M.., Kontkanen, J., Krejci, R., Kulmala, M., Lavric, J. V., de Leeuw, G., Lehtipalo, K., Machado, L. A. T., McFiggans, G., Franco, M. A. M., Meller, B. B., Morais, F. G., Mohr, C., Morgan, W., Nilsson, M. B., Peichl, M., Petäjä, T., Prass, M., Pöhlker, C., Pöhlker, M. L., Pöschl, U., von Randow, C., Riipinen, I., Rinne, J., Rizzo, L. V., Rosenfeld, D., Silva Dias, M. A. F., Sogacheva, L., Stier, P., Swietlicki, E., Sörgel, M., Tunved, P., Virkkula, A., Wang, J., Weber, B., Yanez-Serrano, A. M., Zieger, P., Mikhailov, E., Smith, J. N., and Kesselmeier, J. (2022) Tropical and boreal forest – atmosphere interactions;; A review. *Tellus B*, 74, 24-163, 2022.
675

Asmi, E., Kivekäs, N., Kerminen, V.-M., Komppula, M., Hyvärinen, A.-P., Hatakka, J., Viisanen, Y., and Lihavainen, H.: Secondary new particle formation in Northern Finland Pallas site between the years 2000 and 2010. *Atmos. Chem. Phys.*, 11, 12959-12972, 2011.
680

Baldocci, D. D.: How eddy covariance flux measurements have contributed to our understanding of global change biology, *Global Change Biol.*, 26, 242-260, <https://doi.org/10.1111/gcb.14807>, 2019.
685

Boden, T. A., Krassovski, M., and Yang, B.: The AmeriFlux data activity and data system: an evolving collection of data management techniques, tools, products and services, *Geosci. Inst. Meth. Data Syst.*, 2, 165-176, <https://doi.org/10.5194/gi-2-165-2013>, 2016.
690

Bauer, P., Stevens, B., and Hazeleger, W.: A digital twin of Earth for the green transition, *Nature Clim. Change*, 11, 80–83, 2021.
695

695 Beck, L., Sarnela, N., Junninen, H., Hoppe, C. J. M., Garmash, O., Bianchi, F., Riva, M., Rose, C., Peräkylä, O., Wimmer, D.,
Kausiala, O., Jokinen, T., Ahonen, L., Mikkilä, J., Hakala, J., He, H.-U., Kontkanen, J., Wolf, K. K. E., Cappelletti, D.,
Mazzola, M., Traversi, R., Petroselli, C., Viola, A. P., Vitale, V., Lange, R., Massling, A., Nojgaard, J. K., Krejci, R., Karlsson,
L., Zieger, P., Jang, S., Lee, K., Vakkari, V., Lampilahti, J., Thakur, R. C., Leino, K., Kangasluoma, J., Duplissy, E.-M.,
Siivola, E., Marbouti, M., Tham, Y. J., Saiz-Lopez, A., Petäjä, T., Ehn, M., Worsnop, D. R., Skov, H., Kulmala, M., Kerminen,
700 V.-M., and Sipilä, M.: Differing mechanisms of new particle formation at two arctic sites, *Geophys. Res. Lett.*, 48,
e2020GL091334, 2021.

Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M.,
Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle,
705 F., Mauritzen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz,
M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative
Forcing of Climate Change, *Reviews of Geophysics*, 58, e2019RG000660, <https://doi.org/10.1029/2019RG000660>, 2020.

Bergman, T., Makkonen, R., Schrödner, R., Swietlicki, E., Phillips, V. T. J., Le Sager, P., and van Noije, T.: Description and
evaluation of a secondary organic aerosol and new particle formation scheme within TM5-MP v1.2, *Geosci. Model Dev.*, 15,
710 683–713, <https://doi.org/10.5194/gmd-15-683-2022>, 2022.

Blichner, S. M., Yli-Juuti, T., Kokkola, H., Mielonen, T., Holopainen, E., Heikkinen, L., Petäjä, T., Pöhlker, C., Artaxo, P.,
Meller, B., Scott, C., Sporre, M., Svenhag, C., Partridge, D., Tovazzi, E., Virtanen, A., and Riipinen, I.: Constraining the
BVOC feedback in models with process based evaluation along the scale chain, 2023, manuscript in prep.

715 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B.,
Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H.,
Zhang, S., Bellouin, N., Gttikundia, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz,
J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender C. S.: Bounding the role of black carbon in the climate system, *J.
Geophys. Res.: Atmos.*, 118, 5380-5552, 2013.

720 Boy, M., Thomson, E. S., Acosta Navarro, J.-C., Arnalds, O., Batchvarova, E., Bäck, J., Berninger, F., Bilde, M., Brasseur,
Z., Dagsson-Waldhauserova, P., Castarède, D., Dalirian, M., de Leeuw, G., Dragosics, M., Duplissy, E.-M., Duplissy, J.,
Ekman, A. M. L., Fang, K., Gallet, J.-C., Glasius, M., Gryning, S.-E., Grythe, H., Hansson, H.-C., Hansson, M., Isaksson, E.,
Iversen, T., Jónsdóttir, I., Kasurinen, V., Kirkevåg, A., Korhola, A., Krejci, R., Kristjansson, J. E., Lappalainen, H. K., Lauri,
725 A., Leppäranta, M., Lihavainen, H., Makkonen, R., Massling, A., Meinander, O., Nilsson, E. D., Olafsson, H., Pettersson, J.
B. C., Prisle, N. L., Riipinen, I., Roldin, P., Ruppel, M., Salter, M., Sand, M., Seland, Ø., Seppä, H., Skov, H., Soares, J., Stohl,
A., Ström, J., Svensson, J., Swietlicki, E., Tabakova, K., Thorsteinsson, T., Virkkula, A., Weyhenmeyer, G. A., Wu, Y., Zieger,

P., and Kulmala, M.: Interactions between the atmosphere, cryosphere, and ecosystems at northern high latitudes, *Atmos. Chem. Phys.*, 19, 2015–2061, <https://doi.org/10.5194/acp-19-2015-2019>, 2019.

730

Brean, J., Beddows, D. C. S., Harrison, R. M., Song, C., Tunved, P., Ström, J., Krejci, R., Freud, E., Massling, A., Skov, H., Asmi, E., Lupi, A., and Dall’Osto, M.: Collective geographical ecoregions and precursor sources driving Arctic new particle formation., *Atmos. Chem. Phys.*, 23, 2183–2198, 2023.

735 Butt, E. W., Turnock, S. T., Rigby, R., Reddington, C. L., Yoshioka, M., Johnson, J. S., Regayre, L. A., Pringle, K. J., Mann, G. W., and Spracklen, D. W.: Global and regional trends in particulate air pollution and attributable health burden over the past 50 years, *Environ. Res. Lett.*, 12, 104017, 2017.

China NHC.: COVID-19 prevention and control plan (6th edition), 2020.

740 <http://www.nhc.gov.cn/jkj/s3577/202003/4856d5b0458141fa9f376853224d41d7.shtml>.

Ciais, P., Bombelli, A., Williams, M., Piao, S. L., Chave, J., Ryan, C. M., Henry, M., Prender, P., and Valentini, R.: The carbon balance of Africa: synthesis of recent research studies, *Phil. Trans. Royal Soc. A.*, 369, <https://doi.org/10.1098/rsta.2010.0328>, 2011.

745

Chu, B., Kerminen, V.-M., Bianchi, F., Yan, C., Petäjä, T., and Kulmala, M.: Atmospheric new particle formation in China, *Atmos. Chem. Phys.*, 19, 115–138, 2019.

COM(2020)21, European Green Deal Investment Plan, Brussels, 14.1.2020,

750 <https://www.eumonitor.eu/9353000/1/j9vvik7m1c3gyxp/vl5bgbajymzx>, 2020.

Dada, L., Ylivinkka, I., Baalbaki, R., Li, C., Guo, Y., Yan, C., Yao, L., Sarnela, N., Jokinen, T., Daellenbach, K. R., Yin, R., Deng, C., Chu, B., Nieminen, T., Wang, Y., Lin, Z., Thakur, R. C., Kontkanen, J., Stolzenburg, D., Sipilä, M., Hussein, T., Paasonen, P., Bianchi, F., Salma, I., Weidinger, T., Pikridas, M., Sciare, J., Jiang, J., Liu, Y., Petäjä, T., Kerminen, V.-M., and

755 Kulmala, M.: Sources and sinks driving sulfuric acid concentrations in contrasting environments: implications on proxy calculations, *Atmos. Chem. Phys.*, 20, 11747–11766, <https://doi.org/10.5194/acp-20-11747-2020>, 2020.

Deng, C., Fu, Y., Dada, L., Yan, C., Cai, R., Yang, D., Zhou, Y., Yin, R., Lu, Y., Li, X., Qiao, X., Fan, X., Nie, W., Kontkanen, J., Kangasluoma, J., Chu, B., Ding, A., Kerminen, V.-M., Paasonen, P., Worsnop, D. R., Bianchi, F., Liu, Y., Zheng, J., Wang, L., Kulmala, M., and Jiang, J.: Seasonal characteristics of new particle formation and growth in Urban Beijing. *Environ. Sci. Technol.*, 54, 8547–8557, 2020.

Doherty, R. M., O'Connor, F. M., and Turnock, S. T.: Projections of future air quality are uncertain. But which source of uncertainty is most important? *J. Geophys. Res.: Atmos.*, 127, e2022JD037948, 2022.

765

Donahue, N. M., Robinson, A. L., and Pandis, S. N.: Atmospheric organic particulate matter: From smoke to secondary particulate matter, *Atmos. Environ.*, 43, 94–106, 2009.

Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arsouze, T., Bergman, T., Bernardello, R., Boussetta, S., Caron, L.-P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblas-Reyes, F. J., Docquier, D., Echevarria, 770 P., Fladrich, U., Fuentes-Franco, R., Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M. P., Keskinen, J.-P., Koenigk, T., Makkonen, R., Massonet, F., Ménégoz, M., Miller, P. A., Moreno-Chamarro, E., Nieradzik, L., van Noije, T., Nolan, P., O'Donnell, D., Ollinaho, P., van den Oord, G., Ortega, P., Prims, O. T., Ramos, A., Reerink, T., Rousset, C., Ruprich-Robert, Y., Le Sager, P., Schmitt, T., Schrödner, R., Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E., Uotila, P., Vancoppenolle, M., Wang, S., Wårlind, D., Willén, U., Wyser, K., Yang, S., Yepes-Arbós, X., and Zhang, Q.: The EC- 775 Earth3 Earth system model for the Coupled Model Intercomparison Project 6, *Geosci. Model Dev.*, 15, 2973–3020, <https://doi.org/10.5194/gmd-15-2973-2022>, 2022.

European Commission, Directorate-General for Research and Innovation, Monitoring the open access policy of Horizon 2020: final report, Publications Office, <https://data.europa.eu/doi/10.2777/268348>, 2021.

780

Ezhova, E., Ylivinkka, I., Kuusk, J., Komsaare, K., Vana, M., Krasnova, A., Noe, S., Arshinov, M., Belan, B., Park, S.-B., Lavric, J. V., Heimann, M., Petäjä, T., Vesala, T., Mammarella, I., Kolari, P., Bäck, J., Rannik, U., Kerminen, V.-M., and Kulmala, M.: Direct effect of aerosols on solar radiation and gross primary production in boreal and hemiboreal forests. *Atmos. Chem. Phys.*, 18, 17863–17881, 2018.

785

Fanourakis, G. S., Kanakidou, M., Nenes, A., Bauer, S. E., Bergman, T., Carslaw, K. S., Grini, A., Hamilton, D. S., Johnson, J. S., Karydis, V. A., Kirkevåg, A., Kodros, J. K., Lohmann, U., Luo, G., Makkonen, R., Matsui, H., Neubauer, D., Pierce, J. R., Schmale, J., Stier, P., Tsigaridis, K., van Noije, T., Wang, H., Watson-Parris, D., Westervelt, D. M., Yang, Y., Yoshioka, M., Daskalakis, N., Decesari, S., Gysel-Beer, M., Kalivitis, N., Liu, X., Mahowald, N. M., Myriokefalitakis, S., Schrödner, R., 790 Sfakianaki, M., Tsimpidi, A. P., Wu, M., and Yu, F.: Evaluation of global simulations of aerosol particle and cloud condensation nuclei number, with implications for cloud droplet formation, *Atmos. Chem. Phys.*, 19, 8591–8617, <https://doi.org/10.5194/acp-19-8591-2019>, 2019.

Fastre, C., Possingham, H. P., Strubbe, D., and Matthysen, E.: Identifying trade-offs between biodiversity conservation and 795 ecosystem services delivery for land-use decisions, *Sci Rep-Uk*, 10, ARTN 7971, 10.1038/s41598-020-64668-z, 2020.

Feng, Z., Zheng, F., Liu, Y., Fan, X., Yan, C., Zhang, Y., Daellenbach, K. R., Bianchi, F., Petäjä, T., Kulmala, M., and Bao, X.: Evolution of organic carbon during COVID-19 lockdown period: possible contribution of nocturnal chemistry, *Sci. Total Environ.*, 808, 152191, 2022.

800 Fischer, L., Breitenlechner, M., Canaval, E., Scholz, W., Striednig, M., Graus, M., Karl, T., Petäjä, T., Kulmala, M., and Hansel, A.: First eddy covariance flux measurements of semi-volatile organic compounds with the PTR3-TOF-MS, *Atmos. Meas. Techn.*, 14, 8019-8039, 2021.

805 Fountoukis, C., Riipinen, I., Denier van der Gon, H. A. C., Charalampidis, P. E., Pilinis, C., Wiedensohler, A., O'Dowd, C. D., Putaud, J. P., Moerman, M., and Pandis, S. N.: Simulating ultrafine particle formation in Europe using a regional CTM: contribution of primary emissions versus secondary formation to aerosol number concentration. *Atmos. Chem. Phys.*, 12, 8663-8677, 2012.

810 Gettelman, A., Lamboll, R., Bardeen C. G., Forster, P. M., and Watson-Paris, D.: Climate impacts of COVID-19 induced emission changes, *Geophys. Res. Lett.*, 48, e2020GL091805, 2020.

Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kurten, A., Lehtipalo, K., Makhtutov, V., Molteni, U., Rissanen, M. P., Stozkhov, Y., Tröstl, J., Tsagkogeorgs, G., Wagner, R., Williamsson, C., Wimmer, D., Winkler, P. M. Yan, C., and Carslaw, K. S.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres. *J. Geophys. Res.: Atmos.*, 122, 8739–8760, 2017.

Green, J. K., Konings, A. G., Alemohammad, S. H., Berry, J., Entekhabi, D., Kolassa, J., Lee, J.-E., and Gentile, P.: Regionally strong feedbacks between the atmosphere and terrestrial biosphere, *Nature Geosci.*, 10, 410-413, 2017.

820 Guo, S., Hu, M., Peng, J., Wu, Z., Zamora, M. L., Shang, D., Du, Z., Zheng, J., Fang, X., Tang, R., Wu, Y., Zeng, L., Shuai, S., Zhang, W., Wang, Y., Ji, Y., Li, Y., Zhang, A. L., Wang, W., Zhang, F., Zhao, J., Gong, X., Wang, C., Molina, M. J., and Zhang, R.: Remarkable Nucleation and Growth of Ultrafine Particles from Vehicular Exhaust, *Proc Natl Acad Sci U A*, 117 (7), 3427–3432, <https://doi.org/10.1073/pnas.1916366117>, 2020.

825 Guo, H.: Steps to the digital Silk Road, *Nature*, 554, 25-27.

Hakala, S., Alghamdi, M. A., Paasonen, P., Vakkari, V., Khoder, M. I., Neitola, K., Dada, L., Abdelmaksoud, A. S., Al-Jeelani, H., Shabbaj, I. I., Almehmadi, F. M., Sundström, A.-M., Lihavainen, H., Kerminen, V.-M., Kontkanen, J., Kulmala, M.,

Hussein, T., and Hyvärinen, A.-P.: New particle formation, growth and apparent shrinkage at a rural background site in western
830 Saudi Arabia, *Atmos. Chem. Phys.*, 19, 10537–10555, <https://doi.org/10.5194/acp-19-10537-2019>, 2019.

Hamed, A., Birmili, W., Joutsensaari, J., Mikkonen, S., Asmi, A., Wehner, B., Spindler, G., Jaatinen, A., Wiedensohler, A.,
Korhonen, H., Lehtinen, K. E. J., and Laaksonen, A.: Changes in the production rate of secondary particles in Central Europe
in view of decreasing SO₂ emissions between 1996 and 2006, *Atmos. Chem. Phys.*, 10, 1071-1091, 2010.

835

Hamill, P., Turco, R.P., Kiang, C.S., Toon, O.B., and Whitten, R.C.: An analysis of various nucleation mechanisms for sulfate
particles in the stratosphere, *J. Aerosol. Sci.*, 13, 561-585, 1982.

Hari, P. and Kulmala, M.: Station for Measuring Ecosystem–Atmosphere Relations (SMEAR II), *Boreal Env. Res.*, 10, 315–
840 322, 2005.

Hari, P., Petäjä, T., Bäck, J., Kerminen, V-M., Lappalainen, H.K. Vihma, T., Laurila, T., Viisanen, Y., Vesala, T., and Kulmala
M.: Conceptual design of a measurement network of the global change, *Atmos. Chem. Phys.*, 16, 1017-1028, doi:10.5194/acp-
16-1017-2016, 2016.

845

He, X.-C., Tham, Y. J., Dada, L., Wang, M., Finkenzeller, H., Stolzenburg, D., Iyer, S. Simon, M., Kürten, A., Shen, J., Rörup,
B., Rissanen, M., Schobesberger, S., Baalbaki, Wang, D. S., Koenig, T. K., Jokinen, T., Sarnela, N. Beck, L., Almeida, J.,
Amanatidis, S., Amorim1, A., Ataei, F., Baccarini, A., Bertozzi, B., Bianchi, F., Brilke, S., Caudillo, L., Chen, D., Chiu, R.,
Chu, B., Dias, A., Ding, A., Dommen, J., Duplissy, J., El Haddad, I., Carracedo, L. G., Granzin, M., Hansell, A., Heinritzi,
850 Hofbauer, V., Junninen, H., Kangasluoma, J., Kemppainen, D., Kim, C., Kong, W., Krechmer, J. E., Kvashin, A., Laitinen,
T., Lamkaddam, H., Lee, C. P., Lehtipalo, K., Leiminger, M., Li, Z., Makhmutov, V., Manninen, H. E., Marie, G., Marten, R.,
Mathot, S., Mauldin, R. L., Mentler, B. Möhler, O., Müller, T., Nie, W., Onnela, A., Petäjä, T., Pfeifer, J., Philippov, M.,
Ranjithkumar, A., Saiz-Lopez, A., Salma, I., Scholz, W., Schuchmann, S., Schulze, B., Steiner, G., Stozhkov, Y., Tauber, C.,
Tomé, A., Thakur, R. C., Väisänen, O., Vazquez-Pufleau, M., Wagner, A. C., Wang, Y., Weber, S. K., Winkler, P. M., Wu,
855 Y., Xiao, M., Yan, C., Ye, Q., Ylisirniö, A., Zauner-Wieczorek, M., Zha, Q., Zhou, P., Flagan, R. C., Curtius, J., Baltensperger,
U., Kulmala, M., Kerminen, V.-M., Kurtén, T., Donahue, N. M., Volkamer, R., Kirkby, J., Worsnop, D. R. and Sipilä, M.:
Role of iodine oxoacids in atmospheric aerosol nucleation. *Science*, 371, 589-595, 2021.

Hellén, H., Praplan, A. P., Tykkä, T., Ylivinkka, I., Vakkari, V., Bäck, J., Petäjä, T., Kulmala, M., and Hakola, H.: Long-term
860 measurements of volatile organic compounds highlight the importance of sesquiterpenes for the atmospheric chemistry of a
boreal forest, *Atmos. Chem. Phys.*, 18, 13839-13863, <https://doi.org/10.5194/acp-18-13839-2018>, 2018.

Holben, B.N., Eck, T.F., Slutsker, I., Tanré, D., Buis, J.P., Setzer, A., Vermote, E., Reagan, J.A., Kaufman, Y.J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET—A Federated Instrument Network and Data Archive
865 for Aerosol Characterization, *Remote Sens. Environ.*, 66, 1-16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.

Isokäntä, S., Kim, P., Mikkonen, S., Kühn, T., Kokkola, H., Yli-Juuti, T., Heikkinen, L., Luoma, K., Petäjä, T., Kipling, Z., Partridge, D., and Virtanen, A.: The effect of clouds and precipitation on the aerosol concentrations and composition in a boreal forest environment, *Atmos. Chem. Phys.*, 22, 11823–11843, <https://doi.org/10.5194/acp-22-11823-2022>, 2022.

870

IPCC: Climate Change 2013: The Physical Science Basis, fifth assessment report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and Midgley, P.M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

875

IPCC: Climate Change 2021: The Physical Science Basis, sixth assessment of the Inter-governmental Panel on Climate Change, edited by: Masson-Delmotte, V., P. Zhai, A., Pirani, S. L., Connors, C., Péan, S., Berger, N., Caud, Y., Chen, L., Goldfarb, M. I., Gomis, M., Huang, K., Leitzell, E., Lonnoy, J. B. R., Matthews, T. K., Maycock, T., Waterfield, O., Yelekçi, R., Yu., and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.

880

Jiang, Z., Shi, H., Zhao, B., Gu, Y., Zhu, Y., Miyazaki, K., Lu, X., Zhang, Y., Bowman, K. W., Sekiya, T., and Liou, K-N.: Modeling the impact of COVID-19 on air quality in southern California: implications for future control policies, *Atmos. Chem. Phys.*, 21, 8693-8708, 2021.

890

Jokinen, T., Sipilä, M., Kontkanen, J., Vakkari, V., Tisler, P., Duplissy, E.-M., Junninen, H., Kangasluoma, J., Manninen, H., Petäjä, T., Kulmala, M., Worsnop, D. R., Kirkby, J., Virkkula, A., and Kerminen, V.-M.: Ion induced sulfuric acid-ammonia nucleation drives particle formation in coastal Antractica. *Sci. Adv.*, 4, eaat9744, 2018.

Junninen, H., Lauri, A., Keronen, P., Aalto, P., Hiltunen, V., Hari, P., and Kulmala, M.: Smart-SMEAR: on-line data exploration and visualization tool for SMEAR stations, *Boreal Environ Res*, 14, 447-457, 2009.

895

Kalivitis, N., Kerminen, V.-M., Kouvarikis, G., Stavroulas, I., Tzitzikalaki, E., Kalkavouras, P., Daskalakis, N., Myriokefalitakis, S., Bougiatioti, A., Manninen, H. E., Roldin, P., Petäjä, T., Boy, M., Kulmala, M., Kanakidou, M., and Mihalopoulos, N.: Formation and growth of atmospheric nanoparticles in the eastern Mediterranean: results from long-term measurements and process simulation. *Atmos. Chem. Phys.*, 19, 2671-2686, 2019.

895

Kerminen, V.-M., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M., and Bianchi F.: Atmospheric new particle formation and growth: review of field observations, Environ. Res. Lett., 13, 103003, 2018.

Kezoudi, M., Keleshin, C., Antoniou, P., Biskos, G., Bronz, M., Constantinides, C., Desservetaz, M., Gao, R.-S., Girdwood, J., Harnetiaux, J., Kandler, K., Leonidou, A., Liu, Y., Lelieveld, J., Marenco, F., Mihalopoulos, N., Mocnik, G., Neitola, K., Paris, J.-D., Pikridas, M., Sarda-Esteve, R., Stopford, C., Unga, F., Vrekoussis, M. and Sciare, J.: The Unmanned Systems Research Laboratory (USRL): A new facility for UAV-based atmospheric observations, Atmosphere 12, 1042, <https://doi.org/10.3390/atmos12081042>, 2021.

905 Khadir, T., Riipinen, I., Isokääntä, S., Heslin-Rees, D., Pöhlker, C., Rizzo, L., Machado, L., Franco, M., Kremper, L. A., Artaxo, P., Petäjä, T., Kulmala, M., Tunved, P., Ekman, A. M. L., Krejci, R., and Virtanen, A.: Sink, source or something in-between? Net effects of precipitation on aerosol particle populations, 2023, submitted manuscript.

Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagne, S., Ickes, L., Kurten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkila, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petaja, T., Schnitzhofer, R., Seinfeld, J. H., Sipila, M., Stozhkov, Y., Stratmann, F., Tome, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, Nature, 476, 429-U477, 10.1038/nature10343, 2011.

920 Kulmala, M., Arola, A., Nieminen, T., Riuttanen, L., Sogacheva, L., de Leeuw, G., Kerminen, V.-M., and Lehtinen, K. E. J.: The first estimates of global nucleation mode aerosol concentrations based on satellite measurements, Atmos. Chem. Phys., 11, 10791–10801, <https://doi.org/10.5194/acp-11-10791-2011>, 2011.

925 Kulmala, M., Nieminen, T., Chellapermal, R., Makkonen, R., Bäck, J., and Kerminen, V.-M.: Climate feedbacks linking the increasing atmospheric CO₂ concentration, BVOC emissions, aerosols and clouds in forest ecosystems. In: Biology, Controls and Models of Tree Volatile Organic Compound Emissions (edited by Ü Niinemets and R. K Monson), Springer, pp.489-508, 2013.

Kulmala, M., Nieminen, T., Nikandrova, A., Lehtipalo, K., Manninen, H. E., Kajos, M. K., Kolari, P., Lauri, A., Petäjä, T., Krejci, R., Hansson, H.-C., Swietlicki, E., Lindroth, A., Christensen, T. R., Arneth, A., Hari, P., Bäck, J., Vesala, T., and

930 Env. Res., 19, suppl. B, 122-131, 2014a.

Kulmala, M., Petäjä, T., Ehn, M., Thornton, J., Sipilä, M., Worsnop, D. R., and Kerminen, V.-M.: Chemistry of atmospheric nucleation: On the recent advances on precursor characterization and atmospheric cluster composition in connection with atmospheric new particle formation, Annu. Rev. Phys. Chem., 65, 21-37, 2014b.

935

Kulmala, M.: China's chocking air cocktail, Nature, 526, 497–499, doi:10.1038/526497a, 2015.

Kulmala, M., Lappalainen, H. K., Petäjä, T., Kurten, T., Kerminen, V.-M., Viisanen, Y., Hari, P., Sorvari, S., Bäck, J., Bondur,

V., Kasimov, N., Kotlyakov, V., Matvienko, G., Baklanov, A., Guo, H. D., Ding, A., Hansson, H.-C., and Zilitinkevich, S.:

940 Introduction: The Pan-Eurasian Experiment (PEEX) – multidisciplinary, multiscale and multicomponent research and capacity-building initiative, Atmos. Chem. Phys., 15, 13085–13096, <https://doi.org/10.5194/acp-15-13085-2015>, 2015.

Kulmala, M.: Build a global Earth observatory. Nature, 553, 21-23, doi: 10.1038/d41586-017-08967-y, 2018.

945 Kulmala, M., Kerminen, V.-M., Petäjä, T., Ding, A. J., and Wang, L.: Atmospheric gas-to-particle conversion: why NPF events are observed in megacities?, Faraday Discuss., 200, 271-288, doi:10.1039/c6fd00257a, 2017.

Kulmala, M., Ezhova, E., Kalliokoski, T., Noe, S., Vesala, T., Lohila, A., Liski, J., Makkonen, R., Bäck, J., Petäjä, T., and Kerminen, V.M.: CarbonSink plus - Accounting for multiple climate feedbacks from forests, Boreal Environment Research,

950 25, 145-159, 2020.

Kulmala, M., Junninen, H., Dada, L., Salma, I., Weidinger, T., Thén, W., Vörösmarty, M., Komsaare, K., Stolzenburg, D., Cai, R., Yan, C., Li, X., Deng, C., Jiang, J., Petäjä, T., Nieminen, T., and Kerminen, V.-M.: Quiet New Particle Formation in the Atmosphere. Front. Environ. Sci., 10, 912385, doi: 10.3389/fenvs.2022.912385, 2022a.

955

Kulmala, M., Cai, R., Stolzenburg, D., Zhou, Y., Dada, L., Guo, Y., Yan, C., Petäjä, T., Jiang, J. and Kerminen, V.-M.: The contribution of new particle formation and subsequent growth to haze formation, Environ. Sci.: Atmos., 2, 352-361, 2022b.

Kulmala, M., Lintunen, A., Ylivinkka, I., Mukkala, J., Rantanen, R., Kujansuu, J., Petäjä, T., and Lappalainen, H.K.:

960 Atmospheric and ecosystem big data providing key contributions in reaching United Nations' sustainable development goals, Big Earth Data, 5, 3, 277-305, doi: 10.1080/20964471.2021.1936943, 2021.

Kulmala, M., Kokkonen, T., Ezhova, E., Baklanov, A., Mahura, A., Mammarella, I., Bäck J., Lappalainen, H., Tyuryakov, S., Kerminen, V.-M., Zilitinkevich, S., and Petäjä, T. Aerosols, Clusters, Greenhouse Gases, Trace Gases and Boundary-Layer
965 Dynamics: on Feedbacks and Interactions. *Boundary-Layer Meteorology*, 186, 475–503, 2023.

Kriegler, E., Luderer, G., Bauer, N., Baumstark, L., Fujimori, S., Popp, A., Rogelj, J., Strefler, J., and van Vuuren, D. P.: Pathways limiting warming to 1.5 degrees C: a tale of turning around in no time?, *Philos T R Soc A*, 376, ARTN 20160457
10.1098/rsta.2016.0457, 2018.

970

Kroll, J. H., Heald, C. L., Cappa, C. D., Farmer, D. K., Fry, J. L., Murphy, J. G., and Steiner, A. L.: The complex chemical effects of COVID-19 shutdowns on air quality, *Nature Chem.*, 12, 777–779, <https://doi.org/10.1038/s41557-020-0535-z>, 2020.

975

Kyrö, E.-M., Väänänen, R., Kerminen, V.-M., Virkkula, A., Petäjä, T., Dal Maso, M., Nieminen, T., Juhola, S., Shcherbinin, A., Riipinen, I., Lehtipalo, K., Keronen, P., Aalto, P. P., Hari, P., and Kulmala, M.: Trends in new particle formation in eastern Lapland, Finland: effect of decreasing sulfur emissions from Kola Peninsula, *Atmos. Chem. Phys.*, 14, 4383-4396, 2014.

980

Köberle, A. C., Vandyck, T., Guivarch, C., Macaluso, N., Bosetti, V., Gambhir, A., Tavoni, M., and Rogelj, J.: The cost of mitigation revisited, *Nat Clim Change*, 11, 1035-1045, 10.1038/s41558-021-01203-6, 2021.

980

985

Lappalainen, H. K., Kerminen, V. M., Petäjä, T., Kurten, T., Baklanov, A., Shvidenko, A., Bäck, J., Vihma, T., Alekseychik, P., Andreae, M. O., Arnold, S. R., Arshinov, M., Asmi, E., Belan, B., Bobylev, L., Chalov, S., Cheng, Y., Chubarova, N., de Leeuw, G., Ding, A., Dobrolyubov, S., Dubtsov, S., Dyukarev, E., Elansky, N., Eleftheriadis, K., Esau, I., Filatov, N., Flint, M., Fu, C., Glezer, O., Gliko, A., Heimann, M., Holtslag, A. A. M., Hörrak, U., Janhunen, J., Juhola, S., Järvi, L., Järvinen, H., Kanukhina, A., Konstantinov, P., Kotlyakov, V., Kieloaho, A. J., Komarov, A. S., Kujansuu, J., Kukkonen, I., Duplissy, E. M., Laaksonen, A., Laurila, T., Lihavainen, H., Lisitzin, A., Mahura, A., Makshtas, A., Mareev, E., Mazon, S., Matishov, D., Melnikov, V., Mikhailov, E., Moisseev, D., Nigmatulin, R., Noe, S. M., Ojala, A., Pihlatie, M., Popovicheva, O., Pumpanen, J., Regerand, T., Repina, I., Shcherbinin, A., Shevchenko, V., Sipilä, M., Skorokhod, A., Spracklen, D. V., Su, H., Subetto, D. A., Sun, J., Terzhevik, A. Y., Timofeyev, Y., Troitskaya, Y., Tynkkynen, V. P., Kharuk, V. I., Zaytseva, N., Zhang, J., Viisanen, Y., Vesala, T., Hari, P., Hansson, H. C., Matvienko, G. G., Kasimov, N. S., Guo, H., Bondur, V., Zilitinkevich, S., and Kulmala, M.: Pan-Eurasian Experiment (PEEX): towards a holistic understanding of the feedbacks and interactions in the land-atmosphere-ocean-society continuum in the northern Eurasian region, *Atmos. Chem. Phys.*, 16, 14421-14461, 10.5194/acp-16-14421-2016, 2016.

995

Lappalainen, H. K., Petäjä, T., Vihma, T., Räisänen, J., Baklanov, A., Chalov, S., Esau, I., Ezhova, E., Leppäranta, M., Pozdnyakov, D., Pumpanen, J., Andreae, M. O., Arshinov, M., Asmi, E., Bai, J., Bashmachnikov, I., Belan, B., Bianchi, F.,

Biskaborn, B., Boy, M., Bäck, J., Cheng, B., Chubarova, N., Duplissy, J., Dyukarev, E., Eleftheriadis, K., Forsius, M., Heimann, M., Juhola, S., Konovalov, V., Konovalov, I., Konstantinov, P., Köster, K., Lapshina, E., Lintunen, A., Mahura, A., Makkonen, R., Malkhazova, S., Mammarella, I., Mammola, S., Buenrostro Mazon, S., Meinander, O., Mikhailov, E., Miles, 1000 V., Myslenkov, S., Orlov, D., Paris, J.-D., Pirazzini, R., Popovicheva, O., Pulliainen, J., Rautiainen, K., Sachs, T., Shevchenko, V., Skorokhod, A., Stohl, A., Suhonen, E., Thomson, E. S., Tsidilina, M., Tynkkynen, V.-P., Uotila, P., Virkkula, A., Voropay, N., Wolf, T., Yasunaka, S., Zhang, J., Qiu, Y., Ding, A., Guo, H., Bondur, V., Kasimov, N., Zilitinkevich, S., Kerminen, V.-M., and Kulmala, M.: Overview: Recent advances in the understanding of the northern Eurasian environments and of the urban air quality in China – a Pan-Eurasian Experiment (PEEX) programme perspective, *Atmos. Chem. Phys.*, 22, 4413–4469, 1005 <https://doi.org/10.5194/acp-22-4413-2022>, 2022.

Laughner, J. L., Neu, J. L., Schimel, D., Wennberg, P. O., Barsanti, K., Bowman, K. W., Chatterjee, A., Croes, B. E., Fitzmaurice, H. L., Henze, D. K., Kim, J., Kort, E. A., Liu, Z., Miyazaki, K., Turner, A. J., Anenberg, S., Avise, J., Cao, H., Crisp, D., de Gouw, J., Eldering, A., Fyfe, J. C., Goldberg, D. L., Gurney, K. R., Hasheminassab, S., Hopkins, F., Ivey, C. E., 1010 Jones, D. B. A., Liu, J., Lovenduski, N. S., Martin, R. V., McKinley, G. A., Ott, L., Poulter, B., Ru, M., Sander, S. P., Swart, N., Yung, Y. L., and Zeng, Z.-C.: Societal shifts due to COVID-19 reveal large-scale complexities and feedbacks between atmospheric chemistry and climate change, *Proceedings of the National Academy of Sciences*, 118, e2109481118, doi:10.1073/pnas.2109481118, 2021.

Lehtipalo, K., Yan, C., Dada, L., Bianchi, F., Xiao, M., Wagner, R., Stolzenburg, D., Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bernhamme,r A.-K., Breitenlechner, M., Brilke, S., Buckholz, A., Buenrostro Mazon, S., Chen, D., Chen, X., Dias, A., Dommen, J., Draper, D. C., Duplissy, J., Ehn, M., Finkenzeller, H., Fischer, L., Frege, C., Fuchs, C., Garmash, O., Gordon, H., Hakala, J., He, X., Heikkinen, L., Heinrizi, M., Helm, J. C., Hofbauer, V., Hoyle, C. R., Jokinen, T., Kangasluoma, J., Kerminen, V.-M., Kim, C., Kirkby, J., Kontkanen, J., Kürten, A., Lawler, M. J., Mai, H., 1015 Mathot, S., Mauldin III, R. L., Molteni, U., Nichman, L., Nie, W., Nieminen, T., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Piel, F., Pospisilova, V., Quéléver, L. L. J., Rissanen, M. P., Rose, C., Sarnela, N., Schallhart, S., Schuchmann, S., Sengupta, K., Simon, M., Sipilä, M., Tauber, K., Tomé, A., Tröstl, J., Väisänen, O., Vogel, A. L., Volkamer, A., Wagner, A. C., Wang, M., Weitz, L., Wimmer, D., Ye, P., Ylisirniö, A., Zha, Q., Carslaw, K. S., Curtius, J., Donahue, N. M., Flagan, R. C., Hansel, A., Riipinen, I., Virtanen, A., Winkler, P. M., Baltensperger, U., Kulmala, M., and Worsnop, D. R.: 1020 Multicomponent new particle formation from sulfuric acid, ammonia, and biogenic vapors. *Sci. Adv.*, 4, eaau5363, 2018.

Leinonen, V., Kokkola, H., Yli-Juuti, T., Mielonen, T., Kühn, T., Nieminen, T., Heikkinen, S., Miinalainen, T., Bergman, T., Carslaw, K., Decesari, S., Fiebig, M., Hussein, T., Kivekäs, N., Krejci, R., Kulmala, M., Leskinen, A., Massling, A., Mihalopoulos, N., Mulcahy, J. P., Noe, S. M., van Noije, T., O'Connor, F. M., O'Dowd, C., Olivie, D., Pernov, J. B., Petäjä, T., Selander, Ø., Schulz, M., Scott, C. E., Skov, H., Swietlicki, E., Tuch, T., Wiedensohler, A., Virtanen, A., and Mikkonen, S.: 1030

Comparison of particle number size distribution trends in ground measurements and climate models, *Atmos. Chem. Phys.*, 22, 12873–12905, <https://doi.org/10.5194/acp-22-12873-2022>, 2022.

Lenton, T. M., Rockström, J., Gaffney, O., Rahmstorf, S., Richardson, K., Steffen, W., and Schellnhuber, H. J.: Climate tipping points - too risky to bet against, *Nature*, 575, 592-595, DOI 10.1038/d41586-019-03595-0, 2019.

Lintunen, A., Aalto, J., Asmi, A., Aurela, M., Bäck, J., Ehn, M., Ezhova, E., Hakola, H., Hartonen, K., Heinonsalo, J., Hellén, H., Hölttä, T., Jokinen, T., Järvi, L., Järvinen, H., Kangasluoma, J., Kerminen, V.-M., Köster, K., Köster, E., Kulmala, L., Kurten, T., Laaksonen, A., Lappalainen, H.K., Laurila, T., Lehtipalo, K., Lihavainen, H., Lohila, A., Mäkelä, A., Mäki, M.,
1040 Makkonen, R., Mikkonen, S., Moisseev, D., Ojala, A., Petäjä, T., Pihlatie, M., Porcar-Castell, A., Praplan, A.P., Pulliainen, J., Pumpanen, J., Rantala, P., Riekola, M.-L., Rissanen, K., Romakkaniemi, S., Ruiz-Jimenez, J., Sarnela, N., Schallhart, S., Schiestl-Aalto, P., Rinne, J., Tuittila, E.-S., Vehkämäki, H., Viisanen, Y., Virtanen, A., Ylivinkka, I., Hari, P., and Kulmala, M.: The Center of Excellence in Atmospheric Science (2002-2019) - from molecular and biological processes to the global climate. *Boreal Env. Ress*, 28, 15-80, 2023.

1045 Liu, Y., Yan, C., Feng, Z., Zheng, F., Fan, X., Zhang, Y., Li, C., Zhou, Y., Lin, Z., Guo, Y., Zhang, Y., Ma, L., Zhou, W., Liu, Z., Dada, L., Dällenbach, K., Kontkanen, J., Cai, R., Chan, T., Chu, B., Du, W., Yao, L., Wang, Y., Cai, J., Kangasluoma, J., Kokkonen, T., Kujansuu, J., Rusanen, A., Deng, C., Fu, Y., Yin, R., Li, X., Lu, Y., Liu, Y., Lian, C., Yang, D., Wang, W., Ge, M., Wang, Y., Worsnop, D. R., Junninen, H., He, H., Kerminen, V.-M., Zheng, J., Wang, L., Jiang, J., Petäjä, T., Bianchi, F.,
1050 and Kulmala, M.: Continuous and comprehensive atmospheric observations in Beijing: a station to understand the complex urban atmospheric environment, *Big Earth Data*, 4, 295-321, 10.1080/20964471.2020.1798707, 2020.

Loescher, H. W., Vargas, R., Mirtl, M., Morris, B., Pauw, J., Yu, X., Kutsch, W., Mabee, P., Tang, J., Ruddell, B.L., Pulsifer, P., Bäck, J., Zacharias, S., Grant, M., Feig, G., Zheng L., Waldmann C., and Genazzio, M. A.: Building a Global Ecosystem Research Infrastructure to address global grand challenges for macrosystem ecology, *Earth's Future*, 10, e2020EF001696,
1055 <https://doi.org/10.1029/2020EF001696>, 2022.

Mäkelä, J. M., Aalto, P., Jokinen, V., Pohja, T., Nissinen, A., Palmroth, S., Markkanen, T., Seitsonen, K., Lihavainen, H., and Kulmala, M.: Observations of ultrafine aerosol particle formation and growth in boreal forest. *Geophys. Res. Lett.*, 24, 1219-1222, 1997.

Malila, J.: On the early studies recognizing the role of sulfuric acid in atmospheric haze and new particle formation, *Tellus*, 70, 1471913, 2018.

- 1065 Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global CCN, Atmos. Chem. Phys., 9, 8601–8616, 2009.
- Mizoguchi, Y., Miyata, A., Ohtani, Y., Hirata, R., and Yuta, S.: A review of tower flux observation sites in Asia, J. Forest. Res., 14, 1-9, <https://doi.org/10.1007/s10310-008-0101-9>, 2008.
- 1070 Motlagh, N.H., Irjala, M., Zuniga, A., Lagerspetz, E., Rantala, V., Flores, H., Nurmi, P., and Tarkoma, S.: Toward Blue Skies: City-Scale Air Pollution Monitoring Using UAVs, IEEE Consum. Electron. Mag., 12, 21-31, doi: 10.1109/MCE.2022.3167800, 2023.
- 1075 Msemburi, W., Karlinsky, A., Knutson, V., Aleshin-Guendel, S., Chatterji, S., and Wakefield, J.: The WHO estimates of excess mortality associated with COVID-19 pandemic, Nature, 613, 130-137, 2023.
- Neefjes, I., Laapas, M., Liu, Y., Medus, E., Miettunen, E., Ahonen, L., Quelever, L., Aalto, J., Bäck, J., Kerminen, V.-M., Lampilahti, J., Luoma, K., Mäki, M., Mammarella, I., Petäjä, T., Räty, M., Sarnela, N., Ylivinkka, I., Hakala, S., Kulmala, M.,
1080 Nieminen, T., and Lintunen, A.: 25 years of atmospheric and ecosystem measurements in a boreal forest – Seasonal variation and responses to warm and dry years, Boreal Env. Res., 27, 1-31, 2022.
- Nieminen, T., Asmi, A., Dal Maso, M., Aalto, P. P., Keronen, P., Petäjä, T., Kulmala, M., and Kerminen, V.-M.: Trends in atmospheric new-particle formation: 16 years of observations in a boreal-forest environment, Boreal Env. Res., 19, Suppl. B,
1085 191-214, 2014.
- Nieminen, T., Kerminen, V.-M., Petäjä, T., Aalto, P. P., Arshinov, M., Asmi, E., Baltensperger, U., Beddows, D. S. C., Beukes, J. P., Collins, D., Ding, A., Harrison, R. M., Henzing, B., Hooda, R., Hu, M., Horrak, U., Kivekäs, N., Komsaare, K., Krejci, R., Kristensson, A., Laakso, L., Laaksonen, A., Leaitch, W. R., Lihavainen, H., Mihalopoulos, N., Nemeth, Z., Nie, W.,
1090 O'Dowd, C. D., Salma, I., Sellegrí, K., Svenssonsson, B., Swietlicki, E., Tunved, P., Ulevicius, V., Vakkari, V., Vana, M., Wiedensohler, A., Wu, Z., Virtanen, A., and Kulmala M.: Global analysis of continental boundary layer new particle formation based on long-term measurements, Atmos. Chem. Phys., 18, 14737-14756, 2018.
- van Noije, T., Bergman, T., Le Sager, P., O'Donnell, D., Makkonen, R., Gonçalves-Ageitos, M., Döscher, R., Fladrich, U.,
1095 von Hardenberg, J., Keskinen, J.-P., Korhonen, H., Laakso, A., Myriokefalakis, S., Ollinaho, P., Pérez García-Pando, C., Reerink, T., Schrödner, R., Wyser, K., and Yang, S.: EC-Earth3-AerChem: a global climate model with interactive aerosols and atmospheric chemistry participating in CMIP6 , Geosci. Model Dev., 14, 5637–5668, <https://doi.org/10.5194/gmd-14-5637-2021>, 2021.

1100 Nolan, C., Overpeck, J. T., Allen, J. R. M., Anderson, P. M., Betancourt, J. L., Binney, H. A., Brewer, S., Bush, M. B., Chase, B. M., Cheddadi, R., Djamali, M., Dodson, J., Edwards, M. E., Gosling, W. D., Haberle, S., Hotchkiss, S. C., Huntley, B., Ivory, S. J., Kershaw, A. P., Kim, S. H., Latorre, C., Leydet, M., Lezine, A. M., Liu, K. B., Liu, Y., Lozhkin, A. V., McGlone, M. S., Marchant, R. A., Momohara, A., Moreno, P. I., Muller, S., Otto-Bliesner, B. L., Shen, C. M., Stevenson, J., Takahara, H., Tarasov, P. E., Tipton, J., Vincens, A., Weng, C. Y., Xu, Q. H., Zheng, Z., and Jackson, S. T.: Past and future global
1105 transformation of terrestrial ecosystems under climate change, *Science*, 361, 920-923, 10.1126/science.aan5360, 2018.

Paasonen P., Nieminen T., Asmi E., Manninen H. E., Petäjä T., Plass-Dülmer C., Flentje H., Birmili W., Wiedensohler A., Horak U., Metzger A., Hamed A., Laaksonen A., Facchini M. C., Kerminen V.-M., and Kulmala M.: On the roles of sulphuric acid and low-volatility organic vapours in the initial steps of atmospheric new particle formation, *Atmos. Chem. Phys.*, 10,
1110 11223-11242, 2010.

Paasonen, P., Asmi, A., Petäjä, T., Kajos, M. K., Äijälä, M., Junninen, H., Holst, T., Abbatt J. P. D., Arneth, A., Birmili, W., Denier van der Gon, H., Hamed, A., Hoffer, A., Laakso L., Laaksonen, A., Leaitch, W. R., Plass-Dulmer, C., Pryor, S. C., Räisänen, P., Swietlicki, E., Wiedensohler, A., Worsnop, D. R., Kerminen, V.-M., and Kulmala, M.: Warming-induced
1115 increase in aerosol number concentration likely to moderate climate change, *Nat. Geosci.*, 6, 438–442. doi: 10.1038/ngeo1800, 2013.

Petäjä, T., Mauldin, I. R. L., Kosciuch, E., McGrath, J., Nieminen, T., Paasonen, P., Boy, M., Adamov, A., Kotiaho, T., and Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest site, *Atmos. Chem. Phys.*, 9, 7435-7448, 10.5194/acp-9-
1120 7435-2009, 2009.

Petäjä, T., Tabakova, K., Manninen, A., Ezhova, E., O'Connor, E., Moisseev, D., Sinclair, V. A., Backman, J., Levula, J., Luoma, K., Virkkula, A., Paramonov, M., Räty, M., Äijälä, M., Heikkinen, L., Ehn, M., Sipilä, M., Yli-Juuti, T., Virtanen, A., Ritsche, M., Hickmon, N., Pulik, G., Rosenfeld, D., Worsnop, D. R., Bäck, J., Kulmala, M., and Kerminen, V.-M.: Influence
1125 of biogenic emissions from boreal forests on aerosol–cloud interactions, *Nat. Geosci.* 15, 42–47, 2022.

Praske, E., Otkjaer, R. V., Crounse, J. D., Hethcox, J. C., Stoltz, B. M., Kjaergaard, H. G., and Wennberg, P. O.: Atmospheric Autoxidation Is Increasingly Important in Urban and Suburban North America, *Proc Natl Acad Sci U A*, 115 (1), 64–69,
<https://doi.org/10.1073/pnas.1715540115>, 2018.

1130

Pseftogkas, A., Koukouli, M.-E., Segers, A., Manders, A., Geffen, Jv., Balis, D., Meleti, C., Stavrakou, T., and Eskes H.: Comparison of S5P/TROPOMI Inferred NO₂ Surface Concentrations with In Situ Measurements over Central Europe, *Remote Sensing*, 14, 4886, <https://doi.org/10.3390/rs14194886>, 2022.

1135 Qi, X., Ding, A., Roldin, P., Xu, Z., Putian, Z., Sarnela, N., Nie, W., Huang, X., Rusanen, A., Ehn, M., Rissanen, M. P., Petäjä, T., Kulmala, M., and Boy, M.: Modelling studies of HOMs and their contributions to new particle formation and growth: comparison of boreal forest in Finland and a polluted environment in China, *Atmos. Chem. Phys.*, 18, 1179-11791, 2018.

Raes, F., Liao, H., Chen, W.-T., and Seinfeld, J. H. J.: Atmospheric chemistry-climate feedbacks, *J. Geophys. Res.*, 115, 1140 D12121, doi:10.1029/2009Jd013300, 2010.

Rantala, P., Aalto, J., Taipale, R., Ruuskanen, T. M., and Rinne J.: Annual cycle of volatile organic compound exchange between a boreal pine forest and the atmosphere, *Biogeosci.*, 12, 5753-5770, 2015.

1145 Rap, A., Scott, C. E., Reddington, C. L., Mercado, L., Ellis, R. J., Garraway, S., Evans, M. J., Beerling, D. J., MacKenzie, A. R., Hewitt, C. N., and Spracklen, D. V.: Enhanced global primary production by biogenic aerosol via diffuse radiation fertilization, *Nature Geoscience*, 11, 640-644, 10.1038/s41561-018-0208-3, 2018.

Räty, M., Sogacheva, L., Keskinen, H.-M., Kerminen, V.-M., Nieminen, T., Petäjä, T., Ezhova, E., and Kulmala, M.: Dynamics 1150 of aerosol, humidity, and clouds in air masses travelling over Fennoscandian boreal forests, *Atmos. Chem. Phys. (in press)*, 2023.

Rebeiro-Hargrave, A., Fung, P. L., Varjonen, S., Huertas, A., Sillanpää, S., Luoma, K., Hussein, T., Petäjä, T., Timonen, H., Limo, J., Nousiainen, V., and Tarkoma, S.: City Wide Participatory Sensing of Air Quality, *Frontiers in Environmental 1155 Science*, 9, 10.3389/fenvs.2021.773778, 2021.

Ren, J. Chen, L., Fan, T., Liu, J., Jiang, S., and Zhang, F.: The NPF effect on CCN number concentration: a review and re-evaluation of observations from 35 sites worldwide, *Geophys. Res. Lett.*, 48, e2021GL095190, 2021.

1160 Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F. S., Lambin, E. F., Lenton, T. M., Scheffer, M., Folke, C., Schellnhuber, H. J., Nykvist, B., de Wit, C. A., Hughes, T., van der Leeuw, S., Rodhe, H., Sörlin, S., Snyder, P. K., Costanza, R., Svedin, U., Falkenmark, M., Karlberg, L., Corell, R. W., Fabry, V. J., Hansen, J., Walker, B., Liverman, D., Richardson, K., Crutzen, P., and Foley, J. A.: A safe operating space for humanity, *Nature*, 461, 472-475, 10.1038/461472a, 2009.

Rönkkö, T., Kuuluvainen, H., Karjalainen, P., Keskinen, J., Hillamo, R., Niemi, J. V., Pirjola, L., Timonen, H. J., Saarikoski, S., Saukko, E., Järvinen, A., Silvennoinen, H., Rostedt, A., Olin, M., Yli-Ojanpera, J., Nousiainen, P., Kousa, A., and Dal Maso, M.: Traffic is a major source of atmospheric nanocluster aerosol, Proc. Natl. Acad. Sci. USA, 114, 7549–7554, <https://doi.org/10.1073/pnas.1700830114>, 2017.

Saha, P. K., Robinson, E., S., Shah, R., U., Zimmerman, N., Apté J. S., Robinson, A. L., and Presto, A. A.: Reduced ultrafine particle concentrations in urban air: Changes in nucleation and anthropogenic emissions, Environ. Sci. Technol., 52, 6798-6806, 2018.

Schreck, J. S., Becker, C., Gagne, D. J., Lawrence, K., Wang, S., Mouchel-Vallon, C., Choi, J., and Hodzic, A.: Neural Network Emulation of the Formation of Organic Aerosols Based on the Explicit GECKO-A Chemistry Model, J Adv Model Earth Sy, 14, e2021MS002974, <https://doi.org/10.1029/2021MS002974>, 2022.

Semeniuk, K., and Dastoor, A.: Current state of aerosol nucleation parameterizations for air-quality and climate modeling, Atmos. Environ., 179, 77-106, 2018.

Sihto, S.-L., Kulmala, M., Kerminen, V.-M., Dal Maso, M., Petäjä, T., Riipinen, I., Korhonen, H., Arnold, F., Janson, R., Boy, M., Laaksonen, A., and Lehtinen, K. E. J.: Atmospheric sulphuric acid and aerosol formation: implications from atmospheric measurements for nucleation and early growth mechanisms, Atmos. Chem. Phys., 6, 4079-4091, 2006.

Sipilä, M., Sarnela, N., Jokinen, T., Henschel, H., Junninen, H., Kontkanen, J., Richters, S., Kangasluoma, J., Franchin, A., Peräkylä, O., Rissanen, M. P., Ehn, M., Vehkamäki, H., Kurten, T., Berndt, T., Petäjä, T., Worsnop, D., Ceburnis, D., Kerminen, V.-M., Kulmala, M., and O'Dowd, C. D.: Molecular-scale evidence of aerosol particle formation via sequential addition of HIO_3 . Nature, 537, 532-534, doi:10.1038/nature19314, 2016.

Smith, P., Davies, C. A., Ogle, S., Zanchi, G., Bellarby, J., Bird, N., Boddey, R. M., McNamara, N. P., Powson, D., Cowie, A., van Nordwijk, M., Davis, S. C., Richter, D. D. B., Kryzanowski, L., van Wijk, M. T., Stuart, J., Kirton, A., Eggar, D., Newton-Cross, G., Adhya, T. K., and Braimoh, A. K.: Towards an integrated global framework to assess the impacts of land use and management change on soil carbon: current capability and future vision, Global Change Biol., 18, 2089-2101, doi: 10.1111/j.1365-2486.2012.02689.x, 2012.

Smith, C., Baker, J. C. A., and Spracklen, D. V.: Tropical deforestation causes large reductions in observed precipitation, Nature, <https://doi.org/10.1038/s41586-022-05690-1>, 2023.

Sokhi, R. S., Singh, V., Querol, X., Finardi, S., Targino, A. C., Andrade, M. d. F., Pavlovic, R., Garland, R. M., Massagué, J., Kong, S., Baklanov, A., Ren, L., Tarasova, O., Carmichael, G., Peuch, V.-H., Anand, V., Arbilla, G., Badali, K., Beig, G.,
1200 Belalcazar, L. C., Bolignano, A., Brimblecombe, P., Camacho, P., Casallas, A., Charland, J.-P., Choi, J., Chourdakis, E., Coll, I., Collins, M., Cyrys, J., da Silva, C. M., Di Giosa, A. D., Di Leo, A., Ferro, C., Gavidia-Calderon, M., Gayen, A., Ginzburg, A., Godefroy, F., Gonzalez, Y. A., Guevara-Luna, M., Haque, S. M., Havenga, H., Herod, D., Hörrak, U., Hussein, T., Ibarra, S., Jaimes, M., Kaasik, M., Khaiwal, R., Kim, J., Kousa, A., Kukkonen, J., Kulmala, M., Kuula, J., La Violette, N., Lanzani, G., Liu, X., MacDougall, S., Manseau, P. M., Marchegiani, G., McDonald, B., Mishra, S. V., Molina, L. T., Mooibroek, D.,
1205 Mor, S., Moussiopoulos, N., Murena, F., Niemi, J. V., Noe, S., Nogueira, T., Norman, M., Pérez-Camaño, J. L., Petäjä, T., Piketh, S., Rathod, A., Reid, K., Retama, A., Rivera, O., Rojas, N. Y., Rojas-Quincho, J. P., San José, R., Sánchez, O., Seguel, R. J., Sillanpää, S., Su, Y., Tapper, N., Terrazas, A., Timonen, H., Toscano, D., Tsegas, G., Velders, G. J. M., Vlachokostas, C., von Schneidemesser, E., Vpm, R., Yadav, R., Zalakeviciute, R., and Zavala, M.: A global observational analysis to understand changes in air quality during exceptionally low anthropogenic emission conditions, *Environment International*,
1210 157, 106818, <https://doi.org/10.1016/j.envint.2021.106818>, 2021.

Spracklen, D. V., Carslaw, K. S., Merikanto, J., Mann, G. W., Reddington, C. L., Pickering, S., Ogren, J. A., Andrews, E., Baltensperger, U., Weingartner, E., Boy, M., Kulmala, M., Laakso, L., Lihavainen, H., Kivekäs, N., Komppula, M., Mihalopoulos, N., Kouvarakis, G., Jennings, S. G., O'Dowd, C., Birmili, W., Wiedensohler, A., Weller, R., Gras, J., Laj, P.,
1215 Sellegri, K., Bonn, B., Krejci, R., Laaksonen, A., Hamed, A., Minikin, A., Harrison, R. M., Talbot, R., and Sun, J.: Explaining global surface aerosol number concentrations in terms of primary emissions and particle formation, *Atmos. Chem. Phys.*, 10, 4775–4793, 10.5194/acp-10-4775-2010, 2010.

Sporre, M. K., Blichner, S. M., Karset, I. H. H., Makkonen, R., and Berntsen, T. K.: BVOC–aerosol–climate feedbacks
1220 investigated using NorESM, *Atmos. Chem. Phys.*, 19, 4763–4782, 2019.

Sporre, M. K., Blichner, S. M., Schrödner, R., Karset, I. H. H., Berntsen, T. K., van Noije, T., Bergman, T., O'Donnell, D., and Makkonen, R.: Large difference in aerosol radiative effects from BVOC-SOA treatment in three Earth system models, *Atmos. Chem. Phys.*, 20, 8953–8973, 2020.

1225 Stevens, B., Bernier, N., Prein, A. F., Baehr, J., Bauer, P., Bockelmann, H., Bony, S., Bresch, D., Brunet, G., Doblas-Reyes, F. J., Ewen, C., Farrell, D., Gruber, N., Hazeleger, W., Hoefler, T., Jacob, D., Jakob, C., Krishnan, R., Li, C., Luterbacher, J., Kulmala, M., Manninen, P., Marotzke, J., Palmer, T., Ramaswamy, V., Rauser, F., Satoh, M., Schulthess, T., Schulz, J., Stammer, D., Shukla, J., Slingo, J., Sobel, A., Stocker, T., Teutsch, G., Tompkins, A., and Zho, T.: Earth Virtualization Engines (EVE), A draft concept paper for public comment, Conveners of the Berlin Summit for EVE, June 5, 26 pages, 2023. (manuscript)

Stocker, B. D., Roth, R., Joos, F., Spahni, R., Steinacher, M., Zaehle, S., Bouwman, L., Ri X., and Prentice I. C.: Multiple greenhouse-gas feedbacks from the land biosphere under future climate change scenarios, *Nat. Clim. Change*, 3, 666-671,

1235 2013.

Stolzenburg, D., Cai, R., Blichner, S. M., Kontkanen, J., Zhou, P., Makkonen, R., Kerminen, V.-M., Kulmala, M., Riipinen, I., and Kangasluoma, J.: Atmospheric nanoparticle growth, *Rev. Mod. Phys.*, 2023 (accepted).

1240 Sundström, A.-M., Nikandrova, A., Atlaskina, K., Nieminen, T., Vakkari, V., Laakso, L., Beukes, J. P., Arola, A., van Zyl, P. G., Josipovic, M., Venter, A. D., Jaars, K., Pienaar, J. J., Piketh, S., Wiedensohler, A., Chiloane, E. K., de Leeuw, G., and Kulmala, M.: Characterization of satellite-based proxies for estimating nucleation mode particles over South Africa, *Atmos. Chem. Phys.*, 15, 4983–4996, <https://doi.org/10.5194/acp-15-4983-2015>, 2015.

1245 Taipale, R., Kajos, M. K., Patokoski, J., Rantala, P., Ruuskanen, T. M., and Rinne, J.: Role of de novo biosynthesis in ecosystem scale monoterpane emissions from a boreal Scots pine forest, *Biogeosci.*, 8, 2247–2255. <https://doi.org/10.5194/bg-8-2247-2011>, 2011.

Taleb, N. N.: *The black swan: the impact of the highly improbable*, 2nd, Random House Trade Paperbacks, New York, 444 p., 2010.

1250

Tarvainen, V., Hakola, H., Hellen, H., Bäck, J., Hari, P., and Kulmala M.: Temperature and light dependence of the VOCemissions of Scots pine, *Atmos. Chem. Phys.* 5, 6691-6718, 2005.

1255 Thornhill, G., Collins, W., Olivie', D., Skeie, R. B., Archibald, A., Bauer, S., Checa-Garcia, R., Fiedler, S., Folberth, G., Gjermundsen, A., Horowitz, L., Lamarque, J.-F., Michou, M., Mulcahy, J., Nabat, P., Naik, V., O'Connor, F. M., Paulot, F., Schulz,M., Scott, C. E., Se'fe'rian, R., Smith, C., Takemura, T., Tilmes, S., Tsigaridis, K., and Weber, J.: Climate-driven chemistry and aerosol feedbacks in CMIP6 Earth system models. *Atmospheric Chemistry and Physics*, 21(2):1105–1126, 2021.

1260 Tuovinen, S., Cai S., Kerminen, V.-M., Jiang, J., Yan, C., Kulmala, M., and Kontkanen, J.: Survival properties of atmospheric particles: comparison based on theory, cluster population simulations, and observations in Beijing, *Atmos. Chem. Phys.*, 22, 15071-15091, 2022.

1265 UNEP. The Adaptation Finance Gap Report 2016. United Nations Environment Programme (UNEP), Nairobi, Kenya, 2016.

UNEP. An independent expert review on Solar Radiation Modification research and deployment, 2023.

Unger N.: Human land-use-driven reduction of forest volatiles cools global climate, *Nature Clim. Change*, 4, 907-910, 2014.

- 1270 Vicente-Saez, R. and Martinez-Fuentes, C.: Open Science now: A systematic literature review for an integrated definition, *J. Business. Res.*, 88, 428-436, <https://doi.org/10.1016/j.jbusres.2017.12.043>, 2018.

- Viatte, C., Petit, J.-E., Yamanouchi, S., Van Damme, M., Doucerain, C., Germain-Piaulenne, E., Gros, V., Favez, O., Clarisse, L., Coheur, P.-F., Strong, K., and Clerbaux, C.: Ammonia and PM_{2.5} Air Pollution in Paris during the 2020 COVID 1275 Lockdown. *Atmosphere*, 12, 160, <https://doi.org/10.3390/atmos12020160>, 2021.

Wandji Nyamsi, W., Lipponen, A., Sanchez-Lorenzo, A., Wild, M., and Arola, A.: A hybrid method for reconstructing the historical evolution of aerosol optical depth from sunshine duration measurements, *Atmos. Meas. Tech.*, 13, 3061–3079, <https://doi.org/10.5194/amt-13-3061-2020>, 2020.

- 1280 Wang, N., Xu, J., Pei, C., Tang, R., Zhou, D., Chen, Y., Li, M., Deng, X., Deng, T., Huang, X., and Ding A.: Air quality during COVID-19 lockdown in the Yangtze River Delta and the Pearl River Delta: two different responsive mechanisms to emission reduction in China, *Environ. Sci. Technol.*, 55, 5721-5730, 2021.

- 1285 Wang, S. Foster, A., Lenz, E. A., Kessler, J. D., Stroeve, J. C., Anderson L. O., Turetsky, M., Betts, R., Zou, S., Liu, W., Boos, W. R., and Hausfather, Z.: Mechanisms and impacts of Earth system tipping elements, *Rev. Geophys.*, 61, e2021RG000757, 2023.

- 1290 Wang, Z., Birmila, W., Hamed A., Wehner, B., Spindler, G., Pei X., Wu, Z., Chaen, Y., Su, H., and Wiedensohler, A.: Contributions of volatile and nonvolatile compounds (at 300 °C) to condensational growth of atmospheric nanoparticles: An assessment based on 8.5 years of observations at the Central Europe background site Melpitz, *J. Geophys. Res.: Atmos.*, 122, 485-497, 2017.

- 1295 Wanner, H., Beer, J., Bütkofer, J., Crowley, T. J., Cubasch, U., Flückiger, J., Goosse, H., Grosjean, M., Joos, F., Kaplan, J., Küttel, M., Müller, S. A., Prentice, I. C., Solomina, O., Stocker, T. F., Tarasov, P., Wagner, M., and Widmann, M.: Mid-to Late Holocene climate change: an overview, *Quaternary Science Reviews*, 27, 1791-1828, <https://doi.org/10.1016/j.quascirev.2008.06.013>, 2008.

Wijnands, J. S., Nice, K. A., Seneviratne, S., Thompson, J., and Stevenson, M.: The impact of the COVID-19 pandemic on air
1300 pollution: A global assessment using machine learning techniques, *Atmos. Poll. Res.*, 13, 101438, 2022.

Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva
Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C.
T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hooft,
1305 R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M.,
van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei,
J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR
Guiding Principles for scientific data management and stewardship, *Sci Data*, 3, 160018, 10.1038/sdata.2016.18, 2016.

1310 WMO Global Atmosphere Watch (GAW) Implementation Plan: 2016-2023, GAW Report No. 228, 2017.

WMO 2022 GCOS Implementation Plan (GCOS-244), [GCOS](#)- No. 244, 2022

Wong, T. E., Cui, Y., Royer, D. L., and Keller, K.: A tighter constraint on Earth-system sensitivity from long-term temperature
1315 and carbon-cycle observations, *Nature Communications*, 12, 3173, 10.1038/s41467-021-23543-9, 2021.

Yan, C., Nie, W., Vogel, A. L., Dada, L., Lehtipalo, K., Stolzenburg, D., Wagner, R., Rissanen, M. P., Xiao, M., Ahonen, L.,
Fischer, L., Rose, C., Bianchi, F., Gordon, H., Simon, M., Heinritzi, M., Garmash, O., Roldin, P., Dias, A., Ye, P., Hofbauer,
V., Amorim, A., Bauer, P. S., Bergen, A., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Buchholz, A., Mazon, S. B.,
1320 Canagaratna, M. R., Chen, X., Ding, A., Dommen, J., Draper, D. C., Duplissy, J., Frege, C., Heyn, C., Guida, R., Hakala, J.,
Heikkinen, L., Hoyle, C. R., Jokinen, T., Kangasluoma, J., Kirkby, J., Kontkanen, J., Kürten, A., Lawler, M. J., Mai, H.,
Mathot, S., Mauldin, R. L., Molteni, U., Nichman, L., Nieminen, T., Nowak, J., Ojdanic, A., Onnela, A., Pajunoja, A., Petäjä,
T., Piel, F., Quéléver, L. L. J., Sarnela, N., Schallhart, S., Sengupta, K., Sipilä, M., Tomé, A., Tröstl, J., Väisänen, O., Wagner,
A. C., Ylsirniö, A., Zha, Q., Baltensperger, U., Carslaw, K. S., Curtius, J., Flagan, R. C., Hansel, A., Riipinen, I., Smith, J.
1325 N., Virtanen, A., Winkler, P. M., Donahue, N. M., Kerminen, V.-M., Kulmala, M., Ehn, M., and Worsnop, D. R.: Size-
dependent influence of NO_x on the growth rates of organic aerosol particles, *Science Advances*, 6, eaay4945,
doi:10.1126/sciadv.aay4945, 2020.

Yan, C., Yin, R., Lu, Y., Dada, L., Yang, D., Fu, Y., Kontkanen, J., Deng, C., Garmash, O., Ruan, J., Baalbaki, R., Schervish,
1330 M., Cai, R., Bloss, M., Chan, T., Chen, T., Chen, Q., Chen, X., Chen, Y., Chu, B., Dällenbach, K., Foreback, B., He, X.,
Heikkinen, L., Jokinen, T., Junninen, H., Kangasluoma, J., Kokkonen, T., Kurppa, M., Lehtipalo, K., Li, H., Li, H., Li, X.,
Liu, Y. Ma, Q., Paasonen, P., Rantala, P., Pileci, R. E., Rusanen, A., Sarnela, N., Simonen, P., Wang, S., Wang, W., Wang,

Y. Xue, M., Yang, G., Yao, L., Zhou, Y., Kujansuu, J., Petäjä, T., Nie, W., Ma, N., Ge, M., He, H., Donahue, N. M., Worsnop, D. R., Kerminen, V.-M., Wang, L., Liu, Y., Zheng, J., Kulmala, M., Jiang, J., and Bianchi, F.: The synergistic role of sulfuric acid, bases, and oxidized organics governing new-particle formation in Beijing, *Geophys. Res. Lett.*, 48, e2020GL091944, 2021.

Yan, C., Shen, Y., Stolzenburg, D., Dada, L., Qi, X., Hakala, S., Sundström, A. M., Guo, Y., Lipponen, A., Kokkonen, T. V., Kontkanen, J., Cai, R., Cai, J., Chan, T., Chen, L., Chu, B., Deng, C., Du, W., Fan, X., He, X. C., Kangasluoma, J., Kujansuu, J., Kurppa, M., Li, C., Li, Y., Lin, Z., Liu, Y., Liu, Y., Lu, Y., Nie, W., Pulliainen, J., Qiao, X., Wang, Y., Wen, Y., Wu, Y., Yang, G., Yao, L., Yin, R., Zhang, G., Zhang, S., Zheng, F., Zhou, Y., Arola, A., Tamminen, J., Paasonen, P., Sun, Y., Wang, L., Donahue, N. M., Liu, Y., Bianchi, F., Daellenbach, K. R., Worsnop, D. R., Kerminen, V. M., Petäjä, T., Ding, A., Jiang, J., and Kulmala, M.: The effect of COVID-19 restrictions on atmospheric new particle formation in Beijing, *Atmos. Chem. Phys.*, 22, 12207-12220, 10.5194/acp-22-12207-2022, 2022.

Yu, G.-R., Wen, X.-F., Sun, X.-M., Tanner, B. D., Lee, X., and Chen, J.-Y.: Overview of ChinaFLUX and evaluation of its eddy covariance measurement, *Agric. Forest Met.*, 137, 125-137, <https://doi.org/10.1016/j.agrformet.2006.02.011>, 2006.

Yu, F., Luo, G., Pryor, S. c., Pillai, P. R., Lee, S. H., Ortega, J., Schwab, J. J., Hallar, A. G., Leaitch, W. R., Aneja, V. P., Smith, J. N., Walker, J. T., Hogrefe, O., and Demirjian, K. L.: Spring and summer contrast in new particle formation over nine fest areas in North America, *Atmos. Chem. Phys.*, 15, 13993-14003, 2015.

Yli-Juuti, T., Mielonen, T., Heikkinen, L., Arola, A., Ehn, M., Isokäntä, S., Keskinen, H. M., Kulmala, M., Laakso, A., Lipponen, A., Luoma, K., Mikkonen, S., Nieminen, T., Paasonen, P., Petäjä, T., Romakkaniemi, S., Tonttila, J., Kokkola, H., and Virtanen, A.: Significance of the organic aerosol driven climate feedback in the boreal area, *Nature Communications*, 12, ARTN 5637, 10.1038/s41467-021-25850-7, 2021.

Ylivinkka, I., Kaupinmäki, S., Virman, M., Peltola, M., Taipale, D., Petäjä, T., Kerminen, V. M., Kulmala, M., and Ezhova, E.: Clouds over Hytylä, Finland: an algorithm to classify clouds based on solar radiation and cloud base height measurements, *Atmos Meas Tech*, 13, 5595-5619, 10.5194/amt-13-5595-2020, 2020.

Zhao, S. P., Yu, Y., Yin, D. Y., and Qin, H.: Contrasting response of ultrafine particle number and PM2.5 mass concentrations to clean air actions in China, *Geophys. Res. Lett.*, 48, e2921GL093886, 2021.

Zhao, Y., Saleh, R., Saliba, G., Presto, A. A., Gordon, T. D., Drozd, G. T., Goldstein, A. H., Donahue, N. M., and Robinson, A. L.: Reducing Secondary Organic Aerosol Formation from Gasoline Vehicle Exhaust, *Proc. Natl. Acad. Sci.*, 114 (27), 6984, <https://doi.org/10.1073/pnas.1620911114>, 2017.

Zhu, Y., Shen, Y., Meng, H., Sun, Y., Yao, X., Gao, H., Xue, L., and Wang, W.: Investigation of particle number concentrations and new particle formation with largely reduced air pollutant emissions at a coastal semi-urban site in Northern China, *J.*

1370 *Geophys. Res.: Atmos.*, 126, e2021JD035419, 2021.