

Impact of seawater sulfate concentration on sulfur concentration and isotopic composition in calcite of two cultured benthic foraminifera

Caroline Thaler^{1,2}, Guillaume Paris³, Marc Dellinger², Delphine Dissard⁴, Sophie Berland⁵, Arul Marie²,
5 Amandine Labat², Annachiara Bartolini¹

¹CR2P UMR 7207 MNHN CNRS SU, F-75005 Paris France

²MCAM UMR 7245 MNHN CNRS, F-75005 Paris France

³Université de Lorraine-CNRS, CRPG UMR 7358, F-54000 Nancy France

⁴LOCEAN UMR 7159 IRD SU CNRS MNHN, F-75005 Paris France/ Nouméa New Caledonia

10 ⁵BOREA UMR 8067 MNHN CNRS SU, F-75005 Paris France

Correspondence to: Caroline Thaler (thaler.caroline@gmail.com)

Abstract. Marine sediments can be used to reconstruct the evolution of seawater $[\text{SO}_4^{2-}]$ and $\delta^{34}\text{S}$ over time, two key parameters that contribute to refine our understanding of the sulfur cycle and thus of Earth's redox state. $\delta^{34}\text{S}$ evolution can be measured from carbonates, barites and sulfate evaporites. $[\text{SO}_4^{2-}]$ variations can be reconstructed using fluid inclusions in halites, a method that
15 only allows a low-resolution record. Reconstruction of the past sulfur cycle could be improved if carbonates allowed to track both seawater $\delta^{34}\text{S}$ and $[\text{SO}_4^{2-}]$ variations in a sole, continuous sedimentary repository. However, most primary carbonates formed in the ocean are biogenic, and organisms tend to overprint the geochemical signatures of their carbonates through a combination of processes often collectively referred to as vital effects. Hence, calibrations are needed to allow seawater $\delta^{34}\text{S}$ and $[\text{SO}_4^{2-}]$ reconstructions based on biogenic carbonates. Because foraminifera are important marine **calcifiers**, we opted to focus on calcite
20 synthesized by individuals of **rosalinid benthic foraminifera** cultured in laboratory under controlled conditions, with varying seawater $[\text{SO}_4^{2-}]$ (ranging from 0 mM to 180 mM). Our experimental design allowed us to obtain foraminiferal asexual reproduction over several generations. We measured bulk carbonate associated sulfate (CAS) content and sulfur isotopic composition ($\delta^{34}\text{S}_{\text{CAS}}$) on samples of tens to hundreds of specimens **from a selection of culture media**, where $[\text{SO}_4^{2-}]$ varied from 5 to 60 mM. Increasing or decreasing $[\text{SO}_4^{2-}]$ with respect to modern-day seawater concentration (28 mM) impacted foraminiferal population size dynamics
25 and the total amount of bioprecipitated carbonate. Foraminiferal CAS concentration increased proportionally with $[\text{SO}_4^{2-}]$ concentration from 5 mM up to **28 mM, and then showed a plateau from 28 to 60 mM**. The existence of a threshold **at 28 mM** is interpreted as the result of a control on the precipitation fluid chemistry that foraminifera exert on the carbonate precipitation loci. **However, at high seawater sulfate concentrations (> 40 mM) the formation of sulfate complexes with other cations, may partially contribute to the non-linearity of the CAS concentration in foraminiferal tests at high increases in $[\text{SO}_4^{2-}]$.** Yet, despite the
30 significant effect of **seawater $[\text{SO}_4^{2-}]$ on foraminiferal reproduction** and on CAS incorporation, the isotopic fractionation between CAS and seawater remains stable through varying seawater $[\text{SO}_4^{2-}]$. Altogether, these results illustrate that CAS in biogenic calcite could constitute a good proxy for both seawater $[\text{SO}_4^{2-}]$ and $\delta^{34}\text{S}$ and **suggests that sulfate likely plays a role in** foraminiferal biomineralization and biological activity.

1 Introduction

35

In the modern ocean, marine organisms control the precipitation of most calcium carbonates through the biomineralization of calcite or aragonite, the two main CaCO_3 polymorphs. Biogenic calcium carbonates from the sedimentary record have been used for decades to reconstruct past environmental conditions. At modern sulfate and magnesium concentrations in seawater (about 28

mM and 50 mM respectively), aragonite precipitates preferentially over calcite in abiotic conditions [at room temperature](#) (Bots et al., 2011; Barkan et al., 2020; Goetschl et al., 2019). Seawater sulfate and magnesium concentrations varied over the last 550 Myr, ranging from ~5 mM to ~28 mM (Horita et al., 2002), and from ~44 to ~55 mM, respectively (Lowenstein et al., 2001; Brennan et al., 2004). Lower and higher seawater sulfate and magnesium concentrations have been shown to match calcitic and aragonitic oceans, where calcite or aragonite forming organisms were favored respectively (Lowenstein et al., 2003; Algeo et al., 2015; Lin et al., 2018, Goetschl et al., 2019). In the modern aragonitic ocean (Sandberg et al., 1983) as well as through parts of the geological past of Earth's history, the occurrence of calcitic organisms (e.g. foraminifera, coccolithophorids, some mollusks, bryozoans and coralline algae) could thus appear as a paradox. These calcitic organisms growing in aragonite oceans with high sulfate content [would](#) then have developed adaptive strategies and exerted a high degree of biological control in calcite bioprecipitation and sulfate incorporation, which need to be better understood.

Among the main calcite synthesizers, foraminifera are unicellular eukaryotes that build mainly calcitic (rare aragonitic species exist) shells named "tests", that accumulate on the ocean seafloor (Schiebel 2002; Langer 2008). As foraminifera build their tests, trace elements present in seawater get incorporated in the biomineral structure. Sulfur is assumed to be incorporated in the calcium carbonate lattice structure as SO_4^{2-} by replacing a CO_3^{2-} group (Kontrec et al. 2004; Fernandez-Diaz et al. 2010) and is referred to as CAS, for Carbonate Associated Sulfate. This has been illustrated by an increase in S/Ca in benthic foraminiferal calcite [as a function of seawater \$\[\text{CO}_3^{2-}\]\$ decrease](#) (van Dijk et al., 2017). Paris et al. (2014) evidenced that the planktic species *Orbulina universa* faithfully records the $[\text{SO}_4^{2-}]/[\text{Ca}^{2+}]$ ratio of the seawater in which it grew for $[\text{SO}_4^{2-}]$ values from 18 mM to 28 mM. These encouraging results, however, needed to be tested on benthic species and on a wider range of $[\text{SO}_4^{2-}]$, to cover deep time oceanic values, which varied from less than 5 mM [up](#) to 28 mM nowadays (Algeo et al. 2015) and potentially beyond, during large volcanic events in the past, or in the vicinity of sulfate-rich volcanic hydrothermal fluids on the seabed (Gamo et al. 1997; Laakso et al. 2020). Furthermore, the possibility that foraminiferal calcite could serve both as $[\text{SO}_4^{2-}]$ and $\delta^{34}\text{S}$ record needs to be further validated. While so far measurements in biogenic carbonates showed that sulfur isotopes are systematically fractionated by $\pm 1\%$ from seawater (Kampschulte et al., 2001; Paris et al., 2014; Present et al., 2015; Rennie et al., 2018), recent experiments of abiotic CaCO_3 precipitation showed that a 2-5% fractionation of sulfur isotopes between aqueous sulfate and CAS in calcite covary with $[\text{SO}_4^{2-}]$ and, to a lesser extent, with precipitation rates (Barkan et al., 2020). There is thus a contrasting abiotic-biotic behavior that needs to be elucidated in order to determine whether calcitic foraminiferal tests could be used as a paleoenvironmental archive for the sulfur cycle, and [interrogates](#) the possibility that seawater $[\text{SO}_4^{2-}]$ variations impact foraminiferal biocalcification and carbonate production.

To answer these questions, we grew two strains of Rosalinidae (Fig. 1), which are asymbiotic benthic foraminifera, at constant temperature, pH and salinity over a range [from 0 to 180 mM](#) of seawater $[\text{SO}_4^{2-}]$. Compared to planktic foraminifera, benthic foraminifera have two advantages: (i) they cover deeper geological times and (ii) they can reproduce more easily in experimental conditions.

In general, in both planktic and benthic foraminiferal culture experiments performed to calibrate geochemical proxies, populations of individuals captured in the wild do not have the time to adapt to the experimental conditions because maintaining foraminiferal reproductions over several generations is a complicated task. Therefore, measurements of geochemical proxies are usually performed either on the few test chambers that precipitated in the experimental medium (e.g. Dissard et al., 2010 a and b; van Dijk et al., 2017; Schmidt et al., 2022), or on whole tests that include the initial chambers grown in the natural marine environment prior to collection (e.g. Paris et al., 2014; Le Houedec et al., 2021). Our experiment was carefully designed to obtain several generations grown over several weeks in each experimental medium, ensuring both acclimatization and full precipitation of the test in the medium. [Only live individuals that had fully grown under the experimental conditions were collected for analysis, as empty shells](#)

of dead individuals were discarded at each previous water change. We analyzed population size dynamics, as well as shell $\delta^{34}\text{S}$ and $[\text{SO}_4^{2-}]$ in bulk samples of tens to hundreds of specimens in each medium to shed light on the mechanisms of sulfate incorporation in benthic foraminiferal calcite.

2 Materials and methods

2.1 Culture conditions

2.1.1 Long term culture with asexual reproduction

Culture experiments were conducted at the French National Museum of Natural History (Muséum national d'Histoire naturelle, MNHN) in the free living protist collection facilities (collection group: Biological Resources of Living and Cryopreserved Cells; Collection of Unicellular Eukaryotes) on two previously cloned foraminiferal strains adapted to *in vitro* cell culture in 90 mm diameter Petri dishes with natural sea water (NSW) and fed with *Chlorogonium sp.* (specimen MNHN-CEU-2016-0001), a freshwater microalga. Two strains namely, For1C1 (specimen MNHN-CEU-2016-0075) and C1Tg (specimen MNHN-CEU-2016-0075) (Fig. 1 and Fig. 2), were isolated from the top layer of sediments collected from Banyuls sea shore (Mediterranean French coast) in 2006, and Concarneau (Atlantic French coast) in 2011, respectively.

Both strains were maintained through asexual reproduction (Fig. 3), using the following method: foraminifera were cultured in 90 mm diameter Petri dishes filled with 0.22 μm filtered NSW from Banyuls, France for For1C1 strain, or Concarneau, France for C1Tg strains (Fig. 1).

The NSW was kept in a cold room for at least a month and its pH (NBS scale) was adjusted to 8.2 through addition of NaOH and/or HCl, before use. The Petri dishes were kept at 22°C in an incubator (Memmert IPP 110 plus) equipped with cold white light modules (5,500 K) with a 12h day-12h night cycle. Water in the Petri dishes was changed once a week and foraminifera fed with living freshwater algae (*Chlorogonium sp.*). The algae were cultured in Basal Bold medium at 25°C under medium light intensity, and suspended in sterile pH 8.2 NSW after 3 steps of rinsing with NSW. Live algae can have a major impact on the seawater carbonate chemistry system by reproducing and consuming CO_2 through photosynthesis. As freshwater algae, the *Chlorogonium* cells died immediately in seawater, without undergoing lysis. This prevents those not eaten by foraminifera from spreading and/or being metabolically active and thus they do not influence the seawater chemistry conditions within the Petri dishes. The use of live freshwater instead of seawater algae to feed foraminifera is therefore an innovative approach that is particularly suited to long term culture experiments for the calibration of foraminiferal geochemical proxies, where seawater chemical conditions must be kept under control. Every other week a new Petri dish was set up with a dozen of new juveniles (pre-adults below the age of asexual reproduction, characterized by test with ~ 10 chambers). Live cultures were discarded after a month to prevent bacterial or fungal spread.

2.1.2 Culture in artificial seawater with varying $[\text{SO}_4^{2-}]$

In 2016, the two foraminiferal strains (For1C1 and C1Tg) were transferred to 0.22 μm filtered artificial seawater (ASW) mimicking NSW (Fig. 1). The ASW was prepared following Kester et al. 1967. The total salinity was 35.06 g/L and the main ionic concentrations were as follow, in mM: Cl^- 543.9, Na^+ 467.3, SO_4^{2-} 28.2, Mg^{2+} 53.1, Ca^{2+} 9.9, K^+ 10.0, HCO_3^- 2.3, Br^- 0.8, H_3BO_3 0.4, Sr^{2+} 0.1, F^- 0.1. After equilibration with the atmosphere for 2 to 3 hours, the pH of ASW was adjusted to pH 8.2 by the addition

of NaOH and HCl. ASW and NSW were sterilized by filtration on a 0.22 μm filter. The acclimation to ASW lasted approximately
120 for a year (with foraminifera being transferred to new 90 mm diameter Petri dishes monthly) without any noticeable effect on the
foraminiferal life cycle and morphology. Over this period of time, batches of several hundreds of foraminifera of each species
(strains For1C1 and C1Tg), cultured either in ASW or NSW, were sampled for $[\text{SO}_4^{2-}]$ and $\delta^{34}\text{S}$ composition measurements. The
C1Tg strain was only used for $[\text{SO}_4^{2-}]$ and $\delta^{34}\text{S}$ composition measurements of specimens from media in ASW and NSW at the
current seawater average $[\text{SO}_4^{2-}]$ of 28 mM, whereas the For1C1 strain was also used for $[\text{SO}_4^{2-}]$ and $\delta^{34}\text{S}$ composition
125 measurements of specimens from media with different $[\text{SO}_4^{2-}]$.

To produce media with different $[\text{SO}_4^{2-}]$, we created an ASW without SO_4^{2-} (hereafter ASW[0]) and another with $[\text{SO}_4^{2-}] = 180$
mM (hereafter ASW[180]). The amount of NaCl in those two media was adjusted to keep the total salinity constant (35.06 g/L).
Na⁺ concentrations for ASW[0] and ASW[180] were 479 mM and 402 mM, respectively and while the Cl⁻ concentrations were
612 mM and 175 mM, respectively. ASW[0] and ASW[180] were mixed in various proportions in order to obtain 8 other ASW
130 with the following $[\text{SO}_4^{2-}]$: 1, 5, 10, 35, 40, 50, 60, 90 and 120 mM. Each of these media had the same salinity as ASW (35.06
g/L), pH (8.2), DIC (Dissolved Inorganic Carbon: $[\text{CO}_2] + [\text{H}_2\text{CO}_3] + [\text{HCO}_3^-] + [\text{CO}_3^{2-}]$) and ALK (Alkalinity). For1C1 was the sole
strain grown under different SO_4^{2-} concentrations (Fig. 1).

In a first set of experiments (Set 1), 17 to 31 For1C1 individuals (Fig. 1, Table 1) with ~ 10 chambers each were transferred from
ASW to new 60 mm diameter Petri dishes filled with the following media: ASW (hereafter ASW[28]), ASW[0], ASW[5],
135 ASW[10], ASW[60], ASW[120] and ASW[180] and then cultured for 34 days. In parallel, 17 individuals so far cultured in natural
seawater were moved to a new 60 mm diameter Petri dish containing NSW from Banyuls and were cultured for 39 days.

In a second set of experiment designed to refine the concentration step between 0 and 90 mM (Set 2), 6 individuals of For1C1,
also presenting ~10 chambers each, were transferred from ASW to new 60 mm diameter Petri dishes and were cultured for 33 days
in the following media ASW[28], ASW[1], ASW[10], ASW[35], ASW[40], ASW[50], and ASW[90] (Fig. 1, Table 1).

140 For populations of more than approximately 300 individuals, as obtained in media with concentration ranging from 5 to 35mM
sulfate, the specimens were distributed over several 60 mm diameter Petri dishes (up to 3) to avoid problems associated with
overpopulation. In both sets of experiment, *Chlorogonium* fed to foraminifera were rinsed and suspended in the media
corresponding to each Petri dishes prior to their addition. We counted live individuals each week, for each medium. Since the
studied species live attached to a substrate, individuals that no longer stick to the Petri dishes were considered dead, even though
145 rare dead individuals (empty tests or no reticulopodial activity) may remain attached and few living adults can detach themselves
from the substrate as well. After counting and discarding the dead individuals, we sampled 6 mL of water through a 0.2 μm filtered
for DIC and $[\text{SO}_4^{2-}]$ measurements in gastight Exetainer© tubes full to the brim and stored at 5 °C. Consecutively, pH was measured
using a Hach PHC281101 probe calibrated following the three points procedure (Hach singlet solutions calibrated against NIST
standards, precision of ± 0.01 pH unit). Finally, the old water was completely replaced by fresh sterile water.

150

2.1.3 DIC analyses

DIC analyses were performed using 3 mL samples of seawater that were slowly withdrawn from each assay through the Exetainer©
rubber septa using needles syringes. Ultra-pure helium gas was injected in each vial during sampling to ease solution withdrawal
155 and to prevent atmospheric CO₂ contamination. Each 3 mL sample was injected into a new Exetainer© vial, previously flushed
with ultra-pure helium gas (2.5 bar) and loaded with 0.3 mL of 100% H₃PO₄. Acidification with pure H₃PO₄ converts the total DIC
of the sample into gaseous CO₂ which was allowed to degas and mix with the helium gas overnight under shaking. The CO₂ and
the He mix was then sampled with an autosampler and sent to a Dual Inlet FinniganTM DeltaPlus XP isotope ratio mass

spectrometer (Thermo Fisher Scientific) (reproducibility = $\pm 0.05\%$) at IPGP, Paris. [DIC] was quantified using the linear relationship between DIC concentration and intensity of the m/z 44 ($^{12}\text{C}^{16}\text{O}^{16}\text{O}$) peak provided by the mass spectrometer (Assayag et al., 2006). This linear relationship was established based on repeated analyses of internal laboratory carbonate standards calibrated against international standards (100% calcite), run in different aliquots. The reproducibility for [DIC] measurements was $\pm 5\%$ of the measured values (1σ).

2.2 Collection and rinsing procedure of the tests for geochemical analyses

At the end of the culture experiments (that varied between 34 and 39 days), all live individuals of the strain For1C1, as specified above, those still attached to the substrate from each Petri dish were recovered for geochemical analyses. Individuals from set 1 and set 2, grown under the same conditions (same medium [SO_4^{2-}]), were not combined for analysis. They were measured separately. Each sample typically weighed few milli-grams. The collected tests were rinsed 3 times in MQ water (basified to pH 9.5 with NH_4OH) to remove all traces of salts without dissolving the carbonate phase. In order to remove fresh organic matter, foraminifera were cleaned following Paris et al. (2014): foraminifera were bathed in a NaOH (0.5M) + H_2O_2 (15%) solution at 60°C for 30 minutes. They were then rinsed three times in basified MQ water, and dried overnight at 50°C in a drying oven. All samples were then dissolved at CRPG (Nancy) in 0.5 ml of 1 to 2% HCl . In addition, in order to determine how the remaining traces of organic matter could affect $\delta^{34}\text{S}$ measurements, some individuals from both For1C1 and C1Tg strains were dissolved in Aqua Regia (a 50/50 mix of concentrated HNO_3 and HCl), without prior cleaning in NaOH and H_2O_2 . They were left overnight at 120°C and dried down. All acids were distilled at CRPG and the 18.2 M Ω water purified through a Helga device (Veolia).

2.3 Geochemical analyses

180

2.3.1 CAS concentration analysis

In order to determine the $\text{SO}_4^{2-}/\text{Ca}^{2+}$ ratio of the tests, two dissolved foraminiferal calcite aliquots of 50 μl were used to independently measure the sulfate and calcium concentrations of the samples. To measure [SO_4^{2-}], one of the 50 μl aliquot was diluted in 200 μl of 18.2 M Ω water and ran on a Metrohm ion chromatography system (ICS). The calcium content of the samples was measured using a X-series II ICP-MS using the second aliquot that was dried down and taken up in 3 ml of 2% HNO_3 . For the latter, data were measured in groups of 5 bracketed by a 5.3 ppm Ca standard solution and bracketed assuming linear drift between two standards. In both cases, a calibration line was established to convert the signal to concentrations using home-made concentration standards. The typical reproducibility for sulfate and calcium concentrations is better than 2% based on multiple measurements of a diluted seawater solution for sulfate and of the standard solution for calcium.

190

2.3.2 $\delta^{34}\text{S}$ analysis

Sulfate isolation from the carbonate matrix was performed by ionic chromatography using the anionic resin Biorad AG1X8 (Paris et al., 2014) using precleaned disposable Biorad columns. Each column was prepared by loading 0.6 ml of resin and rinsed with 10% V/V HNO_3 (2x10CV – 1 CV = 1 column volume = 0.6 ml), 33% V/V HCl (2x10CV), 0.5N HCl (1x10CV). After introducing the dissolved carbonates sample on the resin, the column was rinsed with ultrapure water (5x5CV) to remove cations. SO_4^{2-} was then eluted with 0.45 M HNO_3 (3x2CV). Each batch of columns included a sample of 50 μl of seawater as a reference and total

195

200 procedure blanks. After elution, samples were dried down on hotplates with open lids (105°C). The total procedure blank was measured at $0.12 \text{ nmol S} \pm 47 \% \text{ RSD}$ with a $\delta^{34}\text{S}$ value of $6.1 \pm 3.5 \text{ ‰ } 1\sigma$, while the analysed samples contained between at least 23 nmol S (For1C1 grown in ASW[60]). Overall, a blank correction modifies values only within error bars and is thus not applied here.

205 Purified samples were analysed on the ThermoScientific Neptune Plus MC-ICP-MS at the CRPG using a standard-sample bracketing method (Paris et al, 2013). Samples were run at high resolution using an Aridus-II desolvating membrane to decrease oxide and hydride interferences. Isotopic ratios were collected at m/z 32 and 34 as 1 block of 50 cycles of 4.2 seconds each. Data were corrected offline for instrumental fractionation, drift and background following Paris et al. 2013. Each sample was measured twice on the Neptune and the value provided is an average of both measured ratios. The bracketing Na_2SO_4 solution had been previously calibrated against international standard IAEA S1 and checked against IAEA S2 and S3. Seawater samples ran during each Neptune sessions ensure that the data are not biased. Seawater external replicates were measured in association with those samples. They yield an average $\delta^{34}\text{S}$ value of $21.1 \pm 0.2 \text{ ‰}$ (n=4), in full agreement with published values (e.g. Paris et al., 2013; Present et al., 2015; Rennie et al., 2018). Because carbonate samples were too small to measure full external replicates, we assume the reproducibility for all $\delta^{34}\text{S}$ measurements to be the same as seawater ($\pm 0.2\text{‰}$; 2σ), with the exception of For1C1 grown in ASW [40] and [60]. In these two cases, the reproducibility is calculated based on the weighted mean of the internal errors multiplied by the standard deviation of the External Normalized Deviates (Paris et al., 2013), yielding a 2σ smaller than 0.2 ‰ except for these two samples (0.25 and 0.35 ‰ respectively).

3 Results

3.1 Population growth in each medium

220 Individuals For1C1 are morphologically similar to *Rosalina* (Fig. 2). They reproduced asexually when their tests reached a development of 11-12 chambers (Fig. 3 and Fig. 4). Under standard culture conditions of low cell density (i.e. where cells do not compete for food), which in our case was less than about 300 individuals per Petri dish, the reproductive cycle lasted ~ 12 to 15 days and individuals “died” after asexual reproduction by dividing themselves, usually into 20 to 40 viable juveniles, leaving an empty test (Fig. 3). Adult specimens were smaller than the traditional foraminiferal fraction obtained from sieving (through $>125 \mu\text{m}$ mesh) in geochemical studies, they thus may well be common and rarely collected because of their size. A morphological and taxonomic description of the cultured strains is available in the Appendix A. The weekly number of accumulated live individuals incremented by reproduction is given in Fig. 5 and Table 1.

230 The number of accumulated individuals can trace the population size dynamics for each medium and depends on reproduction rate, number of juveniles produced by individual and mortality. However, while the increase in the number of individuals clearly shows that living cells are being produced, no certainty about their viability can be drawn when the number of individuals stagnates or decreases, as no vital staining has been performed. It was therefore not always easy to distinguish between inactive and dead cells. We inferred mortality of foraminifera still adhering to the petri dishes from the cessation of reticulopodial activity and cytoplasmic streaming, as well as from the change in cell color (Bernhard, 2000). In the media with no sulfate or sulfate concentrations above 235 60 mM, we observed little to no reproduction, cell inactivity and probably mortality. As a result, the number of attached foraminifera remained constant and/or decreased over time (Fig. 5, Table 1). The most dramatic reactions were observed within a few hours in the media with highest $[\text{SO}_4^{2-}]$ (ASW[120] and ASW[180]), where individuals did not reproduce nor even show any reticulopodial activity. In ASW[90] and ASW[1], only one reproduction cycle was observed and after few days all the cells were

inactive (Fig. 5, Table 1). Overall, the highest numbers of individuals at the end of the experiment were obtained in the ASW[28],
240 NSW (Banyuls), ASW[5], ASW[10] and ASW[35] media (Fig. 5). Two media configurations, ASW[10] and ASW[28], from set
1 experiments were replicated in set 2 experiments. If the abundances for condition ASW[10] are of the same order of magnitude
in both sets, the abundances for condition ASW[28] are much lower in set 2 compared to set 1. This was related to the reproduction
rate in set 2, which slowed down drastically after 15 days. This decrease can be explained by a microbial bloom in the media that
was observed in no other media (Appendix, Fig. C1). The microbial spread could not be reduced by the weekly water change, and
245 any transfer and rinsing of foraminifera or antibiotic treatment would have constituted an additional experimental modification.
We thus kept counting foraminifera and sampling seawater, but did not take into account any results collected in that media after
day 15.

3.2 pH and DIC evolution

250

pH variations remained within ± 0.3 pH units during each experiment (Table 2).

pH drifted from the starting point between 8.1 and 8.2 towards more acidic values (7.83 minimum) and was reset close to 8.2 at
each medium change for the first 15 days and then remained rather stable with values varying between 8.19 and 8.07. DIC ranged
from 3.2 ± 0.2 mM (2σ) to 4.2 ± 0.3 mM (2σ) (Table 3). These concentrations are higher than the theoretical initial concentration of
255 2.8 mM using the recipe of Kester et al. 1967. While in Kester et al.'s recipe, the targeted 8.2 pH is achieved after 2h equilibration
with the CO₂ in the atmosphere, we had to proceed to NaOH addition despite a similar equilibration time. It is possible that higher
CO₂ dissolution at the atmospheric pressure of the year we performed the experiments (407 ppm in the atmosphere and probably
more in the lab against 322 ppm in 1967), led to an increase in DIC. In addition, DIC probably built up in the Petri dishes each
week as the foraminifera consumed the algae. pH and DIC variations for cultures in ASW[28] and ASW[10] are shown in Fig. 6.

260

3.3 CAS concentration

CAS concentration in foraminiferal calcite was performed for the media ASW[5], ASW[10], ASW[28], ASW[35], ASW[40] and
ASW[60], as the other samples were unfortunately lost during the manipulations or were below the detection limits. The obtained
265 values are presented in Fig. 7 and Table 4. Each datapoint was obtained using hundred to several hundreds of foraminifera for each
medium. CAS concentration (sulfate to calcite ratio) increased from 3320 ppm to ~ 14000 ppm $\text{SO}_4^{2-}/\text{CaCO}_3$ ($\pm 5\%$, 2σ) in
proportion to total SO_4^{2-} concentrations in artificial seawater which increased from 5 mM to 28 mM. While it seems that at sulfate
concentrations above the modern seawater concentration of 28 mM, the foraminiferal CAS concentration is no longer proportional
to that of the seawater and remains on a plateau between seawater $[\text{SO}_4^{2-}]$ 28 mM and 60 mM (Fig. 7). This suggests that a threshold
270 (~ 14000 ppm) is probably reached at about 28 mM $[\text{SO}_4^{2-}]$ in seawater. Because we only have a natural seawater replicate for the
sulfate concentration of modern seawater (28 mM), we observe a scatter at 28 mM that makes it difficult to determine precisely
when the plateau starts. The slight decrease in foraminiferal CAS to 9740 ppm at ASW[60] is actually part of the variability of
CAS values at 28 mM $[\text{SO}_4^{2-}]$ in seawater (Fig. 7 and Table 4). The foraminiferal CAS values from the ASW60 configuration can
therefore be considered as part of a plateau (Fig. 7).

275

3.4 Sulfur isotopic composition

The $\delta^{34}\text{S}$ values of the foraminiferal CAS from the different media are plotted in Fig. 8 and listed in Table 4. Measurements were performed both on foraminiferal samples from the C1Tg and For1C1 strains cultured in NSW or ASW[28] during the acclimation period and in those coming from the $[\text{SO}_4^{2-}]$ variation experiment (For1C1 strain only), from a selection of culture media in which $[\text{SO}_4^{2-}]$ varied from 5 to 60 mM (Table 4). NSW $\delta^{34}\text{S}$ composition was measured before ($21.1 \pm 0.2\text{‰}$) and 7 days after adding the algae ($19.9 \pm 0.2\text{‰}$). There was a difference beyond error bars between the two values. Considering that algae $\delta^{34}\text{S}$ composition are of $7.0 \pm 0.2\text{‰}$ the difference may be explained by the isotopically depleted sulfate added resulting of algae decomposition, lowering the average $\delta^{34}\text{S}$ of the media. This effect was not detectable in ASW possibly because the $\delta^{34}\text{S}$ values of medium ($9.1 \pm 0.2\text{‰}$ for ASW[28] and $0.1 \pm 0.2\text{‰}$ for ASW[5], ASW[10], ASW[35], ASW[40] and ASW[60]) was closer to that of the algae (Appendix B Table B1). Considering that algae were added at each water change and degraded within 1 or 2 days, and that foraminifera entered into a chamber formation sequence after feeding (Fig. 4), we consider that the seawater $\delta^{34}\text{S}$ that prevailed during chamber formation is the value measured after several days of culture with algae, $19.9 \pm 0.2\text{‰}$ in NSW and $9.1 \pm 0.2\text{‰}$ or $0.1 \pm 0.2\text{‰}$ in ASW[28] (Appendix B Table B1). A $\delta^{34}\text{S}_{\text{CAS}} - \delta^{34}\text{S}_{\text{sw}}$ fractionation value of $1.6 \pm 0.3\text{‰}$ (as observed for For1C1 pool (1 sd, 8 samples in total coming from all $[\text{SO}_4^{2-}]$ concentrations) while it was $1.4 \pm 0.2\text{‰}$ for C1Tg specimens (1 sd, 5 samples in total coming from NSW or ASW[28]), which is indistinguishable within the error range (Fig. 8). Samples for which organic matter was preserved yielded $\delta^{34}\text{S}$ values of $1.1 \pm 0.2\text{‰}$ (For1C1 in ASW[28]) $0.4 \pm 0.2\text{‰}$ (For1C1 in NSW) $1.4 \pm 0.2\text{‰}$ (C1Tg in NSW) and $0.5 \pm 0.2\text{‰}$ (C1Tg in ASW[28]) lower than the value that was obtained for the For1C1 and C1Tg tests from which organic matter had been oxidatively removed (Table 4).

4 Discussion

4.1 $[\text{SO}_4^{2-}]$ changes in seawater can affect foraminiferal biology

Our results highlight that a change in seawater $[\text{SO}_4^{2-}]$ concentration can affect foraminiferal cellular activity, reproduction and population size.

Reticulopodial activity stopped few hours after the transfer of individuals of the For1C1 strain from 28 mM of sulfate (ASW[28]) to concentrations above 120 mM (ASW[120] and ASW[180]) or without sulfate (ASW[0]). Dissolved sulfate and food were the only sources of sulfur in this experiment, which is essential for life. Since ASW[0] prevented any reproduction and induced cellular inactivity, we infer that sulfur from food appears insufficient and that dissolved sulfate in seawater is necessary for cellular activity in foraminifera. At the other extreme, toxic impact of the highest $[\text{SO}_4^{2-}]$ (ASW[120] and ASW[180]) can explain the non-reproduction and the cellular inactivity of individuals after a few hours. For1C1 individuals survived and even reproduced once in the ASW[1] and ASW[90] media (Fig. 5). Thus, our results suggest that foraminifera can reproduce and show pseudopodial activity only within a certain range of $[\text{SO}_4^{2-}]$, from 1 to 90 mM, extreme values at which the cellular activity is already very low. Individuals appear to tolerate these extreme conditions for only the first week and then cease all reproductive activity. They appear to be well adaptable, beyond the modern oceanic $[\text{SO}_4^{2-}]$ (28.2 mM) to a range of seawater $[\text{SO}_4^{2-}]$ from 5 to 35 mM, as shown by the high number of accumulated live individuals incremented by reproduction at the end of set 1 and set 2 experiments (Fig. 5). As already mentioned, the low number of individuals at the end of the second set of experiments is due to a bacterial infection after 3 weeks. Population size decreased above $[\text{SO}_4^{2-}]$ of 35 Mm and below 5 Mm, suggesting a foraminiferal reproduction sensitivity to $[\text{SO}_4^{2-}]$ variations. In particular high seawater $[\text{SO}_4^{2-}]$ (> 35 Mm) inhibit the foraminiferal proliferation by an undetermined toxic effect on the cellular physiology.

The effect of changes in seawater $[\text{SO}_4^{2-}]$ on **foraminiferal reproduction** highlights a possible mechanism by which changes in the composition of seawater can affect the carbonate record. It has previously been hypothesized that seawater Mg/Ca ratio and $\text{SO}_4^{2-}/\text{Ca}$ ratio, control the switch between calcite and aragonite dominance in the sedimentary record, as a high SO_4^{2-} and Mg concentrations in seawater inhibit calcite precipitation and promote aragonite precipitation, as shown by inorganic precipitation experiments (Bots et al., 2011; Barkan et al., 2020). Here, we show that a change in seawater $[\text{SO}_4^{2-}]$ may also affect foraminiferal **reproduction**, population size and **hence** their calcite accumulation in the sediment. **However, this appears to be for seawater $[\text{SO}_4^{2-}]$ variations far below and above the range (~5-30 mM) thought to be involved in long-term secular variations in the Phanerozoic, suggesting an adaptation of foraminifera in this range of variations. Indeed, under conditions that mimic the Phanerozoic range of $[\text{SO}_4^{2-}]$ variations, reproduction and population growth appear to be rather optimal.**

4.2 Foraminifer CAS concentration versus seawater $[\text{SO}_4^{2-}]$ or $\text{S}/\text{CO}_3^{2-}$

Our cultured foraminifera contain high levels of sulfates, similar to high Mg-calcite foraminifera previously grown during culture experiments, and significantly higher than low Mg-calcite foraminifera (Paris et al., 2014; van Dijk et al., 2017; 2019) (Fig. 7, Appendix B, Table B2). Similar to previous results of foraminiferal culture experiments comparing CAS content with seawater sulfate concentration, we note an increase in foraminiferal CAS content with seawater sulfate concentration increase. More specifically, our results show that CAS concentration in foraminiferal calcite grown in experimental seawater linearly increases with seawater $[\text{SO}_4^{2-}]$ concentration from 5 to 28 mM (Fig. 7), similarly to what is observed in inorganic carbonates (Busenberg and Plummer, 1985; Fernandez-Diaz et al., 2010; Barkan et al., 2020) or previous foraminiferal investigation (Paris et al., 2014). At $[\text{SO}_4^{2-}]$ higher than 28 mM in seawater, the incorporation of sulfate in the foraminiferal calcite seems to reach a saturation point (Fig. 7). It is remarkable to note that foraminifera can reproduce and thus calcify at $[\text{SO}_4^{2-}]$ as high as 90 mM (Fig. 5), concentrations at which no inorganic calcite precipitation occurs (Bots et al., 2011; Barkan et al., 2020). **However as discussed before, their reproduction is limited to the first week, which strongly suggests that they could only tolerate brief exposure to such a high level of sulfates in their environment.**

A geochemical modeling of experiments in which CAS, pH and DIC was measured is available in Appendix D, and permitted us to extract CO_3^{2-} concentrations. Overall, we observe an increase and a plateau, whether we compare our CAS content to seawater sulfate concentration or $\text{S}/\text{CO}_3^{2-}$ ratios. When we replace total sulfate (the sum of free SO_4^{2-} and its major complexed forms (NaSO_4 , CaSO_4 and MgSO_4) by only free sulfate, the linearity of the 5mM to 28mM CAS accumulation trend is maintained. However, the plateau from 35 to 60 mM is less visible, potentially evidencing the role of complexes formation in the lower SO_4^{2-} incorporation in the tests. As shown in the appendix, the sole formation of complexes cannot explain the plateau observed in figure 7.

To understand this evolution of sulfate content, we must first describe where sulfur is located in the test. Two options are possible:

i) CAS: Sulfate is incorporated into both inorganic and biogenic CaCO_3 minerals as CAS within the growing mineral structure, the larger tetrahedral sulfate substituting to the smaller trigonal-planar carbonate ion (Busenberg and Plummer, 1985; Kontrec et al. 2004; Balan et al., 2014; Tamenori et al. 2014; Perrin et al. 2017; Tamenori and Yoshimura 2018).

ii) S_{org} : Sulfur present in the organic matrix used by biomineralizing organisms to initiate calcification and orient the growing crystals (e.g. Cuif et al., 2004; Richardson et al., 2019; de Noojier et al., 2014). The organic matrix contains over-sulfated glycosaminoglycans and proteins (Weiner and Erez, 1984; Langer 1992). **The benthic foraminifera Rosalinidae belong to the order Rotaliida and likely share the same mechanisms of biomineralisation and test construction.** In the case of rotaliid test, the calcareous

360 wall growth of each new chamber results from the bioprecipitation of two calcite layers, on either side of an organic matrix (Bé et al., 1979; de Nooijer et al., 2014; Nagai et al., 2018) referred to as the Primary Organic Sheet (POS, Erez, 2003). However, since we applied an oxidative cleaning to the foraminiferal tests to destroy the organic matter, we assume that most of the measured $[\text{SO}_4^{2-}]$ in the tests are linked to the CAS concentration, although a small contribution might be still associated with S_{org} within the biomineralized calcite (Burdett et al., 1989; Cuif et al., 2003; Paris et al. 2014). In the following discussion, we will thus assume that our measured sulfate content reflects structurally-bound CAS.

Several hypotheses can be formulated to explain this $\text{SO}_4^{2-}/\text{CaCO}_3$ incorporation pattern:

365 i) Foraminifera may be able to regulate $[\text{SO}_4^{2-}]$ at the site of calcification (SOC) during calcite precipitation through active transmembrane transport, removing excess sulfate and lowering it in the precipitating fluid, enabling calcite nucleation and precipitation, as sulfate in high concentration inhibits calcite precipitation and makes it more soluble (Busenberg and Plummer, 1985; Bots et al., 2011; Barkan et al., 2020). In fact, under our experimental conditions the amount of CAS incorporated in foraminiferal calcite correlates with seawater SO_4^{2-} concentration, from 5 up to a plateau that starts at 28 mM. The mere fact that calcite precipitates therefore suggests that sulfate is at least partially removed from the precipitating fluid, altering the local SO_4^{2-} concentration. The correlation suggests that this removal is partial and, to some extent, proportional to the concentration of SO_4^{2-} in seawater.

370 ii) An increase in the carbonate ion concentration may help maintaining a constant $\text{SO}_4^{2-}/\text{CO}_3^{2-}$. Previous investigations demonstrated that it is more appropriated to reason in terms of $\text{SO}_4^{2-}/\text{CO}_3^{2-}$ ratio of the calcifying fluid rather than $[\text{SO}_4^{2-}]$ as sulfate substitutes for CO_3^{2-} in the forming mineral (van Dijk et al., 2019; Barkan et al., 2020). Another way to maintain $\text{SO}_4^{2-}/\text{CO}_3^{2-}$ constant while $[\text{SO}_4^{2-}]$ increases would be to proportionally increase CO_3^{2-} . Like other calcifying organisms, benthic foraminifera modify the pH of the precipitating fluid to promote calcite formation (Erez, 2003; de Nooijer et al., 2008; Rollion-Bard and Erez, 2010; Toyofuku et al., 2017). Foraminifera most probably actively pump protons out of the SOC (Sabbatini et al., 2014; Toyofuku et al., 2017). Intensifying this process in case of elevated $[\text{SO}_4^{2-}]$ would induce an increase in carbonate ion concentration (and the saturation state) and could therefore help to maintain a constant $\text{SO}_4^{2-}/\text{CO}_3^{2-}$ when $[\text{SO}_4^{2-}]$ reaches values between 28 and 90 mM, allowing calcite bio-precipitation. This mechanism, like that of active sulfate transmembrane transport mentioned above, would cease to function at seawater $[\text{SO}_4^{2-}]$ levels equal to or greater than 120 mM, when the foraminiferal cells become inactive.

385 iii) A preferential sequestering of sulfate in some organic rich layers at the incipient phase of biocalcification might allow to decrease the $[\text{SO}_4^{2-}]$ in the remaining liquid and thereby prevent further sulfate incorporation into foraminiferal calcite above 28 mM $[\text{SO}_4^{2-}]$ in seawater. High resolution sulfur nano-mapping on transversal section of perforate foraminiferal tests (such as Rosalinidae or *Orbulina*) showed a banded heterogeneity in sulfur distribution across the multi-layer structure (Paris et al., 2014; van Dijk et al., 2019). XRF intra-test mapping revealed a preferential incorporation of metals and sulfur in the POS zone, the organic incipient stage of the build-up of the wall of a new chamber of test (Lemelle et al., 2020). In our case, organic matter has been oxidized, and most of the “stored” SO_4^{2-} was likely removed.

390 iv) A kinetic effect could also explain the non-linearity of the CAS concentration in foraminiferal tests with corresponding increases in $[\text{SO}_4^{2-}]$ above 28 Mm, as inorganic calcite precipitation experiments suggest a reduction in crystal growth rates at higher $[\text{SO}_4^{2-}]$ (Busenberg and Plummer, 1985). However, it is worth noting that a decrease in precipitation rate can also be associated to a lower CAS content in inorganic calcite (Barkan et al., 2020). As a result, one could imagine that the change in sulfate concentration reflects a change in precipitation rate induced by different sulfate concentration in seawater and/or in the biomineralizing fluid. However, as calcite is more soluble and precipitates less easily at high sulfate concentration, we would expect an effect opposite

395 to what we observe in the 5-40 mM part of our results. There could nonetheless be a contribution of the rate effect to the plateau we observe.

v) Finally, another possibility to explain why the CAS in foraminiferal tests does not increase linearly with corresponding increase in the [SO₄²⁻] beyond 28 Mm, could be that at such concentrations in solution, sulfate might complex more easily with other cations such as Ca²⁺, K⁺, Mg²⁺, Na⁺, Sr²⁺, etc. (Garrels and Thompson, 1962). Such complexes cannot be effectively incorporated into the mineral lattice structure. This might influence the amount of SO₄²⁻ substituted in carbonates, and thus the CAS in foraminiferal tests. A geochemical model, available in the Appendix D, taking into account our media configurations where we had CAS, pH and DIC data, shows that the model CAS concentration follows linearly the seawater SO₄²⁻/CO₃²⁻ and CaSO₄/CaHCO₃ concentrations, which in turn depend mainly on the [SO₄²⁻] in solution, with a dip between 40 and 60 mM, likely related to the formation of complexes. However the sole formation of complexes cannot explain the plateau observed beyond 28 mM.

The putative mechanisms for sulfate regulation could have been adopted by foraminifera as evolutionary strategies to maintain carbonate precipitation despite potential variation in [SO₄²⁻]. Indeed, at [SO₄²⁻] greater than 8 mM abiotic calcite nucleation and precipitation is inhibited, and aragonite precipitates from saturated solutions (Kitano and Hood, 1962; Kitano et al., 1975; Bots et al., 2011). This inhibition is also true in the lack in magnesium (Barkan et al., 2020) and thus sulfate alone can affect calcite precipitation. Mechanisms such as increasing calcium concentration, pH and/or saturation state (e.g. Zeebe and Sanyal, 2002; Nehrke et al., 2013; Evans et al., 2018), as well as the presence of organics, could help overcome such high concentration of sulfate. However, when it comes to magnesium, active removal is also an option (Bentov and Erez, 2006). Calcitic foraminifera, which first appeared during the Devonian (Vachard et al., 2010) in a range of low-sulfate seawaters of ~3-15 mM (Algeo et al., 2015), might have progressively adopted such strategies in order to precipitate calcite in high-sulfate (~28 mM) seawaters such as those in the present-day ocean, and retain the capacity to precipitate calcite at concentrations reaching 90mM as evidenced here. However, in addition to active biological control to remove sulfate from the calcification site, a reduction in sulfate uptake in the tests at high seawater sulfate concentrations (> 40 mM) is likely to be due to the formation of sulfate complexes with other cations, explaining also the non-linearity of the CAS concentration in foraminiferal tests at high increases in [SO₄²⁻].

420 4.3 Sulfur isotope fractionation

The isotopic composition of CAS remains constants through our experiments. Sulfur isotopic fractionation of CAS in benthic foraminifera (Rosalinidae) is not sensitive to the variation in [SO₄²⁻] in seawater (Fig. 8), thus confirming the earlier observation on planktic foraminifera by Paris et al., 2014. This result by itself is important and confirms that foraminiferal CAS constitutes a reliable proxy of seawater δ³⁴S. This result, together with the correlation between SO₄/CaCO₃ and seawater [SO₄²⁻] (Fig. 7), supports that CAS in foraminiferal tests is of inorganic origin.

More importantly, the fractionation observed here is clearly different from the inorganic fractionation measured in the inorganic calcite (Barkan et al., 2020) highlighting the involvement of some biological isotopic fractionation. Considering that the algae's organic sulfur source had a fixed sulfur composition (7 ‰) while the seawater δ³⁴S varied from one medium to the other (from -0.1 to 20.0‰), our isotopic measurements on S_{org}+CAS allow to infer the origin of S_{org} as well. Mass balance calculation permit to determine that the isotopic composition of S_{org} varies with seawater δ³⁴S value, pointing towards mainly an inorganic source for S_{org} (Fig. 8). This is consistent with our observation that no cellular activity of foraminifera was possible in medium with zero [SO₄²⁻], even in the presence of algae as food and possible source of S_{org}. The δ³⁴S value of the combined S pool (S_{org} + CAS) is

0.4 to 1.4 ‰ more negative than the $\delta^{34}\text{S}$ value of CAS alone, which points towards the involvement of some biological
435 fractionation or vital effect associated to the incorporation of sulfur.

4.4 Implication for Paleoenvironmental reconstructions

Sulfur isotopic composition in the sedimentary record, through sulfur species redox reactivity and multiple deposition form, records
440 several paleoenvironmental processes occurring in the atmosphere and the ocean (Farquhar et al., 2000; Farquhar and Wing, 2003;
Crockford et al., 2019). This makes sulfur one of the most studied elements for the surface processes. And yet, in order to
investigate the sulfur cycle, it is necessary to interrogate different sedimentary archives: carbonates, barites, evaporites as well as
pyrites (Paytan et al., 1998; Algeo et al., 2015; Halevy et al., 2012; Present et al., 2020) to reconstruct variations in both $\delta^{34}\text{S}$ and
445 $[\text{SO}_4^{2-}]$. The work to match the sedimentary record of both $\delta^{34}\text{S}$ and $[\text{SO}_4^{2-}]$ is laborious and requires calibrations. **Our results show
that benthic foraminifera (Rosalinidae) incorporate CAS in their test proportionally to the $[\text{SO}_4^{2-}]$ in seawater, at least in the 5-28
mmol/L range, confirming previous experiments on planktic foraminifers that foraminiferal CAS can serve as a proxy for variations
of both $\delta^{34}\text{S}_{\text{CAS}}$ and $[\text{SO}_4^{2-}]$ in seawater (Paris et al., 2014). However, they also highlight that above the seawater $[\text{SO}_4^{2-}]$ of 28 mM,
it might not be possible to confidently determine the seawater $[\text{SO}_4^{2-}]$ using foraminiferal CAS, as the previous linear correlation
no longer holds. This limitation means that foraminiferal CAS could be used to trace deep time secular changes in seawater $[\text{SO}_4^{2-}]$,
450 $[\text{SO}_4^{2-}]$, which varies from about 5 mM to 28 mM today (Algeo et al. 2015), but not to trace past seawater $[\text{SO}_4^{2-}]$ enrichments above 28
mM, such as those that could be caused by large volcanic eruptions or sulfate-rich volcanic hydrothermal fluids on the seafloor.
Future works are therefore important to confirm whether or not the seawater $[\text{SO}_4^{2-}]$ threshold of 28 mM for CAS incorporation
can be applied to other benthic and planktonic foraminifera, or whether it is restricted to Rosalinidae.**

The use of CAS concentration as a marine $[\text{SO}_4^{2-}]$ record is **still promising, despite the limitation discussed above**, but will require
455 calibration on various types of carbonates and species that may each have their own fractionation factor. The preservation of that
dual $\delta^{34}\text{S}/[\text{SO}_4^{2-}]$ in foraminiferal calcite has to be evaluated in the carbonate record, as diagenesis has the capacity to affect $[\text{SO}_4^{2-}]$
in carbonates (e.g. Gill et al., 2008; Marenco et al., 2008; Rennie and Turchyn, 2014).

Additionally, it has been previously supported that S/Ca can work as a proxy for CO_3^{2-} concentration (van Dijk et al., 2017). Our
results complement this finding under the condition that it is applied on timescales where seawater $[\text{SO}_4^{2-}]$ are constant.

460 The other major **implication** of our results for the interpretation of the geological record is that changes in seawater $[\text{SO}_4^{2-}]$, **could**
affect the production of carbonate by affecting the reproduction/survival of **at least some calcifying organisms, as the benthic
foraminifera studied in this work**. In theory, the increase in seawater $[\text{SO}_4^{2-}]$ is expected to have a purely “abiotic” effect on calcite
production as sulfate thermodynamically inhibits calcite formation and makes calcite more soluble. As a result, higher sulfate
content in the living medium would generate a decrease in calcification intensity for a given individual. In this experiment we
465 showed that $[\text{SO}_4^{2-}]$ higher in the medium than those of the modern ocean can also decrease the amount of accumulated calcite by
affecting foraminiferal population size, suggesting that their biological activity is harmed by such sulfate concentrations. As a
result, the decrease the total amount of calcification would be explained partly by a decrease in biological activity. This work
illustrated **how variations in seawater composition can have a dual effect on biomineralizing organisms. Conditions that inhibit
calcite formation such as increases in marine concentrations of Mg^{2+} or SO_4^{2-} , could have chemical “abiotic” effects on carbonates
470 formation but could also affect biological processes involved in biomineralization.**

5 Conclusion

475 We cultured rotaliid foraminifera in media with $[\text{SO}_4^{2-}]$ spanning from 0 mM to 180 mM, stable salinity and fixed seawater $\delta^{34}\text{S}$.
[SO_4^{2-}] changes in seawater affected foraminiferal **reproduction**, population size dynamics and calcite accumulation. Foraminifera
kept precipitating calcite in media reaching $[\text{SO}_4^{2-}] = 90$ mM. Sulfate from seawater is necessary for the cellular activity of
foraminifera, but at concentrations **equal and** above 90 mM it becomes toxic to them, **as evidenced by cellular inactivity and**
480 **reproductive arrest**. Sulfur concentration in CAS varied proportionally to seawater $[\text{SO}_4^{2-}]$ between 1 and 28.2 mM and then
stabilizes. Our results highlight that isotope fractionation between CAS and seawater does not depend on seawater $[\text{SO}_4^{2-}]$. Overall,
similarly to planktic foraminifera the $\delta^{34}\text{S}_{\text{CAS}}$ value of a given species of benthic foraminifera is a reliable way to reconstruct
seawater $\delta^{34}\text{S}$, despite variations of $[\text{SO}_4^{2-}]$ in seawater.

Author contribution: All authors participated in designing and interpreting the experiments. CT carried out culture experiments
485 under MD and AB supervision and technical support from AL. MD isolated the strains and developed the cultivation protocol.
GP performed all isotopic and $\text{SO}_4/\text{CaCO}_3$ analyses. CT designed the figures and wrote the paper with contributions from all co-
authors.

Competing interests:

490 The authors declare no competing interests

Acknowledgements

This research was funded by the LabEx BCDiv project « SULFOR, Impact of SO_2 on ocean acidification and foraminiferal
biocalcification » (PI AB); the CNRS INSU INTERRVIE project « Impact of sulfate variations on the biocalcification of
495 foraminifera » (PI AB). **Sylvain Pont (MNHN, IMPMC UMR 7590, France) produced the SEM imaging. The authors thank the**
reviewers Julien Richirt and David Evans for their very constructive criticism, which helped to improve the article.

References

- Algeo, T. J., Luo, G. M., Song, H. Y., Lyons, T. W., and Canfield, D. E.: Reconstruction of secular variation in seawater sulfate
500 concentrations, *Biogeosciences*, 12, 2131-2151, <https://doi.org/10.5194/bg-12-2131-2015>, 2015.
- Assayag, N., Rivé, K., Ader, M., Jézéquel, D., and Agrinier, P.: Improved method for isotopic and quantitative analysis of dissolved
inorganic carbon in natural water samples, *Rapid Communications in Mass Spectrometry*, 20, 2243-2251, 2006.
- Balan, E., Blanchard, M., Pinilla, C., and Lazzeri, M.: First-principles modeling of sulfate incorporation and $^{34}\text{S}/^{32}\text{S}$ isotopic
fractionation in different calcium carbonates, *Chemical Geology*, 374-375, 84-91, <https://doi.org/10.1016/j.chemgeo.2014.03.004>,
505 2014.
- Barkan, Y., Paris, G., Webb, S. M., Adkins, J. F., and Halevy, I.: Sulfur isotope fractionation between aqueous and carbonate-
associated sulfate in abiotic calcite and aragonite, *Geochimica et Cosmochimica Acta*, 280, 317-339,
<https://doi.org/10.1016/j.gca.2020.03.022>, 2020.
- Bé, A. W., Hemleben, C., Anderson, O. R., and Spindler, M.: Chamber formation in planktonic foraminifera. *Micropaleontology*,
510 25, 294-307, <https://doi.org/10.2307/1485304>, 1979.
- Bentov, S., Erez, J., 2006. Impact of biomineralization processes on the Mg content of foraminiferal shells: A biological
perspective. *Geochemistry, Geophysics, Geosystems* 7, Q01P08. <https://doi.org/10.1029/2005gc001015>**

- Bernhard, J. M.: Distinguishing live from dead foraminifera: Methods review and proper applications, *Micropaleontology*, 46, Suppl. 1, 38-46, 2000.
- 515 Bots, P., Benning, L., Rickaby, R., and Shaw, S.: The role of SO₄ in the switch from calcite to aragonite seas, *Geology*, 39, 331-334, <https://doi.org/10.1130/G31619.1>, 2011.
- Brennan, S. T., Lowenstein, T. K., and Horita, J.: Seawater chemistry and the advent of biocalcification, *Geology* 32, 473-476, <https://doi.org/10.1130/G20251.1>, 2004.
- Burdett, J. W., Arthur, M. A., and Richardson, M.: A Neogene seawater sulfur isotope age curve from calcareous pelagic
520 microfossils, *Earth and Planetary Science Letters*, 94, 189-198, [https://doi.org/10.1016/0012-821X\(89\)90138-6](https://doi.org/10.1016/0012-821X(89)90138-6), 1989.
- Busenberg, E. and Plummer, L. N.: Kinetic and thermodynamic factors controlling the distribution of SO₃²⁻ and Na⁺ in calcites and selected aragonites, *Geochimica et Cosmochimica Acta*, 49, 713-725, [https://doi.org/10.1016/0016-7037\(85\)90166-8](https://doi.org/10.1016/0016-7037(85)90166-8), 1985.
- Crockford, P. W., Kunzmann, M., Bekker, A., Hayles, J., Bao, H., Halverson, G. P., Peng, Y., Bui, T. H., Cox, G. M., Gibson, T. M., Wörndle, S., Rainbird, R., Lepland, A., Swanson-Hysell, N. L., Master, S., Sreenivas, B., Kuznetsov, A., Krupenik, V., and
525 Wing, B. A.: Claypool continued: Extending the isotopic record of sedimentary sulfate. *Chemical Geology*, 513, 200-225, <https://doi.org/10.1016/j.chemgeo.2019.02.030>, 2019.
- Cuif, J. P., Dauphin, Y., Doucet, J., Salome, M., and Susini, J.: XANES mapping of organic sulfate in three scleractinian coral skeletons, *Geochimica et Cosmochimica Acta*, 67, 75-83, [https://doi.org/10.1016/S0016-7037\(02\)01041-4](https://doi.org/10.1016/S0016-7037(02)01041-4), 2003.
- Cuif, J. P., Dauphin, Y., Berthet, P., and Jegoudez, J.: Associated water and organic compounds in coral skeletons: quantitative
530 thermogravimetry coupled to infrared absorption spectrometry, *Geochemistry, Geophysics, Geosystems*, 5, Q11011, <https://doi.org/10.1029/2004GC000783>, 2004.
- Dettmering, C., Rottger, R., Hoheneqger, J., and Schrnaljohann, R.: The Trimorphic Life Cycle in Foraminifera: Observations from Cultures allow New Evaluation, *Europ. J. Protistol.* 34, 363-368, 1998.
- Dissard, D., Nehrke, G., Reichart, G. J., Bijma, J.: The impact of salinity on the Mg/Ca and Sr/Ca ratio in the benthic foraminifera
535 *Ammonia tepida*: results from culture experiments, *Geochimica et Cosmochimica Acta*, 74, 3-74, doi:10.1016/j.gca.2009.10.040.
- Dissard, D., Nehrke, G., Reichart, G.-J., Bijma, J.: Impact of seawater pCO₂ on calcification and Mg/Ca and Sr/Ca ratios in benthic foraminifera calcite: results from culturing experiments with *Ammonia tepida*, *Biogeosciences*, 7, 81-93, doi:10.5194/bg-7-81-2010.
- Erez, J.: The source of ions for biomineralization in foraminifera and their implications for paleoceanographic proxies, in:
540 *Biomineralization*, edited by: Dove, P. M., De Yoreo J. J., and Weiner, S., De Gruyter, Berlin, Boston, *Reviews in Mineralogy & Geochemistry*, 54, 115-149, <https://doi.org/10.1515/9781501509346-010>, 2003.
- Evans, D., Müller, W., Erez, J., 2018. Assessing foraminifera biomineralisation models through trace element data of cultures under variable seawater chemistry. *Geochimica et Cosmochimica Acta* 236, 198-217. <https://doi.org/10.1016/j.gca.2018.02.048>
- Farquhar J, Bao, H. M. and Thiemens, M.: Atmospheric influence of Earth's earliest sulfur cycle, *Science* 289, 756-759, DOI:
545 10.1126/science.289.5480.756, 2000.
- Farquhar, J. and Wing, B. A.: Multiple sulfur isotopes and the evolution of the atmosphere, *Earth and Planetary Science Letters*, 213, 1-13, [https://doi.org/10.1016/S0012-821X\(03\)00296-6](https://doi.org/10.1016/S0012-821X(03)00296-6), 2003.
- Fernández-Díaz, L., Fernández-González, Á., and Prieto, M.: The role of sulfate groups in controlling CaCO₃ polymorphism, *Geochimica et Cosmochimica Acta*, 74, 6064-6076, <https://doi.org/10.1016/j.gca.2010.08.010>, 2010.
- 550 Gamo, T., Okamura, K., Charlou, J.-L., Urabe, T., Auzende, J.-M., Ishibashi, J., Shitashima, K., and Chiba, H. : Acidic and sulfate-rich hydrothermal fluids from the Manus back-arc basin, Papua New Guinea, *Geology*, 25, 139-142, 1997.

- Garrels, R. M. and Thompson, M. E.: A chemical model for sea water at 25 degrees C and one atmosphere total pressure, *American Journal of Science*, 260, 57-66, <https://doi.org/10.2475/ajs.260.1.57>, 1962.
- Gill, B. C., Lyons, T. W., and Frank, T. D.: Behavior of carbonate-associated sulfate during meteoric diagenesis and implications for the sulfur isotope paleoproxy, *Geochimica et Cosmochimica Acta*, 72, 4699-4711, <https://doi.org/10.1016/j.gca.2008.07.001>, 2008.
- Goetschl, K. E., Purgstaller, B., Dietzel, M., and Mavromatis, V.: Effect of sulfate on magnesium incorporation in low-magnesium calcite, *Geochimica et Cosmochimica Acta*, 265, 505-519, <https://doi.org/10.1016/j.gca.2019.07.024>, 2019.
- Halevy, I., Peters, S. E., and Fischer, W. W.: Sulfate burial constraints on the Phanerozoic sulfur cycle, *Science*, 337, 331-334, DOI: 10.1126/science.1220224, 2012.
- Horita, J., Zimmermann, H., and Holland, H.D.: Chemical evolution of seawater during the Phanerozoic: Implications from the record of marine evaporites, *Geochimica et Cosmochimica Acta* 66, 3733–3756, 2002.
- Holzmann, M. and Pawlowski, J.: An updated classification of rotaliid foraminifera based on ribosomal DNA phylogeny, *Marine Micropaleontology*, 132, 18–34, <http://dx.doi.org/10.1016/j.marmicro.2017.04.002>, 2017.
- Kampschulte, A., Bruckschen, P., and Strauss, H.: The sulphur isotopic composition of trace sulphates in Carboniferous brachiopods: implications for coeval seawater, correlation with other geochemical cycles and isotope stratigraphy, *Chemical Geology*, 175, 149-173, [https://doi.org/10.1016/S0009-2541\(00\)00367-3](https://doi.org/10.1016/S0009-2541(00)00367-3), 2001.
- Kester, D. R., Duedall, I. W., Connors, D. N., and Pytkowicz, R. M.: Preparation of artificial seawater 1, *Limnology and oceanography*, 12, 176-179, <https://doi.org/10.4319/lo.1967.12.1.0176>, 1967.
- Kitano, Y. and Hood, D. W.: Calcium carbonate crystal forms formed from sea water by inorganic processes, *Journal of the Oceanographical Society of Japan*, 18, 141-145, <https://doi.org/10.5928/kaiyou1942.18.141>, 1962.
- Kitano, Y., Okumura, M., and Idogaki, M.: Incorporation of sodium, chloride and sulfate with calcium carbonate, *Geochemical Journal*, 9, 75-84, <https://doi.org/10.2343/geochemj.9.75>, 1975.
- Kontrec, J., Kralj, D., Brečević, L., Falini, G., Fermani, S., Noethig-Laslo, V., and Miroslavljević, M.: Incorporation of Inorganic Anions in Calcite, *EurJIC*, 2004, 4579-4585, <https://doi.org/10.1002/ejic.200400268>, 2004.
- Langer, M. R.: Biosynthesis of glycosaminoglycans in foraminifera: A review, *Marine Micropaleontology*, 19, 245-255, 1992.
- Langer, M. R.: Assessing the contribution of foraminiferan protists to global ocean carbonate production, *The Journal of Eukaryotic Microbiology*, 55, 163-169, <https://doi.org/10.1111/j.1550-7408.2008.00321.x>, 2008.
- Laakso, T. A., Waldeck, A., Macdonald, F. A., and Johnston, D.: Volcanic controls on seawater sulfate over the past 120 million years, *PNAS*, 117, 21118–21124, <https://doi.org/10.1073/pnas.1921308117>, 2020.
- Le Calvez, J.: Recherches sur les foraminifères. 1. Développement et reproduction. *Archives Zoologie Expérimentale et Générale* 80, 163-333, 1938.
- Le Houedec, S., Erez, J., and Rosenthal, Y.: Testing the influence of changing seawater Ca concentration on Elements/Ca ratios in planktic foraminifera: A culture experiment, *Geochemistry, Geophysics, Geosystems*, 22, e2020GC009496, <https://doi.org/10.1029/2020GC009496>, 2021.
- Lemelle, L., Bartolini, A., Simionovici, A., Tucoulou, R., De Nolf, W., Bassinot, F., and de Garidel-Thoron, T. : Nanoscale trace metal imprinting of biocalcification of planktic foraminifers by Toba's super-eruption, *Scientific Reports*, 10, 10974, <https://doi.org/10.1038/s41598-020-67481-w>, 2020.
- Lin, C. Y., Turchyn, A. V., Steiner, Z., Bots, P., Lampronti, G. I., and Tosca, N. J.: The role of microbial sulfate reduction in calcium carbonate polymorph selection, *Geochimica et Cosmochimica Acta*, 237, 184-204, <https://doi.org/10.1016/j.gca.2018.06.019>, 2018.

- Lowenstein, T. K., Timofeeff, M. N., Brennan, S. T., Hardie, L. A., and Demicco, R.V.: Oscillations in Phanerozoic seawater chemistry: Evidence from fluid inclusions, *Science* 294, DOI: 10.1126/science.1064280, 2001.
- 595 Lowenstein, T. K., Hardie, L. A., Timofeeff, M. N., and Demicco, R. V.: Secular variation in seawater chemistry and the origin of calcium chloride basinal brines, *Geology*, 31, 857-860, <https://doi.org/10.1130/G19728R.1>, 2003.
- Marenco, P. J., Corsetti, F. A., Kaufman, A. J., and Bottjer, D. J.: Environmental and diagenetic variations in carbonate associated sulfate: an investigation of CAS in the Lower Triassic of the western USA, *Geochimica et Cosmochimica Acta*, 72, 1570-1582, <https://doi.org/10.1016/j.gca.2007.10.033>, 2008.
- 600 Nagai, Y., Uematsu, K., Chen, C., Wani, R., Tyszka, J., and Toyofuku, T.: Weaving of biomineralization framework in rotaliid foraminifera: implications for paleoceanographic proxies, *Biogeosciences*, 15, 6773-6789, <https://doi.org/10.5194/bg-15-6773-2018>, 2018.
- Nehrke, G., Keul, N., Langer, G., de Nooijer, L. J., Bijma, J., and Meibom, A. A new model for biomineralization and trace-element signatures of Foraminifera tests, *Biogeosciences*, 10, 6759–6767, <https://doi.org/10.5194/bg-10-6759-2013>, 2013
- 605 de Nooijer, L. J., Toyofuku, T., and Kitazato, H: Foraminifera promote calcification by elevating their intracellular pH, *Proc. Natl Acad. Sci.*, 106, 15374–15378, 2009.
- de Nooijer, L. J., Spero, H. J., Erez, J., Bijma, J., and Reichart, G. J: Biomineralization in perforate foraminifera, *Earth-Science Reviews*, 135, 48-58, <https://doi.org/10.1016/j.earscirev.2014.03.013>, 2014.
- Paris, G., Sessions, A. L., Subhas, A. V., and Adkins, J. F.: MC-ICP-MS measurement of $\delta^{34}\text{S}$ and $\Delta^{33}\text{S}$ in small amounts of dissolved sulfate, *Chemical Geology* 345, 50–61, <http://dx.doi.org/10.1016/j.chemgeo.2013.02.022>, 2013.
- 610 Paris, G., Fehrenbacher, J. S., Sessions, A. L., Spero, H. J., and Adkins, J. F.: Experimental determination of carbonate-associated sulfate $\delta^{34}\text{S}$ in planktonic foraminifera shells. *Geochem., Geophys., Geosyst.*, 15, 1452-1461, doi:10.1002/2014GC005295, 2014.
- Paytan, A., Kastner M., Campbell, D., and Thieme, M. H.: Sulfur Isotopic Composition of Cenozoic Seawater Sulfate, *Science* 282, 1459-1462, DOI: 10.1126/SCIENCE.282.5393.145, 1998.
- 615 Perrin, J., Rivard, C., Vielzeuf, D., Laporte, D., Fonquernie, C., Ricolleau, A., and Floquet, N.: The coordination of sulfur in synthetic and biogenic Mg calcites: The red coral case, *Geochimica et Cosmochimica Acta*, 197, 226-244, <https://doi.org/10.1016/j.gca.2016.10.017>, 2017.
- Present, T. M., Paris, G., Burke, A., and Fischer, W. W.: Large carbonate associated sulfate isotopic variability between brachiopods, *Earth and Planetary Science Letters*, 432, 187-198, <https://doi.org/10.1016/j.epsl.2015.10.005>, 2015.
- Present, T. M., Adkins, J. F., and Fischer, W. W.: Variability in sulfur isotope records of Phanerozoic seawater sulfate, *Geophysical Research Letters*, 47, e2020GL088766, <https://doi.org/10.1029/2020GL088766>, 2020.
- 620 Rennie, V. C. and Turchyn, A. V.: The preservation of $\delta\text{S}^{34}_{\text{SO}_4}$ and $\delta\text{O}^{18}_{\text{SO}_4}$ in carbonate-associated sulfate during marine diagenesis: A 25 Myr test case using marine sediments, *Earth and Planetary Science Letters*, 395, 13-23, <https://doi.org/10.1016/j.epsl.2014.03.025>, 2014.
- Rennie, C. F., Paris, G., Sessions, A. L., Abramovich, S., Turchyn, A. V., and Adkins, J. F.: Cenozoic record of $\delta^{34}\text{S}$ in foraminiferal calcite implies an early Eocene shift to deep-ocean sulfide burial, *Nature Geoscience*, 11, 761-765, <https://doi.org/10.1038/s41561-018-0200-y>, 2018.
- 625 Richardson, J. A., Newville, M., Lanzirrotti, A., Webb, S. M., Rose, C. V., Catalano, J. G., and Fike, D. A.: The source of sulfate in brachiopod calcite: Insights from μ -XRF imaging and XANES spectroscopy, *Chemical Geology*, 529, 119328, <https://doi.org/10.1016/j.chemgeo.2019.119328>, 2019.

- 630 Rollion-Bard, C. and Erez, J.: Intra-shell boron isotope ratios in the symbiont-bearing benthic foraminiferan *Amphistegina lobifera*: Implications for $\delta^{11}\text{B}$ vital effects and paleo-pH reconstructions, *Geochimica et Cosmochimica Acta*, 74, 1530-1536, <https://doi.org/10.1016/j.gca.2009.11.017>, 2010.
- Sabbatini, A., Bédouet, L., Marie, A., Bartolini, A., Landemarre, L., Weber, M. X., Kade Mahardika, I. G. N., Berland, S., Zito, F., Vénec-Peyré, M. T.: Biomineralization of *Schlumbergerella floresiana*, a significant carbonate-producing benthic foraminifer, *Geobiology* 12, 289-307, <https://doi.org/10.1111/gbi.12085>, 2014.
- 635 Sandberg, P. A.: An oscillating trend in Phanerozoic non-skeletal carbonate mineralogy, *Nature*, 305, 19-22, 1983.
- Schiebel, R.: Planktic foraminiferal sedimentation and the marine calcite budget, *Global Biogeochemical Cycles*, 16, 1065, <https://doi.org/10.1029/2001GB001459>, 2002.
- Schmidt, S., Hathorne, E. C., Schönfeld, J., and Garbe-Schönberg, D.: Heavy metal uptake of nearshore benthic foraminifera during multi-metal culturing experiments, *Biogeosciences*, 19, 629–664, <https://doi.org/10.5194/bg-19-629-2022>, 2022.
- 640 Tamenori, Y., Yoshimura, T., Luan, N. T., Hasegawa, H., Suzuki, A., Kawahata, H., and Iwasaki, N.: Identification of the chemical form of sulfur compounds in the Japanese pink coral (*Corallium elatius*) skeleton using $\mu\text{-XRF/XAS}$ speciation mapping, *Journal of Structural Biology*, 186, 214-223, <https://doi.org/10.1016/j.jsb.2014.04.001>, 2014.
- Tamenori, Y., and Yoshimura, T.: Sulfur speciation in growth layers of shell cross section of the long-lived bivalve *Margaritifera laevis* using synchrotron spectromicroscopy analysis, *Geochimica et Cosmochimica Acta*, 237, 357-369, <https://doi.org/10.1016/j.gca.2018.07.002>, 2018.
- 645 Toyofuku, T., Matsuo, M. Y., de Nooijer, L. J., Nagai, Y., Kawada, S., Fujita, K., Reichart, G.-J., Nomaki, H., Tsuchiya, M., Sakaguchi, H., and Kitazato, H.: Proton pumping accompanies calcification in foraminifera, *Nature Communications* 8, 14145, DOI: 10.1038/ncomms14145, 2017.
- 650 Vachard, D., Pille, L., and Gaillot, J.: Palaeozoic Foraminifera: Systematics, palaeoecology and responses to global changes, *Rev. de Micropaleontol.*, 53, 209–254, <https://doi.org/10.1016/j.revmic.2010.10.001>, 2010.
- Van Dijk, I., de Nooijer, L. J., and Reichart, G. J.: Trends in element incorporation in hyaline and porcelaneous foraminifera as a function of pCO_2 , *Biogeosciences*, 14, 497-510, doi:10.5194/bg-14-497-2017, 2017.
- Van Dijk, I., Barras, C., De Nooijer, L. J., Mouret, A., Geerken, E., Oron, S., and Reichart, G. J.: Coupled calcium and inorganic carbon uptake suggested by magnesium and sulfur incorporation in foraminiferal calcite, *Biogeosciences*, 16, 2115-2130, <https://doi.org/10.5194/bg-16-2115-2019>, 2019.
- 655 Weiner, S. and Erez, J.: Organic matrix of the shell of the foraminifer, *Heterostegina depressa*, *The Journal of Foraminiferal Research*, 14, 206-212, <https://doi.org/10.2113/gsjfr.14.3.206>, 1984.
- Zeebe, R.E., Sanyal, A., 2002. Comparison of two potential strategies of planktonic foraminifera for house building: Mg^{2+} or H^+ removal? *Geochimica et Cosmochimica Acta* 66, 1159–1169. [https://doi.org/10.1016/S0016-7037\(01\)00852-3](https://doi.org/10.1016/S0016-7037(01)00852-3)
- 660






Figure 1. Experimental design workflow diagram illustrating the sampling, acclimation, and experiments SET 1 and SET 2. Stars highlight samples where $\delta^{34}\text{S}$ and CAS analysis could be performed.




675

Figure 2. Foraminiferal strains cultured in this study. For1C1: (A) *Rosalina* like morphotype (11-12 chambers) reproducing asexually, dorsal view, (B) same as A, ventral view, (C) Morphotype with more than 12 chambers, starting as *Rosalina* morphotype and then developing annular disposition of the last chambers, dorsal view. C1Tg: (D) *Rosalina* like morphotype, (E) same as D, ventral view, (F) morphotype with annular arrangement of the last chambers, dorsal view. (G and H) ventral view of C1Tg with a broken test permitting to see the layered structure of the test's wall (G) and the foramen position inside of the test (H). Scale bar 50 μm , SEM picture in BSE mode operated at 10 to 22 mPa and 20 000kV.

680




685 **Figure 3.** Asexual reproduction of an individual of the For1C1 strain. (A) Light microscope image of a megalospheric schizont adult that has ~12 visible chambers, and whose cell has divided asexually into viable juveniles (for further detail, see Appendix A). The darker appearance of the juveniles compared to the adult is due to the presence of cellular material. After division, the adult is empty and its test partially dissolved, as shown in the SEM micrograph (B). Scale bar 100 μm .



690 **Figure 4.** Chamber formation during day 2 of the set 1 series of experiment. Two For1C1 individuals (from Banyuls) in (a) ASW[5] (containing 5 mM SO_4^{2-}) and (b) ASW[10] (containing 10 mM SO_4^{2-}) and its schematic representation illustrating the new chamber (n) formation and the surrounding gangue (algal cyst) constituted by the foraminifer by the accumulation of foreign detritus and other materials, confining the new chamber in formation in a microenvironment. In the case of rotaliid foraminifera, the formation of a new chamber begins with the isolation of the chamber volume from the surrounding environment by a structure which probably form the organic scaffolding that shapes the morphology of the chamber and serves as a template for the calcification of the wall (Bé et al., 1979; De Nooijer et al., 2014; Nagai et al., 2018). Precipitation of calcium carbonate takes place on both sides of an organic layer, called primary organic sheet (POS, Erez, 2003), sandwiched between the outer and inner organic layers. Scale bar 100 μm .

700

705



710 **Figure 5.** Evolution of the number of the attached specimens of the For1C1 strain in each culture medium at different $[SO_4^{2-}]$ for (a) Set 1 and (b) Set 2. The number of foraminifera counted (with a precision of ± 3 individual) corresponds to the living ones adhering to the bottom of the Petri dish before the change of medium. The increase in the number of individuals is due to asexual multiplication (see Appendix A). In Set 1, the largest population in terms of size occurs for 28.2 mM (ASW[28] and NSW Banyuls). In Set 2, a microbial bloom occurred after 12 days in medium ASW[28], likely affecting the reproduction rate (Appendix C, Appendix Fig. C1). The y-axes

715 are on log scales.






Figure 6. Evolution of pH and DIC over 35 days of For1c1 culture at 10 mM and 28.2 mM during SET2, with a change of medium on days 7, 14 and 26. Measured DIC in all experiments (Table 3) varies between 3.2 mM and 4.1 mM, it decreases by calcite formation and increases by specimen respiration. The variation of pH in all experiments when measured (Table 2) is between 7.89 and 8.19. The decrease in pH is caused by both respiration (CO₂ production) and calcite precipitation.

720



725 **Figure 7. Left panel:** $SO_4^{2-}/CaCO_3$ and S/Ca ratios on tests of the For1C1 strain at the end of SET1 and SET2 experiments, as a function of seawater $[SO_4^{2-}]$ (5, 10, 28.2, 35, 40 and 60 mM). **Right panel:** $SO_4^{2-}/CaCO_3$ and S/Ca ratios on tests of the For1C1 strain at the end of SET1 and SET2 experiments, as a function of seawater S/CO_3^{2-} . For our experimental results, we report the values using both S as the sum of free and complexed sulfate based on our model results (circles), and as only free sulfate (stars). Each measurement has been performed on a pool of hundred to several hundreds of specimens. Values are compared to other culture experiments of foraminifera targeting specifically the CAS content of the tests (Paris et al., 2014; van Dijk et al., 2017; 2019), and, when available, S/CO_3^{2-} as well. See Appendix B, Table B2 for details.

730



735 **Figure 8.** $\delta^{34}\text{S}$ in foraminiferal test (with or without organic matter) for the two strains For1C1 and C1Tg. The two strains were cultured in several media (For1C1 in ASW[5], ASW[10] ASW[28], ASW[35], ASW[40], ASW[60] and NSW (Banyuls), C1Tg in NSW (Concarneau) and ASW[28]) whose $\delta^{34}\text{S}$ has been also measured. The $\delta^{34}\text{S}$ value of the media depend on the salts used to make the solution. The green line corresponds to the $\delta^{34}\text{S}$ composition of the algae that were fed to the foraminifera, and whose isotopic composition remain stable. 2σ error bars ($\pm 0.2\text{‰}$ to $\pm 0.3\text{‰}$) are smaller than symbols.

Table 1. Weekly number of accumulated live individuals incremented by reproduction for each medium at different [SO₄²⁻]

SET 1		Seawater [SO₄²⁻]							
		0	5	10	NSW	28.2	60	120	180
Date	Days	Number of specimens							
23/03/2018	0	27	30	28	17	25	31	22	23
29/03/2018	6	27	60	61	17	63	36	22	23
04/04/2018	12	36	188	116	128	199	151	21	24
09/04/2018	17	21	227	187	294	317	98	13	17
18/04/2018	27	22	293	322	638	556	104	16	20
25/04/2018	34	20	425	713	732	1312	108	14	19

SET 2		Seawater [SO₄²⁻]						
		1	10	28.2	35	40	50	90
Date	Days	Number of specimens						
07/05/2018	0	6	6	6	6	6	6	6
14/05/2018	7	27	35	27	109	48	64	27
22/05/2018	15	82	109	138	141	97	89	32
28/05/2018	21	76	120	142	173	98	91	29
06/06/2018	31	87	737	151	444	127	97	25
08/06/2018	33	161	1014	159	470	194	117	17

740

Table 2. pH values measured in media all along SET1 and SET2 experiments.

Set 1								
Day	ASW[0]	ASW[5]	ASW[10]	ASW[28]	ASW[60]	ASW[120]	ASW[180]	Banyuls
0	8.09	8.09	8.08	8.1	8.1	8.14	8.14	nd
6	7.94	7.94	7.96	8.06	7.97	7.94	8.02	nd
7	8.17	8.19	8.19	8.19	8.19	8.17	8.2	nd
13	7.89	7.89	7.89	8.01	7.93	7.83	8.02	8.06*
14	8.17	8.19	8.19	8.19	8.19	8.17	8.2	nd
25	8.1	8.1	8.09	8.13	8.15	8.16	8.2	8.07*
35	8.05	8.07	8.07	8.11	8.12	8.14	8.16	8.03*

Set 2							
Day	ASW[1]	ASW[10]	ASW[28]	ASW[35]	ASW[40]	ASW[50]	ASW[90]
7	8.1	8.12	8.14	8.09	8.12	8.12	8.17
16	8.08	8.12	8.15	8.15	8.14	8.13	8.14

***: Foraminiferal culture in seawater from Banyuls started with a delay, making the pH measurements day 12, 22 and 26**

745 **Table 3.** DIC concentration in culture media all along Set 1 and Set 2 experiments

Days	6	12	13	14	22	25	29	35	38	46	52	53
Media	DIC mM +/- 4%											
ASW[1]									3.5			
ASW[5]				4.1								
ASW[10]	3.5		3.8			3.2		3.6				3.7
ASW[28]	3.9		3.9			4.1		3.8				3.7
NSW		3.5			3.7		3.8					
ASW[35]									3.6	3.8	3.7	
ASW[40]									3.6			
ASW[50]									3.5			
ASW[60]						4.1						
ASW[90]											3.9	

Table 4. Sulfate concentration and isotopic composition measured in the foraminiferal calcite

Species	Media	SO ₄ /CaCO ₃ ppm +/- 5%	Δ ³⁴ S*	δ ³⁴ S CAS ±0.2‰	δ ³⁴ S SO ₄ ²⁻ water ±0.2‰	[SO ₄ ²⁻] media (mM)
For1C1	ASW[5]	3320	1.7‰	1.6‰	-0.1‰	5
For1C1	ASW[10]	5780	1.5‰	1.4‰	-0.1‰	10
For1C1	ASW[28]	10800	Nd	Nd	Nd	28.2
For1C1	ASW[28]	12800	0.8‰	9.9‰	9.1‰	28.2
For1C1	NSW	14200	1.6‰	21.5‰	19.9	28.2
For1C1	ASW[35]	13700	1.6‰	1.5‰	-0.1	35
For1C1	ASW[40]	14000	1.3‰ ^a	1.2‰ ^a	-0.1	40
For1C1	ASW[60]	12800	1.8‰ ^b	1.7‰ ^b	-0.1	60
For1C1 + org	NSW	20100	1.2‰	21.2	20.0	28.2
For1C1 + org	ASW[28]	9700	-0.3‰	8.8	9.1	28.2
C1Tg	NSW	15600	1.3‰	21.3	20.0	28.2
C1Tg	NSW	13400	1.2‰	21.1	20.0	28.2
C1Tg	ASW[28]	10700	1.5‰	10.6	9.1	28.2
C1Tg	ASW[28]	10400	1.3‰	10.4	9.1	28.2
C1Tg	ASW[28]	nd	1.9‰	11.0	9.1	28.2
C1Tg + org	NSW	25600	-0.8‰	19.2	20.0	28.2
C1Tg + org	NSW	32900	0.3‰	20.3	20.0	28.2
C1Tg + org	ASW[28]	12400	0.2‰	9.3	9.1	28.2

*Δ³⁴S = δ³⁴S CAS - δ³⁴S SO₄²⁻ water

750 ^a The 2sd value of this sample is estimated to be 0.25 ‰.

^b The 2sd value of this sample is estimated to be 0.35 ‰.

6.1 Appendix A

Foraminifer taxonomy

760

The two selected strains come from two distinct locations, from Banyuls (Mediterranean Sea) and Concarneau (Atlantic Ocean). Morphologically they may be related to the family Rosalinidae (Holzmann and Pawlowski, 2017). They are attached forms with a low trochospiral hyaline calcitic perforate test, with a peripheral low arch aperture on the umbilical side bordered by lips (Fig. 2). Chamber interior is simple (Fig. 2). Two morphotypes are noticeable in both the strains (Fig. 2). Individuals usually reproducing asexually after every 12-15 days when their test reaches a development of 11-12 chambers (Fig. 3), are morphologically very close to the genus *Rosalina* (Fig. 2). Individuals who lived for several weeks adding more than 12 chambers have the last chambers with an annular arrangement (Fig. 2).

765

Individuals with a *Rosalina* like morphology (Fig. 2) probably belong to the schizont generation of their trimorphic life cycle (alternating gamont-agamont-schizont-gamont generations), documented for example in *Planorbulina mediterraneis* and a few dozen other species (Le Calvez, 1938; Dettmering et al., 1998). More precisely, they are diploid megalospheric schizonts that have entered a cycle of successive asexual reproduction (apogamic cycle) (Fig. 3), during which the new generation of schizonts is produced by schizogony, i.e. by multiple fission of a multinucleate parental cytoplasm (Le Calvez, 1938; Dettmering et al., 1998). For this reason, it is not obvious to identify them morphologically at the species level because the morphology of the diploid agamont microspheric and/or of the haploid megalospheric gamont parent generation, on which the description of the species has often been made, is unknown to the best of our knowledge. For now, we leave these forms in open nomenclature and call them by the name of the strains For1C1 and C1Tg. Adult specimens of these strains are smaller than the traditional foraminifer fraction obtained after sieving (through >125 µm mesh) in geochemical studies, they thus may be common, while rarely collected because of their size.

770

780 6.2 Appendix B – supplementary tables

Table B1. Sulfur isotope composition of media and algae cells

Sample	$\delta^{34}\text{S} \pm 0.2 \text{ ‰}$
NSW Banyuls before culture	21.1
NSW after feeding For1C1	19.9
ASW[28] before culture	9.1
ASW[28] after feeding For1C1	9.1
ASW[28] after feeding C1Tg	9.2
ASW(all used concentrations except 28) after feeding For1C1*	-0.1
Algae media	5.4
Algae cells	7.0

* Different salts were used to make all ASW and ASW28

785

Table B2. Comparison between different culture experiments. Our S/CO₃²⁻ ratios are given at 15% RSD based on the replicates of the model results at 28 mM and 10 mM. S* represents free uncomplexed sulfate in the solution.

Species		S/Ca mmol/mol	SO ₄ ²⁻ /CaCO ₃ ppm	Tested parameter	[SO ₄ ²⁻] mM r	S/CO ₃ ²⁻ mol/mmol	S*/CO ₃ ²⁻	Ref.
<i>A. lessonii</i>	min max	1.21 1.73	1161 1660	T°C	28 28	n.a.	n.a.	van Dijk et al., 2019
<i>S. marginalis</i>		8.95 9.6 10.4	8588	pCO ₂	28 28 28	0.12 0.14 0.18	n.a. n.a. n.a.	van Dijk et al., 2017
<i>A. gibbosa</i>		0.95 1.02 1.1 1.3	912 979 1056 1247	pCO ₂	28 28 28 28	0.12 0.14 0.18 0.29	n.a. n.a. n.a. n.a.	van Dijk et al., 2017
<i>A. lessonii</i>	min max	1.27 1.35	1219 1295	salinity	28 28	0.12 0.13	n.a. n.a.	van Dijk et al., 2017
<i>O. universa</i>		1.11 1.21 1.34 1.72	1062 1164 1289 1651	[SO ₄ ²⁻]	18 21 24 28	n.a. n.a. n.a. n.a.	n.a. n.a. n.a. n.a.	Paris et al., 2014
For1C1		3.46 6.02 14.80 11.25 14.22 14.59 13.38	3320 5781 14200 10800 13651 14004 12841	[SO ₄ ²⁻]	5 10 28 28 35 40 60	0.12 0.44 n.a. 0.81 1.03 1.2 1.51	0.04 0.15 n.a. 0.35 0.38 0.45 0.59	This study

790 6.3 Appendix C

Potential experimental bias

We designed Set 2 to replicate the ASW[28] and ASW[10] results as well as to extend the range of concentrations and ran it right after Set 1 (see methods for more details). Two differences can be observed. First the reproduction rate is significantly higher in the ASW[10] media of Set 2 than it was in Set 1 even though the media were identical (Fig. 5). However, it might be related to the starting number of foraminifera for each experiment, in Set 2 we started each culture experiment with 6 foraminifera individuals (instead of 28 as in Set 1), each 6 individuals were chosen more carefully, which could induce a bias and explain a more active behavior during experiment 2. If it were the case, the bias would nonetheless be systematic and similar for each medium (all Petri dishes in set 2 started with 6 individuals) and thus do not prevent comparison of results within set 2.

The second difference is observed in the ASW[28] medium in set 2. The reproduction rate, which was the highest observed, slowed down drastically after 15 days. This decrease can be explained by a microbial bloom in the media that was observed in no other media (Fig. C1). The microbial spread could not be reduced by the weekly water change, and any transfer and rinsing of

foraminifera or antibiotic treatment would have constituted an additional experimental modification. We thus kept counting
805 foraminifera and sampling seawater, but did not take into account any results collected in that media after day 15.




Figure C1. Optical microscopy imaging in dark field, the foraminifera are observed from below, the back ground appears black, algae greenish, and bacterial contamination cloudy white. (a) Foraminifera in ASW[10] during experiment 2 where microbial spread stays limited. (b) Foraminifera in ASW[28] during experiment 2 where microbial bloom was uncontrolled after 15 days and could not be reduced.

6.3 Appendix D

815 Geochemical modeling Parameters



The relative abundances of Ca^{2+} , SO_4^{2-} , $\text{CaSO}_{4(d)}$, $\text{MgSO}_{4(d)}$, $\text{NaSO}_{4(d)}$, CaHCO_3 , CO_3^{2-} , HCO_3^- and $\text{CO}_{2(d)}$ in solution were computed with the geochemical code JCHESS (Van der Lee, 1998). ASW composition from kester et al., 1967 and experimental temperature (20 C) were chosen as input parameters assuming a closed system with no gas-solution exchange, and Cl, Na and SO_4^{2-} were modified as they were in each media of the experiment (Base input in the additional excel table). All DIC is provided
820 as HCO_3^- to the model.

Given that pH was adjusted by NaOH addition, this was reproduced in the model, by adding Na^+ and OH^- (in the same concentration) until reaching pH 8.2.

From this starting point, HCO_3^- was adjusted to the measured DIC value, and OH^- was adjusted to match the measured pH value (Adjusted input, and measured values in the additional excel table). No additional Na^+ was added despite the slight electrical imbalance generated, as Na^+ can form complexes with SO_4^{2-} and no Na^+ was provided to the media after pH has been adjusted.
825 The output data considered are Ca^{2+} , all free and complexed SO_4^{2-} species, free DIC species and CaHCO_3^- . The sole DIC specie present as a complex that was extracted is CaHCO_3^- , because it is the major complex. It is also a species that has been hypothesised to be potentially incorporated into calcite, as CaSO_4 could be.

830 Geochemical modelling results

The $\text{SO}_4^{2-}/\text{CO}_3^{2-}$ concentration increases linearly to a slight inflection point at 60mM, linked to complexes formation. Nevertheless, we do not observe a plateau from 28mM onwards, which could have explained a constant incorporation of SO_4^{2-} in calcite beyond 28mM. Similarly, assuming that SO_4^{2-} incorporation into calcite takes place from CaSO_4 , although an inflection of the $\text{CaSO}_4/\text{CaHCO}_3$ ratio is observable from 40mM, no plateau is observed. These results show that the incorporation of sulfate into
835 the calcite of the foraminifers in our experiments, which plateau above 28mM, cannot be explained by the formation of complexes in seawater. Alternatively, CAS concentration is not a good recorder of either the $\text{SO}_4^{2-}/\text{CO}_3^{2-}$ ratio or the $\text{CaSO}_4/\text{CaHCO}_3$ ratio above a concentration of 28mM.



840

Figure D1. Geochemical modeling results showing $\text{SO}_4^{2-} / \text{CO}_3^{2-}$ ratio and $\text{CaSO}_4 / \text{CaHCO}_3$ ratio in solution as a function of total sulfate concentration in solution. Each point corresponds to a different computational run, for samples where DIC and pH were measured, and were used as constraints to the model. Both ratios increase linearly to a slight inflection point at 60mM, but no plateau is seen between 28mM and 60mM.

845