

GPP and the predictability of CO₂: more uncertainty in what we predict than how well we predict it

István Dunkl^{1,2}, Nicole Lovenduski³, Alessio Collalti⁴, Vivek K. Arora⁵, Tatiana Ilyina¹, and Victor Brovkin^{1,6}

¹Max Planck Institute for Meteorology, Hamburg, Germany

²International Max Planck Research School on Earth System Modelling, Hamburg, Germany

³Department of Atmospheric and Oceanic Sciences and Institute of Arctic and Alpine Research, University of Colorado, Boulder, CO, USA

⁴Forest Modelling Lab., Institute for Agriculture and Forestry Systems in the Mediterranean, National Research Council of Italy (CNR-ISAFOM), Perugia, Italy

⁵Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, University of Victoria, Victoria, British Columbia, Canada

⁶Center for Earth System Research and Sustainability, University of Hamburg, Germany

Correspondence: István Dunkl (istvan.dunkl@mpimet.mpg.de)

Abstract. The prediction of atmospheric CO₂ concentrations is limited by the high interannual variability (IAV) of terrestrial gross primary productivity (GPP). However, there are large uncertainties in the drivers of GPP IAV among Earth system models (ESMs). Here, we evaluate the impact of these uncertainties on the predictability of atmospheric CO₂ in six ESMs. We use regression analysis to determine the role of environmental drivers on (i) the patterns of GPP IAV, and (ii) the predictability of GPP. There are large uncertainties in the spatial distribution of GPP IAV. Although all ESMs agree on the high IAV in the tropics, several ESMs have unique hotspots of GPP IAV. The main driver of GPP IAV is temperature in the ESMs using the Community Land Model, and soil moisture in IPSL-CM6A-LR and MPI-ESM-LR, revealing underlying differences in the source of GPP IAV among ESMs. Between 13% and 24% of the GPP IAV is predictable one year ahead, with four out of six ESMs between 19% and 24%. Up to 32% of the GPP IAV induced by soil moisture is predictable, while only 7% to 13% of the GPP IAV induced by radiation. The results show that while ESMs are fairly similar in their ability to predict themselves, their predicted contribution to the atmospheric CO₂ variability originates from different regions and is caused by different drivers. A higher coherence in atmospheric CO₂ predictability could be achieved by reducing uncertainties of GPP sensitivity to soil moisture, and by accurate observational products for GPP IAV.

1 Introduction

Near-term predictions of atmospheric CO₂ concentrations are an essential step towards the evaluation of climate mitigation efforts and the development of carbon monitoring programs (Ilyina et al., 2021). However, the high interannual variability (IAV) of land-atmosphere carbon fluxes, specifically gross primary productivity (GPP), drives the variability of atmospheric

CO₂ and limits its predictability (Piao et al., 2020). The skilful prediction of GPP is therefore a crucial step towards the real-time verification of anthropogenic carbon emissions and the evaluation of mitigation efforts.

20 The usual approach to evaluate the predictability of an earth system variable is to compare predictions with observed values. In the case of GPP, this is complicated by the uncertainty in GPP observations (Zhang and Ye, 2021). As an alternative to calculating the actual predictability that is based on observations, the potential predictability can be assessed by evaluating how well the models can predict themselves. To do this, an ensemble of simulations with an Earth system model (ESM) is initialized from quasi-identical conditions. In a system with little predictability, the spread among the ensemble members increases quickly, until the predictive capabilities are lost when the ensemble spread reaches the magnitude of the IAV. There are, however, certain processes in the earth system which provide predictability and hinder the divergence of the ensemble members. For example, the El Niño-Southern Oscillation (ENSO) produces predictable climate anomalies that have a sustained impact on GPP (Zeng et al., 2008; Betts et al., 2016). Other processes extend predictability by providing “memory” that maintains the initial conditions. Soils, for example, store initial moisture anomalies by acting as a buffer between the atmosphere and the vegetation (Bellucci et al., 2015). Soil moisture anomalies are further extended through land-atmosphere coupling, which creates a feedback loop that enhances the persistence of these anomalies (Kumar et al., 2020). The initial conditions of the simulations are maintained through the lagged response of plant growth to climatic conditions. Slowly reacting vegetation can cause precipitation anomalies or prolonged drought (Alessandri and Navarra, 2008; Zhang et al., 2021). Given all of these mechanisms of predictability, we find that terrestrial carbon fluxes are predictable for two years (Ilyina et al., 2021).

35 Although several ESMs reproduce the same predictability horizon for globally integrated terrestrial carbon fluxes (Séférian et al., 2018; Ilyina et al., 2021; Spring and Ilyina, 2020; Lovenduski et al., 2019), there are substantial differences in the spatial patterns of GPP IAV (Anav et al., 2015; O’Sullivan et al., 2020). The reason for these differences lies in poorly constrained ecosystem processes that have a large impact on GPP. One of these differences arises from the uncertainty in the sensitivity of GPP to environmental drivers (Ahlström et al., 2015; Jung et al., 2017; Beer et al., 2010; Piao et al., 2020; Collalti et al., 2020). The sensitivity of GPP to temperature and precipitation varies among studies, leading to the ongoing discussion concerning the dominant driver of global carbon fluxes (Piao et al., 2020). The different sensitivity of GPP to precipitation across ESMs is further exacerbated by the large disagreement in water storage anomalies (Wu et al., 2021). The simulated annual cycle of water storage anomalies of major river basins is between 0.1 and 2 times that of the observed variability. These deviations in hydrological variability between models are likely to cause similar deviations in GPP IAV, especially in semi-arid watersheds. Further differences in GPP IAV are due to variations in ecosystem boundaries and the related spatial distribution of plant productivity. The Amazon rainforest, for instance, is a hotspot of land-atmosphere carbon fluxes and provides a large contribution to the predictability of atmospheric CO₂ (Zeng et al., 2008; Séférian et al., 2018; Ilyina et al., 2021). However, the transition zone between the wet tropical forest and semi-arid tropics within the Amazon basin varies among the models due to differences in their representation of land cover (Collier et al., 2018; Hu et al., 2022). Such differences in biome boundaries also modify the impact of ENSO on GPP IAV. ENSO produces a distinct spatial pattern of climatic anomalies which influences the GPP on 32% of the vegetated land area significantly (Zhang et al., 2019). These ENSO-related climate patterns will have a different impact on GPP depending on the type of biomes under their influence. In addition to the spatial variability, many ESMs strug-

gle to reproduce the seasonal variability of carbon fluxes. This can be seen in the large biases in phenology (Song et al., 2021). Several models overestimate the seasonal amplitude of leaf area index (LAI) in the tropics, and mismatch the timing of LAI maxima and minima (Peano et al., 2019, 2021).

All of these uncertainties suggest, that there are large differences in the patterns of GPP IAV among the ESMs, but it is yet unclear how these differences affect the predictability of GPP. With this study, we want to extend our understanding of GPP predictability by considering the different patterns of GPP IAV among the ESMs. In a multi-model analysis, we investigate which processes drive the IAV of GPP and which processes allow the GPP IAV to be predictable. Regression analysis is used to determine the role of three environmental variables (soil moisture, temperature, and radiation) on GPP IAV and GPP predictability. We analyse the cause of differences in GPP predictability across ESMs, identify the areas of large discrepancies, and determine the factors contributing to the attached uncertainties. The aim of this study is to reveal which factors of GPP representation are limiting the predictability of atmospheric CO₂.

2 Methods

2.1 Data sources

We analyse model output from the Decadal Climate Prediction Project (DCPP, Boer et al. 2016). This protocol-driven multi-model approach aims at studying the decadal predictability of the earth system with hindcasts, quasi-real-time forecasts, and case studies on predictability mechanisms. The hindcasts are initialized annually from 1960 to 2017 or 2019 with the starting dates between November and January and at least 10 ensemble members. Simulations are driven by CMIP5 or CMIP6 historical forcing and extended by RCP4.5 or SSP2-4.5 afterwards. The DCPP framework does not prescribe any specific initialization or data assimilation methods and leaves these details to be decided by the respective modelling centres.

We additionally use the CESM2 model output from the Seasonal-to-Multiyear Large Ensemble (SMYLE) prediction system (Yeager et al., 2022). The SMYLE hindcasts ensembles are initialized four times per year with 20 ensemble members between 1970 and 2019. In this study, the November initializations are used to achieve the highest comparability with the DCPP hindcasts.

We compare the spatial GPP IAV patterns of the ESMs with observation-based GPP products. Because of the uncertainty among observations, we include products based on three different sources. The Vegetation Photosynthesis Model (VPM, Zhang et al. 2017) is remote sensing-based product that uses a light use efficiency (LUE) model to calculate GPP. VPM uses satellite data from MODIS and an improved LUE algorithm that considers leaf quality. The second data set is GOSIF (Li and Xiao, 2019) which is based on data from MODIS and the Orbiting Carbon Observatory-2. GOSIF uses solar-induced chlorophyll fluorescence, which is a more recent approach to calculate GPP. Lastly, we use FLUXCOM (version RS + METEO (ERA5), Jung et al. 2019), which uses machine learning to upscale flux tower observations with meteorological and remote sensing data. Because FLUXCOM underestimates the IAV of GPP (Anav et al., 2015; O'Sullivan et al., 2020), it is recommended to scale the data so that the IAV of its integrated fluxes resembles observations (Jung et al., 2019). The VPM, GOSIF, and FLUXCOM

85 data are linearly detrended before calculating the IAV. Due to the long time span, FLUXCOM is detrended over two periods (1979 to 1999, and 2000 to 2018).

CanESM5

The Canadian Earth System Model version 5 (CanESM5; Swart et al. 2019) consists of the Canadian Land Surface Scheme (CLASS) and Canadian Terrestrial Ecosystem Model (CTEM) with a T63 grid of an approximate resolution of 2.8° . The atmosphere is realized with the Canadian Atmospheric Model (CanAM5) with 49 vertical levels. Ocean physics is simulated with
90 CanNEMO, on a tripolar grid with a resolution of 1° to $1/3^\circ$ and 45 vertical levels, and ocean biogeochemistry is represented by the Canadian Model of Ocean Carbon (CMOC).

The CanESM5 hindcast simulations are initialized every January between 1960 and 2017 with 20 members. 3D potential temperature and salinity of the global oceans are nudged toward monthly Ocean Reanalysis System 5 (ORAS5; Zuo et al. 2019).
95 Sea surface temperatures are nudged to Extended Reconstructed Sea Surface Temperature (ERSSTv3; Xue et al. (2003); Smith et al. (2008)) until 1981 and to Optimum Interpolation Sea Surface Temperature (OISST; Banzon et al. 2016) afterwards. Sea ice concentration is nudged to the Hadley Centre Sea Ice and Sea Surface Temperature data set (HadISST.2; Titchner and Rayner 2014), and sea ice thickness to monthly climatology until 1988 and to the SMv3 statistical model of Dirkson et al. (2017) afterwards. For the atmosphere, temperature, horizontal wind components and specific humidity are nudged to ERA40
100 (Uppala et al., 2005) until 1978 and to 6-hourly ERA-Interim data (Dee et al., 2011) afterwards.

CESM1-CAM5

The Community Earth System Model (CESM) version 1.1 (Hurrell et al., 2013) is used to produce 40-member simulations in the Decadal Prediction Large Ensemble (DPLE) project (Yeager et al., 2018). The model components are the Community Land Model version 4 (CLM4; Lawrence et al. 2011) with a 1° resolution, Community Atmosphere Model Version 5 (CAM5) with
105 30 vertical levels, the Parallel Ocean Program (POP2) with 60 vertical levels and sea ice with Community Ice Code (CICE4).

The CESM1-CAM5 hindcasts are initialized every November. There is no direct assimilation of observations to produce the initial conditions, instead, ocean and sea ice are obtained from simulation runs forced by historic atmospheric surface fields (Yeager et al., 2018). Initial conditions for the land and atmosphere components are obtained from ensemble member #34 of the CESM Large Ensemble (Kay et al., 2015; Lovenduski et al., 2019).

110 CESM2

CESM version 2 (Danabasoglu et al., 2020) runs on a 1° horizontal resolution of all components. The atmosphere is simulated by the Community Atmosphere Model Version 6 (CAM6) with 32 vertical levels. The ocean model is the Parallel Ocean Program version 2 (POP2) with 60 vertical levels, with the biogeochemistry from the Marine Biogeochemistry Library and sea ice by CICE version 5.1.2 (CICE5) with 8 vertical layers. The land component is simulated by the Community Land Model

115 version 5 (CLM5; Lawrence et al. 2019), which has several updates to its predecessor CLM4 and CLM4.5, leading to a better representation of the global carbon cycle in benchmarks (Bonan et al., 2019).

Hindcasts are initialized on the 1st of every November, February, May and August, and run for 24 months. Only the November initializations are used in this analysis to increase comparability with the DCPD simulations. Initial conditions for the atmosphere, ocean and sea-ice stem from the Japanese 55-year Reanalysis (JRA-55; Kobayashi et al. (2015), and JRA55do; 120 Tsujino et al. (2018)). The land surface and biogeochemistry are initialized from forced CLM5 simulations.

CMCC-CM2-SR5

The Euro-Mediterranean Centre on Climate Change coupled climate model (CMCC-CM2, Cherchi et al. (2019), Lovato et al. (2022)) is based on CESM and consists of the Community Land Model (CLM4.5) with a 1° resolution, the atmospheric model CAM5.3 with 30 vertical levels. The distinguishing element of CMCC-CM is the ocean, which is simulated by NEMO3.6, 125 while sea ice is modelled by CICE4.

The 10-member hindcast simulations are initialized every November (Nicolì et al., 2022). The ocean initial conditions are from CHOR (Yang et al., 2017) until 2010 and from CGLORSv7 (Storto and Masina, 2016) afterwards. The atmosphere is initialized from ERA-40 until 1978 and by ERA-Interim afterwards. The land surface is initialized using the reanalysis with two different meteorological forcing. For this reason, only the ensemble members 1, 3, 5, 7, 8 and 9 are used, as the other 130 members start from a different state and this would not allow quantifying predictability by ensemble spread.

Because the CMCC-CM2-SR5 fields containing land-atmosphere carbon fluxes are not exported for the DCPD runs, the historical simulations are used to infer the relationship between environmental drivers and GPP.

IPSL-CM6A-LR

The ESM developed by the Institute Pierre Simon Laplace (IPSL; Boucher et al. 2020) uses the ORCHIDEE v2.0 (Cheruy et al., 2020) land surface model (LSM) with an average resolution of 157 km. The atmosphere is simulated at the same 135 resolution by LMDZ6 with 79 vertical levels, the ocean with NEMO-OPA with a 1° resolution and 75 vertical levels and ocean biogeochemistry with PISCESv2.

The hindcast simulations of IPSL-CM6A-LR come from the DCPD project. The 10-member ensembles start annually in January between 1960 and 2016. The hindcasts are initiated from an assimilation run with EN4 sea surface temperatures 140 (Good et al., 2013) and Atlantic sea surface salinity (Estella-Perez et al., 2020). Subsurface ocean, sea ice and atmosphere are not assimilated.

MPI-ESM-LR

MPI-ESM-LR is the Max Planck Earth System Model (MPI-ESM1.1; Giorgetta et al. 2013) used in a low-resolution configuration. The land is simulated by JSBACH with dynamic vegetation (Reick et al., 2013). The ocean component is MPIOM with

145 a horizontal resolution of about 150 km and 40 vertical levels. The atmosphere is simulated by ECHAM at a T63 resolution with 47 vertical layers, and ocean biogeochemistry is represented by HAMOCC.

The utilized hindcast simulations of MPI-ESM-LR are conducted within the MiKlip project (Marotzke et al., 2016). The decadal prediction system are 10-member ensembles starting every January between 1961 and 2014. Ocean temperature and salinity are initialized from the Ocean Reanalysis System 4 (ORAS4; Balmaseda et al. 2013) and the atmosphere by ERA-40
150 from 1960 to 1998 and ERA-Interim from 1990 to 2014.

2.2 Statistical approach




Figure 1. Workflow of the statistical analysis: a) Lead years five to ten of the hindcast simulations are used to train a regression model that calculates the components of GPP caused by the environmental drivers. b) The regression model is applied to (i) lead years five to ten to calculate the IAV of the GPP components ($\sigma GPP_{(IAV)}$) and (ii) to lead year one to calculate the mean ensemble spread of the GPP components ($\sigma GPP_{(LY1)}$).

Overview

An overview of the statistical analysis is shown in Figure 1. Every hindcast simulation is initialized from quasi-identical conditions. With the increasing lead-time, the variability within the hindcast ensemble (standard deviation across the ensemble
155 members for a given time) increases too, until it reaches the IAV. Based on this assumption, the hindcast simulations are split into two groups by lead-time (lead year one, and lead years five to ten). For the lead years five to ten, the effects of ocean and atmosphere initialization are assumed to be negligible. These years are used to calculate the monthly mean climatology, which is removed from both groups to obtain the anomalies. The anomalies of the lead years five to ten are used in a regression analysis to derive the sensitivity of GPP to the environmental drivers i.e. soil moisture, temperature and radiation (Fig.
160 1 a)). The regression model is applied to (i) the anomalies of lead years five to ten to calculate the IAV of all GPP components ($\sigma GPP_{(IAV)}$), and (ii) to the anomalies of lead year one to calculate the ensemble variability of all GPP components

($\sigma GPP_{(LY1)}$, Fig 1 b)). We derive the predictability of GPP by comparing $\sigma GPP_{(LY1)}$ to $\sigma GPP_{(IAV)}$. Because the hindcast simulations are not evaluated against observations, the calculated predictability reflects the potential predictability.

Climatology and sensitivity

165 The monthly mean climatologies are calculated from the lead years five to ten, with a three-year moving window approach for every calendar year. Because the moving window method is not applicable for the first decade of hindcast initializations, the monthly climatology for the 1960s (1970s for CESM2) is calculated based on all lead years five to ten within the 1960s (or 1970s). Anomalies of all input fields are calculated by subtracting the monthly climatologies from the hindcast data. The obtained anomalies of lead years five to ten make up a data set of n simulation years:

$$170 \quad n = 6 \text{ hindcast years} \times \text{No. ensemble members} \times \text{No. initializations}. \quad (1)$$

With 10 to 40 ensemble members and 56 to 58 initializations resulting in sample sizes of 3330 to 13680. Because the hindcast length is only two years for the CESM2 simulations, a different approach is used here. Instead of lead years five to ten, only lead year two is selected and only five random ensemble members are used from every hindcast to reduce the number of simulations with the same initial conditions. To offset the reduced number of data points, five random simulations are added
175 from the hindcast simulations initialized in February, May, and August as well.

The resulting data set of lead year 5 to ten anomalies is used to derive the sensitivity of GPP to the environmental drivers (ENV : soil moisture, temperature and radiation) by fitting a regression model for every grid cell and month of the year. The relationship between GPP and the environmental drivers is frequently non-linear, sometimes due to specific break points in the functional representation of GPP. For this reason, segmented linear regression (SLR) is used to model GPP from the envi-
180 ronmental drivers (Muggeo, 2008). SLR finds breakpoints in the data, splitting it into multiple ranges and fitting an individual regression model to each of the data ranges. Here, a single breakpoint is determined for each of the three predictor variables.

Because environmental drivers have some degree of collinearity, the regression analysis will not be able to fully attribute the GPP anomalies to their specific causes. Therefore, the resulting sensitivities should be taken as a “contributive”, and not a “true” effect of the environmental drivers (Wang et al., 2016).

185 Variability and predictability

The SLR can now be applied to individual simulations to determine the component of GPP anomalies that can be attributed to each of the environmental drivers:

$$\Delta GPP \approx \Delta GPP^{Soil \text{ moisture}} + \Delta GPP^{Temperature} + \Delta GPP^{Radiation}. \quad (2)$$

The three components of GPP anomalies (ΔGPP^{ENV}) are calculated for every simulation within the hindcast lead time five
190 to ten. From the results, we calculate the IAV of the components for every grid cell and month of the year ($\sigma GPP_{(IAV)}^{ENV}$). Similarly, the SLR is applied to the anomalies of lead year one, to calculate the standard deviation for every month within the

hindcast simulations. Averaging over the standard deviations of every hindcast returns the ensemble variability of lead year one ($\sigma GPP_{(LY1)}^{ENV}$).

The predictability is assessed by comparing $\sigma GPP_{(LY1)}$ to $\sigma GPP_{(IAV)}$. A high predictability of an input field means, that the ensemble variability is restricted for some time after the hindcast initialization, and does not reach the IAV immediately (Fig. 2 a)). In this study, we use two metrics to evaluate different aspects of predictability. We calculate the fraction of GPP IAV that is predictable (predictable fraction, pf) to assess the ability of a system to retain memory. Although this metric is useful for quantifying the mechanisms that provide predictability at a local level, pf is not suitable for assessing how GPP predictability allows the predictability of atmospheric CO₂. This is because the regions with a high pf do not necessarily contribute much to the global GPP fluxes. The regions with the highest pf are often in deserts with very little carbon fluxes (Dunkl et al., 2021). To assess the contribution of GPP predictability to atmospheric CO₂ predictability, we calculate the absolute portion of the IAV which can be predicted as the predictable component (pc). The pc is the difference between IAV and ensemble variability and is generally higher in regions that contribute more to CO₂ IAV:

$$pc^{ENV} = \sigma GPP_{(IAV)}^{ENV} - \sigma GPP_{(LY1)}^{ENV}, \quad (3)$$

$$pf^{ENV} = \frac{pc^{ENV}}{\sigma GPP_{(IAV)}^{ENV}}. \quad (4)$$

The use of the two predictability metrics is exemplified in Figure 2 b).

3 Results and Discussion

3.1 Patterns of GPP IAV

In order to understand what the models are predicting, we start by analysing the patterns of GPP IAV. There are differences in the overall magnitude of GPP IAV among ESMs with CanESM5, CMCC-CM2-SR5, and IPSL-CM6A-LR at the lower, and CESM2 and MPI-ESM-LR at the higher end of the IAV spectrum. Factors that could explain some of the differences in the overall magnitude of IAV is the relatively weak ENSO teleconnection in CanESM5 (Swart et al., 2019), or the low total GPP in CMCC-CM2-SR5 (Lovato et al., 2022).

Because we focus on the spatial patterns of IAV rather than absolute differences, the GPP IAV patterns are scaled for better comparison (Fig. 3). We find agreement in the large-scale patterns of GPP IAV, with most of the IAV of the ESMs in the northern Amazon Basin, and the semi-arid tropics like western South America, Southern Africa, South Asia, Australia, and southern North America (Detailed maps of the location of the semi-arid regions in the ESMs are shown in Fig. 5). A closer examination of GPP IAV reveals that the ESMs have less agreement in the regions contributing most to the IAV, especially in the semi-arid tropics. Some ESMs have large hotspots of GPP IAV, which cannot be found in other ESMs. These unique hotspots are the western Amazon Basin (CanESM5), Central South America (CESM2), Southern Africa (MPI-ESM-LR and




Figure 2. a) The exemplary composition of GPP variability and predictability in a tropical forest. The components of GPP IAV are calculated from the lead years five to ten (green bars), and the ensemble variability is calculated from lead year one (red bars). In the exemplified region, most of the variability is caused by soil moisture and radiation, while GPP is not restricted by temperature. Predictability is exclusively provided through soil moisture. b) Demonstrating the need for the two predictability metrics by the example of a tropical savanna and an arid shrubland. The predictable component (pc) is the absolute predictable IAV, and the predictable fraction (pf) is pc scaled by the IAV. While the arid shrubland has a better potential to retain memory (seen by the high pf), these ecosystems contribute little to the variability of atmospheric CO_2 , which can be better assessed by pc .

CanESM5), and Australia (MPI-ESM-LR). We can find the most consistency on the northern coast of South America, which is a high IAV region in most ESMs. The spatial patterns of IAV have an average correlation of 0.47 among the ESMs. The ESM with the lowest correlation values is CESM2 with an average of 0.29. CESM2 stands out with very low IAV in the tropical rainforests of the Amazon and Congo Basins and Southeast Asia.

The correlation among the observational products is 0.65, and although they confirm most of the IAV patterns of the ESMs we find stronger deviations in South America. While many ESMs have IAV hotspots along the northern coast of South America, this is only reproduced in FLUXCOM. However, all observational products show a high GPP IAV in western South America, which can not be found in the ESMs. The spatial patterns of GPP IAV revealed here correspond with the literature, which suggests that the semi-arid tropics, tropical forests, grasslands and croplands are the main drivers of global GPP IAV (Ahlström et al., 2015; Piao et al., 2020; O’Sullivan et al., 2020). These studies also reflect the large uncertainty in the contribution of the individual semi-arid regions to GPP IAV between the models, and in particular the uncertain role of Australia. In an ensemble of eight LSMs, Australia contributed 39%, semi-arid tropical Africa 32%, and Southeast Asia 10% to global GPP IAV, while temperate South America only contributed 2% (Chen et al., 2017). Although Australia has the highest mean model IAV, the variability of IAV between the models is also the largest, with the IAV of GPP ranging between 0.26 and 1.01 Pg C yr^{-1} . While the large role of the dry tropics in driving GPP IAV is not disputed, it is likely that ESMs underestimate GPP IAV in wet tropical forests (O’Sullivan et al., 2020). This results from a limited availability of observations due to few flux towers,




Figure 3. GPP IAV in three observational products (VPM, GOSIF, and FLUXCOM) and six ESMs. a) Spatial patterns of GPP IAV with brighter colours standing for higher values. The data is scaled across ESMs to highlight differences in patterns and not absolute differences. b) The spatial correlations between the products.

and because the quality of remote sensing products is limited in tropical forests due to saturation effects and a high cloud cover

(Kolby Smith et al., 2016). In this study, the low GPP IAV in tropical forests is especially evident for CESM2, where IAV
240 increases abruptly outside the wet tropical forests.

The divergence in GPP IAV across different ESMs is largely caused by three factors: the sensitivity of carbon fluxes to climatic drivers (Piao et al., 2020) (discussed in section 3.2), phenology (Chen et al., 2017; Peano et al., 2019, 2021), and meteorological input (Anav et al., 2015). The role of phenology is crucial because the amount and quality of leaves determines the carbon fluxes between the land and the atmosphere (Peano et al., 2021). Most LSMs tend to have a better representation of
245 the growing season type, and growing season boundaries in the wet than in the semi-arid tropics. Peano et al. (2021) analysed the start- and ending months of growing seasons in eight LSMs under the same climate forcing and found several regions with a wide range of simulated growing seasons. The largest uncertainties in growing season are in the semi-arid tropics, the same regions where we find little agreement in GPP IAV. Growing season start ranges from February to October in Australia, and from March to October in Southern Africa, while growing season end ranged from March to September in Africa between
250 0 and 15°N. The vegetation types with the largest uncertainty in growing season timing are broadleaf deciduous shrubs, which are mostly located in Northern Australia, Southern Hemisphere crops, broadleaf evergreen trees and grasses. The better-performing LSMs have a high number of plant functional types or use more complex phenology schemes that depend on plant functional types and use a larger number of environmental variables to constrain phenology. Its complex phenological scheme puts ORCHIDEE among the better-performing LSMs, and might explain the high correlation of IPSL-CM6A-LR with the GPP
255 IAV of all three observational products.

It is also a misrepresentation of phenology that can explain the overall high IAV in MPI-ESM-LR. JSBACH overestimates the seasonality of LAI in the tropics, as it becomes visible in the strong seasonal cycle of tropical LAI in MPI-ESM-LR (Song et al., 2021). Consequentially, the area of the evergreen tropics is underestimated in JSBACH (Peano et al., 2021). This leads to a larger fraction of semi-arid tropics with a higher GPP IAV. This amplification of the equatorial dry season might lead to
260 the high GPP IAV in the Northern Amazon and contribute to the overall high IAV in MPI-ESM-LR (Wang et al., 2011).

3.2 Drivers of GPP IAV

To determine the drivers of GPP IAV, we analysed the sensitivity of GPP to environmental drivers using regression analysis. The globally averaged contribution of the drivers to GPP IAV is shown as the bars in Figure 4. The CLM family and CanESM5 have similar patterns, with temperature dominating the IAV, or being on par with soil moisture. IPSL-CM6A-LR and MPI-
265 EMS-LR have distinctly different patterns, with soil moisture dominating the IAV and radiation contributing equally or more than temperature. A reason for the large contribution of soil moisture to GPP IAV in IPSL-CM6A-LR and MPI-ESM-LR could be that both ESMs are at the high end of soil moisture IAV for deep soil layers in the Southern Hemisphere (Qiao et al., 2022), where many of the semi-arid ecosystems are located that contribute most to GPP IAV. Another explanation could be that out of eleven ESMs, IPSL-CM6A-LR and MPI-ESM-LR have the lowest warm-season soil moisture (Padrón et al., 2022). This
270 increase in dryness can lead to a larger extent of semi-arid ecosystems with a generally higher GPP IAV. Another effect of the reduced warm-season soil moisture can be an increase in land-atmosphere coupling strength (Santanello et al., 2018). Stronger land-atmosphere coupling would explain the higher correlation between soil moisture and temperature in IPSL-CM6A-LR and

MPI-ESM-LR (Padrón et al., 2022), and make the regression coefficients shift towards the stronger predictor, which is soil moisture.




Figure 4. The contribution of environmental drivers to GPP variability ($\sigma GPP_{(IAV)}^{ENV}$). Colour intensity stands for higher GPP IAV. The data is scaled across ESMs to highlight differences in patterns and not absolute differences. Bars represent the mean contribution of environmental drivers to global GPP IAV ($\text{kg C m}^{-2} \text{s}^{-1} 10^{-13}$).

275 The spatial drivers of GPP IAV show agreement in the wet and arid tropics, while there is little consistency in the semi-arid transition zones (Fig. 4). In many ESMs, the GPP IAV in the wet tropics, and eastern China is induced by radiation, while soil moisture becomes more prevalent along the aridity gradient, and is driving IAV in Southern Africa, Southern South America, and Australia. The IAV of the remaining land surface is driven predominantly by soil moisture in IPSL-CM6A-LR and MPI-ESM-LR, and by a combination of temperature and soil moisture in the remaining ESMs.

280 Some of the differences in GPP sensitivity among the ESMs can be explained by differences in aridity. A higher sensitivity to soil moisture can result from a dryer climate. The distribution of climate zones in the analysed models is shown in Figure 5 as the De Martonne aridity index (Gavrilov et al., 2019). MPI-ESM-LR and CanESM5 show an above average extent of arid and semi-arid regions in Australia and Southern Africa. This could explain the high sensitivity of GPP to soil moisture in these regions. Differences in climate also explain some of the discovered GPP patterns in the Amazon Basin. CESM2 is the model
 285 with the most humid climate in the Amazon Basin, which could be the reason for the low sensitivity of GPP to soil moisture and the generally low GPP IAV in this region. We find CanESM5 on the other side of the spectrum, with a relatively dry Amazon Basin, leading to a higher sensitivity to soil moisture and a high GPP IAV. However, there are also differences in GPP sensitivity which cannot be explained by differences in climate. IPSL-CM6A-LR is more or equally humid in Australia, Southern Africa

and South America than the models of the CLM family, despite having a high sensitivity of GPP to soil moisture in these
 290 regions. These differences are more likely to be caused by differences in their land surface models than climate.




Figure 5. De Martonne aridity index of the analysed models.

The general patterns of GPP sensitivity agree with reported sensitivity patterns in the literature. Multi-model averages and observations of GPP sensitivity agree with the larger role of temperature in tropical forests, radiation in western Amazonia, and the importance of precipitation in the semi-arid tropics (O’Sullivan et al., 2020; Anav et al., 2015). However, the role of water on carbon fluxes increases when soil moisture is used instead of precipitation in sensitivity studies (Piao et al., 2020).
 295 This can be observed in the sensitivity of net biome productivity (NBP), showing a more balanced contribution of soil moisture and temperature in the wet tropical forests (Piao et al., 2020; Padrón et al., 2022). Although the comparison of GPP and NBP imposes limitations, GPP explains the majority of tropical NBP (Ahlström et al., 2015). This suggests, that the low water sensitivity of tropical GPP might explain the lower than expected GPP IAV of tropical forests in ESMs.

3.3 Predictability of GPP

300 To analyse the role of GPP in the predictability of atmospheric CO₂, we assessed GPP predictability using two metrics. The predictable component (*pc*) is calculated as the difference between ensemble variability and IAV and provides a measure of absolute predictable variability. *pc* can be used to assess the predictability of GPP fluxes that contribute to CO₂ variability. The predictable fraction (*pf*) is the ratio of *pc* to IAV and illustrates how well memory is retained in the system. This metric can be used to compare the predictive performance of different biomes, for example.




Figure 6. The contribution of environmental drivers to the predictable component of GPP (pc). The contribution is calculated as the difference between the IAV and the ensemble variability within lead year one of the hindcast experiments. Values are scaled for each ESM. Bars represent the mean contribution of environmental drivers to the pc ($\Delta\sigma$ GPP in $\text{kg C m}^{-2} \text{s}^{-1} 10^{-13}$). Numbers on top of the bars show the predictable fraction (pf), which is the share of the pc to overall IAV. The correlation between GPP IAV and the pc is shown at the bottom of the plots.

305 There is relatively high consistency among the pf of the environmental drivers across the models ($pf^{\text{Soil moisture}} > pf^{\text{Temperature}} > pf^{\text{Radiation}}$, numbers above the bars in Fig. 6). This pattern reflects the anticipated differences in predictability among the drivers. Atmospheric fields, as radiation, have a low persistence, leading to a low predictability of two weeks for most regions (Zeng et al., 2008). Soil hydrology, on the other hand, acts as a low-pass filter which removes the unpredictable high-frequency variability of precipitation and allows a predictability of soil moisture of around two years (Chikamoto et al., 310 2017). Temperature gains most of its predictability through sea surface temperature (SST) forcing in the equatorial regions (Feng et al., 2011), and land-atmosphere coupling in the semi-arid tropics (Seo et al., 2019).

The overall pf of CESM2, CMCC-CM2-SR5 and IPSL-CM6A-LR falls into a narrow window of 0.19 to 0.21. CESM1-CAM5 has the highest pf with a value of 0.24. It is likely that this increased share of predictable IAV is not due to differences in model structure, but due to the large number of 40 ensemble members. Most other ESMs in this study have only ten 315 ensemble members, which is not enough to capture the difference between ensemble variance and IAV, so that an increase in ensemble members leads to an increase in prediction skill (Meehl et al., 2021). However, despite having 20 ensemble members, CanESM5 has the lowest pf among the models. A possible explanation could be the low IAV of deep layer soil moisture in CanESM5 (Qiao et al., 2022). A limited ability to reproduce the full spectrum of soil moisture variability could mean that

the soils have a smaller buffering capacity. As a result, they are not able to simulate the observed persistence of soil moisture anomalies, leading to a reduction of predictability. On the other hand, a high variability of soil moisture does not guarantee a high pf , as seen by the example of MPI-ESM-LR. The low pf of MPI-ESM-LR can be explained by the sensitivity of GPP to radiation. Since only 7% to 12% of the radiation-induced IAV is predictable, a high share of $\sigma\text{GPP}^{\text{Radiation}}$ reduces the predictability of GPP. This becomes evident in MPI-ESM-LR, in which the share of $\sigma\text{GPP}^{\text{Radiation}}$ is 20% higher than in the other ESMs.

We find that the regions contributing to the predictability of atmospheric CO_2 (pc) are highly related to the IAV patterns. The spatial correlation between pc and IAV exceeds 0.79 in all models but CanESM5. Indeed, these high correlations between predictability and IAV align with our understanding. Under a constant pf , pc would grow linearly with increasing IAV, leading to a perfect correlation. These high correlations show that the differences in the predictability of atmospheric CO_2 are determined more by the differences in GPP IAV than the differences in the pf of GPP. While the pf values show that the ESMs have a similar degree of memory retention, there are few overlaps in the spatial distribution of the pc , with an average correlation of 0.38 between the ESMs. For an alternative quantification of this disagreement, we separate the high-predictability grid cells, which are the grid cells contributing to the top 20th quantile of pc . 74% of these high-predictability grid cells are unique to only one ESM, and only 8% of high-predictability grid cells can be found in three or more ESMs.

Although the spatial patterns of the pc broadly resemble the patterns of GPP IAV, there are some slight differences between these fields. The pc is relatively high along the northeastern coast of South America in most ESMs. This could be due to the high climate predictability caused by slowly evolving Atlantic SST patterns (Dirmeyer et al., 2018). Other systematic differences can be explained by the differing pc of the environmental drivers. The most evident is the difference between IAV and predictability in regions where GPP IAV is driven by radiation. This leads to relatively low predictability in the tropical rainforests of the western Amazon basin and the Congo basin. An exception is the predictability provided through radiation on the Southeast Asian islands in IPSL-CM6A-LR and CESM1-CAM5. High predictability in these regions could be explained by the proximity to the ENSO SST region. Strong and predictable SST anomalies in the tropical Pacific that surround the islands can directly influence the cloud cover over land. The predictable component is also higher over areas where IAV is driven by soil moisture rather than temperature. In many ESMs, this leads to a high predictable component in the semiarid regions of South America, Africa and India.

4 Conclusions

We tested the ability of six ESMs to predict terrestrial GPP and determined their similarities and the sources of uncertainties. The ESMs are fairly similar in their ability to retain memory in hindcast simulations and predict their own variability, with the pf values of four of the ESMs falling between 19% and 24%. Most of the GPP pf is provided by soil moisture. Up to 32% of the GPP IAV caused by soil moisture is predictable, while it is only 7% to 12% for the IAV caused by radiation. The differences in the pf among ESMs are due to ensemble size and the sensitivity of GPP to radiation. Further sources of predictability which is not studied here are long-term vegetation dynamics. Specifically, the large and structural changes like

tree mortality (Wigneron et al., 2020) and recruitment (Holmgren et al., 2001). These processes only occur in extreme years and cause shifts in ecosystem states with long-lasting effects. The correct representation of these processes in ESMs allows them to reproduce the low-frequency IAV of vegetation, thereby extending the *pf* of GPP.

355 Although ESMs are similar in the fraction of GPP IAV they can predict, there are substantial differences in the patterns and drivers of GPP IAV. The ESMs have distinct, non-overlapping hotspots of GPP IAV that drive the variability of atmospheric CO₂. We find large disparities in the role of Australia, Southern Africa, and Central South America on GPP IAV. The leading cause of the uncertainties in IAV patterns are differences in the response of GPP to soil moisture and the capability of the ESMs to simulate soil hydrology accurately. These differences materialize through the direct effect of soil moisture on photosynthesis, and through the role of soil moisture on phenology. The inability of ESMs to reproduce GPP IAV also means, that there are regions where the potential predictability of GPP does not resemble the actual predictive skill.

This study shows, that the predictability of atmospheric CO₂ is currently not limited by the processes that provide predictability in the earth system, but rather by the representation of carbon flux variability patterns. The mismatches in GPP IAV imply, that the IAV of atmospheric CO₂ are caused by different regions and by different drivers across the ESMs. Consequentially, when ESMs predict the atmospheric CO₂, the GPP anomalies that constitute the predicted CO₂ growth rate originate from different regions. Because the predicted CO₂ depends more on the distribution of GPP IAV hotspots than actual mechanisms that provide predictability, CO₂ forecast skill is not a suitable metric for studies on carbon flux predictability. An ESM with a high carbon flux predictability can be outperformed in CO₂ forecast skill by a model that has a better representation of IAV patterns. A more suitable measure to assess carbon flux predictability could be the globally averaged anomaly correlation coefficient.

370 With the current uncertainties in GPP IAV patterns, the prediction of atmospheric CO₂ relies less on the prediction of regional climate anomalies, and more on the predictable global climate patterns like ENSO. These global climate anomalies are able to balance out the regional differences in GPP IAV patterns, by affecting large parts of the land surface simultaneously. In order to utilize the benefits of regional climate predictability for the predictability of CO₂, further work ought to focus on constraining GPP IAV, and not on the processes providing predictability. The most limiting aspect in the use of ESMs to predict atmospheric CO₂ is a better understanding of the drivers of carbon flux variability. Whether GPP is limited by moisture, temperature or radiation does not only affect variability patterns but also the predictability of the fluxes. An overestimation of humidity in an ecosystem by an ESM would result in GPP being more controlled by radiation than soil moisture, leading to an underestimation of predictability, or vice-versa for systems that are too dry.

380 The findings of this study also suggest that previous estimations of ESM-based CO₂ forecast skill are underestimating the predictive capabilities of these systems. Various post-processing strategies could help to produce a CO₂ forecast skill that is not obscured by inaccurate IAV patterns but is a closer representation of the actual performance of ESM-based prediction systems. These strategies could include the scaling of carbon flux IAV patterns to resemble the observed IAV patterns. Since there are strong regional differences in the predictive performance among the ESMs, another strategy would be to combine ESM predictions in a way that utilizes these differences. This could be done in a regionally weighted multi-model approach.

The limiting factor to predicting atmospheric CO₂ is the chaotic nature of weather and climate. However, our results show that we have not reached this limitation yet and that we are instead constrained by our understanding of terrestrial carbon flux variability. The development of observational products for terrestrial carbon fluxes, especially in the tropics, remains the main objective on the path of improving the predictability of the global carbon cycle and atmospheric CO₂.

390 *Code and data availability.* [TBD]

Author contributions. The Study was conceptualised by ID, TI, and VB, ID developed the methodology, ran the analysis, and wrote the original draft, NL, AC, VA, TI and VB reviewed and edited the manuscript, and VB and TI supervised.

Competing interests. The authors declare that they have no conflict of interest.

395 *Acknowledgements.* We acknowledge funding from European Union's Horizon 2020 research and innovation programme under Grant Agreement No. 821003 'Climate-Carbon Interactions in the Coming Century CCiCC (4C)'. NL is grateful for funding from the US National Science Foundation (OCE-1752724). The authors wish to thank the modelling groups who have participated in the DCP, SMYLE and MiKlip projects. We also thank Jürgen Bader for reviewing the manuscript.

References

- Ahlström, A., Raupach, M. R., Schurgers, G., Smith, B., Arneth, A., Jung, M., Reichstein, M., Canadell, J. G., Friedlingstein, P., Jain, A. K.,
400 Kato, E., Poulter, B., Sitch, S., Stocker, B. D., Viovy, N., Wang, Y. P., Wiltshire, A., Zaehle, S., and Zeng, N.: The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink, *Science*, <https://doi.org/10.1126/science.aaa1668>, 2015.
- Alessandri, A. and Navarra, A.: On the coupling between vegetation and rainfall inter-annual anomalies: Possible contributions to seasonal rainfall predictability over land areas, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2007GL032415>, 2008.
- Anav, A., Friedlingstein, P., Beer, C., Ciais, P., Harper, A., Jones, C., Murray-Tortarolo, G., Papale, D., Parazoo, N. C., Peylin, P., Piao, S.,
405 Sitch, S., Viovy, N., Wiltshire, A., and Zhao, M.: Spatiotemporal patterns of terrestrial gross primary production: A review, *Reviews of Geophysics*, 53, 785–818, <https://doi.org/10.1002/2015RG000483>, 2015.
- Balmaseda, M. A., Mogensen, K., and Weaver, A. T.: Evaluation of the ECMWF ocean reanalysis system ORAS4, *Quarterly Journal of the Royal Meteorological Society*, 139, 1132–1161, <https://doi.org/10.1002/qj.2063>, 2013.
- Banzon, V., Smith, T. M., Chin, T. M., Liu, C., and Hankins, W.: A long-term record of blended satellite and in situ sea-surface temperature
410 for climate monitoring, modeling and environmental studies, *Earth System Science Data*, 8, 165–176, <https://doi.org/10.5194/essd-8-165-2016>, 2016.
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luysaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal, E., Viovy, N., Williams, C., Woodward, F. I., and Papale, D.: Terrestrial Gross Carbon Dioxide Uptake: Global Distribution and Covariation
415 with Climate, *Science*, <https://doi.org/10.1126/science.1184984>, 2010.
- Bellucci, A., Haarsma, R., Bellouin, N., Booth, B., Cagnazzo, C., Hurk, B. v. d., Keenlyside, N., Koenigk, T., Massonnet, F., Materia, S., and Weiss, M.: Advancements in decadal climate predictability: The role of nonoceanic drivers, *Reviews of Geophysics*, 53, 165–202, <https://doi.org/10.1002/2014RG000473>, 2015.
- Betts, R. A., Jones, C. D., Knight, J. R., Keeling, R. F., and Kennedy, J. J.: El Niño and a record CO₂ rise, *Nature Climate Change*, 6,
420 806–810, <https://doi.org/10.1038/nclimate3063>, 2016.
- Boer, G. J., Smith, D. M., Cassou, C., Doblas-Reyes, F., Danabasoglu, G., Kirtman, B., Kushnir, Y., Kimoto, M., Meehl, G. A., Msadek, R., Mueller, W. A., Taylor, K. E., Zwiers, F., Rixen, M., Ruprich-Robert, Y., and Eade, R.: The Decadal Climate Prediction Project (DCPP) contribution to CMIP6, *Geoscientific Model Development*, 9, 3751–3777, <https://doi.org/10.5194/gmd-9-3751-2016>, 2016.
- Bonan, G. B., Lombardozzi, D. L., Wieder, W. R., Oleson, K. W., Lawrence, D. M., Hoffman, F. M., and Collier, N.: Model Structure and
425 Climate Data Uncertainty in Historical Simulations of the Terrestrial Carbon Cycle (1850–2014), *Global Biogeochemical Cycles*, 33, 1310–1326, <https://doi.org/10.1029/2019GB006175>, 2019.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., Lavergne, C. d., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-
430 A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, E., L., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levvasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luysaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Ottlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont,

- R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR
435 Climate Model, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002 010, <https://doi.org/10.1029/2019MS002010>, 2020.
- Chen, M., Rafique, R., Asrar, G. R., Bond-Lamberty, B., Ciais, P., Zhao, F., Reyer, C. P. O., Ostberg, S., Chang, J., Ito, A., Yang, J., Zeng, N.,
Kalnay, E., West, T., Leng, G., Francois, L., Munhoven, G., Henrot, A., Tian, H., Pan, S., Nishina, K., Viovy, N., Morfopoulos, C., Betts,
R., Schaphoff, S., Steinkamp, J., and Hickler, T.: Regional contribution to variability and trends of global gross primary productivity,
Environmental Research Letters, 12, 105 005, <https://doi.org/10.1088/1748-9326/aa8978>, 2017.
- 440 Cherchi, A., Fogli, P. G., Lovato, T., Peano, D., Iovino, D., Gualdi, S., Masina, S., Scoccimarro, E., Materia, S., Bellucci, A., and Navarra,
A.: Global Mean Climate and Main Patterns of Variability in the CMCC-CM2 Coupled Model, *Journal of Advances in Modeling Earth
Systems*, 11, 185–209, <https://doi.org/10.1029/2018MS001369>, 2019.
- Cheruy, F., Ducharne, A., Hourdin, F., Musat, I., Vignon, , Gastineau, G., Bastrikov, V., Vuichard, N., Diallo, B., Dufresne, J.-L., Ghattas, J.,
Grandpeix, J.-Y., Idelkadi, A., Mellul, L., Maignan, F., Ménégoz, M., Ottlé, C., Peylin, P., Servonnat, J., Wang, F., and Zhao, Y.: Improved
445 Near-Surface Continental Climate in IPSL-CM6A-LR by Combined Evolutions of Atmospheric and Land Surface Physics, *Journal of
Advances in Modeling Earth Systems*, 12, e2019MS002 005, <https://doi.org/10.1029/2019MS002005>, 2020.
- Chikamoto, Y., Timmermann, A., Widlansky, M. J., Balmaseda, M. A., and Stott, L.: Multi-year predictability of climate, drought, and
wildfire in southwestern North America, *Scientific Reports*, 7, 6568, <https://doi.org/10.1038/s41598-017-06869-7>, 2017.
- Collalti, A., Ibrom, A., Stockmarr, A., Cescatti, A., Alkama, R., Fernández-Martínez, M., Matteucci, G., Sitch, S., Friedlingstein, P., Ciais,
450 P., Goll, D. S., Nabel, J. E. M. S., Pongratz, J., Arneth, A., Haverd, V., and Prentice, I. C.: Forest production efficiency increases with
growth temperature, *Nature Communications*, 11, 5322, <https://doi.org/10.1038/s41467-020-19187-w>, 2020.
- Collier, N., Hoffman, F. M., Lawrence, D. M., Keppel-Aleks, G., Koven, C. D., Riley, W. J., Mu, M., and Randerson, J. T.: The International
Land Model Benchmarking (ILAMB) System: Design, Theory, and Implementation, *Journal of Advances in Modeling Earth Systems*, 10,
2731–2754, <https://doi.org/10.1029/2018MS001354>, 2018.
- 455 Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R.,
Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb,
W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., Kampenhout, L. v., Vertenstein,
M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C.,
Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2
460 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001 916, <https://doi.org/10.1029/2019MS001916>, 2020.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer,
P., Bechtold, P., Beljaars, A. C. M., Berg, L. v. d., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haim-
berger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz,
B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., Rosnay, P. d., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis:
465 configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597,
<https://doi.org/10.1002/qj.828>, 2011.
- Dirkson, A., Merryfield, W. J., and Monahan, A.: Impacts of Sea Ice Thickness Initialization on Seasonal Arctic Sea Ice Predictions, *Journal
of Climate*, 30, 1001–1017, <https://doi.org/10.1175/JCLI-D-16-0437.1>, 2017.
- Dirmeyer, P. A., Halder, S., and Bombardi, R.: On the Harvest of Predictability From Land States in a Global Forecast Model, *Journal of
470 Geophysical Research: Atmospheres*, 123, 13,111–13,127, <https://doi.org/10.1029/2018JD029103>, 2018.

- Dunkl, I., Spring, A., Friedlingstein, P., and Brovkin, V.: Process-based analysis of terrestrial carbon flux predictability, *Earth System Dynamics*, 12, 1413–1426, <https://doi.org/10.5194/esd-12-1413-2021>, 2021.
- Estella-Perez, V., Mignot, J., Guilyardi, E., Swingedouw, D., and Reverdin, G.: Advances in reconstructing the AMOC using sea surface observations of salinity, *Climate Dynamics*, 55, 975–992, <https://doi.org/10.1007/s00382-020-05304-4>, 2020.
- 475 Feng, X., DelSole, T., and Houser, P.: Bootstrap estimated seasonal potential predictability of global temperature and precipitation, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2010GL046511>, 2011.
- Gavrilov, M. B., An, W., Xu, C., Radaković, M. G., Hao, Q., Yang, F., Guo, Z., Perić, Z., Gavrilov, G., and Marković, S. B.: Independent Aridity and Drought Pieces of Evidence Based on Meteorological Data and Tree Ring Data in Southeast Banat, Vojvodina, Serbia, *Atmosphere*, 10, 586, <https://doi.org/10.3390/atmos10100586>, number: 10 Publisher: Multidisciplinary Digital Publishing Institute, 2019.
- 480 Giorgetta, M. A., Jungclauss, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *Journal of Advances in Modeling Earth Systems*, 5, 572–597, <https://doi.org/10.1002/jame.20038>, 2013.
- 485 Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *Journal of Geophysical Research: Oceans*, 118, 6704–6716, <https://doi.org/10.1002/2013JC009067>, 2013.
- Holmgren, M., Scheffer, M., Ezcurra, E., Gutiérrez, J. R., and Mohren, G. M. J.: El Niño effects on the dynamics of terrestrial ecosystems, *Trends in Ecology & Evolution*, 16, 89–94, [https://doi.org/10.1016/S0169-5347\(00\)02052-8](https://doi.org/10.1016/S0169-5347(00)02052-8), 2001.
- 490 Hu, Q., Li, T., Deng, X., Wu, T., Zhai, P., Huang, D., Fan, X., Zhu, Y., Lin, Y., Xiao, X., Chen, X., Zhao, X., Wang, L., and Qin, Z.: Intercomparison of global terrestrial carbon fluxes estimated by MODIS and Earth system models, *Science of The Total Environment*, 810, 152 231, <https://doi.org/10.1016/j.scitotenv.2021.152231>, 2022.
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J.-F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D.,
- 495 Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: A Framework for Collaborative Research, *Bulletin of the American Meteorological Society*, 94, 1339–1360, <https://doi.org/10.1175/BAMS-D-12-00121.1>, 2013.
- Ilyina, T., Li, H., Spring, A., Müller, W. A., Bopp, L., Chikamoto, M. O., Danabasoglu, G., Dobrynin, M., Dunne, J., Fransner, F., Friedlingstein, P., Lee, W., Lovenduski, N. S., Merryfield, W. J., Mignot, J., Park, J. Y., Séférian, R., Sospedra-Alfonso, R., Watanabe, M., and Yeager, S.: Predictable Variations of the Carbon Sinks and Atmospheric CO₂ Growth in a Multi-Model Framework, *Geophysical Research Letters*, 48, e2020GL090 695, <https://doi.org/10.1029/2020GL090695>, 2021.
- 500 Jung, M., Reichstein, M., Schwalm, C. R., Huntingford, C., Sitch, S., Ahlström, A., Arneth, A., Camps-Valls, G., Ciais, P., Friedlingstein, P., Gans, F., Ichii, K., Jain, A. K., Kato, E., Papale, D., Poulter, B., Raduly, B., Rödenbeck, C., Tramontana, G., Viovy, N., Wang, Y.-P., Weber, U., Zaehle, S., and Zeng, N.: Compensatory water effects link yearly global land CO₂ sink changes to temperature, *Nature*, 541, 516–520, <https://doi.org/10.1038/nature20780>, 2017.
- 505 Jung, M., Koirala, S., Weber, U., Ichii, K., Gans, F., Camps-Valls, G., Papale, D., Schwalm, C., Tramontana, G., and Reichstein, M.: The FLUXCOM ensemble of global land-atmosphere energy fluxes, *Scientific Data*, 6, 74, <https://doi.org/10.1038/s41597-019-0076-8>, 2019.
- Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S. C., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J.-F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L., and Vertenstein,

- M.: The Community Earth System Model (CESM) Large Ensemble Project: A Community Resource for Studying Climate Change in the
510 Presence of Internal Climate Variability, *Bulletin of the American Meteorological Society*, 96, 1333–1349, <https://doi.org/10.1175/BAMS-D-13-00255.1>, 2015.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 Reanalysis: General Specifications and Basic Characteristics, *Journal of the Meteorological Society of Japan*. Ser. II, 93, 5–48, <https://doi.org/10.2151/jmsj.2015-001>, 2015.
- 515 Kolby Smith, W., Reed, S. C., Cleveland, C. C., Ballantyne, A. P., Anderegg, W. R. L., Wieder, W. R., Liu, Y. Y., and Running, S. W.: Large divergence of satellite and Earth system model estimates of global terrestrial CO₂ fertilization, *Nature Climate Change*, 6, 306–310, <https://doi.org/10.1038/nclimate2879>, number: 3 Publisher: Nature Publishing Group, 2016.
- Kumar, S., Newman, M., Lawrence, D. M., Lo, M.-H., Akula, S., Lan, C.-W., Livneh, B., and Lombardozi, D.: The GLACE-Hydrology Experiment: Effects of Land–Atmosphere Coupling on Soil Moisture Variability and Predictability, *Journal of Climate*, 33, 6511–6529,
520 <https://doi.org/10.1175/JCLI-D-19-0598.1>, 2020.
- Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in Version 4 of the Community Land Model, *Journal of Advances in Modeling Earth Systems*, 3, <https://doi.org/10.1029/2011MS00045>, 2011.
- Lawrence, D. M., Fisher, R. A., Koven, C. D., Oleson, K. W., Swenson, S. C., Bonan, G., Collier, N., Ghimire, B., Kampenhout, L. v.,
525 Kennedy, D., Kluzek, E., Lawrence, P. J., Li, F., Li, H., Lombardozi, D., Riley, W. J., Sacks, W. J., Shi, M., Vertenstein, M., Wieder, W. R., Xu, C., Ali, A. A., Badger, A. M., Bisht, G., Broeke, M. v. d., Brunke, M. A., Burns, S. P., Buzan, J., Clark, M., Craig, A., Dahlin, K., Drewniak, B., Fisher, J. B., Flanner, M., Fox, A. M., Gentine, P., Hoffman, F., Keppel-Aleks, G., Knox, R., Kumar, S., Lenaerts, J., Leung, L. R., Lipscomb, W. H., Lu, Y., Pandey, A., Pelletier, J. D., Perket, J., Randerson, J. T., Ricciuto, D. M., Sanderson, B. M., Slater, A., Subin, Z. M., Tang, J., Thomas, R. Q., Martin, M. V., and Zeng, X.: The Community Land Model Version 5: Description
530 of New Features, Benchmarking, and Impact of Forcing Uncertainty, *Journal of Advances in Modeling Earth Systems*, 11, 4245–4287, <https://doi.org/10.1029/2018MS001583>, 2019.
- Li, X. and Xiao, J.: Mapping Photosynthesis Solely from Solar-Induced Chlorophyll Fluorescence: A Global, Fine-Resolution Dataset of Gross Primary Production Derived from OCO-2, *Remote Sensing*, 11, <https://doi.org/10.3390/rs11212563>, 2019.
- Lovato, T., Peano, D., Butenschön, M., Matera, S., Iovino, D., Scoccimarro, E., Fogli, P. G., Cherchi, A., Bellucci, A., Gualdi, S., Masina,
535 S., and Navarra, A.: CMIP6 Simulations With the CMCC Earth System Model (CMCC-ESM2), *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002814, <https://doi.org/10.1029/2021MS002814>, 2022.
- Lovenduski, N. S., Bonan, G. B., Yeager, S. G., Lindsay, K., and Lombardozi, D. L.: High predictability of terrestrial carbon fluxes from an initialized decadal prediction system, *Environmental Research Letters*, 14, 124074, <https://doi.org/10.1088/1748-9326/ab5c55>, 2019.
- Marotzke, J., Müller, W. A., Vamborg, F. S. E., Becker, P., Cubasch, U., Feldmann, H., Kaspar, F., Kottmeier, C., Marini, C., Polkova, I.,
540 Prömmel, K., Rust, H. W., Stammer, D., Ulbrich, U., Kadow, C., Köhl, A., Kröger, J., Kruschke, T., Pinto, J. G., Pohlmann, H., Reyers, M., Schröder, M., Sienz, F., Timmreck, C., and Ziese, M.: MiKlip: A National Research Project on Decadal Climate Prediction, *Bulletin of the American Meteorological Society*, 97, 2379–2394, <https://doi.org/10.1175/BAMS-D-15-00184.1>, 2016.
- Meehl, G. A., Richter, J. H., Teng, H., Capotondi, A., Cobb, K., Doblas-Reyes, F., Donat, M. G., England, M. H., Fyfe, J. C., Han, W., Kim, H., Kirtman, B. P., Kushnir, Y., Lovenduski, N. S., Mann, M. E., Merryfield, W. J., Nieves, V., Pegion, K., Rosenbloom, N., Sanchez,
545 S. C., Scaife, A. A., Smith, D., Subramanian, A. C., Sun, L., Thompson, D., Ummenhofer, C. C., and Xie, S.-P.: Initialized Earth System

- prediction from subseasonal to decadal timescales, *Nature Reviews Earth & Environment*, 2, 340–357, <https://doi.org/10.1038/s43017-021-00155-x>, 2021.
- Muggeo, V.: Segmented: An R Package to Fit Regression Models With Broken-Line Relationships, *R News*, 8, 20–25, 2008.
- 550 Nicolì, D., Bellucci, A., Ruggieri, P., Athanasiadis, P., Materia, S., Peano, D., Fedele, G., and Gualdi, S.: The CMCC Decadal Prediction System, preprint, *Climate and Earth system modeling*, <https://doi.org/10.5194/gmd-2022-181>, 2022.
- O’Sullivan, M., Smith, W. K., Sitch, S., Friedlingstein, P., Arora, V. K., Haverd, V., Jain, A. K., Kato, E., Kautz, M., Lombardozzi, D., Nabel, J. E. M. S., Tian, H., Vuichard, N., Wiltshire, A., Zhu, D., and Buermann, W.: Climate-Driven Variability and Trends in Plant Productivity Over Recent Decades Based on Three Global Products, *Global Biogeochemical Cycles*, 34, e2020GB006613, <https://doi.org/10.1029/2020GB006613>, 2020.
- 555 Padrón, R. S., Gudmundsson, L., Liu, L., Humphrey, V., and Seneviratne, S. I.: Controls of intermodel uncertainty in land carbon sink projections, *Biogeosciences Discussions*, pp. 1–20, <https://doi.org/10.5194/bg-2022-92>, 2022.
- Peano, D., Materia, S., Collalti, A., Alessandri, A., Anav, A., Bombelli, A., and Gualdi, S.: Global Variability of Simulated and Observed Vegetation Growing Season, *Journal of Geophysical Research: Biogeosciences*, 124, 3569–3587, <https://doi.org/10.1029/2018JG004881>, 2019.
- 560 Peano, D., Hemming, D., Materia, S., Delire, C., Fan, Y., Joetzjer, E., Lee, H., Nabel, J. E. M. S., Park, T., Peylin, P., Wårlind, D., Wiltshire, A., and Zaehle, S.: Plant phenology evaluation of CRESCENDO land surface models – Part 1: Start and end of the growing season, *Biogeosciences*, 18, 2405–2428, <https://doi.org/10.5194/bg-18-2405-2021>, 2021.
- Piao, S., Wang, X., Wang, K., Li, X., Bastos, A., Canadell, J. G., Ciais, P., Friedlingstein, P., and Sitch, S.: Interannual variation of terrestrial carbon cycle: Issues and perspectives, *Global Change Biology*, 26, 300–318, <https://doi.org/10.1111/gcb.14884>, 2020.
- 565 Qiao, L., Zuo, Z., and Xiao, D.: Evaluation of Soil Moisture in CMIP6 Simulations, *Journal of Climate*, 35, 779–800, <https://doi.org/10.1175/JCLI-D-20-0827.1>, 2022.
- Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *Journal of Advances in Modeling Earth Systems*, 5, 459–482, <https://doi.org/10.1002/jame.20022>, 2013.
- Santanello, J. A., Dirmeyer, P. A., Ferguson, C. R., Findell, K. L., Tawfik, A. B., Berg, A., Ek, M., Gentine, P., Guillod, B. P., Heerwaarden, C. v., Roundy, J., and Wulfmeyer, V.: Land–Atmosphere Interactions: The LoCo Perspective, *Bulletin of the American Meteorological Society*, 99, 1253–1272, <https://doi.org/10.1175/BAMS-D-17-0001.1>, 2018.
- 570 Seo, E., Lee, M.-I., Jeong, J.-H., Koster, R. D., Schubert, S. D., Kim, H.-M., Kim, D., Kang, H.-S., Kim, H.-K., MacLachlan, C., and Scaife, A. A.: Impact of soil moisture initialization on boreal summer subseasonal forecasts: mid-latitude surface air temperature and heat wave events, *Climate Dynamics*, 52, 1695–1709, <https://doi.org/10.1007/s00382-018-4221-4>, 2019.
- 575 Smith, T. M., Reynolds, R. W., Peterson, T. C., and Lawrimore, J.: Improvements to NOAA’s Historical Merged Land–Ocean Surface Temperature Analysis (1880–2006), *Journal of Climate*, 21, 2283–2296, <https://doi.org/10.1175/2007JCLI2100.1>, 2008.
- Song, X., Wang, D.-Y., Li, F., and Zeng, X.-D.: Evaluating the performance of CMIP6 Earth system models in simulating global vegetation structure and distribution, *Advances in Climate Change Research*, 12, 584–595, <https://doi.org/10.1016/j.accr.2021.06.008>, 2021.
- 580 Spring, A. and Ilyina, T.: Predictability Horizons in the Global Carbon Cycle Inferred From a Perfect-Model Framework, *Geophysical Research Letters*, 47, e2019GL085311, <https://doi.org/10.1029/2019GL085311>, 2020.
- Storto, A. and Masina, S.: C-GLORSv5: an improved multipurpose global ocean eddy-permitting physical reanalysis, *Earth System Science Data*, 8, 679–696, <https://doi.org/10.5194/essd-8-679-2016>, 2016.

- Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Sigmond, M., Solheim, L., von Salzen, K., Yang, D., and Winter, B.: The Canadian Earth System Model version 5 (CanESM5.0.3), *Geoscientific Model Development*, 12, 4823–4873, <https://doi.org/10.5194/gmd-12-4823-2019>, 2019.
- Séférian, R., Berthet, S., and Chevallier, M.: Assessing the Decadal Predictability of Land and Ocean Carbon Uptake, *Geophysical Research Letters*, 45, 2455–2466, <https://doi.org/10.1002/2017GL076092>, 2018.
- Titchner, H. A. and Rayner, N. A.: The Met Office Hadley Centre sea ice and sea surface temperature data set, version 2: 1. Sea ice concentrations, *Journal of Geophysical Research: Atmospheres*, 119, 2864–2889, <https://doi.org/10.1002/2013JD020316>, 2014.
- Tsujino, H., Urakawa, S., Nakano, H., Small, R. J., Kim, W. M., Yeager, S. G., Danabasoglu, G., Suzuki, T., Bamber, J. L., Bentsen, M., Böning, C. W., Bozec, A., Chassignet, E. P., Curchitser, E., Boeira Dias, F., Durack, P. J., Griffies, S. M., Harada, Y., Ilıcak, M., Josey, S. A., Kobayashi, C., Kobayashi, S., Komuro, Y., Large, W. G., Le Sommer, J., Marsland, S. J., Masina, S., Scheinert, M., Tomita, H., Valdivieso, M., and Yamazaki, D.: JRA-55 based surface dataset for driving ocean–sea-ice models (JRA55-do), *Ocean Modelling*, 130, 79–139, <https://doi.org/10.1016/j.ocemod.2018.07.002>, 2018.
- Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D. C., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Berg, L. V. D., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, L., Janssen, P. a. E. M., Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Quarterly Journal of the Royal Meteorological Society*, 131, 2961–3012, <https://doi.org/10.1256/qj.04.176>, 2005.
- Wang, G., Sun, S., and Mei, R.: Vegetation dynamics contributes to the multi-decadal variability of precipitation in the Amazon region, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL049017>, 2011.
- Wang, J., Zeng, N., and Wang, M.: Interannual variability of the atmospheric CO₂ growth rate: roles of precipitation and temperature, *Biogeosciences*, 13, 2339–2352, <https://doi.org/10.5194/bg-13-2339-2016>, 2016.
- Wigner, J.-P., Fan, L., Ciais, P., Bastos, A., Brandt, M., Chave, J., Saatchi, S., Baccini, A., and Fensholt, R.: Tropical forests did not recover from the strong 2015–2016 El Niño event, *Science Advances*, <https://doi.org/10.1126/sciadv.aay4603>, 2020.
- Wu, R.-J., Lo, M.-H., and Scanlon, B. R.: The Annual Cycle of Terrestrial Water Storage Anomalies in CMIP6 Models Evaluated against GRACE Data, *Journal of Climate*, 34, 8205–8217, <https://doi.org/10.1175/JCLI-D-21-0021.1>, 2021.
- Xue, Y., Smith, T. M., and Reynolds, R. W.: Interdecadal Changes of 30-Yr SST Normals during 1871–2000, *Journal of Climate*, 16, 1601–1612, [https://doi.org/10.1175/1520-0442\(2003\)016<1601:ICOYSN>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<1601:ICOYSN>2.0.CO;2), 2003.
- Yang, C., Masina, S., and Storto, A.: Historical ocean reanalyses (1900–2010) using different data assimilation strategies, *Quarterly Journal of the Royal Meteorological Society*, 143, 479–493, <https://doi.org/10.1002/qj.2936>, 2017.
- Yeager, S. G., Danabasoglu, G., Rosenbloom, N. A., Strand, W., Bates, S. C., Meehl, G. A., Karspeck, A. R., Lindsay, K., Long, M. C., Teng, H., and Lovenduski, N. S.: Predicting Near-Term Changes in the Earth System: A Large Ensemble of Initialized Decadal Prediction Simulations Using the Community Earth System Model, *Bulletin of the American Meteorological Society*, 99, 1867–1886, <https://doi.org/10.1175/BAMS-D-17-0098.1>, 2018.
- Yeager, S. G., Rosenbloom, N., Glanville, A. A., Wu, X., Simpson, I., Li, H., Molina, M. J., Krumhardt, K., Mogen, S., Lindsay, K., Lombardozzi, D., Wieder, W., Kim, W. M., Richter, J. H., Long, M., Danabasoglu, G., Bailey, D., Holland, M., Lovenduski, N., and

- 620 Strand, W. G.: The Seasonal-to-Multiyear Large Ensemble (SMYLE) Prediction System using the Community Earth System Model Version 2, *Geoscientific Model Development Discussions*, pp. 1–42, <https://doi.org/10.5194/gmd-2022-60>, 2022.
- Zeng, N., Yoon, J.-H., Vintzileos, A., Collatz, G. J., Kalnay, E., Mariotti, A., Kumar, A., Busalacchi, A., and Lord, S.: Dynamical prediction of terrestrial ecosystems and the global carbon cycle: A 25-year hindcast experiment, *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2008GB003183>, 2008.
- 625 Zhang, Y. and Ye, A.: Would the obtainable gross primary productivity (GPP) products stand up? A critical assessment of 45 global GPP products, *Science of The Total Environment*, 783, 146 965, <https://doi.org/10.1016/j.scitotenv.2021.146965>, 2021.
- Zhang, Y., Xiao, X., Wu, X., Zhou, S., Zhang, G., Qin, Y., and Dong, J.: A global moderate resolution dataset of gross primary production of vegetation for 2000–2016, *Scientific Data*, 4, 170 165, <https://doi.org/10.1038/sdata.2017.165>, 2017.
- Zhang, Y., Dannenberg, M. P., Hwang, T., and Song, C.: El Niño-Southern Oscillation-Induced Variability of Terrestrial
630 Gross Primary Production During the Satellite Era, *Journal of Geophysical Research: Biogeosciences*, 124, 2419–2431, <https://doi.org/10.1029/2019JG005117>, 2019.
- Zhang, Y., Keenan, T. F., and Zhou, S.: Exacerbated drought impacts on global ecosystems due to structural overshoot, *Nature Ecology & Evolution*, 5, 1490–1498, <https://doi.org/10.1038/s41559-021-01551-8>, 2021.
- Zuo, H., Balmaseda, M. A., Tietsche, S., Mogensen, K., and Mayer, M.: The ECMWF operational ensemble reanalysis–analysis system
635 for ocean and sea ice: a description of the system and assessment, *Ocean Science*, 15, 779–808, <https://doi.org/10.5194/os-15-779-2019>, 2019.