

# ESD Ideas: Translating historical extreme weather events into a warmer world

Ed Hawkins<sup>1</sup>, Gilbert P. Compo<sup>2,3</sup> and Prashant D. Sardeshmukh<sup>2,3</sup>

5 <sup>1</sup> National Centre for Atmospheric Science, Department of Meteorology, University of Reading, Reading, UK

<sup>2</sup> Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, Boulder, USA

<sup>3</sup> NOAA Physical Sciences Laboratory, Boulder, USA

*Correspondence to:* Ed Hawkins (ed.hawkins@ncas.ac.uk)

**Abstract.** A new reanalysis-based approach is proposed to examine how [reconstructions of](#) extreme weather events differ in  
10 [a warmer or cooler counter-factual world](#)s. This approach offers a novel way to develop plausible storylines for some types of  
extreme event that other methods may not be suitable for. As [a proof-of-concept](#), a reanalysis of a severe windstorm that  
occurred in February 1903 is translated into a warmer world where it produces higher wind speeds and increased rainfall,  
suggesting that this storm would be more damaging if it occurred today rather than 120 years ago.

Whenever a severe weather event occurs [and causes with](#) harmful impacts [to in](#) a particular region, [one question that it is often](#)  
15 asked by disaster responders, recovery planners, politicians and journalists [is](#) whether climate change [has](#) caused or affected  
the event. The [cause of the](#) harmful impacts [are caused by](#)is the unusual weather, but climate change may have made the  
weather event more likely, [more more](#)-severe, or [potentially](#) both. In those cases, the harmful impacts may be partly or even  
mostly due to the change in climate. In some cases, the worst consequences may be due to the vulnerability or exposure of  
[those living in the region](#)[the local population or ecosystems](#), or a combination of many [different other](#) factors (e.g. Otto et al.,  
20 2022).

Many methodologies exist to understand how climate change has affected extreme events. These are broadly categorized into  
risk-based approaches and storyline approaches (Stott and Christidis, 2023). The risk-based approaches [examine assess](#) the  
change in likelihood and magnitude of a particular class of event (e.g. Stott et al., 2004), whereas storyline approaches consider  
25 how climate change may have affected a specific event (Trenberth et al., 2015; Shepherd et al., 2018). Other studies consider  
the related question of what a plausible worst-case event might look like in a particular climate (e.g. Thompson et al., 2017).  
Some of these methods are now regularly used to provide attribution statements soon after events occur (van Oldenborgh et  
al., 2021).

Event storyline approaches attempt to quantify how an extreme event would be different in an altered climate. This can be  
achieved by producing reconstructions of the event as it occurred and in [a counter-factual cooler or warmer climate](#)s [to and](#)  
30 [comparing](#) the consequences. Various approaches exist for such analyses, including statistical methods (e.g. Cattiaux et al.,  
2010), [analogues](#) (Ginesta et al., 2023; Faranda et al., 2022), [nudging](#) a climate model (e.g. van Garderen et al., 2021; Sánchez-  
Benítez et al., 2022) and forecast-based (e.g. Wehner et al., 2019; Leach et al., 2021). Here, we propose a complementary

reanalysis-based approach ~~to developing event storylines translate extreme events into different climates using a proof-of-concept analysis of a severe extra-tropical windstorm; this type of extreme event has not been examined using other methods.~~

35 Specifically, we use the NOAA-CIRES-DOE 20th Century Reanalysis v3 system (20CRv3; Slivinski et al., 2021) which assimilates surface pressure observations into a NOAA weather forecast model to reconstruct the atmospheric circulation from 1806-2015 at 0.7° resolution using 80 ensemble members. Boundary conditions of SSTs and sea ice concentration are prescribed in 20CRv3. ~~We can generate, and so counter-factual versions-reconstructions of extreme events can be generated by perturbing the SST boundary conditions and rerunning the reanalysis for those events, assimilating the same surface~~  
40 ~~pressure observations. By using 20CRv3 no other perturbations are required, unlike if this approach used a reanalysis that assimilated other types of observation. As atmospheric temperature changes will be primarily mediated by imposed changes to SSTs (rather than by direct radiative forcing; Compo and Sardeshmukh, 2009), this perturbation translates the reanalysis of the events into a changed climate. In the experiments described here, sea ice and atmospheric greenhouse gas concentrations remain fixed, and the SST perturbation is spatially uniform. A ‘spin-up’ period needs to be run prior to the extreme event to~~  
45 ~~ensure that the atmosphere has adjusted to the counter-factual SSTs; in this example a period of 6 months was used. Sensitivity experiments are planned to explore all these issues.~~

As proof-of-concept, ~~experiment we~~ ~~has~~ ~~applied~~ this approach to a severe extra-tropical windstorm known as Storm Ulysses, which occurred in February 1903, to examine how the consequences of this historical event may have been different had it occurred in a warmer world. Three ~~versions-experiments withof~~ the ~~same~~ reanalysis ~~system~~ are considered: (1) the original  
50 20CRv3, (2) an improved version of 20CRv3 with added historical surface pressure observations and a small change to the data assimilation scheme (Hawkins et al. 2022), and (3) same as (2), but with a ~~spatially uniform uniform~~+2K perturbation added to the sea surface temperature (SST) boundary conditions ~~and assimilating the same surface pressure observations. Sea ice, atmospheric greenhouse gas and aerosol concentrations are unchanged. A 6 month ‘spin-up’ period was run prior to the~~  
55 ~~the sensitivity of the findings to all these different choices. This perturbation plausibly translates the reanalysis of Storm Ulysses into a warmer world.~~

Storm Ulysses is not well represented in the original 20CRv3, but the improved reanalysis shows a far more intense storm, mainly from adding new rescued pressure observations; it is a credible reconstruction of one of the most extreme windstorms to occur over the British & Irish Isles in the past 120 years (Hawkins et al., 2022). Figure 1 (top row) shows the wet-bulb  
60 potential temperature at 850 hPa for one specific time during the storm for the different experiments. The storm in the improved reanalysis has a hook-shaped Shapiro-Keyser-type structure which is not present in the original 20CRv3. As expected, the

~~storm~~ structure of the storm in the warmer world ~~experiment~~ reanalysis is constrained to be ~~looks~~ very similar to the improved reanalysis, except for a roughly 2.5K offset. Specific humidity and surface air temperatures are also higher (not shown).

65 The warmer world ~~version of the reanalysis of the~~ storm generates stronger maximum winds (middle row) and increased rainfall (bottom row) compared to the improved reanalysis, suggesting that the consequences of the same surface circulation pattern would be more severe in a warmer world. The reconstructed circulation has some limited flexibility to vary given the observational coverage and uncertainties. ~~Although~~ While the increase in wind footprint appears small over land (around 1 m/s in some places), ~~this implies it becomes~~ a significant increase in damage (>10%; roughly related to the cube of the wind speed; Klawa and Ulbrich, 2003). The total rainfall over land during the storm increases by 26%, i.e. around 10%/K when using the temperature change at 850 hPa to normalise the rainfall change, which is larger than might be expected due to the Clausius-Clapeyron relationship alone. An animation of the storm is included as a Supplementary Figure.

75 These ~~proof-of-concept~~ experiments have reconstructed an intense windstorm in different climates and demonstrated the potential of using a surface pressure based reanalysis to ~~explore~~ develop heavily-constrained storylines which examine the thermodynamic and local dynamical changes in extreme weather events ~~such as windstorms and potentially heavy rainfall events.~~ A wider range of events and experimental designs need to be considered to examine the robustness and implications of the findings presented, and to build understanding about the types of weather event for which this approach is best suited. Downscaling these simulations would also be possible to provide additional smaller-scale details. ~~Although~~ This approach does not currently consider any large-scale dynamical component to the changes, it may be of interest to policymakers as. ~~The analysis presented highlights how extreme~~ historical (and modern) weather events could be translated into warmer climates to ~~consider~~ construct examples of plausible ‘worst-case’ events to aid planning. This approach also offers the opportunity to attribute changes in observed weather events to human influence by translating reanalyses of modern events into cooler worlds. The 20CRv3 reanalysis currently ends in 2015 but could be extended to run in near-real-time to examine extreme events soon after they occur.

## Acknowledgements

85 Valuable discussions with Philip Brohan, Andrew Schurer, Ted Shepherd, Laura Slivinski, and Rowan Sutton are gratefully acknowledged. Support for the Twentieth Century Reanalysis Project version 3 dataset is provided by the U.S. Department of Energy, Office of Science Biological and Environmental Research (BER), by the National Oceanic and Atmospheric Administration Climate Program Office, and by the NOAA Physical Sciences Laboratory. This research used resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility located at Lawrence Berkeley National Laboratory, operated under Contract No. DE-AC02-05CH11231. EH is supported by the NERC GloSAT project and by the National Centre for Atmospheric Science. GPC and PDS were supported in part by the NOAA Cooperative Agreement with CIRES, NA22OAR4320151, and by NOAA's Physical Sciences Laboratory.

95 **Data availability**

Data available from: <https://github.com/ed-hawkins/ulysses-storm-data>

**Author contribution**

Ed Hawkins: Investigation, Formal Analysis, Writing – Original Draft, Conceptualization

100 Gilbert Compo: Conceptualization, Writing – Review & Editing, Resources

Prashant Sardeshmukh: Conceptualization, Writing – Review & Editing

**Competing interests**

None

105 **References**

Cattiaux, J., Vautard, R., Cassou, C., Yiou, P., Masson-Delmotte, V., and Codron, F.: Winter 2010 in Europe: A cold extreme in a warming climate, *Geophys Res Lett*, 37, 20704, <https://doi.org/10.1029/2010GL044613>, 2010.

Compo, G. P. and Sardeshmukh, P. D.: Oceanic influences on recent continental warming, *Clim Dyn*, 32, 333–342, <https://doi.org/10.1007/s00382-008-0448-9>, 2009.

110 Faranda, D., Bourdin, S., Ginesta, M., Krouma, M., Noyelle, R., Pons, F., Yiou, P., and Messori, G.: A climate-change attribution retrospective of some impactful weather extremes of 2021, *Weather and Climate Dynamics*, 3, 1311–1340, <https://doi.org/10.5194/wcd-3-1311-2022>, 2022.

van Garderen, L., Feser, F., and Shepherd, T. G.: A methodology for attributing the role of climate change in extreme events: A global spectrally nudged storyline, *Natural Hazards and Earth System Sciences*, 21, 171–186, <https://doi.org/10.5194/NHESS-21-171-2021>, 2021.

115 Ginesta, M., Yiou, P., Messori, G., and Faranda, D.: A methodology for attributing severe extratropical cyclones to climate change based on reanalysis data: the case study of storm Alex 2020, *Clim Dyn*, 61, 229–253, <https://doi.org/10.1007/s00382-022-06565-x>, 2023.


Hawkins, E., Brohan, P., Burgess, S., Burt, S., Compo, G. P., Gray, S. L., and et al.: Rescuing historical weather observations improves quantification of severe windstorm risks, *NHESS*, <https://doi.org/10.5194/egusphere-2022-1045>, 2022.

Klawa, M. and Ulbrich, U.: A model for the estimation of storm losses and the identification of severe winter storms in Germany, *Natural Hazards and Earth System Sciences*, 3, 725–732, <https://doi.org/10.5194/nhess-3-725-2003>, 2003.

Leach, N. J., Weisheimer, A., Allen, M. R., and Palmer, T.: Forecast-based attribution of a winter heatwave within the limit of predictability, *Proc Natl Acad Sci U S A*, 118, e2112087118, <https://doi.org/10.1073/pnas.2112087118>, 2021.

- 125 van Oldenborgh, G. J., van der Wiel, K., Kew, S., Philip, S., Otto, F., Vautard, R., King, A., Lott, F., Arrighi, J., Singh, R.,  
and van Aalst, M.: Pathways and pitfalls in extreme event attribution, *Clim Change*, 166, 1–27,  
<https://doi.org/10.1007/S10584-021-03071-7>, 2021.
- Otto, F. and et al.: Climate change likely increased extreme monsoon rainfall, flooding highly vulnerable communities in  
Pakistan, <https://www.worldweatherattribution.org/wp-content/uploads/Pakistan-floods-scientific-report.pdf>, last access: 30  
130 March 2023., 2022.
- Sánchez-Benítez, A., Goessling, H., Pithan, F., Semmler, T., and Jung, T.: The July 2019 European Heat Wave in a Warmer  
Climate: Storyline Scenarios with a Coupled Model Using Spectral Nudging, *J Clim*, 35, 2373–2390,  
<https://doi.org/10.1175/JCLI-D-21-0573.1>, 2022.
- Shepherd, T. G., Boyd, E., Calel, R. A., Chapman, S. C., Dessai, S., Dima-West, I. M., Fowler, H. J., James, R., Maraun, D.,  
135 Martius, O., Senior, C. A., Sobel, A. H., Stainforth, D. A., Tett, S. F. B., Trenberth, K. E., van den Hurk, B. J. J. M., Watkins,  
N. W., Wilby, R. L., and Zenghelis, D. A.: Storylines: an alternative approach to representing uncertainty in physical aspects  
of climate change, *Clim Change*, 151, 555–571, <https://doi.org/10.1007/S10584-018-2317-9>, 2018.
- Slivinski, L. C., Compo, G. P., Sardeshmukh, P. D., Whitaker, J. S., McColl, C., Allan, R. J., Brohan, P., Yin, X., Smith, C.  
A., Spencer, L. J., Vose, R. S., Rohrer, M., Conroy, R. P., Schuster, D. C., Kennedy, J. J., Ashcroft, L., Brönnimann, S., Brunet,  
140 M., Camuffo, D., Cornes, R., Cram, T. A., Domínguez-Castro, F., Freeman, J. E., Gergis, J., Hawkins, E., Jones, P. D., Kubota,  
H., Lee, T. C., Lorrey, A. M., Luterbacher, J., Mock, C. J., Przybylak, R. K., Pudmenzky, C., Slonosky, V. C., Tinz, B., Trewin,  
B., Wang, X. L., Wilkinson, C., Wood, K., and Wyszynski, P.: An evaluation of the performance of the twentieth century  
reanalysis version 3, *J Clim*, 34, <https://doi.org/10.1175/JCLI-D-20-0505.1>, 2021.
- Stott, P. A. and Christidis, N.: Operational attribution of weather and climate extremes: what next?, *Environmental Research:*  
145 *Climate*, 2, 013001, <https://doi.org/10.1088/2752-5295/ACB078>, 2023.
- Stott, P. A., Stone, D. A., and Allen, M. R.: Human contribution to the European heatwave of 2003, *Nature* 2004 432:7017,  
432, 610–614, <https://doi.org/10.1038/nature03089>, 2004.
- Thompson, V., Dunstone, N. J., Scaife, A. A., Smith, D. M., Slingo, J. M., Brown, S., and Belcher, S. E.: High risk of  
unprecedented UK rainfall in the current climate, *Nat Commun*, 8, 107, <https://doi.org/10.1038/s41467-017-00275-3>, 2017.
- 150 Trenberth, K. E., Fasullo, J. T., and Shepherd, T. G.: Attribution of climate extreme events, *Nature Climate Change* 2015 5:8,  
5, 725–730, <https://doi.org/10.1038/nclimate2657>, 2015.
- Wehner, M. F., Zarzycki, C., and Patricola, C.: Estimating the Human Influence on Tropical Cyclone Intensity as the Climate  
Changes, 235–260, [https://doi.org/10.1007/978-3-030-02402-4\\_12](https://doi.org/10.1007/978-3-030-02402-4_12), 2019.

# Storm Ulysses



26-27 February 1903

Figure 1: Translating Storm Ulysses into a warmer world. The columns represent the original 20<sup>th</sup> Century Reanalysis v3 (20CRv3; left), the improved reanalysis (middle) and the warmer world [simulation-reanalysis](#) (right). The wet-bulb potential temperature ( $\theta_w$ ) at 850 hPa at 06 UTC on 27<sup>th</sup> February 1903 (top row) shows the synoptic structure of the storm as it passes over the UK & Ireland. 20CRv3 does not accurately represent the intensity of the storm, but this is corrected in the improved reanalysis, mainly due to adding additional observations (Hawkins et al. 2022). The warmer world [simulation-reanalysis](#) shows a very similar atmospheric structure to the improved reanalysis and is warmer by about 2.5K at 850 hPa. The thick lines indicate particular isotherms as labelled and the black dot represents the centre of the storm. The wind footprint (middle row; maximum ensemble mean wind speed at 10m during the storm) shows stronger peak wind speeds in the improved reanalysis than 20CRv3, which are stronger again in a warmer world. The track of Storm Ulysses is shown with the thick black line. The rainfall during the two days of the storm increases in a warmer world (bottom row). Supplementary Figure 1 is an animated version of this figure which shows the 850 hPa wet-bulb potential temperature, wind speed (with sea level pressure) and rainfall rate, every 3-hours during the storm for the three experiments.