

# Contrasting extremely warm and long-lasting cold air anomalies in the North Atlantic sector of the Arctic during the HALO-( $\mathcal{AC}$ )<sup>3</sup> campaign

Andreas Walbröl<sup>1</sup>, Janosch Michaelis<sup>2,3</sup>, Sebastian Becker<sup>4</sup>, Henning Dorff<sup>5</sup>, Kerstin Ebell<sup>1</sup>, Irina Gorodetskaya<sup>6,7</sup>, Bernd Heinold<sup>8</sup>, Benjamin Kirbus<sup>4</sup>, Melanie Lauer<sup>1</sup>, Nina Maherndl<sup>4</sup>, Marion Maturilli<sup>9</sup>, Johanna Mayer<sup>10</sup>, Hanno Müller<sup>4</sup>, Roel A. J. Neggers<sup>1</sup>, Fiona M. Paulus<sup>1</sup>, Johannes Röttenbacher<sup>4</sup>, Janna E. Rückert<sup>11</sup>, Imke Schirmacher<sup>1</sup>, Nils Slättberg<sup>9</sup>, André Ehrlich<sup>4</sup>, Manfred Wendisch<sup>4</sup>, and Susanne Crewell<sup>1</sup>

<sup>1</sup>Institute for Geophysics and Meteorology, University of Cologne, Cologne, Germany

<sup>2</sup>Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Bremerhaven, Germany

<sup>3</sup>Maritime Klimatologie, Maritim-klimatologische Analysen und Produkte, Deutscher Wetterdienst (DWD), Hamburg, Germany

<sup>4</sup>Leipzig Institute for Meteorology, Leipzig University, Leipzig, Germany

<sup>5</sup>Meteorological Institute of Hamburg, University of Hamburg, Hamburg, Germany

<sup>6</sup>Centre for Environmental and Marine Studies, University of Aveiro, Aveiro, Portugal

<sup>7</sup>Interdisciplinary Centre of Marine and Environmental Research (CIIMAR), University of Porto, Matosinhos, Portugal

<sup>8</sup>Leibniz-Institut für Troposphärenforschung (TROPOS), Leipzig, Germany

<sup>9</sup>Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Potsdam, Germany

<sup>10</sup>German Aerospace Centre (Deutsches Zentrum für Luft- und Raumfahrt (DLR)), Oberpfaffenhofen, Germany

<sup>11</sup>Institute of Environmental Physics, University of Bremen, Bremen, Germany

**Correspondence:** Andreas Walbröl (a.walbroel@uni-koeln.de)

## Abstract.

How air masses transform during meridional transport into and out of the Arctic is not well represented by numerical models. The airborne field campaign HALO-( $\mathcal{AC}$ )<sup>3</sup> applied the High Altitude and Long-range Research Aircraft (HALO) within the framework of the collaborative research project on Arctic amplification ( $\mathcal{AC}$ )<sup>3</sup> to address this question by providing a comprehensive observational basis. The campaign took place from 07 March to 12 April 2022 in the North Atlantic sector of the Arctic, a main gateway of atmospheric transport into and out of the the Arctic. Here, we investigate to which degree the meteorological and sea ice conditions during the campaign align with the long-term climatology (1979–2022). For this purpose, we use the European Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis v5 (ERA5), satellite data, and measurements at Ny-Ålesund, including atmospheric soundings. The observations and reanalysis data revealed two distinct periods with different weather conditions during HALO-( $\mathcal{AC}$ )<sup>3</sup>: The campaign started with a warm period (11–20 March 2022) where strong southerly winds prevailed that caused poleward transport of warm and moist air masses, so-called moist and warm air intrusions (WAIs). Two WAIs events were identified as Atmospheric Rivers (ARs), which are narrow bands of strong moisture transport. These warm and moist air masses caused the highest measured 2m temperatures (5.5°C) and daily precipitation rates (42mm day<sup>-1</sup>) at Ny-Ålesund for March since the beginning of the record (1993). Over the sea ice

15 northwest of Svalbard, ERA5 indicated record-breaking rainfall. After the passage of a strong cyclone on 21 March 2022,  
a cold period followed. Northerly winds advected cold air into the Fram Strait causing marine cold air outbreaks (MCAOs)  
until the end of the campaign. This second phase included one of the longest MCAO events found in the ERA5 record (19  
days). On average, the entire campaign period was warmer than the climatological mean due to the strong influence of the  
ARs. In the Fram Strait, the sea ice concentration was well within the climatological variability over the entire campaign  
20 duration. However, during the warm period, a large polynya opened northeast of Svalbard untypical for this season. Compared  
to previous airborne field campaigns focusing on the evolution of (mixed-phase) clouds, a larger variety of MCAO conditions  
were observed during HALO-( $\mathcal{AC}$ )<sup>3</sup>. In summary, air mass transport into and out of the Arctic were both more pronounced  
than usual, providing exciting prospects to study air mass transformation using HALO-( $\mathcal{AC}$ )<sup>3</sup>.



## 1 Introduction

25 Currently, the Arctic experiences a drastic temperature increase, which is up to 4 times stronger compared to the rest of the globe (Rantanen et al., 2022). This enhanced Arctic warming is one of the most obvious signs of Arctic amplification, which is caused by numerous feedback mechanisms (Serreze et al., 2009; Screen and Simmonds, 2010; Serreze and Barry, 2011). While the contributions of some feedback mechanisms to Arctic amplification can be regarded as scientific consensus (e.g., sea ice–albedo feedback, Serreze et al., 2009), others are not yet sufficiently explored (e.g., influence of clouds, lapse-rate feedback, 30 Wendisch et al., 2023). Especially the role of linkages between Arctic and mid-latitudes is still under debate. Moist and warm air intrusions (WAIs) transport large amounts of heat and moisture into the Arctic through the Atlantic sector (Woods and Caballero, 2016). Over the past decades, the frequency of meridional transports through the North Atlantic has increased due to more frequent atmospheric blocking over the Barents Sea (Mewes and Jacobi, 2019; You et al., 2022). Woods et al. (2013) and Woods and Caballero (2016) found that intense WAIs may have a large effect on the downward thermal-infrared radiation 35 at the surface, contributing to the enhanced warming of the Arctic. Furthermore, WAIs precondition the sea ice for the melting season, resulting in lower sea ice extent at the end of the summer (Kapsch et al., 2013, 2019). Numerical models struggle to accurately represent mixed-phase clouds and the transformation processes of the meridionally transported air masses (Pithan et al., 2014; Cohen et al., 2020).

WAIs are often linked with filaments of strong moisture transport, known as Atmospheric Rivers (ARs, Newell et al., 1992). 40 ARs are responsible for over 90% of the poleward moisture transport across the mid-latitudes and are frequently accompanied by strong winds and precipitation (Nash et al., 2018). While the precipitation related to ARs can cause snow accumulation in the Arctic, the enhanced emission of downward thermal-infrared radiation by the clouds can lead to melting of snow and sea ice (Neff et al., 2014; Komatsu et al., 2018; Mattingly et al., 2018, 2020; Bresson et al., 2022; Viceto et al., 2022). In a warming climate, ARs are expected to shift polewards and to intensify due to the increased moisture load (Ma et al., 2020).

45 Cold air outbreaks (CAOs) are often responsible for severe weather events in the high- and mid-latitudes and mainly occur in winter and spring (Fletcher et al., 2016; Pithan et al., 2018). During marine CAOs (MCAOs), cold and dry air is advected southwards from the sea ice to the ice-free (open) ocean. Over the open ocean, the strong temperature contrast between the surface and the lower tropospheric air leads to intense fluxes of sensible and latent heat, responsible for 60–80% of the oceanic heat losses in that region (Papritz and Spengler, 2017). This heat and moisture transfer into the atmosphere destabilizes the 50 atmospheric boundary layer and leads to the formation of cloud streets, which later develop into open cloud structures. This cloud evolution is difficult to capture by atmospheric models (Pithan et al., 2018) motivating dedicated measurement campaigns (Geerts et al., 2022; Lloyd et al., 2018).

So far, observations of air mass transformations in the Arctic have mostly been conducted from a fixed local position (Eulerian view). Only a few aircraft based samplings of air mass properties over a limited regional area have been reported (e.g., 55 Wendisch et al., 2019; Mech et al., 2022). To observe air mass transformations along their meridional pathway in a Eulerian view, multiple stations that are aligned with the wind direction would be needed. Therefore, Pithan et al. (2018) proposed a quasi-Lagrangian approach following air masses to and from the Arctic. This motivated the field campaign HALO-( $\mathcal{A}\mathcal{C}$ )<sup>3</sup>

within the Transregional Collaborative Research Center TRR 172 "Arctic amplification: Climate Relevant Atmospheric and Surface Processes and Feedback Mechanisms ( $\mathcal{AC}$ )<sup>3</sup>". The campaign was designed to obtain quasi-Lagrangian observational data of air mass transformations during WAIs and MCAOs to gain process understanding and evaluate the performance of weather and climate models (Wendisch et al., 2021, 2024).

During HALO-( $\mathcal{AC}$ )<sup>3</sup>, extensive remote sensing and in situ measurements of surface, cloud, and thermodynamic properties have been performed between the Norwegian Sea and the North Pole from 11 March to 12 April 2022 using three research aircraft (HALO, Polar 5, and Polar 6). The High Altitude and Long-range Research Aircraft (HALO) operated by the German Aerospace Center (Ziereis and Gläßer, 2006; Stevens et al., 2019) is a modified Gulfstream G550. It has an operating range of 9000 km in altitudes up to 15 km, which is beneficial for quasi-Lagrangian air mass observations, and was based in Kiruna during HALO-( $\mathcal{AC}$ )<sup>3</sup>. It was equipped with a similar instrumental payload as during the EUREC<sup>4</sup>A campaign (Stevens et al., 2019, 2021; Konow et al., 2021). Polar 5 and Polar 6 (P5 and P6) were based in Longyearbyen and operated by the Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research (Wesche et al., 2016).


This study aims to investigate whether the atmospheric and sea ice conditions encountered during HALO-( $\mathcal{AC}$ )<sup>3</sup> were suitable for studying air mass transformation. Herein we analyze which types of air mass transport occurred and how representative the conditions were. Therefore, the conditions are analysed in the climatological context. In this way, our study serves as comprehensive reference for future studies analyzing the HALO-( $\mathcal{AC}$ )<sup>3</sup> campaign. After introducing the data and methods in Sect. 2, we provide a general overview of the weather conditions during HALO-( $\mathcal{AC}$ )<sup>3</sup> (Sect. 3), which leads to the identification of a warm (Sect. 4) and a cold period (Sect. 5). Both periods are assessed in the climatological context. For the unusually strong warm phase, we further investigate how the associated precipitation might have influenced sea ice conditions (Sect. 6). Finally, we conclude our study in Sect. 7.

## 2 Data and Methods

Our study concentrates on the North Atlantic sector of the Arctic, the major pathway of WAIs (Johansson et al., 2017; You et al., 2022). Around the Fram Strait, we define a central region including the marginal sea ice zone (Fig. 1) where the air mass transfer from the ocean to the sea ice during WAIs and vice versa during MCAOs is most pronounced. Within this region, most airborne measurements were performed. To better illustrate how the meridional air mass transition progresses, we also include a southerly domain over ocean including the Greenland and Norwegian Seas between Svalbard and Norway and a northerly one in the central Arctic over sea ice. Note that circulation weather type analysis has shown that the flow in this area is generally meridional, while zonal flow hardly occurs (Schirmacher et al., 2023; von Lerber et al., 2022).

### 2.1 Atmospheric measurements

Only very few permanent radiosonde stations exist in the Arctic. Therefore, the French–German AWIPEV research base in Ny-Ålesund (Svalbard), which is the only station within our central domain, plays a key role in observing atmospheric conditions in the region. We use the 6-hourly soundings at Ny-Ålesund (Vaisala RS41, Maturilli, 2020b) performed during HALO-( $\mathcal{AC}$ )<sup>3</sup>



**Figure 1.** Study area with corresponding domains (southern region, central region, and northern region) and mean sea ice concentration (07 March to 12 April 2022) based on MODIS–AMSR2 satellite data (Ludwig et al., 2020). The locations of the dropsondes launched by HALO (P5) are marked as purple (green) circles. Operational radiosonde stations are indicated by yellow triangles.

90 to assess vertical profiles of temperature, relative humidity, pressure and wind. Similar in situ profiling data is available from dropsondes (Hock and Franklin, 1999; George et al., 2021) launched by HALO and the Polar aircrafts. In total 330 dropsondes from HALO and 141 from Polar 5 and 6, distributed over 18 and 13 research flights, respectively, provide profiles in otherwise not sampled areas (Fig. 1).

From the radiosonde profiles, we identified the thermal tropopause according to the WMO definition as the lowest level at 95 which the temperature lapse-rate falls below  $2\text{K km}^{-1}$  and does not exceed this value for the next 2 km. The integrated water vapor (IWV) was calculated from profiles of air pressure and specific humidity. Ground-based meteorological measurements at AWIPEV are available since 1993 and provide an additional view on the environmental conditions during HALO-(AC)<sup>3</sup> (Maturilli et al., 2013; Maturilli, 2020a). Here, the CL–51 ceilometer is used to assess the cloud conditions at Ny-Ålesund (Maturilli and Ebell, 2018; Maturilli, 2022). The detection of a cloud base is used to derive the frequency of cloud occurrence 100 on a daily base. Long-term precipitation observations are performed manually by the Norwegian Meteorological Institute (MET Norway). For Ny-Ålesund (station number SN99910), daily precipitation totals (06 to 06 UTC) are available since 01 January 1974 and have been analyzed in this study for March and April of the years 1974 to 2022.

## 2.2 Satellite observations of sea ice

Sea ice concentration (SIC), i.e., the percentage of a satellite pixel covered by sea ice, is obtained from the Ocean and Sea Ice  
105 (OSI) Satellite Application Facility (SAF) Global Sea Ice Concentration Climate Data Record (SIC CDR v2.0), namely the  
product OSI-450 from 1979 to 2015, and the complementary Interim Climate Data Record OSI-430-b from 2016 onwards  
(OSI SAF, 2017; Copernicus Climate Change Service (C3S), 2020). The product is based on low-frequency passive microwave  
satellite data and is described in Lavergne et al. (2019). The datasets are available on a daily basis with a grid spacing of  
25 km  $\times$  25 km. For a higher spatial resolution (1 km), i.e., for the analysis of how SIC changed during HALO-( $\mathcal{A}$ )<sup>3</sup>, the  
110 merged MODIS-AMSR2 SIC product (Ludwig et al., 2020) is used.

## 2.3 Reanalysis data and diagnostics

The European Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis ERA5 (Hersbach et al., 2018a, b, 2020)  
offers hourly data output since 1950 with a horizontal resolution of 31 km and 137 vertical model levels. For our region,  
ERA5 has been found superior compared to other global reanalysis (Graham et al., 2019a, b; Avila-Diaz et al., 2021). With  
115 very few ground-based stations in our study area, the assimilation of satellite data plays a major role for the quality of ERA5.  
Therefore, our climatological analysis uses the years 1979–2022 starting with the onset of the satellite era. Note that during  
HALO-( $\mathcal{A}$ )<sup>3</sup>, 216 dropsondes were used in the Global Telecommunication System data assimilation, likely resulting in an  
improved reanalysis quality.

Domain averages of the regularly gridded ERA5 data were calculated as area averages weighted by the cosine of the latitude  
120 to respect the increasing data point density with increasing latitudes. Grid points with a land fraction  $> 0$  have been excluded  
from our analysis. ERA5 data serve to detect WAI, ARs, and MCAOs:

- **WAIs** are identified by positive daily means of the vertically integrated meridional moisture flux ( $IVT_{\text{north}}$ ) averaged  
over the central region. Guided by the study of Woods and Caballero (2016) an WAI is considered weak (strong) when  
 $IVT_{\text{north}}$  is below (equal or above)  $100 \text{ kg m}^{-1} \text{ s}^{-1}$ .
- 125 – **ARs** are detected by the global algorithm from Guan and Waliser (2015) in its revised version (Guan et al., 2018), adapted  
to the lower moisture content of the Arctic by reducing moisture transport thresholds. Data set in one-hour resolution is  
available at Lauer et al. (2023).
- **MCAOs** are characterized by their strong temperature decrease with increasing height over the open ocean. The MCAO  
index  $M$  is calculated following Papritz and Spengler (2017) and Dahlke et al. (2022):

$$130 \quad M = \theta_{\text{SKT}} - \theta_{850} \quad (1)$$

with  $\theta_{\text{SKT}}$  ( $\theta_{850}$ ) as the potential skin temperature (potential temperature at 850 hPa). Grid points with skin temperatures  
below 271.5 K (i.e., over sea ice) were excluded from further processing as in Dahlke et al. (2022). Following Papritz  
and Spengler (2017), MCAO conditions are present when  $M > 0 \text{ K}$  and its strength can be classified as weak ( $0 \text{ K} <$   
 $M \leq 4 \text{ K}$ ), moderate ( $4 \text{ K} < M \leq 8 \text{ K}$ ), strong ( $8 \text{ K} < M \leq 12 \text{ K}$ ) or very strong ( $M > 12 \text{ K}$ ).

135 For climatological assessment, we average over each region and compute the temporal mean, 25–75<sup>th</sup> and 10–90<sup>th</sup> percentiles of  $IVT_{\text{north}}$  and  $M$  over the climatology period.

### 3 Overview of the campaign period


#### 3.1 Ny-Ålesund

Radiosoundings from Ny-Ålesund provide the only continuous source of information on the vertical structure of the atmosphere  
140 in our study region and are therefore frequently used to characterize the climate of the whole North Atlantic sector of the Arctic (Maturilli et al., 2013). The temperature and moisture profiles measured by radiosondes at Ny-Ålesund during the HALO-(AC)<sup>3</sup> are shown in Fig. 2 and indicate a high temporal variability at all altitudes. Temperatures near the surface (5 km) vary between  $-20^{\circ}\text{C}$  and  $+5^{\circ}\text{C}$  ( $-45^{\circ}\text{C}$  and  $-20^{\circ}\text{C}$ ), indicating the presence of different air masses. This is confirmed by wind measurements, which reveal episodes of strong southerly winds associated with warm temperatures in the first half of  
145 the campaign, while in the second half of the campaign weaker, mainly northerly winds associated with lower temperatures prevail. The strongest episode of warm air advection on 12–13 March led to an increase of 2 m temperature from about  $-14^{\circ}\text{C}$  to  $+2^{\circ}\text{C}$  within 19 hours (meteorological tower measurements, not shown). This air mass lifted the tropopause up to 13 km (Fig. 2a), and the 2 m temperature remained above freezing for five days. The 2 m temperature even reached a new maximum for March on 15 March ( $5.5^{\circ}\text{C}$ ) since the beginning of the data record in 1993.

150 To investigate whether the observed warm air advection was related to ARs we analyze humidity profiles. The first half of the campaign featured a number of short-term events of high integrated water vapor (IWV) as shown in Fig. 2b. On 15 March, a radiosonde measured an IWV of  $14.6\text{ kg m}^{-2}$ , which is the highest value recorded between 07 March and 12 April from 1993 to 2022. The rapid doubling of IWV within a day already gives an indication of the presence of ARs. Applying the detection algorithm to reanalysis (Sect. 2) confirms that two ARs passed over Ny-Ålesund. AR I arrived at Ny-Ålesund on 12 March  
155 at 13 UTC and lasted until 14 March, 13 UTC. However, in the AR detection algorithm this AR consisted of two parts with enhanced moisture transport and IWV, which can also be seen in the IWV dip on 13 March between two peaks (12 and 14 March) in Fig. 2b. AR II reached Ny-Ålesund on 15 March at 03 UTC and was only detected until 22 UTC of the same day although IWV and the moisture transport stayed on high levels until 16 March, 15 UTC (not shown).

The second half of the campaign was drier, featuring the lowest IWV of less than  $1.2\text{ kg m}^{-2}$  on 24 March at 06 UTC. This  
160 IWV is below the 3<sup>rd</sup> percentile of all radiosondes between 07 March and 12 April from 1993 to 2022 (Maturilli and Kayser, 2016, 2017; Maturilli, 2020b). Afterwards, IWV varied less and stayed below  $4\text{ kg m}^{-2}$  throughout the second half of the campaign. Only on 10 April, enhanced moisture values indicate a weak moist air advection that was relevant for the formation of cirrus clouds over sea ice observed by HALO (not shown).

The ceilometer at Ny-Ålesund reveals high cloud occurrence until 23 March (Fig. 2c). The high cloudiness is accompanied  
165 by precipitation and related to the WAIs and ARs passing over Ny-Ålesund. From 15 March, 06 UTC to 16 March, 06 UTC, 42 mm of precipitation was recorded, which is the highest daily accumulated precipitation for March since the beginning of the measurements. The situation changed during the course of 17 March, when the cloud deck started to dissolve, and the



**Figure 2.** Time series of Ny-Ålesund radiosondes of (a) temperature profiles (shading), height of thermal tropopause (black line), and wind barbs in selected levels, (b) specific humidity profiles (shading) and resulting I WV (black line, right axis). (c) shows daily accumulated precipitation (06 to 06 UTC) from gauge measurements and daily mean cloud occurrence from ceilometer measurements.

2 m temperature dropped below 0 °C. No further precipitation was observed after 23 March, and cloud occurrence generally remained low at Ny-Ålesund until the end of the campaign. However, note that Ny-Ålesund is located within a fjord on the west coast of Svalbard and measurements, especially within the atmospheric boundary layer, may be influenced by the local orography (Gierens et al., 2020). For example, the relatively cloud-free second part of the campaign was associated with lee effects.

### 3.2 Reanalysis


Given the scarcity of measurements, e.g., no permanent radiosonde stations north of 82.5° N (Rinke et al., 2019), reanalysis data (ERA5) are used to characterize the conditions over the whole study domain. However, ERA5 assimilated dropsonde measurements from HALO (details will be given in a dedicated study presenting all HALO-(AC)<sup>3</sup> measurements), influencing

its performance: In Fig. 3, we show the agreement between ERA5 and HALO dropsondes in terms of root mean square deviation (RMSD) and bias profiles. Over ocean, the ERA5 temperature agrees with the dropsonde measurements within 0.5–0.9K (Fig. Fig. 3a, b). Over sea ice, we see the expected warm bias of ERA5 in the lower troposphere, which was also found  
180 by Yu et al. (2021), leading to higher biases and RMSD of up to 1.5K. The RMSD of relative humidity is between 10 and 20%, while wind and pressure deviations are  $1.5\text{--}2\text{m s}^{-1}$  and less than 1 hPa, respectively. Deviations can be higher for single dropsondes because of the spatio-temporal variability of humidity and temperature close to the sea ice edge and ARs, and due to measurement errors of the dropsondes (especially humidity in dry conditions and at low temperatures). The temperature, relative humidity and wind speed deviations found here agree well with those in Graham et al. (2019b). Note that the deviations  
185 would probably be higher if dropsonde measurements were not assimilated. Given the good performance of ERA5, we analyze the environmental conditions in the following.

Time series of area-averaged mean sea level pressure (MSLP), 10m wind, 2m temperature, 850hPa temperature, and IWV (Fig. 4) illustrate that the major features of warm and moist conditions in mid-March and cooler and drier conditions later on observed at Ny-Ålesund also apply to the central domain. We also look at the spatial distribution of mean sea level pressure,  
190 500hPa geopotential height and 850hPa equivalent-potential temperature in the North Atlantic sector of the Arctic (Fig. 5) to investigate the driving conditions for synoptic events, such as ARs and MCAOs.


The strong warming events observed at Ny-Ålesund between 12 and 16 March are related to a low-pressure system over Greenland, which drives southerly winds through the North Atlantic (Fig. 5a). From 13 to 15 March, the zonal pressure gradient across the North Atlantic intensifies, leading to stronger southerly winds in all three domains (Fig. 5b, Fig. 4b). In this  
195 period, the IWV is very high in the central and northern domains with daily area averages of  $12\text{--}14\text{kg m}^{-2}$  and  $6\text{--}8\text{kg m}^{-2}$ , respectively (Fig. 4e). At the same time, ERA5 indicates 2m temperatures above freezing in the central domain, which agrees with the time series observed at Ny-Ålesund (Fig. 4c). Around 20 March, the MSLP gradient between the three domains reverses due to changes in the large-scale pressure constellation (Fig. 5c). The central Arctic now shows typical conditions with the highest MSLP of all regions and the strongest static stability as can be judged from the difference between temperatures at  
200 850hPa and at 2m (Fig. 4a, c, d). The high MSLP of the central Arctic agrees well with the climatological pressure shown for April in Fig. 4.8 of Serreze and Barry (2014). In this second part of the campaign, northerly winds typical for MCAO activity led to extremely dry conditions with IWV down to  $1.4\text{kg m}^{-2}$  in the northern domain (Fig. 4e), clearly indicating a polar air mass.

Motivated by the clear differences between the first and second half of the campaign, we separate the campaign into two  
205 major periods based on the northward component of the integrated water vapor transport ( $IVT_{\text{north}}$ ) and the MCAO index ( $M$ ) in the central region (see Fig. 6). From 11 to 20 March, from here on called the warm period,  $IVT_{\text{north}}$  is positive, indicating warm air advection into the Arctic. In the subsequent period (21 March to 12 April, hereafter referred to as cold period),  $IVT_{\text{north}}$  turns slightly negative, indicating a moisture flux out of the Arctic. This is connected to MCAO activity as quantified by the positive MCAO index in the cold period. In the following, we assess the weather conditions of warm and cold periods  
210 in the climatological context separately.



**Figure 3.** ERA5 errors (bias and root mean square deviation (RMSD)) of (a,b) temperature (T), (c,d) relative humidity (rh), (e,f) wind speed, and (g,h) pressure (p) using dropsonde observations as reference. The errors are separated into open ocean and sea ice using the ERA5 sea ice concentration. Additionally, we added the number of available dropsonde observations in each height level. Note, that the number of dropsondes in the wind speed comparison is lower than for other variables due to gaps in the GPS communication necessary for determining the sonde position.






**Figure 4.** Regional averages (colored lines) and standard deviation (shading) of ERA5-based daily means of (a) mean sea level pressure, (b) 10m wind speed (wind barbs), (c) 2 m temperature, (d) 850 hPa temperature and (e) integrated water vapor (IWV). Regional averages are performed for the southern (S, red), central (C, green), and northern (N, blue) regions shown in Fig. 1. Vertical black lines indicate days shown in Fig. 5.


#### 4 Warm air intrusions

To answer what caused the occurrence of the strong WAIs we analyze the general circulation pattern. A high surface pressure system over Scandinavia was connected to a ridge in the 500hPa geopotential height, resulting in a blocking situation (Fig. 5a, b). The blocking is evident in Fig. 7b (and also Fig. B1a in Appendix B), showing the strong anomalies of more than 10hPa in the MSLP field over the whole warm period with lower pressure over Greenland and the central Arctic, and higher pressure over Scandinavia. The strength of the pressure anomalies over Scandinavia suggests the presence of an unusual blocking situation. Intense WAIs are often connected to blocking situations over the eastern border of a large basin (here, North Atlantic), redirecting cyclones northward as the typical eastward propagation is blocked (Woods et al., 2013). Consequently,



**Figure 5.** Maps of mean sea level pressure (white contour lines with black outlines), 500hPa geopotential height (black contour lines), and 850hPa equivalent-potential temperature (shading and grey contours) from ERA5 data for representative days of the main weather conditions at 12 UTC. The 15% sea ice concentration from AMSR2 is displayed as blue contour line. "H" and "L" illustrate the centres of high and low pressure systems, respectively.

warm and moist air masses originating from the North Atlantic were transported towards the Fram Strait, driven by several  
 220 low-pressure systems that formed between Iceland and eastern Greenland. All days of the warm period passed the criteria for




**Figure 6.** Daily mean time series of (a) northward component of integrated water vapor transport ( $IVT_{\text{north}}$ ), (b) Marine cold air outbreak (MCAO) index ( $M$ ) based on ERA5 and averaged over the central region for HALO- $(\mathcal{A}C)^3$  (black line). The mean, as well as the 10–90<sup>th</sup> and 25–75<sup>th</sup> percentiles of the climatology (1979–2022), are illustrated as grey lines, and light and dark grey shadings. The red (blue) box indicates the warm (cold) period dominated by moist and warm air intrusions (marine cold air outbreaks). Vertical dashed black lines mark the days shown in Fig. 5.

WAI occurrence (Sect. 2), and from 12 to 17 March,  $IVT_{\text{north}}$  exceeded the 90<sup>th</sup> percentile of the ERA5 climatology on each day (Fig. 6a).

Not surprisingly, temperatures at 2m and 850hPa show positive anomalies over all domains during the warm period (Fig. 7e, h). However, with far above 8K, the strongest 2m temperature anomalies occur over sea ice surfaces. Together with the weaker positive anomaly at 850hPa (up to 7K in the three domains), this demonstrates the decreased stability of the lower atmosphere during this period over sea ice. The warm period was also moister than average as indicated by the positive I WV anomalies over all domains (Fig. 7k). Note that although the highest I WV anomalies occurred in the southern region, the relative effect increases with higher latitudes as I WV mean values generally decline towards the North along with the decreasing temperatures. The highest relative I WV anomalies (up to 90%) occurred over sea ice northwest and northeast of Svalbard (not shown). Note that this latitudinal effect also makes it difficult to diagnose ARs based on I WV only.

The two AR events detected at Ny-Ålesund also affected the entire measurement area of HALO- $(\mathcal{A}C)^3$ . However, the timing might be different, depending on the exact location. AR I passed through the measurement regions on 12–14 March (Fig. 2b



**Figure 7.** From top to bottom: Anomaly maps of mean sea level pressure (MSLP), 2 m temperature ( $\Delta T_{2m}$ ), 850hPa temperature ( $\Delta T_{850hPa}$ ), and integrated water vapor ( $\Delta IWV$ ) based on ERA5 for the entire campaign (left), the warm (middle), and the cold (right) period. The long—term mean (1979–2022) is subtracted from the mean of the respective time period. Dashed black isolines describe the long—term climatology for the respective period in each panel. The three measurement regions (shown in Fig. 1) are illustrated as black boxes.


and Fig. 5a). In the northern domain, IWV reached its campaign maximum ( $8 \text{ kg m}^{-2}$ ) on 13 March and declined afterwards while the 850 hPa and 2 m temperature in the northern domain continued to increase until the following day (Fig. 4c–e). AR 235 II, which arrived on 15 March (Fig. 5b), had slightly less IWV over the central domain (Fig. 4e). This shows that Ny-Ålesund is not representative of the whole region because here, AR II had the highest IWV. The strong northward moisture transport related to these ARs can be seen in area averages of  $\text{IVT}_{\text{north}}$ , which reached  $225 \text{ kg m}^{-1} \text{ s}^{-1}$  due to AR II in the central domain (Fig. 6a).

We compare the strength of the ARs during HALO-( $\mathcal{AC}$ )<sup>3</sup> to the long-term climatology in Fig. 8. For this purpose, the 240 mean IVT over the detected AR area and the AR’s central latitude are shown in relation for all six-hourly ERA5 time steps since 1979. Note that a slightly different region over the North Atlantic and the Arctic Ocean was selected to extend the view southwards to cover the major pathways of ARs (Guan and Waliser, 2017; Nash et al., 2018). The number of strong AR events decreases meridionally (Fig. 8) because of two effects: Firstly, along their northward propagation, ARs generally decline in intensity. Secondly, within the Arctic circle, the likelihood of new AR formation strongly decays, as the moisture uptake from 245 the ocean is substantially reduced (Papritz et al., 2022).

AR I and AR II both represent strong cases in terms of mean IVT as they partly lie outside the 25<sup>th</sup> percentile in latitude–IVT space (Fig. 8). At its northernmost position with a central latitude of  $80^\circ \text{ N}$ , AR I had a stronger mean IVT than 90% of all ARs in the climatology. However, AR II was characterized by even stronger moisture transport but did not reach as far north as AR I. When AR II was at its northernmost position (centered at  $76^\circ \text{ N}$ ), its area-averaged IVT was just below 250  $200 \text{ kg m}^{-1} \text{ s}^{-1}$  and therefore also higher than 90% of all ARs in the climatology (Fig. 8). The maximum IVT of AR II (meridional and zonal) slightly exceeded  $400 \text{ kg m}^{-1} \text{ s}^{-1}$  on 15 March in the central region (Fig. A1 in Appendix A). HALO’s dropsonde measurements showed a maximum IVT of  $490 \text{ kg m}^{-1} \text{ s}^{-1}$  on 15 March, suggesting that ERA5 underestimates the moisture flux at the local scale. As all AR events had meridionally elongated structures, the outflow region reached far up to the central Arctic while their centers were located at  $68\text{--}80^\circ \text{ N}$  (Fig. 8). During their poleward propagation, the moisture transport 255 decreased so that they no longer fulfilled the detection requirements.

The northward transport of warm and moist air was associated with anomalous amounts of precipitation (Fig. 9). Particularly when the ARs made landfall in Scandinavia, Svalbard, and eastern Greenland, strong anomalies of up to  $8 \text{ mm day}^{-1}$  compared to the climatology were observed. The strongest anomalies found on the east coast of Greenland relate to 6 times higher precipitation compared to the climatology. For Svalbard, the ERA5 data are in line with the record-precipitation measured at 260 Ny-Ålesund (Fig. 2) coming from AR II. The area of increased precipitation reaches up to the North Pole, where precipitation is enhanced by more than a factor of 2.

The phase of the precipitation is highly important for its climate effect, and the transition of Arctic precipitation from snow-fall to rain is heavily discussed (Serreze et al., 2021). While we note the difficulties of correctly differentiating precipitation phase, it is important to note that northwest of Svalbard, ERA5 showed record-breaking rainfall connected with near surface 265 temperatures above zero. Liquid precipitation at high latitudes over sea ice was also observed by the cloud radar onboard HALO as we detected a distinct bright band in the linear depolarization ratio at about 0.5–1 km height on 13 March (Fig. 10).



**Figure 8.** Six-hourly climatological (1979–2022) distribution of central latitudes of Atmospheric Rivers (ARs) as a function of mean AR integrated water vapor transport (IVT) using an ERA5-based AR catalog based on Guan and Waliser (2015). The analyzed area covers 60–90° N and 60° W–40° E. Blue dots mark ARs whose center was within the latitudes and longitudes of the central domain. Cases categorized as ARs during HALO- $(\mathcal{A}C)^3$  are illustrated by colored squares. Black (dashed) lines indicate the 25<sup>th</sup> and 75<sup>th</sup> (10<sup>th</sup> and 90<sup>th</sup>) percentiles of a kernel density estimation to visualize the shape of the histogram. The green rectangle marks the latitudes of the central domain.


Thus, the observations confirm the presence of liquid precipitation at least in some regions over sea ice. This motivated us to examine the impact of this AR on the sea ice later on (Sect. 6).

In summary, the warm period observed during HALO- $(\mathcal{A}C)^3$  featured ARs with averaged IVT values higher than 90% of all ARs in the climatology. This transport was related to a blocking event affecting the Arctic up to the North Pole with heat and anomalously high precipitation, including record-breaking rainfall rates over the sea ice northwest of Svalbard.


## 5 Cold air outbreaks

The warm period was followed by a cold and dry period from 21 March until the end of the campaign, where  $IVT_{\text{north}}(M)$  was almost exclusively negative (positive) (Fig. 6). The transition from the warm to the cold period was initiated on 21 March, when an intense cyclone centered east of Svalbard began to draw cold air from the central Arctic into the Fram Strait (Fig.





**Figure 9.** Hourly averaged total precipitation (sum of snowfall and rain rate) ( $\text{mm h}^{-1}$ ) (top row) and absolute deviations from the climatology (bottom row) derived from ERA5 for (a, c) the warm period, and (b, d) the cold period.



**Figure 10.** (a) Linear depolarization ratio measured by HALO's radar during research flight 3 on 13 March 2022 and the collocated sea ice concentration (SIC) from AMSR2, which is included in the published radar data. (b) Map showing the flight track (white line with black outline) and the sea ice concentration from the MODIS-AMSR2 product. The segment of the flight track shown in (a) is highlighted in blue.

5c), which caused snowfall on the west coast of Svalbard. In its aftermath, the MSLP anomaly reversed compared to the warm period, with positive values in the northwest and negative values in the southeast (Fig. 7c). Consequently, the southward flow of cold Arctic air led to cold and dry anomalies at 850 hPa over the open ocean (Fig. 7i, l).

280 Within the cold period, two periods of strong MCAO conditions (21–26 March and 01–02 April) were identified whose strength exceeded the 90<sup>th</sup> (25 March) and 75<sup>th</sup> (2 April) percentile of the climatology (Fig. 6b). During these strong MCAO conditions, we found the maximum values of turbulent surface sensible and latent heat fluxes, which coincided with the highest wind speeds and lowest temperatures of the cold period (not shown). Low-level cloud cover remained high in the central domain until 07 April, reflecting the presence of the convective boundary layer clouds typically associated with MCAOs. Interestingly, while most parameters have reversed anomalies compared to the warm period, precipitation in the central domain also shows a positive anomaly. ERA5 produces an extensive region of enhanced precipitation over open water extending far east and being highest at the Scandinavian coast (Fig. 9).

285 To compare the MCAOs encountered in the cold period to the climatology we apply a metric established by Knudsen et al. (2018) and characterize each event by its strength and duration in Fig. 11. Herein the duration is defined as the time the MCAO index stays above zero, and the strength is given by its maximum MCAO index  $M$  as indicated in the time series of  $M$  in Fig. 11b. Note that we used the central domain for the computation of  $M$  while Knudsen et al. (2018) considered a smaller area that lies within our central domain.


The statistics over the long-term ERA5 climatology (Fig. 11a) generally show that longer MCAO events are also more intense (higher peak  $M$ ). During the HALO-( $\mathcal{AC}$ )<sup>3</sup> period, distinct MCAO events could be identified, separated by short phases with  $M < 0$  K visible in high temporal resolution only. The first and third event, starting on 18 March and 11 April, respectively, were relatively short-lived (2–3 days) and had a moderate strength ( $M$  of about 5 K). Most notably, the second event, which started on 21 March after the passage of the strong cyclone, was unusually long (19 days) and had a maximum  $M$  of about 11 K. The longevity of the MCAO conditions can be explained by the persistence of low pressure over Scandinavia, the Barents Sea and Russia, and high pressure over Greenland and the central Arctic during the cold period (Fig. B1b in Appendix B). Analyzing the time series with hourly resolution (Fig. 11b), the 21 March MCAO event consists of multiple waves of strong MCAO conditions, which were associated with the presence of cyclones near Svalbard influencing the atmosphere over the Fram Strait (e.g., Fig. 5c, e, f). Lower tropospheric winds were stronger and more northerly during the stronger MCAOs. When the cyclones dissipated or propagated away from the area, winds were weaker, and the MCAO strength decreased (e.g., Fig. 5d).

Since 1979, only 9 of the 179 events over the HALO-( $\mathcal{AC}$ )<sup>3</sup> period had a longer duration.

## 305 **6 Effect of WAI on sea ice conditions**

The different atmospheric conditions likely affected the sea ice conditions in the measurement area. To understand the impact on sea ice, we examine the spatial distribution of SIC before (09–11 March) and during (14–16 March) the WAIs and ARs, as well as after the MCAO period (10–12 April) as shown in Fig. 12a–c. SIC was > 90% in almost the entire northern region in the first period (Fig. 12a). In the central region, the sea ice covered 30–40% of the area with a southwest–northeast oriented edge. The southern region was almost completely ice-free over the entire campaign duration (Fig. 12a–c).





**Figure 11.** a) Two-dimensional histogram showing the ERA5-based duration and strength of MCAOs in the central domain for the period 1979–2022. Cases categorized during HALO- $(AC)^3$  are illustrated by colored squares and the legend entries of these cases indicate the start date. Black (dashed) lines indicate the 25<sup>th</sup> and 75<sup>th</sup> (10<sup>th</sup> and 90<sup>th</sup>) percentiles of a kernel density estimation to visualize the shape of the histogram. b) MCAO index time series with higher temporal resolution and indicating the MCAO periods marked in a).

The WAIs and ARs caused a reduction of SIC in the northwestern part of the central region, resulting in the lowest SIC values of the entire campaign period (Fig. 12b, d). While the main sea ice boundary stayed at the same location, the SIC in the marginal ice zone declined. SIC was below the climatological mean but remained within the 10–90<sup>th</sup> percentiles. Furthermore, an unusually large polynya opened from north of Svalbard to Franz Josef Land, which resulted in the below-average static stability and stronger turbulent surface heat fluxes during the cold period in this region (Fig. 12b). We assume that ice dynamics related to strong winds caused the SIC decrease in the central region, but ice melt cannot be excluded as well because temperatures were above freezing, and liquid precipitation was observed over sea ice (Sect. 4). Rapid ice melt has been attributed to warm air advections and induced increases of heat flux in summer (Tjernström et al., 2015; Woods and Caballero,

2016). Liquid precipitation on snow alters the signal of the microwave radiometry and increases the uncertainty of SIC prod-  
320 ucts (Stroeve et al., 2022; Rückert et al., 2023). However, SIC reduction was obvious in visual satellite images as well (e.g.,  
NASA Worldview, not shown). After the WAIs, the sea ice conditions recovered as temperatures fell below the freezing point  
and northerly winds were established during the MCAOs (Fig. 12c, d). The polynya was also closed again, mostly due to sea  
ice transport from the central Arctic.


With respect to the long-term climatology, SIC during HALO-( $\mathcal{AC}$ )<sup>3</sup> was mostly above the climatological mean in our  
325 central domain (Fig. 12d). The positive SIC anomaly was related to a further east expansion of the sea ice off Greenland, which  
overlaps well with the negative 2 m temperature anomalies shown in Fig. 7d, f. In contrast, the west coast of Svalbard used  
to be covered by ice in the climatology but was completely ice-free during HALO-( $\mathcal{AC}$ )<sup>3</sup>. Similarly, southwest of 75° N and  
0° E, as well as northeast and southeast of Svalbard, SIC anomalies were negative (Fig. 12e) in line with the strongest warm  
anomalies of the campaign.

330 Note that uncertainties of derived SIC in the marginal ice zone are especially large as a result of temporal and spatial  
interpolation (smearing uncertainty), and due to so-called weather filters. Weather filters remove the atmospheric contribution  
from the satellite signal to remove false sea ice in open-water regions but run the risk of removing true sea ice as well,  
especially in the MIZ. This mainly affects the central region where the estimated uncertainties of the OSI-SAF SIC product  
(total standard uncertainty) can reach up to 40% at the ice edge, mostly due to smearing uncertainty, and different satellite  
335 footprints at different frequency channels.

## 7 Summary

In this study, we analyzed the atmospheric and sea ice conditions during the HALO-( $\mathcal{AC}$ )<sup>3</sup> campaign, whose focus lies on  
studying transformation processes of air masses on their way into and out of the Arctic. The campaign took place in the North  
Atlantic sector of the Arctic from 07 March to 12 April 2022. Within this area, we defined three domains over which we  
340 investigate whether the conditions were well suited for the campaign’s research objectives. The central domain, which lies  
in the marginal sea ice zone west of Svalbard, represents the area with the highest research flight activity of the campaign.  
For our analysis, we relate the conditions during HALO-( $\mathcal{AC}$ )<sup>3</sup> to the long-term record of Ny-Ålesund measurements, ERA5  
reanalysis, and satellite-derived sea ice conditions.

The radiosonde and surface observations at Ny-Ålesund showed two main periods with different atmospheric conditions.  
345 The beginning of the campaign was dominated by extremely warm and moist air masses, resulting in new maximum 2 m tem-  
peratures, record-breaking precipitation and integrated water vapor (IWV) for March since the beginning of the measurements.  
The second half of the campaign was much colder and drier, with mostly clear sky conditions at Ny-Ålesund because of oro-  
graphic effects. Reanalyses data, which was shown to agree well with dropsonde measurements from HALO-( $\mathcal{AC}$ )<sup>3</sup>, confirmed  
these general conditions with northward transport of warm and moist air in the first part and a southward flow of cold Arctic air  
350 in the second part of the campaign. Based on this clear temporal change of conditions, we divided the campaign into a warm  
period (11–20 March) and a cold period (21 March–12 April) and analyzed both in detail.



**Figure 12.** Average sea ice concentration (SIC) from the MODIS–AMSR2 product at 1 km grid resolution and 10 m wind speed (in knots) from ERA5 for (a) 09–11 March (prior to the first AR event), (b) 14–16 March (during the second AR event), (c) 10–12 April (at the end of the campaign). Short (long) wind barbs represent wind speeds of 5 (10) knots. (d): SIC time series averaged over the central region for the campaign period, as well as the mean, and 10–90<sup>th</sup> percentiles of the 1979–2022 climatology. (e): SIC anomalies from the climatological mean averaged over the entire campaign period. Note that only differences larger than 5% are considered due to the uncertainties of the satellite product in the marginal ice zone. Data for (d) and (e) is from the OSI–SAF sea ice concentration climate data record.

During the warm period, an atmospheric blocking situation diverted cyclones northwards into the Arctic, resulting in strong northward heat and moisture transports (moist and warm air intrusions, WAIs). Two WAIs were identified as Atmospheric Rivers (ARs). At their northernmost positions, both ARs were stronger than 90% of all ARs in the latitudes of the central domain in the climatology. The ARs led to unusually high precipitation amounts at the coasts of Greenland and Svalbard. Over the sea ice northwest of Svalbard, the rainfall indicated by ERA5 was a record-breaking event for mid-March.

The cold period was initialized after the passage of a strong cyclone on 21 March. Northerly winds pulled cold and dry air from the central Arctic into the North Atlantic, resulting in marine cold air outbreaks (MCAOs). We identified three MCAO events, separated by negative values of the MCAO index  $M$ . The second MCAO event, which started at the beginning of

360 the cold period, lasted for 19 days, making it the tenth longest MCAO event in the climatology. The strength of the MCAO conditions in the central domain was related to the cyclone activity in the vicinity of this region.

Despite the longer cold period, the entire campaign was on average warmer than the climatological mean. Thus, the short-lived ARs and WAIs dominated the anomaly signal, highlighting the importance of these events for the warming of the Arctic (Johansson et al., 2017). However, it must be noted that the ERA5 climatology may have systematic differences in 2022  
365 compared to previous years as measurements from HALO-( $\mathcal{A}\mathcal{C}$ )<sup>3</sup> dropsondes were assimilated. Furthermore, the quality and quantity of satellite measurements to be included in the assimilation improved over the years. Nevertheless, due to the low density of observations in the Arctic, using reanalyses is currently a well suited option for climatological comparisons.

Interestingly, precipitation was higher than the climatology for both periods as MCAOs also led to enhanced precipitation in the Fram Strait. Precipitation associated with MCAOs has been discussed as being responsible for differences between climate  
370 models and observations (von Lerber et al., 2022). However, precipitation is also difficult to measure accurately as MCAOs are mostly within the blind zone of satellite measurements (Schirmacher et al., 2023). Therefore, the quality of precipitation in reanalyses is of high interest. The detailed HALO-( $\mathcal{A}\mathcal{C}$ )<sup>3</sup> remote sensing and in situ measurements can help to further constrain the representation of precipitation, including its phase, in the reanalysis.

Finally, we looked at the sea ice conditions and how they changed due to the different circulation patterns during the cam-  
375 paign. Overall, the marginal ice zone in the Fram Strait had higher sea ice concentrations (SIC) compared to the climatological mean, while the regions around Svalbard had lower SIC. WAIs and ARs resulted in a reduction of SIC in the marginal ice zone below the climatological mean. The reduction was probably dominated by ice dynamics associated with the strong southerly winds, but melt was also possible because of temperatures above freezing. During the long period of MCAO conditions, the northerly winds and low temperatures led to a recovery of the SIC.


380 Compared to previous aircraft campaigns in the Arctic within ( $\mathcal{A}\mathcal{C}$ )<sup>3</sup> that focused on the evolution of (mixed-phase) clouds (ACLOUD; AFLUX and MOSAiC-ACA, Wendisch et al., 2019; Mech et al., 2022), we observed a larger variety of MCAO conditions during HALO-( $\mathcal{A}\mathcal{C}$ )<sup>3</sup>. The long phase of MCAOs with varying strength and different wind regimes provides opportunities for detailed MCAO studies making use of airborne measurements. Also, the sea ice edge was closer to Svalbard than during AFLUX (March–April 2019) so that ocean–ice transects could be performed more easily. HALO-( $\mathcal{A}\mathcal{C}$ )<sup>3</sup> captured sev-  
385 eral WAIs / ARs with unusual or even record-breaking strength. With regard to the changing climate when exchanges between the mid–latitudes and the Arctic become more frequent, the campaign provides a unique opportunity to study stronger than average WAIs / ARs. Thus, the weather conditions were well suited to achieve the objectives of the HALO-( $\mathcal{A}\mathcal{C}$ )<sup>3</sup> campaign.

*Code and data availability.* All codes used for the analyses presented in this study have been published on Zenodo for public access (Mayer et al., 2024). Sea ice concentration climatology data is found on OSI SAF (2017) and Copernicus Climate Change Service (C3S) (2020). The  
390 high resolution sea ice concentration dataset used for 07 March–12 April 2022 is based on the product from the Institute of Environmental Physics, University of Bremen. The data is available at [https://seaice.uni-bremen.de/data/modis\\_amsr2](https://seaice.uni-bremen.de/data/modis_amsr2). ERA5 data on single and pressure levels can be accessed through Hersbach et al. (2018b) and Hersbach et al. (2018a). Ny-Ålesund radiosondes and ceilometer data have been

published on PANGAEA (Maturilli and Kayser, 2016, 2017; Maturilli, 2020b, 2022). The Ny-Ålesund precipitation data is provided by MET Norway and has been downloaded from <https://seklima.met.no> (last access 2023-11-30). Near-surface meteorology data from Ny-Ålesund is available on PANGAEA as well (Maturilli, 2020a). Radar measurements from HALO during HALO-( $\mathcal{A}C$ )<sup>3</sup> is available on PANGAEA (Dorff et al., 2023). Dropsonde measurements during HALO-( $\mathcal{A}C$ )<sup>3</sup> are currently processed and will be published on PANGAEA, searchable via the tag "HALO-( $\mathcal{A}C$ )<sup>3</sup>". A full description of the data set is in preparation.

## Appendix A: Integrated water vapor transport for moist and warm air intrusions

We show the total IVT from ERA5 for two snapshots of the ARs on 12–15 March, and for the weak WAI on 10 April in  
400 Fig. A1. While the ARs on 12–15 March were meridionally aligned, the WAI on 10 April had a rather zonal orientation. The  
strongest total IVT is found on 15 March, exceeding  $400 \text{ kg m}^{-1} \text{ s}^{-1}$ .




**Figure A1.** Total integrated water vapor transport (IVT, zonal and meridional component) from ERA5 (colours) and dropsondes (circular marker) for moist and warm air intrusions / Atmospheric Rivers: (a) 13 March 2022 16 UTC, (b) 15 March 2022 16 UTC, (c) 10 April 2022 14 UTC. Quivers indicate the flow direction and strength, while black line with white outline shows the HALO flight track. The orange (brown) line indicate the 15% (85%) sea ice concentration isoline. Grey dash-dotted contours show the mean sea level pressure. Data is based on ERA5.



## Appendix B: Pressure constellation during the warm and the cold period

To illustrate the contrast of the two periods, we show an Arctic-wide average pressure constellation for the warm and cold period  
405 in Fig. B1. The warm period featured this extended 500 hPa geopotential height trough and several low pressure systems over  
Greenland and the geopotential height ridge and high pressure over Scandinavia. During the cold period, strong highs over  
Greenland and the central Arctic persisted while low pressure systems were located over Scandinavia, Northern Russia or the  
Barents Sea.

(a) Warm period



(b) Cold period



**Figure B1.** Maps of mean sea level pressure (white contour lines with black outlines) and 500 hPa geopotential height (shading and black contour lines) from ERA5 data averaged over (a) the warm period and (11–20 March 2022) and (b) the cold period (21 March–12 April 2022). The 15% sea ice concentration from ERA5 is displayed as blue contour line. "H" and "L" illustrate the centres of high and low pressure systems, respectively. The measurement domains are displayed as purple boxes

*Author contributions.* SC, AE, IG, JanM, MW, and AW conceptualized the manuscript. MW and AW formulated the introduction with comments and input from AE, JanM, and SC. HD, BK, ML, NM, MM, JanR, and AW prepared the description of the data and methods chapter. SB, HD, BK, ML, JanM, HM, RN, FP, JohR, JanR, IS, NS, and AW provided visualisations and analysed figures. JohM and AW collected codes from co-authors and made them publicly available. AW is the main author of this manuscript and ensured validation. All co-authors reviewed the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We gratefully acknowledge the funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) for the ArctiC amplification: Climate Relevant Atmospheric and SurfaCe Processes, and Feedback Mechanisms ( $\mathcal{AC}$ )<sup>3</sup> Project Number 268020496 – TRR 172 within the Transregional Collaborative Research Center. We are further grateful for funding of project grant no. 316646266 by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) within the framework of Priority Programme SPP 1294 to promote research with HALO. We thank the Institute of Environmental Physics, University of Bremen for the provision of the merged MODIS-AMSR2 sea-ice concentration data at [https://seaice.uni-bremen.de/data/modis\\_amr2](https://seaice.uni-bremen.de/data/modis_amr2) (last access 2023-04-04). Hersbach et al. (2018a) and Hersbach et al. (2018b) were downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store. The results contain modified Copernicus Climate Change Service information 2022. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains. We thank the Alfred Wegener Institute for providing and operating the two aircraft (Polar 5 and Polar 6), the crew, and also the technicians of the Polar 5 and Polar 6 aircraft. We are also grateful for the research aircraft HALO, the pilots and technicians provided and operated by the German Aerospace Centre (Deutsches Zentrum für Luft- und Raumfahrt).



## References

- Avila-Diaz, A., Bromwich, D. H., Wilson, A. B., Justino, F., and Wang, S.-H.: Climate Extremes across the North American Arctic in Modern Reanalyses, *J. Climate*, 34, 2385–2410, <https://doi.org/10.1175/JCLI-D-20-0093.1>, 2021.
- 430 Bresson, H., Rinke, A., Mech, M., Reinert, D., Schemann, V., Ebell, K., Maturilli, M., Viceto, C., Gorodetskaya, I., and Crewell, S.: Case study of a moisture intrusion over the Arctic with the ICOSahedral Non-hydrostatic (ICON) model: resolution dependence of its representation, *Atmos. Chem. Phys.*, 22, 173–196, <https://doi.org/10.5194/acp-22-173-2022>, 2022.
- Cohen, J., Zhang, X., Francis, J., Jung, T., Kwok, R., Overland, J., Ballinger, T. J., Bhatt, U. S., Chen, H. W., Coumou, D., Feldstein, S., Gu, H., Handorf, D., Henderson, G., Ionita, M., Kretschmer, M., Laliberte, F., Lee, S., Linderholm, H. W., Maslowski, W., Peings, Y., Pfeiffer, K., Rigor, I., Semmler, T., Stroeve, J., Taylor, P. C., Vavrus, S., Vihma, T., Wang, S., Wendisch, M., Wu, Y., and  
435 Yoon, J.: Divergent consensus on Arctic amplification influence on midlatitude severe winter weather, *Nat. Clim. Change*, 10, 20–29, <https://doi.org/10.1038/s41558-019-0662-y>, 2020.
- Copernicus Climate Change Service (C3S): Sea ice concentration daily gridded data from 1979 to present derived from satellite observations, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/CDS.3CD8B812>, 2020.
- Dahlke, S., Solbès, A., and Maturilli, M.: Cold Air Outbreaks in Fram Strait: Climatology, Trends, and Observations During an Extreme  
440 Season in 2020, *J. Geophys. Res.: Atmos.*, 127, <https://doi.org/10.1029/2021JD035741>, 2022.
- Dorff, H., Aubry, C., Ewald, F., Hirsch, L., Jansen, F., Konow, H., Mech, M., Ori, D., Ringel, M., Walbröl, A., Crewell, S., Ehrlich, A., Wendisch, M., and Ament, F.: Unified Airborne Active and Passive Microwave Measurements over Arctic Sea Ice and Ocean during the HALO-(AC)<sup>3</sup> Campaign in Spring 2022, <https://doi.org/10.1594/PANGAEA.963250>, 2023.
- Fletcher, J., Mason, S., and Jakob, C.: The Climatology, Meteorology, and Boundary Layer Structure of Marine Cold Air Outbreaks in Both  
445 Hemispheres\*, *J. Climate*, 29, 1999–2014, <https://doi.org/10.1175/JCLI-D-15-0268.1>, 2016.
- Geerts, B., Giangrande, S. E., McFarquhar, G. M., Xue, L., Abel, S. J., Comstock, J. M., Crewell, S., DeMott, P. J., Ebell, K., Field, P., et al.: The comble campaign: A study of marine boundary layer clouds in arctic cold-air outbreaks, *Bulletin of the American Meteorological Society*, 103, E1371–E1389, 2022.
- George, G., Stevens, B., Bony, S., Pincus, R., Fairall, C., Schulz, H., Kölling, T., Kalen, Q. T., Klingebiel, M., Konow, H., Lundry, A., Prange, M., and Radtke, J.: JOANNE: Joint dropsonde Observations of the Atmosphere in tropical North Atlantic meso-scale Environments, *Earth  
450 Syst. Sci. Data*, 13, 5253–5272, <https://doi.org/10.5194/essd-13-5253-2021>, 2021.
- Gierens, R., Kneifel, S., Shupe, M. D., Ebell, K., Maturilli, M., and Löhnert, U.: Low-level mixed-phase clouds in a complex Arctic environment, *Atmospheric Chemistry and Physics*, 20, 3459–3481, <https://doi.org/10.5194/acp-20-3459-2020>, 2020.
- Graham, R. M., Cohen, L., Ritzhaupt, N., Segger, B., Graversen, R. G., Rinke, A., Walden, V. P., Granskog, M. A., and Hudson, S. R.: Evaluation of Six Atmospheric Reanalyses over Arctic Sea Ice from Winter to Early Summer, *J. Climate*, 32, 4121–4143,  
455 <https://doi.org/10.1175/JCLI-D-18-0643.1>, 2019a.
- Graham, R. M., Hudson, S. R., and Maturilli, M.: Improved Performance of ERA5 in Arctic Gateway Relative to Four Global Atmospheric Reanalyses, *Geophys. Res. Lett.*, 46, 6138–6147, <https://doi.org/10.1029/2019GL082781>, 2019b.
- Guan, B. and Waliser, D. E.: Detection of atmospheric rivers: Evaluation and application of an algorithm for global studies: Detection of  
460 Atmospheric Rivers, *J. Geophys. Res.: Atmos.*, 120, 514–12 535, <https://doi.org/10.1002/2015JD024257>, 2015.
- Guan, B. and Waliser, D. E.: Atmospheric rivers in 20 year weather and climate simulations: A multimodel, global evaluation, *J. Geophys. Res.: Atmos.*, 122, 5556–5581, <https://doi.org/10.1002/2016JD026174>, 2017.

- Guan, B., Waliser, D. E., and Ralph, F. M.: An Intercomparison between Reanalysis and Dropsonde Observations of the Total Water Vapor Transport in Individual Atmospheric Rivers, *J. Hydrometeorol.*, 19, 321–337, <https://doi.org/10.1175/JHM-D-17-0114.1>, 2018.
- 465 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on pressure levels from 1959 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.bd0915c6>, 2018a.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1959 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2018b.
- 470 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Q. J. R. Meteorolog. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hock, T. F. and Franklin, J. L.: The NCAR GPS Dropwindsonde, *Bull. Am. Meteorol. Soc.*, 80, 407–420, [https://doi.org/10.1175/1520-0477\(1999\)080<0407:TNGD>2.0.CO;2](https://doi.org/10.1175/1520-0477(1999)080<0407:TNGD>2.0.CO;2), 1999.
- Johansson, E., Devasthale, A., Tjernström, M., Ekman, A. M. L., and L'Ecuyer, T.: Response of the lower troposphere to moisture intrusions into the Arctic, *Geophys. Res. Lett.*, 44, 2527–2536, <https://doi.org/10.1002/2017GL072687>, 2017.
- 480 Kapsch, M.-L., Graversen, R. G., and Tjernström, M.: Springtime atmospheric energy transport and the control of Arctic summer sea-ice extent, *Nat. Clim. Change*, 3, 744–748, <https://doi.org/10.1038/nclimate1884>, 2013.
- Kapsch, M.-L., Skific, N., Graversen, R. G., Tjernström, M., and Francis, J. A.: Summers with low Arctic sea ice linked to persistence of spring atmospheric circulation patterns, *Clim. Dyn.*, 52, 2497–2512, <https://doi.org/10.1007/s00382-018-4279-z>, 2019.
- Knudsen, E. M., Heinold, B., Dahlke, S., Bozem, H., Crewell, S., Gorodetskaya, I. V., Heygster, G., Kunkel, D., Maturilli, M., Mech, M., Viceto, C., Rinke, A., Schmithüsen, H., Ehrlich, A., Macke, A., Lüpkes, C., and Wendisch, M.: Meteorological conditions during the ACLOUD/PASCAL field campaign near Svalbard in early summer 2017, *Atmospheric Chemistry and Physics*, 18, 17995–18022, <https://doi.org/10.5194/acp-18-17995-2018>, 2018.
- 485 Komatsu, K. K., Alexeev, V. A., Repina, I. A., and Tachibana, Y.: Poleward upgliding Siberian atmospheric rivers over sea ice heat up Arctic upper air, *Sci. Rep.*, 8, 2872, <https://doi.org/10.1038/s41598-018-21159-6>, 2018.
- 490 Konow, H., Ewald, F., George, G., Jacob, M., Klingebiel, M., Kölling, T., Luebke, A. E., Mieslinger, T., Pörtge, V., Radtke, J., Schäfer, M., Schulz, H., Vogel, R., Wirth, M., Bony, S., Crewell, S., Ehrlich, A., Forster, L., Giez, A., Göttsche, F., Groß, S., Gutleben, M., Hagen, M., Hirsch, L., Jansen, F., Lang, T., Mayer, B., Mech, M., Prange, M., Schnitt, S., Vial, J., Walbröl, A., Wendisch, M., Wolf, K., Zinner, T., Zöger, M., Ament, F., and Stevens, B.: EUREC<sup>4</sup>A's HALO, *Earth Syst. Sci. Data*, 13, 5545–5563, <https://doi.org/10.5194/essd-13-5545-2021>, 2021.
- 495 Lauer, M., Mech, M., and Guan, B.: Global Atmospheric Rivers catalog for ERA5 reanalysis, <https://doi.org/10.1594/PANGAEA.957161>, 2023.
- Lavergne, T., Sørensen, A. M., Kern, S., Tonboe, R., Notz, D., Aaboe, S., Bell, L., Dybkjær, G., Eastwood, S., Gabarro, C., Heygster, G., Killie, M. A., Brandt Kreiner, M., Lavelle, J., Saldo, R., Sandven, S., and Pedersen, L. T.: Version 2 of the EUMETSAT OSI SAF and ESA CCI sea-ice concentration climate data records, *The Cryosphere*, 13, 49–78, <https://doi.org/10.5194/tc-13-49-2019>, 2019.

- 500 Lloyd, G., Choulaton, T. W., Bower, K. N., Gallagher, M. W., Crosier, J., O'shea, S., Abel, S. J., Fox, S., Cotton, R., and Boutle, I. A.: In situ measurements of cloud microphysical and aerosol properties during the break-up of stratocumulus cloud layers in cold air outbreaks over the North Atlantic, *Atmospheric Chemistry and Physics*, 18, 17 191–17 206, 2018.
- Ludwig, V., Spreen, G., and Pedersen, L. T.: Evaluation of a New Merged Sea-Ice Concentration Dataset at 1 km Resolution from Thermal Infrared and Passive Microwave Satellite Data in the Arctic, *Remote Sens.*, 12, 3183, <https://doi.org/10.3390/rs12193183>, 2020.
- 505 Ma, W., Chen, G., and Guan, B.: Poleward Shift of Atmospheric Rivers in the Southern Hemisphere in Recent Decades, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2020GL089934>, 2020.
- Mattingly, K. S., Mote, T. L., and Fettweis, X.: Atmospheric River Impacts on Greenland Ice Sheet Surface Mass Balance, *J. Geophys. Res.: Atmos.*, 123, 8538–8560, <https://doi.org/10.1029/2018JD028714>, 2018.
- Mattingly, K. S., Mote, T. L., Fettweis, X., van As, D., Van Tricht, K., Lhermitte, S., Pettersen, C., and Fausto, R. S.: Strong Summer  
510 Atmospheric Rivers Trigger Greenland Ice Sheet Melt through Spatially Varying Surface Energy Balance and Cloud Regimes, *J. Climate*, 33, 6809–6832, <https://doi.org/10.1175/JCLI-D-19-0835.1>, 2020.
- Maturilli, M.: Continuous meteorological observations at station Ny-Ålesund (2011-08 et seq), <https://doi.org/10.1594/PANGAEA.914979>, 2020a.
- Maturilli, M.: High resolution radiosonde measurements from station Ny-Ålesund (2017-04 et seq), PANGAEA - Data Publisher for Earth  
515 & Environmental Science [data set], <https://doi.org/10.1594/PANGAEA.914973>, 2020b.
- Maturilli, M.: Ceilometer cloud base height from station Ny-Ålesund (2017-08 et seq), PANGAEA - Data Publisher for Earth & Environmental Science [data set], <https://doi.org/10.1594/PANGAEA.942331>, 2022.
- Maturilli, M. and Ebell, K.: Twenty-five years of cloud base height measurements by ceilometer in Ny-Ålesund, Svalbard, *Earth Syst. Sci. Data*, 10, 1451–1456, <https://doi.org/10.5194/essd-10-1451-2018>, 2018.
- 520 Maturilli, M. and Kayser, M.: Homogenized radiosonde record at station Ny-Ålesund, Spitsbergen, 1993-2014, PANGAEA - Data Publisher for Earth & Environmental Science [data set], <https://doi.org/10.1594/PANGAEA.845373>, 2016.
- Maturilli, M. and Kayser, M.: Homogenized radiosonde record at station Ny-Ålesund, Spitsbergen, 2015-2016, PANGAEA - Data Publisher for Earth & Environmental Science [data set], <https://doi.org/10.1594/PANGAEA.875196>, 2017.
- Maturilli, M., Herber, A., and König-Langlo, G.: Climatology and time series of surface meteorology in Ny-Ålesund, Svalbard, *Earth Syst. Sci. Data*, 5, 155–163, <https://doi.org/10.5194/essd-5-155-2013>, 2013.
- 525 Mayer, J., Walbröl, A., Becker, S., Dorff, H., Kirbus, B., Lauer, M., Müller, H., Rückert, J. E., and Slättberg, N.: Codes for: Contrasting extremely warm and long- lasting cold air anomalies in the North Atlantic sector of the Arctic during the HALO-(AC)3 campaign, <https://doi.org/10.5281/zenodo.11300722>, 2024.
- Mech, M., Ehrlich, A., Herber, A., Lüpkes, C., Wendisch, M., Becker, S., Boose, Y., Chechin, D., Crewell, S., Dupuy, R., Gourbeyre, C.,  
530 Hartmann, J., Jäkel, E., Jourdan, O., Kliesch, L.-L., Klingebiel, M., Kulla, B. S., Mioche, G., Moser, M., Risse, N., Ruiz-Donoso, E., Schäfer, M., Stapf, J., and Voigt, C.: MOSAiC-ACA and AFLUX - Arctic airborne campaigns characterizing the exit area of MOSAiC, *Sci. Data*, 9, 790, <https://doi.org/10.1038/s41597-022-01900-7>, 2022.
- Mewes, D. and Jacobi, C.: Heat transport pathways into the Arctic and their connections to surface air temperatures, *Atmos. Chem. Phys.*, 19, 3927–3937, <https://doi.org/10.5194/acp-19-3927-2019>, 2019.
- 535 Nash, D., Waliser, D., Guan, B., Ye, H., and Ralph, F. M.: The Role of Atmospheric Rivers in Extratropical and Polar Hydroclimate, *J. Geophys. Res.: Atmos.*, 123, 6804–6821, <https://doi.org/10.1029/2017JD028130>, 2018.

- Neff, W., Compo, G. P., Martin Ralph, F., and Shupe, M. D.: Continental heat anomalies and the extreme melting of the Greenland ice surface in 2012 and 1889: Melting of Greenland in 1889 and 2012, *J. Geophys. Res.: Atmos.*, 119, 6520–6536, <https://doi.org/10.1002/2014JD021470>, 2014.
- 540 Newell, R. E., Newell, N. E., Zhu, Y., and Scott, C.: Tropospheric rivers? - A pilot study, *Geophys. Res. Lett.*, 19, 2401–2404, <https://doi.org/10.1029/92GL02916>, 1992.
- OSI SAF: Global Sea Ice Concentration Climate Data Record v2.0 - Multimission, EUMETSAT SAF on Ocean and Sea Ice [data set], [https://doi.org/10.15770/EUM\\_SAF\\_OSI\\_0008](https://doi.org/10.15770/EUM_SAF_OSI_0008), 2017.
- Papritz, L. and Spengler, T.: A Lagrangian Climatology of Wintertime Cold Air Outbreaks in the Irminger and Nordic Seas and Their Role  
545 in Shaping Air–Sea Heat Fluxes, *J. Climate*, 30, 2717–2737, <https://doi.org/10.1175/JCLI-D-16-0605.1>, 2017.
- Papritz, L., Hauswirth, D., and Hartmuth, K.: Moisture origin, transport pathways, and driving processes of intense wintertime moisture transport into the Arctic, *Weather and Climate Dynamics*, 3, 1–20, <https://doi.org/10.5194/wcd-3-1-2022>, 2022.
- Pithan, F., Medeiros, B., and Mauritsen, T.: Mixed-phase clouds cause climate model biases in Arctic wintertime temperature inversions, *Clim. Dyn.*, 43, 289–303, <https://doi.org/10.1007/s00382-013-1964-9>, 2014.
- 550 Pithan, F., Svensson, G., Caballero, R., Chechin, D., Cronin, T. W., Ekman, A. M. L., Neggers, R., Shupe, M. D., Solomon, A., Tjernström, M., and Wendisch, M.: Role of air-mass transformations in exchange between the Arctic and mid-latitudes, *Nat. Geosci.*, 11, 805–812, <https://doi.org/10.1038/s41561-018-0234-1>, 2018.
- Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun. Earth Environ.*, 3, 168, <https://doi.org/10.1038/s43247-022-00498-3>,  
555 2022.
- Rinke, A., Segger, B., Crewell, S., Maturilli, M., Naakka, T., Nygård, T., Vihma, T., Alshawaf, F., Dick, G., Wickert, J., and Keller, J.: Trends of Vertically Integrated Water Vapor over the Arctic during 1979–2016: Consistent Moistening All Over?, *Journal of Climate*, 32, 6097–6116, <https://doi.org/10.1175/JCLI-D-19-0092.1>, 2019.
- Rückert, J. E., Rostosky, P., Huntemann, M., Clemens-Sewall, D., Ebell, K., Kaleschke, L., Lemmetyinen, J., Macfarlane, A. R., Naderpour, R., Stroeve, J., Walbröl, A., and Spreen, G.: Sea ice concentration satellite retrievals influenced by surface changes due to warm air intrusions: A case study from the MOSAiC expedition, *Elem. Sci. Anth.*, EarthArXiv [preprint], <https://doi.org/10.31223/X5VW85>, 2023-03-24, 2023.
- Schirmacher, I., Kollias, P., Lamer, K., Mech, M., Pfitzenmaier, L., Wendisch, M., and Crewell, S.: Assessing Arctic low-level clouds and precipitation from above – a radar perspective, *Atmospheric Measurement Techniques*, 16, 4081–4100, <https://doi.org/10.5194/amt-16-4081-2023>, 2023.  
565 4081-2023, 2023.
- Screen, J. A. and Simmonds, I.: The central role of diminishing sea ice in recent Arctic temperature amplification, *Nature*, 464, 1334–1337, <https://doi.org/10.1038/nature09051>, 2010.
- Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research synthesis, *Global Planet. Change*, 77, 85–96, <https://doi.org/10.1016/j.gloplacha.2011.03.004>, 2011.
- 570 Serreze, M. C. and Barry, R. G.: The Arctic Climate System, Cambridge Atmospheric and Space Science Series, Cambridge University Press, 2 edn., 2014.
- Serreze, M. C., Barrett, A. P., Stroeve, J. C., Kindig, D. N., and Holland, M. M.: The emergence of surface-based Arctic amplification, *The Cryosphere*, 3, 11–19, <https://doi.org/10.5194/tc-3-11-2009>, 2009.

Serreze, M. C., Gustafson, J., Barrett, A. P., Druckenmiller, M. L., Fox, S., Voveris, J., Stroeve, J., Sheffield, B., Forbes, B. C., Rasmus, S.,  
575 et al.: Arctic rain on snow events: bridging observations to understand environmental and livelihood impacts, *Environmental Research Letters*, 16, 105009, 2021.

Stevens, B., Ament, F., Bony, S., Crewell, S., Ewald, F., Gross, S., Hansen, A., Hirsch, L., Jacob, M., Kölling, T., Konow, H., Mayer, B.,  
Wendisch, M., Wirth, M., Wolf, K., Bakan, S., Bauer-Pfundstein, M., Brueck, M., Delanoë, J., Ehrlich, A., Farrell, D., Forde, M., Göttsche,  
F., Grob, H., Hagen, M., Jäkel, E., Jansen, F., Klepp, C., Klingebiel, M., Mech, M., Peters, G., Rapp, M., Wing, A. A., and Zinner, T.:  
580 A High-Altitude Long-Range Aircraft Configured as a Cloud Observatory: The NARVAL Expeditions, *Bull. Am. Meteorol. Soc.*, 100,  
1061–1077, <https://doi.org/10.1175/BAMS-D-18-0198.1>, 2019.

Stevens, B., Bony, S., Farrell, D., Ament, F., Blyth, A., Fairall, C., Karstensen, J., Quinn, P. K., Speich, S., Acquistapace, C., Aemisegger, F.,  
Albright, A. L., Bellenger, H., Bodenschatz, E., Caesar, K.-A., Chewitt-Lucas, R., De Boer, G., Delanoë, J., Denby, L., Ewald, F., Fildier,  
B., Forde, M., George, G., Gross, S., Hagen, M., Hausold, A., Heywood, K. J., Hirsch, L., Jacob, M., Jansen, F., Kinne, S., Klocke, D.,  
585 Kölling, T., Konow, H., Lathon, M., Mohr, W., Naumann, A. K., Nuijens, L., Olivier, L., Pincus, R., Pöhlker, M., Reverdin, G., Roberts,  
G., Schnitt, S., Schulz, H., Siebesma, A. P., Stephan, C. C., Sullivan, P., Touzé-Peiffer, L., Vial, J., Vogel, R., Zuidema, P., Alexander,  
N., Alves, L., Arixi, S., Asmath, H., Bagheri, G., Baier, K., Bailey, A., Baranowski, D., Baron, A., Barrau, S., Barrett, P. A., Batier,  
F., Behrendt, A., Bendinger, A., Beucher, F., Bigorre, S., Blades, E., Blossey, P., Bock, O., Böing, S., Bossler, P., Bourras, D., Bouruet-  
Aubertot, P., Bower, K., Branellec, P., Branger, H., Brennek, M., Brewer, A., Brilouet, P.-E., Brüggemann, B., Buehler, S. A., Burke, E.,  
590 Burton, R., Calmer, R., Canonici, J.-C., Carton, X., Cato Jr., G., Charles, J. A., Chazette, P., Chen, Y., Chilinski, M. T., Choulaton, T.,  
Chuang, P., Clarke, S., Coe, H., Cornet, C., Coutris, P., Couvreux, F., Crewell, S., Cronin, T., Cui, Z., Cuyper, Y., Daley, A., Damerell,  
G. M., Dauhut, T., Deneke, H., Desbios, J.-P., Dörner, S., Donner, S., Douet, V., Drushka, K., Dütsch, M., Ehrlich, A., Emanuel, K.,  
Emmanouilidis, A., Etienne, J.-C., Etienne-Leblanc, S., Faure, G., Feingold, G., Ferrero, L., Fix, A., Flamant, C., Flatau, P. J., Foltz,  
G. R., Forster, L., Furtuna, I., Gadian, A., Galewsky, J., Gallagher, M., Gallimore, P., Gaston, C., Gentemann, C., Geyskens, N., Giez, A.,  
595 Gollop, J., Gouirand, I., Gourbeyre, C., De Graaf, D., De Groot, G. E., Grosz, R., Güttler, J., Gutleben, M., Hall, K., Harris, G., Helfer,  
K. C., Henze, D., Herbert, C., Holanda, B., Ibanez-Landeta, A., Intrieri, J., Iyer, S., Julien, F., Kalesse, H., Kazil, J., Kellman, A., Kidane,  
A. T., Kirchner, U., Klingebiel, M., Körner, M., Kremper, L. A., Kretzschmar, J., Krüger, O., Kumala, W., Kurz, A., L'Hégaret, P., Labaste,  
M., Lachlan-Cope, T., Laing, A., Landschützer, P., Lang, T., Lange, D., Lange, I., Laplace, C., Lavik, G., Laxenaire, R., Le Bihan, C.,  
Leandro, M., Lefevre, N., Lena, M., Lenschow, D., Li, Q., Lloyd, G., Los, S., Losi, N., Lovell, O., Luneau, C., Makuch, P., Malinowski,  
600 S., Manta, G., Marinou, E., Marsden, N., Masson, S., Maury, N., Mayer, B., Mayers-Als, M., Mazel, C., McGeary, W., McWilliams, J. C.,  
Mech, M., Mehlmann, M., Meroni, A. N., Mieslinger, T., Minikin, A., Minnett, P., Möller, G., Morfa Avalos, Y., Muller, C., Musat, I.,  
Napoli, A., Neuberger, A., Noisel, C., Noone, D., Nordsiek, F., Nowak, J. L., Oswald, L., Parker, D. J., Peck, C., Person, R., Philippi, M.,  
Plueddemann, A., Pöhlker, C., Pörtge, V., Pöschl, U., Pologne, L., Posylniak, M., Prange, M., Quiñones Meléndez, E., Radtke, J., Ramage,  
K., Reimann, J., Renault, L., Reus, K., Reyes, A., Ribbe, J., Ringel, M., Ritschel, M., Rocha, C. B., Rochetin, N., Röttenbacher, J., Rollo,  
605 C., Royer, H., Sadoulet, P., Saffin, L., Sandiford, S., Sandu, I., Schäfer, M., Schemann, V., Schirmacher, I., Schlenczek, O., Schmidt, J.,  
Schröder, M., Schwarzenboeck, A., Sealy, A., Senff, C. J., Serikov, I., Shohan, S., Siddle, E., Smirnov, A., Späth, F., Spooner, B., Stolla,  
M. K., Szkółka, W., De Szoek, S. P., Tarot, S., Tetoni, E., Thompson, E., Thomson, J., Tomassini, L., Totems, J., Ubele, A. A., Villiger,  
L., Von Arx, J., Wagner, T., Walther, A., Webber, B., Wendisch, M., Whitehall, S., Wiltshire, A., Wing, A. A., Wirth, M., Wiskandt, J.,  
Wolf, K., Worbes, L., Wright, E., Wulfmeyer, V., Young, S., Zhang, C., Zhang, D., Ziemann, F., Zinner, T., and Zöger, M.: EUREC4A,  
610 Earth System Science Data, 13, 4067–4119, <https://doi.org/10.5194/essd-13-4067-2021>, 2021.

- Stroeve, J., Nandan, V., Willatt, R., Dadic, R., Rostosky, P., Gallagher, M., Mallett, R., Barrett, A., Hendricks, S., Tonboe, R., McCrystall, M., Serreze, M., Thielke, L., Spreen, G., Newman, T., Yackel, J., Ricker, R., Tsamados, M., Macfarlane, A., Hannula, H.-R., and Schneebeli, M.: Rain on snow (ROS) understudied in sea ice remote sensing: a multi-sensor analysis of ROS during MOSAiC (Multidisciplinary drifting Observatory for the Study of Arctic Climate), *The Cryosphere*, 16, 4223–4250, <https://doi.org/10.5194/tc-16-4223-2022>, 2022.
- 615 Tjernström, M., Shupe, M. D., Brooks, I. M., Persson, P. O. G., Prytherch, J., Salisbury, D. J., Sedlar, J., Achtert, P., Brooks, B. J., Johnston, P. E., Sotiropoulou, G., and Wolfe, D.: Warm-air advection, air mass transformation and fog causes rapid ice melt: WARM-AIR ADVECTION, FOG AND ICE MELT, *Geophys. Res. Lett.*, 42, 5594–5602, <https://doi.org/10.1002/2015GL064373>, 2015.
- Viceto, C., Gorodetskaya, I. V., Rinke, A., Maturilli, M., Rocha, A., and Crewell, S.: Atmospheric rivers and associated precipitation patterns during the ALOUD and PASCAL campaigns near Svalbard (May–June 2017): case studies using observations, reanalyses, and a regional climate model, *Atmos. Chem. Phys.*, 22, 441–463, <https://doi.org/10.5194/acp-22-441-2022>, 2022.
- 620 von Lerber, A., Mech, M., Rinke, A., Zhang, D., Lauer, M., Radovan, A., Gorodetskaya, I., and Crewell, S.: Evaluating seasonal and regional distribution of snowfall in regional climate model simulations in the Arctic, *Atmospheric Chemistry and Physics*, 22, 7287–7317, <https://doi.org/10.5194/acp-22-7287-2022>, 2022.
- Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Dethloff, K., Velasco, C. B., Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Engler, C., Eppers, O., Gehrman, M., Gong, X., Gottschalk, M., Gourbeyre, C., Griesche, H., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L., Leroy, D., Maturilli, M., Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T., Notholt, J., Palm, M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder, K., Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P.,
- 630 Shupe, M. D., Siebert, H., Spreen, G., Stapf, J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M., and Zeppenfeld, S.: The Arctic Cloud Puzzle: Using ALOUD/PASCAL Multiplatform Observations to Unravel the Role of Clouds and Aerosol Particles in Arctic Amplification, *Bull. Am. Meteorol. Soc.*, 100, 841–871, <https://doi.org/10.1175/BAMS-D-18-0072.1>, 2019.
- Wendisch, M., Handorf, D., Tegen, I., Neggers, R., and Spreen, G.: Glimpsing the Ins and Outs of the Arctic Atmospheric Cauldron, *Eos*, 102, <https://doi.org/10.1029/2021EO155959>, 2021.
- 635 Wendisch, M., Brückner, M., Crewell, S., Ehrlich, A., Notholt, J., Lüpkes, C., Macke, A., Burrows, J. P., Rinke, A., Quaas, J., Maturilli, M., Schemann, V., Shupe, M. D., Akansu, E. F., Barrientos-Velasco, C., Bärfuss, K., Blechschmidt, A.-M., Block, K., Bougoudis, I., Bozem, H., Böckmann, C., Bracher, A., Bresson, H., Bretschneider, L., Buschmann, M., Chechin, D. G., Chylik, J., Dahlke, S., Deneke, H., Dethloff, K., Donth, T., Dorn, W., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Eppers, O., Gerdes, R., Gierens, R., Gorodetskaya, I. V., Gottschalk, M., Griesche, H., Gryanik, V. M., Handorf, D., Harm-Altstädter, B., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Höschel, I., Hofmann, Z., Hölemann, J., Hünnerbein, A., Jafariserajehlou, S., Jäkel, E., Jacobi, C., Janout, M., Jansen, F., Jourdan, O., Jurányi, Z., Kalesse-Los, H., Kanzow, T., Käthner, R., Kliesch, L. L., Klingebiel, M., Knudsen, E. M., Kovács, T., Körtke, W., Krampe, D., Kretzschmar, J., Kreyling, D., Kulla, B., Kunkel, D., Lampert, A., Lauer, M., Lelli, L., von Lerber, A., Linke, O., Löhnert, U., Lonardi, M., Losa, S. N., Losch, M., Maahn, M., Mech, M., Mei, L., Mertes, S., Metzner, E., Mewes, D., Michaelis, J., Mioche, G., Moser, M., Nakoudi, K., Neggers, R., Neuber, R., Nomokonova, T., Oelker, J., Papakonstantinou-Presvelou, I., Pätzold, F., Pefanis, V., Pohl, C., van Pinxteren, M., Radovan, A., Rhein, M., Rex, M., Richter, A., Risse, N., Ritter, C., Rostosky, P., Rozanov, V. V., Donoso, E. R., Saavedra Garfias, P., Salzmann, M., Schacht, J., Schäfer, M., Schneider, J., Schnierstein, N., Seifert, P., Seo, S., Siebert, H., Soppa, M. A., Spreen, G., Stachlewska, I. S., Stapf, J., Stratmann, F., Tegen, I., Viceto, C., Voigt, C., Vountas, M., Walbröl, A., Walter, M., Wehner, B., Wex, H., Willmes, S., Zanatta, M., and Zeppenfeld, S.: Atmospheric and Surface Processes, and Feedback

- Mechanisms Determining Arctic Amplification: A Review of First Results and Prospects of the (AC)<sup>3</sup> Project, *Bull. Am. Meteorol. Soc.*, 650 104, E208–E242, <https://doi.org/10.1175/BAMS-D-21-0218.1>, 2023.
- Wendisch, M., Crewell, S., Ehrlich, A., Herber, A., Kirbus, B., Lüpkes, C., Mech, M., Abel, S. J., Akansu, E. F., Ament, F., Aubry, C., Becker, S., Borrmann, S., Bozem, H., Brückner, M., Clemen, H.-C., Dahlke, S., Dekoutsidis, G., Delanoë, J., De La Torre Castro, E., Dorff, H., Dupuy, R., Eppers, O., Ewald, F., George, G., Gorodetskaya, I. V., Grawe, S., Groß, S., Hartmann, J., Henning, S., Hirsch, L., Jäkel, E., Joppe, P., Jourdan, O., Jurányi, Z., Karalis, M., Kellermann, M., Klingebiel, M., Lonardi, M., Lucke, J., Luebke, A., Maahn, M., Maherndl, 655 N., Maturilli, M., Mayer, B., Mayer, J., Mertes, S., Michaelis, J., Michalkov, M., Mioche, G., Moser, M., Müller, H., Neggers, R., Ori, D., Paul, D., Paulus, F., Pilz, C., Pithan, F., Pöhlker, M., Pörtge, V., Ringel, M., Risse, N., Roberts, G. C., Rosenburg, S., Röttenbacher, J., Rückert, J., Schäfer, M., Schäfer, J., Schemann, V., Schirmacher, I., Schmidt, J., Schmidt, S., Schneider, J., Schnitt, S., Schwarz, A., Siebert, H., Sodemann, H., Sperzel, T., Spreen, G., Stevens, B., Stratmann, F., Svensson, G., Tatzelt, C., Tuch, T., Vihma, T., Voigt, C., Volkmer, L., Walbröl, A., Weber, A., Wehner, B., Wetzel, B., Wirth, M., and Zinner, T.: Overview: Quasi-Lagrangian observations 660 of Arctic air mass transformations – Introduction and initial results of the HALO–(AC)<sup>3</sup> aircraft campaign, *EGUsphere*, 2024, 1–46, <https://doi.org/10.5194/egusphere-2024-783>, 2024.
- Wesche, C., Steinhage, D., and Nixdorf, U.: Polar aircraft Polar5 and Polar6 operated by the Alfred Wegener Institute, *JLSRF*, 2, A87, <https://doi.org/10.17815/jlsrf-2-153>, 2016.
- Woods, C. and Caballero, R.: The Role of Moist Intrusions in Winter Arctic Warming and Sea Ice Decline, *J. Climate*, 29, 4473–4485, 665 <https://doi.org/10.1175/JCLI-D-15-0773.1>, 2016.
- Woods, C., Caballero, R., and Svensson, G.: Large-scale circulation associated with moisture intrusions into the Arctic during winter: Moisture Intrusions during Arctic Winter, *Geophys. Res. Lett.*, 40, 4717–4721, <https://doi.org/10.1002/grl.50912>, 2013.
- You, C., Tjernström, M., Devasthale, A., and Steinfeld, D.: The Role of Atmospheric Blocking in Regulating Arctic Warming, *Geophys. Res. Lett.*, 49, <https://doi.org/10.1029/2022GL097899>, 2022.
- 670 Yu, Y., Xiao, W., Zhang, Z., Cheng, X., Hui, F., and Zhao, J.: Evaluation of 2-m Air Temperature and Surface Temperature from ERA5 and ERA-I Using Buoy Observations in the Arctic during 2010–2020, *Remote Sens.*, 13, 2813, <https://doi.org/10.3390/rs13142813>, 2021.
- Ziereis, H. and Gläßer, M.: HALO - Global Player für die Atmosphärenforschung, Deutsches Zentrum für Luft- und Raumfahrt (DLR), <https://elib.dlr.de/51508/>, 2006.