



Modeling atmosphere-land interactions at a rainforest site - a case study using Amazon Tall Tower Observatory (ATTO) measurements and reanalysis data

Amelie U. Schmitt^{1,2}, Felix Ament¹, Alessandro C. de Araújo^{3,4}, Marta Sá⁴, and Paulo Teixeira⁴

¹Meteorological Institute, Center for Earth System Research and Sustainability (CEN), Universität Hamburg, Hamburg, Germany

²Climate Service Center Germany (GERICS), Helmholtz-Zentrum Hereon, Hamburg, Germany

³Empresa Brasileira de Pesquisa Agropecuária (EMBRAPA), Belém, Brazil

⁴Large Scale Biosphere-Atmosphere Experiment in Amazonia (LBA), Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil

Correspondence: Amelie Schmitt (amelie.schmitt@hereon.de)

Abstract. Modeling the interactions between atmosphere and soil at a forest site remains a challenging task. Using tower measurements from the Amazon Tall Tower Observatory (ATTO) in the rainforest, we evaluated the performance of the land model JSBACH focusing especially on processes influenced by the forest canopy.

As a first step, we analyzed whether ERA5 and MERRA-2 reanalysis data are suitable to be used as land model forcing. Comparing five years of ATTO measurements to near-surface reanalysis data, we found a substantial underestimation of wind speeds by about 1 m s^{-1} . ERA5 captures monthly mean temperatures quite well but overestimates annual mean precipitation by 30%. Contrarily, MERRA-2 overestimates monthly mean temperatures in the dry season (August - October) by more than 1 K, while mean precipitation biases are small.

To test how much the choice of reanalysis data set and the reanalysis biases affect the results of the land model we performed spin-up and model runs using either ERA5 or MERRA-2 and with and without a bias correction for precipitation and wind speed and compared the results. The choice of reanalysis data set results in large differences of up to 1.3 K for soil temperatures and 20% for soil water content, which are non-negligible especially in the first weeks after spin-up. Correcting wind speed and precipitation biases also notably changes the land model results - especially in the dry season.

Based on these results, we constructed an optimized forcing data set using bias-corrected ERA5 data for the spin-up period and ATTO measurements for a model run of two years and comparing the results to observations to identify model shortcomings. Generally, the shape of the soil water profile is not reproduced correctly, which might be related to a lack of vertical variability of soil properties or of the root density. The model also shows a positive soil temperature bias and overestimates the penetration depth of the diurnal cycle. This problem could possibly be addressed by including a separate canopy layer into the model to improve the processes related to storage and vertical transport of energy within the model.



20 1 Introduction

The presence of vegetation - especially in forest canopies with tall trees - alters the exchange processes between land surface and atmosphere. Forests influence the shape of the wind profile (Yi, 2008; de Souza et al., 2016; Santana et al., 2017) and the structure of turbulence within and above the canopy layer (Chor et al., 2017; Dias-Júnior et al., 2019; Zahn et al., 2016). In tall forests, the air close to the ground can even become decoupled from the air above the forest (Santana et al., 2018). Additional
25 heat storage in trees and the air within the canopy changes the surface energy balance (e.g. Oliphant et al., 2004; Lindroth et al., 2010) and the transmission of radiation is affected by the leaf area and the canopy structure (e.g. Hardy et al., 2004). Forests also have an impact on global carbon and water cycles. Evapotranspiration is particularly enhanced in rainforests (de Oliveira et al., 2018; Costa et al., 2010), which leads to a moistening of the atmosphere and increased cloud cover (Wright et al., 2017).

These processes at the canopy scale are very complex and thus difficult to represent in coupled atmosphere-land models. A
30 few land surface models (LSMs) - such as CLM (Bonan et al., 2018) or ORCHIDEE (Chen et al., 2016), for example - have implemented parameterizations of vertically resolved processes in forest canopies in the past years. However, many LSMs still incorporate rather simple parameterizations of canopy effects. The aim of this study is to evaluate the performance of an LSM without a resolved canopy layer in a region with a tall forest canopy. For this purpose, we choose the land model JSBACH and compare model results to measurements from the Amazon Tall Tower Observatory (ATTO) located at a rainforest site.

Observations from tropical forests have been previously used to evaluate specific aspects of LSMs, for example to test
35 different parameterizations of rainfall intercepted by forest canopies (Wang et al., 2007; Fan et al., 2019). Anwar et al. (2022) used measurements from different FLUXNET sites in the Amazon rainforest to test the performance of two land-surface hydrology schemes in the CLM model. They found that both schemes were able to reproduce the shape of the seasonal cycle of turbulent fluxes but failed in capturing their correct magnitude. Other studies used measurements from canopies in the
40 mid-latitudes - which are dominated by deciduous broadleaf and evergreen needleleaf forests - for example to evaluate the performance of multi-layer canopy representations in LSMs (Ma and Liu, 2019; Bonan et al., 2021). Studies using specifically the JSBACH model have focused on calibrating stomatal conductance using FLUXNET data from evergreen needleleaf forests (Mäkelä et al., 2019) or on the global evaluation of a canopy heat storage parametrisation (Heidkamp et al., 2018). In this study, we use a site-level setup of the JSBACH LSM to evaluate the model performance at a rainforest site with a special focus on
45 canopy processes.

LSMs are not only used in coupled model setups but can also be run offline using external data as atmospheric forcing. Feedbacks between land and atmosphere in coupled models make it more difficult to assess single processes and thus it is beneficial to use offline simulations to isolate processes when evaluating LSMs (e.g. Decharme et al., 2019). In addition, uncertainties of the LSM originating from errors the atmospheric model can be avoided by carefully choosing a forcing data
50 set. Besides observations, reanalysis data are frequently used for offline and site-level simulations, which aim at testing the model's performance with respect to different parameterizations (Brun et al., 2013; Knauer et al., 2017) or initialisation data sets (Ardilouze et al., 2017), for example. To minimize the impact of forcing data on the model results, some studies also use bias corrected reanalysis data sets, such as the WFDEI meteorological forcing data set (Weedon et al., 2014). When



evaluating the global performance of the ISBA land model, Decharme et al. (2019) used atmospheric forcings based two
55 different reanalysis data sets. The results showed large differences with respect to hydrological variables, which underlines the
importance of the choice of forcing data sets for the evaluation of land surface models.

After initialization, an LSM undergoes an adjustment process until an equilibrium between external forcing and the simulated
land surface fluxes is reached. The length of this so-called spin-up period can reach several years Yang et al. (1995) but requires
shorter time spans for locations with large precipitation amounts (Lim et al., 2012; Yang et al., 2011). Here, we use a spin-up
60 period of ten years following the example of Heidkamp et al. (2018) who evaluated land surface fluxes in the LSM JSBACH.
For our JSBACH site-level simulations we also intend to partly use reanalysis data since ATTO measurements contain many
data gaps. We use two different high-resolution global reanalyses (ERA5 and MERRA-2) to compare and minimize the impact
of the forcing data set on the model results. As a first step, we compare near-surface meteorological data reanalyses to ATTO
measurements to answer the following questions: How well do the reanalyses reproduce meteorological conditions - ranging
65 from hourly to yearly scales - at the ATTO site? And are they thus generally suitable as forcing data sets for the land model?
How does the choice of forcing data set affect the results of the land model? For this purpose, we compare model results using
the two different reanalyses as forcing for the model spin-up run. Also, we test how a correction of reanalysis biases changes
the model results. Based on the findings from these preparative analyses we perform a model run using an optimized forcing
data set (consisting of air temperature and humidity, wind speed, precipitation and incoming long- and shortwave radiation)
70 and compare the results to ATTO measurements focusing on the following questions: How well does JSBACH reproduce the
temporal evolution of the model output variables soil water content, soil temperatures and turbulent fluxes? Can we identify
specific model shortcoming, which could be improved in future model versions?

Observations from the ATTO site, data from the ERA5 and MERRA-2 reanalyses and the JSBACH land model are described
in Sect. 2. Results of the comparison between reanalysis data and ATTO measurements are presented in Sect. . Section con-
75 tains results of JSBACH model runs used for sensitivity studies (Sect. 3.2.1 and 3.2.2) and to identify model shortcomings
(Sect. 3.2.3), followed by a summary and conclusions in Sect. 4.

2 Data and model description

2.1 Amazon Tall Tower Observatory

2.1.1 Measurements at the ATTO site

80 The Amazon Tall Tower Observatory (ATTO) is a scientific research facility in Brazil with a focus on interactions between
the rainforest and the atmosphere. An extensive description of the characteristics of the ATTO site can be found in Andreae
et al. (2015). Here, we summarize the most important features. The ATTO site is located in the central Amazon at an altitude
of 130 m, roughly 150 km north-east of Manaus (Fig. 1). The area is covered by terra firme forest with an average canopy
height of about 37.5 m. In this study, we use measurements from the 81 m high walk-up tower, which is located at 2.144° S
85 and 59.002° W.




Figure 1. Location of the ATTO site (blue triangle) and reanalysis grid boxes used for comparison in this study. The background shows GLCC land cover types derived using the BATS scheme for 1992 data (Loveland et al., 2000).

We use five years of ATTO measurements from 2014 to 2018. For an overview of the measurements used in this study see Table 1. Incoming shortwave and longwave radiation, precipitation, pressure, as well as turbulent sensible and latent heat fluxes are measured at the top part of the tower. Air temperature, relative humidity and wind speed are measured at several heights above and within the forest canopy. Since the reanalyses used for comparison in Sect. 3.1 do not have a separate resolved canopy layer, we use only measurements above the canopy top (≤ 36 m) for the comparison. Additionally, air temperatures within the canopy at 1.5 m above the ground are used to calculate temperature differences between the soil and atmosphere (see Sect. 3.2.3). Measurements of soil moisture and temperature are conducted close to the walk-up tower and are available at several depths (see Table 1).

2.1.2 Data preparation

For comparison with reanalysis data and for model forcing ATTO measurements are aggregated to hourly values. We apply a linear interpolation to fill small data gaps of up to 10 minutes for temperature, humidity, wind, pressure and radiation and up to 1 minute for precipitation. Larger data gaps were caused by interruptions of the power supply or by general sensors issues.



Table 1. Observations at the ATTO site used in this study.

Variables	Instruments	Measurement heights (m)	Accuracy
Air temperature and relative humidity	Termohygrometer (CS215, Rotronic Measurement Solutions,UK)	1.5, 36.3, 40.2, 55.3	Temperature: 0.3 °C (at 25 °C), 0.4 °C (5 to 40 °C); relative humidity: 4 % (0-100 % range), 2 % (10-90 % range) at 25 °C
Wind speed and direction	2-D sonic anemometer (WindSonic, Gill Instruments Ltd., UK)	43.1*, 50.8, 66.0, 73.7	Wind speed uncertainty: 2 % at 12 m s ⁻¹
Turbulent sensible and latent heat flux	Wind, temperature: 2014-2017: 3D Sonic Anemometer (WindMaster, Gill Instruments Ltd., UK); 2017-2018: 3D Sonic Anemometer (CSAT3, Campbell Scientific, Inc., USA) Humidity: Infrared Gas Analyzer (IRGA LI-7500A, LI-COR Inc., USA)	82.0	WindMaster: wind speed: <1.5 % RMS at 12 m s ⁻¹ CSAT3: wind speed <2 % (wind vector within ±5° of horizontal) temperature: 0.01 K H ₂ O content: <2 %
Precipitation	Rain gauge (TB4, Hydrological Services Pty. Ltd., Australia)	81.3	Rainfall per tip: 0.254 mm; uncertainty: 2 % (<250 mm h ⁻¹), 3 % (250-500 mm h ⁻¹)
Atmospheric pressure	Barometer (PTB101B, Vaisala, Finland)	81.0	0.5 hPa at 20 °C; 1.5 hPa (0-40 °C)
Shortwave radiation	Pyranometer (CMP21, Kipp & Zonen, Netherlands)	75.6	Expected daily uncertainty <2 %
Longwave radiation	Pyrgeometer (CGR4, Kipp & Zonen, Netherlands)	75.6	Uncertainty <3 % for daily totals
Soil water content	Water content reflectometer (CS615, Campbell Scientific Inc., USA)	-0.1, -0.2, -0.3, -0.4, -0.6, -1.0	2 % manufacturers standard calibration
Soil temperature	Thermistor (108, Campbell Scientific Inc., USA)	-0.1, -0.2, -0.4	0.2 K

* wind measurements at 43.1 m height are only available for the years 2014 and 2015

To account for these gaps we mask out times when any of the considered ATTO data are missing in the reanalysis data as well (Sect. 3.1). Wind measurements at 43.1 m height are only available for the years 2014 and 2015 and were therefore not used for the general statistical analysis.



For comparison of the humidity with reanalysis data we convert the relative humidity RH to specific humidity q with

$$q = \frac{0.622 \text{ Pa}}{p} \left(\frac{RH}{100\%} e_{\text{sat}}(\vartheta) \right), \quad (1)$$

where p is the air pressure in Pa and e_{sat} is the saturation water vapor pressure calculated using the Magnus equation with ϑ being the air temperature in °C. Since air pressure is only measured at 81 m height, we extrapolate the values downward to the measurement heights of humidity using the barometric formula:

$$p_2 = p_1 \cdot \exp \left(-\frac{g (z_1 - z_2)}{R_l T_v} \right). \quad (2)$$

p_i is the pressure at height z_i , $g = 9.81 \text{ ms}^{-1}$ is the gravity of Earth, $R_l = 287.1 \text{ Jkg}^{-1}\text{K}^{-1}$ is the specific gas constant for dry air and the virtual temperature is calculated as $T_v = T(1 + 0.61q)$. Air pressure is calculated stepwise downwards and if data required for the calculations are missing at the considered height, values are replaced by applying Eq. 2 to the levels above.

110 2.1.3 Seasonality

Being located at around 2° S, the climate at ATTO is tropical humid and strongly influenced by the location of the Inter-Tropical Convergence Zone, leading to pronounced wet and dry seasons. While Andreae et al. (2015) strictly divide the year into a wet and a dry season, Saturno et al. (2018) also consider transition zones between the seasons.

In this study we calculate seasonal means and thus aim for relatively homogeneous meteorological conditions within a season. Average annual cycles of air temperature, incoming shortwave radiation and precipitation are shown in Fig. 2. For our purposes, we define the wet season as the months from January to April, which are characterized by large precipitation sums exceeding 200 mm per month as well as monthly mean temperatures and shortwave radiation below the annual means. Accordingly, the dry season is defined as the months from August to October with precipitation sums below 100 mm per month. Air temperatures in the dry season are on average more than 2 K higher than in the wet season. The wind speed shows no distinct annual cycle and seasonal changes are small. Due to the latitude there are only small variations of less than 15 minutes of the day length, which facilitates the interpretation and comparison of average diurnal cycles.

2.2 Reanalysis data

2.2.1 ERA5

ERA5 is the fifth generation reanalysis produced by ECMWF based on version Cy41r2 of the IFS model (Hersbach et al., 2020). Model output is available every hour on a 0.25°x 0.25° grid with 137 vertical levels. Mean total precipitation rate, incoming longwave and shortwave radiation, sensible and latent heat flux from the ERA5 surface data set (Hersbach et al., 2018) and air temperature, specific humidity, horizontal wind components, and pressure from ERA5 on model levels (Hersbach et al., 2017) for the years 2007 to 2018 were downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store. Since ATTO is located close to the border of two intersecting ERA5 grid boxes (see Fig. 1), we average the values of those grid boxes using inverse distance squared weighting.




Figure 2. Average annual cycles based on ATTO measurements of the years 2014 to 2018. (a) Mean temperature and average daily maximum and minimum temperatures measured at 36 m height (blue) and incoming shortwave radiation (red). (b) Precipitation (dark green) and fraction of missing data for all parameters displayed in this figure (bars). Shaded areas denote the wet (blue) and dry (yellow) seasons.

The IFS model calculates exchange processes at the land surface using the Tiled ECMWF Scheme for Surface Exchanges over Land with revised Hydrology component (HTESSEL; e.g. van den Hurk et al., 2000; Balsamo et al., 2009). Each grid box consists of separate tiles - e.g. for water, bare ground, high and low vegetation (ECMWF, 2016b). The tile fractions are based on the USGS Global Land Cover Characteristics (GLCC) data set with a 1-km resolution, which is derived from one year of remote sensing data from the Advanced Very High Resolution Radiometer (AVHRR) in 1992 (Loveland et al., 2000). The GLCC data are shown as background in Fig. 1. The grid boxes used for comparison with ATTO measurements are mostly covered by evergreen broadleaf trees with fractions of 97.5 % and 99.6 % for the northern and southern grid box, respectively.

We calculate the geometrical height of the model levels using the surface pressure, as well as temperature and specific humidity on model levels according to Eq. 2.22 of the IFS model documentation (ECMWF, 2016a). For the study period and location the heights of the lowest model levels are approximately 10, 32 and 57 m. Some of the ERA5 variables are provided



as instantaneous values. For comparison with ATTO and MERRA-2, we average two successive values to obtain an estimate of the hourly mean.

2.2.2 MERRA-2

MERRA-2 is the second version of the Modern-Era Retrospective analysis for Research and Applications (Gelaro et al., 2017) produced using the GEOS-5 atmospheric general circulation model by GMAO (NASA's Global Modeling and Assimilation Office). Model output is available every hour on a 0.5° -latitude and 0.625° -longitude grid with 72 vertical levels. The following variables from MERRA-2 for the years 2007 to 2018 were obtained from Global Modeling and Assimilation Office (GMAO) (2015): Incoming longwave and shortwave radiation; sensible and latent heat flux; pressure; air temperature, specific humidity and horizontal wind components at 10 and 50 m height; bias corrected precipitation.

The location of the MERRA-2 model grid box used for comparison with ATTO measurements is displayed in Fig. 1. As for ERA5, MERRA-2 land cover is based on the GLCC data set (Loveland et al., 2000). We calculate the pressure at model levels using the pressure thickness variable and apply the same method as for ERA5 to calculate the geometrical height of the model levels. For the study period and location the height of the lowest model levels is approximately 68 m.

To improve the soil moisture calculations in MERRA-2 the model-generated precipitation is corrected using observations (Reichle et al., 2017). In the ATTO region, gauge-based precipitation estimates from the CPCU data set (Unified Gauge-Based Analysis of Global Daily Precipitation) are used to replace the model precipitation. However, the number of rain gauges in the Amazon region has decreased in the last decades, resulting in an uncertainty increase of the CPCU data set (Reichle et al., 2017).

2.3 Modeling setup

2.3.1 The land surface model JSBACH

We use the land surface model JSBACH (Giorgetta et al., 2013) to simulate fluxes of energy, water and momentum between the land surface and the atmosphere. We perform our model simulations using JSBACH4, which is a more flexible re-implementation of JSBACH3 and can be used within the global models MPI-ESM1.2 (Mauritsen et al., 2019) or ICON-A (Giorgetta et al., 2018; Jungclaus et al., 2022), as well as a stand-alone model forced by external data (e.g. Nabel et al., 2020).

For comparison with point measurements at ATTO, we use a site-level setup of JSBACH consisting of a single grid box. The required meteorological variables for model forcing are air temperature, specific or relative humidity, precipitation, wind speed, as well as incoming longwave and shortwave radiation. We use ATTO measurements, as well as reanalysis data from ERA5 and MERRA-2 as forcing data sets to obtain the results in Sect. 3.2. The forcing is applied at each model time step of 1 hour.

JSBACH uses a fractional approach to represent vegetation classes. Each grid cell is divided into tiles with 11 different vegetation classes. The grid box used for the model simulations at the ATTO location contains 5.6% lake and river areas and the remaining land fraction is covered by 99.3% tropical evergreen trees with a root depth of 1.95 m. The soil depth at this grid

box is 2.23 m, covering the upper 4 of 5 model layers. These characteristics are based on the data set of land surface parameters derived by Hagemann (2002) on a T63 grid, which corresponds to a grid cell size of about 200 km at the considered latitude.

175 Vertical soil water transport within the model is calculated using the one-dimensional Richards equation (see e.g. Ekici et al., 2014, , Eq. 2), which incorporates vertical diffusion, percolation from gravitational drainage and sources and sinks. The only depth-dependent sink term is transpiration, which depends on the root density. The surface energy balance in JSBACH is closed within the uppermost soil layer - meaning that the soil temperature between 0 and 6.5 cm is considered as the surface temperature.

180 **2.3.2 Preparation of model forcing data**

We perform two separate 10-year model spin-up runs using ERA5 data from the lowest model level of about 10 m height and MERRA-2 data from 10 m height of the years 2007 to 2016. After the spin-up, we perform model runs for the years 2017 and 2018 using forcing data from either ERA5, MERRA-2, or ATTO measurements from 4 m (temperature and specific humidity), 13 m (wind), and roughly 40 m (radiation and precipitation) above the forest top.

185 ATTO measurements contain data gaps that need to be filled before the data can be used as model forcing. We apply the following methods: 1) Small gaps of up to 4 hours (2 hours for precipitation and shortwave radiation) are filled by linear interpolation. 2) Medium sized gaps of up to 6 days are filled with data from the day before and after the gap. Only days with at least 12 hours of valid measurements are used for filling. 3) Large gaps are filled with a monthly mean diurnal cycle, which we calculate by averaging only days without missing values for each calendar month of the years 2017 and 2018. Time periods
190 during which missing data have been interpolated are masked out in the interpretation of the model results in Sect. 3.2.

3 Results and discussion

3.1 Comparison of reanalysis data to ATTO measurements

To evaluate whether reanalysis data are suitable forcing data sets for a land model we compare ATTO measurements closely above the forest top to reanalysis data at the corresponding heights. For this purpose, ATTO measurement heights are specified
195 as heights above forest (a.f.) relative to the forest top of about 37.5 m height. Here, we focus on the forcing data required for JSBACH, which are air temperature, specific humidity, wind speed, precipitation, and incoming shortwave and longwave radiation. As described in Sect. 2.1.2 hours with measurement gaps are also masked out in the reanalysis data sets.

As a first step, we consider annual mean temperatures and precipitation sums, which are widely used to characterize local climates. ERA5 annual mean temperatures at 10 m height between 2014 and 2018 agree almost perfectly with the ATTO values
200 of 26.1 °C measured at 18 m above the forest. MERRA-2 is generally too warm with annual average temperatures of 26.9 °C. For annual precipitation sums we see the opposite, with very good agreement between ATTO (1560 mm) and MERRA-2 (1540 mm) and a strong overestimation of ERA5 with 2030 mm per year. A more detailed comparison of average annual and




Figure 3. Monthly means of the years 2014 to 2018 for (a) air temperature, (b) specific humidity, (c) wind speed, (d) precipitation, (e) incoming shortwave radiation and (f) incoming longwave radiation. Red and yellow lines are values at 10 m height or at the surface (sfc) derived from ERA5 and MERRA-2 reanalysis data, respectively. Blue lines denote ATTO measurements at different heights above the forest top (a.f.) with measurement uncertainties represented as shading.

diurnal cycles is presented in Sect. 3.1.1 for wind, in Sect. 3.1.2 for precipitation and radiation and in Sect. 3.1.3 for temperature and humidity.

205 3.1.1 Wind

The most apparent feature of the wind speed annual cycle is the large underestimation by both reanalysis data sets (Fig. 3c). Averaged over the years 2014 to 2018 ERA5 shows a negative bias of about -0.8 ms^{-1} and MERRA-2 an even larger one of -1.2 ms^{-1} . The bias only shows small changes for larger heights up to 36 m above the forest (Fig 4a) and does not change significantly during the course of the day (Fig 4b). The shape of the diurnal cycle is well reproduced by both reanalyses, while
210 ERA5 does a better job a capturing the shape of the annual cycle.

Our results indicate large biases at heights close to the forest canopy and could thus simply be related to the shape of the wind profile at these heights. To analyze whether the bias extends to larger heights, we examined measurements from the top of the ATTO tall tower at 285 m above the forest top, which is available from the ATTO data portal (attodata.org) for the time




Figure 4. Profiles (a) and diurnal cycles (b) of wind speed averaged over the years 2014 to 2018 based on ERA5 (red) and MERRA-2 (yellow) reanalysis data and ATTO measurements (blue). Measurement uncertainty is denoted as blue errorbars and shading, respectively. The grey line in (a) shows a profile with additional heights, including wind measurements at 6 m above the forest, which are only available for the years 2014 and 2015. Gray vertical lines in (b) mark the times of sunrise and sunset, respectively.

period from March 2018 to the end of 2019. At this higher altitude we also found negative biases in the order of -1 ms^{-1} for both reanalyses (not shown), indicating that wind speed biases are a significant issue within the lower parts of the ABL in this region.

Wind speed biases of the same order of magnitude have also been found in various previous studies. Jourdier (2020) compared ERA5 and MERRA-2 wind speeds to measurements between 55 m and 100 m height at eight locations in France. They found that ERA5 wind speeds were generally about 1 to 1.5 ms^{-1} lower than those from MERRA-2. ERA5 wind speeds agreed better with measurements from the northern flat terrain, while those from MERRA-2 agreed better in southern moun-



tainous regions. The dependence of the sign of the wind speed bias on the region has also been found in a study by Staffell and Pfenninger (2016). Their results for MERRA-2 indicate a line dividing Europe with negative wind speed biases in the Mediterranean region and positive biases around the North and Baltic Seas. Carvalho (2019) compared MERRA-2 to wind speed measurements all over the globe and also found that the sign of the 10-year mean bias over land varied for different regions. In the northern Amazon region, the found a general underestimation of about -1 ms^{-1} , which is in line with our results. Gualtieri (2021) compared mast measurements to ERA5 wind speeds and found large negative biases of up to -3 ms^{-1} at mountain sites and a positive bias in a forest site. They concluded that reanalyses with a spatial resolution of several tens of kilometers have problems accurately reproducing wind speeds at sites with high variation of topography and land use.

Using these reanalysis data, which underestimate the observed wind speed, to force a land model we expect to see an impact on turbulent fluxes and thus also on related soil quantities.

3.1.2 Precipitation and radiation

Both ERA5 and MERRA-2 are able to reproduce the seasonality of precipitation and incoming shortwave radiation (Fig. 3d,e). The amplitude of the annual cycle of longwave radiation is small and biases for both reanalyses are within the ATTO measurement uncertainty (Fig. 3f). Precipitation sums from ERA5 are too large in all seasons with an overestimation of about 20 % in the wet season and of 45 % in the dry season. This is in line with a general underestimation of incoming shortwave radiation indicating exaggerated cloud formation. MERRA-2 captures monthly precipitation sums quite well, which can be attributed to the correction of precipitation biases over land within the model using a gauge-based data set (see Sect. 2.2.2).

The diurnal cycle of precipitation measured at ATTO in the wet season (Fig. 5a) shows two distinct maxima - one shortly after sunrise and one in the afternoon around the time of the temperature maximum. A similar shape can be found in the dry season but with much smaller amplitudes (not shown). The afternoon peak in continental rainfall is related to the diurnal cycle of solar forcing with land surface heating causing an ABL growth and the formation of convective clouds (see e.g. Yang and Smith, 2006). MERRA-2 captures the amplitude of the afternoon peak quite well, while ERA5 overestimates the maximum rain rate by about 40 %. ERA5 also shows a negative shortwave radiation bias during the day (Fig. 5b), which indicates an overestimation of the cloud cover.

The early morning precipitation peak can be observed more frequently over ocean regions. A common explanation is that nighttime cooling of the cloud-top causes a thermal destabilisation of the upper cloud, which then increases convection and precipitation (see e.g. Randall et al., 1991). MERRA-2 fails to reproduce this early morning peak. ERA5 does produce a secondary rainfall maximum, however it occurs earlier in the night between 2:00 and 6:00 local time. To further analyze the occurrence of the early morning peak we evaluated rainfall data from the IMERG data set in a larger region around ATTO (for details see Appendix A). Our results indicate that nighttime precipitation occurs more frequently in the region north-east of ATTO, while precipitation after sunrise between 6:00 and 8:00 is more patchy. The location of the morning maximum is also quite variable between the considered years (not shown). This gives a hint that MERRA-2 does not necessarily fail to reproduce early morning precipitation in general but it might just produce it at the wrong location. Further evidence can be




Figure 5. Average diurnal cycles for (a) rain rate and (b) incoming shortwave radiation bias. The curves represent hourly means from the wet season (JFMA) of the years 2014 to 2018. Shaded areas denote the ATTO measurement uncertainty. Gray vertical lines mark the times of sunrise and sunset, respectively.

found in a study by Sato et al. (2009) who showed that the timing of local precipitation maxima strongly depends on the model resolution.

3.1.3 Temperature and humidity

Monthly mean annual cycles of near-surface air temperature and specific humidity based on ATTO measurements and reanalysis data are shown in Fig. 3a and b, respectively. Generally, ERA5 reproduces the shape of both annual cycles well and the temperature bias is negligible. Specific humidity is underestimated in all months by roughly 1 g kg⁻¹. MERRA-2 shows a good agreement with only small biases in the wet season, while model results in the dry season are too warm (+1.1 K) and too dry (-0.9 g kg⁻¹).




Figure 6. Average diurnal cycles for (a) air temperature and (b-c) specific humidity closely above the forest top. The curves represent hourly means from the dry (a-b) and the wet (c) seasons of the years 2014 to 2018. Shaded areas denote the ATTO measurement uncertainties. Gray vertical lines mark the times of sunrise and sunset, respectively.

It is evident from the diurnal cycles in Fig. 6a that the positive temperature bias for MERRA-2 in the dry season is mainly driven by too high daily maximum temperatures. MERRA-2 maximum temperatures exceed the measurements by about 2.5 K, while nighttime temperatures agree well with measurements. Also, MERRA-2 maximum temperatures in the wet season show a much better agreement with ATTO measurements (not shown), which is in line with the smaller differences of monthly means (Fig. 3a) in this season. This overestimation of the maximum temperatures could be related to an underestimation of

265



cloud cover by MERRA-2. However, incoming shortwave radiation from MERRA-2 does not show a considerable bias in the dry season. Another hypothesis is that the bias of daily maximum temperature is related to the vertical structure of the ABL, i.e. vertical mixing and stability. Testing this hypothesis, however, would require more investigations with measurements spanning
270 the whole ABL column, e.g. from radiosonde measurements.

The diurnal cycles of specific humidity (Fig. 6b) show distinctly different shapes, with a maximum of the measured humidity in the later morning while the humidity of both reanalyses decreases during the morning with a minimum in the early afternoon. We found similar discrepancies also during the wet season (Fig. 6c). The processes leading to a humidity minimum during
275 midday have been described in detail by Brümmer et al. (2012), for example. Evaporation starting in the morning leads to an increase of the absolute humidity. However, when the atmospheric boundary layer (ABL) grows during the day, entrainment and downward-mixing of dryer air from above cause a decrease in humidity. These mixing processes lead to a humidity minimum that coincides with the time of the temperature maximum.

According to this process explanation the differing shape of the observed diurnal cycle of humidity could have two possible reasons: 1) evapotranspiration is stronger than modeled by the two reanalyses, or 2) vertical mixing-processes within the ABL
280 are weaker than in the reanalyses. To test the first hypothesis we analysed the diurnal cycles of latent heat fluxes (not shown) and found that during the dry season, maximum measured latent heat fluxes were indeed larger than those modeled by the two reanalyses. However, during the wet season ERA5 underestimates the measured fluxes while MERRA-2 shows a strong overestimation of the latent heat fluxes. Since the two reanalyses do not agree on the sign of the flux bias, it is unlikely that the different shapes of the humidity diurnal cycles are mainly related to evapotranspiration.

The shape of the humidity diurnal cycles is also related to the ABL growth. It is possible that vertical mixing within the ABL
285 or entrainment at the ABL top in the reanalyses are too strong and thus the ABL grows too quickly in the morning. Testing this hypothesis would require measurements spanning the whole boundary layer depth. Dias-Júnior et al. (2022) compared ABL heights from ERA5 to those derived from radiosonde and ceilometer measurements at ATTO in November 2015. Their findings indicate that the ERA5 ABL grows faster in the morning and is larger than the measured values after 9:00 local time.
290 The average maximum ABL height from ERA5 exceeds the measured values by more than 200 m. We checked MERRA-2 ABL heights for the corresponding periods and found an almost identical ABL growth rate in the morning with an even larger maximum exceeding ERA5 by about 200 m. These results are well in line with our second hypothesis and thus observed humidity biased are probably related to the growth rates of the ABL.

3.2 JSBACH model simulations

295 The comparison of near-surface atmospheric variables in the previous section revealed a mostly good agreement between reanalysis data and measurements at the ATTO site but also notable biases for certain variables. In this section, we examine the impact of these biases on JSBACH model results if reanalysis data are used for model spin-up (Sect. 3.2.1) or as forcing of the subsequent model run (Sect. 3.2.2). Based on the conclusions of these two sections we then set up a model run with optimized forcing. By comparing model results to ATTO measurements of soil variables and surface fluxes, we aim to identify possible
300 model shortcomings (Sect. 3.2.3).



3.2.1 Impact of the choice of spin-up data set

Land models require a spin-up period of several years to reach an equilibrium state for the soil water content. Since on-site measurements are not always available for such an extended time period, a solution is to use output from other atmospheric models or reanalysis data as spin-up forcing. Here, we evaluate how the choice of spin-up data set affects the subsequent model run. For this purpose, we perform two separate model experiments with a 10 year spin-up of the JSBACH model from to 2007 to 2016 using either data from the ERA5 or the MERRA-2 reanalyses as spin-up forcing. For the following model run we use ATTO data from 2017 and 2018 as model forcing. Since the impact of the spin-up data set might vary by season, we choose two separate starting times for the model run - one in the wet season starting in January 2017 and one in the dry season starting in September 2017.

As expected, the largest differences occur in the deeper soil layers. During the first week of the model run soil temperature differences are 0.4 K in the layer between 0.07 and 0.32 m depth and even 1.3 K at about 2 m depth (Fig. 7a). While the temperature differences in the upper layers decay within a few months, there is still a difference of 0.6 K in the deepest layer after 12 months of model run time.

For soil water, differences are also largest in the deeper layers with values exceeding 5 % between 0.32 and 1.23 m and of up to 20 % at about 2 m depth in the first month after spin-up (Fig. 7d). Differences for soil water and also for sensible and latent fluxes (Fig. 7c) decrease much quicker than for soil temperature and reach negligible values after about two months. When the model run is started in the dry season, there is a longer impact on soil water and fluxes of about six months (not shown). This rather quick decay of the soil water differences is probably specific for the ATTO site, where a relatively shallow soil layer and large precipitation sums in the wet season often cause a saturation of the soil layers with water. Consequently, this finding cannot a priori be transferred to other sites and model grid points.

As a second step, we aim to evaluate whether the size of the biases induced by the different spin-up data sets are relevant compared to general model biases. We quantify the model biases by comparing model runs with an optimized spin-up data set (for details see Sect. 3.2.3) to measurements from the ATTO site. The relative bias is then calculated as the fraction between spin-up bias and model bias. For both soil temperature and water (Fig. 7b,e) the choice of the spin-up data set has a larger impact in the wet season than in the dry season. For soil temperature, values are larger in the layer around 0.7 m depth with spin-up biases in the wet season amounting to up to 75 % of the model bias in the first month after spin-up and of still 20 % after seven months. In the layer above at around 0.2 m spin-up biases are only relevant in the first month.

Model biases for soil water in the wet season are slightly distorted since the soil is often saturated with water (see also Sect. 3.2.3). Results are more meaningful in the dry season, where spin-up biases amount to up to 25 % of the observed model biases. Thus, especially for shorter model runs of only a few days or weeks, the choice of spin-up data set has a notable impact on the model results and is not negligible.




Figure 7. Time series of differences of JSBACH model results using ATTO measurements as forcing for the model run and either ERA5 or MERRA-2 data for spin-up for (a) soil temperature, (d) soil water content at different depths and (c) RMSD for turbulent fluxes. Corresponding differences between ERA5 and MERRA-2 spin-up relative to differences between model results and measured ATTO data are shown in panels (b) for soil temperature and (e) for soil water for two different starting times of the model run after spin-up in the wet (S1: January 2017) and dry (S2: September 2017) seasons. Empty symbols in (b) and (e) indicate that differences between ERA5 and MERRA-2 spin-up are below 0.2 K for soil temperature or below 0.7 % for soil water content.

3.2.2 Sensitivity to wind speed and precipitation biases

The comparisons in Sect. 3.1 revealed significant biases between reanalysis data and ATTO measurements. Here, we quantify the impact of these biases on the JSBACH model results when reanalysis data are used as model forcing. We found the largest annual mean biases for MERRA-2 wind speed and ERA5 precipitation. For the sensitivity tests, we set up a JSBACH model run with a 10-year spin-up period and a model run of one year in 2017. For wind, we compare model results using the original MERRA-2 forcing to those with an added offset of the annual mean wind speed bias of -1.2 ms^{-1} between 2014 and 2018. In a second sensitivity study, we use a factor to adjust the ERA5 maximum annual precipitation bias of 44 % and compare



the results to those using original ERA5 forcing. Figure 8 shows the monthly mean differences between the "original" and the
 340 "adjusted" model results.




Figure 8. Monthly mean differences between the original and the adjusted JSBACH model runs for the MERRA-2 wind (top) and ERA5 precipitation (bottom) sensitivity runs. Panels (a) and (d) show root mean squared differences (RMSD) for sensible (SH) and latent (LH) heat fluxes and biases of net turbulent fluxes. Differences for soil water content (b,d) and soil temperatures (c,e) are presented for different vertical soil layers of the model.

The wind sensitivity runs show the largest soil water differences in the dry season. In the original run, monthly mean soil water content in all depths is overestimated by up to 5 % (Fig. 8b) and in the upper two layers daily mean soil water content is overestimated by up to 10 % (not shown). Soil temperatures of the original run are also slightly too high in the dry season (Fig. 8c). These differences are also related to the turbulent exchange at the surface. Correction of the wind speed bias results in
 345 an increase of the forcing wind speed in the adjusted run, which causes an increase of the absolute value of both turbulent latent and sensible heat fluxes. However, in the dry season these flux changes counteract each other and the resulting net turbulent flux bias is close to zero (Fig. 8a). In the wet season, the largest differences occur for soil temperatures down to a depth of 1.23 m. The original run overestimates monthly mean soil temperatures by about 0.5 K (Fig. 8c) and daily mean temperatures by up to 1.2 K in the upper two layers (not shown). Other than in the dry season, an increased wind speed in the adjusted run an the
 350 resulting stronger evaporation does not decrease the soil water content since the soil is mostly saturated with water in the wet season. As a result, changed latent and sensible heat fluxes do not fully counteract each other resulting in an underestimation of the net turbulent heat fluxes by about $4 W m^{-2}$ by the original run.

For the precipitation sensitivity run all changes in the wet season are negligible. This means that - even by decreasing precipitation sums by 44 % in the adjusted run - the soil is still mostly saturated with water. Changes are more notable in the



355 dry season. The original run overestimates the monthly mean soil water content in the upper two layers by up to 5 % (Fig. 8e),
which is the same order of magnitude as in the wind sensitivity run. An overestimated soil water content results in increased
latent heat fluxes, which in turn cool the surface. This is evident from an underestimation of the soil temperatures in the upper
three layers by up to 0.3 K (Fig. 8f) and also from the differences of the net turbulent heat fluxes in the dry season by about
2 W m⁻². Overall, these sensitivity studies demonstrate that biases of the forcing data sets can have a non-negligible impact
360 on the model results - at the ATTO site especially in the dry season - and should be corrected if possible.

3.2.3 Comparison of model results to ATTO measurements

In this section, we compare ATTO measurements to model results to identify possible shortcomings of the JSBACH land
model. Based on the results of the previous sections we construct an optimized version of the spin-up run with the following
characteristics: We use ERA5 because for most forcing variables the shape of the annual cycles agree better with measurements.
365 Wind speed bias (-0.8 ms⁻¹) and mean precipitation bias (+32%) are corrected and 9 years of model spin-up are performed
(2007-2015). The last spin-up year is performed using ATTO data. Two years of model results forced by ATTO data are
then compared to measurements. By using this setup, we can minimize the impact of the choice of spin-up forcing data set
on the results and thus focus mainly on uncertainties caused by model physics. For the comparisons, missing values of the
measurement data are also flagged in the model results. We evaluate the model performance based on average annual cycles of
370 soil water and temperature and seasonally averaged soil profiles and turbulent fluxes.

ATTO soil measurements are only available for certain model soil layers (see also Tab. 1). Therefore, we compare model
results from the second (0.07 to 0.32 m) and third (0.32 - 1.23 m) soil layers and average the corresponding ATTO measure-
ments, which are 0.1 to 0.3 m and 0.4 to 1.0 m, respectively. Please note that soil water measurements reach deeper than soil
temperature measurements, which are only available at 0.1, 0.2 and 0.4 m. Figure 9 shows annual cycles of soil water content
375 and temperature averaged over the two years of the model run from 2017 to 2018. It is evident that soil temperatures are over-
estimated in the model in all months and for both considered depths (Fig. 9b) with large annual mean biases of -1.1 K in the
upper and -1.3 K in the lower layer.

To identify possible reasons for this bias we analyzed temperature differences between the soil and the air above. The
land model is forced by air temperatures measured roughly 4 m above the forest canopy. It is evident from Fig. 9c that soil
380 temperatures in the model are always higher than air temperatures with a difference of about 0.6 K, which remains almost
constant throughout the year. In contrast, measured soil temperatures are always lower than air temperatures 4 m above the
forest with larger differences in the dry season. However, in reality soil temperature depend more likely on air temperatures
close to the ground within the canopy layer than on temperatures above the forest. We thus also analyzed differences between
ATTO soil temperatures and air temperatures measured at 1.5 m above the ground. Here, we find a much better agreement with
385 the modeled temperature differences throughout the year and even an almost perfect match in the wet season. This indicates
that the observed bias between measured and modeled soil temperatures is likely caused by a cooling effect within the canopy
layer, which is not included in the land model - such as evaporation of rain from stems and leaves.




Figure 9. Annual cycles of soil temperatures (a) and soil water content (b) averaged over the years 2017 and 2018 from ATTO measurements and the JSBACH model. Solid lines represent the second and dashed lines the third model soil layer and the corresponding ATTO measurements height, respectively. (c) Annual cycles of temperature differences between soil layers L1 in (b) and air temperatures measured about 4 m above the forest (a.f.) or within the forest canopy at 1.5 m above the ground (a.g.). Empty symbols represent months with more than 30 % of missing data. Shading denotes the measurement uncertainty.

In contrast, differences between modeled and measured soil water content show larger seasonal dependence (Fig. 9a). In the upper layer, the model overestimates the soil water content with an almost constant monthly bias of about 11 %. In the lower layer, modeled soil water agrees very well with measurements in the wetter season from February to July. However, in the dry months from September to December the model underestimates soil water content - also by about 11 %. The fact that different soil layers exhibit different bias signs hints towards problems of vertical soil water transport in the model. To analyze vertical dynamics more deeply we look at soil profiles in the following.



Figure 10 shows boxplots of daily mean soil temperature and water content for different depths. For soil water content, 395 ATTO measurements show very similar profile shapes for the wet and dry seasons, with lower values in the upper soil layers and higher values in deeper layers. The minimum and maximum can be found at 0.2 m and 0.6 m depth, respectively. However, the JSBACH model produces different profiles in the wet and dry seasons. In the wet season, all soil layers approach a constant value of about 35 % (Fig. 10a), which is close to the field capacity and thus the soil is mostly saturated with water. In the dry season, the maximum soil water content is found in the uppermost two layers until 0.32 m, which is contrasting the 400 measurements, for which a minimum is found in these depths (Fig. 10b).

A possible reason for the discrepancies of the soil water profiles between model and measurements in both seasons could be an overestimation of the soil water storage capacity in the upper two layers. This could either mean that boundary conditions of the soil properties do not represent the actual conditions at ATTO properly. Or - since JSBACH prescribes the same soil properties in all model layers - the soil at the ATTO site might actually be heterogeneous and consist of different soil types in 405 different depths. Another relevant process could be transpiration, which is a sink term in the soil water balance equation and is constant with depth in JSBACH. However, it is well known that root density - and thus water extraction from the soil by plants - varies with depth (e.g. Feddes, 1982; Perrochet, 1987). Improving the representation of roots in JSBACH (based on common approaches used in other land models, see e.g. Zheng and Wang (2007)) would potentially change the soil water profile, which we suggest should be investigated further in the future.

410 Profiles of soil temperature also show larger seasonal differences in the model results than in the observations. In the wet season, the shapes of the profiles agree reasonably well, with slightly increasing temperatures with depth both in the model and observations (Fig. 10c) and biases below 1 K. However, in the dry season, model temperatures are higher in the upper two layers until 0.32 m than below, while the measured temperature profile has on average nearly constant temperatures (Fig. 10d) with biases exceeding 1.5 K in all depths. The whiskers of the boxplots (10 % and 90 % percentiles) also indicate that the 415 day-to-day variability is overestimated in the model.

To get more insight into the temperature variability with depth we look at average diurnal cycles of soil temperature. Since the changes of the overall characteristics with seasons are small, we only present annual mean diurnal cycles in Fig. 11a. We focus on model layer 2 between 0.07 and 0.32 m for which observation depths show the best match. We would expect the best agreement with the observations at 0.2 m depth. However, the amplitude of the diurnal cycle is strongly overestimated and the 420 model agrees better with the amplitude measured at 0.1 m depth (Fig. 11b).

Typically, diurnal and annual soil temperatures have a sinusoidal shape with decreasing amplitude with depth and a phase shift, both depending on the heat conductivity of the soil. Our results indicate that the penetration depth of the diurnal cycle is overestimated in the model, which might be related to an overestimation of the heat conductivity of the soil. This is also in line with our findings of an overestimated soil water content (see above) since wet soil has a larger heat conductivity than dry soil. 425 To further test this hypothesis, we make use of the formula describing the exponentially decreasing soil temperature amplitude A with depth (see e.g. Moene and Dam, 2014):

$$A(z) = A_0 \exp\left(-\frac{z}{d}\right), \quad (3)$$




Figure 10. Vertical profiles of soil water content (a,b) and soil temperatures (c,d) during the wet (left) and dry seasons (right) in 2017 and 2018. Vertical lines denote the median values of daily means for the respective heights and boxplots - showing quartiles (box) and 10%- and 90%-percentiles (whiskers) - are centered at the ATTO measurement depths (blue) and the center of the JSBACH model layer (purple).

where A_0 is the amplitude at the surface and d is the damping depth, which is a function of the heat conductivity of the soil. Using the amplitudes of the diurnal temperature variations from the two uppermost layers z_1 and z_2 , we can determine the damping depth as:

$$d = \frac{z_2 - z_1}{\ln(A(z_1)/A(z_2))}. \quad (4)$$

For both the ATTO measurements (at 0.1 and 0.2 m) and the model results (layers centered at 0.03 and 0.19 m) we obtain a damping depth of about 7 cm. This good agreement indicates that properties concerning heat transport in the soil are well represented in the model, which counteracts the above hypothesis.

The overestimation of the penetration depth could also be related to the amplitude of the temperature at the soil surface A_0 . Using the value of $d = 0.07$ m, we obtain $A_0 = 3.0$ K using ATTO measurements at 0.1 m depth but a much larger value of $A_0 = 8.5$ K using model results from the uppermost model layer. This gives a hint that the forest canopy dampens the soil surface temperature. Since JSBACH does not include a canopy layer explicitly, the model is not able to capture this effect.




Figure 11. (a) Diurnal cycle of soil temperatures averaged over the years 2017 and 2018 in the second JSBACH model layer and the corresponding ATTO measurement depths. (b) Corresponding amplitudes and times of minimum and maximum of the curves in (a).

As a final step, we compared measured turbulent heat fluxes to those from the model. Two major aspects had to be considered before the comparison. First, time series of measured turbulent fluxes contain many gaps - mostly due to rain - and in 2017 and 2018 there is not a single day with full data coverage. Since the model is forced with ATTO measurements, rain occurs at the same time in the model and in reality and those cases can therefore be masked simultaneously. It needs to be kept in mind that data availability is much lower during the day than at night (Fig. 13a,b) and thus the diurnal cycles should be interpreted with care. To obtain more representative monthly means we first calculated average diurnal cycles for each month, which were then averaged to obtain monthly values. By doing so, we reduce the impact of a higher data availability at night on the overall monthly values. The second issue concerns the height differences between the ATTO measurements at about 44 m above the forest and the model results at the surface. In a well mixed atmospheric boundary layer, turbulent fluxes typically decrease linearly with height with a zero crossing at the height of the boundary layer depth. To get a rough estimate, we calculated a height correction factor for the measurements using boundary layer heights from ERA5 and MERRA-2. However,




Figure 12. Annual cycles of turbulent sensible (SH) and latent (LH) heat fluxes (a) and net fluxes (b) averaged over the years 2017 and 2018 from ATTO measurements and the JSBACH model. A positive sign denotes upward fluxes from the soil to the atmosphere.

450 this correction only had a notable impact on a few days but did not change the overall conclusions drawn from the analysis of flux differences between model and measurements and thus, the corrected results are not shown here.

Figure 12 shows average annual cycles of sensible (SH) and latent (LH) heat fluxes and the sums of the two. It is evident that measured LH fluxes are by a factor of about three to four larger than SH fluxes. Generally, modeled and measured fluxes agree quite well. In the dry season, LH fluxes are slightly overestimated by the model, which is however counteracted by underestimated SH fluxes and thus net fluxes show a good agreement.

455

Diurnal cycles of turbulent heat fluxes are presented in Fig. 13. It is notable that LH fluxes in the model show a time shift compared with the measurements with fluxes increasing and decreasing about 2 to 3 hours later in the day (Fig. 13b). On the one hand, this results in an underestimation of LH fluxes in the morning of up to 100 W m^{-2} . As a consequence, the surface warms up too quickly causing an overestimation of the SH fluxes - which does, however, not fully compensate

460 for the overestimation of LH fluxes, resulting in a net bias of about -50 to -100 W m^{-2} . On the other hand, LH fluxes are overestimated in the afternoon with the largest values during the hours before sunset. The impact is most notable in the dry season (Fig. 13d). Median LH fluxes are overestimated by more than 200 W m^{-2} in the model, which leads to an increased cooling of the surface and resulting decreasing SH fluxes. Net turbulent fluxes are overestimated by about 100 W m^{-2} in the




Figure 13. Top: Annual median diurnal cycles of sensible (a) and latent (b) heat fluxes from the JSBACH model and measured at ATTO in 2017 and 2018. Boxplots display differences between the model and the measurements with quartiles as box and 10 %- and 90 %-percentiles as whiskers. Light blue bars indicate the fraction of available measurements. Bottom: Median diurnal cycles of flux differences between model and measurements for sensible and latent heat flux and their sum for the wet (c) and dry (d) seasons. Gray vertical lines mark the times of sunrise and sunset, respectively.

afternoon. Due to the considerable amount of missing measurements, however, it remains difficult to determine whether these
 465 biases have a significant impact on longer time periods.

To summarize, our comparison of JSBACH model results with soil and surface measurements at the ATTO site indicates that
 the following model processes should be evaluated more carefully in the future: Firstly, the model performance could possibly
 be improved by a better representation of the vertical structure of the soil. It would be useful to allow for different soil types
 in different model layers and additionally, more detailed knowledge of soil properties - either from field measurements at the
 470 considered site, or from high resolution data sets of soil types - could be used as boundary conditions. Furthermore, the vertical
 distribution of the root density should be considered to allow for a varying soil water sink with depth due to transpiration.



Secondly, it would be beneficial to include a separate canopy layer into the model to represent the temperature damping effect by the canopy. Such a canopy layer would also be able to include the canopy heat storage process (Heidkamp et al., 2018), which has been shown to be an important process controlling the diurnal cycle of turbulent heat fluxes (Renner et al., 2021).
475 To reduce the soil temperature bias of the model, the representation of additional cooling terms within the canopy layer (e.g. due to evaporation of rainfall intercepted by the vegetation) should be re-evaluated.

4 Summary and conclusions

In this study we used the land model JSBACH in a site-level setup to study its performance at a rainforest site - hereby focusing especially on processes influenced by the forest canopy. Observations at the Amazon Tall Tower Observatory were
480 used to evaluate the model performance and, as a first step, to optimize the external forcing data of the land model.

First, we compared near-surface atmospheric variables from ERA5 and MERRA-2 to five years of ATTO measurements to determine whether reanalysis data are generally suitable to be used as forcing data for the land model. For wind, both reanalyses were able to simulate the shape of the diurnal cycle correctly but strongly underestimated wind speeds at all heights by about 1 m s^{-1} . Biases of the same order of magnitude have been found in other studies, which also report that the sign of the bias
485 varies with region (Jourdier, 2020; Staffell and Pfenninger, 2016; Carvalho, 2019) and terrain (Gualtieri, 2021).

MERRA-2 reproduced the annual cycle of precipitation quite well, while ERA5 overestimated annual mean precipitation by 30%. Since the amount of incoming shortwave radiation was also underestimated, it is likely that the cloud cover in ERA5 was too large. ATTO measurements showed a bimodal distribution of diurnal precipitation with a second peak around the time of sun rise. ERA5 also produced early-morning precipitation but too early at night and MERRA-2 completely failed to capture
490 the second peak.

Contrary to precipitation, ERA5 captured the annual cycle of temperature better than MERRA-2. In the dry season, MERRA-2 overestimated monthly mean temperatures by more than 1 K, which is related to too large daily maximum temperatures in this season. For specific humidity, both reanalyses did not reproduce the shape of the diurnal cycle correctly. While ATTO measurements showed increasing humidity during the morning with a maximum around mid-day, both reanalyses produce a
495 local minimum of humidity during this time, which is possibly caused by an overestimated growth rate of the atmospheric boundary layer.

Next, we tested how much these biases of the reanalysis data affect the results of the land model. Comparing different model simulations, which use either ERA5 or MERRA-2 as forcing for the spin-up period, we found the largest differences of up to 20% for soil water content and up to 1.3 K for soil temperatures in the deepest soil layers. The soil temperature bias remained
500 non-negligible even after one year. For both soil temperature and water the choice of the spin-up data set accounted for more than 10% of the observed model biases during the first three months after spin-up in 0.7 m depth and for more than 50% closer to the surface. Thus - especially for shorter model runs of only a few days or weeks - the choice of spin-up data set is not negligible and can have a large impact on the model results.



505 Correcting biases of the forcing data also changes the land model results. We conducted sensitivity runs for wind speed and precipitation and found the largest changes in the dry season, when the soil is not saturated with water. For both variables, soil water differences amounted to up to 5 % for monthly means and 10 % for daily means in all depths. Too low wind speeds caused an overestimation of soil temperatures by about 0.5 K in all months. These results indicate that biases of the forcing data sets can have a notable impact on the model results and should be checked and corrected beforehand if possible.

510 Based on these results, we performed a model run with optimized forcing. The spin-up consists of nine years of ERA5 data (with corrected wind speed and precipitation biases), followed by one year of ATTO data. Results from a two-year model run forced by ATTO measurements were then compared to the observations of soil water content, soil temperatures and turbulent heat fluxes to identify possible model shortcomings. Comparing profiles of soil water we found that the model generally overestimates the water content in the upper layers until 0.32 m. On the one hand, this could be related to the choice of boundary conditions, like the overall soil type. It could also be caused by a lack of vertical differences of soil types or of the root density, which impacts transpiration.

520 We also found that the model overestimates soil temperatures by about 1 K with a slightly larger bias in the dry season. Comparisons with measured air temperatures above and within the canopy suggest that this is most likely caused by additional cooling within the canopy layer, e.g. by evaporation of rainfall intercepted by the vegetation, which is not sufficiently accounted for in the model. Furthermore, the penetration depth of the diurnal cycle of temperatures is overestimated. A more detailed analysis of the change of temperature amplitude with depth indicates that this is likely related to the temperature dampening effect of the forest canopy, which has not been incorporated into the JSBACH model, yet. The analysis of turbulent fluxes revealed a timing mismatch of the diurnal cycles of latent and sensible heat fluxes. ATTO flux measurements contain a large amount of missing data, however, and thus further analysis of longer and more complete time series would be required to examine the underlying causes in more detail.

525 To conclude, we suggest that future improvements of the JSBACH model could possibly focus on allowing more vertical variability - both of soil type and root density - within the soil column. Furthermore, a separate canopy layer would likely improve processes related to the energy transport within the soil. When implementing and evaluating future improvements of the canopy scheme, we suggest to consider especially the bias and phase shift of the soil temperature. Finally, we showed that the ATTO site provides an ideal framework for testing canopy related processes in LSMs and will certainly be useful in future modeling studies.

Appendix A: Early morning precipitation

530 To study the regional distribution of average precipitation patterns we use the IMERG data set (Integrated Multi-satellitE REtrievals for GPM from the NASA-JAXA Global Precipitation Measurement mission; Huffman et al. (2019)). The data set provides half-hourly estimates of surface precipitation rates on a 0.1° grid. More details can be found in Tan et al. (2019). We calculate the wet season (JFMA) average early morning precipitation sums for the years 2014 to 2018. Panel (a) of Fig. A1




Figure A1. Fraction of nighttime precipitation between (a) 0:00 and 8:00 and (b) 6:00 and 8:00 local time averaged over the wet season (JFMA) from 2014 to 2018 based on the IMERG precipitation data set. The circle marks the ATTO location. Red and orange boxes denote the considered grid boxes for ERA5 and MERRA-2, respectively (see also Fig. 1).

shows late night-early morning (LN-EM) precipitation sums for the period between 0:00 and 8:00 local time and panel (b) shows early morning (EM) precipitation sums after sunrise between 6:00 and 8:00 as a fraction of total daily precipitation.

LN-EM precipitation shows strong local gradients with up to 50 % of daily precipitation falling between 0:00 and 8:00 in the region about 200 km northeast of ATTO. Northeast was also the most common wind direction for this time period with atmospheric flow coming from the 30° - 105° sector more than half of the time (not shown). The fraction of LN-EM precipitation



decreases toward the southeast to only around 25 % in the region around Manaus. EM precipitation shows a band with a local maximum from northwest to southeast of about 50 km width, where precipitation after sunrise accounts for up to 9 % of the daily precipitation. The ATTO location receives about 30 % of its precipitation between 0:00 and 8:00 and about 6 % in the early morning. Even though the IMERG data set does not fully reproduce the early morning precipitation maximum measured
545 at ATTO between 7:00 and 8:00 (see Fig. 5a), it clearly shows that there is a significant amount of EM precipitation in this region with fractions changing significantly within a few hundred kilometers, i.e. within a only a few reanalysis grid cells.

Data availability. Meteorological measurements from the Amazon Tall Tower Observatory are available through the ATTO data portal (www.attodata.org). ERA5 surface data were generated using Copernicus Climate Change Service information [2021] and can be obtained from the Climate Data Store (<https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels?tab=overview>). ERA5 profile
550 data on model levels were obtained from ECMWF's MARS tape archive. MERRA-2 are available through the Goddard Earth Sciences Data and Information Services Center (<https://disc.gsfc.nasa.gov/>)

Author contributions. AS and FA designed the study. AA, MS and PT collected and prepared the measurement data. AS performed the data analysis, conducted the model runs, visualized the results and prepared the manuscript. FA assisted in interpreting the results and editing of the manuscript.

555 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This research was made possible by the German-Brazilian project ATTO, supported by the German Federal Ministry of Education and Research (BMBF contracts 01LB1001A and 01LK1602E) and the Brazilian Ministério da Ciência, Tecnologia e Inovação (MCTI/FINEP contract 01.11.01248.00) as well as the Max Planck Society. We furthermore acknowledge the support by the Amazon State University (UEA), FAPEAM, LBA/INPA, and SDS/CEUC/RDS-Uatumã. This work was partly funded by the Deutsche Forschungsgemein-
560 schaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2037 'CLICCS - Climate, Climatic Change, and Society' – Project Number: 390683824, contribution to the Center for Earth System Research and Sustainability (CEN) of Universität Hamburg. Thanks to ICDC, CEN, University of Hamburg for data support. The results contain modified Copernicus Climate Change Service information (2014-2018). Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains.



565 References

- Andreae, M. O., Acevedo, O. C., Araújo, A., Artaxo, P., Barbosa, C. G. G., Barbosa, H. M. J., Brito, J., Carbone, S., Chi, X., Cintra, B. B. L., da Silva, N. F., Dias, N. L., Dias-Júnior, C. Q., Ditas, F., Ditz, R., Godoi, A. F. L., Godoi, R. H. M., Heimann, M., Hoffmann, T., Kesselmeier, J., Könemann, T., Krüger, M. L., Lavric, J. V., Manzi, A. O., Lopes, A. P., Martins, D. L., Mikhailov, E. F., Moran-Zuloaga, D., Nelson, B. W., Nölscher, A. C., Santos Nogueira, D., Piedade, M. T. F., Pöhlker, C., Pöschl, U., Quesada, C. A., Rizzo, L. V., Ro, C.-U., Ruckteschler, N., Sá, L. D. A., de Oliveira Sá, M., Sales, C. B., dos Santos, R. M. N., Saturno, J., Schöngart, J., Sörgel, M., de Souza, C. M., de Souza, R. A. F., Su, H., Targhetta, N., Tóta, J., Trebs, I., Trumbore, S., van Eijck, A., Walter, D., Wang, Z., Weber, B., Williams, J., Winderlich, J., Wittmann, F., Wolff, S., and Yáñez Serrano, A. M.: The Amazon Tall Tower Observatory (ATTO): overview of pilot measurements on ecosystem ecology, meteorology, trace gases, and aerosols, *Atmospheric Chemistry and Physics*, 15, 10723–10776, <https://doi.org/10.5194/acp-15-10723-2015>, 2015.
- 570 Anwar, S. A., Reboita, M. S., and Llopart, M.: On the sensitivity of the Amazon surface climate to two land-surface hydrology schemes using a high-resolution regional climate model (RegCM4), *International Journal of Climatology*, 42, 2311–2327, <https://doi.org/10.1002/joc.7367>, 2022.
- Ardilouze, C., Batté, L., Bunzel, F., Decremier, D., Déqué, M., Doblas-Reyes, F. J., Douville, H., Fereday, D., Guemas, V., MacLachlan, C., et al.: Multi-model assessment of the impact of soil moisture initialization on mid-latitude summer predictability, *Climate Dynamics*, 49, 3959–3974, <https://doi.org/10.1007/s00382-017-3555-7>, 2017.
- 580 Balsamo, G., Beljaars, A., Scipal, K., Viterbo, P., van den Hurk, B., Hirschi, M., and Betts, A. K.: A Revised Hydrology for the ECMWF Model: Verification from Field Site to Terrestrial Water Storage and Impact in the Integrated Forecast System, *Journal of Hydrometeorology*, 10, 623 – 643, <https://doi.org/10.1175/2008JHM1068.1>, 2009.
- Bonan, G. B., Patton, E. G., Harman, I. N., Oleson, K. W., Finnigan, J. J., Lu, Y., and Burakowski, E. A.: Modeling canopy-induced turbulence in the Earth system: a unified parameterization of turbulent exchange within plant canopies and the roughness sublayer (CLM-ml v0), *Geoscientific Model Development*, 11, 1467–1496, <https://doi.org/10.5194/gmd-11-1467-2018>, 2018.
- 585 Bonan, G. B., Patton, E. G., Finnigan, J. J., Baldocchi, D. D., and Harman, I. N.: Moving beyond the incorrect but useful paradigm: reevaluating big-leaf and multilayer plant canopies to model biosphere-atmosphere fluxes—a review, *Agricultural and Forest Meteorology*, 306, 108435, <https://doi.org/10.1016/j.agrformet.2021.108435>, 2021.
- 590 Brümmer, B., Lange, I., and Konow, H.: Atmospheric boundary layer measurements at the 280 m high Hamburg weather mast 1995–2011: mean annual and diurnal cycles, *Meteorologische Zeitschrift*, pp. 319–335, <https://doi.org/10.1127/0941-2948/2012/0338>, 2012.
- Brun, E., Vionnet, V., Boone, A., Decharme, B., Peings, Y., Valette, R., Karbou, F., and Morin, S.: Simulation of northern Eurasian local snow depth, mass, and density using a detailed snowpack model and meteorological reanalyses, *Journal of Hydrometeorology*, 14, 203–219, <https://doi.org/10.1175/JHM-D-12-012.1>, 2013.
- 595 Carvalho, D.: An assessment of NASA’s GMAO MERRA-2 reanalysis surface winds, *Journal of Climate*, 32, 8261–8281, <https://doi.org/10.1175/JCLI-D-19-0199.1>, 2019.
- Chen, Y., Ryder, J., Bastrikov, V., McGrath, M. J., Naudts, K., Otto, J., Ottlé, C., Peylin, P., Polcher, J., Valade, A., et al.: Evaluating the performance of land surface model ORCHIDEE-CAN v1. 0 on water and energy flux estimation with a single-and multi-layer energy budget scheme, *Geoscientific Model Development*, 9, 2951–2972, <https://doi.org/10.5194/gmd-9-2951-2016>, 2016.



- 600 Chor, T. L., Dias, N. L., Araújo, A., Wolff, S., Zahn, E., Manzi, A., Trebs, I., Sá, M. O., Teixeira, P. R., and Sörgel, M.: Flux-variance and flux-gradient relationships in the roughness sublayer over the Amazon forest, *Agricultural and Forest Meteorology*, 239, 213–222, <https://doi.org/10.1016/j.agrformet.2017.03.009>, 2017.
- Costa, M. H., Biajoli, M. C., Sanches, L., Malhado, A. C., Hutyra, L. R., Da Rocha, H. R., Aguiar, R. G., and de Araújo, A. C.: Atmospheric versus vegetation controls of Amazonian tropical rain forest evapotranspiration: Are the wet and seasonally dry rain forests any different?, *Journal of Geophysical Research: Biogeosciences*, 115, <https://doi.org/10.1029/2009JG001179>, 2010.
- 605 de Oliveira, J. V., Ferreira, D. B. d. S., Sahoo, P. K., Sodré, G. R. C., de Souza, E. B., and Queiroz, J. C. B.: Differences in precipitation and evapotranspiration between forested and deforested areas in the Amazon rainforest using remote sensing data, *Environmental earth sciences*, 77, 1–14, <https://doi.org/10.1007/s12665-018-7411-9>, 2018.
- de Souza, C. M., Dias-Júnior, C. Q., Tóta, J., and de Abreu Sá, L. D.: An empirical-analytical model of the vertical wind speed profile above and within an Amazon forest site, *Meteorological Applications*, 23, 158–164, <https://doi.org/10.1002/met.1543>, 2016.
- 610 Decharme, B., Delire, C., Minvielle, M., Colin, J., Vergnes, J.-P., Alias, A., Saint-Martin, D., Séférian, R., Sénési, S., and Voldoire, A.: Recent changes in the ISBA-CTRIP land surface system for use in the CNRM-CM6 climate model and in global off-line hydrological applications, *Journal of Advances in Modeling Earth Systems*, 11, 1207–1252, <https://doi.org/10.1029/2018MS001545>, 2019.
- Dias-Júnior, C. Q., Dias, N. L., dos Santos, R. M. N., Sörgel, M., Araújo, A., Tsokankunku, A., Ditas, F., de Santana, R. A., Von Randow, C., Sá, M., et al.: Is there a classical inertial sublayer over the Amazon forest?, *Geophysical Research Letters*, 46, 5614–5622, <https://doi.org/10.1029/2019GL083237>, 2019.
- 615 Dias-Júnior, C. Q., Carneiro, R. G., Fisch, G., D’Oliveira, F. A. F., Sörgel, M., Botía, S., Machado, L. A. T., Wolff, S., Santos, R. M. N. d., and Pöhlker, C.: Intercomparison of Planetary Boundary Layer Heights Using Remote Sensing Retrievals and ERA5 Reanalysis over Central Amazonia, *Remote Sensing*, 14, 4561, <https://doi.org/10.3390/rs14184561>, 2022.
- 620 ECMWF: IFS Documentation CY41R2 - Part III: Dynamics and Numerical Procedures, <https://doi.org/10.21957/83wouv80>, 2016a.
- ECMWF: IFS Documentation CY41R2 - Part IV: Physical Processes, <https://doi.org/10.21957/tr5rv27xu>, 2016b.
- Ekici, A., Beer, C., Hagemann, S., Boike, J., Langer, M., and Hauck, C.: Simulating high-latitude permafrost regions by the JSBACH terrestrial ecosystem model, *Geoscientific Model Development*, 7, 631–647, <https://doi.org/10.5194/gmd-7-631-2014>, 2014.
- Fan, Y., Meijide, A., Lawrence, D. M., Rounsard, O., Carlson, K. M., Chen, H.-Y., Röhl, A., Niu, F., and Knohl, A.: Reconciling canopy interception parameterization and rainfall forcing frequency in the Community Land Model for simulating evapotranspiration of rainforests and oil palm plantations in Indonesia, *Journal of Advances in Modeling Earth Systems*, 11, 732–751, <https://doi.org/10.1029/2018MS001490>, 2019.
- 625 Feddes, R. A.: Simulation of field water use and crop yield, in: *Simulation of plant growth and crop production*, pp. 194–209, Pudoc, 1982.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 630 Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes



- from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *Journal of Advances in Modeling Earth Systems*, 5, 572–597, <https://doi.org/https://doi.org/10.1002/jame.20038>, 2013.
- 640 Giorgetta, M. A., Brokopf, R., Crueger, T., Esch, M., Fiedler, S., Helmert, J., Hohenegger, C., Kornblueh, L., Köhler, M., Manzini, E., Mauritsen, T., Nam, C., Raddatz, T., Rast, S., Reinert, D., Sakradzija, M., Schmidt, H., Schneck, R., Schnur, R., Silvers, L., Wan, H., Zängl, G., and Stevens, B.: ICON-A, the Atmosphere Component of the ICON Earth System Model: I. Model Description, *Journal of Advances in Modeling Earth Systems*, 10, 1613–1637, <https://doi.org/https://doi.org/10.1029/2017MS001242>, 2018.
- Global Modeling and Assimilation Office (GMAO): MERRA-2 V5.12.4, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), `tavg1_2d_slv_Nx: 2d, 1-Hourly,Time-Averaged,Single-Level,Assimilation,Single-Level-Diagnostics` <https://doi.org/10.5067/VJAFPLI1CSIV>; `tavg1_2d_rad_Nx: 2d,1-Hourly,Time-Averaged,Single-Level,Assimilation,Radiation-Diagnostics` <https://doi.org/10.5067/Q9QMY5PBNV1T>; `tavg1_2d_flux_Nx: 2d,1-Hourly,Time-Averaged,Single-Level,Assimilation,Surface Flux-Diagnostics` <https://doi.org/10.5067/7MCPBJ41Y0K6>; `tavg1_2d_lnd_Nx: 2d,1-Hourly,Time-Averaged,Single-Level,Assimilation,Land Surface-Diagnostics` <https://doi.org/10.5067/RKPHT8KC1Y1T>; Accessed: 03/03/2021, 2015.
- 645 Gualtieri, G.: Reliability of ERA5 Reanalysis Data for Wind Resource Assessment: A Comparison against Tall Towers, *Energies*, 14, 4169, <https://doi.org/10.3390/en14144169>, 2021.
- Hagemann, S.: An improved land surface parameter dataset for global and regional climate models, Max-Planck-Institut für Meteorologie, <https://doi.org/10.17617/2.2344576>, 2002.
- 655 Hardy, J., Melloh, R., Koenig, G., Marks, D., Winstral, A., Pomeroy, J., and Link, T.: Solar radiation transmission through conifer canopies, *Agricultural and forest meteorology*, 126, 257–270, <https://doi.org/10.1016/j.agrformet.2004.06.012>, 2004.
- Heidkamp, M., Chlond, A., and Ament, F.: Closing the energy balance using a canopy heat capacity and storage concept—A physically based approach for the land component JSBACHv3. 11, *Geoscientific Model Development*, 11, 3465–3479, <https://doi.org/10.5194/gmd-11-3465-2018>, 2018.
- 660 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: Complete ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate, Copernicus Climate Change Service (C3S) Data Store (CDS). (Accessed on 09-02-2021), 2017.
- 665 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1979 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) (Accessed on 12-02-2021), <https://doi.org/10.24381/cds.adbb2d47>, 2018.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.



- 675 Huffman, G., Stocker, E., Bolvin, D., Nelkin, E., and Tan, J.: GPM IMERG Final Precipitation L3 Half Hourly 0.1 degree x 0.1 degree V06, Greenbelt, MD, Goddard Earth Sciences Data and Information Services Center (GES DISC), last access: 27 August 2021, distributed in netCDF file format by ICDC, CEN, University of Hamburg., <https://doi.org/10.5067/GPM/IMERG/3B-HH/06>, 2019.
- Jourdir, B.: Evaluation of ERA5, MERRA-2, COSMO-REA6, NEWA and AROME to simulate wind power production over France, *Advances in Science and Research*, 17, 63–77, <https://doi.org/10.5194/asr-17-63-2020>, 2020.
- 680 Jungclaus, J. H., Lorenz, S. J., Schmidt, H., Brovkin, V., Brüggemann, N., Chegini, F., Crüger, T., De-Vrese, P., Gayler, V., Giorgetta, M. A., et al.: The ICON earth system model version 1.0, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002813, <https://doi.org/10.1029/2021MS002813>, 2022.
- Knauer, J., Zaehle, S., Reichstein, M., Medlyn, B. E., Forkel, M., Hagemann, S., and Werner, C.: The response of ecosystem water-use efficiency to rising atmospheric CO₂ concentrations: Sensitivity and large-scale biogeochemical implications, *New Phytologist*, 213, 1654–1666, <https://doi.org/10.1111/nph.14288>, 2017.
- 685 Lim, Y.-J., Hong, J., and Lee, T.-Y.: Spin-up behavior of soil moisture content over East Asia in a land surface model, *Meteorology and Atmospheric Physics*, 118, 151–161, <https://doi.org/10.1007/s00703-012-0212-x>, 2012.
- Lindroth, A., Mölder, M., and Lagergren, F.: Heat storage in forest biomass improves energy balance closure, *Biogeosciences*, 7, 301–313, <https://doi.org/10.5194/bg-7-301-2010>, 2010.
- 690 Loveland, T. R., Reed, B. C., Brown, J. F., Ohlen, D. O., Zhu, Z., Yang, L., and Merchant, J. W.: Development of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data, *International Journal of Remote Sensing*, 21, 1303–1330, <https://doi.org/10.1080/014311600210191>, 2000.
- Ma, Y. and Liu, H.: An advanced multiple-layer canopy model in the WRF model with large-eddy simulations to simulate canopy flows and scalar transport under different stability conditions, *Journal of Advances in Modeling Earth Systems*, 11, 2330–2351, <https://doi.org/10.1029/2018MS001347>, 2019.
- 695 Mäkelä, J., Knauer, J., Aurela, M., Black, A., Heimann, M., Kobayashi, H., Lohila, A., Mammarella, I., Margolis, H., Markkanen, T., et al.: Parameter calibration and stomatal conductance formulation comparison for boreal forests with adaptive population importance sampler in the land surface model JSBACH, *Geoscientific Model Development*, 12, 4075–4098, <https://doi.org/10.5194/gmd-12-4075-2019>, 2019.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenez-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von
- 705 Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, <https://doi.org/https://doi.org/10.1029/2018MS001400>, 2019.
- Moene, A. F. and Dam, J. C. v.: *Transport in the Atmosphere-Vegetation-Soil Continuum*, Cambridge University Press, <https://doi.org/10.1017/CBO9781139043137>, 2014.
- 710 Nabel, J. E., Naudts, K., and Pongratz, J.: Accounting for forest age in the tile-based dynamic global vegetation model JSBACH4 (4.20 p7; git feature/forests)–a land surface model for the ICON-ESM, *Geoscientific Model Development*, 13, 185–200, <https://doi.org/10.5194/gmd-13-185-2020>, 2020.



- Oliphant, A., Grimmond, C., Zutter, H., Schmid, H., Su, H.-B., Scott, S., Offerle, B., Randolph, J., and Ehman, J.: Heat storage and energy balance fluxes for a temperate deciduous forest, *Agricultural and Forest Meteorology*, 126, 185–201, <https://doi.org/10.1016/j.agrformet.2004.07.003>, 2004.
- Perrochet, P.: Water uptake by plant roots—A simulation model, I. Conceptual model, *Journal of hydrology*, 95, 55–61, [https://doi.org/10.1016/0022-1694\(87\)90115-6](https://doi.org/10.1016/0022-1694(87)90115-6), 1987.
- Randall, D. A., Dazlich, D. A., et al.: Diurnal variability of the hydrologic cycle in a general circulation model, *Journal of the atmospheric sciences*, 48, 40–62, [https://doi.org/10.1175/1520-0469\(1991\)048<0040:DVOTHC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1991)048<0040:DVOTHC>2.0.CO;2), 1991.
- Reichle, R. H., Liu, Q., Koster, R. D., Draper, C. S., Mahanama, S. P. P., and Partyka, G. S.: Land Surface Precipitation in MERRA-2, *Journal of Climate*, 30, 1643–1664, <https://doi.org/10.1175/JCLI-D-16-0570.1>, 2017.
- Renner, M., Kleidon, A., Clark, M., Nijssen, B., Heidkamp, M., Best, M., and Abramowitz, G.: How well can land-surface models represent the diurnal cycle of turbulent heat fluxes?, *Journal of Hydrometeorology*, 22, 77–94, <https://doi.org/10.1175/JHM-D-20-0034.1>, 2021.
- Santana, R. A., Dias-Júnior, C. Q., da Silva, J. T., Fuentes, J. D., do Vale, R. S., Alves, E. G., dos Santos, R. M. N., and Manzi, A. O.: Air turbulence characteristics at multiple sites in and above the Amazon rainforest canopy, *Agricultural and Forest Meteorology*, 260, 41–54, <https://doi.org/10.1016/j.agrformet.2018.05.027>, 2018.
- Santana, R. A. S. d., Dias-Júnior, C. Q., Vale, R. S. d., Tóta, J., and Fitzjarrald, D. R.: Observing and modeling the vertical wind profile at multiple sites in and above the Amazon rain forest canopy, *Advances in Meteorology*, 2017, <https://doi.org/10.1155/2017/5436157>, 2017.
- Sato, T., Miura, H., Satoh, M., Takayabu, Y. N., and Wang, Y.: Diurnal cycle of precipitation in the tropics simulated in a global cloud-resolving model, *Journal of Climate*, 22, 4809–4826, <https://doi.org/10.1175/2009JCLI2890.1>, 2009.
- Saturno, J., Holanda, B. A., Pöhlker, C., Ditas, F., Wang, Q., Moran-Zuloaga, D., Brito, J., Carbone, S., Cheng, Y., Chi, X., Ditas, J., Hoffmann, T., Hrabe de Angelis, I., Könemann, T., Lavrič, J. V., Ma, N., Ming, J., Paulsen, H., Pöhlker, M. L., Rizzo, L. V., Schlag, P., Su, H., Walter, D., Wolff, S., Zhang, Y., Artaxo, P., Pöschl, U., and Andreae, M. O.: Black and brown carbon over central Amazonia: long-term aerosol measurements at the ATTO site, *Atmospheric Chemistry and Physics*, 18, 12 817–12 843, <https://doi.org/10.5194/acp-18-12817-2018>, <https://acp.copernicus.org/articles/18/12817/2018/>, 2018.
- Staffell, I. and Pfenninger, S.: Using bias-corrected reanalysis to simulate current and future wind power output, *Energy*, 114, 1224–1239, <https://doi.org/10.1016/j.energy.2016.08.068>, 2016.
- Tan, J., Huffman, G. J., Bolvin, D. T., and Nelkin, E. J.: IMERG V06: Changes to the Morphing Algorithm, *Journal of Atmospheric and Oceanic Technology*, 36, 2471 – 2482, <https://doi.org/10.1175/JTECH-D-19-0114.1>, 2019.
- van den Hurk, B., Viterbo, P., Beljaars, A., and Betts, A.: Offline validation of the ERA40 surface scheme, *ECMWF Technical Memoranda*, <https://doi.org/10.21957/9aoaspz8>, 2000.
- Wang, D., Wang, G., and Anagnostou, E. N.: Evaluation of canopy interception schemes in land surface models, *Journal of Hydrology*, 347, 308–318, <https://doi.org/10.1016/j.jhydrol.2007.09.041>, 2007.
- Weedon, G. P., Balsamo, G., Bellouin, N., Gomes, S., Best, M. J., and Viterbo, P.: The WFDEI meteorological forcing data set: WATCH Forcing Data methodology applied to ERA-Interim reanalysis data, *Water Resources Research*, 50, 7505–7514, <https://doi.org/10.1002/2014WR015638>, 2014.
- Wright, J. S., Fu, R., Worden, J. R., Chakraborty, S., Clinton, N. E., Risi, C., Sun, Y., and Yin, L.: Rainforest-initiated wet season onset over the southern Amazon, *Proceedings of the National Academy of Sciences*, 114, 8481–8486, <https://doi.org/10.1073/pnas.1621516114>, 2017.



- 750 Yang, S. and Smith, E. A.: Mechanisms for diurnal variability of global tropical rainfall observed from TRMM, *Journal of climate*, 19, 5190–5226, <https://doi.org/10.1175/JCLI3883.1>, 2006.
- Yang, Y., Uddstrom, M., and Duncan, M.: Effects of short spin-up periods on soil moisture simulation and the causes over New Zealand, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD016121>, 2011.
- Yang, Z.-L., Dickinson, R. E., Henderson-Sellers, A., and Pitman, A. J.: Preliminary study of spin-up processes in land surface models with
755 the first stage data of Project for Intercomparison of Land Surface Parameterization Schemes Phase 1(a), *Journal of Geophysical Research: Atmospheres*, 100, 16 553–16 578, <https://doi.org/10.1029/95JD01076>, 1995.
- Yi, C.: Momentum transfer within canopies, *Journal of Applied Meteorology and Climatology*, 47, 262–275, <https://doi.org/10.1175/2007JAMC1667.1>, 2008.
- Zahn, E., Dias, N. L., Araújo, A., Sá, L. D., Sörgel, M., Trebs, I., Wolff, S., and Manzi, A.: Scalar turbulent behavior in the roughness
760 sublayer of an Amazonian forest, *Atmospheric Chemistry and Physics*, 16, 11 349–11 366, <https://doi.org/10.5194/acp-16-11349-2016>, 2016.
- Zheng, Z. and Wang, G.: Modeling the dynamic root water uptake and its hydrological impact at the Reserva Jaru site in Amazonia, *Journal of Geophysical Research: Biogeosciences*, 112, <https://doi.org/10.1029/2007JG000413>, 2007.