Comparison of Methods to Estimate Aerosol Effective Radiative Forcings in CMIP Models

Mark D. Zelinka¹, Christopher J. Smith^{2,3}, Yi Qin⁴, and Karl E. Taylor¹

¹Lawrence Livermore National Laboratory, 7000 East Avenue, L-103, Livermore, CA 94550, USA
 ²School of Earth and Environment, University of Leeds, LS2 9JT, UK
 ³International Institute for Applied Systems Analysis, Laxenburg 2361, Austria
 ⁴Pacific Northwest National Laboratory, PO Box 999, Richland, WA 99352, USA

Correspondence: Mark D. Zelinka (zelinka1@llnl.gov)

Abstract. Uncertainty in the effective radiative forcing (ERF) of climate primarily arises from the unknown contribution of aerosols, which impact radiative fluxes directly and through modifying cloud properties. Climate model simulations with fixed sea surface temperatures but perturbed atmospheric aerosol loadings allow for an estimate of how strongly the planet's radiative energy budget has been perturbed by the increase in aerosols since pre-industrial times. The approximate partial radiative

- 5 perturbation (APRP) technique further decomposes the contributions to the direct forcing from due to aerosol scattering and absorption, and to the indirect forcing from due to aerosol-induced changes in cloud scattering, amount, and absorption, as well as the effects of aerosols on surface albedo. Here we evaluate previously published APRP-derived estimates of aerosol effective radiative forcings from these simulations conducted in the 6th phase of the Coupled Model Intercomparison Project (CMIP6) and find that they are slightly biased as a result of large but compensating errors two large coding errors that in most
- 10 cases fortuitously compensate. The most notable exception is the direct radiative forcing from absorbing aerosols, which is more than 40% larger averaged across CMIP6 models in the present study. These biases are largest for the aerosol direct effect owing to underestimated aerosol absorption. Correcting these biases eliminates the residuals and leads to better agreement with ground-truth benchmark estimates derived from double-calls to the radiation code. The APRP method – when properly implemented – remains a highly accurate and efficient technique for diagnosing aerosol ERF in cases where double radiation
- 15 calls are not available, and in all cases it provides quantification of the individual contributors to the ERF that are highly useful but not otherwise available.

1 Introduction

The primary source of uncertainty in effective radiative forcing of the climate comes from aerosols, both through their direct impact on radiation and via modifying cloud properties. This uncertainty limits our ability to know how strongly the Earth

20 has been forced over recent decades, which hampers our ability to confidently narrow bounds on climate sensitivity based on the observed temperature record (Sherwood et al., 2020). It also degrades our confidence in predictions of near-term climate evolution, particularly whether and how soon dangerous global mean temperature thresholds will be crossed (Watson-Parris and Smith, 2022; Dvorak et al., 2022), and the committed warming level if emissions rapidly cease (Armour and Roe, 2011) 25 many regions may change (Persad et al., 2022).

30

Despite its importance, aerosol radiative forcing (and forcing in general) has historically been poorly diagnosed in global climate models, though recent efforts have improved this state of affairs. Standard atmosphere-only model experiments to diagnose aerosol radiative forcing have been designed and made part of the CMIP5 and CMIP6 protocols, allowing for a relatively clean method for diagnosing aerosol ERF across models. Diagnostic approaches of various levels of sophistication have also been developed and applied to climate model output to provide consistent estimates of aerosol ERF across models.

A common method for computing aerosol ERF involves additional calls to the radiation code neglecting aerosols in the atmospheric column (Ghan, 2013), as described further below. Gryspeerdt et al. (2020) extended this method to additionally separate the indirect effect into aerosol effects on cloud droplet number concentration (the Twomey effect) and aerosol-induced adjustments of cloud fraction and liquid water path. However, aerosol-free radiation diagnostics are only available in a subset

35 of CMIP6 models that took part in the Radiative Forcing Model Intercomparison Project (RFMIP) (Pincus et al., 2016). Moreover, these diagnostics do not separately quantify the absorption and scattering components of the direct effect or the cloud absorption, scattering, and amount components of the indirect effect.

The approximate partial radiative perturbation (APRP) technique (Taylor et al., 2007) offers another method of computing aerosol ERF. Unlike the Ghan (2013) method this does not require additional aerosol-free radiation calls but rather operates

- 40 on standard monthly resolution model output available across all models. Another advantage is that it allows for a breakdown of the aerosol direct effect into absorption and scattering components and of the indirect effect into absorption, amount, and scattering components. The main disadvantages are that it is an approximate technique that may be biased with respect to more accurate methods, and that it only applies to shortwave (SW) radiation. Fortunately, aerosol direct and indirect effects primarily operate in the SW with much smaller effects in the longwave (LW), except for models with strong aerosol effects
- 45 on high clouds <u>. If a small degradation of absolute accuracy can be tolerated</u>(Zelinka et al., 2014; Smith et al., 2020). Hence, APRP allows for an efficient way to quantify aerosol forcing and its individual components, making it highly valuable for systematically inter-comparing the full suite of models performing aerosol perturbation experiments.

The APRP technique has been used to quantify aerosol ERF in models, revealing the diverse strengths of the various terms comprising it across models taking part in CMIP5 (Zelinka et al., 2014) and CMIP6 (Smith et al., 2020). These studies, however,

- 50 independently implemented the technique as computer code, and made different choices that have quantitative impacts on the results (as described below). In this study, we demonstrate that the implementation of the APRP method in Smith et al. (2020) was erroneous, leading to slightly biased values of aerosol ERF compared to the correct formulation implemented in Zelinka et al. (2014). We explain and quantify the two largely compensating errors that cause the bias. We also compare the two APRP formulations to the double radiation call method to evaluate how well APRP-derive APRP-derived results agree with this
- 55 independent technique. Finally, we provide values of aerosol ERF components that are corrected from those reported in Smith et al. (2020) and supplemented with additional models that have become available since the publication of that paper.

2 Data and Methods

2.1 Climate model simulations used

Our analysis makes use of pairs of idealized atmosphere-only climate model simulations in which sea-surface temperatures

- 60 (SSTs) and sea ice concentrations (SICs) are fixed at model-specific preindustrial climatological values. Aerosol burdens are set to their preindustrial levels in the control experiment and to their present-day levels in the perturbation experiment. In CMIP6, these experiments are known as 'piClim-control' and 'piClim-aer', respectively, and present-day is interpreted as year 2014 (Pincus et al., 2016). In CMIP5, they are known as 'sstClim' and 'sstClimAerosol', repectively, with present-day interpreted as year 2000. These experiments are nominally 30 years long.
- We make use of all available ensemble members of all models that performed these simulations (listed in Tables 1 and 2), with the exception of the EC-Earth3 models. These were model. This was excluded because of spurious up- and down-welling clear-sky shortwave radiative fluxes at the surface ('rsuscs' and 'rsdscs') that are determined to be erroneous and bias the APRP calculations. In particular, we identified numerous examples in which these fluxes exhibited very large values at individual grid cells and months but were surrounded in time and space by near-zero values (e.g., in winter locations with negligible insolation).
- 70 The monthly fields used in our analyses are listed in Table A1, and abbreviations and other nomenclature used throughout the paper are listed in Table A2.

2.2 Aerosol Effective Radiative Forcing Calculations

2.2.1 IPCC AR6 Definitions

One can express the total change in net (downwelling minus upwelling) radiation (ΔR) between a control experiment and an aerosol-perturbed experiment (both with fixed SSTs and SICs) as the sum of effective radiative forcings due to aerosolradiation interactions , (*ERF_{ari}*) and aerosol-cloud interactions , and changes in surface albedo: (*ERF_{aci}*), as well as changes in TOA radiation due to changes in surface albedo (ΔR_{alb}) and surface temperature (ΔR_{T_0}):

$$\Delta R = ERF_{ari} + ERF_{aci} + \underbrace{ERF\Delta R_{alb} + \Delta R_{T_0}}_{(1)}.$$

As defined in IPCC AR6 (Forster et al., 2021), ERF_{ari} comprises the instantaneous radiative forcing (IRF_{ari}), non-cloud atmospheric adjustments that are uncoupled to any change in global surface temperature, and adjustments of clouds due to changes in the thermal structure of the atmosphere caused by absorbing aerosols (also known as the "semidirect effect"):

$$ERF_{ari} = IRF_{ari} + K^T \Delta T + K^q \Delta q + K^C \Delta C_{semidirect},$$
⁽²⁾

where K^{χ} terms quantify the sensitivity of TOA net radiation to infinitesimal perturbations in variable χ :

$$K^{\chi} = \frac{\partial R}{\partial \chi},\tag{3}$$

and here and elsewhere χ may represent temperature (T), humidity (q), surface albedo (α), or clouds (C). It is intended that ΔT and Δq include only changes in tropospheric temperature and water vapor that occur independently of changes in surface temperature, not those that occur in response to any land surface temperature change in these fixed-SST simulations. In practice it is difficult to determine which portion of the change in any field is an adjustment to aerosols versus a response that is mediated by land surface temperature changes or by circulation changes induced by said temperature changes. Hence for

- 90 simplicity we follow a conservative approach, also taken by AR6, and exclude the radiative impact of land temperature changes from ERF_{ari} . This approach leads to positive longwave ERF_{ari} values that are smaller than if this correction is not applied, making the overall net negative ERF about 5% stronger on average across models, consistent with the 5% inflation that IPCC AR6 applied to the ERF values derived in Smith et al. (2020) and Zelinka et al. (2014). Finally, the radiative impact of surface albedo changes – which we also keep separate from the ERF terms in Eq 1 – arises due to deposition of absorbing aerosol on
- 95 snow and ice, changes in snow and ice cover arising from the aerosol-induced change in climate, and aerosol-induced changes in the relative amount of direct versus diffuse radiation reaching the surface.

As defined in IPCC AR6, ERF_{aci} comprises the instantaneous radiative forcing due to changes in cloud liquid and ice particle number concentrations and sizes (IRF_{aci}), and subsequent adjustments of cloud water and coverage ($K^C \Delta C_{adjustments}$):

100
$$ERF_{aci} = IRF_{aci} + K^C \Delta C_{adjustments}.$$
 (4)

Finally, the surface albedo forcing is generally kept separate from the other ERFs,

$$ERF_{alb} = K^{\alpha} \Delta \alpha,$$

and includes deposition of absorbing aerosol on snow and ice as well as changes in snow and ice cover arising from the aerosol-induced change in climate.

Although the above decomposition of direct and indirect forcings is employed by IPCC AR6 (Forster et al., 2021), in the Ghan (2013) and APRP methods described below, there is no way to separately quantify the three cloud-related terms – the semidirect effect which is considered part of ERF_{ari} , the instantaneous component due to changing particle number and size (the Twomey effect), and the subsequent adjustments. Instead, for these two methodologies, the cloud terms are combined into a single term that is included as part of the ERF_{aci} :

110
$$K^C \Delta C = IRF_{aci} + K^C \Delta C_{adjustments} + K^C \Delta C_{semidirect}.$$
 (5)

 $K^{C}\Delta C$ can be further broken down into contributions from changes in cloud amount and optical properties.

2.2.2 Approximate Partial Radiative Perturbation (APRP) Technique

Taylor et al. (2007) developed an efficient method of decomposing perturbations to the TOA SW energy budget from clouds, the cloud-free portion of the atmosphere (here assumed to be dominated by aerosols), and surface albedo. Briefly, the APRP

115 method employs a simple one-layer model of the atmosphere to diagnose the scattering and absorption of SW radiation at the surface and in the atmosphere. This simple model represents the transfer of SW radiation through the atmosphere at every grid point on the globe in terms of a small number of parameters – the insolation, surface albedo, an atmospheric scattering

coefficient, and an atmospheric absorptance coefficient. Given the known SW fluxes at the TOA and surface under both clearand all-sky conditions as well as the total cloud fraction, at each grid point on the globe one can solve for the atmospheric

- 120 scattering and absorption parameters in this simplified representation such that the upwelling and downwelling SW radiative fluxes at the surface and TOA match those produced by the GCM. Then the sensitivity of TOA albedo to these parameters can be determined, allowing one to isolate the individual contributions from changes in non-cloud atmospheric constituents and from changes in cloud properties. This technique is an approximation to the more rigorous but difficult-to-implement partial radiative perturbation (PRP) technique, and was shown to closely agree with PRP-derived SW cloud and surface albedo feedbacks
- 125 (Taylor et al., 2007), with errors that were no larger than 10%. It is uniquely well-suited for quantifying and decomposing aerosol forcing, since aerosols primarily affect scattering and absorption of SW radiation both directly and indirectly through clouds. Code to A more detailed description of APRP and how it is used to estimate the individual aerosol forcing components is provided in Appendix A.2, and code to perform all APRP calculations for this paper is provided in the *Code availability* section.
- APRP (denoted below with a superscript A) provides estimates of SW ERF (superscript SW) that are made up of slightly different term groupings than the IPCC AR6 definition:

$$ERF_{ari}^{A,SW} = IRF_{ari}^{SW} + K^{q,SW}\Delta q = ERF_{ari}^{SW} - K^{C,SW}\Delta C_{semidirect},$$
(6)

and

$$ERF_{aci}^{A,SW} = IRF_{aci}^{SW} + K^{C,SW}\Delta C_{adjustments} + K^{C,SW}\Delta C_{semidirect} = ERF_{aci}^{SW} + K^{C,SW}\Delta C_{semidirect}.$$
⁽⁷⁾

135 The impact of temperature changes on the SW ERF_{ari} can be neglected. Therefore in the SW, APRP's direct effect equals IPCC's direct effect minus the semidirect effect, while APRP's indirect effect equals IPCC's indirect effect plus the semidirect effect. The sum of the direct and indirect SW effects are the same, independent of how the individual components are defined. Thus, from Eqs $\frac{7 \text{ and } 86 \text{ and } 7}{12}$:

$$ERF_{ari}^{A,SW} + ERF_{aci}^{A,SW} = ERF_{ari}^{SW} + ERF_{aci}^{SW}.$$
(8)

140 The APRP surface albedo term is identical to that given in Eq 5:-

 $ERF^{A,SW}_{alb} = K^{\alpha}\Delta\alpha.$

Further insight comes from separating $\text{ERF}_{ari}^{A,SW}$ into absorption and scattering components,

$$ERF_{ari}^{A,SW} = ERF_{ari,abs}^{A,SW} + ERF_{ari,scat}^{A,SW},\tag{9}$$

and from separating $\text{ERF}_{aci}^{A,SW}$ into cloud absorption, amount, and scattering components,

$$145 \quad ERF_{aci}^{A,SW} = ERF_{aci,abs}^{A,SW} + ERF_{aci,amt}^{A,SW} + ERF_{aci,scat}^{A,SW}.$$
(10)

APRP also quantifies the impact of surface albedo changes on TOA SW radiation.

2.2.3 LW ERFs: Proxies Derived from Standard Model Output

There is no equivalent to APRP for LW radiation, so instead we compute proxies (denoted with superscript P) for LW ERFs using standard model output, following Zelinka et al. (2014) and Smith et al. (2020). The LW direct effect is estimated as the change in clear-sky TOA LW radiation excluding the portion due solely to land surface temperature changes:

$$ERF_{ari}^{P,LW} = \Delta R_{cs}^{LW} = -\Delta \underline{OLRR_{cs}^{T_0}}_{cs}, \tag{11}$$

where R^{LW} refers to net (downwelling minus upwelling) LW radiation , *OLR* is outgoing longwave radiation, and the subscript *cs* refers to clear-sky conditions. The change in clear-sky LW radiation due to changes in (land) surface temperature is computed by multiplying the change in surface temperature between the two experiments by the clear-sky surface temperature radiative kernel (Huang et al., 2017):

$$\Delta R_{cs}^{T_0} = K_{cs}^{T_0} \Delta T_0. \tag{12}$$

We compute a proxy for longwave ERF_{aci} as follows: the change in LW cloud radiative effect:

$$ERF_{aci}^{P,LW} = \Delta CRE^{LW} = \Delta R^{LW} - \Delta R_{cs}^{LW} = \Delta OLR_{cs} - \Delta OLR.$$
(13)

As shown in Appendix AB, our proxy for the LW direct effect equals IPCC's direct effect, minus the semidirect effect, minus masking terms that quantify how much the radiative impact of changes in temperature, humidity, and aerosols are attenuated by the presence of clouds. Our proxy for the LW indirect effect equals IPCC's indirect effect, plus the semidirect effect, plus the aforementioned masking terms.

2.2.4 Double Radiation Call Method

Ghan (2013) introduced a method to compute aerosol direct and indirect effects that relies on "aerosol-free" radiative fluxes
under both clear- and all-sky conditions. These are produced by performing additional calls to the radiation code during model integration in which all aerosols are neglected. For the models that provide aerosol-free radiative fluxes, we compute the following quantities, which are given the superscript *G* to indicate Ghan. Ghan (2013) defines the direct forcing as

$$ERF_{ari}^G = \Delta R - \Delta R_{af},\tag{14}$$

the indirect forcing as

150

155

170
$$ERF_{aci}^G = \Delta R_{af} - \Delta R_{af,cs},$$
 (15)

and a third forcing term as ÷

$$ERF_{other}^G = \Delta R_{af,cs},$$
(16)

where the subscript af refers to aerosol-free radiative fluxes and, as above, ΔR represents the change in net radiation between the control and aerosol-perturbed experiment.

- As shown in Appendix AB, Ghan's direct aerosol radiative forcing equals IPCC's instantaneous direct forcing minus masking terms that quantify how much the radiative impact of changes in temperature, humidity, surface albedo, and clouds are attenuated by the presence of aerosols. Ghan's indirect effect equals IPCC's indirect effect plus masking terms that quantify how much the radiative impact of changes in temperature, humidity, and surface albedo are attenuated by the presence of clouds under aerosol-free conditions and how much the radiative impact of changes in clouds are attenuated by the presence of
- aerosols, plus the semidirect effect. Finally, ERF_{other}^G which in the SW Ghan (2013) refers to as the surface albedo forcing – equals $ERF_{atb} \Delta R_{alb}$ plus the aerosol-free clear-sky radiative contributions from changes in humidity, plus a masking term that quantifies how much the radiative impact of changes in surface albedo are attenuated by the presence of both clouds and aerosols.

3 Results

185 **3.1** Errors in APRP Implementation of Smith et al. (2020)

We begin by comparing the sum of all APRP-derived SW ERF components with the total change in SW radiation between the perturbed aerosol and control experiment. Since the majority of the results shown below are derived using APRP, we omit the *A* superscript hereafter. If APRP is correctly implemented, the sum of its components should perfectly reproduce the total change in SW radiation between the control and perturbed experiment. As shown in Figure 1, the APRP implementation in

- 190 the present study has a negligible residual whereas that in Smith et al. (2020) is generally nonzero, ranging from -0.21 to +0.08 W/m² across models. These errors, resulting from a mistake in coding as detailed below, are comparable in magnitude to typical values for the total aerosol direct effect and the cloud absorption and amount components of the indirect effect (shown below).
- The small residual in Smith's implementation of APRP masks the two much larger errors that are nearly perfectly com-195 pensating. The first error – which in isolation leads to underestimated ERF magnitudes – is that net (downwelling minus upwelling) rather than downwelling TOA SW radiation was used in all calculations. The second error – which in isolation leads to overestimated ERF magnitudes – is in how TOA albedo sensitivities are computed, particularly in the calculation of overcast-sky albedo sensitivity to aerosol and cloud absorption and scattering. The code overestimated the impact of changes in scattering or absorption on overcast-sky fluxes because it made use of the raw difference between clear-sky scattering and
- 200 absorption coefficients. The correct implementation, in contrast, scales this difference by the appropriate factor to account for the fact that changes in the non-cloud portion of a column are attenuated by the presence of clouds in the column. This error is described in further detail in Appendix A.3. The second error – which in isolation leads to underestimated ERF magnitudes – is that net (downwelling minus upwelling) rather than downwelling TOA SW radiation was used in all calculations. This error is described in further detail in Appendix A.4.
- 205 Estimates of global mean ERF^{SW}_{ari+aci} across CMIP6 models. The pink bar shows the values derived in this study using the corrected APRP formulation. Overlain on this with dotted gray hatching is the ERF derived if using net rather than downwelling SW radiation, and the ratio between these two is printed. The adjacent black bar shows the ERF derived if using erroneous



Figure 1. Global mean residuals estimated as the difference between the true model-produced change in TOA SW fluxes and those estimated by summing the APRP components, shown for the APRP implementation in this study and in Smith et al. (2020). Each marker represents a different CMIP model (all pink markers overlap).

albedo sensitivities (but correct insolation). Overlain on this with cyan hatching is this same value scaled by the aforementioned ratio, which closely matches the values derived using the APRP formulation of Smith et al. (2020), which are shown with cyan dots.

To quantify the impact of these two errors, we first corrected both errors in the code of Smith et al. (2020), and verified that it produces results identical to those from the original APRP implementation of Zelinka et al. (2014). The correct implementation is shown as pink bars in Figure 2, and serves as the baseline against which subsequent calculations are compared. We then performed the APRP calculations two more times, once reverting back to the original erroneous insolation formulation, and once reverting back to the original erroneous albedo sensitivity formulation. The ERF estimate derived from the APRP

- implementation with erroneous insolation is shown by the dotted hatching overlain on the pink bar. This bar has a smaller magnitude than the true value because it uses net (down- minus up-welling radiation) rather than downwelling radiation. The ratio of these two ERF estimates (shown in printed numbers) is 0.77 averaged across models, with a standard deviation of 0.05. This is consistent with the fact that net radiation is equal to the downwelling radiation times (1 minus planetary albedo) and
 - that planetary albedos vary between about 0.2 and 0.3.

210

215

The ERF estimates derived from the APRP implementation with erroneous albedo sensitivities (but correct insolation) are shown in the black bars in Figure 2. The erroneous albedo sensitivity formulation leads to ERF values that are biased too large in magnitude. This arises almost entirely from the scattering components of ERF_{aci}^{SW} and ERF_{ari}^{SW} , which are overestimated (discussed further below). This occurs because the erroneous code did not properly allow the increase in aerosol scattering to

225 be attenuated by cloud scattering, and did not properly allow the increase in cloud scattering to be attenuated by non-cloud (aerosol) scattering (not shown).



Figure 2. Estimates of global mean ERF^{SW}_{ari±aci} across CMIP6 models. The pink bar shows the values derived in this study using the corrected APRP formulation. Overlain on this with dotted gray hatching is the ERF derived if using net rather than downwelling SW radiation, and the ratio between these two is printed. The adjacent black bar shows the ERF derived if using erroneous albedo sensitivities (but correct insolation). Overlain on this with cyan hatching is this same value scaled by the aforementioned ratio, which closely matches the values derived using the APRP formulation of Smith et al. (2020), which are shown with cyan dots.

The bar with cyan hatching overlain on the black bar in Figure 2 indicates the result if we scale these overestimated ERFs by the ratios diagnosed above to account for the compensating insolation error. This scaled estimate closely matches the original Smith et al. (2020) formulation in which both errors are present (cyan dots).

230

Hence the two errors are individually substantial in magnitude, but act in opposite directions that almost perfectly compensate, leading to total ERF values that are in good agreement with the correct values estimated here using the APRP implementation of Zelinka et al. (2014). This also holds for the geographic structure of the ERF components, which are highly consistent between the two APRP implementations but differ quantitatively (not shown). This appears to be a fortuitous result. Moreover, while this good agreement with the correct values holds for the total ERF, several of its sub-components show larger biases, as 235 discussed next.

9



Figure 3. Estimates of global mean (a) $\text{ERF}_{art+aci}^{SW}$, (b) ERF_{art}^{SW} , (c) ERF_{act}^{SW} , and (d) ΔSW_{alb} in each CMIP5 and CMIP6 model derived using the APRP formulation of Smith et al. (2020) scattered against estimates computed in this study. Cyan markers indicate the values estimated by Smith et al. (2020), gray markers indicate values derived using net rather than downwelling SW radiation (but correct albedo sensitivities), and black markers indicate values derived using erroneous albedo sensitivities (but correct insolation). Gray markers in panel d are overlain by cyan markers. Correlation, slope, and intercept are reported as r, m, and b, respectively, and correlations that are statistically significant at 95% confidence are indicated with an asterisk.

Estimates of global mean (a) $\text{ERF}_{ari+aci}^{SW}$, (b) ERF_{alb}^{SW} , (c) ERF_{ari}^{SW} , and (d) ERF_{aci}^{SW} in each CMIP5 and CMIP6 model derived using the APRP formulation of Smith et al. (2020) scattered against estimates computed in this study. Cyan markers indicate the values estimated by Smith et al. (2020), gray markers indicate values derived using net rather than downwelling SW radiation (but correct albedo sensitivities), and black markers indicate values derived using erroneous albedo sensitivities (but correct insolation). Correlation, slope, and intercept are reported as r, m, and b, respectively, and correlations that are statistically significant at 95% confidence are indicated with an asterisk.

As in Figure 3, but for the scattering and absorption sub-components of ERF_{ari}^{SW} and ERF_{aci}^{SW} .

In Figure 3 we scatter estimates of $\text{ERF}_{ari+aci}^{SW}$, its two sub-components (ERF_{ari}^{SW} and ERF_{aci}^{SW}), and the $\frac{\text{ERF}_{alb}^{SW}}{\Delta SW_{alb}}$ term derived using the APRP implementation of Smith et al. (2020) against those derived using the implementation of Zelinka

et al. (2014). ERF^{SW}_{ari} and ERF^{SW}_{aci} are further broken down into their scattering and absorption sub-components in Figure 4. The aforementioned compensating biases are indicated by the gray and black markers, which show the ERF values derived if correcting each bias in turn. Correcting the albedo sensitivity formulation but keeping the insolation bias leads to ERF values that are too small in magnitude (Figure 3-4, gray markers). This is expected because a given change in albedo will produce a weaker TOA impact when using a smaller (downwelling minus upwelling) SW radiation stream. Further details are provided
in Appendix A.4.



Figure 4. As in Figure 3, but for the scattering and absorption sub-components of ERF_{ari}^{SW} and ERF_{aci}^{SW}

255

Correcting the insolation but keeping the erroneous albedo sensitivity formulation (black markers) has the opposite effect: In most cases it leads to ERF values that are too large in magnitude for reasons described in Appendix A.3. This indicates that using correct values of downwelling (rather than net) SW radiation exposes the erroneous albedo sensitivity formulation wherein changes in scattering or absorption have too-strong an effect on TOA SW radiation. This error primarily manifests itself in the scattering components of ERF_{ari}^{SW} and ERF_{aci}^{SW} (Figure 4a and b, black markers), whereas the $\frac{\text{ERF}_{alb}^{SW}}{\text{Biss vanishes}}$ because it is not affected by the albedo sensitivity formulation error (Figure 3b, black markers), and the absorption components are largely corrected when the insolation is corrected (Figure 4c and d, black markers). The albedo sensitivity formulation error has a larger impact on the scattering components because clouds scatter more <u>SW radiation</u> than they absorbSW radiation. This means that neglecting the attenuation of the change in aerosol scattering by cloud scattering induces a larger error than

- 260 neglecting cloud absorption when computing the impact of a change in aerosol absorption. Finally, the ΔSW_{alb} bias vanishes once the insolation bias is corrected (Figure 3d, black markers). This is because, unlike for the sensitivity of planetary albedo to atmospheric scattering and absorption parameters, the sensitivity of planetary albedo to surface albedo is implemented correctly in Smith et al. (2020).
- For the ERF components that depend only on scattering of radiation, these two biases largely compensate such that the values reported in Smith et al. (2020) agree well with those estimated here (Figure 4a and b, cyan markers). However, for the absorption components of ERF_{ari}^{SW} and ERF_{aci}^{SW} (for which the albedo sensitivity formulation was not as biased) and for the surface albedo term (which does not depend on how the atmospheric scattering and absorption sensitivities are formulated), the insolation error is largely uncompensated. Therefore, the values reported in Smith et al. (2020) remain systematically biased for these components. This is most apparent for $\frac{\text{ERF}_{alb}^{SW}}{\text{ERF}_{alb}^{SW}}$ (Figure 3b ΔSW_{alb} (Figure 3d) and the absorption component of ERF_{ari}^{SW} (Figure 4c).





Figure 5 shows the CMIP6 multi-model mean SW ERF components estimated using the APRP formulation of Smith et al. (2020) and as derived in this study. Note that the Smith et al. values differ slightly from those reported in that study because here we include four additional CMIP6 models not assessed in that study and exclude EC-Earth3 which was used in that study. We find that all but three ERF values are within 5% of those diagnosed by Smith et al. (2020): The very small ΔSW_{alb} component

275 increases in strength from -0.02 to -0.03 W/m². More importantly, the negative ERF_{avi}^{SW} diagnosed here is roughly 20% weaker in magnitude. This is primarily because the positive absorbing component of ERF_{avi}^{SW} is roughly 40% larger in the present study.

3.2 Validation of ERF Estimates Against the Double-Call Method

In Figure 5-6 we compare the SW ERF values estimated using APRP with those derived using the technique of Ghan (2013) in which radiation calculations are performed an additional time with all aerosols neglected. This comparison is done for 10 models (indicated with asterisks in Table 2) that provided the necessary aerosol-free diagnostics. As detailed earlier mentioned earlier and as detailed in Appendix B, APRP and Ghan (2013) define each of these terms differently(see Appendix A), so differences between the two estimates do not necessarily indicate errors. However, one would expect these to be closely related and, when implemented correctly, APRP values closely reproduce the double-call double-call values. Indeed, both APRP implementations are well correlated with the double-call method for each ERF component, but the RMSD values are much smaller when using the values derived in this study.

Even when correctly implemented, APRP-estimated ERF_{ari}^{SW} values are biased towards stronger negative / weaker positive values than those derived via double-call (Figure 5e6b). This is compensated somewhat by the ERF_{aci}^{SW} values, for which the APRP method is biased towards smaller negative values (Figure 5d6c), as also shown for a single model in Zelinka et al.

- 290 (2014). APRP also yields $ERF_{alb}^{SW} \Delta SW_{alb}$ values that are weaker in magnitude than produced by the double-call (Figure 5b6d). As shown in Appendix AB, these differences are expected because the two methods are diagnosing slightly different things for each of the individual components, with the offset arising from small masking terms. For example, Ghan (2013) defines $ERF_{alb}^{SW} \Delta SW_{alb}$ as the change in clear-sky aerosol-free SW fluxes (Eq 1716) whereas the APRP quantifies this as the change in all-sky radiation due to changes in surface albedo. Hence the latter calculation allows the radiative impact of
- 295 changing surface albedo to be attenuated by the presence of clouds and aerosols (i.e., masking effects) and avoids aliasing in contributions from humidity changes that impact SW absorption. Indeed, the double-call $ERF_{alb}^{SW} \Delta SW_{alb}$ values agree closely with the clear-sky surface albedo component diagnosed by APRP scaled by the clear-sky fraction (r=1.0; RMSD=0.02; not shown).

Estimates of global mean (a) ERF^{SW}_{ari+aci}, (b) ERF^{SW}_{alb}, (c) ERF^{SW}_{ari}, and (d) ERF^{SW}_{aci} derived using the APRP method as
implemented in this study (pink) and in Smith et al. (2020) (blue) in each CMIP6 model scattered against estimates computed using the double-call method of Ghan (2013). Note that we are referring to Ghan's ERF^{SW}_{other} as ERF^{SW}_{alb} here even though the former receives contributions from more than just changes in surface albedo (see Appendix A). The correlation and root mean square difference are reported as r and RMSD, respectively, and correlations that are statistically significant at 95% confidence are indicated with an asterisk.

The multi-model mean maps show excellent agreement between the two methodologies in the spatial structure of each component between the correct implementation of APRP and Ghan (2013) methodologies, albeit with quantitative differences (Figure 67). Both methods agree on negative $\text{ERF}_{ari+aci}^{SW}$ over the vast majority of the Northern Hemisphere, with largest negative values over the Northern Indian Ocean and over Southeast Asia. Negative values are also present across the North



Figure 6. Estimates of global mean (a) $\text{ERF}_{ari+aci}^{SW}$ (b) ERF_{ari}^{SW} (c) ERF_{aci}^{SW} , and (d) ΔSW_{alb} derived using the APRP method as implemented in this study (pink) and in Smith et al. (2020) (blue) in each CMIP6 model scattered against estimates computed using the double-call method of Ghan (2013). Note that we are referring to Ghan's ERF_{atber}^{SW} as ΔSW_{alb} here even though the former receives contributions from more than just changes in surface albedo (see Appendix B). The correlation and root mean square difference are reported as r and RMSD, respectively, and correlations that are statistically significant at 95% confidence are indicated with an asterisk.

Pacific and North Atlantic Oceans and over most of the Northern Hemisphere continental regions, and local maxima are present 310 just west of South America and Africa.

Both methods diagnose a positive ERF_{ari}^{SW} over Africa (Figure 67, row 2). The APRP results indicate that this is a region of strong aerosol absorption that is not fully countered by aerosol scattering (not shown), perhaps indicating a role of black carbon from biomass burning. Just east of this, the two methods agree on a negative direct effect over the northern Indian Ocean. Despite the large aerosol emission sources, the ERF_{ari}^{SW} is not particularly large over Southeast Asia because of close

315 compensation between the absorbing and scattering components (not shown). The spatial structure of the total ERF is very consistent with the ERF^{SW}_{aci} map (Figure 67, row 3), highlighting the dominant role played by the indirect effect in models' total aerosol forcing. Weak negative ERF^{SW}_{alb} ΔSW_{alb} values are present over much of the NH continents, likely due to increased



Figure 7. $\text{ERF}_{ari+aci}^{SW}$ (row 1), ERF_{ari}^{SW} (row 2), ERF_{aci}^{SW} (row 3), and $\frac{\text{ERF}_{alb}^{SW}}{\text{alb}}$ (row 4) estimated by the APRP method (column 1), the double-call method (column 2), and their difference (column 3). Area-weighted global mean values are printed in the top right corner of each panel. Results are averaged across all models providing double-call output.

snow cover in response to aerosol-induced cooling (Figure 67, row 4). An exception is a region of large positive $\frac{ERF_{alb}^{SW}}{\Delta SW_{alb}}$ over the Himalayas, which may result from black carbon deposition on snow.

320

325

Consistent with the comparison for a single model shown in Zelinka et al. (2014), locations where APRP overestimates the negative ERF_{ari}^{SW} are co-located with locations where it underestimates ERF_{aci}^{SW} , and vice versa. This is subtle but noticeable over the North Pacific Ocean downwind of the Eastern Asia $\text{ERF}_{ari+aci}^{SW}$ maximum. These opposite-signed errors differences with respect to the double-call double-call method cancel such that the total $\text{ERF}_{ari+aci}^{SW}$ maps are nearly identical (Figure 67, row 1). The global mean root-mean-square difference between APRP and double-call estimates of $\text{ERF}_{ari+aci}^{SW}$ is less than 10% of the standard deviation of either field, confirming that both methods agree as to spatial variations.

Longwave ERF components cannot be derived using APRP, but we can derive proxies for the direct and indirect components from standard model output (Section 2.2.3). Specifically, we use the (negative of the) change in clear-sky OLR as a proxy for the LW direct effect and the change in LW cloud radiative effect (LW CRE) LW radiation with the contribution from changes in surface temperature removed as a proxy for the LW indirect direct effect. Recall that the formulation of Ghan

330 (2013) defines the direct effect as the difference between changes in OLR-LW radiation under aerosol-free conditions and under all-sky conditions (Eq 15). The negative change in clear-sky outgoing longwave radiation (Δ OLR_{cs}) actually shows little correspondence with this definition of ERF^{LW}_{ari} (Figure 7a, blue markers). In contrast, it does closely match 14), As



Figure 8. Estimates of global mean (a) ERF_{ari}^{LW} and (b) ERF_{aci}^{LW} derived using standard output from each CMIP6 model scattered against estimates computed using the double-call method of Ghan (2013). Our proxy for the LW direct effect - the negative change The double-call estimate in elear-sky OLR – is poorly correlated with $\text{ERF}_{ari}^{G,LW}$ (blue markersa) but is well correlated with the sum of $\text{ERF}_{ari}^{G,LW}$ plus and $\text{ERF}_{other}^{G,LW}$ with the surface temperature response removed (orange markerssee Appendix B).

demonstrated mathematically in Appendix B and as shown in Figure 8a, our ERF_ari_proxy is closely related to the sum of Ghan's $\text{ERF}_{ari}^{LW,G}$ and $\text{ERF}_{other}^{LW,G}$ terms , as indicated by the orange markers in Figure 7a and as demonstrated mathematically 335 in Appendix A, with the contribution from changes in surface temperature removed. In other words, ΔOLR_{cs} the change in clear-sky LW radiation is made up of the direct aerosol effect as defined in Ghan (2013) plus an OLR perturbation perturbations due to aerosol-induced fast adjustments of temperature and humidity. These adjustments are clearly larger contributors to both the mean and spread in $-\Delta OLR_{cs}$ than are the direct impact of aerosols on LW radiation. Comparing the vertical spread of the orange and blue markers in Figure 7a, we find that the direct impact of aerosols on LW radiation accounts for less than half as much inter-model spread as the fast atmospheric adjustments. humidity and temperature, including at the surface. 340

The change in LW CRE – our proxy for the LW indirect effect – is very well-correlated with ERF_{aci}^{LW} derived with the double-call method (Figure $\frac{7b8b}{}$), justifying its use as a proxy for the LW indirect effect. The model with large ERF^{LW}_{aci} apparent in Figure 7b-8b is MRI-ESM2-0, whose CMIP5-era counterpart also exhibited a large indirect component in the LW. This model was among only a few in CMIP5 that parameterized aerosol impacts on ice clouds, and it exhibited large changes in 345 the amount and optical depth of high clouds in response to aerosols (Zelinka et al., 2014). This remains the model with largest

aerosol effects on high clouds and therefore on LW cloud-radiative fluxes, as discussed further in Smith et al. (2020).

We conclude from this section that the APRP technique yields aerosol ERF values that agree well with the independent double-call method, both in the global average and in the spatial distribution. Given that it does not require advanced diagnostics that may not be available in many models and experiments, it is an attractive method for systematically diagnosing these values

350

across a broad suite of climate models. The typically smaller LW components of aerosol forcing are likewise well-captured using simpler diagnostics that are widely available.

Table 1. Aerosol ERF_{ari} , ERF_{aci} , and $\text{ERF}_{ari+aci}$ values [W/m²] estimated for individual CMIP5 models, separated into SW, LW, and net components. The ERF_{ari}^{SW} is further separated into scattering and absorption components, and the ERF_{aci}^{SW} is further separated into scattering, absorption, and amount components. Also shown are the multi-model means and across-model standard deviations, both computed using only one ensemble member per model. Forcings are for present-day (year 2000) relative to pre-industrial conditions.

					SW				LW			Net		
Model	ARI			ACI										
	scat	abs	sum	scat	abs	amt	sum	ARI+ACI	ARI	ACI	ARI+ACI	ARI	ACI	ARI+ACI
CSIRO-Mk3-6-0.r1i1p1	-1.13	0.49	-0.64	-0.76	0.08	0.00	-0.68	-1.33	-0.02 0.06	-0.21	-0.23 0.27	-0.66 -0.70	-0.89	-1.55-1.59
CanESM2.r1i1p1	-0.58	0.17	-0.41	-0.52	0.01	-0.01	-0.51	-0.91	0.11- 0.06	-0.04	0.07-0.02	-0.30_0.35	-0.55	-0.85 0.90
FGOALS-s2.r1i1p1	-0.77	0.20	-0.57	0.11	0.01	0.02	0.14	-0.42	0.09- 0.05	-0.06	0.03_0.01	-0.48-0.52	0.08	-0.400.44
GFDL-CM3.r1i1p1	-0.91	0.41	-0.50	-1.00	-0.04	-0.13	-1.17	-1.66	0.11-0.02	0.02	0.13-0.04	-0.39_0.48	-1.15	-1.54-1.63
HadGEM2-A.r1i1p1	-0.53	0.26	-0.27	-0.99	0.06	-0.13	-1.06	-1.33	0.14 0.07	-0.05	0.09-<u>0.02</u>	-0.13_0.20	-1.11	-1.24-1.31
IPSL-CM5A-LR.r1i1p1	-0.66	0.28	-0.38	-0.23	-0.01	0.13	-0.11	-0.48	0.01-0.05	-0.21_0.22	-0.20_0.27	- 0.37_0.43	-0.32_0.33	-0.690.76
MIROC5.r1i1p1	-0.66	0.16	-0.50	-0.93	-0.01	-0.28	-1.22	-1.72	0.22-0.13	0.27	0.49 <u>0.40</u>	-0.28_0.37_	-0.95	-1.231.32
MPI-ESM-LR.r1i1p1	-0.71	0.49	-0.22	0.10	-0.08	-0.01	0.00	-0.22	-0.02-0.07	-0.10	-0.12-0.17	-0.24_0.29	-0.10	-0.34 0.39
MPI-ESM-LR.r1i1p2	-1.15	0.68	-0.47	0.13	-0.07	0.04	0.11	-0.36	-0.03-0.12	-0.17	-0.20 -0.29	-0.50_0.59	-0.06	-0.56 -0.65
MRI-CGCM3.r1i1p1	-0.11	0.12	0.01	-1.79	-0.09	-0.23	-2.12	-2.11	0.00- 0.03	0.95	0.95- 0.98	0.01- 0.04	-1.17	-1.16_ -1.13
NorESM1-M.r1i1p1	-0.63	0.29	-0.34	-0.80	0.03	0.18	-0.59	-0.93	0.13 0.04	-0.17	- 0.04 0.13	-0.21_0.30	-0.76	-0.971.06
bcc-csm1-1.r1i1p1	-0.93	0.23	-0.71	0.23	0.01	0.03	0.26	-0.44	0.06_0.01	-0.05	0.01-0.06	- 0.65 0.72	0.21	-0.440.51
Mean	-0.69	0.28	-0.41	-0.60	-0.00	-0.04	-0.64	-1.05	0.08 0.02	0.03	0.11- 0.05	- 0.34 -0.39	-0.61	- 0.95 1.00
1 - σ	0.25	0.12	0.20	0.58	0.05	0.14	0.68	0.60	0.07- 0.06	0.32	0.33-0.34	0.20 0.21	0.48	0.42

3.3 Summary of Corrected APRP-Derived ERF Values

355

Having established that biases were present in the values of aerosol ERF provided in Smith et al. (2020) and that the values derived herein have negligible residuals and are in better agreement with the double-call method, we now provide the aerosol ERF values for all available models, along with their breakdown into components (Tables 1-2). This includes several CMIP6 models that were not available at the time Smith et al. (2020) was published. Although our APRP implementation is unchanged from that used in Zelinka et al. (2014), we report here CMIP5 results that include three additional models (bcc-csm1-1, FGOALS-s2, and MPI-ESM-LR) that became available after the publication of the were not included in the earlier paper.

Every CMIP6 model agrees on a negative ERF^{SW}_{ari} due to scattering and a positive ERF^{SW}_{ari} due to absorption, but the relative
strengths vary, leading to a lack of agreement on the sign of ERF^{SW}_{ari}. All CMIP6 models have negative ERF^{SW}_{aci} values. This is due to the dominance of strong negative ERF^{SW}_{aci} scattering components, with two exceptions: The r1i1p1f1 and r1i1p1f2 variants of the GISS-E2-1-G model have small positive ERF^{SW}_{aci} cloud scattering components, but anomalously strong negative ERF^{SW}_{aci} cloud amount components. This is consistent with the fact that the 'p1' physics variants of GISS-E2-1-G parameterize aerosol effects on clouds by directly relating anthropogenic aerosol mass to a change in (only) total cloud cover (Miller et al., 2021). In the 'p3' physics variants, in contrast, aerosols act as cloud condensation nuclei and change (only) cloud optical depth;

Table 2. As in Table 1, but for CMIP6 models. Models that provided aerosol-free diagnostics (allowing for the comparisons in Figures 5-7) are indicated with asterisks. Multi-model means and across-model standard deviations are computed using only one ensemble member per model, but we treat the r1i1p1f1 and r1i1p3f1 members of GISS-E2-1-G as separate models. Forcings are for present-day (year 2014) relative to pre-industrial conditions.

					SW				LW			Net		
Model	ARI		ACI											
	scat	abs	sum	scat	abs	amt	sum	ARI+ACI	ARI	ACI	ARI+ACI	ARI	ACI	ARI+ACI
ACCESS-CM2.r1i1p1f1	-0.84	0.42	-0.42	-0.82	-0.01	-0.13	-0.96	-1.37	0.26_0.17	0.05_0.04	0.31-0.21	- 0.16_ 0.25	-0.91_0.92	-1.07_1.17
ACCESS-ESM1-5.r1i1p1f1	-0.44	0.28	-0.16	-1.04	0.05	-0.13	-1.12	-1.28	0.17_0.09	-0.02	0.15- 0.07	0.01-0.07	-1.14	-1.13-1.21
BCC-ESM1.r1i1p1f1	-1.13	0.30	-0.83	-0.62	-0.09	-0.09	-0.79	-1.62	0.16_0.03	0.07-0.06	0.23 0.09	- 0.67 - <u>0.80</u>	-0.72 -0.73	-1.39 -1.53
CESM2.r1i1p1f1	-0.26	0.50	0.23	-1.80	0.04	-0.00	-1.76	-1.53	0.05_0.09	0.10	0.15- 0.19	0.28 0.32	-1.66	-1.38 -1.34
CNRM-CM6-1.rli1p1f2*_rli1p1f2	-0.65	0.29	-0.36	-0.83	-0.07	0.05	-0.85	-1.20	0.15_0.05	-0.05	0.10-0.00	-0.21-0.31	-0.90	-1.11-1.21
CNRM-ESM2-1.r1i1p1f2*_r1i1p1f2	-0.44	0.25	-0.19	-0.53	-0.06	-0.03	-0.62	-0.81	0.10_0.03	-0.02	0.08-<u>0.01</u>	- 0.09_0.16	-0.64	-0.73-0.80
CanESM5. rli1p2f1* r1i1p2f1	-0.64	0.76	0.12	-0.91	0.08	-0.14	-0.98	-0.86	0.06_0.04	-0.08	-0.02 -0.04	0.18_0.16	-1.06	-0.88 -0.90
GFDL-CM4. rli1p1f1* rli1p1f1	-0.68	0.59	-0.09	-0.55	0.00	-0.09	-0.64	-0.73	0.13_0.05	-0.06	0.07_0.01	0.04-0.04	-0.70	-0.66 -0.74
GFDL-ESM4. rli1p1f1* rli1p1f1	-0.68	0.77	0.09	-0.59	0.01	-0.06	-0.64	-0.55	0.06_ 0.00	-0.17	- 0.11-0.17	0.15_0.09	-0.81	- 0.66 0.72
GISS-E2-1-G.r1i1p1f1	-0.94	0.27	-0.67	0.07	0.02	-0.95	-0.86	-1.53	0.19<u>0</u>.12	0.09-<u>0.10</u>	0.28 <u>0.22</u>	-0.48-0.55	-0.77_0.76	-1.25_1.31
GISS-E2-1-G.r1i1p1f2	-0.88	0.20	-0.68	0.07	0.03	-0.84	-0.74	-1.42	0.11_0.06	0.09-<u>0.10</u>	0.20-0.16	- 0.57 - <u>0.62</u>	-0.65 -0.64	-1.22 -1.26
GISS-E2-1-G.r1i1p3f1	-0.99	0.24	-0.75	-0.23	-0.01	-0.07	-0.30	-1.05	0.18 <u>0.12</u>	- 0.06_0.05	0.12-0.07	- 0.57- 0.63	- 0.36 -0.35	-0.93_ 0.98
HadGEM3-GC31-LL.r1i1p1f3	-0.83	0.42	-0.41	-0.78	-0.01	-0.07	-0.86	-1.26	0.20<u>0.12</u>	0.00_0.000	0.20-<u>0.12</u>	- 0.21 0.29	-0.86	-1.07-1.15
IPSL-CM6A-LR-INCA.r1i1p1f1	-0.69	0.20	-0.49	-0.29	-0.01	0.03	-0.28	-0.77	0.09_0.02	-0.07	0.02_0.05	-0.40_0.47	-0.35	-0.75_0.82
IPSL-CM6A-LR.r1i1p1f1	-0.63	0.23	-0.40	-0.27	-0.01	0.09	-0.19	-0.59	0.07_0.00	-0.06 - <u>-0.07</u>	0.01_0.07_0.07_	-0.33_ -0.40	-0.25 -0.26	-0.58_ -0.66
IPSL-CM6A-LR.r2i1p1f1	-0.63	0.23	-0.40	-0.28	-0.01	0.02	-0.28	-0.68	0.06- 0.03	-0.08	- 0.020.11	-0.34-0.43	-0.36	- 0.70 -0.79
IPSL-CM6A-LR.r3i1p1f1	-0.63	0.24	-0.39	-0.25	-0.02	0.10	-0.16	-0.56	0.03 - <u>0.02</u>	-0.05	-0.02_0.07_	- 0.36 -0.41	-0.21	- 0.57_0.62
IPSL-CM6A-LR.r4i1p1f1	-0.63	0.23	-0.40	-0.28	-0.01	0.03	-0.26	-0.66	0.08_0.00	-0.07	0.01_0.07_	-0.320.40	-0.33	-0.650.73
MIROC6.r11i1p1f1	-0.47	0.12	-0.35	-1.13	-0.09	-0.03	-1.24	-1.59	0.19_0.11	0.41	0.60-0.52	- 0.16 -0.24	-0.83	- 0.99_1.07
MIROC6.rli1p1f1	-0.48	0.13	-0.35	-1.13	-0.08	-0.01	-1.22	-1.58	0.17<u>0.10</u>	0.36	0.53-<u>0.46</u>	-0.18-0.25	-0.86	-1.04-1.11
MPI-ESM-1-2-HAM. rlilp1f1* rlilp1f1	-0.26	0.28	0.01	-1.33	0.03	-0.24	-1.54	-1.53	0.16_0.05	0.10	0.26 0.15	0.17 <u>0.06</u>	-1.44	-1.27_ 1.38
MRI-ESM2-0. rli1p1f1*rli1p1f1	-0.76	0.27	-0.48	-1.73	-0.12	-0.38	-2.23	-2.71	0.05_0.01	1.47-<u>1.48</u>	1.52-<u>1.49</u>	-0.43 -0.47	-0.76 -0.75	-1.19 -1.22
NorESM2-LM.rli1p1f1*rli1p1f1	-0.43	0.30	-0.13	-1.16	-0.00	-0.10	-1.27	-1.39	0.07	0.14	0.21	-0.06	-1.13	-1.19
NorESM2-LM. rli1p2f1* rli1p2f1	-0.45	0.31	-0.14	-1.32	0.01	-0.19	-1.50	-1.64	0.09	0.17.0.16	0.26-0.25	-0.05	-1.33_1.34	-1.38_ 1.39
NorESM2-MM. rli1p1f1* rli1p1f1	-0.42	0.32	-0.10	-1.19	0.02	-0.17	-1.34	-1.44	0.15<u>0</u>.11	0.04	0.19 <u>0.15</u>	0.05_0.01_	-1.30	-1.25_ -1.29
UKESM1-0-LL. rli1p1f4* rli1p1f4	-0.76	0.48	-0.28	-0.85	-0.01	-0.10	-0.95	-1.23	0.18_0.11	-0.01	0.17- 0.10	- 0.10_0.17	-0.96	-1.06_ 1.13
Mean	-0.65	0.37	-0.28	-0.83	-0.01	-0.13	-0.97	-1.25	0.13_0.07	0.09	0.23 0.16	- 0.15_0.21	-0.88	-1.03_ 1.09
1-σ	0.23	0.17	0.28	0.47	0.05	0.21	0.49	0.47	0.06_0.05	0.34	0.33	0.26 0.28	0.34	0.24

hence the r1i1p3f1 variant of GISS-E2-1-G has a moderate negative ERF_{aci}^{SW} cloud scattering component and weak ERF_{aci}^{SW} cloud amount component (Table 2). In all models, $\text{ERF}_{ari+aci}^{SW}$ is negative.

 ERF_{ari}^{LW} is systematically positive across positive in nearly all CMIP6 models but is small with opposite sign relative to its SW counterpart. ERF_{aci}^{LW} values are generally small except for the models identified in Smith et al. (2020) as parameterizing aerosol effects on ice clouds (CESM2, MIROC6, MRI-ESM2-0 and NorESM2-LM), for which the ERF_{aci}^{LW} values are non-negligible and positive. Finally, the net (LW+SW) $\text{ERF}_{ari+aci}$ is systematically negative across all models, primarily due to the systematically negative indirect component that is generally larger in magnitude than the direct component, which is small or also negative.



Figure 9. Global mean $\text{ERF}_{ari+aci}$ values averaged across (a) CMIP5 and (b) CMIP6 models, separated into ERF_{ari} and ERF_{aci} for LW, SW, and net (LW+SW) radiation. ERF_{ari}^{SW} is further separated into its scattering and absorption components. ERF_{aci}^{SW} , is further separated into its amount, scattering, and absorption components. The sum of terms in each row is indicated by the black dot, with the inter-model standard deviation of each sum indicated by the horizontal error bar.

Figure 8, which shows the CMIP6-

To aid in visualizing the model results and to connect back to Figure 10 in Zelinka et al. (2014) and Figure 10 in Smith et al. (2020), multi-model mean ERF values are shown for both the CMIP5 and CMIP6 models in Figure 9. We strongly caution against over-interpreting differences between the two ensembles for two reasons: First, they differ in how "present-day" is defined (2000 for CMIP5 and 2014 for CMIP6). Second, the fraction of models that represent aerosol effects on cloud optical properties and/or lifetimes differs, with three CMIP5 models (FGOALS-s2, is qualitatively very similar to its counterpart Figure 10 of
Smith et al. (2020), with a few quantitative differences: A weaker negative net ERFMPI-ESM-LR, and bcc-csm1-1) lacking any representation of indirect effects, some models representing only the first indirect effect, and some representing both the first and second indirect effects (Rotstayn et al., 2013; Wilcox et al., 2013). With these caveats in mind, we note that the

negative net (LW+SW) ERF_{aci} is nearly 50% stronger in magnitude in CMIP6, while the smaller net ERF_{ari} is diagnosed here because the positive absorbing component of ERF^{SW}_{ari} is increased by more than 30% in the present study. Conversely, a stronger roughly halved in CMIP6. Close compensation between these two changes mean that the overall net ERF_{ari+aci} is only slightly larger in CMIP6. The negative ERF_{aci} is diagnosed because the negative scattering component of ERF^{SW}_{aci} has strengthened slightly, stronger in CMIP6 due primarily to stronger SW cooling from a greater sensitivity of cloud scattering and amount to aerosols (partly related to the fact that three models in CMIP5 do not incorporate indirect effects). The negative ERF_{ari} is weaker in CMIP6 due primarily to stronger SW heating from absorbing aerosols. These same conclusions hold even

390 if we compare CMIP5 and CMIP6 averages based only on models from centers that contributed to both phases (not shown). Global mean $\text{ERF}_{ari+aci}$ values averaged across CMIP6 models, separated into ERF_{ari} and ERF_{aci} for LW, SW, and net (LW+SW) radiation. ERF_{ari}^{SW} is further separated into its scattering and absorption components. ERF_{aci}^{SW} , is further separated into its amount, scattering, and absorption components. The sum of terms in each row is indicated by the black diamond, with the inter-model standard deviation of each sum indicated by the horizontal error bar.

395 4 Conclusions

Accurate values of radiative forcings across a broad suite of climate models is a prerequisite for proper understanding of the drivers of inter-model differences in climate response. Model-to-model differences in aerosol radiative forcing are particularly large and come from both aerosol direct and indirect components, each of which has competing contributions from changes in scattering and absorption of SW radiation. In this study we corrected estimates of aerosol effective radiative forcing derived

- 400 in Smith et al. (2020) from a collection of CMIP6 models that performed idealized aerosol perturbation experiments. We also provided values from additional CMIP6 models that became available subsequent to its publication (as well as CMIP5 models previously reported, for completeness). The errors in the previous study resulted from two larger errors that, fortuitously, largely cancelled in the global mean, though for certain sub-components these errors do not cancel and are non-negligible. Most notably, the direct radiative forcing from absorbing aerosols averaged across CMIP6 models is more than 40% larger
- 405 <u>in the present study</u>. Aerosol effective radiative forcings derived herein have negligible residuals and agree well with values derived using an independent double radiation call technique, both in the global mean and in geographic structure. Code to perform the accurate APRP method is provided at the link in the *Code availability* section.

Code availability. Code to perform the calculations in this study is available at https://doi.org/10.5281/zenodo.7809085 (Zelinka, 2021).

Data availability. All CMIP climate model data used in this study is available from the Earth System Grid Federation (https://esgf.llnl.gov/).

410 *Author contributions.* All analyses in the paper were performed by MDZ. The first draft of the manuscript was written by MDZ and all authors commented on subsequent versions of the manuscript. All authors read and approved the final manuscript.

Competing interests. The authors declare no competing interests.

Acknowledgements. We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP. We thank the climate modeling groups for producing and making available their model output, the Earth

- 415 System Grid Federation (ESGF) for archiving the data and providing access, and the multiple funding agencies who support CMIP and ESGF. The work of MDZ and KET was MDZ, YQ, and KET were supported by the U.S. Department of Energy (DOE) Regional and Global Model Analysis program areaand. CJS was supported by a NERC/IIASA Collaborative Research Fellowship (NE/T009381/1). The work of MDZ and KET and was performed under the auspices of the U.S. DOE by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. CJS was supported by a NERC/IIASA Collaborative Research Fellowship (NE/T009381/1). The Pacific Northwest
- 420 National Laboratory is operated for the DOE by the Battelle Memorial Institute under Contract DE-AC05-76RL01830. We thank Susanne Bauer for assistance interpreting the GISS results as well as two anonymous reviewers and editor Yuan Wang for helpful comments that improved the manuscript.

Appendix A

A.1 Introduction

425 In this Appendix, we provide further explanation of the APRP technique and how it is to be used correctly. We then explain the two implementation errors in Smith et al. (2020), the first involving the calculation of albedo sensitivities and the second involving usage of incorrect SW fluxes at the top of atmosphere.

A.2 APRP

As detailed in Taylor et al. (2007), the planetary albedo (A) can be written in terms of the surface albedo (α), an atmospheric transmittance coefficient (μ), and an atmospheric scattering coefficient (γ):

$$A = \mu\gamma + \frac{\mu\alpha(1-\gamma)^2}{1-\alpha\gamma}$$
(A1)

Surface albedo is computed as the ratio of upwelling to downwelling SW fluxes at the surface:

$$\alpha = SW_{SFC}^{\uparrow} / SW_{SFC}^{\downarrow}.$$
(A2)

Atmospheric transmittance is expressed as

435
$$\mu = A + \hat{Q}_{SFC}^{\downarrow}(1-\alpha),$$
 (A3)

where

$$\hat{Q}_{SFC}^{\downarrow} = \frac{SW_{SFC}^{\downarrow}}{SW_{TOA}^{\downarrow}} \tag{A4}$$

is the ratio of surface to TOA incident SW flux, and the scattering coefficient is expressed as

$$\gamma = \frac{\mu - \hat{Q}_{SFC}^{\downarrow}}{\mu - \alpha \hat{Q}_{SFC}^{\downarrow}}.$$
(A5)

440 All of the terms given in A2 - A5 can be computed for both clear-sky fluxes and for overcast-sky fluxes. This separation into clear- and overcast-sky conditions, along with some assumptions, allow for the effects of clouds to be isolated from other atmospheric constituents (primarly aerosols in the present study). Specifically, the scattering coefficient in the overcast portion of a scene can be expressed as a combination of the scattering coefficients due to non-cloud constituents and the cloud itself:

$$(1 - \gamma_{oc}) = (1 - \gamma_{clr})(1 - \gamma_{cld}),\tag{A6}$$

445 where subscripts *clr* and *oc* refer to clear- and overcast-sky conditions. Similarly, the transmissivity coefficients are related as:

$$\mu_{oc} = \mu_{clr} \mu_{cld}. \tag{A7}$$

This allows total planetary albedo to be expressed as the sum of clear-sky albedo scaled by the clear-sky portion of the scene and overcast-sky albedo scaled by the cloudy portion of the scene:

$$A = (1-c)A_{clr} + cA_{oc},\tag{A8}$$

450 where c is the total cloud fraction. Hence, the planetary albedo is a function of seven parameters:

$$A = A(c, \alpha_{clr}, \alpha_{oc}, \mu_{clr}, \mu_{cld}, \gamma_{clr}, \gamma_{cld}).$$

The derivation of these expressions (and the assumptions made in doing so) are detailed in Taylor et al. (2007), and so are not further explained here.

(A9)

Having determined all of the terms upon which planetary albedo depends, we can now substitute in values of individual 455 coefficients from the perturbed climate experiment to isolate their individual impacts on albedo. For example, to determine the impact on planetary albedo from the change in aerosol scattering, we difference the albedo computed with (only) γ_{clr} taken

from the perturbed experiment (indicated with superscript 'pert') and the albedo computed with all fields set to their control state (no superscript):

$$\frac{\partial A}{\partial \gamma_{clr}} \Delta \gamma_{clr} = A(c, \alpha_{clr}, \alpha_{oc}, \mu_{clr}, \mu_{cld}, \gamma_{clr}^{pert}, \gamma_{cld}) - A(c, \alpha_{clr}, \alpha_{oc}, \mu_{clr}, \mu_{cld}, \gamma_{clr}, \gamma_{cld}).$$
(A10)

460 In practice, we do this calculation twice – once as a forward calculation as shown above and once as a backward calculation in which all fields are set to their perturbed value and the isolated field is set to its control state – and then these are averaged.

A.3 The Albedo Sensitivity Error in Smith et al. (2020)

Smith et al. (2020) did not use A9 to estimate their components but rather followed a different procedure. In the example considered above, they compute the impact on planetary albedo from the change in aerosol scattering as the sum of two terms:

$$465 \quad \frac{\partial A}{\partial \gamma_{clr}} \Delta \gamma_{clr} = (1-c) \frac{\partial A_{clr}}{\partial \gamma_{clr}} \Delta \gamma_{clr} + c \frac{\partial A_{oc}}{\partial \gamma_{clr}} \Delta \gamma_{clr}, \tag{A11}$$

where the responses of A_{clr} and A_{oc} to perturbations in γ_{clr} are computed by substituting directly into Eq A1. The error arises for the latter term, which Smith et al. compute as:

$$\frac{\partial A_{oc}}{\partial \gamma_{clr}} \Delta \gamma_{clr} = A(\alpha_{oc}, \mu_{oc}, \gamma_{clr}^{pert}) - A(\alpha_{oc}, \mu_{oc}, \gamma_{clr}).$$
(A12)

Note that in A12 the *clear-sky* scattering coefficients are erroneously used to compute *overcast-sky* albedo. The correct calculation is:

$$\frac{\partial A_{oc}}{\partial \gamma_{clr}} \Delta \gamma_{clr} = A(\alpha_{oc}, \mu_{oc}, \gamma_{oc*}^{pert}) - A(\alpha_{oc}, \mu_{oc}, \gamma_{oc}), \tag{A13}$$

where we have defined γ_{gc*}^{pert} via A6:

$$(1 - \gamma_{oc*}^{pert}) = (1 - \gamma_{clr}^{pert})(1 - \gamma_{cld}).$$
(A14)

This modified overcast-sky scattering coefficient allows changes in the clear-sky scattering coefficient to independently affect

- 475 the overcast-sky scattering coefficient holding the cloud scattering fixed. Hence it incorporates the impact on atmospheric scattering under overcast-sky conditions from (solely) changes in clear-sky (aerosol) scattering, which is the coefficient needed to correctly estimate the influence of changes in clear-sky (aerosol) scattering on overcast-sky albedo. This is already implicitly done in the correct method involving substitution into Eq A10. It is only when substituting coefficients directly into Eq A1 (as done by Smith et al.) that one has to separately account for this.
- 480 Why does this error cause an overestimate of the sensitivity of TOA albedo to aerosols relative to the correct implementation of APRP? The difference between scattering coefficients in the erroneous expression (A12) is:

$$\Delta \gamma_{smith} = (\gamma_{clr}^{pert} - \gamma_{clr}). \tag{A15}$$

In contrast, after some algebra one can show that the difference between scattering coefficients in the corrected expression (A13) is:

$$485 \quad \underline{\Delta\gamma_{true} = \gamma_{oc*}^{pert} - \gamma_{oc} = (1 - \gamma_{cld})(\gamma_{clr}^{pert} - \gamma_{clr}),} \tag{A16}$$

which is equal to $\Delta \gamma_{smith}$ scaled by $1 - \gamma_{cld}$. This scaling factor, which is less than 1, represents the attenuation of aerosol effects by the presence of clouds: The larger the cloud scattering coefficient, the greater the attenuation and hence the weaker the aerosol influence on TOA albedo. Therefore the correct formulation of APRP, which accurately accounts for this attenuation, has a weaker sensitivity of TOA albedo to changes in clear-sky (aerosol) scattering. In other words, Smith's scattering coefficient changes are overestimated by the factor

$$\frac{\Delta\gamma_{smith}}{\Delta\gamma_{true}} = \frac{1}{1 - \gamma_{cld}} \tag{A17}$$

We note that similar coding errors are present in Smith et al.'s calculation of the sensitivity of overcast-sky albedo to aerosol absorption, cloud scattering, and cloud absorption. We find that the errors in the sensitivity of overcast-sky albedo to aerosol and cloud absorption are small in practice because cloud absorption of SW radiation is small. We do not detail these here for the sake of brevity.

A.4 The Insolation Error in Smith et al. (2020)

All of the aforementioned APRP calculations yield sensitivities of planetary albedo (A) to perturbations in the seven parameters noted in A9. These need to be multiplied by the insolation at the TOA in order to determine the impact on the TOA SW energy budget. The net absorbed SW radiation at the TOA is expressed as

500
$$SW_{abs} = S(1-A),$$

490

495

(A18)

where S is the SW flux at the top of the atmosphere and A is the planetary albedo. Its sensitivity to A is therefore

$$\frac{\partial SW_{abs}}{\partial A} = -S. \tag{A19}$$

In Smith et al.'s formulation, S is

$$S_{smith} = (SW_{TOA}^{\downarrow} - SW_{TOA}^{\uparrow}) \tag{A20}$$

505 whereas it should be

$$S_{true} = SW_{TOA}^{\downarrow} \tag{A21}$$

By using the net (downwelling minus upwelling) SW flux, Smith et al's values of the TOA radiative impact are therefore underestimated by the factor

$$S_{smith}/S_{true} = 1 - \frac{SW_{TOA}^{\uparrow}}{SW_{TOA}^{\downarrow}} = 1 - A, \tag{A22}$$

510 which is roughly 70%.

Appendix B

B.1 Introduction

In this Appendix, we describe in greater detail how the various aerosol effective radiative forcings relate to each other and to the IPCC AR6 definitions (Forster et al., 2021). Specifically, we relate the LW ERFs defined by IPCC to the LW ERF proxies

515 derived using standard model output in Section AB.2and... We then relate the IPCC SW and LW ERF proxies to the ERFs derived using the Ghan (2013) double-call method in Section A(Section B.3. We then relate.). Finally, the ERFs derived by the Ghan (2013) double-call method are related to those derived using the Taylor et al. (2007) APRP method (Section AB.4) and to the LW ERF proxies derived using standard model output (Section AB.5).

B.2 LW ERFs: Proxies Derived from Standard Model Output

520 One can expand our expression for $ERF_{ari}^{P,LW}$ as:

$$ERF_{ari}^{P,LW} = \Delta R_{cs}^{LW} - \Delta R_{cs}^{T_0} = IRF_{ari,cs}^{LW} + K_{cs}^T \Delta T + K_{cs}^{q,LW} \Delta q.$$
(B1)

Given the IPCC definition of direct effective radiative forcing (Eq 2), we can express our proxy as

$$ERF_{ari}^{P,LW} = ERF_{ari}^{LW} - K^{C,LW}\Delta C_{semidirect} - M_{cld}^{LW},$$
(B2)

where

525
$$M_{cld}^{LW} = (IRF_{ari}^{LW} - IRF_{ari,cs}^{LW}) + (K^T - K_{cs}^T)\Delta T + (K^{q,LW} - K_{cs}^{q,LW})\Delta q.$$
 (B3)

Therefore in the LW, our proxy for the direct effect equals IPCC's direct effect, minus the semidirect effect, minus masking terms that quantify how much the radiative impact of rapid changes in temperature, humidity, and aerosols are attenuated by the presence of clouds.

Turning now to the LW indirect effect, we note that the change in all-sky TOA net LW radiation is given by:

530
$$\Delta R^{LW} = IRF_{ari}^{LW} + K^T \Delta T + K^{q,LW} \Delta q + K^{C,LW} \Delta C.$$
(B4)

Therefore,

$$ERF_{aci}^{P,LW} = K^{C,LW}\Delta C + M_{cld}^{LW},\tag{B5}$$

where M_{cld}^{LW} is defined above. Putting this in terms of IPCC nomenclature:

$$ERF_{aci}^{P,LW} = ERF_{aci}^{LW} + K^{C,LW}\Delta C_{semidirect} + M_{cld}^{LW}.$$
(B6)

535 Therefore in the LW, our proxy for the indirect effect equals IPCC's indirect effect, plus the semidirect effect, plus masking terms that quantify how much the radiative impact of changes in temperature, humidity, and aerosols are attenuated by the presence of clouds. The sum of the direct and indirect LW effects are the same, independent of how the individual components are defined. Thus, from Eqs A2 and A6 B2 and B6

$$ERF_{ari}^{P,LW} + ERF_{aci}^{P,LW} = ERF_{ari}^{LW} + ERF_{aci}^{LW}.$$
(B7)

540 B.3 Double Radiation Call Method

B.3.1 Ghan's Direct Effect

Expanding Eq 1514, one can express the change in TOA net radiation as

$$\Delta R = ERF_{ari}^G + K_{af}^T \Delta T + K_{af}^q \Delta q + K_{af}^\alpha \Delta \alpha + K_{af}^C \Delta C, \tag{B8}$$

or, equivalently,

545
$$\Delta R = IRF_{ari} + K^T \Delta T + K^q \Delta q + K^\alpha \Delta \alpha + K^C \Delta C$$
(B9)

Combining the previous two equations yields an expression for Ghan's ERF_{ari}^G in terms of the true instantaneous aerosol direct forcing:

$$ERF_{ari}^G = IRF_{ari} - M_{aer},\tag{B10}$$

where the aerosol masking is given by:

550
$$M_{aer} = (K_{af}^T - K^T)\Delta T + (K_{af}^q - K^q)\Delta q + (K_{af}^\alpha - K^\alpha)\Delta \alpha + (K_{af}^C - K^C)\Delta C.$$
 (B11)

Therefore Ghan's direct aerosol radiative forcing equals IPCC's instantaneous direct forcing minus masking terms that quantify how much the radiative impact of changes in temperature, humidity, surface albedo, and clouds are attenuated by the presence of aerosols.

B.3.2 Ghan's Indirect Effect

555 Turning now to the indirect effect, let us write the change in TOA energy budget change as:

$$\Delta R = ERF_{ari}^G + ERF_{aci}^G + K_{af,cs}^T \Delta T + K_{af,cs}^q \Delta q + K_{af,cs}^\alpha \Delta \alpha.$$
(B12)

Combining Equations A9 and A11 Defining

$$M_{aer,cld} = (K_{af,cs}^T - K^T)\Delta T + (K_{af,cs}^q - K^q)\Delta q + (K_{af,cs}^\alpha - K^\alpha)\Delta\alpha, \tag{B13}$$

which measures how much the radiative impact of changes in temperature, humidity, and surface albedo are masked by clouds
 and aerosols, and combining Eqs B9 and B11 yields an expression for the total cloud-induced radiation anomalies in terms of ERF^G_{aci}:

$$K^C \Delta C = ERF_{ari}^G + ERF_{aci}^G - IRF_{ari} + M_{aer,cld}.$$
(B14)

where

$$M_{aer,cld} = (K_{af,cs}^T - K^T)\Delta T + (K_{af,cs}^q - K^q)\Delta q + (K_{af,cs}^\alpha - K^\alpha)\Delta \alpha$$

565 Given Eq A10Given Eq B10, we can therefore write:

$$K^C \Delta C = ERF^G_{aci} - M_{aer} + M_{aer,cld} \tag{B15}$$

or equivalently,

$$K^{C}\Delta C = ERF_{aci}^{G} + (K_{af,cs}^{T} - K_{af}^{T})\Delta T + (K_{af,cs}^{q} - K_{af}^{q})\Delta q + (K_{af,cs}^{\alpha} - K_{af}^{\alpha})\Delta \alpha + (K^{C} - K_{af}^{C})\Delta C.$$
(B16)

We can now express Ghan's ERF_{aci}^G in terms of IPCC's aerosol indirect forcing:

570
$$ERF_{aci}^G = ERF_{aci} + M_{aer} - M_{aer,cld} + K^C \Delta C_{semidirect}.$$
 (B17)

Therefore Ghan's indirect effect equals IPCC's indirect effect plus masking terms that quantify how much the radiative impact of changes in temperature, humidity, and surface albedo are attenuated by the presence of clouds under aerosol-free conditions and how much the radiative impact of changes in clouds are attenuated by the presence of aerosols, plus the semidirect effect.

B.3.3 Ghan's Other Forcing Term

575 Finally, let us separate the third forcing term defined by Ghan (2013) into its LW and SW components:

$$ERF_{other}^{G,LW} = K_{af,cs}^T \Delta T + K_{af,cs}^{q,LW} \Delta q, \tag{B18}$$

and

$$ERF_{other}^{G,SW} = K_{af,cs}^{q,SW} \Delta q + K_{af,cs}^{\alpha} \Delta \alpha.$$
(B19)

Given Eq.5, we can express-We can expand $ERF_{other}^{G,SW}$ in terms of IPCC's surface albedo forcing: as:

580
$$ERF_{other}^{G,SW} = \underbrace{ERF \Delta R^{SW}}_{af,cs} Aq + (K^{\alpha}_{af,cs} - K^{\alpha})\Delta\alpha.$$
(B20)

Therefore $ERF_{other}^{G,SW}$ – which Ghan (2013) refers to as the surface albedo forcing – equals $ERF_{alb} \Delta R_{alb}^{SW}$ plus the aerosolfree clear-sky radiative contributions from changes in humidity, plus a masking term that quantifies how much the radiative impact of changes in surface albedo are attenuated by the presence of both clouds and aerosols.

B.4 Relating SW ERF Terms: Ghan (2013) vs APRP

585 Combining Eqs 7 and A10 B10 yields the relationship between Ghan- and APRP-derived SW direct radiative forcing:

$$ERF_{ari}^{A,SW} = ERF_{ari}^{G,SW} + M_{aer}^{SW} + K^{q,SW}\Delta q,$$
(B21)

Because $K^{C,SW}\Delta C$ is equivalent to $ERF_{aci}^{A,SW}$ (cf Eqs 6 and 8), 5 and 7), a simple substitution into Eq B15 (applied to the shortwave) yields the relationship between the relation between APRP and Ghan (2013) expressions for representations of the SW indirect effectis already expressed in Eq A15:

590
$$ERF_{aci}^{A,SW} = ERF_{aci}^{G,SW} - M_{aer}^{SW} + M_{aer,cld}^{SW}$$
 (B22)

Similarly the relation between APRP's surface albedo forcing component and Ghan's equivalent $(ERF_{other}^{G,SW})$ is already expressed in Eq A20:-B20:

$$\underline{ERF_{alb}} \underline{\Delta R}^{A,SW} \underline{alb} = ERF_{other}^{G,SW} - K_{af,cs}^{q,SW} \Delta q - (K_{af,cs}^{\alpha} - K^{\alpha}) \Delta \alpha.$$
(B23)

B.5 Relating LW ERF Terms: Ghan (2013) vs Proxies

595 Given Eqs 13 Subtracting Eqs 14 and 16 from Eq. 11 and rearranging terms, 15, and 17, we can write:

$$ERF_{ari}^{P,LW} = (ERF_{ari}^{G,LW} + ERF_{other}^{G,LW}) + (\Delta R_{cs}^{LW} - \Delta R_{af,cs}^{LW}) - (\Delta R^{LW} - \Delta R_{af}^{LW})_{cs}^{T_0} + \epsilon,$$
(B24)

Because-where

$$\epsilon = (\Delta R_{cs}^{LW} - \Delta R^{LW}) - (\Delta R_{af,cs}^{LW} - \Delta R_{af}^{LW}) = \Delta CRE^{LW} - \Delta CRE_{af}^{LW}$$
(B25)

is very small because the change in LW TOA radiative fluxes should not depend on the presence or absence of aerosols, ΔR_{af}^{LW} 600 is equivalent to ΔR^{LW} (and similarly for the clear-sky conditions). Therefore,

$$ERF_{ari}^{P,LW} = ERF_{ari}^{G,LW} + ERF_{other}^{G,LW}$$

whereas-

$$ERF_{ari}^{P,LW} = ERF_{ari}^{G,LW} - \Delta R^{LW} + \Delta R_{cs}^{LW} + \Delta R_{af}^{LW}.$$

cloud radiative effect is roughly the same under aerosol-free and aerosol-present conditions. This means that our proxy for longwave ERF_{ari} is more closely related to $(ERF_{ari}^G + ERF_{other}^G)$ than to ERF_{ari}^G alone, as demonstrated in Figure 7adirect effect equals the sum of Ghan's direct effect and "other" term plus adjustments that account for the radiative impact of surface temperature change and differences in the CRE response under aerosol-free and aerosol-present conditions. Combining Eqs A6 and A17-B6 and B17 yields the relationship between the Ghan- and proxy-derived estimates of the LW indirect effect:

$$ERF_{aci}^{P,LW} = ERF_{aci}^{G,LW} - M_{aer}^{LW} + M_{aer,cld}^{LW} + M_{cld}^{LW}.$$
(B26)

610 In the LW, M_{aer} is zero, so

$$ERF_{aci}^{P,LW} = ERF_{aci}^{G,LW} + M_{aer,cld}^{LW} + M_{cld}^{LW}.$$
(B27)

This means that our proxy for longwave indirect effect equals Ghan's indirect effect plus masking terms quantifying how strongly clouds attenuate the LW impact of changes in temperature, humidity, and aerosols and how strongly clouds and aerosols together attenuate the LW impact of changes in temperature and humidity.

Table A1. Climate model output used in this study. APRP requires the first eight fields. LW ERF components are estimated using the following two fields. These ten fields are routinely diagnosed in CMIP5 and CMIP6 models. For the double-call method, we rely upon aerosol-free radiative fluxes, which are the final four fields. These are only available for a subset of CMIP6 models. Aerosol-free upwelling SW radiation fluxes at the top of the atmosphere from the NorESM2-LM and NorESM2-MM models were found to be actually the *net* (downwelling minus upwelling) aerosol-free SW fluxes, and were corrected prior to usage.

Description	Variable name
Total cloud fraction	clt
TOA downwelling SW radiation	rsdt
TOA upwelling SW radiation	rsut
TOA upwelling SW radiation under clear-sky conditions	rsutes
Surface downwelling SW radiation	rsds
Surface downwelling SW radiation under clear-sky conditions	rsdscs
Surface upwelling SW radiation	rsus
Surface upwelling SW radiation under clear-sky conditions	rsuscs
TOA outgoing longwave radiation	rlut
TOA outgoing longwave radiation under clear-sky conditions	rlutcs
TOA upwelling SW radiation under aerosol-free conditions	rsutaf
TOA upwelling SW radiation under aerosol-free clear-sky conditions	rsutcsaf
TOA outgoing LW radiation under aerosol-free conditions	rlutaf
TOA outgoing LW radiation under aerosol-free clear-sky conditions	rlutcsaf

615 References

- Armour, K. C. and Roe, G. H.: Climate commitment in an uncertain world, Geophysical Research Letters, 38, L01707, https://doi.org/10.1029/2010GL045850, 2011.
- Dvorak, M. T., Armour, K. C., Frierson, D. M. W., Proistosescu, C., Baker, M. B., and Smith, C. J.: Estimating the timing of geophysical commitment to 1.5 and 2.0 °C of global warming, Nature Climate Change, 12, 547–552, https://doi.org/10.1038/s41558-022-01372-y,

620

0 number: 6 Publisher: Nature Publishing Group, 2022.

- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D. J., Mauritsen, T., Palmer, M. D., Watanabe, M., Wild, M., and Zhang, X.: The Earth's energy budget, climate feedbacks, and climate sensitivity, in: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M.,
- Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, , Yu, R., and Zhou, B., Cambridge University Press, 2021.

Abbreviation	Description						
$\stackrel{\Delta}{\sim}$	Change between pre-industrial and present-day						
\approx	Surface albedo						
APRP	Approximate partial radiative perturbation						
$\underset{\sim}{C}$	Cloud fraction						
\underbrace{CRE}	Cloud radiative effect						
\underbrace{ERF}	Effective radiative forcing						
ERFaci	ERF due to aerosol-cloud interactions						
ERFaciabs	ERFaci due to cloud absorption						
ERFaci.amt	ERFaci due to cloud amount						
$\underbrace{ERF_{aci,scat}}$	ERFaci due to cloud scattering						
ERFari	ERF due to aerosol-radiation interactions						
ERFari,abs	ERFari due to aerosol absorption						
$\underbrace{ERF_{ari,scat}}$	ERFari due to aerosol scattering						
$\underset{\sim}{IRF}$	Instantaneous radiative forcing						
$\underset{\sim}{\overset{K^{\chi}}{\sim}}$	Sensitivity of top of atmosphere radiation to χ						
$\underset{\sim}{LW}$	Longwave radiation						
q_{\sim}	Specific humidity						
$\stackrel{R}{\sim}$	Net (downwelling minus upwelling) top of atmosphere radiation						
ΔR_{alb}	Change in top of atmosphere radiation due to surface albedo changes						
$\underset{\sim}{SW}$	Shortwave radiation						
\sum_{∞}	Temperature						
TOA	Top of atmosphere						
A	Superscript indicating value was computed via APRP technique						
G	Superscript indicating value was computed via Ghan (2013) technique						
$\stackrel{P}{\sim}$	Superscript indicating value is a proxy for ERF derived from standard model output						
af_{\sim}	Subscript indicating aerosol-free radiative fluxes						
$\stackrel{cs}{\sim}$	Subscript indicating clear-sky radiative fluxes						

Ghan, S. J.: Technical Note: Estimating aerosol effects on cloud radiative forcing, Atmospheric Chemistry and Physics, 13, 9971–9974, https://doi.org/10.5194/acp-13-9971-2013, 2013.

Gryspeerdt, E., Mülmenstädt, J., Gettelman, A., Malavelle, F. F., Morrison, H., Neubauer, D., Partridge, D. G., Stier, P., Takemura, T.,

630 Wang, H., Wang, M., and Zhang, K.: Surprising similarities in model and observational aerosol radiative forcing estimates, Atmospheric Chemistry and Physics, 20, 613–623, https://doi.org/10.5194/acp-20-613-2020, publisher: Copernicus GmbH, 2020.

- Huang, Y., Xia, Y., and Tan, X. X.: On the pattern of CO2 radiative forcing and poleward energy transport, Journal of Geophysical Research-Atmospheres, 122, 10578–10593, https://doi.org/10.1002/2017jd027221, 2017.
- Miller, R. L., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Kelley, M., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer,
 M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi,
- G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezuman, K., Murray, L. T., Oinas, V., Orbe, C., Pérez García-Pando, C., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: CMIP6 Historical Simulations (1850–2014) With GISS-E2.1, Journal of Advances in Modeling Earth Systems, 13, e2019MS002 034, https://doi.org/10.1029/2019MS002034, _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2019MS002034, 2021.
- Persad, G. G., Samset, B. H., and Wilcox, L. J.: Aerosols must be included in climate risk assessments, Nature, 611, 662–664, https://doi.org/10.1038/d41586-022-03763-9, bandiera_abtest: a Cg_type: Comment Number: 7937 Publisher: Nature Publishing Group Subject_term: Climate change, Atmospheric science, Policy, 2022.

Pincus, R., Forster, P. M., and Stevens, B.: The Radiative Forcing Model Intercomparison Project (RFMIP): experimental protocol for CMIP6,

645 Geoscientific Model Development, 9, 3447–3460, https://doi.org/10.5194/gmd-9-3447-2016, 2016.

- Rotstayn, L. D., Collier, M. A., Chrastansky, A., Jeffrey, S. J., and Luo, J. J.: Projected effects of declining aerosols in RCP4.5: unmasking global warming?, Atmos. Chem. Phys., 13, 10883–10905, https://doi.org/10.5194/acp-13-10883-2013, 2013.
 - Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., Hegerl, G., Klein, S. A., Marvel, K. D., Rohling, E. J., Watanabe, M., Andrews, T., Braconnot, P., Bretherton, C. S., Foster, G. L., Hausfather, Z., Heydt, A. S. v. d.,
- 650 Knutti, R., Mauritsen, T., Norris, J. R., Proistosescu, C., Rugenstein, M., Schmidt, G. A., Tokarska, K. B., and Zelinka, M. D.: An Assessment of Earth's Climate Sensitivity Using Multiple Lines of Evidence, Reviews of Geophysics, 58, e2019RG000678, https://doi.org/https://doi.org/10.1029/2019RG000678, 2020.
 - Smith, C. J., Kramer, R. J., Myhre, G., Alterskjær, K., Collins, W., Sima, A., Boucher, O., Dufresne, J.-L., Nabat, P., Michou, M., Yukimoto,
 S., Cole, J., Paynter, D., Shiogama, H., O'Connor, F. M., Robertson, E., Wiltshire, A., Andrews, T., Hannay, C., Miller, R., Nazarenko,
- 655 L., Kirkevåg, A., Olivié, D., Fiedler, S., Lewinschal, A., Mackallah, C., Dix, M., Pincus, R., and Forster, P. M.: Effective radiative forcing and adjustments in CMIP6 models, Atmospheric Chemistry and Physics, 20, 9591–9618, https://doi.org/10.5194/acp-20-9591-2020, publisher: Copernicus GmbH, 2020.
 - Taylor, K. E., Crucifix, M., Braconnot, P., Hewitt, C. D., Doutriaux, C., Broccoli, A. J., Mitchell, J. F. B., and Webb, M. J.: Estimating Shortwave Radiative Forcing and Response in Climate Models, J. Climate, 20, 2530–2543, https://doi.org/10.1175/JCLI4143.1, 2007.
- 660 Watson-Parris, D. and Smith, C. J.: Large uncertainty in future warming due to aerosol forcing, Nature Climate Change, 12, 1111–1113, https://doi.org/10.1038/s41558-022-01516-0, number: 12 Publisher: Nature Publishing Group, 2022.
 - Wilcox, L. J., Highwood, E. J., and Dunstone, N. J.: The influence of anthropogenic aerosol on multi-decadal variations of historical global climate, Environmental Research Letters, 8, 024 033, https://doi.org/10.1088/1748-9326/8/2/024033, 2013.

Zelinka, M.: mzelinka/aprp: Sep 17, 2021 Release, https://doi.org/10.5281/zenodo.5514142, 2021.

665 Zelinka, M. D., Andrews, T., Forster, P. M., and Taylor, K. E.: Quantifying components of aerosol-cloud-radiation interactions in climate models, Journal of Geophysical Research-Atmospheres, 119, 7599–7615, https://doi.org/10.1002/2014jd021710, 2014.