



5 **The Atmospheric Oxidizing Capacity in China: Part 1. Roles of different photochemical processes**

Jianing Dai^a, Guy P. Brasseur^{a,e,f}, Mihalis Vrekoussis^{b,g,h}, Maria Kanakidou^{b,d}, Kun Qu^b, Yijuan Zhang^b, Hongliang Zhang^c, Tao Wang^f

10 ^a Environmental Modelling Group, Max Planck Institute for Meteorology, Hamburg, 20146, Germany

^b Institute of Environmental Physics (IUP), University of Bremen, Bremen, 28359, Germany

^c Department of Environmental Science and Engineering, Fudan University, 200433, China

^d Environmental Chemical Processes Laboratory, Department of Chemistry, University of Crete, Heraklion, 71003, Greece

15 ^e National Center for Atmospheric Research, Boulder, Colorado, 80307, USA

^f Department of Civil and Environmental Engineering, The Hong Kong Polytechnic University, Hong Kong, China

^g Center of Marine Environmental Sciences (MARUM), University of Bremen, Germany

^h Climate and Atmosphere Research Center (CARE-C), The Cyprus Institute, Nicosia, Cyprus

20

Correspondence to: Guy P. Brasseur (guy.brasseur@mpimet.mpg.de)

25

30

35

40



Abstract

The atmospheric oxidation capacity (*AOC*) characterizes the ability of the atmosphere to scavenge
45 air pollutants. However, it is not well understood in China, where anthropogenic emissions have
changed dramatically in the past decade. A detailed analysis of different parameters that determine
the *AOC* in China is presented on the basis of numerical simulations performed with the regional
chemical-meteorological model WRF-Chem. The model results, with the aerosol effects of extinc-
tion and heterogeneous processes taken into account, show that the presence of aerosols leads to a
50 decrease in surface ozone of approximately 8-10 ppbv in NO_x-limited rural areas and an increase
of 5-10 ppbv in VOC-limited urban areas. The ozone reduction in NO_x-sensitive regions is due to
the combined effect of nitrogen dioxide and peroxy radical uptake on particles and of the light
extinction by aerosols, which affects the photodissociation rates. The ozone increase in VOC-sen-
sitive areas is attributed to the uptake of NO₂ by aerosols, which is offset by the reduced ozone
55 formation associated with HO₂ uptake and with the aerosol extinction. Our study concludes that
more than 90% of the daytime *AOC* is due to the reaction of the hydroxyl radical with VOCs and
carbon monoxide. In urban areas, during summertime, the main contributions to daytime *AOC* are
the reactions of OH with alkene (30-50%), oxidized volatile organic compounds (OVOCs) (33-
45%), and carbon monoxide (20-45%). In rural areas, the largest contribution results from the
60 reaction of OH with alkenes (60%). Nocturnal *AOC* is dominantly attributed to the nitrate radical
(50-70%). Our results shed light on the contribution of aerosol-related NO_x loss and the high re-
activity of alkenes for photochemical pollution. With the reduction of aerosols and anthropogenic
ozone precursors, the chemistry of nitrogen and temperature-sensitive VOCs will become increas-
ingly important. More attention needs to be paid to the role of photodegradable OVOCs and noc-
65 turnal oxidants in the formation of secondary pollutants.

Keywords: O₃, Atmospheric Oxidation Capacity, Heterogeneous Reactions.

70

75

80



1. Introduction

85 With the drastic actions initiated by the Chinese authorities to improve air quality, specifically to
reduce the emissions of primary pollutants, including nitrogen oxides (NO_x), volatile organic com-
pounds (VOCs), carbon monoxide (CO), sulfur dioxide (SO_2), and the concentration of particulate
matter (PM) suspended in the atmosphere, the level of several secondary pollutants including near-
90 North China Plain (e.g., Lu et al., 2018; Liu and Wang, 2020., Wang et al., 2020). Several papers
have documented the observed trends of O_3 in China (Lu et al., 2018; Wang et al., 2022). In some
cases, studies have provided some explanation for the cause of these trends (Li et al., 2019a., Liu
and Wang et al., 2020), specifically in the most polluted areas, or have proposed some mitigation
strategies (Li et al., 2019b). Among the formulated hypotheses to explain these trends, the most
95 feasible explanation is the reduction in the level of NO_x in the polluted planetary boundary layer
(PBL) with a related reduction in the rate at which O_3 is titrated by nitric oxide (NO) in VOC-
limited areas. Another potential cause for the observed O_3 increase is the reduction in the atmos-
pheric aerosol burden and hence in the rate at which peroxy radicals (HO_2 and RO_2) that contribute
to O_3 formation are removed by heterogeneous processes (Li et al., 2019a; Liu and Wang, 2020).

100

Alleviating O_3 pollution requires a quantitative understanding of the different chemical processes
that contribute to the photochemical formation and destruction of secondary species. It also re-
quires detailed investigation of the budget of fast-reacting radicals that are directly involved in
photochemical oxidation processes. Recent observational studies have documented and analyzed
105 the evolution of several reactive species, such as OH, HO_2 and RO_2 radicals and O_3 . On the basis
of observations, generally at a single location with routine measurements lasting for several years
(Liu et al., 2022, Tan et al., 2019., Zhu et al., 2021., Wang et al., 2022). The sensitivity of ozone,
particulate matter and oxidative processes to precursor emissions has also been studied in previous
work (Liu et al., 2010; Xing et al., 2017) under highly polluted conditions. Such studies need to
110 be repeated for current conditions characterized by reduced pollutant emissions and aerosol load-
ing.

The purpose of the present study is to provide a quantitative estimate of the different factors that
affect the oxidation capacity of the atmosphere in the entire geographical area covered by China.
115 The concept of atmospheric oxidation capacity (AOC) has been introduced several decades ago
(e.g., Thompson, 1992; Prinn, 2003) to highlight the existence of self-cleansing processes in the
atmosphere. These processes allow the removal of most primary pollutants, including methane
(CH_4), non-methane hydrocarbons (NMHCs), CO, NO_x , SO_2 , as well as the formation of second-
ary species, including O_3 , particulate nitrate (NO_3^-), sulfate (SO_4^{2-}), and secondary organic aerosols
120 (SOA). The oxidation capacity is a measure of the ability of the atmosphere to destroy primary



species emitted at the Earth's surface. It is directly linked to the presence of highly reactive radicals, including OH and nitrate radical (NO_3). It is therefore influenced by processes such as photolysis generated by solar radiation, temperature, emission, scavenging processes, atmospheric transport, and other meteorological factors. It is characterized by different factors, including the atmospheric production rate of RO_x radicals (with RO_x defined as $\text{OH} + \text{HO}_2 + \text{RO}_2$, where R represents an organic chain) and the OH reactivity.

Based on model simulation, the analysis presented in this paper (Part 1) assesses the relative importance of different photochemical processes that contribute to the formation and destruction of near-surface RO_x and O_3 in different chemical environments encountered in China. A companion paper (Part 2; Dai et al., 2023) focus on the role of emission changes on the photo-oxidative species and parameters. This paper is structured as follows. Section 2 first provides some theoretical considerations on which our analysis is based. The adopted regional chemical-meteorological model, described in Sect. 3, is driven by reanalyzed meteorology for the year 2018 and by regional surface emissions that account for the contribution of different sectors. The analysis of the model simulations performed for different conditions is presented in Sect. 4-6. Specifically, the budget of oxidants (RO_x and O_3) is discussed in Sect. 4. The effect of heterogeneous chemical processes in the presence of aerosols is addressed in Sect. 5. Section 6 provides a quantitative estimate of the different indicators that describe the oxidation capacity of the atmosphere. A summary of the principal findings is provided in Sect. 7. Additional information, including the validation of the model simulations, can be found in the Supplementary Information.

2. Theoretical considerations

2.1. RO_x radicals

As shown in the pioneering paper by Levy (1971), the fate of many primary atmospheric species (CO , CH_4 , and NMHCs) and the formation of secondary species, including tropospheric O_3 , are associated with cycling chain reactions involving OH, HO_2 , and RO_2 . In the theoretical description of the key chemical processes presented here, we refer to the simplified reaction scheme listed in Table 1, but the chemical mechanism adopted in our model is considerably more detailed.

The production of RO_x radicals in the troposphere results primarily from the photolysis of O_3 (reactions R1 and R8, see Table 1), of nitrous acid (HONO) (reaction R5), and of different oxygenated volatile organic compounds (OVOCs) such as formaldehyde (HCHO) (reaction R3), larger aldehydes, acetone, etc. (reaction R4). The ozonolysis of alkenes (Alk) (reaction R11) is an additional source of RO_x , which is believed to play a relatively minor role. When considering the sources due to the photolysis of OVOCs, we single out formaldehyde due to the major contribution of this species to RO_x production. OVOCs stand, therefore, for the remaining non-HCHO OVOCs. Thus, we express the production rate of RO_x as



$$P(\text{RO}_x) = 2 k_8 [\text{O}(^1\text{D})] [\text{H}_2\text{O}] + J_{\text{HONO}} [\text{HONO}] + J_{\text{HCHO}} [\text{HCHO}] + \sum_i r_i J_i [\text{OVOC}_i] + \sum_i s_i k_{11,i} [\text{Alk}_i] [\text{O}_3],$$

(1)

165

Here and in other expressions below, factors k_i represent the reaction coefficients for the reaction i in Table 1, and J is the photolysis frequency for the chemical species under consideration. The brackets stand for the number densities of species generally expressed in molecules or radicals per cm^3 . Coefficient r_i represents the number of RO_x produced by the photolysis of each OVOCs species and s_i is the number of RO_x produced by each alkene ozonolysis reaction. These coefficients are specific to a reaction involving the photolysis of OVOCs and the ozonolysis of the alkenes. A more explicit form for the two last terms in the above expression depends on the adopted chemical mechanism. The RO_x source term provides an estimate for the availability of radicals that initialize photooxidation processes in the troposphere.

170

175

The destruction of RO_x radicals results from the termination reactions R15-18 between different RO_x radicals (L_H), reactions R22-24 between RO_x radicals and nitric oxide (L_N) and the heterogeneous uptake of HO_2 (reaction R28) on aerosol surfaces (L_{het}). Thus, the total destruction rate of RO_x can be expressed as

180

$$D(\text{RO}_x) = L_H + L_N + L_{het},$$

(2)

with

185

$$L_H = \{ 2 k_{15} [\text{OH}] + 2 k_{16} [\text{HO}_2] + 2 \sum_i k_{17} [\text{RO}_{2,i}] \} [\text{HO}_2] + 2 \sum_{i,j} k_{18,i,j} [\text{RO}_{2,i}] [\text{RO}_{2,j}]$$

$$L_N = \sum_i k_{22,i} [\text{RO}_{2,i}] [\text{NO}] + k_{23} [\text{OH}] [\text{NO}_2] + k_{24} [\text{OH}] [\text{NO}]$$

190

$$L_{het} = k_{28} [\text{HO}_2]$$

In the above expressions, we assume that, near the surface, the RO_x termination reactions between HO_2 and NO_2 (that produce nitrous acid, HO_2NO_2) and between acetyl peroxy radicals and NO_2 (that produce peroxyacetyl nitrate, PAN) are balanced by the regeneration of RO_x resulting from the photolysis of HO_2NO_2 and the thermal decomposition of PAN (equilibrium conditions), respectively. Therefore, the related reaction rates do not appear explicitly in the above expressions.

195

2.2. Odd Oxygen

200



In the troposphere, odd oxygen ($O_x = O_3 + NO_2$) is produced through complex recurrent radical reaction chains involving the oxidation of hydrocarbons in the presence of NO_x . The O_3 molecule is formed by the rapid photolysis of NO_2 (reaction R2) followed by the recombination of atomic oxygen (reaction R7). Reaction R2 is balanced by reactions R19 and R20, which reproduce NO_2 .

205 A production of odd oxygen occurs only if NO is converted to NO_2 without consuming O_3 , i.e., by reactions R20 and R21 with a peroxy radical (HO_2 , CH_3O_2 , and other higher-order organic radicals) provided by the oxidation of methane, other hydrocarbons (HC), and by carbon monoxide (reactions R9, R10 and R12). The resulting production rate of odd oxygen can be expressed with a good approximation by

210

$$P(O_x) = k_{20} [HO_2] [NO] + \sum_i k_{21,i} [RO_2]_i [NO],$$

(3)

215 This equation highlights the nonlinear nature of $P(O_x)$ since the concentration of nitric oxide and peroxy radicals are dependent on each other.

The photochemical destruction of O_x results from several processes, including the photolysis of O_3 (reaction R1) followed by the reaction R8 between the electronically excited oxygen atom $O(^1D)$ and water vapor (H_2O). Other O_x loss mechanisms involve the reactions of ozone with OH (reaction R13), HO_2 (reaction R14), and different alkenes (Alk) (R11). In the presence of NO_x , an additional loss mechanism is provided by the titration of O_3 by NO (R19), followed by the conversion of NO_2 to nitric acid (HNO_3 ; R23). The total destruction rate of O_x is therefore expressed as

225

$$D(O_x) = k_8 [O^1D][H_2O] + \{\sum_i k_{11} [Alk_i] + k_{13} [OH] + k_{14} [HO_2] + k_{19} [NO]\} [O_3] + k_{23} [OH][NO_2],$$

(4)

230 The dominant pathways leading to the formation and destruction rates of O_x and hydroxyl radicals vary according to chemical environments. Under relatively clean conditions with low levels of NO_x , the production of odd oxygen, as provided by reactions R20 and R21, is limited by the availability of NO_x , while the loss of the RO_x radicals is dominated by the peroxy-radical self-reaction R16 that leads to the formation of hydrogen peroxide (H_2O_2) (Song et al., 2021). Under these conditions, a reduction in the emissions of NO_x tends to reduce the ground-level O_3 concentration.

235 In polluted areas, including many urban centers, the level of NO_x is so high that saturation conditions prevail. In this environment, the formation of O_3 is determined by the availability of VOCs, and the loss of RO_x is dominated by reaction R23 between NO_2 and OH, which produces HNO_3 . In this case, a reduction in NO_x tends to increase the concentration of O_3 , while a reduction in VOCs is expected to reduce its near-surface abundance (Wang et al., 2022). Furthermore, since
240 highly polluted environments are generally characterized by elevated aerosol loads, the effect of



heterogeneous processes on the abundance of reactive species becomes particularly important. Among these reactions, we consider more specifically the uptake of HO₂, NO₂, NO₃, and N₂O₅ on the atmospheric aerosol surfaces (reactions R28-31). The heterogeneous destruction of peroxy radicals on the surface of aerosols tends to inhibit the formation of O₃ by reducing the rate of reactions
245 R20 and R21. Under high levels of aerosols, this reaction may become important for O₃ formation, which has led Ivatt et al. (2022) to define a third O₃ sensitivity regime called the aerosol-inhibited photochemical O₃ regime. The uptake of NO₂ leads to the formation of HONO, whose photolysis represents a significant source of OH. It also forms NO, which reacts with HO₂ and RO₂ to produce O₃. In short, heterogeneous processes may either favor or inhibit the formation of odd oxygen in
250 polluted areas.

The lifetime of O_x in the PBL is sufficiently long (one or two days) that additional processes besides photochemical production and destruction need to be taken into consideration. Among them is the additional loss of O₃ and NO₂ resulting from dry deposition on the vegetation. With a deposition
255 velocity of about 1 cm s⁻¹ (Wesely et al., 2000), the corresponding odd oxygen loss rate in the boundary layer is close to 1 ppbv h⁻¹, if one assumes that the depth of the mixing layer is in the order of 1 km. In addition, vertical mixing in the convective PBL and advective horizontal transport tend to disperse locally produced O₃ and balance its net production, for example, in urban centers.

260

2.3. Formulation of aerosol uptake

As stated above, heterogeneous chemical reactions can substantially influence the concentrations of RO_x radicals and O₃. The reactions under consideration in our analysis are reactions R28 to
265 R31, listed in Table 1. The first-order reaction rate constant on aerosols k_a [s⁻¹] for species i associated with these reactions is expressed by (Schwartz (1986)):

$$k_{a,i} = A_a \left[\frac{a}{D} + \frac{4}{\gamma_{a,i} v_i} \right]^{-1},$$

(5)

270

where A_a [cm² cm⁻³] is the aerosol surface area density, a [cm] is the mean radius of the particles, D [0.247 cm² s⁻¹] is the gas-phase diffusion coefficient (Mozurkewich et al., 1987), γ_a is the dimensionless reaction-dependent uptake coefficient for species i , and v_i [cm s⁻¹] is the mean thermal velocity of species i given as a function of temperature T [K] and molecular mass m_i by

275

$$v_i = \left[\frac{8k_b T}{\pi m_i} \right]^{\frac{1}{2}},$$

(6)

with k_b (1.38×10^{-23} J K⁻¹) being the Boltzmann constant.



280

The chemical substances produced as a result of the HO₂ uptake onto aerosol surfaces are not clearly established and could be H₂O or H₂O₂ (Song et al., 2021). Here, to determine the maximum effect of this reaction, we assume that the HO₂ uptake onto aerosols (reaction R28) represents a terminal reaction of the hydrogen radical chain. Hence, water molecules rather than peroxide molecules are assumed to be formed. The corresponding uptake coefficient (γ_{HO_2}) is chosen to be 0.1 in this study, a value lower by a factor of 2 than used in some earlier model studies ($\gamma_{\text{HO}_2} = 0.2$; Tie et al., 2001, 2005; Martin et al., 2003; Liu and Wang, 2020; Ivatt et al. 2022), but consistent with the conclusions reached by Gaubert et al. (2020) from their model simulations. These lower values are also adopted by Yang et al. (2022) and are consistent with the measurements of Lakey et al. (2015), Tan et al. (2020), and Song et al. (2020). Specifically, on the basis of observations made in the Beijing-Tianjin-Hebei area during the summer of 2014, Song et al. (2020) conclude that the best fit for the value of γ_{HO_2} is a value of 0.116 ± 0.086 , which is close to the value adopted in the present study.

285

290

295

The heterogeneous uptake of N₂O₅ by aerosol particles leads to the formation of nitric acid molecules (reaction R29). In the present study, we neglect the possible formation of ClNO₂ followed by its photolysis into Cl and NO₂. This process associated with the presence of chloride ions in the bulk of the particles is a source of additional radicals and hence could have an influence on O₃ (Thornton et al., 2010; Dai et al., 2020). Here, for the uptake of N₂O₅, we adopt the first-order rate constant as expressed by Bertram et al. (2009) and modified by Yu et al. (2020) with the surface concentrations of the particle chloride and nitrate ions taken from the MOSAIC estimates. This parameterization has been used in simulating the concentration of N₂O₅ in several Chinese sites (Yu et al., 2020), and the simulated levels are in good agreement with the observed N₂O₅ values (Dai et al., 2020).

300

305

The rate of heterogeneous conversion of NO₃ (reaction R30) is calculated by Eq. (5) with a value of the uptake coefficient equal to 10⁻³ (Jacob, 2000; Xue et al., 2014; Liu and Wang et al., 2020).

310

Finally, the heterogeneous uptake of NO₂ on aerosol surfaces leads to the production of HONO and HNO₃ (reaction R31), and, as HONO is rapidly photolyzed after sunrise, this heterogeneous process represents a source of OH radicals. The process also converts NO₂ into NO. Here, according to Zhang et al. (2021), we express the first order rate constant by Eq. (5) with a value of the uptake coefficient equal to 8×10⁻⁶ during nighttime and to 1×10⁻³ × (J/J_{max}) during daytime (Li et al., 2010; Czader et al., 2012; Fu et al., 2019), with J representing the light intensity [W m^{-2}] and J_{max} [W m^{-2}] the peak value of light intensity (chosen to be 400 W m^{-2} in this study).

315

2.4. Other HONO sources



320 For the particular heterogeneous reaction involving NO_2 , which leads to the formation of nitrous
acid, we also consider the additional contribution of the uptake on flat surfaces, specifically on
bare soils, including asphalt in urban areas. This effect is believed to play a significant role partic-
ularly in urban areas (Zhang et al., 2016; Li et al., 2018). Zhang et al. (2021) claim that the meas-
ured vertical nighttime profile of this species suggests that the dominant formation nighttime
325 process, which is only crudely represented here, we assume that the first-order rate constant k_g [s^{-1}]
for this process is given by Liu et al. (2019) for nighttime conditions

$$k_g = \frac{1}{8} \gamma_g v_{\text{NO}_2} A_g, \quad (7)$$

330 where A_g is the surface area density over the bare soil and urban surfaces; γ_g is the uptake coeffi-
cient on the ground. Here, according to Zhang et al. (2021), we express a value of the uptake
coefficient equal to 4×10^{-6} during nighttime and to $6 \times 10^{-5} \times (J/J_{\text{max}})$ during daytime, where J and
 J_{max} [W m^{-2}] represent the solar intensity and its maximum value. Following the suggestion of
335 Vogel et al. (2003) adopted, for example, by Zhang et al. (2021), we express the surface density
over the ground by $1.7/h$, where h [m] represents the height of the model layer adjacent to the
ground; the 1.7 value represents an effective factor per ground surface area in the first layer.

In this model case, we also account for the gas-phase reactions of HONO (R24-R27) as well as the
340 direct transportation HONO emissions. The latter are assumed to be equal to 0.8% of the traffic
emission of NO (Dai et al., 2021). In this study, we neglect the direct HONO emissions from soil
and the daytime HONO source from the photolysis of NO_3^- , which may lead to an underestimation
of HONO concentration in rural areas and during daytime (Zhang et al., 2016, Fu et al., 2021;
Zhang et al., 2021).

345

2.5. Photochemical reactivity and *AOC*

To characterize the oxidation capacity of the atmosphere in China, we consider several indicators
that have proven to be useful for developing O_3 -controlling strategies. These include the OH reac-
350 tivity associated with the action of volatile organic compounds (VOC^R) and nitrogen oxides
(NO_x^R), the radical chain length (*ChL*), the ozone production efficiency (*OPE*), and the atmos-
pheric oxidation capacity (*AOC*).

Since NO_x , VOCs, and CO are oxidized by the OH radical as part of a cyclic chain process that
355 initiates the O_3 formation, an estimate of the OH reactivity (expressed in s^{-1}) allows us to under-
stand the factors that determine the photochemical budget of O_3 and more generally the factors
that characterize the atmosphere's oxidizing capacity. The OH reactivity by the different VOCs
and CO and by NO_x is defined as



360 $VOC^R = \sum_i k_{10,i} [VOC_i] + k_{12} [CO],$
 (8a)

$NO_x^R = k_{23} [NO_2],$
 (8b)

365

The radical chain length ChL provides a measure of the number of cycles affecting RO_x radicals before these radicals undergo a termination process. It can therefore be expressed by the ratio between the conversion rate between RO_x radicals, including the conversion by NO of HO_2 to OH (reaction R20) and of RO_2 to HO_2 (reaction R21), and the destruction rate of RO_x (or equivalently by the production rate of RO_x). Thus, adopting here the definition of Martinez et al. (2003), Mao et al. (2010), and Zhu et al. (2020), we write

370

$$ChL = \frac{k_{20}[HO_2][NO] + \sum_i k_{21,i}[RO_{2,i}][NO]}{D(RO_x)},$$

(9)

375

From this adopted definition Eq. (9), and assuming that the RO_x production and destruction rates are in balance, we can write

380 $P(O_x) \cong P(RO_x)ChL,$
 (10)

which shows that O_x production is proportional to the RO_x production rate and is favored by a large number of radical regenerations.

385 The Ozone Production Efficiency (OPE) is used to quantify the efficiency of O_3 molecules formed per NO_x molecule oxidized. It is defined as the ratio between the O_3 production rate and NO_x loss rate

390 $OPE = \frac{P(O_3)}{D(NO_x)} \cong \frac{P(O_3)}{P(HNO_3)},$
 (11)

and represents the efficiency of NO_x . To a good approximation, this expression can be expressed as

395 $OPE \approx \frac{k_{20}[HO_2][NO]}{k_{23}[OH][NO_2]},$
 (12)



As the instantaneous value of OPE depends on the HO_2/OH and NO/NO_2 concentration ratios, it accounts for the couplings between RO_x and NO_x cycles. One can show that this factor is usually
400 highest in the remote atmosphere or low- NO_x environments (Ridley, 1999). Note that, under VOC-limited conditions (polluted areas) when the production rate of odd oxygen can be expressed as (Kleinman et al., 2002)

$$P(O_x) = \sum_i \varphi_i [VOC_i][OH],$$

405 (13)

where φ_i represents the O_3 yield from the production of HO_2 radicals, OPE can be approximated by

$$OPE \approx \bar{\varphi} \frac{VOC^R}{NO_x^R},$$

410 (14)

where $\bar{\varphi}$ represents an average yield value. In other words, under VOC-limited situations, the ratio between VOC^R and NO_x^R has some similarities with the odd oxygen production efficiency.

415 Finally, the atmospheric oxidizing capacity (AOC ; expressed in $cm^{-3} s^{-1}$), a parameter introduced by Geyer et al. (2001) to account for the contribution of all oxidants, is derived here as the rate at which CO , CH_4 , and NMHCs (all species are noted here as Y_i) are oxidized by OH , O_3 , and NO_3 (noted as X_j) (Geyer et al., 2001; Elshorbany et al., 2009; Xue et al., 2016; Wang et al., 2022; Yang et al., 2022). Thus, when considering all combinations of the different primary pollutants and atmospheric oxidants, we write

$$AOC = \sum_i^j k_{i,j} [Y_i][X_j],$$

420 (15)

425 As stated by Wang et al. (2022) and Yang et al. (2022), AOC is a parameter well-suited to describe the removal rate of primary pollutants and the formation of secondary species including O_3 and secondary $PM_{2.5}$. It is, therefore, an indicator used to design control policies for these secondary species. During daytime, the largest contribution to AOC is due to the oxidation of pollutants by the OH radical (Li et al., 2017; Liu et al., 2022). At night, the oxidizing capacity is due to the
430 oxidation by NO_3 and O_3 .

3. Model description and validation

435 3.1. Modeling setting



To characterize the chemical budget of reactive species, photochemical parameters and *AOC* in China, we use the version 4.1.2 of the WRF-Chem model (Skamarock et al., 2019) to simulate the meteorological fields as well as the regional transport, the chemical and physical transformations of trace gasses and aerosols. We adopt the MOZART-4 gas-phase chemical mechanism documented and evaluated by Emmons et al. (2010), which includes 108 chemical species and 235 gas-phase reactions. This scheme is coupled to the MOSAIC aerosol module described by Fast et al. (2006), Zaveri et al. (2008), and Lu et al. (2021). A list of detailed RO₂, VOCs and aerosol species included in the model is provided in Table S1 in the Supplementary information.

We select one month in the winter (1st to 31st January) and the summer (1st to 31st July) of 2018 respectively to analyze the calculated distributions of chemical species. The horizontal resolution adopted in the present study is 36 km × 36 km over the entire domain that covers East and Southeast Asia (from 15° S to 60° N in latitude and 60° E to 150° E in longitude). Initial meteorological conditions are taken from the NCEP reanalysis dataset FNL (<http://rda.ucar.edu/datasets/ds083.2/>). Initially chemical boundary conditions are constrained by the results of the global CAM-chem model (<https://www.acom.ucar.edu/cam-chem/>). The different modules used to represent physical processes are provided in Table S2.

For the anthropogenic emissions of air pollutants, we adopt the surface emissions provided by the Multi-resolution Emission Inventory for China (MEIC v1.3; <http://www.meicmodel.org/>) derived for the year 2017 (Zhang et al., 2009; Zheng et al., 2018). This inventory covers the anthropogenic emissions for the geographical area of mainland China. For the remaining areas of Asia, we use the anthropogenic emissions provided by the 2018 global inventory of the Copernicus Atmosphere Monitoring Service (CAMS)-GLOB-ANT_v4.2 (Elguindi et al., 2020; Granier et al., 2019). Biogenic emissions are calculated online by the Model of Emission of Gas and Aerosols from Nature (MEGAN) version 2.1 (Guenther et al., 2006). The dust and sea-salt emissions are calculated online by the Global Ozone Chemistry Aerosol Radiation and Transport (GOCART) module (Chin et al., 2002).

The availability of several observational datasets allows us to evaluate the meteorological parameters and air pollutant concentrations derived by our regional model. The meteorological data used to validate the model simulations, including the wind direction, wind speed, surface temperatures, and specific humidity, are obtained from the NOAA National Climatic Data Center (NCDC). Conventional air pollutant data, including SO₂, NO₂, CO, O₃, and PM_{2.5}, are obtained from the surface stations of China's Ministry of Ecology and Environment (MEE; <https://www.mee.gov.cn/>). To validate the model results, we calculate the mean bias, the normalized mean bias, the normalized mean error, the root mean square errors, and the correlation coefficient. The equations for these statistical parameters are found in the paper by Dai et al. (2020).



In our analysis presented in the subsequent sections, we examine in more detail the calculated concentrations of photochemical parameters at urban sites in four big cities in China: Beijing, Shanghai, Guangzhou, and Chengdu (Fig. 1). We also provide these parameters at four rural observational sites for which detailed observational analysis is available. These include the relatively
480 polluted site of Wangdu in the suburban region near Beijing, the Atmospheric Supersite of Heshan located 50 km to the southwest of Guangzhou, the remote free site of Waliguan at about 3800 m altitude, and the coastal site of Hok Tsui in Hong Kong. Detailed information on the selected sites in the present study is listed in Table 2.

485 3.2. Design of numerical experiments

Table 3 lists the different sensitivity cases designed for this study. The baseline case, called *Het-all*, accounts for all heterogeneous reactions referred to in Table 1 and includes all identified sources of HONO mentioned in Sect. 2.4. The *Het-all* case is used to evaluate the performance of
490 the model relative to observations. To quantify the specific aerosol effects through the uptake of HO₂, N₂O₅, NO₃ and NO₂ and extinction, and NO₂ uptake over the ground on surface O₃ concentration, other nine sensitivity cases are considered based on different assumptions. The details on these sensitivity cases are given as follows.

495 For the specific effects of aerosol uptake, the respective importance of these processes is determined by subtracting the baseline results from the results in sensitivity cases in which specific heterogeneous reactions are ignored: HO₂ (*No-HetHO2-Aero*), NO₃ (*No-HetNO3-Aero*), N₂O₅ (*No-HetN2O5-Aero*), NO₂ (*No-HetNO2-Aero*). The case labeled *No-Het-Aero* ignores all the above heterogeneous reactions on aerosols. The difference between *Het-All* and *No-Het-Aero* represents the combined effects of these heterogeneous reactions on particles. The case denoted as
500 *No-Phot* ignores the radiative effects of aerosols on the calculation of the photodissociation coefficients. The difference between *Het-All* and *No-Phot* represents the effect of aerosol radiation. An additional case labeled as *No-Het-Aero-Phot* ignores the above-mentioned heterogeneous reactions on aerosols and aerosol effects on light extinction and photodissociation to quantify the combined effect of aerosol uptake and radiation.
505

In order to quantify the contribution of HONO sources added to the model, we consider a case labeled *No-HONO* in which the heterogeneous uptake of NO₂ by aerosols, bare soils, urban surfaces as well as the homogeneous formation, and surface emissions of HONO are all ignored. The
510 difference between the results of the *Het-All* case and the *No-HONO* case represents the effect of all HONO sources. The final case, denoted as *No-Het-HONO-Phot*, ignores all the above heterogeneous reactions on aerosols, other HONO sources, and the radiative effects of aerosols on the calculation of the photodissociation coefficients. The difference between *Het-All* and *No-Het-HONO-Phot* provides quantitative measures of the effects resulting from all heterogeneous reactions (aerosols and ground effects), other HONO sources, and aerosol radiation.
515



3.3. Model validation

In Fig. S1, we show the spatial validation of the calculated surface concentrations of the Maximum
520 Daily 8-hour average (MDA8) O₃, as well as the monthly averages NO₂, CO, and PM_{2.5} (*Het-all*
case) with available observational data from MEE for January and July 2018. In most cases, this
comparison shows a good performance of the model with, however, some discrepancies: an over-
estimation of summertime O₃ in central and western China associated with an underestimation of
NO₂ in these regions, an underestimation of summertime O₃ in eastern China with a slight overes-
525 timation of NO₂ (Fig. S1). In the case of CO and PM_{2.5}, the calculated concentrations are higher
than the measured values in central China in both seasons. The diurnal variation of NO₂, O₃, CO,
and PM_{2.5} in January and July for the four metropolitan areas selected in our study are compared
with measurements from monitoring stations in Fig. S2 and S3. The model successfully simulates
the diurnal variations of these chemicals. However, an overestimation of summertime NO₂ is pre-
530 sented in these urban areas, with an underestimation of O₃ at night and an overestimation during
daytime. These discrepancies can be explained by the relatively lower NO₂ uptake coefficients
used in our studies (Liu et al., 2019; Fu et al., 2019). The simulated CO is slightly overestimated,
which can be attributed to uncertainties in chemical boundary conditions and emissions (Liu and
Wang et al., 2020). An overestimation of PM_{2.5} is found in summer, which can be partially due to
535 uncertainties in emissions and the mechanisms of secondary aerosol formation (Li et al., 2022).

Base estimates of the NO, HONO, HCHO, OH, HO₂, NO₃, isoprene, ethane, and ethene mixing
ratios are found in Fig. S4-S6. Calculated diurnal variations of surface NO, HONO, OH, HO₂, and
NO₃ are provided in Fig. S8-S11. Generally, our simulated concentrations of OH, HO₂, HONO,
540 and HCHO matched relatively well with the observational data. The calculated aerosol surface
area density is shown in Fig. S12. The values calculated in eastern China are considerably higher
during wintertime (2.5 to 3×10^{-5} cm² cm⁻³) than during the summer (0.7 to 1.0×10^{-5} cm² cm⁻³).
A more detailed discussion and validation of the base model results are provided in the Supple-
mentary Information.

545

4. The budget of oxidants

In order to highlight the regional differences in the existing photochemical regimes, we first show
the distributions of areas where the O₃ formation is either NO_x- or VOC-limited. Two indicators
550 define these areas: the concentration between H₂O₂ and HNO₃ and between HCHO and NO₂. An
area is considered as NO_x-limited if $[H_2O_2]/[HNO_3] > 0.2$ (Zhang et al., 2009) or $[HCHO]/[NO_2] >$
 1.0 (Jing et al., 2021). It is assumed to be VOC-limited if $[H_2O_2]/[HNO_3] < 0.06$ (Zhang et al.,
2009) or $[HCHO]/[NO_2] < 0.55$ (Jing et al., 2021). Regions in between these bounds correspond
to an intermediate situation. Abdi-Oskouei et al. (2022) also define NO_x-sensitive region by the
555 criterion $L_H/L_N < 0.35$.



560 According to the definition based on the HCHO to NO₂ concentration ratios, we see in Fig. 2 that, during summertime, the ozone formation in southern and western China as well as in the surrounding oceanic regions, is primarily NO_x-sensitive. Exceptions are found in the south-eastern coast with the metropolitan regions of Guangzhou and Hong Kong. The formation of ozone in a large fraction of northern China, specifically in the areas to the south of Beijing and in the region near Shanghai, is VOC-sensitive. A broader area surrounding Beijing and Shanghai is in an intermediate situation. This latter condition prevails in urban hotspots such as Seoul and Tokyo. During winter, the ozone formation in most of eastern and northern China is VOC-limited. We note that
565 the geographical area of the VOC-limited regime is larger when adopting the [HCHO]/[NO₂] criteria than when adopting the L_H/L_N ratio, but smaller than when adopting the [H₂O₂]/[HNO₃] concentration ratio (see Fig. S13).

4.1. The budget of RO_x

570

Figure 3 shows the geographical distribution of the average daytime (08:00-19:00 Local Standard Time) production rate of RO_x ($P(\text{RO}_x)$) contributed by the photolysis of HONO, HCHO, non-HCHO OVOCs and O₃ for the January and July of 2018.

575 In winter, the mean daytime production rate of the radicals is small in less populated western China. In the eastern plain, its value associated with the HONO photolysis is typically 0.5-0.8 ppbv h⁻¹ in rural areas and reaches 1-2 ppbv h⁻¹ in polluted urban areas. The contribution of the HCHO photolysis to $P(\text{RO}_x)$ is of the order of 0.15 ppbv h⁻¹ in most areas, with values as high as 0.5 ppbv h⁻¹ in Guangzhou, which benefits from sufficient solar radiation during winter. Correspondingly,
580 the contribution of other OVOCs is around 0.2 ppbv h⁻¹ in southern China, with a similar distribution of the photolysis of HCHO. The mean daytime $P(\text{RO}_x)$ associated with O₃ photolysis is small (less than 0.5 ppbv h⁻¹) over the entire Chinese territory except in the very south of the country, where it reaches 0.2 ppbv h⁻¹. The contribution by alkene ozonolysis is negligible (Fig. S14).

585 In summer, the mean $P(\text{RO}_x)$ by HONO photolysis reaches 2 to 3 ppbv h⁻¹ in the regions surrounding Beijing, Shanghai, Guangzhou, and Chengdu but is considerably smaller (less than 0.5 ppbv h⁻¹) in the outskirts rural areas. The photolysis of HCHO reaches values ranging from 0.5 to 1.0 ppbv h⁻¹ in the rural areas of eastern China, with no particular maximum in metropolitan areas. The high value of $P(\text{RO}_x)$ contributed by non-HCHO OVOCs photolysis, ranging from 0.3 to 0.8
590 ppbv h⁻¹, with obvious peak values in city areas. The daytime averaged value of $P(\text{RO}_x)$ resulting from the photolysis of O₃ is of the order of 1.0 ppbv h⁻¹ in eastern and southern China. The peak spot in the Sichuan basin is due to the high water vapor contributed by heavy rainfall in summer, which leads to high OH radical (Xia et al., 2021).



595 In summary, and for the present conditions, our model suggests a higher value of $P(\text{RO}_x)$ in sum-
mer than in winter. The higher summertime $P(\text{RO}_x)$ in eastern and southern China is associated
with the photolysis of formaldehyde and O_3 . In urban areas, the dominant contribution to the higher
 $P(\text{RO}_x)$ in summer is provided by the photolysis of HONO. The spatial distributions that vary in
winter and summer are also related to the seasonal variations in meteorological parameters, such
600 as surface temperature and water vapor (Fig. S15).

The diurnal variation of the $P(\text{RO}_x)$ in four different metropolitan areas (Beijing, Shanghai, Guang-
zhou, and Chengdu) at two relatively polluted rural (Wangdu and Heshan) and two clean remote
sites (Hok Tsui and Waliguan) is shown for summertime in Fig. 4. The graph shows the contribu-
605 tion of the HONO (green area), HCHO (red area), non-HCHO OVOCs (dark green area), and O_3
(yellow green area) photolysis, as well as the effect of alkene ozonolysis (blue area). In the four
urban centers, the maximum values of $P(\text{RO}_x)$ in the early afternoon range between 5 ppbv h^{-1} in
Shanghai and $6\text{-}8 \text{ ppbv h}^{-1}$ in the three other cities. In the early morning, as the sun rises, the largest
contribution is due to the photolysis of HONO. A few hours later, the contribution of the photolysis
610 of HCHO and other OVOCs becomes large. The value of $P(\text{RO}_x)$ from the photolysis of O_3 is
small in the early morning and peaks in the mid-afternoon. In the four sites of the rural areas, the
maximum value of $P(\text{RO}_x)$ is close to 5 ppbv h^{-1} in Wangdu (suburban site southwest of Beijing),
 1.3 ppbv h^{-1} in Heshan (rural site close to Guangzhou), and less than 0.3 ppbv h^{-1} in Hok Tsui
(coastal site) and Waliguan (western China). The contribution of O_3 photolysis generally domi-
615 nates at these rural sites.

A comparison between the values of $P(\text{RO}_x)$ derived from local observations is performed with
model estimates in Table 4. At four city sites, our simulated values of the maximum of $P(\text{RO}_x)$
and the contribution by the photolysis of HONO, HCHO and O_3 reproduce satisfactorily the ob-
620 servations. However, at the Heshan site, the calculated value of $P(\text{RO}_x)$ (1.1 ppbv h^{-1}) in our study
is much smaller than the observed value (4.0 ppbv h^{-1}) (Tan et al., 2019). One reason for this
discrepancy may be the missing soil HONO emission, which leads to an underestimation of HONO
concentration at this site (Table S4) and the contributed value to $P(\text{RO}_x)$ by the photolysis of
HONO (0.5 ppbv h^{-1} v.s. 2.0 ppbv h^{-1}). Another uncertainty is the contribution of non-HCHO
625 OVOCs photolysis to the $P(\text{RO}_x)$. Wang W. et al. (2022), based on the measurement data in
Guangzhou, reported that a model without constraints of non-HCHO OVOCs would lead to an
underestimation in the production rate of RO_x and O_3 . Owing to the lack of specific OVOCs meas-
urements, it is difficult to quantify the contributions of different OVOC species. Additional efforts
regarding the OVOC measurements are needed to understand the specific contribution of OVOCs
630 to the atmospheric oxidation capacity.

The diurnal variations of $P(\text{RO}_x)$ are displayed in Fig. S16. The maximum value of the $P(\text{RO}_x)$ is
close to 3 ppbv h^{-1} in Beijing, Chengdu and Shanghai and is about twice as large in the region of
Guangzhou. In Wangdu and Heshan, the peak values are comparable to those in urban areas, while



635 in Hok Tsui and Waligan, they are lower than 1 and 0.1 ppbv h⁻¹, respectively. In most cases, the major contribution to $P(\text{RO}_x)$ is provided by the photolysis of HONO.

The spatial distribution of the daytime averaged destruction rate for RO_x ($D(\text{RO}_x)$) in January and July is shown in Fig. 5. In the two seasons under consideration, the loss rate is the largest in the eastern and southern regions of China. In January, this quantity is largest in polluted metropolitan areas, with daytime mean values surpassing 2 ppbv h⁻¹. In July, the total daytime average value of $D(\text{RO}_x)$ is of the order of 3 ppbv h⁻¹ in the rural areas of eastern China and reaches about 6 ppbv h⁻¹ in the urban and industrialized areas. In Tibet, the value of this quantity is small (~0.1 ppbv h⁻¹ in winter and 1.2 ppbv h⁻¹ in summer).

645 Interestingly, the relative importance of the different photochemical mechanisms involved in the $D(\text{RO}_x)$ varies considerably with the season. In January, the $D(\text{RO}_x)$ due to the reaction between OH radicals and NO_x (L_N in Eq. 2) dominates in most parts of eastern China (contribution of typically 90%) except in Tibet, where the largest loss (contribution of 70-80%) is due to the recombination of hydrogen radicals (L_H in Eq. 2). In July, it is this last type of loss (L_H) that plays the dominant role (typically 90%), except in the eastern plain of China where the level of NO_x is highest, and L_N (contribution of 70-80%) is, therefore, larger than L_H (contribution of 20%). In both seasons, the destruction of RO_x by the uptake of HO_2 is relatively small (generally less than 15%). The highest contribution occurs during winter in southwestern China (contribution of 30%) and in the Ganges River Valley of India (contribution of 40%).

The diurnal variation of $D(\text{RO}_x)$ in July is presented in Fig. 6 in four urban areas (Beijing, Shanghai, Guangzhou, and Chengdu) and in four selected locations in rural areas (Wangdu, Heshan, Hoktsui, and Waliguan, see Fig. 1). In the summertime, the value of $D(\text{RO}_x)$ around noon reaches about 13 ppbv h⁻¹ in the urban areas except in Shanghai, where it reaches only 8 ppbv h⁻¹. In winter, the corresponding maximum values are closer to 3-4 ppbv h⁻¹ except in the southern city of Guangzhou, where the maximum loss rate is closer to 6 ppbv h⁻¹. In all these cases, the dominant contribution to $D(\text{RO}_x)$ is attributed to the reactions involving the presence of NO_x (L_N). In rural areas, the value of $D(\text{RO}_x)$ is considerably smaller. In July, it is of the order or smaller than 2 ppbv h⁻¹ and is dominated by the HO_x recombination (L_H). In the wintertime, the peak loss is smaller than 0.2 ppbv h⁻¹ except in Chengdu, where it reaches 0.4 ppbv h⁻¹. The major contribution is due to the reactions involving NO_x (L_N).

670 The experimental study of Whalley et al. (2021) in Beijing during the summer of 2018 provides for $D(\text{RO}_x)$ a maximum daytime value of 7 ppbv h⁻¹ with the following contributions: 4 ppbv h⁻¹, 1.3 ppbv h⁻¹, and 1 ppbv h⁻¹ for the $\text{NO}_2 + \text{OH}$, $\text{NO} + \text{OH}$, and $\text{RO}_2 + \text{OH}$ reactions, respectively. Our simulated value matches well the reported experimental data, and the corresponding values in July are 5 ppbv h⁻¹, 0.8 ppbv h⁻¹, and 1.2 ppbv h⁻¹, respectively. Yang et al. (2021) report diurnal variations in $D(\text{RO}_x)$ as derived from their observation in Chengdu during the autumn of 2018.



675 The peak value of this quantity is about 7 ppbv, which is lower than our calculated value of 12.5
ppbv h⁻¹ in July. The higher level in our study is due to the overestimated concentration of sum-
mertime NO₂ in Chengdu (Fig. S2). This overestimation also leads to a higher contribution of the
NO₂ + OH reaction by 55% in our study than the reported 35% in the literature.

680 4.2. The budget of odd oxygen

The production rate of odd oxygen ($P(O_x)$) with its two contributions (reaction of NO with hydro-
genated and organic peroxy radicals (HO₂ and RO₂) shown in Eq. 3) is shown in Fig. 7 for January
and July 2018. In the Northern China Plain and other urbanized areas, the production rate is of the
685 order of 4–6 ppbv h⁻¹ during winter (January), while in the rural areas of southern China, it is larger
than 20 ppbv h⁻¹ during summer (July) and 6–10 ppbv h⁻¹ during winter. The value of $P(O_x)$ is very
small in the western part of China. The relative contributions of both step-limiting processes to the
total $P(O_x)$ are of the same order of magnitude, although the reaction involving the hydrogenated
peroxy radicals seems to slightly dominate, particularly outside densely populated areas.

690 The diurnal variations in $P(O_x)$ are depicted in Fig. 8 for specific areas of China in July. This graph
highlights the maximum values found during the early afternoon in urban areas: 115 ppbv h⁻¹ in
Beijing, 40 ppbv h⁻¹ in Shanghai, 110 ppbv h⁻¹ in Guangzhou, and 70 ppbv h⁻¹ in Chengdu. In
Beijing, Whalley et al. (2021) derived from their observations in the summer of 2018 a maximum
695 O_x production rate of 100 ppbv h⁻¹. These high values must be contrasted by the considerably lower
values found in rural areas: 1.5 ppbv h⁻¹ at Mount Walinguan and Hok Tsui. Intermediate maxi-
mum values are found at the sites located in the vicinity of large metropolitan areas: 40 ppbv h⁻¹
in Wangdu and only 7 ppbv h⁻¹ in Heshan. The graph also shows the relative contribution of the
hydrogen and organic peroxy radicals. Both radicals contribute about equally to the odd oxygen
700 production rate. The contribution of the organic peroxy radical is determined by anthropogenic
emissions of hydrocarbons in the cities and by biogenic hydrocarbons in rural areas. A similar
representation of the factors contributing to the formation of O₃ during winter is shown in Fig.
S17.

705 Finally, a quantitative estimate of Eq. (4) is provided in Fig. 9. In the eastern regions of China, the
largest contribution to the diurnal mean value of $D(O_x)$ is due to the reaction between NO₂ and
OH, particularly in winter. This chemical path remains, however, the dominant loss channel during
summer in the polluted northern plain between Shanghai and Beijing. In urbanized area, the reac-
tion between H₂O and O(¹D) also play a relatively considerable role on the value of $D(O_x)$ in
710 summer. The relative contribution of ozonolysis reaction with alkene to the value of $D(O_x)$ is dis-
played in the southern China, which is associated with the high level of alkene in this area. In rural
area, the highest contribution is from the reaction between H₂O and O(¹D) in summer, followed
by the reaction between HO₂ and O₃.



715 5. Effects of aerosols on oxidants

The presence of aerosols in the atmosphere affects the abundance of atmospheric oxidants primarily through two different processes: (1) changes in the heterogeneous reaction rates associated with the uptake of several species by the particles (Tan et al., 2020; 2022), and (2) changes in the photolysis rate associated with enhanced extinction of solar light (Tie et al., 2001; 2005; Xing et al., 720 2017; Tan et al., 2022). Here we assess the relative importance of these two different mechanisms and derive the combined effect on the concentration of surface O₃.

5.1. Effects due to heterogeneous reactions

725 Figure S18 summarizes the response of surface NO, NO₂, OH, and HO₂ concentrations to the introduction of the added heterogeneous chemical reactions (R28-31 in Table 1) in the model. The concentration of NO_x species decreases due to the enhanced conversion of NO_x into HNO₃. In the eastern plain of China, we derive a reduction of up to 9 ppbv for NO₂ and 3 ppbv for NO in winter, 730 with a summertime decrease of 6 ppbv for NO₂ and 2 ppbv for NO. At the same time, the concentration of HO_x increases due to the enhanced formation of HONO, which is a source of HO_x in the presence of sunlight. This process overrides the expected reduction in HO₂ due to its uptake by the aerosol. We find an increase of up to 0.15 pptv for OH and 5 pptv for HO₂ in winter, and 0.3 pptv for OH and 8 pptv for HO₂ in summer.

735 We now examine how the uptake of HO₂, N₂O₅, and NO₂ on the surface of particles modifies the surface concentration of surface O₃ (Fig. 10). As shown by Fig. 10a, the uptake of HO₂ onto aerosols in January leads to a reduction in the surface concentration of ozone of about 3-4 ppbv, with the large decrease concentrated in Sichuan Basin and central China. In July (Fig. 10b), the highest 740 ozone changes are found in the North China Plain, especially in the vicinity of Beijing (about 3 ppbv). The varied high spots of aerosols effect of HO₂ uptake on ozone is associated with the distribution of aerosol surface area density. In winter (Fig. S12), the simulated high value of aerosol surface area density is derived in Sichuan Basin and central China, while, in summer, the high value is calculated in Beijing and surrounding areas. This indicates the high sensitivity of the effect 745 of HO₂ uptake on particles to the aerosol geometric parameters (Song et al., 2020).

The response of ozone to the uptake of N₂O₅ by aerosols is negative during winter when the competing photochemical conversion of NO_x to HNO₃ by the OH radical is very slow. The heterogeneous conversion of N₂O₅ to HNO₃ tends to reduce ozone by up to 3-4 ppbv in southern China 750 during winter (Fig. 10c), with limited effects in the summertime (Fig. 10d).

The uptake of NO₂ by aerosols tends to increase the wintertime ozone concentration by 8-9 ppbv in eastern China and in large urban areas of southern China (Fig. 10e) since the photolysis of HONO (formed from the heterogeneous NO₂ conversion) leads to enhanced concentrations of NO



755 and OH. As the simulated value of wintertime HO₂ is low (below 0.5 pptv) in large parts of China, the production of HO_x, from the photolysis of HONO, dominantly controls the value of $P(\text{RO}_x)$ (Fig. S14) and the formation of O₃ in this season. In summer (Fig. 10f), the concentration of ozone is reduced by 3-6 ppbv in the NO_x-sensitive rural areas of eastern and central China but is enhanced by 6-7 ppbv in VOC-sensitive urban areas. During this season, the high value of the HO₂ density
760 weakens the contribution of the HO_x produced by the HONO photolysis. However, the lower level of summertime NO_x strengthens the effect of the NO₂ loss resulting from the NO₂ uptake on particles.

The effect on near-surface ozone of the heterogeneous conversion of NO₃ by aerosols, also considered in the present model study (not shown), has been found to be very small.
765

The lowest panels of Fig. 10 show the change in ozone resulting from all four heterogeneous processes on aerosol surfaces. When combining the effects of all HO₂, NO₂, NO₃, and N₂O₅ heterogeneous reactions, we derive an ozone increase of 6-8 ppbv in winter and a decrease of 6-8 ppbv
770 in summer. However, ozone increases up to 8 ppbv in the VOC-limited metropolitan areas of Beijing, Shanghai, Guangzhou, and Chengdu. Comparison of Fig. 10 e, f with Fig. 10 g, h, respectively, suggests that heterogeneous loss of NO₂ on atmospheric particles discussed above dominates the impact of the studied heterogeneous reactions on surface O₃.

775 The O₃ response to the uptake of HO₂ and NO₂ is complex. The HO₂ + O₃ reaction provides a direct destruction mechanism for ozone, and the heterogeneous uptake of HO₂ contributes, therefore, to ozone enhancement. At the same time, the HO₂ uptake reduces the ozone production resulting from the reaction between HO₂ and NO, a photochemical process that is most efficient during summertime. The conversion of NO₂ to HNO₃ tends to reduce the O₃ formation in NO_x-
780 limited areas due to the loss of NO₂ by particles. However, in VOC-limited areas, the loss of NO₂ leads to an increase in the O₃ concentration. Moreover, the photolysis of HONO, which results from NO₂ uptake, produces NO and OH, which further affects the formation of O₃.

Our model simulation in July suggests that the presence of aerosol leads to a decrease of O₃ in
785 NO_x-limited areas and an increase of O₃ in VOC-limited areas. In other words, the continuous reduction in aerosol emissions observed in the past years should have led to an increased ozone concentration in NO_x-limited areas and a reduced ozone concentration in VOC-limited areas. The O₃ decrease in VOC-limited areas is the result of two opposite effects: the ozone decrease due to reduced NO₂ uptake (increased NO_x densities and enhanced ozone titration), and the ozone increase from reduced HO₂ uptake (increased HO₂ concentration and enhanced rate of the HO₂ +
790 NO reaction). The importance of HO₂ uptake by aerosols on ozone formation has been highlighted by several modeling studies (Li et al., 2021; Liu and Wang, 2021; Ivatt et al., 2022). However, recent studies based on field measurements (Tan et al., 2020; 2022; Dyson et al., 2022; Yang et



795 al., 2021) made in the urban/rural areas of northern and southern China (Wangdu, Beijing, Shen-
zhen, and Chengdu) during the summertime of 2014, 2017, 2018, and 2019, highlighted the minor
importance of HO₂ uptake for radical chemistry and O₃ formation, and showed the increasing im-
portance for the ozone production of the reduced NO₂ uptake by particles. One potential reason
for changes in the conclusions of these studies could be attributed to the sharp reduction in the
800 emissions of pollutants in China (Zheng et al., 2018), including the reduction in the aerosol load
and in the anthropogenic NO_x emissions. Another possible explanation is the adoption for the
analyses of different values for the uptake coefficients and for the aerosol geometric parameters
associated with the heterogeneous reactions affecting NO₂ and HO₂.

5.2. Effects due to photolysis

805

The presence of aerosols in the atmosphere tends to enhance the absorption and scattering of in-
coming solar radiation with direct impacts on the photolysis rates and hence on the abundance of
chemical species. Figure S19 shows a model estimate of the resulting effects on the surface con-
centrations of NO₂, NO, OH, and HO₂.

810

In the month of January, during which the aerosol burden is high, and the solar intensity is low,
the effect of light reduction by the aerosols through changes in the photolysis rates tends to increase
the surface concentration of NO_x, especially in the most populated and polluted urban areas (Bei-
jing, Shanghai, and Chengdu) where an increase in the concentration of NO₂ typically 0.5 ppbv is
815 derived. In these urban areas, the concentration of NO is increased by 0.5 ppbv. A reduction in
surface OH (about 0.1 pptv) and HO₂ concentrations (about 1.5 pptv) is derived, with the largest
effect occurring in the southeastern regions of China. The surface O₃ decreases by up to 4-5 ppbv
(Fig. 11a), with the highest decrease found in the Sichuan basin. In July, the aerosol burden is
lower, while the solar intensity is higher. The effect of light reduction by the aerosols tends to
820 increase the surface concentration of NO₂ by 1 ppbv in the North China Plain. In the case of OH
and HO₂, a decrease of 0.05-0.1 pptv and 2-3 pptv is found in the North China Plain. A decrease
of O₃ by up to 3-4 ppbv is derived in the Beijing and surrounding area (Fig. 11b).

5.3. Combined aerosol effects on ozone (uptake effects and photolysis).

825

When all heterogeneous processes affecting HO₂, N₂O₅, and NO₂ are simultaneously taken into
account, we derive for January an increase in the surface ozone concentration that is generally of
the order of 6-8 ppbv (Fig. 11c) in the middle of the country. The change in the photolysis rates
reduces the ozone concentration by 2-4 ppbv (Fig. 11a) and compensates to some extent the in-
830 crease due to aerosol uptake. Such a compensation mechanism was highlighted by Tan et al. (2022)
based on observations made in Shenzhen in the autumn of 2018. The combined effect in winter is
therefore limited, with ozone values increasing by less than 4 ppbv in most regions of China and
by less than 6 ppbv in the urban area.



835 In July, when combining the photolysis and uptake effects (Fig. S20), we derive a decrease in the
concentration of NO_2 (up to 10 pptv) and NO (up to 5 pptv) and an increase in the concentration
of OH (0.05 pptv) and HO_2 (up to 10 pptv) in eastern China. The response of ozone (Fig. 11b and
d) is characterized by a reduction in the surface concentration of about 10 ppbv (or 15%) in most
840 NO_x -limited regions of China. In the metropolitan areas of Shanghai and Beijing, an increase of
about 8 ppbv (or about 12%) is calculated.

These results highlight that, during summertime, the presence of aerosol particles leads to a decrease of the surface O_3 concentrations in NO_x -limited areas whereas it produces an increase in the ozone level in the VOC-limited (metropolitan) areas. Figure 12 presents a schematic view of different pathways that characterize the effects of aerosols on ozone concentrations. Specifically, this figure suggests that the reduction in the aerosol burden that has occurred in China in recent years should have produced an increase in surface ozone concentrations in NO_x -limited areas and a decrease in VOC-limited areas. The cause of the ozone increase in NO_x -sensitive areas should not be attributed exclusively to a reduction of the HO_2 uptake but to a combination of the different uptake processes and a reduction in the light extinction by the aerosols. The ozone decrease in VOC-limited areas should be mainly attributed to the NO_2 uptake with a counteracting effect by the HO_2 aerosol uptake and by the light extinction by the particles. Our results imply that, if the aerosol loading continues to decrease in the future, the ozone formation will increase so that the air quality measures currently implemented will become less efficient in NO_x -limited areas. This does not imply that ozone will necessarily increase in VOC-limited areas. With a further reduction in the NO_x emissions, which tends to shift the O_3 formation regimes from VOC-limited to NO_x -limited (Tan et al., 2022), the O_3 response to aerosol effects may gradually reverse in these geographical areas.

860 5.4. Effects of other HONO sources on ozone.

Figure 11e-f shows the changes in the surface ozone concentration due to all sources of HONO considered in the model, including direct emissions from transportation, gas phase production, and heterogeneous reactions of NO_2 uptake on aerosol surfaces and on ground surfaces. In January, the increase of the ozone concentration due to all these different processes reaches 9 ppbv and is more pronounced than in the “aerosol-only” case shown in Fig. 10e. In July, the decrease in the ozone concentration in eastern China (8-10 ppbv) and the increase in the metropolitan regions of Beijing and Shanghai (6-8 ppbv) are about 50% larger than when only the aerosol uptake of NO_2 is taken into account (see Fig. 10f).

870 Figure 11g-h also shows the changes in the surface ozone concentrations when all heterogeneous reactions involving HO_2 , NO_2 , NO_3 , and N_2O_5 as well as all sources of HONO are included in the model calculation. These two panels (g and h) must be compared with panels (c and d) of Fig. 11.



875 With the additional formation processes of HONO, surface ozone is increased by about 6-8 ppbv
in southeastern China during winter. For summer conditions, surface ozone concentration is re-
duced by up to 10 ppbv in the eastern and southern parts of China, but is increased by about 6-8
ppbv in the two major metropolitan centers.

880 6. Quantification of the oxidizing capacity of the atmosphere in China

6.1. OH reactivity

885 The model results presented above allow us to quantify the different factors that characterize the
oxidizing capacity of the atmosphere in China. We first analyze the geographical distribution of
the OH reactivity (Eq. 8a and 8b) resulting from the reaction of this radical with VOCs and CO
(noted VOC^R (Fig. 13a-b)) as well as NO_x (noted NO_x^R (Fig. 13c-d)). These quantities, and partic-
ularly the VOC^R/NO_x^R ratio (Fig. 13e-f), can be viewed as a proxy representing the competition
between radical production and destruction (see Kirchner et al., 2001).

890 During winter, the calculated value of the daytime averaged VOC^R ranges from typically 2 s^{-1} ,
mostly in rural areas, to 10 s^{-1} in the North China Plain between the urban areas of Shanghai and
Beijing, as well as in the area of Chengdu. The high value of calculated VOC^R in urbanized areas
is consistent with high values in the spatial distribution of wintertime VOCs, such as ethene (Fig.
S5), ethane (Fig. S5), and HCHO (Fig. S4). The values derived for the daytime averaged NO_x^R are
895 of the order of 10 s^{-1} in the North China Plain and most metropolitan areas of the country. Values
are close to 1 s^{-1} in rural areas. The VOC^R/NO_x^R ratio is of the order of 2 in most regions of China
except in the polluted areas where values close to 0.6 to 1 are derived.

900 During summer, the daytime averaged VOC^R parameter reaches values close to or higher than 10 s^{-1}
in southern China. This distribution of these high values is consistent with the spatial distribu-
tion of isoprene (Fig. S5) and HCHO (Fig. S4). The value of NO_x^R is smaller than in wintertime
with values generally close to 5 s^{-1} in the North China Plain, and approaching 10 s^{-1} inside the
cities of Beijing, Shanghai, Guangzhou, and Chengdu. The VOC^R/NO_x^R ratio is larger than 2 in the
entire spatial domain, except in a small polluted area of the North China Plain and in the urban
905 areas of Guangzhou and Shanghai where it is close to 1. Figure S21 shows the diurnal variation of
the simulated value of the VOC^R to NO_x^R ratio at different sites in January and July of 2018. The
daytime values range from 0.5 to 1.8 in urban sites in both two months, with the highest daytime
value shown in the Wangdu site in July (by the value of 5).

910 The diurnal variations of the OH reactivities due to different organic compounds, CO and nitrogen
dioxide in January and July are shown in Fig. 14. These calculated values need to be compared
with the data provided by the observations. In city sites, for example, Whalley et al. (2021) reported
diurnal variations of the OH reactivity in Beijing during the summer of 2018 with values of 25-35



915 s^{-1} to be compared to our model values of 23-42 s^{-1} in July. Based on measurements, the contribu-
tion to the OH reactivity of NO_x reactions is 40-50%, and of VOCs reactions 40-50%. The corre-
sponding values derived in our model study in July are 50% and 45%, respectively. In Shanghai
during summertime, Zhu et al. (2021) derived values of 10-25 s^{-1} , where the contribution of the
reaction of OH with NO_x is 33%, with CO is 26%, with OVOCs is 18%, and with alkenes is 15%.
920 The reactivity value derived by our model is 10-17 s^{-1} , with the highest contribution from the
reaction of OH with NO_x (50-60%). The values measured by Tan et al. (2018) in Chengdu (Sep-
tember 2016) are in the range of 15-30 s^{-1} to be compared to our July model values of 12-32 s^{-1} .
During the campaign that took place in Shenzhen in the autumn of 2018, Yang et al. (2022) derived
a total OH reactivity value that varied between 10 and 25 s^{-1} , which is relatively lower than our
calculated value by 20-45 s^{-1} in Guangzhou in summer.

925

Several other attempts have been made to derive the OH reactivity from in situ observations in
rural sites. At Wangdu during summertime, Tan et al. (2017) derived a reactivity value of 12-23 s^{-1} ,
and our model study provides a value of 8-22 s^{-1} . Tan et al. (2019), refer to the campaign con-
ducted at the Heshan site in the autumn of 2014, derived experimentally mean daytime OH reac-
930 tivities that range from 20 to 40 s^{-1} , with the contributions with CO, NMHCs, and NO_x , are 10%
(2-4 s^{-1}), 20% (4-8 s^{-1}), and 14% (3-6 s^{-1}), respectively. The value of VOC^R and NO_x^R in our model
account for approximately 1 to 5 s^{-1} and 2 to 5 s^{-1} , respectively. The underestimation of the HONO
concentration in our model (Table S5), contributes to the underestimation of the calculated value
of the OH radical and the OH reactivity.

935

Generally, in winter (Fig. 14a), the dominant contribution to OH reactivity in urban/rural sites is
through the reaction of OH and NO_x (40~60%), while, at a remote site (Waliguan) is from the
reaction between OH and CO (80%). In summer (Fig. 14b), the contribution of NO_x to OH reac-
tivity is still high in urban/rural sites (40~70%), with one exception at the Heshan site. At this site,
940 the largest reactivity of OH results from the reaction with alkene (40~50%), which is associated
with the relatively low value of NO_x (Fig. S1d) and high value of isoprene (Fig. S5) at this site.

We also show in Fig. 15 three other indicators; these describe the catalytic cycling of NO_x leading
to ozone formation until the chain reaction is interrupted: the radical chain length ChL (defined by
945 Eq. 9), the ozone production efficiency OPE (defined by Eq. 11), and the atmospheric oxidation
capacity AOC (defined by Eq. 15).

6.2. ChL

950 The value of the daytime averaged ChL , which, according to our definition, increases with the
atmospheric concentration of RO_x radicals and NO, is of the order of 3 to 5 cycles in remote areas.
In January (Fig. 15a), ChL reaches values as high as 8 to 10 cycles in the southern area of China.
The ChL values are low in northern and western China, where the NO concentrations are low



955 during winter. In July (Fig. 15b), the highest values of ChL (about 10 cycles) are found only in the metropolitan areas, where the HO_x concentrations are high. Since this parameter can be viewed as the ratio between the ozone and RO_x production rate (see Eq. 11), the polluted areas tend to favor ozone production for a given value of the RO_x formation rate. Examples of the diurnal variation of the chain length are provided in Fig. S22 in the Supplementary Information.

960 A few experimental estimates are available to quantify the value of the chain length: Zhu et al. (2021) derived a value of 2-6 for daytime ChL in Shanghai during the warm season of 2018 to be compared to the values of 4-5 provided by our model (Fig. S22). Yang et al. (2021) derive daytime values of 2-4.5 in Chengdu during summer, while our model provides values close to 4-5. In Guangzhou during summer, Wang et al. (2022) derive daytime values of 3-12 in fair agreement
965 with our calculated values of 6-10.

6.3. Ozone production efficiency (OPE)

The Ozone Production Efficiency (OPE) represents the number of ozone molecules produced by
970 NO_x before NO_x is further oxidized to form more stable nitrogen reservoirs or removed from the atmosphere by deposition. The daytime averaged OPE values are highest (larger than 30) in the southwest of China and are of the order of 25-30 above the Tibetan plateau (Fig. 15C-d) in both seasons. OPE values typically ranges between 3 to 15 in the eastern plain during both months under consideration. We note the similarities suggested by Eq. (14) between the distributions of
975 OPE and of the VOC^R/NO_x^R ratio in the VOC-limited regions of the China Northern Plain and over the Eastern China Sea where the effects of ship emissions are visible. Daytime OPE values are usually low at urban sites. More details on the diurnal variation of the OPE are provided in Fig. S23 of the Supplementary Information.

980 A few experimental data characterizing the OPE are available: Wang et al. (2017) summarize the summertime OPE values derived in China between 2006 and 2015 with values ranging from 2.1 to 20.2, which is comparable with our results that range from 1 to 30.

6.4. Atmospheric Oxidizing Capacity (AOC)

985 Finally, we show in Fig. 15e-f the spatial distribution of the daytime averaged value of the atmospheric oxidizing capacity AOC in January and July, respectively. In winter, the highest daytime values of AOC are found in the southern part of China, especially in the Pearl River Delta region ($3 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$). During nighttime (Fig. S24), the AOC values are lower than $0.2 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$,
990 with maximum values found at the southern coast of China. These high AOC values are associated with the spatial distribution of wintertime formaldehyde (Fig. S4), isoprene (Fig. S5), and of the HO_x and NO_3 radicals (Fig. S6). In summer, the values of daytime AOC are highest in the metropolitan urban areas (up to $10 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$), particularly in the vicinity of Beijing. The nocturnal



AOC values are lower than $2 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$ (Fig. S24) with high spots are found in urban areas..
995 The distribution of summertime *AOC* has some resemblance with the distribution of nitrogen species, including NO_2 (Fig. S1) and HONO (Fig. S4).

Figure 16 shows the diurnal evolution of the *AOC* and the dominant photochemical processes that contribute to this quantity in different urban and rural areas for January and July, respectively.
1000 First, we note that, in January, the noontime value of *AOC* does not supersede $6 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$ in urban areas and $3 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$ at rural sites (where the value is lower than $1.5 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$ in most cases). At the remote high-altitude station of Waliguan in western China (remote conditions), the maximum *AOC* value is lower than $0.2 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$. In July, as expected, the oxidizing capacity is larger than in winter, with noontime values reaching $(15-20) \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$ in metropolitan areas
1005 but limited increase in rural and remote areas. The increasing summertime *AOC* in city sites is attributed to the larger value of *AOC* contributed by OVOCs and alkene, which is associated with the higher value of summertime OVOCs (Fig. S4) and isoprene (Fig. S6) in urban areas.

Additional information on the relative contribution of different photochemical processes as a function of the time of the day and for two seasons is provided in Fig. 17. During winter, in Beijing during daytime, the major contributions to *AOC* are provided by the reaction of OH with alkenes, aromatics, and carbon monoxide. The situation is similar in the other three metropolitan areas under consideration. In Chengdu, however, the relative contribution of CO is larger as is the case at rural sites where the concentration of hydrocarbons is generally low. At night, the largest contribution in Beijing, Shanghai, and Chengdu is provided by the oxidation of hydrocarbons by ozone.
1015

During summertime, the daytime value of *AOC* in urban regions is determined by the reaction of OH with alkenes and with OVOCs. In remote areas, the dominant contribution is attributed to the reaction of OH with OVOCs and with CO. OVOCs are produced as a result of the oxidation of biogenic hydrocarbons such as isoprene, primarily but not exclusively in rural areas, and by the oxidation of anthropogenic hydrocarbons, mostly in urban and industrialized areas. Thus, any analysis of the processes that determine the value of *AOC*, particularly during summertime, must take into account the role played by OVOCs. Li et al. (2023) reported that the OVOCs have a significant impact on the atmospheric oxidative capacity in the Yangtze River Delta region, through reactions with OH during daytime and with NO_3 at night. Due to the limited amount of measurement data available (Wang W. et al., 2022), and the high uncertainties in the emissions of organic species (Li et al., 2023), the contributions of specific OVOCs to the oxidative process are still not unclear. More work is needed to obtain a better understanding of the impact of OVOCs on the oxidative processes in China.
1020
1025
1030

In all regions under consideration, during summertime, a dominant nighttime contribution to the value of *AOC* is provided by the oxidation of hydrocarbons by NO_3 . As the production of NO_3



1035 occurs through the reaction of O_3 and NO_2 , the higher summertime O_3 concentrations and temperature lead to the larger formation of NO_3 than during winter. Wang H. (2023) highlights the increasingly critical role of NO_3 -related nighttime oxidative chemistry associated with the positive trend in particulate nitrate abundance and in the formation of O_3 and other secondary pollutants. With the reduction in NO_x emissions, the importance of NO_3 radicals may become more notable.

1040 The model study confirms that, during daytime, more than 90% of *AOC* is due to reactions of chemical species with the OH radical. During nighttime, the oxidation processes are considerably slower and are due principally to the reactions of hydrocarbons with the nitrate radical (NO_3) and alkenes with ozone (O_3). In urban areas, the dominant daytime contributions to *AOC* during summertime are the reactions with carbon monoxide (20%), alkenes (35%), aromatics (10%), and
1045 OVOCs (30%). During winter, the corresponding numbers are 25% for carbon monoxide, 40% for alkenes, 15% for aromatics, and 15% for OVOCs, respectively. These approximate values vary somewhat from city to city. At night, during summer, the major contributions in urban areas are the oxidation by NO_3 (60% in Beijing, 45% in Chengdu, 15% in Shanghai and Guangzhou) and ozone (20% in Beijing, 35% in Shanghai, and 25% in Chengdu). At the very remote station of
1050 Waliguan, the relative daytime contributions to *AOC* in summer are 25% due to CO, 20% due to methane, 40% due to OVOCs, and 10% due to alkenes. The corresponding contributions during winter are 60% for CO, 20% for methane, 10% for OVOCs, and 5% for alkenes.

These model values can be compared, for example, with values calculated by Zhu et al. (2020)
1055 from experimental data obtained in Shanghai. The peak value around noontime in summer is about $(5-10) \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$ in fair agreement with our estimates. In the wintertime, the values are between $(5-8) \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$, i.e., slightly higher than our calculated *AOC*. Feng et al. (2021) report a maximum value of about $1.7 \times 10^8 \text{ cm}^{-3} \text{ s}^{-1}$ for the *AOC* at urban sites in Beijing during the summertime of 2014. This number is close to our calculated value of *AOC* by $1.8 \times 10^8 \text{ cm}^{-3} \text{ s}^{-1}$ in July. The peak
1060 *AOC* value of $2.1 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$ reported by Liu et al. (2021) for the winter of 2018 in Beijing is close to the model value of $3 \times 10^7 \text{ cm}^{-3} \text{ s}^{-1}$.

7. Summary and Conclusions

1065 The oxidizing capacity of the atmosphere can be characterized by different parameters including the production and destruction rates of ozone and other oxidants, the ozone production efficiency, the OH reactivity, and the length of the reaction chain responsible for the formation of ozone and RO_x . The value of these parameters depends on whether ozone formation is limited by the availability of NO_x or VOCs. It is also affected by the aerosol burden in the atmosphere, specifically by
1070 the rate at which heterogeneous chemical reactions take place in the atmosphere. In the present study, we have used a regional chemical transport model with a detailed chemical scheme to quantify these parameters in several chemical environments in China. Such studies should be helpful



1075 in determining the factors that are responsible for the documented changes in the oxidation capacity of the atmosphere and hence the mean concentration of surface ozone in different regions of the country.

1080 Our study shows that during winter, the formation of ozone in most of the eastern China Plain is VOC-limited. The ozone formation in remote western regions of the country, however, are NO_x-limited. In the south, an intermediate situation prevails, except in the Pearl River Delta area, where the formation of ozone is VOC-limited. In summer, ozone formation is NO_x-limited in most regions of China except in the urban areas of Beijing, Shanghai, Guangzhou, and Chengdu.

1085 Our model calculations conducted for the summer season show that the largest contribution to the formation of RO_x radicals in rural areas is due to the photolysis of ozone followed by the reaction between the electronically excited oxygen atom with water vapor. The second largest contribution is provided by the photolysis of formaldehyde. In urban and suburban areas, the formation of RO_x starts in the early morning with the photolysis of HONO, followed by the photolysis of HCHO and other OVOCs and finally of ozone. In polluted areas, the contribution of oxygenated VOCs is important and needs to be included in any oxidant budget analysis.

1090 The summertime destruction of RO_x radicals in the rural regions is principally due to radical-radical reactions, including HO₂ + HO₂, HO₂ + RO₂, and HO₂ + OH. In urban and suburban areas, the main destruction processes are associated with reactions between NO₂ and OH, and between RO₂ and NO. The destruction of radicals associated with the uptake of HO₂ on aerosol surfaces plays a limited role.

1100 At all the considered/studied sites, the production rate of ozone is with a relatively equal contribution of the reactions of NO with HO₂ and RO₂, respectively. The source of RO₂ varies according to the region; it is mostly anthropogenic in urban areas and biogenic in remote areas. Values of ozone production rate are substantially higher in metropolitan areas than in remote areas: the maximum value in the early afternoon reaches 100 ppbv h⁻¹ in Beijing and Shanghai but is less than 2 ppbv h⁻¹ at the rural sites.

1105 Our model simulations suggest that heterogeneous chemistry together with the effect of aerosol on light extinction (photolysis) contribute to an increase in the surface ozone concentrations by 4-6 ppbv during wintertime. In summer, the presence of the aerosol burden derived by our model leads to a reduction in surface ozone of up to 8 ppbv in the NO_x-limited areas of the central and eastern parts of China. The ozone concentrations, however, are enhanced by about 5-7 ppbv in the VOC-limited regions near Shanghai and Beijing. The reduction in the aerosol burden, resulting from the measures taken by Chinese authorities, will therefore lead to an increase of the O₃ density in the NO_x-limited rural areas and affect the efficiency of O₃ pollution control. A decrease of the O₃ concentration is expected in the VOC-limited areas. However, with the continuous reduction in



NO_x emissions, the NO_x-limited areas tend to geographically expand, and an O₃ increase should therefore occur in a broader area.

1115

The daytime averaged OH reactivity due to NO_x varies from less than 1 s⁻¹ in the western part of the country to 10 s⁻¹ in the North China Plain during winter. It is closer to 3 s⁻¹ during summer except in large urban areas where it is close to 10 s⁻¹. The reactivity due to VOCs is very small in the western regions of the country, but varies from 2 to 10 s⁻¹ in eastern China, with somewhat higher values in winter. The VOC^R-to-NO_x^R ratio is higher than 5 over the Tibetan Plateau, but less than 1.5 in the eastern regions of the country and even smaller in NO_x-rich regions of the most polluted areas.

1120

The number of cycles affecting RO_x radicals during daytime before they undergo a termination process is around 3 to 5 in remote areas. In January, it reaches as high as 8 to 10 cycles in the eastern plain and in the southern area of China. The values are low in northern and western China, where the NO concentration is low during winter. In the metropolitan areas during July, about 10 cycles are performed before the radicals are lost.

1125

Finally, our model simulations suggest that the daytime oxidizing capacity is mostly influenced by the reaction of the OH radical with alkenes, carbon monoxide, and oxygenated VOCs, and to a lesser extent with aromatics. The relative contribution of different chemical processes varies with locations (urban versus rural) and with seasons (winter versus summer). During nighttime, the largest contributions are due to the oxidation of hydrocarbons by NO₃ and ozone.

1130

1135

With the reduction of NO_x emissions observed in China, explicit consideration of nocturnal oxidative chemistry taking into account, the effect of nitrate radicals will become increasingly crucial for the assessment of air quality. With the reduction in the anthropogenic emissions of VOCs in China, the role of natural VOCs with high reactivity, such as isoprene, will become increasingly important regarding oxidative processes, especially in scenarios with increasing temperature and extreme weather associated with climate change. To understand the contribution of photodegradable OVOCs to the oxidative capacity of the atmosphere and to the formation of secondary pollutants, additional studies that include systematic measurements of OVOCs and more accurate time-dependent estimates of emissions will be needed.

1140

1145

Code and data availability. The WRF-Chem model is public available at <https://www2.mmm.ucar.edu/wrf/users/>. The air quality data at surface station are public available at the website of Ministry of Ecology and Environment of the People's republic of China at <http://english.mee.gov.cn/>.

1150



1155 *Author contributions.* JD and GB designed the structure of the manuscript, performed the numerical experiments, analyzed the results, and wrote the manuscript. JD analyzed the data and established the figures. All co-authors provided comments and reviewed the manuscript.

1160 *Competing interests.* The authors declare that they have no conflict of interest.

1165 *Acknowledgments.* The present joint Sino-German study was supported by the German Research Foundation (Deutsche Forschungsgemeinschaft DFG) and the National Science Foundation of China (NSFC) under the Air-Changes Project number 4487-20203. The National Center for Atmospheric Research (NCAR) is sponsored by the US National Science Foundation. We would like to acknowledge the high-performance computing support from NCAR Cheyenne.

1170 **References**

1175 Abdi-Oskouei, M., Roozitalab, B., Stanier, C. O., Christiansen, M., Pfister, G., Pierce, R. B., et al. The impact of volatile chemical products, other VOCs, and NO_x on peak ozone in the Lake Michigan region. *Journal of Geophysical Research: Atmospheres*, 127, e2022JD037042. <https://doi.org/10.1029/2022JD037042>, 2022.

1180 Bertram, T. H. and Thornton, J. A.: Toward a general parameterization of N₂O₅ reactivity on aqueous particles: the competing effects of particle liquid water, nitrate, and chloride, *Atmos. Chem. Phys.*, 9, 8351–8363, <https://doi.org/10.5194/acp-9-8351-2009>, 2009.

1185 Czader, B. H., Rappenglück, B., Percell, P., Byun, D. W., Ngan, F., and Kim, S., Modeling nitrous acid and its impact on ozone and hydroxyl radical during the Texas Air Quality Study 2006, *Atmos. Chem. Phys.*, 12, 6939-6951, <https://doi.org/10.5194/acp12-6939-2012>, 2012.

Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., Nakajima, T. Tropospheric Aerosol Optical Thickness from the GOCART Model and Comparisons with Satellite and Sun Photometer Measurements, *Journal of the Atmospheric Sciences*, 59(3), 461-483., [https://doi.org/10.1175/1520-0469\(2002\)059](https://doi.org/10.1175/1520-0469(2002)059), 2002.

1190

Dai, J., Brasseur, P., Vrekoussis, M., Kanakidou, M., Qu, K., Zhang, Y., Zhang, H., Wang, T.,: The atmospheric oxidizing capacity in China: roles of emission changes, in preparation, 2023.



- 1195 Dai, J., Liu, Y., Wang, P., Fu, X., Xia, M., Wang, T. The impact of sea-salt chloride on ozone through heterogeneous reaction with N₂O₅ in a coastal region of south China. *Atmospheric environment*, 236, 117604. <https://doi.org/10.1016/j.atmosenv.2020.117604>, 2020.
- 1200 Dai, J. and Wang, T.: Impact of international shipping emissions on ozone and PM_{2.5} in East Asia during summer: the important role of HONO and ClNO₂, *Atmos. Chem. Phys.*, 21, 8747–8759, <https://doi.org/10.5194/acp-21-8747-2021>, 2021.
- Dyson J. E., et al., Impact of HO₂ aerosol uptake on radical levels and O₃ production during summertime in Beijing, *Atmos. Chem. Phys. Discussions*, <https://doi.org/10.5194/acp-2022.800>, 2022.
- 1205 Elshorbany, Y. F., R. Kurtenbach, P. Wiesen, E. Lissi, M. Rubio, G. Villena, E. Gramsch, A. R. Rickard, M. J. Pilling, and J. Kleffmann, Oxidation capacity of the city air of Santiago, Chile, *Atmos. Chem. Phys.*, 9, 2257–2273, <https://doi.org/10.5194/acp-9-2257-2009>, 2009.
- 1210 Elguindi, N., Granier, C., Stavrou, T., Darras, S., Bauwens, M., Cao, H., et al. Intercomparison of magnitudes and trends in anthropogenic surface emissions from bottom-up inventories, top-down estimates, and emission scenarios. *Earth's Future*, 8, e2020EF001520. <https://doi.org/10.1029/2020EF001520>, 2020.
- 1215 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- 1220 Fast, J. D., Gustafson, W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E., Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, *J. Geophys. Res.*, 111, D21305, <https://doi.org/10.1029/2005JD006721>, 2006.
- 1225 Feng, T., Zhao, S., Hu, B., Bei, N., Zhang, X., Wu, J., Li, G. Assessment of Atmospheric Oxidizing Capacity Over the Beijing-Tianjin-Hebei (BTH) Area, China. *Journal of Geophysical Research: Atmospheres*, 126(7), <https://doi.org/10.1029/2020JD033834>, 2021.
- 1230 Fu, X., Wang, T., Zhang, L., Li, Q., Wang, Z., Xia, M., Yun, H., Wang, W., Yu, C., Yue, D., Zhou, Y., Zheng, J., and Han, R.: The significant contribution of HONO to secondary pollutants during a severe winter pollution event in southern China, *Atmos. Chem. Phys.*, 19, 1–14, <https://doi.org/10.5194/acp-19-1-2019>, 2019.



- 1235 Fu, X., Wang, T., Gao, J., Wang, P., Liu, Y., Wang, S., Persistent heavy winter nitrate pollution driven by increased photochemical oxidants in northern China. *Environmental Science & Technology*, 54(7), 3881–3889. <https://doi.org/10.1021/acs.est.9b07248>, 2020.
- 1240 Gaubert, B., L. K. Emmons, K. Raeder., Correcting model biases of CO in East Asia: impact on oxidant distributions during KORUS-AQ, *Atmos. Chem. Phys.*, 20, 14617-14647, <https://doi.org/105194/acp-20-14617-2020>, 2020.
- 1245 Granier, C., Darras, S., Denier van der Gon, H., Doubalova, J., Elguindi, N., Galle, B., et al., The Copernicus atmosphere monitoring service global and regional emissions. Copernicus Atmosphere Monitoring Service (CAMS). <https://doi.org/10.24380/d0bn-kx16>, 2019.
- 1250 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., & Wang, X. The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): An extended and updated framework for modeling biogenic emissions. *Geo-scientific Model Development*, 5, 1471–1492. <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- 1255 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., & Eder, B. Fully coupled “online” chemistry within the WRF model. *Atmos. environ.*, 39(37), 6957-6975. <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.
- 1260 Geyer, A., Alicke, B., Konrad, S., Schmitz, T., Stutz, J. and Platt, U.: Chemistry and oxidation capacity of the nitrate radical in the continental boundary layer near Berlin, *J. Geophys. Res.*, 106, 8013–8025, <https://doi.org/10.1029/2000JD900681>, 2001.
- 1265 Ivatt, P. D., M. J. Evans, and A. C. Lewis, Suppression of surface ozone by an aerosol-inhibited photochemical ozone regime, *Nature Geoscience*, <https://doi.org/10.1038/s41561-022-00972-9>, 2022
- 1270 Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34, 2131-2159, [https://doi.org/10.1016/S1352-2310\(99\)00462-8](https://doi.org/10.1016/S1352-2310(99)00462-8), 2000.
- 1275 Kirchner, F., Jeanneret, F., Clappier, A., Krüger, B., van den Bergh, H., Calpini, B. Total VOC reactivity in the planetary boundary layer: 2. A new indicator for determining the sensitivity of the ozone production to VOC and NO_x. *J. Geophys. Res.*, 106 (D3), 3095-3110. <https://doi.org/10.1029/2000JD900603>, 2001.
- 1280 Liu, J., Li, X., Tan, Z., Wang, W., Yang, Y., Zhu, Y., Yang, S., Song, M., Chen, S., Wang, H., Lu, K., Zeng, L., and Zhang, Y.: Assessing the ratios of formaldehyde and glyoxal to NO₂ as Indicators



- of O₃-NO_x-VOC sensitivity, *Environ. Sci. Technol.*, 55, 10935–10945, <https://doi.org/10.1021/acs.est.0c07506>, 2021
- 1275 Lakey, P. S. J., I. J. George, L. K. Whalley, M. T. Baeza-Romero, and D. E. Heard, Measurements of the HO₂ uptake coefficients onto single component organic aerosols, *Environ. Sci., Technol.*, 49, 4878–4885, <https://doi.org/10.1021/acs.est.5b00948>, 2015.
- 1280 Levy, H., Normal atmosphere: large radical and formaldehyde concentrations predicted, *Science*, 173, 141–143, <https://doi.org/10.1126/science.173.3992.14>, 1971.
- Li, G., W. Lei, M. Zavala, R. Volkamer, S. Dusanter, P. Stevens, and L. T. Molina, Impacts of HONO sources on the photochemistry in Mexico City during the MCMA-2006/MILOGO Campaign, *Atmos. Chem. Phys.* 10, 6551–6567, <https://doi.org/10.5194/acp-10-655102010>, 2010.
- 1285 Li, D. D., Xue, L. K., Wen, L., Wang, X. F., Chen T. S., Mellouki, A. Chen J. M., and Wang W. X., Characteristics and source of nitrous acid in an urban atmosphere in northern China: Results from 1-yr continuous observations, *Atmos. Environ.*, 182, 296–306, <https://doi.org/10.1016/j.atmosenv.2018.03.033>, 2018.
- 1290 Li, K.; Jacob, D. J.; Liao, H.; Shen, L.; Zhang, Q.; Bates, K. H., Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China. *Proc. Natl. Acad. Sci.*, 116 (2), 422–427, <https://doi.org/10.1073/pnas.1812168116>, 2019a.
- 1295 Li, K.; Jacob, D. J.; Liao, H.; Zhu, J.; Shah, V.; Shen, L.; Bates, K.H.; Zhang, Q.; Zhai, S. A two-pollutant strategy for improving ozone and particulate air quality in China. *Nat. Geosci.* 12, 906–910. <https://doi.org/10.1038/s41561-019-0464-x>, 2019b.
- 1300 Li, J., Li, X.L., Wang, X., Wang, H., Jing, S., Ying, Q., Qin, M., Hu, J. Fate of oxygenated volatile organic compounds in the Yangtze River Delta Region: source contributions and impacts on the atmospheric oxidation capacity. *Environ. Sci. Technol.* 56, 11212–11224. <https://doi.org/10.1021/acs.est.2c00038>, 2022.
- 1305 Li, Z., Xue, L., Yang, X., Zha, Q., Tham, Y. J., Yan, C., ... & Wang, W.. Oxidizing capacity of the rural atmosphere in Hong Kong, Southern China. *Science of the total environment*, 612, 1114–1122. <https://doi.org/10.1016/j.scitotenv.2017.08.310>, 2018.
- 1310 Liu, P., Xue, C., Ye, C., Liu, C., Zhang, C., Wang, J., Mu, Y. The Lack of HONO Measurement May Affect the Accurate Diagnosis of Ozone Production Sensitivity. *ACS Environ. Au*, <https://doi.org/10.1021/acsenvironau.2c00048>, 2022.



- 1315 Liu, T., Hong, Y., Li, M., Xu, L., Chen, J., Bian, Y., Yang, C., Dan, Y., Zhang, Y., Xue, L., Zhao, M., Huang, Z., and Wang, H.: Atmospheric oxidation capacity and ozone pollution mechanism in a coastal city of southeastern China: analysis of a typical photochemical episode by an observation-based model, *Atmos. Chem. Phys.*, 22, 2173–2190, <https://doi.org/10.5194/acp-22-2173-2022>, 2022.
- 1320 Liu, Y. and Wang Tao, Worsening urban ozone pollution in China from 2013 to 2017 – Part 2: The effects of emission changes and implications for multi-pollutant control, *Atmos. Chem. Phys.*, 20, 6323–6337, <https://doi.org/10.5194/acp-206323>, 2020.
- 1325 Liu Y., Lu K., Li, X., Dong H., Tan Z. Wang, H., Zou, Q., Wu, Y., Hu, M., Min, K.-E., Kecorius, S., Wiendensohler, A., and Zhang Y., A comprehensive model test of the HONO sources constrained to field measurements at rural North China Plain, *Environ. Sci., Technol.* 53, 3517–3525, <https://doi.org/10.1021/acs.est.8b06367>, 2019.
- 1330 Liu, Z., Wang, Y., Hu, B., Lu, K., Tang, G., Ji, D., Zhang, Y., Elucidating the quantitative characterization of atmospheric oxidation capacity in Beijing, China. *Science of the Total Environment*, 771, 145306., <https://doi.org/10.1016/j.scitotenv.2021.145306>, 2021.
- 1335 Lu, X., Hong, J. Y., Zhang, L., Cooper, O. R., Schultz, M. G., Xu, X. B., Wang, T., Gao M., Zhao, Y. H., and Zhang, Y. H., Severe surface ozone pollution in China: A global perspective, *Environ. Sci. Tech Lett.*, 5, 487–494, <https://doi.org/10.1021/acs.estlett.8b00366>, 2018.
- Ma, X., Tan, Z., Lu, K., Yang, X., Liu, Y., Li, S., Zhang, Y., Winter photochemistry in Beijing: Observation and model simulation of OH and HO₂ radicals at an urban site. *Sci. of the Total Environ.*, 685: 85–95, <https://doi.org/10.1016/j.scitotenv.2019.05.329>, 2019.
- 1340 Mao, J., Ren, X., Chen, S., Brune, W. H., Chen, Z., Martinez, M., Leuchner, M. Atmospheric oxidation capacity in the summer of Houston 2006: Comparison with summer measurements in other metropolitan studies. *Atmospheric Environment*, 44(33), 4107–4115. <https://doi.org/10.1016/j.atmosenv.2009.01.013>, 2010.
- 1345 Martinez, M., Harder, H., Kovacs, T. A., Simpasa, J. B., Bassis, J., Leshner, R., Zamora, R. J.. OH and HO₂ concentrations, sources, and loss rates during the Southern Oxidants Study in Nashville, Tennessee, summer 1999. *Journal of Geophysical Research: Atmospheres*, 108(D19).<https://doi.org/10.1029/2003JD003551>, 2003.
- 1350 Martin, R. V., D., J., Jacob, R. M. Yantosca, M. Chin and P. Ginoux, Global and regional decreases in tropospheric oxidants from photochemical effects of aerosols, *J. Geophys. Res.*, 108, <https://doi.org/10.1029/2002JD002622>, 2003.



- 1355 Mozurkewich, M., McMurry, P. H., Gupta, A., and Calvert, J. G.: Mass accommodation coefficient of HO₂ on aqueous particles, *J. Geophys. Res.*, 92, 4163–4170, <https://doi.org/10.1029/JD092iD04p04163>, 1987.
- 1360 Prinn, R. G., The cleansing capacity of the atmosphere, *Annu. Rev. Environ. Resour.*, 28, 29-57, <https://doi:10.1146/annurev.energy.28.011503.163425>, 2003.
- Schwarz, H. A., Bielski, B. H., Reactions of hydroperoxy and superoxide with iodine and bromine and the iodide (I₂⁻) and iodine atom reduction potentials. *The Journal of Physical Chemistry*, 90(7), 1445-1448. <https://doi/pdf/10.1021/j100398a045>, 1986.
- 1365 Schwartz, S. E., Mass-transport considerations pertinent to aqueous phase reactions of gases in liquid water clouds, in *Chemistry of Multiphase Atmospheric System*, NATO ASI Ser. Ser. G., vol6, edited by W. Jaeschke, pp. 415-471, Springer, New York. https://doi/10.1007/978-3-642-70627-1_16, 1986.
- 1370 Song, H., X. Chen, K. Lu et al., Influence of aerosol copper on HO₂ uptake: a novel parameterized equation, *Atmos. Chem. Phys.*, 20 15835-15850, <https://doi.org/10.5194/acp-20-15835-2020>, 2020.
- 1375 Song, H., Lu, K., Dong, H., Tan, Z., Chen, S., Zeng, L., & Zhang, Y. Reduced aerosol uptake of hydroperoxyl radical may increase the sensitivity of ozone production to volatile organic compounds. *Environmental Science & Technology Letters*, 9(1), 22-29. <https://doi.org/10.1021/acs.estlett.1c00893>, 2021.
- 1380 Ridley, B., Nitrogen compounds, in Brasseur G. P., J. J. Orlando and G. S. Tyndall, *Atmospheric Chemistry and Global Change*, Oxford University Press, pp.235-289, 1999.
- 1385 Skamarock, W.C.; Klemp, J.B.; Dudhia, J.; Gill, D.O.; Liu, Z.; Berner, J.; Wang, W.; Powers, J.G.; Duda, M.G.; Barker, D.M.; et al. *A Description of the Advanced Research WRF Model Version 4*; Mesoscale and Microscale Meteorology Laboratory NCAR: Boulder, CO, USA, 2019.
- 1390 Tan, Z., Fuchs, H., Lu, K., Hofzumahaus, A., Bohn, B., Broch, S., Zhang, Y. Radical chemistry at a rural site (Wangdu) in the North China Plain: observation and model calculations of OH, HO₂ and RO₂ radicals. *Atmos. Chem. Phys.*, 17(1), 663-690., <https://doi.org/10.5194/acp-17-663-2017>, 2017.
- Tan, Z., Lu, K., Jiang, M., Su, R., Dong, H., Zeng, L., Zhang, Y. Exploring ozone pollution in Chengdu, southwestern China: A case study from radical chemistry to O₃-VOC-NO_x sensitivity.



- Science of the Total Environment, 636, 775-786. <https://doi.org/10.1016/j.scitotenv.2018.04.286>, 2018.
- 1395
- Tan, Z., Lu, K., Hofzumahaus, A., Fuchs, H., Bohn, B., Holland, F., Liu, Y., Rohrer, F., Shao, M., Sun, K., Wu, Y., Zeng, L., Zhang, Y., Zou, Q., Kiendler-Scharr, A., Wahner, A., and Zhang, Y.: Experimental budgets of OH, HO₂, and RO₂ radicals and implications for ozone formation in the Pearl River Delta in China 2014, *Atmos. Chem. Phys.*, 19, 7129–7150, 1400 <https://doi.org/10.5194/acp-19-7129-2019>, 2019.
- Tan, Z., A. Hofzumahaus, K. Lu, S. S. Brown, F. Holland, L. G. Huey, A. Kiendler-Scharr, X. Li, X. Liu, and N. Ma, No evidence for a significant impact of heterogeneous chemistry on radical concentrations in the northern China Plain in summer 2014, *Environ. Sci., Technol.*, 54, 5973-1405 5979, <https://doi.org/10.1021/acs.est.0c00525>, 2020.
- Tan, Z., Lu, K., Ma, X., Chen, S., He, L., Huang, X., Zhang, Y. Multiple Impacts of Aerosols on O₃ Production Are Largely Compensated: A Case Study Shenzhen, China. *Environmental Science & Technology*, 56(24), 17569-17580, <https://doi.org/10.1021/acs.est.2c06217>, 2022.
- 1410
- Tie, X., G. Brasseur, L. Emmons, L. Horowitz, and D. Kinnison, Effects of aerosols on tropospheric oxidants: A global model study, *J. Geophys. Res.*, 106, 2931-2964, <https://doi.org/10.1029/2001JD900206>, 2001.
- 1415
- Tie, X., S. Madronich, S. Walters, D. P. Edwards, P. Ginoux, N. Mahowald, R. Y. Zhang, C. Lou and G. Brasseur, Assessment of the global impact of aerosols on tropospheric oxidants, *J. Geophys. Res.*, 110, D03204, <https://doi.org/10.1029/2004JD005359>, 2005.
- 1420
- Thornton, J. A., Kercher, J. P., Riedel, T. P., Wagner, N. L., Cozic, J., Holloway, J. S., ... & Brown, S. S. A large atomic chlorine source inferred from mid-continental reactive nitrogen chemistry. *Nature*, 464(7286), 271-274. <https://doi.org/10.1038/nature08905>, 2010.
- 1425
- Thompson, A. M., The oxidizing capacity of the earth's atmosphere: Probable past and future changes, *Science*, 256, 1157-1165, <https://doi.org/10.1126/science.256.5060.1157>, 1992.
- 1430
- Tonnesen, G. S. & Dennis, R. L. Analysis of radical propagation efficiency to assess ozone sensitivity to hydrocarbons and NO_x: 1. Local indicators of instantaneous odd oxygen production sensitivity. *Journal of Geophysical Research*, 105, 9213–9225. <https://doi.org/10.1029/1999jd900371>, 2000.
- Whalley, L. K., Slater, E. J., Woodward-Massey, R., Ye, C., Lee, J. D., Squires, F., Heard, D. E.,. Evaluating the sensitivity of radical chemistry and ozone formation to ambient VOCs and NO_x in



- Beijing. Atmospheric Chemistry and Physics, 21(3), 2125-2147. <https://doi.org/10.5194/acp-21-2125-2021>, 2021.
- 1435 Wang, H., Wang, H., Lu, X., Lu, K., Zhang, L., Tham, Y. J., Zhang, Y. Increased night-time oxidation over China despite widespread decrease across the globe. Nature Geoscience, 1-7. <https://doi.org/10.1038/s41561-022-01122-x>, 2023.
- 1440 Wang, P., Zhu, S., Vrekoussis, M., Brasseur, G. P., Wang, S., & Zhang, H. Is atmospheric oxidation capacity better in indicating tropospheric O₃ formation, Front. Environ. Sci., Eng., 16(5): 65, <https://doi.org/10.1007/s11783-022-1544-5>, 2022.
- 1445 Wang, W., Yuan, B., Peng, Y., Su, H., Cheng, Y., Yang, S., Shao, M.. Direct observations indicate photodegradable oxygenated volatile organic compounds (OVOCs) as larger contributors to radicals and ozone production in the atmosphere. Atmospheric Chemistry and Physics, 22(6), 4117-4128. <https://doi.org/10.5194/acp-22-4117-2022>, 2022.
- 1450 Wang T, Xue L, Brimblecombe P, Lam YF, Li L, Zhang L. Ozone pollution in China: A review of concentrations, meteorological influences, chemical precursors, and effects. Sci Total Environ. 575:1582-1596. <https://doi.org/10.1016/j.scitotenv.2016.10.081>, 2017.
- 1455 Wang, T., Xue, L., Feng, Z., Dai, J., Zhang, Y., Tan, Y. Ground-level ozone pollution in China: a synthesis of recent findings on influencing factors and impacts. Environmental Research Letters, 17(6), 063003. <https://doi.org/10.1088/1748-9326/ac69fe>, 2022.
- Wesely, M.L., Hicks, B.B., A review of the current status of knowledge in a dry deposition. Atmos. Environ., 34, 2261-2282, [https://doi.org/10.1016/S1352-2310\(99\)00467-7](https://doi.org/10.1016/S1352-2310(99)00467-7), 2000.
- 1460 Xia, R., Luo, Y., Zhang, D. L., Li, M., Bao, X., Sun, J., On the diurnal cycle of heavy rainfall over the Sichuan Basin during 10–18 August 2020., Advances in Atmospheric Sciences, 38, 2183-2200, <https://doi.org/10.1007/>, 2021.
- 1465 Xu, X., Ge, B., Lin, W. Progresses in the research of ozone production efficiency (OPE)., Adv. in Earth Science, 24(8), 845-853, <https://doi.org/10.11867/j.issn.1001-8166.2009.08.0845>, 2009.
- Xue, L., et al., Oxidative capacity and radical chemistry in the polluted atmosphere of Hong Kong and Pearl delta River region: analysis of a severe photochemical smog episode, Atmos. Chem. Phys., 16, 9891-9903, <https://doi.org/10.5194/acp-16-9891-2016>, 2016.
- 1470 Yang L. H., D. J. Jacob, N. K. Colombia., Tropospheric NO₂ vertical profiles over South Korea and their relation to oxidant chemistry: Implications for geostationary satellite retrievals and the



- observation of NO₂ diurnal variation from space, <https://doi.org/10.5194/egusphere-2022-1309>, 2022.
- 1475 Yang, X., Lu, K., Ma, X., Liu, Y., Wang, H., Hu, R., Zhang, Y. Observations and modeling of OH and HO₂ radicals in Chengdu, China in summer 2019. *Science of The Total Environment*, 772, 144829. <https://doi.org/10.1016/j.scitotenv.2020.144829>, 2021.
- 1480 Yang, X., Lu, K., Ma, X., Gao, Y., Tan, Z., Wang, H., ... & Zhang, Y. Radical chemistry in the Pearl River Delta: observations and modeling of OH and HO₂ radicals in Shenzhen in 2018. *Atmos. Chem. Phys.*, 22(18), 12525-12542. <https://doi.org/10.5194/acp-22-12525-2022>, 2022.
- 1485 Yu, C., Wang, Z., Xia, M., Fu, X., Wang, W., Tham, Y. J., Chen, T., Zheng, P., Li, H., Shan, Y., Wang, X., Xue, L., Zhou, Y., Yue, D., Ou, Y., Gao, J., Lu, K., Brown, S. S., Zhang, Y., and Wang, T.: Heterogeneous N₂O₅ reactions on atmospheric aerosols at four Chinese sites: improving the model representation of uptake parameters, *Atmos. Chem. Phys.*, 20, 4367–4378, <https://doi.org/10.5194/acp-20-4367-2020>, 2020.
- 1490 Yu, D., Tan, Z., Lu, K., Ma, X., Li, X., Chen, S., Zhang, Y. An explicit study of local ozone budget and NO_x-VOCs sensitivity in Shenzhen China. *Atmospheric Environment*, 224, 117304. <https://doi.org/10.1016/j.atmosenv.2020.117304>, 2020.
- 1495 Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K., Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *J. Geophys. Res.*, 113, D13204, <https://doi.org/10.1029/2007JD008782>, 2008.
- 1500 Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131–5153, <https://doi.org/10.5194/acp-9-5131-2009>, 2009.
- 1505 Zhang, S., Sarwar, G., Xing, J., Chu, B., Xue, C., Sarav, A., Ding, D., Zheng, H., Mu, Y., Duan, F., Ma, T., and He, H.: Improving the representation of HONO chemistry in CMAQ and examining its impact on haze over China, *Atmos. Chem. Phys.*, 21, 15809–15826, <https://doi.org/10.5194/acp-21-15809-2021>, 2021.
- 1505 Zhang, Y., Wen, X. Y., Wang, K., Vijayaraghavan, K., Jacobson, M. Z., Probing into regional O₃ and particulate matter pollution in the United States: 2. An examination of formation mechanisms through a process analysis technique and sensitivity study. *Journal of Geophysical Research*, 114. <https://doi.org/10.1029/2009jd011900>, 2009.



1510 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L.,
Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions
since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095–14111,
<https://doi.org/10.5194/acp-18-14095-2018>, 2018

1515 Zhu, J. S. Wang, H. Wang, S. Jing, S. Lou, A. Saiz-Lopez, and B. Zhou, Observationally con-
strained modeling of atmospheric oxidation capacity and photochemical reactivity in Shanghai,
China, *Atmos. Chem. Phys.*, 20, 1217-1232, <https://doi.org/10.5194/acp-20-1217-2020>, 2020.

1520

1525

1530

1535

1540

1545



1550

Table 1. Simplified chemical mechanisms used for the interpretation of model results

Reaction Number	Reactions	Reaction rate constant
Photolysis reactions		
R1	$O_3 + h\nu \rightarrow O(^1D) + O_2$	(Emmons et al., 2010)
R2	$NO_2 + h\nu \rightarrow NO + O$	
R3	$HCHO + h\nu + O_2 \rightarrow 2 HO_2 + CO$	
R4	$OVOC_i + h\nu \rightarrow RO_{x,i}$	
R5	$HONO + h\nu \rightarrow NO + OH$	
Thermal reactions		
R6	$O(^1D) + M \rightarrow O(^3P) + M$	(Emmons et al., 2010)
R7	$O + O_2 + M \rightarrow O_3 + M$	
R8	$O(^1D) + H_2O \rightarrow 2 OH$	
R9	$CH_4 + OH + O_2 \rightarrow RO_2 (= CH_3O_2) + H_2O$	
R10	$HC_i + OH + O_2 \rightarrow \alpha HO_2 + \beta RO_{2,i} + \text{oxidation products}$	
R11	$Alk_i + O_3 \rightarrow OH, HO_2, RO_{2,i}$	
R12	$CO + OH + O_2 \rightarrow CO_2 + HO_2$	
R13	$OH + O_3 \rightarrow HO_2 + O_2$	
R14	$HO_2 + O_3 \rightarrow OH + 2 O_2$	



R15	$\text{OH} + \text{HO}_2 \rightarrow \text{H}_2\text{O} + \text{O}_2$	
R16	$\text{HO}_2 + \text{HO}_2 + \text{M} \rightarrow \text{H}_2\text{O}_2 + \text{O}_2 + \text{M}$	
R17	$\text{HO}_2 + \text{RO}_{2,i} \rightarrow \text{ROOH}_i + \text{O}_2$	
R18	$\text{RO}_{2,i} + \text{RO}_{2,j} \rightarrow \text{products}$	
R19	$\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2$	
R20	$\text{NO} + \text{HO}_2 \rightarrow \text{NO}_2 + \text{OH}$	
R21	$\text{NO} + \text{RO}_{2,i} + \text{O}_2 \rightarrow \text{carbonyl} + \text{NO}_2 + \text{HO}_2$	
R22	$\text{NO} + \text{RO}_{2,i} \rightarrow \text{RONO}_{2,i}$	
R23	$\text{NO}_2 + \text{OH} + \text{M} \rightarrow \text{HNO}_3 + \text{M}$	
R24	$\text{NO} + \text{OH} \rightarrow \text{HONO}$	
R25	$\text{NO} + \text{NO}_2 + \text{H}_2\text{O} \rightarrow 2 \text{HONO}$	
R26	$\text{HONO} + \text{HONO} \rightarrow \text{NO} + \text{NO}_2 + \text{H}_2\text{O}$	(Zhang et al., 2021)
R27	$\text{HONO} + \text{OH} \rightarrow \text{NO}_2 + \text{H}_2\text{O}$	
Heterogeneous reactions		
R28	$\text{HO}_2 \rightarrow \text{H}_2\text{O}$	(Gaubert et al., 2020)
R29	$\text{N}_2\text{O}_5 \rightarrow 2 \text{HNO}_3$	(Bertram et al., 2009; Yu et al., 2020)
R30	$\text{NO}_3 \rightarrow \text{HNO}_3$	(Liu and Wang., 2020)



R31	$\text{NO}_2 \rightarrow 0.5 \text{HONO} + 0.5 \text{HNO}_3$	(Fu et al., 2019; Zhang et al., 2021)
-----	--	--

1555

1560

1565

1570

1575

1580

1585

1590



Table 2. Location of the observational sites or areas referred to in the present study

Site name	Latitude	Longitude	Site type
Beijing	39.90° N-40.10° N	116.30° E-116.50° E	Urban sites
Shanghai	31.10° N-31.30° N	121.40° E-121.60° E	Urban sites
Guangzhou	23.00° N-23.20° N	113.10° E-113.30° E	Urban sites
Chengdu	30.50° N-30.70° N	103.90° E-104.10° E	Urban sites
Wangdu	38.66° N	115.25° E	Rural site
Heshan	22.73° N	112.93° E	Rural site
Waliguan	36.17° N	100.54° E	Mountainous/Background site
Hok Tsui	22.22° N	114.25° E	Coastal/Background site

1595

1600

1605

1610

1615



1620

Table 3. Sensitivity experiments

Modeling cases	Description
<i>Het-All</i>	With all heterogeneous reactions and all other HONO sources
<i>No-HetHO2-Aero</i>	Without HO ₂ uptake on aerosols
<i>No-HetNO3-Aero</i>	Without NO ₃ uptake on aerosols
<i>No-HetN2O5-Aero</i>	Without N ₂ O ₅ uptake on aerosols
<i>No-HetNO2-Aero</i>	Without NO ₂ uptake on aerosols
<i>No-Het-Aero</i>	Without HO ₂ , NO ₃ , N ₂ O ₅ , and NO ₂ uptake reactions on aerosols
<i>No-Phot</i>	Without aerosol effects on light extinction and photodissociation
<i>No-Het-Aero-Phot</i>	Without HO ₂ , NO ₃ , N ₂ O ₅ , and NO ₂ uptake reactions on aerosols and aerosol effects on light extinction and photodissociation
<i>No-HONO</i>	Without HONO sources
<i>No-Het-HONO-Phot</i>	Without HO ₂ , NO ₃ , N ₂ O ₅ , and NO ₂ uptake reactions on aerosols, other HONO sources and aerosol effects on light extinction and photodissociation

1625



1630

1635

Table 4. Comparison between values of the production rate of RO_x [Units: ppbv h⁻¹] derived from local observations and calculated by our regional model in July 2018.

Location	Period	Me. ^a	Ca. ^b	Me. ^a	Ca. ^b	Me. ^a	Ca. ^b	Me. ^a	Ca. ^b
		Peak value ^c		HONO ^d		HCHO ^e		Ozone ^f	
Beijing ^g	Spring 2018	7.0	7.5	4.0	4.0	1.5	2.0	1.5	1.5
Shanghai ^h	Summer 2021	4.5	5.0	2.5	1.0	1.0	1.0	1.0	1.5
Guangzhou ⁱ	Autumn 2018	5.6	6.0	2.0	3.0	1.5	1.0	1.0	1.5
Chengdu ^j	Summer 2019	7.0	7.5	2.0	2.5	2.0	1.5	1.5	2.0
Wangdu ^k	Summer 2014	5.0	4.8	2.0	2.0	1.0	2.0	1.5	1.0
Heshan ^l	Autumn 2014	4.0	1.1	2.0	0.5	1.5	0.4	0.5	0.5

1640 ^a Measured value in relevant periods; ^b Calculated value in our study in July 2018; ^c Peak value of
 production rate of RO_x; ^{d,e,f} Peak value of the production rate of RO_x from the photolysis of ^d
 HONO, ^e HCHO and ^f O₃; ^{g,h,i,j,k,l} Observations from the studies of ^g Whalley et al. (2021), ^h Zhu et
 al. (2021), ⁱ Wang et al. (2022), ^j Yang et al., (2022), ^k Tan et al., (2017) and, ^l Tan et al. (2019).

1645

1650





Figure 1. Location of sites (stars) considered in our analysis (from © Google Maps).

1655

1660

1665

1670




1675 **Figure 2.** Display of regions in which the ozone production is limited by the availability of nitro-
gen oxides (NO_x-limited, in green), and volatile organic carbon (VOC-limited, in red) in January
1680 (a) and July (b), 2018. The regions with intermediate conditions (Transition) are shown in blue.
The indicator used to define these regions is the concentration ratio between formaldehyde
(HCHO) and nitrogen dioxide (NO₂).

1680

1685

1690


1695



1700

Figure 3. Spatial distribution of the production rate of RO_x (RO_2+HO_2+OH) [$P(RO_x)$, Unit: ppbv h^{-1}] (*Het-All* case) from the photolysis of nitrous acid (HONO) (a, b), formaldehyde (HCHO) (c, d), non-HCHO oxidized volatile organic compounds (OVOCs) (g, h) and O_3 (Reactions between O^1D and H_2O ; e, f) in the daytime (08:00-19:00 Local Standard Time (LST)) of January (left column: a, c, e, g) and July (right column: b, d, f, h). Note the difference in scales among panels.

1705



1710 **Figure 4.** Diurnal variation of the production rate of RO_x [$P(RO_x)$], Unit: $ppbv\ h^{-1}$] in different regions of China calculated for July 2018.






Figure 5. Spatial distribution of destruction rate of RO_x [$D(\text{RO}_x)$, Unit: ppbv h⁻¹] (a, b) and the relative contribution $L_N/D(\text{RO}_x)$ (c, d), $L_H/D(\text{RO}_x)$ (e, f) and $L_{hel}/D(\text{RO}_x)$ (g, h) in the daytime of January (left column: a, c, e, g) and July (right column: b, d, f, h).



1720 **Figure 6.** Diurnal variation of the photochemical destruction rate of RO_x [$D(RO_x)$, Unit: $ppbv\ h^{-1}$] in eight sites of China for July 2018. The contributions to the destruction rate are the following: L_H accounts for the following reactions: $HO_2 + HO_2$, $HO_2 + RO_2$, $RO_2 + RO_2$, and $OH + HO_2$. L_N accounts for $OH + NO_2$ and $RO_2 + NO$. L_{HO_2} to the uptake of HO_2 by particles.



1725

Figure 7. Spatial distribution of production rate of odd oxygen [$P(O_x)$, Unit: ppbv h^{-1}] (a, b) and the relative contributions from the reactions between HO_2 and NO (c, d) and RO_2 and NO (e, f) in the daytime of January (left column: a, c, e) and July (right column: b, d, f).

1730





Figure 8. Diurnal variation of the O_x production rate [$P(O_x)$, Unit: ppbv h⁻¹] in different regions of China for July 2018.



1735

Figure 9. Spatial distribution of destruction rate of odd oxygen (a, b) [$D(O_x)$, Unit: ppbv h⁻¹] (*Het-All* case) and the relative contributions from the reactions of OH with NO₂ (c, d), O(¹D) with H₂O (e, f), Alkene with O₃ (g, h) and HO₂ with O₃ (i, j) in the daytime of January (left column: a, c, e, g, i) and July (right column: b, d, f, h, j).

1740




Figure 10. Spatial distribution of the response of the monthly average surface O_3 concentration [Unit: ppbv] to the aerosol uptake by HO_2 (a, b; *Het-All minus No-HetHO2-Aero*), N_2O_5 (c, d, *Het-All minus No-HetN2O5-Aero*), NO_2 (e, f, *Het-All minus No-HetNO2-Aero*), and to the uptake by all these processes (g, h, *Het-All minus No-Het-Aero*). The results are shown for the daytime of January (left column: a, c, e, g) and July (right column: b, d, f, h) of 2018.

1745






Figure 11. Changes in the surface concentrations of daytime O₃ [Unit: ppbv] resulting from the effect of aerosol-related solar light extinction on photolysis for January (a) and July (b) (*Het-All* minus *No-Phot*), the combined effect of photolysis and aerosol uptake for January (c) and July (d) (*Het-All* relative to *No-Het-Aero-Phot*), effects of NO₂ uptake by aerosols and on the surfaces as well as direct HONO emissions from traffic and gas phase formation for January (e) and July (f) (*Het-All* minus *No-HONO*), and from all NO₂, N₂O₅, NO₃ and HO₂ uptake processes photolysis effects and other HONO sources for January (g) and July (h) (*Het-All* minus *No-Het-HONO-Phot*).



1755


Figure 12. Schematic for the impact of aerosol through aerosol extinction of solar radiation and heterogeneous processes on ozone concentration. Arrows represent the changes in chemicals and photolysis rate associated with the reduction of aerosols.

1760



1765

Figure 13. Spatial distribution of reactivity of VOCs (VOC^R) [Unit: s^{-1}] and NO_x (NO_x^R ; c, d) [Unit: s^{-1}], VOC^R to NO_x^R ratio (e, f) (*Het-All* case) for daytime conditions in January (left column: a, c, e) and July (right column: b, d, f).




1770

Figure 14. Diurnal variation of different contributions to the OH reactivity [Unit: s⁻¹] in cities and remote sites. The values associated with alkenes include the contribution of biogenic isoprene and terpenes. The two upper rows refer to January (a) and the two lower rows to July (b).

1775

1780



1785

Figure 15. Spatial distribution of radical chain length (ChL) in January (a) and July(b), ozone production efficiency (OPE) in January (c) and July(d), and atmospheric oxidation capacity (AOC [Unit: 10^7 molecular $cm^{-3} s^{-1}$]) in January (e) and July(f) for daytime conditions extracted from the
1790 *Het-All* case.

1795





Figure 16. Diurnal variation of the atmospheric oxidizing capacity (AOC) [Unit: 10^7 molecules $\text{cm}^{-3} \text{s}^{-1}$] in cities and remote sites. The effect of alkenes includes the contribution of biogenic isoprene and terpenes, while the effect of OVOCs includes the contribution of formaldehyde. The

1800 two upper rows refer to January (a) and the two lower rows to July (b).



1805

Figure 17. Same as Fig. 16, but expressed in relative terms [Unit: %] and highlighting the nighttime contribution to the *AOC* value. The two upper rows refer to January (a) and the two lower rows to July (b).