



# Technical note: Bimodal Parameterizations of in situ Ice Cloud Particle Size Distributions

Irene Bartolomé García<sup>1,a</sup>, Odran Sourdeval<sup>2</sup>, Reinhold Spang<sup>1</sup>, and Martina Krämer<sup>1,3</sup>

<sup>1</sup>Institute for Energy and Climate Research (IEK-7), Research Center Jülich, D-52425 Jülich, Germany

<sup>2</sup>University of Lille, CNRS, UMR 8518-LOA-Laboratoire d'Optique Atmosphérique, F-59000 Lille, France

<sup>3</sup>Institute for Physics of the Atmosphere (IPA), Johannes-Gutenberg University, D-55122 Mainz, Germany

<sup>a</sup>Now at: Institute of Geophysics and Meteorology, University of Cologne, Cologne, Germany

**Correspondence:** ibartolo@uni-koeln.de, m.kraemer@fz-juelich.de

**Abstract.** The cloud particle size distribution (PSD) is a key parameter for the retrieval of micro-physical and optical properties from remote sensing instruments, which in turn are necessary for determining the radiative effect of clouds. Current representations of PSDs for ice clouds rely on parameterizations that were largely based on in situ measurements where the distribution of small ice crystals were uncertain. This makes current parameterizations deficient to simulate remote sensing observations sensitive to small ice, such as from lidar and thermal infrared instruments. In this study we fit the in situ PSDs of ice crystals from the JULIA (JÜLich In situ Aircraft data set) database, which consists of 11 campaigns covering the tropics, mid-latitudes and the Arctic, consistently processes and considered more robust in their measurements of small ice. For the fitting, we implement an established approach to PSD parameterizations, which consists in finding an adequate set of parameters for a modified gamma function after normalization of both PSD axes. These parameters are constrained to match in-situ measurements when predicting microphysical properties from the PSDs, via a cost function minimization method. We selected the ice water content and the ice crystal number concentration, which are currently key parameters for modern satellite retrievals and model microphysics schemes. We found that a bimodal parameterization yields better results than a monomodal one. The bimodal parameterization has a lower spread for almost all ice crystal sizes over the entire range of analyzed temperatures and fits better the observations, especially for particles between 20 and about 110  $\mu\text{m}$  at temperatures between  $-60$  and  $-20$   $^{\circ}\text{C}$ . For this temperature range, the root mean square error for the retrieved  $N_{\text{ice}}$  is reduced from 0.36 to 0.20. This demonstrates a clear advantage to considering the bimodality of PSDs, e.g. for satellite retrievals.

## 1 Introduction

Ice clouds play an important role in Earth's radiative budget, as their radiative effects can either contribute to a warming or a cooling of the surface (Liou, 1986; Stephens et al., 1990). The balance between these two effects depends on their macro- and micro-physical properties, which stem from an array of very complex processes governing ice crystal formation and growth (Zhang et al., 1999; Krämer et al., 2020). Despite extensive research over the past decades most of these processes remain highly uncertain, making ice clouds major unknowns in current climate studies (Bellouin et al., 2020). Ice clouds are particularly challenging for satellite remote-sensing techniques, in great part due to the complexity and variety of their



microphysical and optical properties (Baran, 2009). In turn, this lack of accurate global observational constraints leads to critical shortfalls for evaluation efforts of predictions of ice cloud properties and processes in numerical weather forecast and climate models (Lohmann et al., 2007).

The particle size distribution (PSD), which describes the number concentration of ice crystals as a function of their size, impacts most microphysical and radiative properties of clouds. This makes the PSD a central parameter in remote-sensing retrieval techniques (Yang et al., 2001; Vidot et al., 2015). PSD shapes can greatly vary, as they are influenced by formation and growth mechanisms dictated by the environment in which the cloud developed (Heymsfield and McFarquhar, 2002). For instance, the cloud origin (e.g. in-situ or liquid-origin; Krämer et al., 2016) was identified as having a major influence on PSD shapes for cirrus (Luebke et al., 2016). However, the exact drivers to PSD shapes remain poorly understood and their global variability only addressed by few modelling studies (e.g. Gasparini and Lohmann, 2016). For this reason, retrieval techniques must make critical simplifications regarding PSD shapes, which are commonly reflected by the use of fixed parameterizations that are universally applied globally and for all cloud types. Such recent parameterizations rely on advanced normalization procedures that aim to make PSDs more representative of a large sample of ice clouds (Delanoë et al., 2005; Field et al., 2007, also see Section 3). While this allows for more accurate representations of ice clouds in retrieval methods (e.g. Sourdeval et al., 2015, 2016), current parameterizations still have important issues. For instance, they largely rely on in-situ measurements where the distribution of small ice crystals (sizes less than 100  $\mu\text{m}$  diameter) are at best very uncertain (Korolev et al., 2011), which makes them deficient to simulate remote-sensing observations sensitive to small ice and leads to erroneous retrievals, especially for estimations of the ice crystal number concentration ( $N_{\text{ice}}$ ). Reaching more accurate estimations of  $N_{\text{ice}}$  is particularly of great importance to understand ice cloud formation mechanisms and for improving their predictions in models, since this parameter is often used as a prognostic variable to predict the evolution of clouds (e.g. Seifert and Beheng, 2006).

An example of widely used application of normalized PSD parameterizations in satellite retrievals is found in the DARDAR algorithm (raDAR/liDAR; Delanoë and Hogan, 2008, 2010). DARDAR retrieves vertical profiles of ice cloud microphysical properties by using a combination of coincident measurements from the Cloud Profiling Radar (CPR) onboard CloudSat and from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP). Although the original algorithm does not include retrievals of  $N_{\text{ice}}$ , Sourdeval et al. (2018) explored the capabilities of its framework to retrieve this parameter (DARDAR-Nice). However, comparisons of  $N_{\text{ice}}$  retrievals by DARDAR-Nice to recent in-situ measurements highlighted strong limitations linked to its used parameterizations of PSDs, which are in specific cases not suited to retrieve  $N_{\text{ice}}$  (Sourdeval et al., 2018; Krämer et al., 2020). A main reason is that the current parameterization only represents a single ice crystals size mode, which can result in PSDs being significantly misrepresented when they happen to be strongly bimodal, i.e. consisting of two size modes. The bimodality of the PSDs can often occur in cirrus clouds and is a function of temperature and location within the cloud (Zhao et al., 2011), having warmer clouds more often bimodal PSDs (Jackson et al., 2015). Associated with the study of bimodality is the analysis of the effect of shattering ice crystals at the inlets of the in situ instruments that can artificially increase the concentration of small particles. This problem has been widely discussed and as a result, new inlets were designed and correction algorithms applied to minimize this effect (Korolev et al., 2011; Lawson, 2011; Krämer et al., 2020).




The present study investigates the benefits of considering a second mode in existing PSD parameterizations methods. It follows the framework proposed by Delanö et al. (2005, 2014) (hereafter D05 and D14), which is applied on an extensive database of in-situ ice cloud properties from airborne campaigns (JULIA; (Krämer et al., 2016)), which is considered less sensitive to shattering effects. This study proposes a new set of parameterizations, based on single- and double-mode PSDs, that will be useful to improve remote-sensing retrieval methods. Section 2 describes the JULIA database and Section 3 details the PSD normalization method. Section 4 presents and analyze the newly developed parameterizations by comparison to the original in-situ data. Section 5 concludes this study.

## 65 2 The JULIA Database

JULIA (JÜLich In situ Airborne database) is a compilation of in situ measurements of cirrus, mixed-phase and liquid clouds, water vapor and also other trace gases, collected at the Research Center Jülich starting in 2008 (Schiller et al., 2008; Luebke et al., 2013; Costa et al., 2017; Krämer et al., 2016; Krämer et al., 2018; Krämer et al., 2020) and continued until 2021. The measurements were taken onboard of different research aircraft and cover the tropics, mid-latitudes and the Arctic (Fig. 1). For our study, we focus only on the data of ice particles, which includes data from 11 field campaigns with measurements of ice crystals of diameters from 3 to 1000  $\mu\text{m}$  taken every second. The analyzed data, i.e., the  $N_{\text{ice}}$  and IWC, was sampled with (or computed from) the NIXE-CAPS (New Ice eXperiment: Cloud and Aerosol Particle spectrometer), which is a combination of CAS (Cloud and Aerosol Spectrometer) and a CIPg (Cloud Imaging Probe - grayscale), and combinations of the FCDP (Fast Cloud Droplet Probe) and 2D-S (Two-Dimensional Stereo), CDP (Cloud Droplet Probe) and 2D-C (Two-Dimensional Cloud) instruments (the respective used size ranges are listed in Table2). These instruments and their data processing procedures are described in the literature (e.g., Lawson et al., 2006; McFarquhar et al., 2007; Lawson, 2011; Krämer et al., 2016; Luebke et al., 2016; Baumgardner et al., 2017; Afchine et al., 2018; Krämer et al., 2020). Since the widths of the size bins of each pair of instruments are different, Krämer et al. (2022) synchronized all PSDs to the same grid with logarithmically equidistant size bins in order to facilitate inter-comparisons and the interpretation of the results. The data included in JULIA has undergone an extensive quality check process to warranty its validity. In the following section, a brief description of the campaigns and the instruments is given.

### 2.1 Selected airborne campaigns

The analyzed data covers a wide range of meteorological conditions and extend from the tropics to the polar region. In total there are 5 campaigns in the tropics (ACRIDICON-CHUVA, ATTREX, CONTRAST, POSIDON and STRATOCLIM), 3 in mid-latitudes (COALESC, ML-CIRRUS and START), one covering both mid and high-latitudes (CIRRUS-HL) and 2 in high latitudes (RACEPAC and VERDI). ACRIDICON consisted on flights over the Amazonian forest in September 2014 with the aim of studying the interaction between aerosols and deep convective clouds (Wendisch et al., 2016). During ATTREX (February-March 2014), measurements inside cirrus clouds were taken over the West tropical Pacific, around the tropical tropopause layer, avoiding convection (Thornberry et al., 2017). CONTRAST, based on Guam, took also place in the West



**Figure 1.** Location of all campaigns of the JULIA data base. In different colors, the campaigns used for this study, other campaigns in grey. (Figure by Nicole Spelten).

90 tropical Pacific (January-February 2014) to study tropical convection, oceanic processes and ozone chemistry in the upper troposphere - lowermost stratosphere (UTLS) (Pan et al., 2017). POSIDON (October 2016) was also based on Guam and among its objectives were the study of cirrus clouds and dehydration in the tropical UTLS (POSIDON: last accessed: 29 July 2022). During POSIDON three flights were near an active typhoon. The last campaign in the tropics is STRATOCLIM (July-August 2017), based in Nepal and focused on water vapor variations and upper tropospheric clouds (Krämer et al., 2020; 95 Khaykin et al., 2022). COALESC took place during February and March 2011 and the observations were taken over the South-East coast of England and Wales. Its main focus was on stratocumulus clouds, but there are also plenty of observations within mixed-phase and cirrus clouds (Osborne et al., 2014). During March and April 2014, ML-CIRRUS took place, centered in observations over Europe and the North Atlantic focusing on processes involving cirrus (Voigt et al., 2017). START was based in Colorado, USA during April-June 2008 (Pan et al., 2010). CIRRUS-HL is a campaign from July-August 2021, based in 100 Oberpfaffenhofen, Germany, that investigated microphysical properties and climate impact of ice clouds in high latitudes, as well as the effect of aviation (CIRRUS-HL, last accessed: 29 July 2022). VERDI (April-May 2012, Inuvik, Canada) (Costa



**Table 1.** Instruments used during the field campaigns of the JULIA data base (NIXE-CAPS as a combination of CAS: Cloud and Aerosol Spectrometer and CIPg: Cloud Imaging Probe greyscale. CDP: Cloud Droplet Probe, FCDP: Fast Cloud Droplet Probe, 2D-C: Two-Dimensional Cloud spectrometer, 2D-S: Two-Dimensional Stereo Cloud spectrometer) \*The range of the instruments might be larger.

Instruments	Campaigns	Aircraft	Years	Used ranges*
CDP + 2D-C	START, CONTRAST	HIAPER	2008, 2014	3 – 50 $\mu\text{m}$ + 60 – 1000 $\mu\text{m}$
FCDP + 2D-S	ATTREX, POSIDON	Global Hawk, WB-57	2014, 2016	3 – 25 $\mu\text{m}$ + 25 – 1000 $\mu\text{m}$
NIXE-CAPS	COALESC, VERDI, RACEPAC	BAe-146, Polar-5/6,	2011, 2012, 2014	3 – 25 $\mu\text{m}$ + 25 – 930 $\mu\text{m}$
NIXE-CAPS	ML-CIRRUS, ACRIDICON-CHUVA	HALO	2014	3 – 25 $\mu\text{m}$ + 25 – 930 $\mu\text{m}$
NIXE-CAPS	STRATOCLIM, CIRRUS-HL	Geophysica, HALO	2017, 2021	3 – 25 $\mu\text{m}$ + 25 – 930 $\mu\text{m}$

et al., 2017) aimed to investigate, among other goals, the radiation budget of ice clouds in the Arctic and the influence of convective transport in the upper troposphere. During this campaign a persistent anticyclone was present and favored the formation of persistent stratus (Klingebiel et al., 2015). RACEPAC (Costa et al., 2017) is the follow up campaign of VERDI and took place in April-May 2014.

In Table 1 a summary of the used instruments for each campaign is given. The NIXE-CAPS is formed by CAS and CIPg that combined cover particles with diameters from 0.61 to 930  $\mu\text{m}$ . To avoid overlap of particle sizes, the PSDs from NIXE-CAPS are merged between 20 and 25  $\mu\text{m}$ . Detailed explanations about the data analysis methods, including position of the instrument in the aircraft and shattering effects, are given by Meyer (2013); Krämer et al. (2016); Luebke et al. (2016); Costa (2017). Other instruments are the light scattering sensors CDP that measures concentration for particles with diameters between 2 and 50  $\mu\text{m}$  (McFarquhar et al., 2007) and the FCDP for particles between 1 and 50  $\mu\text{m}$  (Baumgardner et al., 2017). The optical imaging cloud probe 2D-C spectrometer (Baumgardner et al., 2017) and the 2D-S spectrometer are used to reconstruct cloud particle shapes and sizes between 25 – 800  $\mu\text{m}$  and 5 – 1280  $\mu\text{m}$ , respectively. The 2D-S includes tips and software to reduce shattering effects (Lawson et al., 2006; Lawson, 2011). FCDP and 2D-S PSDs are merged between 20 and 25  $\mu\text{m}$  and the ones from CDP and 2D-C at 55  $\mu\text{m}$ . For more information about the campaigns and the instruments the reader is also referred to Costa (2017); Krämer et al. (2020).

## 2.2 Ice crystal number concentration and ice water content computation

$N_{ice}$  is the sum of the ice crystal number concentration of each size bin. The IWC is obtained following:

$$IWC = \sum_{i=1}^n m_i \cdot \Delta N_{ice, i}, \quad (1)$$

where  $m_i$  and  $\Delta N_{ice, i}$  are the mass and number concentration of the ice crystals in the  $i$ th size bin, respectively. The mass  $m$  is computed using the mass - dimension ( $m - D$ ) relation of the form  $m = a \cdot D^b$  described in Krämer et al. (2016), which is based on the modified  $m - D$  relation of Mitchell et al. (2010). The coefficients  $a$  and  $b$  depend on the ice crystal size. The used



$m - D$  relation was compared in Afchine et al. (2018) with other  $m - D$  relations from the literature showing good agreement for cirrus clouds.

### 125 3 Normalization method

The method used to fit the PSDs is based on the methodology described by D05 and D14, which is an adaptation of the normalization method for raindrop spectra introduced by Testud et al. (2001). It consists of computing several moments of the measured in situ PSDs, use them to normalize the in situ PSDs and then fit the normalized PSDs to a certain function. This universal function can then be used to obtain microphysical and optical properties of cirrus clouds. Here, we present the key points of the method, for a complete description, the reader is referred to the aforementioned references.

The first step is to compute the equivalent melted diameter ( $D_{eq}$ ) following:

$$D_{eq} = \left( \frac{6 \cdot m(D)}{\pi \rho_w} \right)^{1/3} [m], \quad (2)$$

where  $m(D)$  is obtained with the  $m - D$  relation indicated in Sect. 2.2.

Using the measured PSD, the moments of the distribution are computed as follows:

$$135 \quad M_n = \int_0^{\infty} D^n N_{ice}(D) dD \approx \sum_{D=min}^{D=max} D^n N_{ice}(D) \Delta D, \quad (3)$$

with  $n$  the moment order,  $N_{ice}(D)$  the ice number concentration for the size bin  $D$  and  $\Delta D$  the width of the corresponding size bin.

D05 and D14 use the volume-weighted diameter  $D_m$  to scale the PSD in the size space and the intercept parameter  $N_0^*$  in the number concentration space. These parameters can be defined in terms of the third and fourth moment of the PSD by:

$$140 \quad D_m = \frac{\int_0^{\infty} N_{ice}(D_{eq}) D_{eq}^4 dD_{eq}}{\int_0^{\infty} N_{ice}(D_{eq}) D_{eq}^3 dD_{eq}}, \quad \text{and} \quad (4)$$

$$N_{ice}(D_{eq}) = N_0^* F(D_{eq}/D_m), \quad (5)$$

considering that  $N_{ice}(D_{eq}) dD_{eq} = N_{ice}(D) dD$  for a given size bin.  $F$  is a modified gamma function that describes the shape of the normalized PSDs and, as first given by D05 is:

$$F(\alpha, \beta, X) = \beta \frac{\Gamma(4) \Gamma(\frac{\alpha+5}{\beta})^{4+\alpha}}{4^4 \Gamma(\frac{\alpha+4}{\beta})^{5+\alpha}} X^\alpha \exp \left[ - \left( X \frac{\Gamma(\frac{\alpha+5}{\beta})}{\Gamma(\frac{\alpha+4}{\beta})} \right)^\beta \right] \quad (6)$$

145  $F$  is defined by four parameters.  $N_0^*$  and  $D_m$  (through  $X = D_{eq}/D_m$ ) that change for each PSD and  $\alpha, \beta$  that are fixed and can be found by computing a cost function. The normalization is applied to each individual PSD for all campaigns.



The  $\alpha$  and  $\beta$  that best fit the normalized measured PSD are chosen using an in situ database (in our study the JULIA dataset) and a least square regression linear fit on moments of the PSD (Field et al., 2005, 2007; Delanoë et al., 2014). Following D14, we use a combination of a low and a high moment of the PSD. However, unlike previous studied studies, we here aim at improving the direct prediction of physical parameters of the PSDs and therefore minimize  $N_{ice}$  and IWC via the cost function. This cost function,  $J$ , is commonly used to quantify the consistency between predicted and in-situ parameters. Considering tendency of both IWC and  $N_{ice}$  to follow a log-normal distribution, we used the logarithmic values of these two parameters when computing  $J$ :

$$J = \sum_{i=1}^n (J_{N_{ice}} + J_{IWC}) \quad (7a)$$

with  $J_{N_{ice}}$  and  $J_{IWC}$  as:

$$J_{N_{ice}} = \left( 1 - \frac{\log(N_{param}(\alpha, \beta))}{\log(N_{insitu})} \right)^2 \quad (7b)$$

$$J_{IWC} = \left( 1 - \frac{\log(IWC_{param}(\alpha, \beta))}{\log(IWC_{insitu})} \right)^2 \quad (7c)$$

where  $n$  is the total number of PSDs,  $N_{insitu}$  is the sum of the ice crystal number concentration from the in-situ database in each size bin and  $IWC_{insitu}$  is derived using Eq. 1.  $N_{param}$  is computed by integration of the size distribution  $N_{ice}(\alpha, \beta, D)$  obtained from the normalized function  $F$  and using only the size bins that are present in the in situ data.  $IWC_{param}$  is derived by applying Eq. 1. The objective of the cost function approach is to find the optimal coefficient pair of  $\alpha$  and  $\beta$  that will minimize  $J$ .

## 4 Results

### 4.1 Original and normalized PSDs

The analyzed in-situ data is limited to temperatures lower than 255 K. This choice was made to minimize the selection of water droplets in mixed-phase clouds, since below this temperature most of the water droplets are glaciated. No differentiation between contrail and natural cirrus was made in our analysis and, in total, 542 719 PSDs were analyzed ( $\approx 151$  h). Of the total number of analyzed PSDs, about 9.8 % are found in temperatures between 235 K and 255 K (i.e. mixed-phase regime). The individual contribution of each campaign is registered in Table 2.

Figure 2 (right column) shows how PSDs contain information about the characteristics of clouds and their history and may differ according to dynamical conditions. These differences can be seen in the frequency distributions of the concentrations of the PSDs of each campaign. For example, very high number concentrations ( $> 1 \text{ cm}^{-3}$ ) of the smallest ice crystals can be due to the presence of contrails (e.g. during COALESC, ML-CIRRUS or CIRRUS-HL). Another example, during several flights of




**Table 2.** Contribution of each campaign considering only  $T < 255$  K. Total number of PSDs: 542 719. Total number of cumulative hours: 150.76.

Campaign	Hours	Num. PSDs	%
ACRIDICON	11.62	41833	7.71
ATTREX	30.45	109605	20.19
CIRRUS-HL	29.38	105410	19.42
COALESC	11.2	40325	7.43
CONTRAST	25.34	91220	16.81
ML-CIRRUS	20.54	73934	13.62
POSIDON	11.72	42178	7.77
RACEPAC	0	15	0.002
START	4.02	14473	2.67
STRATOCLIM	6.03	21697	4
VERDI	0.56	2029	0.37

ACRIDICON-CHUVA and CIRRUS-HL strong convection was present (i.e. fast updrafts). During fast updrafts, air masses can be lifted and reach temperatures lower than 235 K and if the conditions are favourable, ice nucleation will take place and result in the formation of small ice crystals. Also during strong updrafts small supercooled liquid water droplets glaciate, giving as a result a large number of small ice crystals. In the PSDs that is translated in an increase in number concentration between  $1 \times 10^{-2} - 1 \text{ cm}^{-3}$  for ice crystal sizes smaller than  $\approx 20 \mu\text{m}$ . Another indicator of cirrus that have their origin at  $T < 235$  K is the presence of large ice crystals that come from heterogeneous drop freezing ( $\approx 200 \mu\text{m}$ , more noticeable for particles  $> 500 \mu\text{m}$  that reach a number concentration of  $\approx 1 \times 10^{-4} \text{ cm}^{-3}$ ). A more detailed analysis of the characteristics of the PSDs according to dynamical conditions will be given in Krämer et al. (2023).

Figure 2 (left column) shows the results of applying the normalization method described in Sect. 3 for a selection of campaigns (see figure legend for detail). As described in D05 and D14, the normalization approach removes some natural variability of PSDs and therefore narrows the data down to smaller size and concentration areas (mostly around  $D_{eq}/D_m = 1$ ), in comparison to the original observed PSD. Another visible feature of the normalization is that the normalized spectra look similar for all campaigns, although there are some differences depending on the measurement location and the cloud processes involved (Delanoë et al., 2005; Field et al., 2007). Overall, these features are also visible in our study, including as well the larger variability of the PSDs tail, which is linked to the temperature. In D14 it is explained that for very cold temperatures (less than  $-60^\circ\text{C}$ ) the tail vanishes and for warmer temperatures the variability of  $D_m$  is larger, since the range of possible ice crystal sizes increases due to ice particles coming from different microphysical processes.





**Figure 2.** The right column shows size-resolved occurrence frequencies of ice crystals of the observed PSDs and the left column their corresponding normalized PSDs for a selection of campaigns covering the tropics (ACRIDICON-CHUVA, STRATOCLIM), mid-latitudes (COALESC, ML-CIRRUS, CIRRUS-HL) and polar region (CIRRUS-HL); the bottom panels show the combination of all campaigns. In the panels of the left column, the black dashed and solid lines corresponds to the proposed parameterization by D05 and D14, respectively.



## 4.2 Investigation of optimal PSD parameters


As described in Sect. 3, a cost function minimization approach is used in this study to find an optimal pair of  $\alpha$  and  $\beta$  coefficients  
195 needed to defined the normalized PSD parameterizations that will fit in-situ measurements from JULIA. We seek to optimize  
the representation of physical properties, IWC and  $N_{ice}$  (see Eq. 7), as opposed to optical properties as for instance in D05 and  
D14. This is done with the intent to produce a PSD parameterization that can be universally used for retrievals of  $N_{ice}$  and IWC  
using a wide variety of instruments.

In D05 and D14, whose parameterizations are here used for comparison purposes, the lidar extinction coefficient and radar  
200 reflectivity factor are used to minimize the cost function. The selected  $\alpha$  and  $\beta$  pairs in that study yield very similar normalized  
PSD shapes for each campaign, with almost a complete overlap of the normalized function for  $D_m/D_{eq}$  between 0.5 and  
1.5 (see Fig. 9 of D14). Following Fig. 9 of D14, we also plotted the normalized function for each campaign (i.e.  $N_{ice}/N_0^*$   
vs  $D_m/D_{eq}$ ) (Fig. 3). Figure. 3a) shows the normalized function for each campaign when selecting the optimal parameters  
using in the cost function both the  $\log(IWC)$  and the  $\log(N_{ice})$ . Figure 3b shows the result of only using  $\log(IWC)$  for the cost  
205 function and Fig. 3c the result of only using  $\log(N_{ice})$ . Figure 3d is an overview of the selected optimal parameters  $\alpha$  and  $\beta$   
for each campaign (in different colors) and for all together (black circle, JULIA 1M in Table 3). In our study, although the  
normalized functions also cluster around  $D_m/D_{eq}$  values between 0.5 and 1.5, the overlap is not as pronounced as in D14  
and the spread in the  $\alpha$  and  $\beta$  values is also larger (Fig. 3d). To try to obtain a more compact cluster of selected coefficient  
pairs, we divided the PSDs into ice crystals smaller and larger than  $50\mu m$  (not shown) (JULIA small and JULIA large in  
210 Table 3). Splitting the PSDs modifies the spread, but our cluster in any case looks as compact as in D14. This might be due  
to the selected parameters to compute  $J$ . Whereas D14 uses optical parameters, i.e., visible extinction and reflectivity, we use  
physical parameters, i.e.,  $N_{ice}$  (which is very sensitive to temperature) and IWC. Moreover, although we follow the indication  
of applying one moment sensitive to the small particles and another one to the large particles, the selected moments are not  
the same. In D14 the parameters are proportional to the second and approximately sixth moment of the distribution and in our  
215 study to the zeroth moment and between the second and third moment. Table 3 gives an overview of the selected coefficients  
for each parameterization and in Sect. 4.3 we discuss the characteristics of each them.



**Table 3.** Best  $\alpha$  and  $\beta$  coefficients for each parameterization. D05 is from Delanoë et al. (2005) and D14 from Delanoë et al. (2014). D14 was obtained by Delanoë et al. (2014) using extinction and reflectivity in the cost function. All other parameterizations are based on the JULIA database and the use of Eq.7. JULIA 1M refers to one mode, JULIA large to  $D \geq 50 \mu\text{m}$ , JULIA small to  $D < 50 \mu\text{m}$  and JULIA 2M to the combination of the small and large modes.

Parameterization	$\alpha$	$\beta$
D05	-1	3
D14	-0.262	1.754
JULIA 1M	-0.945	0.886
JULIA large	0.968	3.307
JULIA small	-0.968	5.225
JULIA 2M	JULIA small + JULIA large	



**Figure 3.** (a-c) normalized functions for each campaign after selecting the best pair of coefficients. (a) is the result of computing the cost function  $J$  using IWC and  $N_{ice}$ , (b) of using only IWC to compute  $J$  and (c) of using only  $N_{ice}$ . (d) shows the selection of best coefficients pair for each campaign and the coefficients for the parameterization by D05 (black upwards triangle), the parameterization proposed in D14 (black downwards triangle) and the new one (left triangle). For this case, the horizontal line symbol in (d) is coincidental with the symbol of the crosses. JULIA 1M refers to the monomodal parameterization obtained in this study. A detailed overview of the coefficients of each parameterization is given in Table 3. Only PSDs measured at  $T < 255$  K are considered.



### 4.3 One mode vs. two modes parameterization

Ice crystals smaller than about  $50\ \mu\text{m}$  present a greater variability in the PSDs as the larger ice crystals (see Fig. 2, right column), because they correspond to the regime of newly formed ice crystals that quickly grow to larger, sedimenting sizes. In our study we compare the results of using one or two size modes, i.e., a parameterization for small particles ( $D < 50\ \mu\text{m}$ ) combined with another parameterization for large particles ( $D \geq 50\ \mu\text{m}$ ).

Figure 4 shows a comparison between the median of the differences between the single PSDs (observed PSD from JULIA - parameterized PSD) for different parameterizations. In the upper row, panel a) shows the comparison between the parameterizations JULIA 1M and JULIA 2M proposed in this study. Panel b) shows the parameterizations from the literature and the JULIA 2M. The bottom row (c-d) is the same as the upper row but adding in shaded color the corresponding  $1\sigma$  of each parameterization to illustrate their variability. As expected, the larger variability is associated with the smaller size bins. From panels (a) and (c) it is clear, that the bimodal parameterization (JULIA 2M: black) is closer to the observations than the proposed mono-modal (JULIA 1M: yellow) and that the variability is reduced for the bimodal parameterization. Between the bimodal parameterization and the ones from the literature (panels (b) and (d)), the clearer improvement is for ice crystals between  $20$  and  $\approx 110\ \mu\text{m}$ . For the small particles (smaller than  $20\ \mu\text{m}$ ), D14 (dark blue) presents the highest deviation. This parameterization underestimate the number concentration of particles in this size range. In the case of D05 (red), the number concentration is also underestimated, but in less extent. The bimodal JULIA 2M (black) is the only parameterization that slightly overestimates the number concentration of the smaller particles. Panel (d) shows that the main improvement of the bimodal parameterization regarding sizes smaller than  $20\ \mu\text{m}$  is the reduced variability. A detailed analysis by temperature of the comparison from Fig. 4b, presented in Fig. 5, shows that the coldest temperatures ( $-90^\circ\text{C}$  and  $-60^\circ\text{C}$ ) have the lowest deviation between observations and parameterizations for the smaller particles and that all parameterizations deliver similar results, especially for the larger particles. For temperatures between  $-60$  and  $-50^\circ\text{C}$ , where bimodality starts playing an important role, the parameterization from D05 (red) adjust slightly better to the smaller size bins but overestimates the number concentration for particles between  $20$  and  $100\ \mu\text{m}$  ( $\text{PSD}_{\text{obs}} - \text{PSD}_{\text{param}} < 0$ ). This overestimation is also observed between  $-50^\circ\text{C}$  and  $-20^\circ\text{C}$ , but for this range the bimodal parameterization is closer to the measured number concentrations of small particles. Moreover, the bimodal parameterization is closer to the observations for particles between  $20$  and  $110\ \mu\text{m}$  for temperatures between  $-60^\circ\text{C}$  and  $-20^\circ\text{C}$ .


In terms of IWC (not shown), correlation plots show that there are no significant differences between the results of the parameterizations (all parameterizations have a correlation factor of 1.0 and a RMSE of 0.18 or 0.19), but show a slight underestimation of the IWC for values between about  $\log_{10} = -5$  and  $-7$  and an overestimation between about  $\log_{10} = -3$  and 0. Since all parameterizations have a similar behaviour for the large particles and IWC is sensitive to large particles, this result was expected.

In terms of  $N_{\text{ice}}$ , correlation plots between the measurements and parameterizations (Fig. 6) show that the JULIA 2M parameterization (lower row) significantly reduces the spread seen in the monomodal D05 parametrization (upper row) and that the highest frequency for D05 is found slightly above the one-to-one line. Considering temperatures lower than  $255\ \text{K}$ , the




root mean square error (RMSE) is 0.34 for D05 and 0.21 for the JULIA 2M parameterization (panels (c) and (f), respectively). The bimodal parameterization presents a slightly higher correlation factor (0.98 vs. 0.96) than D05. Subsetting these analyses by 10-K temperature bins between  $-90^{\circ}\text{C}$  and  $-60^{\circ}\text{C}$  (Fig. 6a, b, d, e), where there is only one mode, JULIA 2M shows the lowest RMSE, with 0.22 vs 0.30 of the monomodal D05. For temperatures between  $-60^{\circ}\text{C}$  and  $-20^{\circ}\text{C}$ , where two modes are clearly visible, D05 has a RMSE of 0.36 vs. a RMSE of 0.20 of the two-modes parameterization. A comparison in temperature ranges of  $10^{\circ}\text{C}$  (not shown), gives the same conclusions as already discussed, similar results for the coldest temperatures, being the two modes parameterization slightly closer to the observations, and better results of the two-modes parameterization for the warmer temperatures.

These analysis confirm what was hinted by Sourdeval et al. (2018), i.e., defining a two-modes parameterization instead of one-mode improves the reconstruction of PSDs and retrieval of  $N_{\text{ice}}$  for warmer temperatures because it adjusts better to the bimodal shape of the PSDs.




**Figure 4.** Median of the difference between the single parameterized PSDs and their corresponding observed PSDs. (a, c) Comparison between one mode (JULIA 1M: yellow) and two mode parameterization (JULIA 2M: black); (b, d) comparison between parameterizations from the literature (D05: red, D14: blue) and the two-modes parameterization (JULIA 2M: black) of this study. The shadow region in the lower panels correspond to  $1\sigma$ . y-axis is the median of the difference in number concentration between the observed (from JULIA database) and the parameterized PSDs in  $\text{cm}^{-3}$  and the x-axis represent the size bins in  $\mu\text{m}$ . A detailed overview of the coefficients of each parameterization is given in Table 3.



**Figure 5.** Median of the difference between the single observed PSDs of all campaigns and their corresponding parametrized PSDs, as in Fig. 4c, but for 10K temperature ranges. The parameterizations from the literature are: D05 in red and D14 in blue. JULIA 2M (in black) is the proposed new bimodal parameterization. y-axis is the median of the difference in number concentration between the observed and the parametrized PSDs in  $\text{cm}^{-3}$  and the x-axis is the size bins in  $\mu\text{m}$ . A detailed overview of the coefficients of each parameterization is given in Table 3.





**Figure 6.** Comparison between the in-situ  $N_{ice}$  and parameterized  $N_{ice}$  for temperatures between  $-90^{\circ}\text{C}$  and  $-60^{\circ}\text{C}$  (a) D05, (d) JULIA 2M), temperatures between  $-60^{\circ}\text{C}$  and  $-20^{\circ}\text{C}$  (b) D05, (e) JULIA 2M) and temperatures lower than 255 K (c) D05, (f) JULIA 2M). The y-axis is the logarithm of the parameterized variable and the x-axis the logarithm of the observed one. Color code indicates frequency (%). The black line corresponds to the 1 to 1 line, the solid red line to the regression line and the dashed red lines to the regression line shifted a factor of  $\pm 2$ . The black ellipse is the  $2\sigma$  area. A detailed overview of the coefficients of each parameterization is given in Table 3.



## 5 Conclusions

In the present study, recent airborne in-situ measurements of ice crystals PSDs from the JULIA database were used to assess the ability of existing PSD parameterizations to represent  $N_{\text{ice}}$  and IWC, and to investigate the added-value of considering two-mode PSDs. One of the main advantages of JULIA with respect to other dataset used for constructing current PSD parameterizations is that it includes observations of ice crystals down to  $3\ \mu\text{m}$ , which are consistently processed for all field campaigns in particular to minimize the impact of ice shattering effects. To find a new parameterization, we have followed the method by Delanoë et al. (2014), which consists of normalizing the PSDs and fitting them to a modified gamma function whose coefficients are chosen by minimizing a cost function  $J$ . The variables chosen to define the cost function were IWC and  $N_{\text{ice}}$ . We found that considering a possible bimodality of PSDs by combining two parameterizations, one for particles with  $D < 50\ \mu\text{m}$  and one for  $D \geq 50\ \mu\text{m}$ , yields better results when compared to the observations than the hitherto used monomodal parameterizations. The variability of the retrieved around the observed PSDs is reduced across all analyzed temperatures and there is a better fit, especially for ice particles between 20 and  $\approx 110\ \mu\text{m}$  and for temperatures between  $-60$  and  $-20^\circ\text{C}$ . For this temperature range, the RMSE for the retrieved  $N_{\text{ice}}$  is reduced from 0.36 to 0.20. In conclusion, we propose here a new bimodal ice particle PSD parameterization including ice crystals smaller than  $50\ \mu\text{m}$ . An important next step would be to test the feasibility of implementing two parameterizations, one for smaller particles and another for larger particles, in the retrieval algorithms of remote sensing instruments.

*Code and data availability.* Please, contact the contact author

*Author contributions.* OS designed the study, IB performed the analyses, MK provided the insitu database. All co-authors contributed to the discussion of the results, improvement of the analysis and the revision of the manuscript.

*Competing interests.* The authors declare that there are no competing interests.

*Acknowledgements.* The authors thank the PIRE research initiative and the Procope-Mobilität-2021 program that made possible the progress of this study. Odran Sourdeval acknowledges support by CNES, focused on EarthCare and IASI.



## 285 References

- Afchine, A., Rolf, C., Costa, A., Spelten, N., Riese, M., Buchholz, B., Ebert, V., Heller, R., Kaufmann, S., Minikin, A., Voigt, C., Zöger, M., Smith, J., Lawson, P., Lykov, A., Khaykin, S., and Krämer, M.: Ice particle sampling from aircraft – influence of the probing position on the ice water content, *Atmospheric Measurement Techniques*, 11, 4015–4031, <https://doi.org/10.5194/amt-11-4015-2018>, 2018.
- Baran, A. J.: A Review of the Light Scattering Properties of Cirrus, *J. Quant. Spectrosc. Radiat. Transfer*, 110, 1239–1260, <https://doi.org/10.1016/j.jqsrt.2009.02.026>, 2009.
- 290 Baumgardner, D., Abel, S., Axisa, D., Cotton, R., Crosier, J., Field, P., Gurganus, C., Heymsfield, A., Korolev, A., Krämer, M., Lawson, P., Mefarqhar, G., Ulanowski, Z., and Um, J.: Cloud Ice Properties - In Situ Measurement Challenges, *Meteorological Monographs*, 58, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0011.1>, 2017.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change, *Reviews of Geophysics*, 58, e2019RG000660, <https://doi.org/10.1029/2019RG000660>, 2020.
- 315 CIRRUS-HL, last accessed: 29 July 2022: CIRRUS-HL: project description, <https://halo-research.de/science/previous-missions/cirrus-hl/>, 2021.
- Costa, A.: Mixed-phase and ice cloud observations with NIXE-CAPS, Dissertation, Universität Wuppertal, Jülich, <https://juser.fz-juelich.de/record/829847>, universität Wuppertal, Diss., 2017, 2017.
- Costa, A., Meyer, J., Afchine, A., Luebke, A., Günther, G., Dorsey, J. R., Gallagher, M. W., Ehrlich, A., Wendisch, M., Baumgardner, D., Wex, H., and Krämer, M.: Classification of Arctic, midlatitude and tropical clouds in the mixed-phase temperature regime, *Atmospheric Chemistry and Physics*, 17, 12 219–12 238, <https://doi.org/10.5194/acp-17-12219-2017>, 2017.
- 330 Delanoë, J. and Hogan, R. J.: A variational scheme for retrieving ice cloud properties from combined radar, lidar, and infrared radiometer, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/https://doi.org/10.1029/2007JD009000>, 2008.
- Delanoë, J. and Hogan, R. J.: Combined CloudSat-CALIPSO-MODIS retrievals of the properties of ice clouds, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/https://doi.org/10.1029/2009JD012346>, 2010.
- 345 Delanoë, J., Protat, A., Testud, J., Bouniol, D., Heymsfield, A. J., Bansemmer, A., Brown, P. R. A., and Forbes, R. M.: Statistical properties of the normalized ice particle size distribution, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/https://doi.org/10.1029/2004JD005405>, 2005.
- Delanoë, J. M. E., Heymsfield, A. J., Protat, A., Bansemmer, A., and Hogan, R. J.: Normalized particle size distribution for remote sensing application, *Journal of Geophysical Research: Atmospheres*, 119, 4204–4227, <https://doi.org/https://doi.org/10.1002/2013JD020700>, 315 2014.
- Field, P. R., Hogan, R. J., Brown, P. R. A., Illingworth, A. J., Choullarton, T. W., and Cotton, R. J.: Parametrization of ice-particle size distributions for mid-latitude stratiform cloud, *Quarterly Journal of the Royal Meteorological Society*, 131, 1997–2017, <https://doi.org/https://doi.org/10.1256/qj.04.134>, 2005.
- Field, P. R., Heymsfield, A. J., and Bansemmer, A.: Snow Size Distribution Parameterization for Midlatitude and Tropical Ice Clouds, *Journal of the Atmospheric Sciences*, 64, 4346 – 4365, <https://doi.org/10.1175/2007JAS2344.1>, 2007.
- 320



- Gasparini, B. and Lohmann, U.: Why Cirrus Cloud Seeding Cannot Substantially Cool the Planet, *J. Geophys. Res. Atmos.*, 121, 4877–4893, <https://doi.org/10.1002/2015JD024666>, 2016.
- Heymsfield, A. J. and McFarquhar, G.: Mid-latitude and tropical cirrus microphysical properties, in: *Cirrus*, edited by Lynch, D., chap. 4, pp. 78–101, Oxford University Press, 2002.
- 325 Jackson, R. C., McFarquhar, G. M., Fridlind, A. M., and Atlas, R.: The dependence of cirrus gamma size distributions expressed as volumes in  $N_0$ - $\lambda$ - $\mu$  phase space and bulk cloud properties on environmental conditions: Results from the Small Ice Particles in Cirrus Experiment (SPARTICUS), *Journal of Geophysical Research: Atmospheres*, 120, 10–351, 2015.
- Khaykin, S. M., Moyer, E., Krämer, M., Clouser, B., Bucci, S., Legras, B., Lykov, A., Afchine, A., Cairo, F., Formanyuk, I., Mitev, V., Matthey, R., Rolf, C., Singer, C. E., Spelten, N., Volkov, V., Yushkov, V., and Stroh, F.: Persistence of moist plumes from  
330 overshooting convection in the Asian monsoon anticyclone, *Atmospheric Chemistry and Physics (highlight article)*, 22, 3169–3189, <https://doi.org/10.5194/acp-22-3169-2022>, 2022.
- Klingebiel, M., de Lozar, A., Molleker, S., Weigel, R., Roth, A., Schmidt, L., Meyer, J., Ehrlich, A., Neuber, R., Wendisch, M., and Borrmann, S.: Arctic low-level boundary layer clouds: in situ measurements and simulations of mono- and bimodal supercooled droplet size distributions at the top layer of liquid phase clouds, *Atmospheric Chemistry and Physics*, 15, 617–631, <https://doi.org/10.5194/acp-15-617-2015>,  
335 2015.
- Korolev, A., Emery, E., Strapp, J., Cober, S., Isaac, G., Wasey, M., and Marcotte, D.: Small Ice Particles in Tropospheric Clouds: Fact or Artifact? Airborne Icing Instrumentation Evaluation Experiment, *Bulletin of the American Meteorological Society*, 92, 967–973, <https://doi.org/10.1175/2010BAMS3141.1>, 2011.
- Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zöger, M., Smith, J., Herman, R. L., Buchholz, B., Ebert,  
340 V., Baumgardner, D., Borrmann, S., Klingebiel, M., and Avallone, L.: A microphysics guide to cirrus clouds – Part 1: Cirrus types, *Atmospheric Chemistry and Physics*, 16, 3463–3483, <https://doi.org/10.5194/acp-16-3463-2016>, 2016.
- Krämer, M., Rolf, C., Spelten, N., and Riese, M.: The JÜLich In-situ Airborne Data Base JULIA: Water Vapor, Clouds & other Trace Substances in the UT/LS, in: *AGU Fall Meeting Abstracts*, vol. 2018, pp. A510–2404, 2018.
- Krämer, M., Rolf, C., Spelten, N., Afchine, A., Fahey, D., Jensen, E., Khaykin, S., Kuhn, T., Lawson, P., Lykov, A., Pan, L. L., Riese,  
345 M., Rollins, A., Stroh, F., Thornberry, T., Wolf, V., Woods, S., Spichtinger, P., Quaas, J., and Sourdeval, O.: A microphysics guide to cirrus – Part 2: Climatologies of clouds and humidity from observations, *Atmospheric Chemistry and Physics (highlight article)*, 20, 12 569–12 608, <https://doi.org/10.5194/acp-20-12569-2020>, 2020.
- Krämer, M., Spelten, N., Afchine, A., and Spang, R.: Occurrence patterns of cloud particles sizes in cirrus and mixed-phase clouds, *EGU General Assembly 2022, Vienna, Austria, 23-27 May 2022*, EGU22-5119, <https://doi.org/https://doi.org/10.5194/egusphere-egu22-5119>,  
350 2022.
- Krämer, M., Rolf, C., Spang, R., Afchine, A., and Spelten, N.: A Microphysics Guide to Cirrus – Part III: Occurrence patterns of ice cloud particles, *Atmospheric Chemistry and Physics*, in preparation, p. xx, xx, 2023.
- Lawson, R. P.: Effects of ice particles shattering on the 2D-S probe, *Atmospheric Measurement Techniques*, 4, 1361–1381, <https://doi.org/10.5194/amt-4-1361-2011>, 2011.
- 355 Lawson, R. P., O’Connor, D., Zmarzly, P., Weaver, K., Baker, B., Mo, Q., and Jonsson, H.: The 2D-S (STEREO) PROBE: Design and preliminary tests of a new airborne, high-speed, high-resolution particle imaging probe, *Journal of Atmospheric and Oceanic Technology*, 23, 1462–1477, <https://doi.org/10.1175/jtech1927.1>, 2006.



- Liou, K.-N.: Influence of Cirrus Clouds on Weather and Climate Processes: A Global Perspective, *Monthly Weather Review*, 114, 1167 – 1199, [https://doi.org/10.1175/1520-0493\(1986\)114<1167:IOCCOW>2.0.CO;2](https://doi.org/10.1175/1520-0493(1986)114<1167:IOCCOW>2.0.CO;2), 1986.
- 360 Lohmann, U., Quaas, J., Kinne, S., and Feichter, J.: Different Approaches for Constraining Global Climate Models of the Anthropogenic Indirect Aerosol Effect, *Bulletin of the American Meteorological Society*, 88, 243 – 250, <https://doi.org/10.1175/BAMS-88-2-243>, 2007.
- Luebke, A., Avallone, L., Schiller, C., Meyer, J., Rolf, C., and Krämer, M.: Ice water content of Arctic, midlatitude, and tropical cirrus – Part 2: Extension of the database and new statistical analysis, *ACP*, 13, 6447–6459, 2013.
- Luebke, A. E., Afchine, A., Costa, A., Groß, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.:  
365 The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, *Atmospheric Chemistry and Physics*, 16, 5793–5809, <https://doi.org/10.5194/acp-16-5793-2016>, 2016.
- McFarquhar, G. M., Um, J., Freer, M., Baumgardner, D., Kok, G. L., and Mace, G.: Importance of small ice crystals to cirrus properties: Observations from the Tropical Warm Pool International Cloud Experiment (TWP-ICE), *Geophysical Research Letters*, 34, <https://doi.org/https://doi.org/10.1029/2007GL029865>, 2007.
- 370 Meyer, J.: Ice Crystal Measurements with the New Particle Spectrometer NIXE-CAPS, Dr. (univ.), Universität Wuppertal, Jülich, <https://juser.fz-juelich.de/record/22871>, record converted from VDB: 12.11.2012; Universität Wuppertal, Diss., 2012, 2013.
- Mitchell, D. L., d’Entremont, R. P., and Lawson, R. P.: Inferring Cirrus Size Distributions through Satellite Remote Sensing and Microphysical Databases, *Journal of the Atmospheric Sciences*, 67, 1106 – 1125, <https://doi.org/10.1175/2009JAS3150.1>, 2010.
- Osborne, S., Abel, S., Boutle, I., and Marengo, F.: Evolution of Stratocumulus Over Land: Comparison of Ground and Aircraft Observations  
375 with Numerical Weather Prediction Simulations, *Boundary-Layer Meteorology*, 153, 165–193, <https://doi.org/10.1007/s10546-014-9944-0>, 2014.
- Pan, L. L., Bowman, K. P., Atlas, E. L., Wofsy, S. C., Zhang, F., Bresch, J. F., Ridley, B. A., Pittman, J. V., Homeyer, C. R., Romashkin, P., and Cooper, W. A.: The Stratosphere-Troposphere Analyses of Regional Transport 2008 (START08) Experiment, *Bulletin of the American Meteorological Society*, 91, 327–342, <https://doi.org/10.1175/2009BAMS2865.1>, 2010.
- 380 Pan, L. L., Atlas, E. L., Salawitch, R. J., Honomichl, S. B., Bresch, J. F., Randel, W. J., Apel, E. C., Hornbrook, R. S., Weinheimer, A. J., Anderson, D. C., Andrews, S. J., Baidar, S., Beaton, S. P., Campos, T. L., Carpenter, L. J., Chen, D., Dix, B., Donets, V., Hall, S. R., Hanisco, T. F., Homeyer, C. R., Huey, L. G., Jensen, J. B., Kaser, L., Kinnison, D. E., Koenig, T. K., Lamarque, J.-F., Liu, C., Luo, J., Luo, Z. J., Montzka, D. D., Nicely, J. M., Pierce, R. B., Riemer, D. D., Robinson, T., Romashkin, P., Saiz-Lopez, A., Schauffler, S., Shieh, O., Stell, M. H., Ullmann, K., Vaughan, G., Volkamer, R., and Wolfe, G.: Convective Transport of Active Species in the Tropics  
385 (CONTRAST) experiment, *Bull. Amer. Meteor. Soc.*, 98, 106–128, <https://doi.org/10.1175/BAMS-D-14-00272.1>, 2017.
- POSIDON: last accessed: 29 July 2022: POSIDON: project description, [https://espo.nasa.gov/posidon/content/POSIDON\\_0](https://espo.nasa.gov/posidon/content/POSIDON_0), 2016.
- Schiller, C., Krämer, M., Afchine, A., Spelten, N., and Sitnikov, N.: Ice water content in Arctic, midlatitude and tropical cirrus, *J. Geophys. Res.*, 113, D24208, <https://doi.org/10.1029/2008JD010342>, 2008.
- Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 1: Model description,  
390 *Meteor. Atmos. Phys.*, 92, 45–66, <https://doi.org/10.1007/s00703-005-0112-4>, 2006.
- Sourdeval, O., C. Labonnote, L., Baran, A. J., and Brogniez, G.: A methodology for simultaneous retrieval of ice and liquid water cloud properties. Part I: Information content and case study, *Quart. J. Roy. Meteor. Soc.*, 141, 870–882, <https://doi.org/10.1002/qj.2405>, 2015.
- Sourdeval, O., C. Labonnote, L., Baran, A. J., Mülmenstädt, J., and Brogniez, G.: A methodology for simultaneous retrieval of ice and liquid water cloud properties. Part 2: Near-global retrievals and evaluation against A-Train products, *Quart. J. Roy. Meteor. Soc.*, 142,  
395 3063–3081, <https://doi.org/10.1002/qj.2889>, 2016.



- Sourdeval, O., Gryspeerdt, E., Krämer, M., Goren, T., Delanoë, J., Afchine, A., Hemmer, F., and Quaas, J.: Ice crystal number concentration estimates from lidar–radar satellite remote sensing – Part 1: Method and evaluation, *Atmospheric Chemistry and Physics*, 18, 14 327–14 350, <https://doi.org/10.5194/acp-18-14327-2018>, 2018.
- Stephens, G. L., Tsay, S.-C., Stackhouse, Paul W., J., and Flatau, P. J.: The Relevance of the Microphysical and Radiative Properties of Cirrus Clouds to Climate and Climatic Feedback., *Journal of Atmospheric Sciences*, 47, 1742–1754, [https://doi.org/10.1175/1520-0469\(1990\)047<1742:TROTMA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<1742:TROTMA>2.0.CO;2), 1990.
- Testud, J., Oury, S., Black, R. A., Amayenc, P., and Dou, X.: The Concept of “Normalized” Distribution to Describe Raindrop Spectra: A Tool for Cloud Physics and Cloud Remote Sensing, *Journal of Applied Meteorology*, 40, 1118 – 1140, [https://doi.org/10.1175/1520-0450\(2001\)040<1118:TCONDNT>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<1118:TCONDNT>2.0.CO;2), 2001.
- 405 Thornberry, T. D., Rollins, A. W., Avery, M. A., Woods, S., Lawson, R. P., Bui, T. V., and Gao, R.-S.: Ice water content-extinction relationships and effective diameter for TTL cirrus derived from in situ measurements during ATTREX 2014, *Journal of Geophysical Research: Atmospheres*, 122, 4494–4507, <https://doi.org/https://doi.org/10.1002/2016JD025948>, 2017.
- Vidot, J., Baran, A. J., and Brunel, P.: A new ice cloud parameterization for infrared radiative transfer simulation of cloudy radiances: Evaluation and optimization with IIR observations and ice cloud profile retrieval products, *J. Geophys. Res. Atmos.*, 120, 6937–6951, <https://doi.org/10.1002/2015JD023462>, 2015.
- Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrmann, S., Boettcher, M., Bucuchholz, B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Doernbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., Fuetterer, D., Giez, A., Graf, K., Grooss, J.-U., Gross, S., Heimerl, K., Heinold, B., Hueneke, T., Jaervinen, E., Jurkat, T., Kaufmann, S., Kenntner, M., Klingebiel, M., Klimach, T., Kohl, R., Krämer, M., Krisna, T. C., Luebke, A., Mayer, B., Mertes, S., Molleker, S., Petzold, A., Pfeilsticker, K., Port, M., Rapp, M., Reutter, P., Rolf, C., Rose, D., Sauer, D., Schaefer, A., Schlage, R., Schnaiter, M., Schneider, J., Spelten, N., Spichtinger, P., Stock, P., Walser, A., Weigel, R., Weinzierl, B., Wendisch, M., Werner, F., Wernli, H., Wirth, M., Zahn, A., Ziereis, H., and Zöger, M.: ML-Cirrus the airborne experiment on natural cirrus and contrail cirrus with the high-altitude long-range research aircraft HALO, *Bulletin of the American Meteorological Society*, 98, 271–288, <https://doi.org/10.1175/BAMS-D-15-00213.1>, 2017.
- Wendisch, M., Pöschl, U., Andreae, M. O., Machado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D., Martin, S. T., Abdelmonem, A., Afchine, A., Araújo, A. C., Artaxo, P., Aufmhoff, H., Barbosa, H. M. J., Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Fütterer, D., Heckl, C., Heidelberg, F., Hüneke, T., Jäkel, E., Järvinen, E., Jurkat, T., Kanter, S., Kästner, U., Kenntner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Kölling, T., Krämer, M., Krüger, M., Krisna, T. C., Lavric, J. V., Longo, K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Münch, S., Nillius, B., Pfeilsticker, K., Pöhlker, C., Roiger, A., Rose, D., Rosenow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., de Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walser, A., Walter, D., Weigel, R., Weinzierl, B., Werner, F., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zöger, M.: ACRIDICON–CHUVA Campaign: Studying Tropical Deep Convective Clouds and Precipitation over Amazonia Using the New German Research Aircraft HALO, *Bulletin of the American Meteorological Society*, 97, 1885 – 1908, <https://doi.org/10.1175/BAMS-D-14-00255.1>, 2016.
- 425 Yang, P., Gao, B.-C., Baum, B. A., Wiscombe, W. J., Hu, Y. X., Nasiri, S. L., Soulen, P. F., Heymsfield, A. J., McFarquhar, G. M., and Miloshevich, L. M.: Sensitivity of cirrus bidirectional reflectance to vertical inhomogeneity of ice crystal habits and size distributions for two Moderate-Resolution Imaging Spectroradiometer (MODIS) bands, *J. Geophys. Res. Atmos.*, 106, 17 267–17 291, <https://doi.org/10.1029/2000JD900618>, 2001.
- 430

<https://doi.org/10.5194/egusphere-2023-754>

Preprint. Discussion started: 22 May 2023

© Author(s) 2023. CC BY 4.0 License.



- Zhang, Y., Macke, A., and Albers, F.: Effect of crystal size spectrum and crystal shape on stratiform cirrus radiative forcing, *Atmospheric Research*, 52, 59–75, [https://doi.org/10.1016/S0169-8095\(99\)00026-5](https://doi.org/10.1016/S0169-8095(99)00026-5), 1999.
- 435 Zhao, Y., Mace, G. G., and Comstock, J. M.: The Occurrence of Particle Size Distribution Bimodality in Midlatitude Cirrus as Inferred from Ground-Based Remote Sensing Data, *Journal of the Atmospheric Sciences*, 68, 1162 – 1177, <https://doi.org/10.1175/2010JAS3354.1>, 2011.