

# Evaluation and optimisation of the soil carbon turnover routine in the MONICA model (version 3.3.1)

Konstantin Aiteew<sup>1</sup>, Jarno Rouhiainen<sup>1</sup>, Claas Nende<sup>2,3</sup>, René Dechow<sup>1</sup>

<sup>1</sup>Thünen Institute of Climate-Smart Agriculture, Bundesallee 68, Braunschweig, Germany

5 <sup>2</sup>Leibniz Centre for Agricultural Landscape Research (ZALF), Eberswalder Straße 84, Müncheberg, Germany

<sup>3</sup>Institute of Biochemistry and Biology, University of Potsdam, Am Mühlenberg 3, Potsdam, Germany

*Correspondence to:* Konstantin Aiteew ([konstantin.aiteew@thuenen.de](mailto:konstantin.aiteew@thuenen.de))

**Abstract.** Simulation models are tools commonly used to predict changes in soil carbon stocks. Prior validation is essential, however, for determining the reliability and applicability of model results. In this study, the process-based biogeochemical model MONICA (Model of Nitrogen and Carbon dynamics on Agro-ecosystems) was evaluated with respect to soil organic carbon (SOC) using long-term monitoring data from 46 German agricultural sites. A revision and parameterisation of equations, encompassing crop and fertiliser-specific C contents and the abiotic factors of soil temperature, soil water and clay content, were undertaken and included in the model. The modified version was also used for a Morris elementary effects screening method, which confirmed the importance of environmental and management factors to the model's performance. The model was then calibrated by means of Bayesian inference using the Metropolis-Hastings algorithm. The performance of the MONICA model was compared with that of five established carbon turnover models (CCB, CENTURY, C-TOOL, ICBM and RothC). The original MONICA model systematically overestimated SOC decomposition rates and produced on average a ~17 % greater mean absolute error (MAE) than the other models. The modification and calibration significantly improved its performance, reducing the MAE by ~30 %. Consequently, MONICA outperformed CENTURY, CCB and C-TOOL, and produced results comparable with ICBM and RothC. Use of the modified model allowed mostly adequate reproduction of site-specific SOC stocks, while the availability of a nitrogen, plant growth and water submodel enhanced its applicability compared with models that only describe carbon dynamics.

## 1 Introduction

The historic conversion from native to agriculturally managed soils released 116 Pg of soil organic carbon (SOC) as CO<sub>2</sub> to the atmosphere (Sanderman et al., 2018). If no measures are taken, current projections estimate a continuation in the decline of SOC stocks, further intensifying the release of CO<sub>2</sub> from soils (Riggers et al., 2021; Zhao et al., 2021). However, the sequestration of atmospheric CO<sub>2</sub> in the soil is a climate change mitigation strategy that has substantial potential (Poepflau and Don, 2015; Fuss et al., 2018; Amelung et al., 2020). In particular, croplands are promising landscapes since they are depleted in SOC and generally accessible for cultivation practices (Poepflau and Don, 2015; Sanderman et al., 2017). Researchers have proposed various agronomic systems, including agroforestry, reduced tilling, and perennial and cover cropping, to optimise

agricultural opportunities for sequestering carbon and thus partly restore the historically lost SOC (Smith et al., 2005). Estimates of the global capacity of agricultural soils for sequestering carbon range from 2 to 5 Pg C a<sup>-1</sup> (Fuss et al., 2018). However, identifying and promoting site-specific measures in a highly diverse agricultural landscape encompassing more than 570 million farms globally and affecting ~36 % of the earth's terrestrial surface is a challenge (FAO, 2014; Amundson and 35 Biardeau, 2018; FAO, 2022). Despite the uncertainty in their practicability, international initiatives such as the '4 per mille' and scientific working groups advise the rapid modification of agricultural systems so that they act as carbon sinks (Minasny et al., 2017; Amelung et al., 2020). In contrast, some scientists focus on the potential for underestimated detrimental effects and advocate a well-thought-out approach to sequestering SOC on a large scale (Sommer and Bossio, 2014; Lugato et al., 2014; Amundson and Biardeau, 2018; Murphy, 2020).

40 Process-based models are tools that can be used to assess mitigation capabilities of agricultural systems for different environmental and management conditions and to analyse disagreements and accuracy of alternative estimation approaches. They are designed to represent the behaviour of natural systems based on a selection of mathematical formulations of generally accepted biophysical principles. Carbon turnover models predict the dynamics of SOC in response to management and climate. Agro-ecosystem models combine carbon and nitrogen turnover routines with algorithms that represent soil water dynamics, 45 greenhouse gas emissions and crop growth and allow the simultaneous prediction of multiple variables from different domains of the soil-plant-atmosphere continuum. Due to the explicit simulation of crop growth in response to weather, site conditions and crop management, agroecosystem models exhibit a clear advantage over simple soil carbon turnover models, as they can translate any changes in conditions that affect biomass input to soil, e.g., climate or crop management, directly to responses in SOC. Potentially, they could also simulate any feedback from those changing SOC dynamics to crop growth, e.g. via nitrogen 50 release or immobilization. The disadvantage of this is that complex models require an accurate reproduction of the carbon input variability, whereas simpler carbon models estimate the carbon inputs from measured yields. Processes that are not described by the model, such as the effects of phosphorus and other nutrients and the effects of plant diseases on plant growth, could further constrain the ability of complex models to simulate adequate carbon inputs. Additionally, complex agro-ecosystem models are inherently black boxes that make it difficult to understand the various factors and feedback loops. They 55 have yet rarely been tested for their ability to do exactly this, just as little as they have been compared to simple soil carbon turnover models to identify whether their advantage in applicability comes with a reduced predictive power (Smith et al., 1997; Farina et al., 2021). In any case, to ensure the adequate predictive capability of such models, complex or simple, a comprehensive validation process using measured data with different soil types, management practices and weather conditions is required.


60 MONICA (Model of Nitrogen and Carbon dynamics on Agro-ecosystems; Nendel et al., 2011; 2013) is a process-based agro-ecosystem model that simulates the effects of spatiotemporal environmental variabilities and agricultural practices on crops yields and soil conditions (Nendel et al., 2014). It is a successor of the HERMES model, which was initially developed to estimate N dynamics in the soil-crop system (Kersebaum, 1995; 2007). A comprehensive SOC turnover submodule based on the Daisy model was added, replacing the simpler approach in the HERMES model (Hansen et al., 1991; Abrahamson and

65 Hansen, 2000). The holistic principle of the MONICA model supports research questions on the interlinkage between site conditions, plant development and C and N dynamics in a changing climate. The performance of the MONICA model showed the capability to adequately reproduce crop yields for different crops (Asseng et al., 2013; Bassu et al., 2014; Kothari et al., 2022; Rötter et al., 2012; Salo et al., 2016) and more complex crop rotations (Kollas et al., 2015; Kostková et al., 2021). For SOC dynamics under bare fallow treatment, MONICA performed similar to simple C turnover models (Farina et al., 2021),  
70 but furthermore demonstrated also good performance when simulating short-term high-resolution CO<sub>2</sub> exchange in a soil-plant system (Specka et al., 2016). The Daisy model, another agro-ecosystem model, had already been tested against long-term soil carbon experiments (Smith et al., 1997) and performed comparably well to soil carbon turnover models, such as RothC (Coleman & Jenkinson, 1996) or CANDY (Franko, 1996).

The question now is whether the MONICA model with its carbon turnover submodule is able to adequately simulate changes  
75 in SOC stocks considering different agronomic practices. It is also important to establish how the performance of a complex model like MONICA compares with simple C turnover models such as RothC, ICBM, C-TOOL, CENTURY and CCB. To answer these questions, we move along the following four objectives: i) to validate the SOC submodule of the MONICA model with data from 46 agricultural long-term monitoring sites in Germany, ii) if necessary to enhance its applicability in view of the diversification of the dataset used for validation, iii) to identify the main drivers of C sequestration and decomposition in  
80 the model, and iv) if necessary to improve model performance.

## 2 Materials and Methods

### 2.1 Data composition and site characteristics



**Figure 1: Location (a) and soil classification (b) of the 46 agricultural long-term monitoring sites (BDF) in Lower Saxony, Germany.**

For calibration and validation of the SOC turnover subroutine of the MONICA model, long-term monitoring data collected by the State Office for Mining, Energy and Geology (LBEG) were used. In 1990, the state government of Lower Saxony (Germany) established a programme to permanently observe agricultural and forest soils in the federal territory. Its goal was to monitor soil changes and identify adverse ecologic and economic implications, and thus predict the risk of imminent damage for the sustainable utilisation of soils.

The examined area is part of the North German Plain, with highlands up to 971 m in its southern region. It is located at latitudes between 51° and 54° North and longitudes between 7° and 11° East (Fig. 1a). It is characterised by a maritime climate, with mean annual temperatures ranging from 8.9 °C to 10.4 °C. Average annual precipitation ranges from 574 mm to 887 mm, with a positive climatic water balance in spring, autumn and winter, and a negative balance in summer.

The database consists of 48 agricultural sites, of which 42 are conventional agricultural systems and six are organic. Two sites were removed due to their observation periods being less than 10 years, leaving 46 entries for the analysis. The observation data consist of 33 sites that have been used for agriculture for more than a century. The remaining 13 sites were converted from forests, meadows or peatlands less than five decades before the start of the observation period. Soil sampling was generally conducted at regular intervals of between one and five years (Höper and Meesenburg, 2021). They were sampled at a depth of up to ~30 cm and consisted of four replicates, which were combined as mixed samples.

In addition, soil horizons were sampled every 10 years at a minimum depth of 1 m. During the soil monitoring programme, a total of 333 soil carbon measurements (replicates not included) were taken at the selected sites from 1992 until the end of

2015. Between the sites, the mineral composition was diverse, with soil textures predominantly sand (20 sites) or silt (21 sites), and less frequently clay soils (five sites; Fig. 1b). The key properties of the examined soils were highly variable between the sites, with SOC ranging from 0.84 % to 4.91 %, C/N from 5.25 to 23.66, bulk density from 1.14 g cm<sup>3</sup> to 1.76 g cm<sup>3</sup>, and pH from 3.9 to 7.9. A special distinctiveness of the investigated area is the occurrence of so-called black sands, which are characterised by comparably high, stable C contents in sand-dominated locations (Vos et al., 2018). Site-specific details are summarised in the appendix (Table A1).

The soil and crop management at the sites was undertaken by the landowners or their associates. Therefore, there is a high degree of variability in the agronomic practices performed at the sites, such as tillage depth and type, fertiliser application, irrigation, crop rotation, harvest and crop residue management. The reported management reflects common agricultural practices for the site-specific regions (more details are given by Höper and Meesenburg (2021)). Control treatments were not available for the sites. Simulated soil temperatures and moisture contents were evaluated using data from 11 German long-term field experiments (Table A2; Flessa et al., 1995; Frohling et al., 1998; Schmädeke, 1998; Leidel et al., 2000; Wolf et al., 2014; Flessa et al., 2017; Mallast et al., 2021; Winkhart et al., 2022).

## 2.2 Model description

The MONICA model is comprised of subroutines that characterise plant growth, C and N dynamics, soil water and soil temperature (for a more detailed description see Nendel (2013)). Execution of the programme requires comprehensive input data, including weather, soil and management information, as well as parameter values describing model settings. Every model iteration simulates soil and plant conditions in a daily time step for soil depths up to two metres. Mathematically, the model is one-dimensional (point model) and the spatial representativeness of the model output depends primarily on the calibration data. Generally, the spatial representation ranges between one m<sup>2</sup> and one km<sup>2</sup>. The soil profile is partitioned into 20 layers, each with 10 cm thickness of homogenous soil conditions. Soil input data are uniformly extrapolated to the relevant depth if the entered soil horizons total less than two metres.




Figure 2. Conceptual SOC pool distribution in MONICA, subdivided into the groups added organic matter (AOM), soil microbial biomass (SMB), soil organic matter (SOM) and inert organic matter (IOM). Except for IOM, each group consists of a slow (s) and a fast (f) pool.

According to the Daisy model, the SOC content is partitioned into three groups composed of dead native soil organic matter (SOM), living soil microbial biomass (SMB) and added organic matter (AOM), each of which contains one rapidly (SOMf, SMBf, AOMf) and one slowly decomposing subpool (Fig. 2; SOMs, SMBs, AOMs; Hansen et al., 1991; Bruun et al., 2003). These compartments are calculated separately for each soil layer. MONICA is usually initialised using a standard distribution of the SOC pools, which should correspond to

an equilibrium at the beginning of the simulation (Bruun and Jensen, 2002). During the initialisation, inert SOC (IOM) is subtracted from the total SOC, based on equations adopted from Falloon et al. (1998). Then the SOM and SMB pools are determined by multiplying the manually entered SOC contents by the corresponding distribution parameters. The AOM pools are initialised and altered by added C contents from crop residues and organic fertilisers. Organic fertiliser and crop types are parameterised with specific N contents ( $\text{NH}_4^+$  and  $\text{NO}_3^-$ ), decomposition rate coefficients and pool distributions. Furthermore, plant biomass accumulation is divided into five groups: root, leaf, shoot, fruit and sugar. For perennial plants, an additional permanent structure is considered. Each group has its own parameters, controlling the growth rate and the biomass that can potentially be harvested or added to the AOM pools. Following a harvest command, accrued crop residues left on an agricultural field are immediately distributed to the AOM pools. Root remains are fragmented into the soil layers, depending on the root mass distribution in the soil profile. During tillage, the SOM content is distributed homogeneously to the soil layers affected by tillage depth.

The initialised pools are managed by first-order degradation due to decomposition, each controlled by related parameters. This process follows a decay rate that is proportional to the pool size (Hansen et al., 1991). In addition, the decomposition rate of organic matter is affected by soil temperature, soil water and clay contents (Bruun et al., 2003; Farina et al., 2021). These environmental conditions influence each pool as rate-modifying factors, with the exception of clay, which solely affects SMBs. Decomposed AOMf and AOMs are subsequently transferred to the SMB pools, from where fractions are respired to the atmosphere as  $\text{CO}_2$  or alternatively converted to SOMs and SOMf, following a pool-specific carbon use efficiency coefficient. Biologically degraded SOMs and SOMf goes through a similar process and are either respired or returned to the SMB pools. A detailed description of the SOM turnover subroutine is depicted in Hansen et al. (1991) and Nendel (2013).

### 2.3 Model modification

Sixteen crop plants (amaranth, asparagus, buckwheat, carrot, flaxseed, field bean, hemp, lupine, serradella, spelt, strawberry, summer onion, sunflower, turnip, vicia) and eight types of organic fertilisers (biogas slurry, calf slurry, dried poultry manure, hoof and horn meal, lime-stabilised sewage sludge, sheep manure, vinasse) were added to the model. This enabled the reproduction of all occurring land management characteristics in the dataset. Parameter values for crop growth (Tesar, 1984; Jeuffroy and Ney, 1997; Allen et al., 1998), fertiliser nutrient contents (Möller and Schultheiß, 2015; Wiesler et al., 2016), crop residues and organic fertiliser decomposition and partitioning coefficients (Gilmour et al., 1998; Thuriès et al., 2001; Lashermes et al., 2009; Peltre et al., 2012; Semenov et al., 2019) were taken from the literature. Furthermore, a specific C content parameter (*CorgContent*) was defined for each crop and organic fertiliser, providing a more adequate replication of C inputs in comparison with the original constant (Möller and Schultheiß, 2015; Wiesler et al., 2016; Ma et al., 2018). In addition, an overhaul of existing crop and fertiliser types was undertaken, ensuring distinct and more reasonable values for essential parameters. Crop growth was calibrated according to the measured yield data, while the residue-to-crop product ratio was adapted from the literature and an ensemble of allometric functions (Bolinder et al., 2007; Franko et al., 2011; Jacobs et al.,

165 2020; Taghizadeh-Toosi, 2020). The differences between the allometric functions served as indicators of uncertainty in the estimation of C inputs from crop residues.

The functions that describe the influence of soil temperature, water and clay content on decomposition were revised and parameterised. This enabled all relevant carbon turnover factors to be tested for uncertainty and subsequently calibrated. The optimal values of the parameters were used as the evidential basis for the prior probability distributions, which is necessary  
170 for model calibration by means of Bayesian inference. For further explanations, see Section 2.5 and Table 1.

Extensive research provided evidence of the influence of temperature on physical soil processes, microbial metabolisms and enzymatic activities (Pietikäinen et al., 2005; Haddix et al., 2006; Meyer et al., 2018). The novel temperature function is described by the following function Eq. (1):

$$f_{TOD} = \frac{1}{(1+e^{Td-(e+T_{opt})})^{Q_{10}/(T_{opt}/\pi)}(-1+Q_{10}^{15.76})^{Td}}, \quad (1)$$

175 where  $T$  is the daily soil temperature in °C,  $Q_{10}$  is the  $Q_{10}$  temperature coefficient specified by the parameter  $QTenFactor$ , and  $T_{opt}$  is the optimal temperature for decomposition in °C defined by the parameter  $TempDecOptimal$  (Fig. B1). From literature analysis, a range of  $38.2 \pm 7.5$  °C was identified for the optimal temperature (Pietikäinen et al., 2005; Richardson et al., 2012; Menichetti et al., 2015; Liu et al., 2018; Čapek et al., 2019) and  $2.6 \pm 0.6$  for the  $Q_{10}$  temperature coefficient (Haddix et al., 2006; Vanhala et al., 2007; Conant et al., 2008; Gillabel et al., 2010; Meyer et al., 2018).

180 Microbial aerobic respiration is often driven by changing soil water contents up to a point where water becomes a restricting property for the availability of oxygen (Curiel Yuste et al., 2007; Schimel et al., 2007). Based on analysis of incubation and field experiments, a Gaussian function was proposed, described by the following Eq (2):

$$f_{MOD} = e^{-18(WFPS_d - M_{opt})^2}, \quad (2)$$

where  $WFPS_d$  is the daily water-filled pore space in % and  $M_{opt}$  is the optimal water-filled pore space (WFPS) for  
185 decomposition in % defined by the parameter  $MoistureDecOptimal$  (Fig. B2). Optimal WFPS for respiration was determined to be around  $59.5 \pm 9.9$  % (Linn and Doran, 1984; Doran et al., 1990; Liebig et al., 1995; Aon et al., 2001; Gabriel and Kellman 2014; Zhang et al., 2015).

Several studies have indicated that increasing soil clay contents have a negative effect on respiration due to the reinforced inaccessibility of SOM for microorganisms (Six and Paustian, 2014; Churchman et al., 2020; Fomina and Skorchod, 2020).  
190 In the MONICA model, the modulating effect of clay on the decomposition rates of SMBs is by default adjustable through the parameter  $LimitClayEffect$ . An inverse logistic function was proposed, described by the following Eq (3):

$$f_{COD} = \frac{(1-LCE)}{(1+e^{-\pi+C*16})+LCE}, \quad (3)$$

where  $C$  is the site-specific clay content in  $\text{kg kg}^{-1}$  and  $LCE$  is the limit to which the clay content affects the mineralisation defined by the parameter  $LimitClayEffect$  (Fig. B3). Laboratory evidence suggests that soils amended with clay minerals reduce  
195 mineralisation rates by  $\sim 1.5 \pm 1.1$  % for every percent of clay added to the soil, followed by a gradual reduction in effectiveness

after a certain clay content is exceeded (Nguye and Marschner, 2014; Schapel et al., 2018; Riaz and Marschner, 2020; Liddle et al., 2020).

## 2.4 Screening of variables relevant to soil carbon turnover

For the purpose of understanding the relationship between input variables, parameters and the predictive proficiency of the MONICA model, a comprehensive sensitivity analysis was conducted using an improved Morris elementary effects screening method (Fig. C1; Morris, 1991; Campolongo et al., 2007; Pujol, 2009). The algorithm demonstrates the capability of determining the importance of parameters in relatively short computing times (Campolongo et al., 2007; Confalonieri et al., 2010). The effect of 51 variables was examined, of which 15 were environmental, six were related to the crop, four to the management and 26 to SOC turnover. Of the turnover parameters, six were unique for each fertiliser and crop residue type. Depending on the tangible properties of the variables (factual or conceptual), the upper and lower bounds of each factor were either derived from the literature or intuitively inferred from model characteristics. After building the parameter space, all the factors were scaled to a range from zero to one. During each iteration, one variable was changed to a random value in the defined space (levels = 5, grid.jump = 3) while the remaining variables were unaltered. Subsequently, the effect of the modified factor on SOC stocks was assessed by comparing the present iteration with the previous one. Based on the magnitude of the changes and the degree of the interaction with other factors, a parameter importance ranking was identified using the two measures  $\mu^*$  and  $\sigma$  (Campolongo et al., 2007).

Common recommendations for the number of model iterations are between 10 to 50 for each variable considered (Campolongo et al., 2007). However, several studies suggest a much higher number of iterations to achieve robust results (Sarrazin et al., 2016; Vanuytrech et al., 2014; Pianosi et al., 2016). As a base value,  $10^3$  iterations and three runs were used in total to accomplish adequate coverage of possible input spaces and to estimate the uncertainty in the elementary effects.

The programming language R (version 4.1.0) with the graphical user interface RStudio (version 1.4.1717) and the sensitivity package (Iooss et al., 2020; version 1.26.0) were used for screening of variables relevant to soil carbon turnover in the MONICA model.

## 2.5 Parameter calibration

Bayesian inference techniques are an efficient approach for the calibration of process-based models (Vrugt, 2016; Van de Schoot et al., 2021). Common methods for prior elicitation, likelihood specification and posterior distributions have been extensively researched (Van de Schoot et al., 2021). For instance, Bayesian calibration has been applied successfully to forest biomass growth (Van Oijen et al., 2005; Svenson et al., 2008) and agro-ecosystem models (Lehuger et al., 2009; Gurung et al., 2020).

As a first step, the experimental data were divided into one subset for calibration and one for validation. In the process, the data were clustered into three groups based on their initial SOC stocks in order to achieve a general representation of the variability between the sites. From each of the groups, two thirds of the sites were randomly sampled for model calibration



(totalling 31 sites) and one third for model validation (totalling 15 sites). The parameter selection process for the calibration was adapted from the results of the sensitivity analysis and the recommendations from Minunno et al. (2013), excluding local and 25 % of the least important parameters.

The calibration of the 14 selected parameters was undertaken by means of Bayesian statistical inference. In Bayesian calibration, the determination of possible parameter values is based on evidential probabilities. Accordingly, it was important that the upper and lower limits of the parameters were based on observed, realistic values. Therefore, a comprehensive literature analysis combined with expert knowledge provided the basis for the prior probability distributions, as shown in Section 2.3 (Oakley and O'Hagan, 2007). A weakly informative prior probability distribution was constructed, with the mean of all identified measurements as the most likely value and the observed lower and upper bounds as the variance for each parameter (Table 1; Van de Schoot et al., 2021). At the beginning of the calibration, a random initial value (within the prior probability distribution) was generated for each parameter, creating a parameter vector for each iteration. Selection for subsequent parameter values was limited by the minimum and maximum value of the prior probability distribution and a randomly determined step size corresponding to 1-25 % of the parameter space (Van Oijen et al., 2005). Accordingly, the subsequent parameter value changes from the previous value by a maximum of 25 % of the entire parameter space. During each iteration, MONICA was executed using the created parameter vector. Afterwards, the generated output was compared with the measured data using a log-Laplace likelihood function suggested by Vrugt (2016), estimating the association between simulated and observed values. This association was a representation of the posterior probability of each parameter vector. The Laplace distribution showed robustness against outliers and random variations, resulting in less biased and more consistent parameter estimates (Schoups and Vrugt, 2010).

A Markov Chain Monte Carlo (MCMC) method, namely the Metropolis-Hastings algorithm, was used for random sampling from proposed values to estimate the posterior probability distribution (Fig. D1; Metropolis et al., 1954; Hastings, 1970; Lehuger et al., 2009). In this process, the likelihood function serves as an indicator to generate a chain of accepted vectors. A sample was accepted in the sequence if the current proposal divided by the previous proposal was greater than or equal to a random number between 0 and 1.

**Table 1. Description of the 14 parameters chosen for the calibration of the C submodel in MONICA. The prior distribution depicts the upper and lower bounds of possible parameter values inferred from the literature; the posterior distribution describes the median value of the posterior and the corresponding standard deviation.**

Parameter		Prior distribution				Posterior distribution	
Notation	Unit	Default value	Minimum value	Maximum value	Reference	Mean	Standard deviation
AOM_FastUtilizationEfficiency	%	0.1	0.1	0.65	Manzoni et al. (2012), Spohn et al. (2016), Qiao et al. (2019), Schroeder et al. (2020), Agumas et	0.27	1.2 <sup>-3</sup>
AOM_SlowUtilizationEfficiency	%	0.4	0.1	0.65	Linn and Doran (1984), Doran et al. (1990), Liebig et al. (1995), Aon et al. (2001), Gabriel and Kellman	0.16	0.01
MoistureDecOptimal	%	NA	0.3	0.76	Nguye and Marschner (2014), Schapel et al. (2018), Riaz and Marschner (2020), Liddle et al.	0.45	3.1 <sup>-3</sup>
LimitClayEffect	%	0.25	0.01	0.5	Haddix et al. (2006), Vanhala et al. (2007), Conant et al. (2008), Gillabel et al. (2010), Meyer et al.	0.34	2.2 <sup>-3</sup>
QTenFactor	Q <sub>10</sub>	NA	1.6	4.2	Kindler et al. (2006), Liang and Balsler (2011), Miltner et al. (2012), Kallenbach et al. (2016)	3.15	0.04
PartSMB_Fast_to_SOM_Fast	%	0.6	0.4	0.8	McGill et al. (1986), Joergensen et al. (1990), Throckmorton et al. (2012), Throckmorton et al. (2012), Creamer et al. (2019), Buckeridge et al. (2020), Geyer et al. (2020), Paustian et al. (1992), Trumbore (2000), Gleixner (2013), Wang et al. (2016), Shi et al. (2020)	0.66	2.1 <sup>-3</sup>
PartSMB_Slow_to_SOM_Fast	%	0.6	0.4	0.8	Manzoni et al. (2012), Saifuddin et al. (2019), Qiao et al. (2019), Schroeder et al. (2020), Agumas et	0.61	1.3 <sup>-3</sup>
SMB_FastDeathRateStandard	d <sup>-1</sup>	0.01	0.01	0.25		0.12	1.1 <sup>-3</sup>
SMB_SlowDeathRateStandard	d <sup>-1</sup>	1.0 <sup>-3</sup>	3.0 <sup>-4</sup>	0.01		5.4 <sup>-3</sup>	5.6 <sup>-5</sup>
SMB_UtilizationEfficiency	d <sup>-1</sup>	0.0	0.1	0.88		0.43	6.6 <sup>-3</sup>
SOM_FastDecCoeffStandard	d <sup>-1</sup>	1.4 <sup>-4</sup>	1.0 <sup>-5</sup>	2.8 <sup>-4</sup>		8.6 <sup>-5</sup>	1.9 <sup>-5</sup>
SOM_SlowDecCoeffStandard	d <sup>-1</sup>	4.3 <sup>-5</sup>	6.4 <sup>-7</sup>	5.0 <sup>-5</sup>		1.3 <sup>-5</sup>	4.0 <sup>-7</sup>
SOM_FastUtilizationEfficiency	%	0.5	0.1	0.88		0.35	6.8 <sup>-3</sup>
SOM_SlowUtilizationEfficiency	%	0.4	0.1	0.88		.49	2.2 <sup>-3</sup>

Three chains with 10<sup>5</sup> iterations and different starting values were created to obtain a reliable inference on the posterior distribution (Van de Schoot et al., 2021). Following the recommendation of Van Oijen et al. (2005), the first ten percent of the accepted vectors were discarded as unrepresentative “burn-in” of the chains. Stationary distribution of the chains was verified using the empirical method developed by Gelman and Rubin (1992), which compares the variance between and within the

260 chains. An adequate convergence was obtained if the Gelman-Rubin shrink factor was  $< 1.2$  (Brooks and Gelman, 1998). The R package coda (Plummer et al., 2006; version 0.19-04) was used for random sampling from the posterior distribution and verification of convergence.

## 2.6 Quantitative methods and model evaluation specifics

The statistical methods for evaluating the sampling distributions and model performances were conducted with the R packages hydroGOF (Zambrano-Bigiarini, 2020; version 0.4-0) and car (Fox et al., 2019; version 3.0-12). Shapiro-Wilk (Shapiro and Wilk, 1965) and Levene's tests (Levene, 1960) were used on the measured fluxes to test the population for normality and variance equality. The Shapiro-Wilk test resulted in rejection of the null hypothesis for the variables ( $p < 0.001$ ) inferring that the examined datasets were not normally distributed. The Levene's test resulted in acceptance of the null hypothesis, implying that the variance in the samples was homoscedastic ( $p = 0.88$ ). Using the interquartile range criterion (Jargowsky and Yang, 270 2005), 19 outliers were identified in the data, which represents  $\sim 6\%$  of the cases. Q-Q plots confirmed the assumption that the data points were nonlinear distributed.

To improve representativeness, the years 1992 to 2000 were excluded from the analysis, reducing the observed number of cases from 333 to 223. The reason for this approach was justified by the inherent risk of statistical biases from overfitting of measured and simulated values due to the initialisation of simulated SOC stocks based on measured values, as significant 275 changes in SOC stocks usually take several years. Furthermore, a spin-up period of 20 simulated years was included to ensure a realistic reflection of available fresh matter in the soil, since the AOM pools were initialised equal to zero. The chosen period corresponds to the maximum duration of mineralisation of most AOM (Thuriès et al., 2001; Semenov et al., 2019).

Based on the reduced sample size and the abovementioned statistical assumptions, Kendall's rank correlation coefficient (T; Kendall, 1948) was selected to assess the statistical dependence between the observed and simulated values. Kendall's Tau is 280 the preferred method for measuring the ordinal association in data consisting of small sample sizes and outliers, and is more robust and efficient than the Pearson correlation coefficient and Spearman's rank correlation coefficient (Bonett and Wright, 2000). In addition, the model results were evaluated with the statistical methods mean absolute error (MAE; Willmott and Matsuura, 2005) for analysis of the average error between observed and simulated pairs, and with the Nash-Sutcliffe model efficiency coefficient (NSE; Nash and Sutcliffe, 1970) to assess the predictive skill of the simulation model (Fig. E1). 285 Furthermore, the slope-intercept was considered in order to better assess the behaviour of the models.

## 3 Results

### 3.1 Simulation of soil and plant growth variables

Soil temperatures were well predicted, with an average error of just under  $\sim 1.8\text{ }^{\circ}\text{C}$  (Table 2). Low temperatures were slightly overestimated and high temperatures were underestimated. In comparison, soil water contents were predicted somewhat less 290 favourably, with an average error of  $\sim 5.4\%$ . Nonetheless, the predictive power was better than the mean of the observations.

The model tended to slightly overestimate moderate water contents and underestimate the measured variance. A comparison of the default with the calibrated plant growth parameters was not performed, since new crops were added to the model in order to simulate all possible systems. The calibrated plant growth model achieved a good representation of measured yields, with an average error of less than  $\sim 1.6$  Mg DM ha<sup>-1</sup> for all crops. Overall, there was a linear relationship between the calibrated and measured yields. However, there was a high variability in the estimation quality between different crop types. Some crops, such as triticale and potatoes, were modelled less adequately than silage maize and sugar beet, which was partly explained by the differences in the number of observations and the variance in yields. The C inputs were compared with the estimated mean of the allometric functions. The model produced an average error that was similar to the calculated difference between the allometric functions ( $\sim 1$  Mg DM ha<sup>-1</sup>).

**Table 2. Statistics describing the performance of the MONICA model regarding soil and plant growth variables. The statistics depicted: Kendall's rank correlation coefficient (T), Nash-Sutcliffe model efficiency coefficient (NSE), mean absolute error (MAE), slope and intercept for characterising the relationship between measured values (except values for C inputs, which were calculated with allometric functions), and simulated values.**


Variables	T	NSE	MAE	Slope	Intercept
Soil temperature (n = 24576)	0.83***	0.86	1.84 [°C]	0.93	1.17
Soil water content (n = 9636)	0.63***	0.39	5.37 [%]	0.84	7.83
Yield (n = 846)	0.65***	0.81	1.62 [Mg DM ha <sup>-1</sup> ]	0.99	0.07
C-Input (n = 846)	0.4***	0.32	0.98 [Mg DM ha <sup>-1</sup> ]	0.88	0.38

\*\*\* p < .001, \*\* p < .01, \* p < .05

### 3.2 Quantification of uncertainty and parameter calibration

A strong relationship was identified between SOC stock change rates and C inputs through organic fertilisation and crop residues (Table F1; *OrganicFertilisation*, *AOM\_DryMatterContent*, *CorgContent*), given the upper and lower bounds of possible values. Additionally, these factors were less affected by other variables, noticeably by the lower  $\sigma$  on average in comparison with  $\mu^*$ . The climate variables temperature (*Temperature*) and precipitation (*Daily\_Precipitation*) were also of greater importance to the SOC turnover rate, but had a higher standard deviation of elementary effects. Except for solar irradiance (*Global\_Radiation*), temperature and precipitation, all other climatic factors had an insignificant effect on SOC stocks. Of all the variables that were initialised at the beginning and remained static during the model run, the initial SOC stock (*SoilOrganicMatter*) was the most influential, followed by the site-specific field capacity (*FieldCapacity*), soil bulk density (*SoilBulkDensity*) and pore volume (*PoreVolume*). Other site-specific variables (*Sand*, *Clay*, *PermanentWiltingPoint*, *pH*, *Skeleton*) were comparably less important. The management factors tillage (*Tillage*), mineral fertilisation (*MineralFertilisation*), irrigation (*Irrigation*) and N content in the added organic matter (*CN\_Ratio\_AOM\_Slow*, *CN\_Ratio\_AOM\_Fast*, *CN*, *AOM\_NH4Content*, *AOM\_NO3Content*) had a small effect on the SOC turnover rate.  $\sigma$  values were generally higher than the overall importance of the input factors, indicating that there is probably an interaction and non-

linear effect between the environmental and management variables. In comparison, the sensitivity of the turnover variables  
 320 was less pronounced, and revealed three particularly important parameters. One C pool parameter  
 (*SOM\_SlowDecCoeffStandard*) and two weather-regulated decomposition factors (*QTenFactor*, *MoistureDecOptimal*). They  
 were similar in significance to precipitation, initial SOC and field capacity. Changes in the parameters *QTenFactor* and  
*MoistureDecOptimal* were more important than in *LimitClayEffect*, which was in accordance with the influence of clay being  
 limited to the SMBs pool. The effect of the factor describing the optimal temperature for decomposition (*TempDecOptimal*)  
 325 was negligible, but considerably gained in importance when a higher range of possible values for the variable temperature was



**Figure 3.** Posterior probability distribution (vertical bars) of 14 C submodel parameters calibrated over the range of the prior probability distribution (density curve). The posterior is depicted as the distribution of accepted values determined by random sampling from  $10^5$  proposed values. The vertical red lines represent the optimal values as the median of the posterior and the blue lines the initial values of each chain.

included. Crop and fertiliser-specific turnover parameters were also sensitive and coincided with the importance of C inputs for SOC stock change rates. Less pronounced sensitivities involved the parameters describing the microbial dynamic (besides the parameter *SMB\_UtilizationEfficiency*), distinguishable by the SMB in the name. Non-linear effects and interactions were more strongly pronounced in the SOC turnover parameters.

330 Figure 3 shows the results of the calibration process, with the posterior distribution curve representing the frequency of accepted values and the vertical red line indicating the optimal value as the median of the posterior. Bayesian inference reduced the uncertainty for all parameters, converging the posterior distribution to a normal or log-normal distribution. Throughout the random sampling, the entire plausible parameter space was covered and curve progressions showed that the optimal values were probably within the specified limits. During the calibration process, the posterior distribution of most parameters became  
335 narrower in comparison with the prior, confirming the convergence to the optimal values. Similar to Lehuger et al. (2009), the acceptance rate for the proposed parameter values was ~25%. Based on the results of the Gelman-Rubin diagnostics, adequate convergence was achieved for all parameters (Fig. G1), signifying the unbiased outcome of the MCMC. The correlations across parameters remained below a coefficient of 0.2, with the exception of *QTenFactor* and *SOM\_SlowDecCoeffStandard* (Fig. H1).

### 340 **3.3 Model performances**

The statistical methods showed that the default MONICA model ( $MONICA_{def}$ ) was inferior to the RothC, ICBM and C-TOOL models, and produced results similar to CENTURY and CCB (Table 3; Fig. 4; Fig. I1). The mean absolute error was the highest of all the models, but was within the margin of error compared with CENTURY and CCB for all sites and with CCB for the validation sites. Kendall's rank correlation coefficient and Nash-Sutcliffe model efficiency coefficient values were  
345 within the margin of error for all models and sites, and therefore showed insignificant differences in the predictive skill of the models.


$MONICA_{def}$  estimations deviated on average by more than ~10% from the measurements. Inclusion of a C content parameter for different crop residues and organic fertilisers in the MONICA model ( $MONICA_{mod}$ ) improved the accuracy of predictions for C stock changes by ~16% for all sites and by ~8% for the validation sites, while the additional global parameter calibration  
350 ( $MONICA_{cal}$ ) improved MONICA's average estimation error by ~29% for all sites and by ~30% for the validation sites. Compared to  $MONICA_{cal}$ , only RothC achieved a lower MAE for all sites. The differences between RothC and  $MONICA_{cal}$  were within the margin of error for the validation, but not for all sites.  $MONICA_{cal}$ ,  $MONICA_{mod}$ , RothC, ICBM and C-TOOL produced errors within the average standard deviation between the replicates of the C measurements for the validation (7.76 Mg C ha<sup>-1</sup>) and for all sites (5.01 Mg C ha<sup>-1</sup>). In contrast,  $MONICA_{def}$ , CENTURY and CCB had errors that were not within  
355 the average standard deviation for all sites. Each model was capable of producing results corresponding to the 95% confidence interval for the validation sites (6.4 Mg C ha<sup>-1</sup>). No model achieved the same for all measurements (3.3 Mg C ha<sup>-1</sup>).

360 **Table 3. Statistics describing the performance of the default (MONICA<sub>def</sub>), the modified, uncalibrated (MONICA<sub>mod</sub>), and the modified, calibrated (MONICA<sub>cal</sub>) soil carbon submodel in MONICA in comparison with five C turnover models for all sites. The first ten years were excluded from the analysis (n = 223). The following statistics are depicted are Kendall's rank correlation coefficient (T), Nash-Sutcliffe model efficiency coefficient (NSE), mean absolute error (MAE) and the slope and intercept for characterising the relationship between measured and simulated C stocks.**

Model	T	NSE	MAE [Mg C ha <sup>-1</sup> ]	Slope	Intercept
MONICA <sub>def</sub>	0.83***	0.91	5.54	1.07	0.43
MONICA <sub>mod</sub>	0.83***	0.93	4.68	1.11	-3.8
MONICA <sub>cal</sub>	0.85***	0.95	3.91	1.02	-0.83
RothC	0.85***	0.96	3.69	1.03	-0.86
C-TOOL	0.83***	0.94	4.43	1.04	-0.23
ICBM	0.85***	0.94	3.99	0.93	3.43
CENTURY	0.79***	0.92	5.31	1.1	-4.19
CCB	0.8***	0.92	5.44	0.96	-1

\*\*\* p < 0.001, \*\* p < 0.01, \* p < 0.05

365 Based on the outcomes of the NSE method, all models performed better than the mean of the measurements, with MONICA<sub>def</sub> achieving the lowest and RothC the highest scores. The differences between MONICA<sub>cal</sub> and MONICA<sub>def</sub> were within the margin of error. The Kendall rank coefficient displayed a significant statistical dependence between the simulated and measured data, with MONICA<sub>cal</sub>, RothC and ICBM achieving the best results and CCB and CENTURY the weakest results. Overall the T coefficients were high, with MONICA<sub>def</sub> achieving average results and MONICA<sub>cal</sub> above average results in contrast with the other models. Of the six models, MONICA<sub>cal</sub> generally produced better results than CENTURY, C-TOOL and CCB. The slope and intercept of the statistical analysis indicated that MONICA<sub>def</sub> generally overestimated decomposition, while MONICA<sub>cal</sub> produced results closer to the measured data. Overall, MONICA<sub>def</sub> underestimated the SOC stocks at 24 sites and overestimated them at two sites, leaving 20 sites that were simulated within the standard deviation of the measurements. In comparison, MONICA<sub>cal</sub> underestimated the SOC stocks at six sites and overestimated them at nine sites, leaving 31 sites that were simulated within the standard deviation of the measurements. The calibration resulted in a model 375 behaviour that was comparable with RothC.



**Figure 4. Linear regression between simulated and observed SOC stocks for 46 BDF sites, all years and 333 measurements for the MONICA (a), RothC (b), C-TOOL (c), ICMB (d), CENTURY (e) and CCB (f) models.**

## 4 Discussion

### 4.1 Differences in models and performances

The MONICA model featured a plant growth, nitrogen and soil water submodel, while RothC, CCB, ICMB, CENTURY and C-TOOL did not. Smith et al. (1997) have argued that coupling C turnover with agro-ecosystems submodels will be necessary to simulate the impact of global change on agricultural production and SOC changes. However, they also assumed that simpler



models have an advantage in simulating various land uses due to the lack of adequate data and the necessary parameterisation of complex models. Other literature shows that C turnover models coupled with sophisticated plant growth models do not result in a better performance, highlighting that complex models are error prone (Mueller et al., 1996; Parton, 1996; Li et al., 1997). This also applied to the data requirements of the MONICA model, which led to the necessary parameterisation of new  
385 crop and fertiliser types in order to simulate all sites. Since there was no additional information in the database about the organic fertiliser and crop residue contents, allometric functions and standard values had to be used to compare ingoing C inputs. This could have led to an inaccurate reproduction of C inputs, since the specific properties of organic fertilisers and crop residues are characterised by high standard deviations (Möller and Schultheiß, 2015; Wiesler et al., 2016; Ma et al., 2018). However, the results of the present study did not confirm the assumption that model complexity is an indicator of predictive  
390 ability in process-based models. Notable differences were driven by model characteristics that were partially controlled by parameterisation. Each model showed weaknesses depending on the specific conditions at each site (Fig. J1 to J7). For instance, MONICA<sub>def</sub> overestimated decomposition at ~52 % at all sites in the dataset, but managed to predict the C losses at the Fuhrberg and Vinnhorst sites better than MONICA<sub>cal</sub>. These sites had undergone recent land-use change and had on average higher initial SOC stocks and losses over the measured period. In temperate agro-ecosystems, conversion from grassland or  
395 forest to arable land is considered to be one of the main reasons for C release (Vos et al., 2019; Poeplau et al., 2020). Correspondingly, sites with recent land-use change showed the strongest decline in C stocks and were therefore a good indicator of the downward trend in SOC stocks. Removing sites with a cultivation period < 50 years (n = 10) resulted in a balanced C budget (0.02 t C ha<sup>-1</sup> a<sup>-1</sup>), which also produced an even better performance by the calibrated MONICA model. The magnitude of changes in SOC stocks for all sites was comparable with other studies conducted in temperate climates,  
400 representing a general trend for German arable land (Steinmann et al., 2016; Sanderman et al., 2017; Vos et al., 2019; Seitz et al., 2022).

All the models underestimated the magnitude of variation in the natural SOC dynamics. The best simulation qualities were generally achieved at sites with slowly changing SOC stocks. Some sites distinguished by dramatic changes in SOC were especially problematic for the models to simulate, in particular the Dinklage site (BDF033\_L), which was characterised by  
405 high C losses, and in contrast the Reinhausen site (BDF051\_L), which was characterised by increasing C stocks. CENTURY and CCB, which underperformed in comparison with the other models, were the best at predicting these highly variable sites. The other models overestimated the SOC stock for BDF033\_L and underestimated it for BDF051\_L. In MONICA<sub>cal</sub>, these two sites caused ~10 % of the mean absolute error.

At site BDF033\_L, the farmer was cultivating winter barley and corn as energy crops, but experienced declining corn yields  
410 over the measured period. This probably led to a negative C balance, even though the field was extensively fertilised with pig and bull slurry. In MONICA, organic fertilisation was the best indicator of SOC changes, while there was no significant relationship between the amount of organic amendments and SOC stock changes in the measured data. Accounting for the quantitative variations in manure application supports the assumption that organic fertilisation is a weak predictor of SOC turnover rates, and correspondingly was overestimated in the model. However, numerous studies clearly show a positive effect

415 of organic amendments on SOC stocks (Koishi et al., 2020; Gross and Glaser, 2021; Roß et al., 2022). Since the BDF data did not feature control treatments, it was not possible to undertake a comparative analysis of the individual sites on the effect of organic inputs. Another consideration concerned site-specific uncertainties in the necessary model input. The properties of organic fertilisers and residues were unknown, and thus had to be supplemented with standard values from the literature. Each fertiliser and crop type had fixed C ratios in the models, while in reality these inputs have a highly variable composition and  
420 can vary greatly depending on farm management, animal diet, climate and soil conditions (Schnug et al., 1996; McCartney et al., 2006; Cajamarca et al., 2019). There are three possible reasons for the disparity between the models and observations at site BDF033\_L: i) the fertiliser and crop-specific nutrient contents were inaccurate, leading to an overestimation of C inputs, ii) the C losses from harvesting were underestimated, or iii) the site and environmental characteristics and their effect on SOC decomposition were incorrectly simulated.

425 In comparison, at site BDF051\_L the farmer had transitioned to a management with reduced tillage intensities and no cultivation of root crops, and managed to increase the C content in the upper soil layer by  $\sim 30 \text{ Mg C ha}^{-1}$  during the measured time period. Although the effect of conservational tillage is still being debated, numerous studies support the effect measured at site BDF051\_L (Freibauer et al., 2004; Luo et al., 2010). In the MONICA model, tillage uniformly distributed SOC in the affected soil layers. This was supported by the sensitivity analysis, which showed that tillage had an effect when there was  
430 heterogeneity in the soil profile. However, based on this study's data, it is likely that the model underestimated the effect of tillage on SOC decomposition. Possible reasons for the disparity were either an imbalance of C inputs into the topsoil or missing functions that describe the processes involved, such as the effect of tillage on soil aggregation, hydraulic properties and root growth (Mondal et al., 2020). The effect of tillage on SOC stocks is currently absent from other C turnover models, such as RothC. Jordon and Smith (2022) suggest modifying the decomposition rate to achieve a realistic representation of the  
435 effect of tillage on SOC, although they assume a small effect of tillage on SOC stocks.

Another probable reason for the differences between the model and measurements were the climate variables. In the data, only precipitation correlated significantly with SOC stock change rates, while in MONICA this was important, but less so than temperature. Sierra et al. (2015) argue that functions in process-based models tend to overestimate the temperature effect on decomposition rates and this imbalance cannot be offset by the moisture effect. This was also likely in the MONICA model,  
440 since the effect of temperature on decomposition was generally twice as strong as moisture. The importance of climate variables is still being debated, with Luo et al. (2017) and Carvalhais et al. (2014) finding major effects on SOC changes, while only minor effects have been established by Fujisaki et al. (2018) and Vos et al. (2019).

#### **4.2 Implications for improving MONICA's soil carbon turnover model**

Most biogeochemical models were developed many decades ago. This is also true of the C turnover submodel in MONICA,  
445 which is based on the Daisy model (Hansen et al., 1991; Abrahamsen and Hansen, 2000). Functions in C turnover models need to be improved in order to estimate SOC dynamics across a wide range of spatiotemporal and agronomic conditions. Correspondingly, general insights into soil processes are needed to justify the

introduction of additional functions. There needs to be further examination of existing mechanisms and the interactive effects of site-specific characteristics, but also missing factors that affect microbial abundance, diversity and activity (Louis et al., 2016). Microorganisms in particular play a significant role in the soil carbon cycle by controlling the rate of mineralisation and promoting plant growth by improving mineral nutrition or stimulating immune and stress reactions in crops (Tardy et al., 2015; Berg et al., 2017). However, microbial models as reviewed in Chandel et al. (2023) vary widely in terms of model structure, processes considered and parameters required, which makes it all the clearer that there is currently no consensus on which approach is best suited to represent the biosphere.

This study attempted to introduce two changes to the default MONICA model: i) a revision, parameterisation and calibration of the functions that describe the influence of soil temperature, water and clay content on decomposition, and ii) an integration of a crop and fertiliser-specific C content parameter. Alteration of the temperature function involved mathematically reformulating the original exponential function into a surge function. The reason for this change was that the default equations did not consider the decreasing activity of soil microorganisms at temperatures above 30 °C (Liu et al., 2018; Fang and Moncrieff, 2001). This could lead to an overestimation of mineralisation, especially under globally increasing temperatures caused by climate change (DWD, 2020). Further improvements could be achieved by incorporating changes in the optimal temperature for decomposition based on site-specific varieties of microbial communities and C qualities (Fierer et al., 2006). The clay function was changed to an inverse logistic function on the basis of the original equation. Many studies suggest that increasing soil clay contents have an adverse effect on SOC decomposition due to the reinforced inaccessibility of organic substances for decomposers (Six and Paustian, 2014; Churchman et al., 2020; Liddle et al., 2020). Adversely, several studies have found no effect or even a positive effect (Wei et al., 2014; Fissore et al., 2016; Zhang et al., 2019). There are multiple explanations for this disparity, e.g. different clay types, residue qualities and microbial diversities (Nguye and Marschner, 2014; Fissore et al., 2016). Without knowing the exact underlying processes to explain the variability and limitation of the available data, only minor changes were made when modifying the clay function. Removing the clay factor might reduce the complexity of the model, but could also negatively impact the predictive capability of the SOC turnover routine. Further evaluation is needed to assess the importance of the clay factor on decomposition and therefore change the function comprehensively.

Furthermore, the soil moisture factor, which was originally based on the pressure potential derived from the van Genuchten model (Van Genuchten, 1980; Abrahamsen and Hansen, 2000), was simplified. The default moisture function was directly controlled by soil water content, soil texture, bulk density and SOC stocks, while the revised Gaussian function was only regulated by soil water content. It is argued that the effect of texture on the decomposing community could be overestimated, since clay as a factor was already included in a function limiting mineralisation. In addition, the influence of soil texture on decomposition proved to be inconsistent, therefore requiring a better understanding of underlying processes (Doran et al., 1990; Scott et al., 1996), while the influence of bulk density was limited by model characteristics as it remained static after initialisation, hampering the reproducibility of management-related soil disturbances on the moisture function. Linn and Doran (1984) showed that the effect of ploughing on bulk density increased the microbial activity and decreased the optimum soil

moisture content, while other studies have also determined an impact of soil structure changes on the mineralisation rate (Franzuebbers, 1999; Vilkiene et al., 2016). Another particularity of the default moisture function was that the optimal water-filled pore space for microbial activity was estimated to be between 10-25 % for sandy soils and 30-35 % for clay soils, while  
485 in the literature a general value of ~60 % is considered the most probable (Linn and Doran, 1984; Doran et al., 1990; Liebig et al., 1995; Aon et al., 2001; Gabriel and Kellman 2014; Zhang et al., 2015). Nevertheless, the simplification is probably less error-prone and easier to understand, but at the same time not optimal for depicting the variance in microbial diversity and their environmental preferences. The inclusion of a more complex moisture function should be based on definite results. As shown with the crop and fertiliser-specific C content parameter, increasing the complexity of a model can be important for  
490 improving performance. According to the sensitivity analysis and the results, the addition of a C content parameter was necessary for the representation of realistic C inputs. However, the limited availability of data hinders the determination of precise C contents in plant residues and organic fertilisers. As mentioned in Section 4.1, values adopted from the literature could have a detrimental effect on the model's predictive capability. Therefore, measurements of C contents in crop residues and organic fertilisers should be part of the experimental design when investigating changes in SOC stocks. To further optimise  
495 the model, it is recommended that the C inputs from crop residues be divided into aboveground and belowground biomass. Currently there is no distinction between roots, shoot and leaves in the AOM pool, even though there is good evidence that roots in particular have a much longer mean residence time in the soil (Poeplau et al., 2021; Sokol and Bradford, 2019).

## 5 Conclusions

The evaluation of the original MONICA model revealed weaknesses in estimating SOC trends. Using the default  
500 parameterisation, decomposition was overestimated and the statistical performance was inferior compared with the C turnover models RothC, ICBM and C-TOOL. MONICA's capabilities of simulating SOC trends increased when more realistic functions were introduced describing the effects of soil temperature, moisture and clay on decomposition, coupled with a better specification of C inputs from organic fertilisation and crop residues. A sensitivity analysis displayed the importance of C inputs and the environmental factors of temperature and precipitation, as well as the corresponding parameters on SOC  
505 variation in the model. A successful calibration by means of Bayesian inference improved the parameterisation of the C turnover subroutine. The modified and calibrated MONICA model outperformed CENTURY, C-TOOL and CCB and produced similar results in comparison to RothC and ICBM. With the exception of certain sites, adequate reproduction of SOC stock change rates was achieved. However, none of the investigated models was capable of simulating each site satisfactorily. All the models underestimated the variation in measured SOC dynamics that occurred at sites with particular properties and  
510 management practices. This paper reveals that even complex, biogeochemical models are capable of performing similarly or even better than the more simplistic C turnover models. It is likely that with better data availability and continuing optimisation of functions representing natural processes, MONICA's performance can be further enhanced.

## Appendix A. General properties of each monitoring and experimental site

515 **Table A1. Description of the long-term monitoring sites (n = 46) used for calibration and validation of the carbon turnover routine in the MONICA model. SOC = soil organic carbon, MAT = mean annual temperature, MAP = mean annual precipitation.**

Site	Location	Start date	End date	Sand [%]	Clay [%]	Bulk density [g cm <sup>-3</sup> ]	SOC [%]	C/N ratio	MAT [°C]	MAP [mm]	Average fertiliser amendments [kg N yr <sup>-1</sup> ]**	Land-use change
BDF001_L	Timmerlah	1992	2014	6	16	1.34	1.2	8.6	9.8	632	151 UAN 27 POM	1760
BDF002_L	Druette	1994	2014	3	15	1.35	1.2	11.0	9.8	631	181 UAN 20 POM	1760
BDF003_L	Ehmen	1993	2014	70	7	1.37	1.4	10.6	9.9	689	151 CAN 17 POM	1945
BDF004_L	Hemmendorf	1995	2014	2	16	1.38	1.1	10.6	9.8	769	139 UAN, CAN 40 PIS	1780
BDF005_L	Reinshof	1991	2014	11	16	1.24	1.0	10.0	9.3	632	183 UAN 4 LFR	1775
BDF006_L*	Mariental	1991	2014	60	16	1.48	0.9	10.4	9.6	601	132 CAN	1760
BDF007_L	Barum	1996	2014	21	7	1.4	1.0	10.0	9.4	674	155 UAN, U 54 POM	1776
BDF008_L*	Hofschwicheldt	1991	2014	17	38	1.33	2.3	9.3	9.8	617	169 UAN, U 33 PIS	1833 1992 drainage
BDF009_L*	Hornburg	1991	2014	2	23	1.29	1.5	9.4	9.4	666	158 UAN 26 POM	1760
BDF010_L*	Uesen	1996	2013	86	4	1.36	2.7	17.7	9.7	680	113 CAN, AS 22 LFR	1897 1997 drainage
BDF012_L	Buehren	1991	2014	14	26	1.35	1.6	10.7	10.0	740	180 UAN 46 PIS	1775
BDF013_L	Ottenstein	1995	2014	4	21	1.15	1.6	7.9	8.9	759	160 CAN, AS 22 PIM	1760
BDF014_L*	Neuhauserfelde	1991	2011	38	11	1.34	0.9	9.3	9.3	887	177 CAN, AS, U 44 LFR	1878
BDF016_L	Tetendorf	1992	2014	81	4	1.43	1.4	17.0	9.3	796	119 CAN, U 148 LFR, SS	1897
BDF017_L*	Lueder	1993	2014	82	6	1.46	1.1	12.0	9.2	735	102 CAN, AS 81 LFR	1777

BDF019_L	Ganderkese	1992	2014	75	6	1.37	2.7	15.7	9.7	693	66 UAN	1780
											184 CAS, LFR	
BDF021_L	Groenheimer Moor	1997	2014	84	5	1.45	2.8	18.3	10.0	770	79.8 POM	1945
												< 1990 = pasture
BDF022_L*	Voxtrup	1993	2014	44	16	1.33	1.4	9.9	10.0	830	82 CAN, AS	1920
											101 CAS, PIS	
BDF024_L*	Dalumer Moor	1992	2014	82	4	1.18	4.7	26.3	10.3	783	129 CAN, AS	1978
											171 CAS	
BDF026_L	Vechtel	1995	2014	95	4	1.34	2.0	13.1	10.1	766	112 CAN, AS	1935
											151 CAS, PIS	
BDF027_L	Barrien	1996	2014	32	5	1.26	1.5	11.5	9.7	715	134 UAN, U	1897
											166 SS	
BDF031_L	Vinnhorst	1992	2011	37	16	1.14	4.0	13.5	9.9	645	120 CAN	1980
BDF032_L	Markhausen	1992	2015	81	4	1.39	3.3	54.8	9.9	777	47 UAN, CAN	1957
											172 CAS, PIS	
BDF033_L	Dinklage	1992	2014	92	3	1.32	1.8	13.1	9.8	765	19 CAN	1973
											199 PIS, CAS	
BDF036_L*	Stuetensen	1993	2014	78	7	1.56	0.9	12.9	9.4	628	175 PIM, CAM	1780
BDF037_L	Schladen	1994	2014	6	31	1.42	2.3	7.9	9.6	708	165 UAN, U	1901
											17 CAS	
BDF039_L	Handeloh	1994	2014	86	5	1.63	1.7	17.3	9.5	824	142 CAN, AS, UAN	1901
											96 PIS, LFR	
BDF042_L	Fuhrberg	1995	2014	87	8	1.33	3.1	10.3	10.0	692	35 UAN, CAN, AS	1950
											54 LFR	< 1988 = pasture
BDF043_L	Oldershausen	1996	2014	2	18	1.38	1.0	9.1	9.3	841	178 U, UAN, AS	1784
											35 SS	
BDF045_L*	Riddagshausen	1994	2014	58	3	1.66	0.8	8.7	9.8	631	12 CAS	1759
BDF046_L*	Rodewald	1997	2014	31	31	1.42	2.0	9.9	10.3	687	114 CAN	1965
											48 PIS, LFR	
BDF047_L	Hiddestorf	1994	2014	2	12	1.37	0.9	8.7	10.0	705	141 UAN, U	1781
											28 CAM	
BDF049_L	Glissen	1994	2014	87	6	1.45	1.5	12.8	10.0	730	65 CAN	1955
											43 PIS	
BDF050_L	Bockheber	1994	2014	78	5	1.43	1.3	12.1	9.3	790	27 CAN	1776

BDF051_L	Reinhausen	1995	2014	12	49	1.43	1.3	6.1	9.3	628	58 SHM 169 UAN, AS 28 CAM	1784
BDF052_L	Suestedt	1996	2014	14	11	1.29	1.5	10.9	9.7	715	109 UAN 101 PIS	1899
BDF056_L*	Meinersen	1996	2014	96	2	1.42	1.3	18.9	9.8	625	62 CAN, UAN 6 LFR	1945
BDF057_L	Starkshorn	1997	2014	81	5	1.55	3.3	15.8	9.2	807	118 CAN, AS 53 POM	1777
BDF058_L	Kuingdorf	1996	2014	12	12	1.42	1.2	9.0	9.8	832	130 UAN, AS 62 CAS, PIS	1846
BDF059_L	Wuelferode	1997	2014	64	12	1.76	0.9	9.0	10.1	684	42 CP	1781
BDF063_L*	Meyenburg	1998	2014	4	36	1.3	2.4	10.3	10.0	818	213 U, AS 27 POM	1980 1982 drainage
BDF064_L	Hohenzethen	1998	2014	79	7	1.44	1.0	12.9	9.7	574	145 UAN, CAN 401 CAS, LFR	1998
BDF065_L*	Juehnde	1998	2014	3	41	1.39	2.6	9.6	9.5	642	130 CAN, U 34 PIS, LFR	1775
BDF067_L*	Listrup	1998	2014	83	6	1.48	1.2	15.5	10.4	770	114 CAN, AN, UAN 116 PIS	1985
BDF069_L	Wendhausen	1998	2014	6	22	1.44	1.7	9.7	10.0	698	1.3 HOM	1845
BDF070_L	Sehlde	2001	2014	9	25	1.48	1.9	8.8	9.6	753	164 U, CAN 5 CAS	1845

\* sites used for model validation (n = 15).

\*\* predominantly applied fertilisers, N values for organic amendments were estimated; UAN = urea ammonium nitrate; U = urea; CAN = calcium ammonium nitrate; AS = ammonium sulfate; AN = ammonium nitrate; POM = poultry manure; PIS = pig slurry; PIM = pig manure; LFR = liquid fermentation residues; SS = sewage sludge; CAS = cattle slurry; CAM = cattle manure; SHM = sheep manure; CP = compost; HOM = horse manure.


**Table A2. Description of the long-term experimental sites (n = 11) used for validation of simulated soil temperatures and water contents. SOC = soil organic carbon, MAT = mean annual temperature, MAP = mean annual precipitation.**

Site	Start date	End date	Sand [%]	Clay [%]	Bulk density [g cm <sup>-3</sup> ]	SOC [%]	pH	C/N ratio	MAT [°C]	MAP [mm]
Bornim	2010	2016	75	6	1.65	1.0	6.6	9.8	8.7	503
Braunschweig	2008	2011	62	8	1.55	0.9	6.1	12.2	9.1	617
Dedelow	2010	2016	59	10	1.45	0.8	6.8	7.5	8.4	485
Göttingen	1992	1996	5	30	1.25	1.6	7.5	8.2	8.7	634
Hennef	1997	2000	8	22	1.31	0.9	6.5	7.5	10.3	837
Hohenheim	2010	2016	3	19	1.3	1.7	7.1	8.4	8.3	688
Hohenschulen	2010	2016	60	11	1.43	1.8	6.7	15.2	8.9	732
Merbitz	2010	2016	16	16	1.42	1.2	7.4	10.9	9.0	520
Rostock	1993	1998	70	7	1.5	0.9	6.5	1.1	9.3	593
Scheyern	1989	1996	22	23	1.25	1.9	6.1	9.4	7.4	833
Viehhausen	2011	2022	14	22	1.4	1.06	6.1	5.1	7.5	797




## Appendix B. Revised environmental decomposition factors

530 Figure B1. The effect of soil temperature on the soil organic carbon decomposition rate. The solid red line describes the standard function that is integrated in the MONICA model. The solid blue line describes the revised function as the best fit curve to the observed data, while the dashed blue lines describe the possible properties of the equation according to the prior probability distribution of the parameters QTenFactor and TempDecOptimal.




535 Figure B2. The effect of soil moisture on the soil organic carbon decomposition rate. The solid red line describes the standard function for a clay soil (clay = 60 %, sand = 10 %, bulk density = 1.4, soil organic carbon = 2 %), while the solid yellow line describes the standard function for a sandy soil (clay = 4 %, sand = 80 %, bulk density = 1.4, soil organic carbon = 1 %). The solid blue line describes the revised function as the best fit curve to the observed data, while the dashed blue lines describe the possible properties of the equation according to the prior probability distribution of the parameter MoistureDecOptimal.



540

Figure B3. The effect of clay on the soil organic carbon decomposition rate. The solid red line describes the standard function that is integrated in the MONICA model. The solid blue line describes the revised function as the best fit curve to the observed data, while the dashed blue lines describe the possible properties of the equation according to the prior probability distribution of the parameter LimitClayEffect.



### Appendix C. Morris elementary effects screening method

545

Table C1. Mathematical formulation of the sensitivity measures of the Morris elementary effects screening method, where  $r$  is the number of input factors and  $EE$  is the elementary effect of each input.

Description	Equation	Properties
Ranking of the input factor (Campolongo et al., 2007)	$\mu_j^* = \frac{1}{r} \sum_{i=1}^r  EE_{ij} $	Mean estimates of the distribution of the absolute values of each input
Standard deviation of the input factor (Morris, 1991)	$\sigma_j = \sqrt{\frac{1}{r} \sum_{i=1}^r (EE_{ij} - \frac{1}{r} \sum_{i=1}^r (EE_{ij})^2)}$	Standard deviation estimates of the absolute values of each input

## Appendix D. Metropolis-Hastings algorithm

550 **Table D1. Mathematical formulation of the Metropolis-Hastings algorithm and the log-Laplace likelihood function, where  $\theta$  is the parameter vector,  $Y$  the dataset used for calibration,  $t$  the number of iterations,  $\sigma$  the standard deviation inferred jointly from the error between observation and simulation,  $\omega$  the model input,  $f(\omega; \theta)$  the model run with the parameter vector  $\theta$ ,  $\alpha$  the acceptance ratio and  $\delta$  a random vector from a multivariate distribution.**

Description	Equation	Properties
Log-Laplace likelihood function (Vrugt, 2016)	$\mathcal{L}(\theta Y) = - \sum_{i=1}^t \log\left(\frac{1}{2\sigma_i}\right) - \sum_{i=1}^t \left(\frac{e^{\bar{Y}-f(\omega_i; \theta)}}{\sigma_i}\right)$	Continuous probability distribution, assumes all error residuals equally
Metropolis-Hastings algorithm (Metropolis et al., 1953; Hastings, 1970)	$\theta^* = \theta_{i-1} + \delta$ $\alpha = \sum_{i=1}^t \frac{\mathcal{L}(\theta^* Y)}{\mathcal{L}(\theta_{i-1} Y)} = \sum_{i=1}^t \frac{\mathcal{L}(Y \theta^*)\mathcal{L}(\theta^*)}{\mathcal{L}(Y \theta_{i-1})\mathcal{L}(\theta_{i-1})}$	Accept $\theta^*$ if $\alpha \geq u$ where $u$ is a uniform random value between 0 and 1, otherwise if $\alpha < u$ reject $\theta^*$

## Appendix E. Model evaluation methods

555 **Table E1. Quantitative methods used for data and model evaluation, where  $S$  is the predicted value,  $O$  is the observed value,  $n$  is the number of cases and  $k$  is the number of divided subgroups from  $n$ .**

Statistical method	Equation	Value range	Properties
Kendall rank correlation coefficient (Kendall, 1938; 1948)	$T = \frac{\sum_{i < j} (\text{sgn}(S_i - S_j) \cdot \text{sgn}(O_i - O_j))}{\frac{n(n-1)}{2}}$	-1 - 1	If $T > 0$ positive association between two variables, otherwise if $T = 0$ no association, or if $T < 0$ negative association
Mean absolute error (Willmott and Matsuura, 2005)	$MAE = \sqrt{\frac{1}{n} \sum_{i=1}^n (S_i - O_i)^2}$	> 0	If $MAE = 0$ perfect agreement between two variables, otherwise if $MAE > 0$ decreasing agreement
Nash-Sutcliffe model efficiency coefficient (Nash and Sutcliffe, 1970)	$NSE = 1 - \frac{\sum_{i=1}^n (S_i - O_i)^2}{\sum_{i=1}^n (O_i - \bar{O})^2}$	< 1	If $NSE = 1$ perfect agreement between two variables, otherwise if $NSE < 1$ decreasing agreement

## Appendix F. Results of the Morris elementary effects screening method

560

Table F1. Results of the Morris method depicting the effects of 51 variables on soil carbon changes. The variables are subdivided into site, management, crop and fertiliser-specific and carbon turnover factors. The parameters *CorgContent* (C content of crop residues and organic fertilisers), *MoistureDecOptimal* (optimal water-filled pore space for decomposition), *TempDecOptimal* (optimal temperature for decomposition) and *QTenFactor* ( $Q_{10}$  temperature coefficient) were included as modifications to the model. The overall importance of an input factor is specified by two sensitivity measures: the mean ( $\mu^*$ ) and the standard deviation of the elementary effect ( $\sigma$ ) of each parameter. Higher values in the two sensitivity measures imply a greater importance of the variable on changes in soil carbon stocks. The uncertainty is presented as the standard deviation between the results of three separate Morris elementary effects screening runs.


565

Variables	Absolute mean of elementary effect $\mu^*$ [g C kg <sup>-1</sup> ]	Standard deviation of elementary effect $\sigma$ [g C kg <sup>-1</sup> ]
Site factor		
Temperature	11.43 ± 0.46	15.9 ± 0.58
SoilOrganicCarbon	4.72 ± 0.14	8.08 ± 0.26
SoilBulkDensity	4.44 ± 0.14	4.96 ± 0.04
Daily_Precipitation	4.16 ± 0.28	8.55 ± 0.71
FieldCapacity	3.7 ± 0.07	7.15 ± 0.49
PoreVolume	2.46 ± 0.13	4.85 ± 0.49
Global_Raditation	1.48 ± 0.05	3.37 ± 1.28
Clay	0.62 ± 0.04	1.4 ± 0.11
CN	0.53 ± 0.05	2.77 ± 0.77
PermanentWiltingPoint	0.24 ± 0.01	0.87 ± 0.12
Sand	0.19 ± 0.02	0.54 ± 0.18
pH	0.0 ± 0.0	0.02 ± 0.02
Wind_Speed	0.0 ± 0.0	0.0 ± 0.0
Relative_Humidity	0.0 ± 0.0	0.0 ± 0.0
Skeleton	0.0 ± 0.0	0.0 ± 0.0
Management factor		
OrganicFertilization	18.36 ± 0.68	16.43 ± 0.44
MineralFertilization	1.03 ± 0.05	3.39 ± 0.26
Irrigation	0.37 ± 0.03	1.01 ± 0.12
Tillage	0.22 ± 0.03	1.42 ± 0.42
Crop and fertiliser factor		
AOM_DryMatterContent	17.75 ± 0.32	16.25 ± 0.39
CorgContent	7.74 ± 0.41	8.67 ± 0.3
PartAOM_to_AOM_Slow	2.82 ± 0.11	4.78 ± 0.11
PartAOM_to_AOM_Fast	2.41 ± 0.12	4.23 ± 0.15
AOM_SlowDecCoeffStandard	1.62 ± 0.14	4.84 ± 0.5

AOM_FastDecCoeffStandard	1.51 ± 0.07	4.94 ± 0.79
PartAOM_Slow_to_SMB_Slow	0.57 ± 0.06	2.05 ± 0.44
CN_Ratio_AOM_Fast	0.44 ± 0.06	1.89 ± 0.57
PartAOM_Slow_to_SMB_Fast	0.44 ± 0.04	1.18 ± 0.39
CN_Ratio_AOM_Slow	0.43 ± 0.06	2.53 ± 0.29
AOM_NH4Content	0.18 ± 0.03	0.95 ± 0.14
AOM_NO3Content	0.18 ± 0.03	0.72 ± 0.15
Carbon turnover factor		
QTenFactor	4.77 ± 0.08	8.56 ± 0.1
MoistureDecOptimal	4.02 ± 0.15	6.71 ± 0.15
SOM_SlowDecCoeffStandard	3.73 ± 0.18	7.11 ± 0.14
SOM_FastUtilizationEfficiency	2.5 ± 0.1	4.44 ± 0.33
AOM_FastUtilizationEfficiency	2.06 ± 0.03	4.87 ± 0.63
PartSOM_Fast_to_SOM_Slow	1.96 ± 0.04	3.5 ± 0.04
SOM_FastDecCoeffStandard	1.75 ± 0.09	3.05 ± 0.16
SMB_UtilizationEfficiency	1.63 ± 0.11	3.59 ± 0.24
AOM_SlowUtilizationEfficiency	1.34 ± 0.15	3.35 ± 0.16
PartSMB_Fast_to_SOM_Fast	1.25 ± 0.09	2.37 ± 0.36
PartSMB_Slow_to_SOM_Fast	0.94 ± 0.03	2.24 ± 0.25
SOM_SlowUtilizationEfficiency	0.81 ± 0.01	1.65 ± 0.07
SMB_FastDeathRateStandard	0.65 ± 0.02	2.0 ± 0.09
SMB_SlowDeathRateStandard	0.53 ± 0.08	1.69 ± 0.49
SMB_FastMaintRateStandard	0.38 ± 0.04	1.29 ± 0.48
SMB_SlowMaintRateStandard	0.37 ± 0.02	0.92 ± 0.07
LimitClayEffect	0.25 ± 0.02	0.63 ± 0.13
PartSOM_to_SMB_Slow	0.1 ± 0.0	0.13 ± 0.0
PartSOM_to_SMB_Fast	0.04 ± 0.01	0.13 ± 0.13
TempDecOptimal	0.0 ± 0.01	0.1 ± 0.17


---

Figure G1. Evolution of the Gelman and Rubin shrink factor. The 50 % (solid line) and 97.5 % (dashed red line) quantiles show the convergence of the calibration.



## Appendix H. Correlations between C turnover parameters

575 **Figure H1.** Correlation matrix quantifying the statistical relationship between the prior probability distributions of the SOM submodel parameters.



## Appendix I. Model performance

580 **Table II. Statistics describing the performance of the default (MONICA<sub>def</sub>), the modified, uncalibrated (MONICA<sub>mod</sub>) and the modified, calibrated (MONICA<sub>cal</sub>) soil carbon submodel in MONICA in comparison with five C turnover models for 15 validation sites. The first ten years were excluded from the analysis (n = 70). The statistics depicted are Kendall's rank correlation coefficient (T), Nash-Sutcliffe model efficiency coefficient (NSE), mean absolute error (MAE) and slope and intercept for characterising the relationship between measured and simulated C stocks.**


Description	T	NSE	MAE (Mg C ha <sup>-1</sup> )	Slope	Intercept
MONICA <sub>def</sub>	0.85***	0.93	5.41	1.07	0.54
MONICA <sub>mod</sub>	0.81***	0.94	4.97	1.08	-2.14
MONICA <sub>cal</sub>	0.83***	0.97	3.81	1.02	-0.41
RothC	0.83***	0.97	3.68	1.04	-0.57
C-TOOL	0.85***	0.95	4.63	1	2.68
ICBM	0.87***	0.95	3.6	0.91	4.86
CENTURY	0.84***	0.95	4.32	0.88	4.37
CCB	0.8***	0.93	4.95	0.89	3.29

585 \*\*\* p < 0.001, \*\* p < 0.01, \* p < 0.05




## Appendix J. Model performances


Figure J1. Performance of the default MONICA model for each individual site. The first 10 measured years were excluded.




590 **Figure 12. Performance of the modified and calibrated MONICA model for each individual site. The first 10 measured years were excluded.**




**Figure 13. Performance of the CCB model for each individual site. The first 10 measured years were excluded.**




**Figure I4. Performance of the CENTURY model for each individual site. The first 10 measured years were excluded.**




**Figure 15. Performance of the C-TOOL model for each individual site. The first 10 measured years were excluded.**



**Figure 16. Performance of the ICBM model for each individual site. The first 10 measured years were excluded.**



**Figure 17. Performance of the RothC model for each individual site. The first 10 measured years were excluded.**



## Data availability

The described long-term soil monitoring data is available from the State Office for Mining, Energy and Geology of the Lower Saxony state government. Restrictions apply to the availability of these data, which were used under license for this study.

- 615 Data are available from the State Office for Mining, Energy and Geology of the Lower Saxony state government from the website: [https://www.lbeg.niedersachsen.de/boden\\_grundwasser/bodenmonitoring/bodendauerbeobachtung/das-bodendauerbeobachtungsprogramm-von-niedersachsen-572.html](https://www.lbeg.niedersachsen.de/boden_grundwasser/bodenmonitoring/bodendauerbeobachtung/das-bodendauerbeobachtungsprogramm-von-niedersachsen-572.html) and mail: [bodenkundlicheberatung@lbeg.niedersachsen.de](mailto:bodenkundlicheberatung@lbeg.niedersachsen.de)
- Additional data is openly available at <https://doi.org/10.5281/zenodo.8380332>. Further information on the Berge, Dedelow, Hohenheim, Hohenschulen and Merbitz data is comprehensively described in the publication by Mallast et al. (2021), mail: [janine.mallast@thuenen.de](mailto:janine.mallast@thuenen.de). The Deutsche Wetterdienst weather data is openly available under the following website: [https://opendata.dwd.de/climate\\_environment/CDC/](https://opendata.dwd.de/climate_environment/CDC/).
- 620

## Code availability

The original and modified MONICA model source code is openly available at <https://doi.org/10.5281/zenodo.8380341>.

## Author's contributions

- 625 Konstantin Aiteew prepared the data, developed and implemented the modifications and executed the MONICA model simulations. He developed the scripts for model performance evaluation, sensitivity analysis and calibration and also wrote the manuscript.
- Jarmo Rouhiainen gave advice in developing the sensitivity analysis algorithm, helped interpret MONICA's source code and supported the drafting of the manuscript.
- 630 Claas Nendel advised on the execution of the MONICA model and supported in drafting of the manuscript.
- René Dechow prepared the simulation of the C turnover models CCB, CENTURY, C-TOOL, ICBM and RothC. He also gave advice, material and support in drafting of the manuscript.

## Competing interests

The authors declare that they have no conflict of interest.

## 635 Acknowledgements

We are very grateful to the state government of Lower Saxony and the State Office for Mining, Energy and Geology for collecting and providing the data used in this model evaluation and optimisation. We also like to thank the Chair of Organic



Farming and Agronomy, TUM School of Life Sciences, Technical University of Munich and the Deutscher Wetterdienst for providing additional data. We are also grateful to Michael Berg-Mohnicke for his support regarding questions about the  
640 MONICA model. We also thank our colleagues PD. Dr. Axel Don, Dr. Christopher Poeplau, Dr. Felix Seidel, Dr. Florian Schneider, Dr. Julia Schröder and Dr. Balazs Grosz for their support and comments on the manuscript. We would specifically like to thank the German Federal Ministry of Food and Agriculture and the Agency for Renewable Resources for financially supporting the MASTER project and our work.

## 645 References

- Abrahamsen, P. and Hansen, S.: Daisy: an open soil-crop-atmosphere system model, *Environmental Modelling & Software*, 15, 313-330, [https://doi.org/10.1016/S1364-8152\(00\)00003-7](https://doi.org/10.1016/S1364-8152(00)00003-7), 2000.
- Agumas, B., Blagodatsky, S., Balume, I., Musyoki, M. K., Marhan, S., and Rasche, F.: Microbial carbon use efficiency during plant residue decomposition: Integrating multi-enzyme stoichiometry and C balance approach, *Applied Soil Ecology*, 159, 103820, <https://doi.org/10.1016/j.apsoil.2020.103820>, 2021.
- 650 Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56, Fao, Rome, 300, D05109, 1998.
- Amelung, W., Bossio, D., de Vries, W., Kögel-Knabner, I., Lehmann, J., Amundson, R., Bol, R., Collins, C., Lal, R., Leifeld, J., Minasny, B., Pan, G., Paustian, K., Rumpel, C., Sanderman, J., van Groenigen, J. W., Mooney, S., van Wesemael, B., Wander, M., and Chabbi, A.:
- 655 Towards a global-scale soil climate mitigation strategy, *Nature Communications*, 11, 5427, 10.1038/s41467-020-18887-7, 2020.
- Amundson, R. and Biardeau, L.: Opinion: Soil carbon sequestration is an elusive climate mitigation tool, *Proceedings of the National Academy of Sciences*, 115, 11652 - 11656, 2018.
- Aon, M. A., Cabello, M. N., Sarena, D., Colaneri, A., Franco, M., Burgos, J., and Cortassa, S.: I. Spatio-temporal patterns of soil microbial and enzymatic activities in an agricultural soil, *Applied Soil Ecology*, 18, 239-254, 2001.
- 660 Asseng, S., Ewert, F., Rosenzweig, C., Jones, J. W., Hatfield, J. L., Ruane, A. C., Boote, K. J., Thorburn, P. J., Rötter, R. P., Cammarano, D., Brisson, N., Basso, B., Martre, P., Aggarwal, P. K., Angulo, C., Bertuzzi, P., Biernath, C., Challinor, A. J., Doltra, J., Gayler, S., Goldberg, R., Grant, R., Heng, L., Hooker, J., Hunt, L. A., Ingwersen, J., Izaurralde, R. C., Kersebaum, K. C., Müller, C., Naresh Kumar, S., Nendel, C., O'Leary, G., Olesen, J. E., Osborne, T. M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M. A., Shcherbak, I., Steduto, P., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M., Waha, K., Wallach, D., White, J. W., Williams, J. R., and Wolf, J.:
- 665 Uncertainty in simulating wheat yields under climate change, *Nature Climate Change*, 3, 827-832, 10.1038/nclimate1916, 2013.
- Bassu, S., Brisson, N., Durand, J.-L., Boote, K., Lizaso, J., Jones, J. W., Rosenzweig, C., Ruane, A. C., Adam, M., Baron, C., Basso, B., Biernath, C., Boogaard, H., Conijn, S., Corbeels, M., Deryng, D., De Sanctis, G., Gayler, S., Grassini, P., Hatfield, J., Hoek, S., Izaurralde, C., Jongschaap, R., Kemanian, A. R., Kersebaum, K. C., Kim, S.-H., Kumar, N. S., Makowski, D., Müller, C., Nendel, C., Priesack, E., Pravia, M. V., Sau, F., Shcherbak, I., Tao, F., Teixeira, E., Timlin, D., and Waha, K.: How do various maize crop models vary in their responses to climate change factors?, *Global Change Biology*, 20, 2301-2320, <https://doi.org/10.1111/gcb.12520>, 2014.
- 670 Berg, G., Köberl, M., Rybakova, D., Müller, H., Grosch, R., and Smalla, K.: Plant microbial diversity is suggested as the key to future biocontrol and health trends, *FEMS Microbiology Ecology*, 93, fix050, 10.1093/femsec/fix050, 2017.
- Bolinder, M. A., Janzen, H. H., Gregorich, E. G., Angers, D. A., and VandenBygaart, A. J.: An approach for estimating net primary productivity and annual carbon inputs to soil for common agricultural crops in Canada, *Agriculture, Ecosystems & Environment*, 118, 29-42, <https://doi.org/10.1016/j.agee.2006.05.013>, 2007.
- 675 Bonett, D. G. and Wright, T. A.: Sample size requirements for estimating pearson, kendall and spearman correlations, *Psychometrika*, 65, 23-28, 10.1007/BF02294183, 2000.
- Brooks, S. P. and Gelman, A.: General Methods for Monitoring Convergence of Iterative Simulations, *Journal of Computational and Graphical Statistics*, 7, 434-455, 10.1080/10618600.1998.10474787, 1998.
- 680 Bruun, S. and Jensen, L. S.: Initialisation of the soil organic matter pools of the Daisy model, *Ecological Modelling*, 153, 291-295, 2002.
- Bruun, S., Christensen, B. T., Hansen, E. M., Magid, J., and Jensen, L. S.: Calibration and validation of the soil organic matter dynamics of the Daisy model with data from the Askov long-term experiments, *Soil Biology and Biochemistry*, 35, 67-76, [https://doi.org/10.1016/S0038-0717\(02\)00237-7](https://doi.org/10.1016/S0038-0717(02)00237-7), 2003.
- Buckeridge, K. M., Mason, K. E., McNamara, N. P., Ostle, N., Puissant, J., Goodall, T., Griffiths, R. I., Stott, A. W., and Whitaker, J.:
- 685 Environmental and microbial controls on microbial necromass recycling, an important precursor for soil carbon stabilization, *Communications Earth & Environment*, 1, 36, 10.1038/s43247-020-00031-4, 2020.
- Cajamarca, S. M., Martins, D., da Silva, J., Fontenelle, M. R., Guedes, Í. M., de Figueiredo, C. C., and Pacheco Lima, C. E.: Heterogeneity in the Chemical Composition of Biofertilizers, Potential Agronomic Use, and Heavy Metal Contents of Different Agro-Industrial Wastes, 10.3390/su11071995, 2019.
- 690 Campolongo, F., Cariboni, J., and Saltelli, A.: An effective screening design for sensitivity analysis of large models, *Environmental Modelling & Software*, 22, 1509-1518, <https://doi.org/10.1016/j.envsoft.2006.10.004>, 2007.
- Čapek, P., Starke, R., Hofmockel, K. S., Bond-Lamberty, B., and Hess, N.: Apparent temperature sensitivity of soil respiration can result from temperature driven changes in microbial biomass, *Soil Biology and Biochemistry*, 135, 286-293, <https://doi.org/10.1016/j.soilbio.2019.05.016>, 2019.
- 695 Carvalhais, N., Forkel, M., Khomik, M., Bellarby, J., Jung, M., Migliavacca, M., Mu, M., Saatchi, S., Santoro, M., Thurner, M., Weber, U., Ahrens, B., Beer, C., Cescatti, A., Randerson, J. T., and Reichstein, M.: Global covariation of carbon turnover times with climate in terrestrial ecosystems, *Nature*, 514, 213-217, 10.1038/nature13731, 2014.
- Chandel, A. K., Jiang, L., and Luo, Y.: Microbial Models for Simulating Soil Carbon Dynamics: A Review, *Journal of Geophysical Research: Biogeosciences*, 128, e2023JG007436, <https://doi.org/10.1029/2023JG007436>, 2023.

- 700 Churchman, G. J., Singh, M., Schapel, A., Sarkar, B., and Bolan, N.: Clay minerals as the key to the sequestration of carbon in soils, *Clays and Clay Minerals*, 68, 135-143, 10.1007/s42860-020-00071-z, 2020.
- Coleman, K. and Jenkinson, D. S.: RothC-26.3 - A Model for the turnover of carbon in soil, *Evaluation of Soil Organic Matter Models*, Berlin, Heidelberg, 1996//, 237-246.
- 705 Conant, R. T., Drijber, R. A., Haddix, M. L., Parton, W. J., Paul, E. A., Plante, A. F., Six, J., and Steinweg, J. M.: Sensitivity of organic matter decomposition to warming varies with its quality, *Global Change Biology*, 14, 868-877, 2008.
- Confalonieri, R., Bellocchi, G., Bregaglio, S., Donatelli, M., and Acutis, M.: Comparison of sensitivity analysis techniques: A case study with the rice model WARM, *Ecological Modelling*, 221, 1897-1906, <https://doi.org/10.1016/j.ecolmodel.2010.04.021>, 2010.
- Creamer, C. A., Foster, A. L., Lawrence, C., McFarland, J., Schulz, M., and Waldrop, M. P.: Mineralogy dictates the initial mechanism of microbial necromass association, *Geochimica et Cosmochimica Acta*, 260, 161-176, <https://doi.org/10.1016/j.gca.2019.06.028>, 2019.
- 710 Curiel Yuste, J., Baldocchi, D. D., Gershenson, A., Goldstein, A., Misson, L., and Wong, S.: Microbial soil respiration and its dependency on carbon inputs, soil temperature and moisture, *Global Change Biology*, 13, 2018-2035, <https://doi.org/10.1111/j.1365-2486.2007.01415.x>, 2007.
- Doran, J., Mielke, L., and Power, J.: Microbial activity as regulated by soil water-filled pore space, *Transactions 14th International Congress of Soil Science*, Kyoto, Japan, August 1990, Volume III., 94-99.
- 715 Falloon, P., Smith, P., Coleman, K., and Marshall, S.: Estimating the size of inert organic matter pool from total soil organic carbon content for use the Rothamsted Carbon Model, *Soil Biology & Biochemistry - SOIL BIOL BIOCHEM*, 30, 1207-1211, 10.1016/S0038-0717(97)00256-3, 1998.
- FAO: The State of Food and Agriculture, 2014.
- FAO: Land use statistics and indicators. Global, regional and country trends, 2000-2020, 2022.
- 720 Farina, R., Sándor, R., Abdalla, M., Álvaro-Fuentes, J., Bechini, L., Bolinder, M. A., Brilli, L., Chenu, C., Clivot, H., De Antoni Migliorati, M., Di Bene, C., Dorich, C. D., Ehrhardt, F., Ferchoud, F., Fitton, N., Francaviglia, R., Franko, U., Giltrap, D. L., Grant, B. B., Guenet, B., Harrison, M. T., Kirschbaum, M. U. F., Kuka, K., Kulmala, L., Liski, J., McGrath, M. J., Meier, E., Menichetti, L., Moyano, F., Nendel, C., Recous, S., Reibold, N., Shepherd, A., Smith, W. N., Smith, P., Soussana, J.-F., Stella, T., Taghizadeh-Toosi, A., Tsutskikh, E., and Bellocchi, G.: Ensemble modelling, uncertainty and robust predictions of organic carbon in long-term bare-fallow soils, *Global Change Biology*, 27, 904-928, <https://doi.org/10.1111/gcb.15441>, 2021.
- 725 Fierer, N., Colman, B. P., Schimel, J. P., and Jackson, R. B.: Predicting the temperature dependence of microbial respiration in soil: A continental-scale analysis, *Global Biogeochemical Cycles*, 20, <https://doi.org/10.1029/2005GB002644>, 2006.
- Fissore, C., Jurgensen, M. F., Pickens, J., Miller, C., Page-Dumroese, D., and Giardina, C. P.: Role of soil texture, clay mineralogy, location, and temperature in coarse wood decomposition—a mesocosm experiment, *Ecosphere*, 7, e01605, <https://doi.org/10.1002/ecs2.1605>, 2016.
- 730 Flessa, H., Dörsch, P., and Beese, F.: Seasonal variation of N<sub>2</sub>O and CH<sub>4</sub> fluxes in differently managed arable soils in southern Germany, *Journal of Geophysical Research: Atmospheres*, 100, 23115-23124, <https://doi.org/10.1029/95JD02270>, 1995.
- Flessa, H., Fuß, R., Andres, M., Augustin, J., Christen, O., Dittert, K., Hegewald, H., Heilmann, H., Huth, V., Kage, H., Kern, J., Kesenheimer, K., Knieß, A., Köbke, S., Lewandowski, I., Mallast, J., Moffat, A. M., Mühling, K.-H., Öhlschläger, G., and Stichnothe, H.: Minderung von Treibhausgasemissionen im Rapsanbau unter besonderer Berücksichtigung der Stickstoffdüngung, *Johann Heinrich von Thünen-Institut, Braunschweig* 2017.
- 735 Fomina, M. and Skorochood, I.: Microbial Interaction with Clay Minerals and Its Environmental and Biotechnological Implications, 10.3390/min10100861, 2020.
- Franko, U., Oelschlägel, B., and Schenk, S.: Simulation of temperature-, water- and nitrogen dynamics using the model CANDY, *Ecological Modelling*, 81, 213-222, [https://doi.org/10.1016/0304-3800\(94\)00172-E](https://doi.org/10.1016/0304-3800(94)00172-E), 1995.
- 740 Franko, U., Kolbe, H., Thiel, E., and Ließ, E.: Multi-site validation of a soil organic matter model for arable fields based on generally available input data, *Geoderma*, 166, 119-134, <https://doi.org/10.1016/j.geoderma.2011.07.019>, 2011.
- Franzluebbers, A. J.: Microbial activity in response to water-filled pore space of variably eroded southern Piedmont soils, *Applied Soil Ecology*, 11, 91-101, [https://doi.org/10.1016/S0929-1393\(98\)00128-0](https://doi.org/10.1016/S0929-1393(98)00128-0), 1999.
- Freibauer, A., Rounsevell, M. D. A., Smith, P., and Verhagen, J.: Carbon sequestration in the agricultural soils of Europe, *Geoderma*, 122, 1-23, <https://doi.org/10.1016/j.geoderma.2004.01.021>, 2004.
- 745 Froking, S. E., Mosier, A. R., Ojima, D. S., Li, C., Parton, W. J., Potter, C. S., Priesack, E., Stenger, R., Haberbosch, C., Dörsch, P., Flessa, H., and Smith, K. A.: Comparison of N<sub>2</sub>O emissions from soils at three temperate agricultural sites: simulations of year-round measurements by four models, *Nutrient Cycling in Agroecosystems*, 52, 77-105, 10.1023/A:1009780109748, 1998.
- Fujisaki, K., Chevallier, T., Chapuis-Lardy, L., Albrecht, A., Razafimbelo, T., Masse, D., Ndour, Y. B., and Chotte, J.-L.: Soil carbon stock changes in tropical croplands are mainly driven by carbon inputs: A synthesis, *Agriculture, Ecosystems & Environment*, 259, 147-158, <https://doi.org/10.1016/j.agee.2017.12.008>, 2018.
- 750 Fuss, S., Lamb, W. F., Callaghan, M. W., Hilaire, J., Creutzig, F., Amann, T., Beringer, T., de Oliveira Garcia, W., Hartmann, J., Khanna, T., Luderer, G., Nemet, G. F., Rogelj, J., Smith, P., Vicente, J. L. V., Wilcox, J., del Mar Zamora Dominguez, M., and Minx, J. C.: Negative emissions—Part 2: Costs, potentials and side effects, *Environmental Research Letters*, 13, 063002, 10.1088/1748-9326/aabf9f, 2018.

- 755 Gabriel, C.-E. and Kellman, L.: Investigating the role of moisture as an environmental constraint in the decomposition of shallow and deep mineral soil organic matter of a temperate coniferous soil, *Soil Biology and Biochemistry*, 68, 373-384, <https://doi.org/10.1016/j.soilbio.2013.10.009>, 2014.
- Gelman, A. and Rubin, D. B.: Inference from iterative simulation using multiple sequences, *Statistical science*, 457-472, 1992.
- 760 Geyer, K., Schnecker, J., Grandy, A. S., Richter, A., and Frey, S.: Assessing microbial residues in soil as a potential carbon sink and moderator of carbon use efficiency, *Biogeochemistry*, 151, 237-249, 10.1007/s10533-020-00720-4, 2020.
- Gillabel, J., Cebrian-Lopez, B., Six, J., and Merckx, R.: Experimental evidence for the attenuating effect of SOM protection on temperature sensitivity of SOM decomposition, *Global Change Biology*, 16, 2789-2798, <https://doi.org/10.1111/j.1365-2486.2009.02132.x>, 2010.
- Gilmour, J. T., Norman, R. J., Mauromoustakos, A., and Gale, P. M.: Kinetics of Crop Residue Decomposition: Variability among Crops and Years, *Soil Science Society of America Journal*, 62, 750-755, <https://doi.org/10.2136/sssaj1998.03615995006200030030x>, 1998.
- 765 Gleixner, G.: Soil organic matter dynamics: a biological perspective derived from the use of compound-specific isotopes studies, *Ecological research*, 28, 683-695, 2013.
- Gross, A. and Glaser, B.: Meta-analysis on how manure application changes soil organic carbon storage, *Scientific Reports*, 11, 5516, 10.1038/s41598-021-82739-7, 2021.
- 770 Gurung, R. B., Ogle, S. M., Breidt, F. J., Williams, S. A., and Parton, W. J.: Bayesian calibration of the DayCent ecosystem model to simulate soil organic carbon dynamics and reduce model uncertainty, *Geoderma*, 376, 114529, <https://doi.org/10.1016/j.geoderma.2020.114529>, 2020.
- Haddix, M., Steinweg, M., Conant, R., Plante, A., Paul, E., and Six, J.: Effect of Temperature on the Dynamics of Different Soil Organic Matter Fractions, *The 18th World Congress of Soil Science*,
- 775 Hansen, S., Jensen, H., Nielsen, N., and Svendsen, H.: Simulation of nitrogen dynamics and biomass production in winter wheat using the Danish simulation model DAISY, *Fertilizer research*, 27, 245-259, 1991.
- Hastings, W. K.: Monte Carlo sampling methods using Markov chains and their applications, *Biometrika*, 57, 97-109, 10.1093/biomet/57.1.97, 1970.
- Höper, H. and Meesenburg, H.: 30 Jahre Bodendauerbeobachtung in Niedersachsen, *GeoBerichte*, 39, 2021.
- 780 Jacobs, A., Poeplau, C., Weiser, C., Fahrion-Nitschke, A., and Don, A.: Exports and inputs of organic carbon on agricultural soils in Germany, *Nutrient Cycling in Agroecosystems*, 118, 249-271, 10.1007/s10705-020-10087-5, 2020.
- Jargowsky, P. A. and Yang, R.: Descriptive and Inferential Statistics, in: *Encyclopedia of Social Measurement*, edited by: Kempf-Leonard K., Elsevier, New York, 659-668, <https://doi.org/10.1016/B0-12-369398-5/00145-6>, 2005.
- Jeuffroy, M.-H. and Ney, B.: Crop physiology and productivity, *Field Crops Research*, 53, 3-16, [https://doi.org/10.1016/S0378-4290\(97\)00019-1](https://doi.org/10.1016/S0378-4290(97)00019-1), 1997.
- 785 Joergensen, R. G., Brookes, P. C., and Jenkinson, D. S.: Survival of the soil microbial biomass at elevated temperatures, *Soil Biology and Biochemistry*, 22, 1129-1136, [https://doi.org/10.1016/0038-0717\(90\)90039-3](https://doi.org/10.1016/0038-0717(90)90039-3), 1990.
- Jordon, M. W. and Smith, P.: Modelling soil carbon stocks following reduced tillage intensity: A framework to estimate decomposition rate constant modifiers for RothC-26.3, demonstrated in north-west Europe, *Soil and Tillage Research*, 222, 105428, <https://doi.org/10.1016/j.still.2022.105428>, 2022.
- 790 Kallenbach, C. M., Frey, S. D., and Grandy, A. S.: Direct evidence for microbial-derived soil organic matter formation and its ecophysiological controls, *Nature Communications*, 7, 13630, 10.1038/ncomms13630, 2016.
- Kendall, M. G.: A New Measure of Rank Correlation, *Biometrika*, 30, 81-93, 10.2307/2332226, 1938.
- Kendall, M. G.: Rank correlation methods, 1948.
- 795 Kersebaum, K. C.: Application of a simple management model to simulate water and nitrogen dynamics, *Ecological Modelling*, 81, 145-156, [https://doi.org/10.1016/0304-3800\(94\)00167-G](https://doi.org/10.1016/0304-3800(94)00167-G), 1995.
- Kersebaum, K. C.: Modelling nitrogen dynamics in soil-crop systems with HERMES, *Nutrient Cycling in Agroecosystems*, 77, 39-52, 2006.
- Kindler, R., Miltner, A., Richnow, H.-H., and Kästner, M.: Fate of gram-negative bacterial biomass in soil—mineralization and contribution to SOM, *Soil Biology and Biochemistry*, 38, 2860-2870, <https://doi.org/10.1016/j.soilbio.2006.04.047>, 2006.
- 800 Koishi, A., Bragazza, L., Maltas, A., Guillaume, T., and Sinaj, S.: Long-Term Effects of Organic Amendments on Soil Organic Matter Quantity and Quality in Conventional Cropping Systems in Switzerland, 10.3390/agronomy10121977, 2020.
- Kollas, C., Kersebaum, K. C., Nendel, C., Manevski, K., Müller, C., Palosuo, T., Armas-Herrera, C. M., Beaudoin, N., Bindi, M., Charfeddine, M., Conradt, T., Constantin, J., Eitzinger, J., Ewert, F., Ferrise, R., Gaiser, T., Cortazar-Atauri, I. G. d., Giglio, L., Hlavinka, P., Hoffmann, H., Hoffmann, M. P., Launay, M., Manderscheid, R., Mary, B., Mirschel, W., Moriondo, M., Olesen, J. E., Öztürk, I., Pacholski, A., Ripoche-Wachter, D., Roggero, P. P., Roncossek, S., Rötter, R. P., Ruget, F., Sharif, B., Trnka, M., Ventrella, D., Waha, K.,
- 805 Wegehenkel, M., Weigel, H.-J., and Wu, L.: Crop rotation modelling—A European model intercomparison, *European Journal of Agronomy*, 70, 98-111, <https://doi.org/10.1016/j.eja.2015.06.007>, 2015.
- Kostková, M., Hlavinka, P., Pohanková, E., Kersebaum, K. C., Nendel, C., Gobin, A., Olesen, J. E., Ferrise, R., Dibari, C., Takáč, J., Topaj, A., Medvedev, S., Hoffmann, M. P., Stella, T., Balek, J., Ruiz-Ramos, M., Rodríguez, A., Hooogenboom, G., Shelia, V., Ventrella, D., Giglio, L., Sharif, B., Öztürk, I., Rötter, R. P., Balkovič, J., Skalský, R., Moriondo, M., Thaler, S., Žalud, Z., and Trnka, M.: Performance of 13 crop

- 810 simulation models and their ensemble for simulating four field crops in Central Europe, *The Journal of Agricultural Science*, 159, 69-89, 10.1017/S0021859621000216, 2021.
- Kothari, K., Battisti, R., Boote, K. J., Archontoulis, S. V., Confalone, A., Constantin, J., Cuadra, S. V., Debaeke, P., Faye, B., Grant, B., Hoogenboom, G., Jing, Q., van der Laan, M., Macena da Silva, F. A., Marin, F. R., Nehbandani, A., Nendel, C., Purcell, L. C., Qian, B., Ruane, A. C., Schoving, C., Silva, E. H. F. M., Smith, W., Soltani, A., Srivastava, A., Vieira, N. A., Slone, S., and Salmerón, M.: Are soybean models ready for climate change food impact assessments?, *European Journal of Agronomy*, 135, 126482, <https://doi.org/10.1016/j.eja.2022.126482>, 2022.
- 815 Lashermes, G., Nicolardot, B., Parnaudeau, V., Thuriès, L., Chaussod, R., Guillotin, M. L., Linères, M., Mary, B., Metzger, L., Morvan, T., Tricaud, A., Villette, C., and Houot, S.: Indicator of potential residual carbon in soils after exogenous organic matter application, *European Journal of Soil Science*, 60, 297-310, <https://doi.org/10.1111/j.1365-2389.2008.01110.x>, 2009.
- 820 Lehuger, S., Gabrielle, B., Oijen, M. v., Makowski, D., Germon, J. C., Morvan, T., and Hénault, C.: Bayesian calibration of the nitrous oxide emission module of an agro-ecosystem model, *Agriculture, Ecosystems & Environment*, 133, 208-222, <https://doi.org/10.1016/j.agee.2009.04.022>, 2009.
- Leidel, S., Augustin, J., Köppen, D., and Merbach, W.: Einfluss Unterschiedlicher Organisch-Mineralischer N-Düngung auf die Lachgas- und Methanemission eines ackerbaulich genutzten standortes Norddeutschlands, *Archives of Agronomy and Soil Science*, 45, 453-469, 10.1080/03650340009366141, 2000.
- 825 Levene, H.: Robust Test for Equality of Variances, in: *Contributions to Probability and Statistics: Essays in Honor of Harold Hotelling* edited by: Olkin, I., Stanford University Press, Palo Alto, 278-292, 1960.
- Li, C., Frolking, S., Crocker, G. J., Grace, P. R., Klír, J., Körchens, M., and Poulton, P. R.: Simulating trends in soil organic carbon in long-term experiments using the DNDC model, *Geoderma*, 81, 45-60, [https://doi.org/10.1016/S0016-7061\(97\)00080-3](https://doi.org/10.1016/S0016-7061(97)00080-3), 1997.
- 830 Liang, C. and Balsler, T. C.: Microbial production of recalcitrant organic matter in global soils: implications for productivity and climate policy, *Nature Reviews Microbiology*, 9, 75-75, 2011.
- Liddle, K., McGonigle, T., and Koiter, A.: Microbe Biomass in Relation to Organic Carbon and Clay in Soil, *Soil Systems*, 4, 41, 2020.
- Liebig, M., Jones, A., Doran, J., and Mielke, L.: Potential soil respiration and relationship to soil properties in ridge tillage, *Soil Science Society of America Journal*, 59, 1430-1435, 1995.
- 835 Linn, D. M. and Doran, J. W.: Effect of water-filled pore space on carbon dioxide and nitrous oxide production in tilled and nontilled soils, *Soil Science Society of America Journal*, 48, 1267-1272, 1984.
- Liu, Y., He, N., Wen, X., Xu, L., Sun, X., Yu, G., Liang, L., and Schipper, L. A.: The optimum temperature of soil microbial respiration: Patterns and controls, *Soil Biology and Biochemistry*, 121, 35-42, <https://doi.org/10.1016/j.soilbio.2018.02.019>, 2018.
- 840 Louis, B. P., Maron, P.-A., Viaud, V., Leterme, P., and Menasseri-Aubry, S.: Soil C and N models that integrate microbial diversity, *Environmental Chemistry Letters*, 14, 331-344, 10.1007/s10311-016-0571-5, 2016.
- Lugato, E., Bampa, F., Panagos, P., Montanarella, L., and Jones, A.: Potential carbon sequestration of European arable soils estimated by modelling a comprehensive set of management practices, *Global Change Biology*, 20, 3557-3567, <https://doi.org/10.1111/gcb.12551>, 2014.
- Luo, Z., Wang, E., and Sun, O. J.: Can no-tillage stimulate carbon sequestration in agricultural soils? A meta-analysis of paired experiments, *Agriculture, Ecosystems & Environment*, 139, 224-231, <https://doi.org/10.1016/j.agee.2010.08.006>, 2010.
- 845 Luo, Z., Feng, W., Luo, Y., Baldock, J., and Wang, E.: Soil organic carbon dynamics jointly controlled by climate, carbon inputs, soil properties and soil carbon fractions, *Global Change Biology*, 23, 4430-4439, <https://doi.org/10.1111/gcb.13767>, 2017.
- Ma, S., He, F., Tian, D., Zou, D., Yan, Z., Yang, Y., Zhou, T., Huang, K., Shen, H., and Fang, J.: Variations and determinants of carbon content in plants: a global synthesis, *Biogeosciences*, 15, 693-702, 10.5194/bg-15-693-2018, 2018.
- Mallast, J., Stichnothe, H., Flessa, H., Fuß, R., Lucas-Moffat, A., Petersen-Schlapkohl, U., Augustin, J., Hagemann, U., Kesenheimer, K., Ruser, R., Suárez, T., Prochnow, A., Dittert, K., Köbke, S., Huth, V., Glatzel, S., Rübiger, T., Knieß, A., Kage, H., and Christen, O.: Multi-variable experimental data set of agronomic data and gaseous soil emissions from maize, oilseed rape and other energy crops at eight sites in Germany, *Open Data Journal for Agricultural Research*, 7, 11-19, 10.18174/odjar.v7i0.16124, 2021.
- 850 Manzoni, S., Taylor, P., Richter, A., Porporato, A., and Ågren, G. I.: Environmental and stoichiometric controls on microbial carbon-use efficiency in soils, *New Phytologist*, 196, 79-91, <https://doi.org/10.1111/j.1469-8137.2012.04225.x>, 2012.
- 855 McCartney, D. H., Block, H. C., Dubeski, P. L., and Ohama, A. J.: Review: The composition and availability of straw and chaff from small grain cereals for beef cattle in western Canada, *Canadian Journal of Animal Science*, 86, 443-455, 10.4141/A05-092, 2006.
- McGill, W. B., Cannon, K. R., Robertson, J. A., and Cook, F. D.: Dynamics of soil microbial biomass and water-soluble organic C in Breton L after 50 years of cropping to two rotations, *Canadian journal of soil science*, 66, 1-19, 1986.
- Menichetti, L., Reyes Ortigoza, A. L., García, N., Giagnoni, L., Nannipieri, P., and Renella, G.: Thermal sensitivity of enzyme activity in tropical soils assessed by the Q10 and equilibrium model, *Biology and Fertility of Soils*, 51, 299-310, 10.1007/s00374-014-0976-x, 2015.
- 860 Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., and Teller, E.: Equation of state calculations by fast computing machines, *The journal of chemical physics*, 21, 1087-1092, 1953.
- Meyer, N., Welp, G., and Amelung, W.: The Temperature Sensitivity (Q10) of Soil Respiration: Controlling Factors and Spatial Prediction at Regional Scale Based on Environmental Soil Classes, *Global Biogeochemical Cycles*, 32, 306-323, <https://doi.org/10.1002/2017GB005644>, 2018.
- 865

- Miltner, A., Bombach, P., Schmidt-Brücken, B., and Kästner, M.: SOM genesis: microbial biomass as a significant source, *Biogeochemistry*, 111, 41-55, 10.1007/s10533-011-9658-z, 2012.
- 870 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong, S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke, S., Richer-de-Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., van Wesemael, B., and Winowiecki, L.: Soil carbon 4 per mille, *Geoderma*, 292, 59-86, <https://doi.org/10.1016/j.geoderma.2017.01.002>, 2017.
- 875 Minunno, F., Van Oijen, M., and Pereira, J.: Selecting Parameters for Bayesian Calibration of a Process-Based Model: A Methodology Based on Canonical Correlation Analysis, *Journal on Uncertainty Quantification*, 1, 370-385, 10.1137/120891344, 2013.
- Möller, K. and Schultheiß, U.: Chemical characterization of commercial organic fertilizers, *Archives of Agronomy and Soil Science*, 61, 989-1012, 10.1080/03650340.2014.978763, 2015.
- Mondal, S., Chakraborty, D., Bandyopadhyay, K., Aggarwal, P., and Rana, D. S.: A global analysis of the impact of zero-tillage on soil physical condition, organic carbon content, and plant root response, *Land Degradation & Development*, 31, 557-567, <https://doi.org/10.1002/ldr.3470>, 2020.
- 880 Morris, M. D.: Factorial Sampling Plans for Preliminary Computational Experiments, *Technometrics*, 33, 161-174, 10.1080/00401706.1991.10484804, 1991.
- Mueller, T., Jensen, L. S., Hansen, S., and Nielsen, N. E.: Simulating soil carbon and nitrogen dynamics with the soil-plant-atmosphere system model DAISY, *Evaluation of Soil Organic Matter Models*, Berlin, Heidelberg, 1996//, 275-281,
- 885 Murphy, B.: Soil Carbon Sequestration as an Elusive Climate Mitigation Tool,
- Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I — A discussion of principles, *Journal of Hydrology*, 10, 282-290, [https://doi.org/10.1016/0022-1694\(70\)90255-6](https://doi.org/10.1016/0022-1694(70)90255-6), 1970.
- Nendel, C.: MONICA: A Simulation Model for Nitrogen and Carbon Dynamics in Agro-Ecosystems, in, 389-405, 10.1007/978-3-319-01017-5\_23, 2013.
- 890 Nendel, C., Melzer, D., and Thorburn, P. J.: The nitrogen nutrition potential of arable soils, *Scientific Reports*, 9, 5851, 10.1038/s41598-019-42274-y, 2019.
- Nendel, C., Kersebaum, K. C., Mirschel, W., and Wenkel, K. O.: Testing farm management options as climate change adaptation strategies using the MONICA model, *European Journal of Agronomy*, 52, 47-56, <https://doi.org/10.1016/j.eja.2012.09.005>, 2014.
- Nendel, C., Berg, M., Kersebaum, K. C., Mirschel, W., Specka, X., Wegehenkel, M., Wenkel, K., and Wieland, R.: The MONICA model: Testing predictability for crop growth, soil moisture and nitrogen dynamics, *Ecological Modelling*, 222, 1614-1625, 2011.
- 895 Ngyue, T.-T. and Marschner, P.: Respiration in mixes of sandy and clay soils: influence of clay type and addition rate, *Journal of soil science and plant nutrition*, 14, 881-887, 2014.
- Oakley, J. E. and O'Hagan, A.: Uncertainty in prior elicitation: a nonparametric approach, *Biometrika*, 94, 427-441, 10.1093/biomet/asm031, 2007.
- 900 Parton, W. J.: Ecosystem model comparisons: science or fantasy world?, *Evaluation of Soil Organic Matter Models*, Berlin, Heidelberg, 1996//, 133-142,
- Paustian, K., Parton, W. J., and Persson, J.: Modeling soil organic matter in organic-amended and nitrogen-fertilized long-term plots, *Soil science society of America journal*, 56, 476-488, 1992.
- Peltre, C., Christensen, B., Dragon, S., Icard, C., Kätterer, T., and Houot, S.: RothC simulation of carbon accumulation in soil after repeated application of widely different organic amendments, *Soil Biology & Biochemistry*, 52, 49-60, 10.1016/j.soilbio.2012.03.023, 2012.
- 905 Pianosi, F., Beven, K., Freer, J., Hall, J. W., Rougier, J., Stephenson, D. B., and Wagener, T.: Sensitivity analysis of environmental models: A systematic review with practical workflow, *Environmental Modelling & Software*, 79, 214-232, <https://doi.org/10.1016/j.envsoft.2016.02.008>, 2016.
- Pietikäinen, J., Pettersson, M., and Bååth, E.: Comparison of temperature effects on soil respiration and bacterial and fungal growth rates, *FEMS Microbiol Ecol*, 52, 49-58, 10.1016/j.femsec.2004.10.002, 2005.
- 910 Plummer, M. B., Nicky; Cowles, Kate; Vines, Karen; Sarkar, Deepayan; Bates, Douglas; Almond, Russel; Magnuson, Arni: CODA: Convergence Diagnosis and Output Analysis for MCMC [code], 2020.
- Poeplau, C. and Don, A.: Carbon sequestration in agricultural soils via cultivation of cover crops – A meta-analysis, *Agriculture, Ecosystems & Environment*, 200, 33-41, <https://doi.org/10.1016/j.agee.2014.10.024>, 2015.
- 915 Poeplau, C., Don, A., and Schneider, F.: Roots are key to increasing the mean residence time of organic carbon entering temperate agricultural soils, *Global Change Biology*, 27, 4921-4934, <https://doi.org/10.1111/gcb.15787>, 2021.
- Poeplau, C., Jacobs, A., Don, A., Vos, C., Schneider, F., Wittnebel, M., Tiemeyer, B., Heidkamp, A., Prietz, R., and Flessa, H.: Stocks of organic carbon in German agricultural soils—Key results of the first comprehensive inventory, *Journal of Plant Nutrition and Soil Science*, 183, 665-681, <https://doi.org/10.1002/jpln.202000113>, 2020.
- 920 Pujol, G.: Simplex-based screening designs for estimating metamodels, *Reliability Engineering & System Safety*, 94, 1156-1160, <https://doi.org/10.1016/j.ress.2008.08.002>, 2009.

- Qiao, Y., Wang, J., Liang, G., Du, Z., Zhou, J., Zhu, C., Huang, K., Zhou, X., Luo, Y., Yan, L., and Xia, J.: Global variation of soil microbial carbon-use efficiency in relation to growth temperature and substrate supply, *Scientific Reports*, 9, 5621, 10.1038/s41598-019-42145-6, 2019.
- 925 Riaz, M. and Marschner, P.: Sandy Soil Amended with Clay Soil: Effect of Clay Soil Properties on Soil Respiration, Microbial Biomass, and Water Extractable Organic C, *Journal of Soil Science and Plant Nutrition*, 20, 2465-2470, 10.1007/s42729-020-00312-z, 2020.
- Richardson, J., Chatterjee, A., and Darrel Jenerette, G.: Optimum temperatures for soil respiration along a semi-arid elevation gradient in southern California, *Soil Biology and Biochemistry*, 46, 89-95, <https://doi.org/10.1016/j.soilbio.2011.11.008>, 2012.
- Riggers, C., Poelplau, C., Don, A., Frühauf, C., and Dechow, R.: How much carbon input is required to preserve or increase projected soil organic carbon stocks in German croplands under climate change?, *Plant and Soil*, 460, 417-433, 10.1007/s11104-020-04806-8, 2021.
- 930 Roß, C.-L., Baumecker, M., Ellmer, F., and Kautz, T.: Organic Manure Increases Carbon Sequestration Far beyond the &ldquo;4 per 1000 Initiative&rdquo; Goal on a Sandy Soil in the Thyrow Long-Term Field Experiment DIV.2, 10.3390/agriculture12020170, 2022.
- Rötter, R. P., Palosuo, T., Kersebaum, K. C., Angulo, C., Bindi, M., Ewert, F., Ferrise, R., Hlavinka, P., Moriondo, M., and Nendel, C.: Simulation of spring barley yield in different climatic zones of Northern and Central Europe: a comparison of nine crop models, *Field Crops Research*, 133, 23-36, 2012.
- 935 Saifuddin, M., Bhatnagar, J. M., Segrè, D., and Finzi, A. C.: Microbial carbon use efficiency predicted from genome-scale metabolic models, *Nature Communications*, 10, 3568, 10.1038/s41467-019-11488-z, 2019.
- Salo, T. J., Palosuo, T., Kersebaum, K. C., Nendel, C., Angulo, C., Ewert, F., Bindi, M., Calanca, P., Klein, T., and Moriondo, M.: Comparing the performance of 11 crop simulation models in predicting yield response to nitrogen fertilization, *The Journal of Agricultural Science*, 154, 1218-1240, 2016.
- 940 Sanderman, J., Hengl, T., and Fiske, G. J.: Correction for Sanderman et al., Soil carbon debt of 12,000 years of human land use, *Proceedings of the National Academy of Sciences*, 115, E1700-E1700, 10.1073/pnas.1800925115, 2018.
- Sarrazin, F., Pianosi, F., and Wagener, T.: Global Sensitivity Analysis of environmental models: Convergence and validation, *Environmental Modelling & Software*, 79, 135-152, <https://doi.org/10.1016/j.envsoft.2016.02.005>, 2016.
- Schapel, A., Marschner, P., and Churchman, J.: Clay amount and distribution influence organic carbon content in sand with subsoil clay addition, *Soil and Tillage Research*, 184, 253-260, <https://doi.org/10.1016/j.still.2018.08.001>, 2018.
- 945 Schimel, J., Balsler, T. C., and Wallenstein, M.: MICROBIAL STRESS-RESPONSE PHYSIOLOGY AND ITS IMPLICATIONS FOR ECOSYSTEM FUNCTION, *Ecology*, 88, 1386-1394, <https://doi.org/10.1890/06-0219>, 2007.
- Schmädeke, F.: Lachgas- und Methaneinflüsse eines Gley-Auenbodens unter dem Einfluss einer Rapsfruchtfolge und in Abhängigkeit von der N-Düngung, <http://dx.doi.org/10.53846/goediss-2312>,
- 950 Schnug, E., Oswald, P., and Haneklaus, S.: Organic manure management and efficiency: Role of organic fertilizers and their management practices, in: *Fertilizers and Environment: Proceedings of the International Symposium "Fertilizers and Environment"*, held in Salamanca, Spain, 26-29, September, 1994, edited by: Rodriguez-Barrueco, C., Springer Netherlands, Dordrecht, 259-265, 10.1007/978-94-009-1586-2\_44, 1996.
- Schoups, G. and Vrugt, J. A.: A formal likelihood function for parameter and predictive inference of hydrologic models with correlated heteroscedastic, and non-Gaussian errors, *Water Resources Research*, 46, <https://doi.org/10.1029/2009WR008933>, 2010.
- 955 Schroeder, J., Jannoura, R., Beuschel, R., Pfeiffer, B., Dyckmans, J., Murugan, R., Chavannavar, S., Wachendorf, C., and Joergensen, R. G.: Carbon use efficiency and microbial functional diversity in a temperate Luvisol and a tropical Nitisol after millet litter and N addition, *Biology and Fertility of Soils*, 56, 1139-1150, 10.1007/s00374-020-01487-4, 2020.
- Scott, N. A., Cole, C. V., Elliott, E. T., and Huffman, S. A.: Soil Textural Control on Decomposition and Soil Organic Matter Dynamics, *Soil Science Society of America Journal*, 60, 1102-1109, <https://doi.org/10.2136/sssaj1996.0361599500600040020x>, 1996.
- 960 Seitz, D., Fischer, L. M., Dechow, R., Wiesmeier, M., and Don, A.: The potential of cover crops to increase soil organic carbon storage in German croplands, *Plant and Soil*, 10.1007/s11104-022-05438-w, 2022.
- Semenov, V., Pautova, N., Lebedeva, T., Khromyckina, D., Semenova, N., and Lopes de Gerenyu, V.: Plant Residues Decomposition and Formation of Active Organic Matter in the Soil of the Incubation Experiments, *Eurasian Soil Science*, 52, 1183-1194, 10.1134/S1064229319100119, 2019.
- 965 Shapiro, S. S. and Wilk, M. B.: An Analysis of Variance Test for Normality (Complete Samples), *Biometrika*, 52, 591-611, 10.2307/2333709, 1965.
- Shi, Z., Allison, S. D., He, Y., Levine, P. A., Hoyt, A. M., Beem-Miller, J., Zhu, Q., Wieder, W. R., Trumbore, S., and Randerson, J. T.: The age distribution of global soil carbon inferred from radiocarbon measurements, *Nature Geoscience*, 13, 555-559, 10.1038/s41561-020-0596-z, 2020.
- 970 Shoemaker, W. R., Jones, S. E., Muscarella, M. E., Behringer, M. G., Lehmkuhl, B. K., and Lennon, J. T.: Microbial population dynamics and evolutionary outcomes under extreme energy limitation, *Proceedings of the National Academy of Sciences*, 118, e2101691118, 2021.
- Sierra, C. A., Trumbore, S. E., Davidson, E. A., Vicca, S., and Janssens, I.: Sensitivity of decomposition rates of soil organic matter with respect to simultaneous changes in temperature and moisture, *Journal of Advances in Modeling Earth Systems*, 7, 335-356, <https://doi.org/10.1002/2014MS000358>, 2015.
- 975

- Six, J. and Paustian, K.: Aggregate-associated soil organic matter as an ecosystem property and a measurement tool, *Soil Biology and Biochemistry*, 68, A4-A9, <https://doi.org/10.1016/j.soilbio.2013.06.014>, 2014.
- Smith, P., Andr n, O., Karlsson, T., Per l , P., Regina, K., Rounsevell, M., and Van Wesemael, B.: Carbon sequestration potential in European croplands has been overestimated, *Global Change Biology*, 11, 2153-2163, <https://doi.org/10.1111/j.1365-2486.2005.01052.x>, 2005.
- Smith, P., Smith, J., Powelson, D., McGill, W., Arah, J., Chertov, O., Coleman, K., Franko, U., Froking, S., and Jenkinson, D.: A comparison of the performance of nine soil organic matter models using datasets from seven long-term experiments, *Geoderma*, 81, 153-225, 1997.
- Sommer, R. and Bossio, D.: Dynamics and climate change mitigation potential of soil organic carbon sequestration, *Journal of environmental management*, 144C, 83-87, 10.1016/j.jenvman.2014.05.017, 2014.
- Specka, X., Nendel, C., Hagemann, U., Pohl, M., Hoffmann, M., Barkusky, D., Augustin, J., Sommer, M., and van Oost, K.: Reproducing CO<sub>2</sub> exchange rates of a crop rotation at contrasting terrain positions using two different modelling approaches, *Soil and Tillage Research*, 156, 219-229, <https://doi.org/10.1016/j.still.2015.05.007>, 2016.
- Spohn, M., P tsch, E. M., Eichorst, S. A., Woebken, D., Wanek, W., and Richter, A.: Soil microbial carbon use efficiency and biomass turnover in a long-term fertilization experiment in a temperate grassland, *Soil Biology and Biochemistry*, 97, 168-175, <https://doi.org/10.1016/j.soilbio.2016.03.008>, 2016.
- Steinmann, T., Welp, G., Wolf, A., Holbeck, B., Gro e-R schkamp, T., and Amelung, W.: Repeated monitoring of organic carbon stocks after eight years reveals carbon losses from intensively managed agricultural soils in Western Germany, *Journal of Plant Nutrition and Soil Science*, 179, 355-366, <https://doi.org/10.1002/jpln.201500503>, 2016.
- Svensson, M., Jansson, P.-E., Gustafsson, D., Kleja, D. B., Langvall, O., and Lindroth, A.: Bayesian calibration of a model describing carbon, water and heat fluxes for a Swedish boreal forest stand, *Ecological Modelling*, 213, 331-344, <https://doi.org/10.1016/j.ecolmodel.2008.01.001>, 2008.
- Taghizadeh-Toosi, A., Cong, W.-F., Eriksen, J., Mayer, J., Olesen, J. E., Keel, S. G., Glendining, M., K tterer, T., and Christensen, B. T.: Visiting dark sides of model simulation of carbon stocks in European temperate agricultural soils: allometric function and model initialization, *Plant and Soil*, 450, 255-272, 10.1007/s11104-020-04500-9, 2020.
- Tardy, V., Spor, A., Mathieu, O., L v que, J., Terrat, S., Plassart, P., Regnier, T., Bardgett, R. D., van der Putten, W. H., Roggero, P. P., Seddaiu, G., Bagella, S., Lemanceau, P., Ranjard, L., and Maron, P.-A.: Shifts in microbial diversity through land use intensity as drivers of carbon mineralization in soil, *Soil Biology and Biochemistry*, 90, 204-213, <https://doi.org/10.1016/j.soilbio.2015.08.010>, 2015.
- Tesar, M. B.: Physiological basis of crop growth and development,
- Throckmorton, H. M., Bird, J. A., Dane, L., Firestone, M. K., and Horwath, W. R.: The source of microbial C has little impact on soil organic matter stabilisation in forest ecosystems, *Ecology letters*, 15, 1257-1265, 2012.
- Thuri s, L., Pansu, M., Feller, C., Herrmann, P., and R my, J. C.: Kinetics of added organic matter decomposition in a Mediterranean sandy soil, *Soil Biology and Biochemistry*, 33, 997-1010, [https://doi.org/10.1016/S0038-0717\(01\)00003-7](https://doi.org/10.1016/S0038-0717(01)00003-7), 2001.
- Trumbore, S.: Age of soil organic matter and soil respiration: Radiocarbon constraints on belowground C dynamics, *Ecological Applications*, 10, 399-411, [https://doi.org/10.1890/1051-0761\(2000\)010\[0399:AOSOMA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0399:AOSOMA]2.0.CO;2), 2000.
- van de Schoot, R., Depaoli, S., King, R., Kramer, B., M rtens, K., Tadesse, M. G., Vannucci, M., Gelman, A., Veen, D., Willemsen, J., and Yau, C.: Bayesian statistics and modelling, *Nature Reviews Methods Primers*, 1, 1, 10.1038/s43586-020-00001-2, 2021.
- van Genuchten, M. T.: A Closed-form Equation for Predicting the Hydraulic Conductivity of Unsaturated Soils, *Soil Science Society of America Journal*, 44, 892-898, <https://doi.org/10.2136/sssaj1980.03615995004400050002x>, 1980.
- Van Oijen, M., Rougier, J., and Smith, R.: Bayesian calibration of process-based forest models: bridging the gap between models and data, *Tree Physiology*, 25, 915-927, 10.1093/treephys/25.7.915, 2005.
- Vanhala, P., Karhu, K., Tuomi, M., Sonninen, E., Jungner, H., Fritze, H., and Liski, J.: Old soil carbon is more temperature sensitive than the young in an agricultural field, *Soil Biology and Biochemistry*, 39, 2967-2970, 2007.
- Vanuytrecht, E., Raes, D., and Willems, P.: Global sensitivity analysis of yield output from the water productivity model, *Environmental Modelling & Software*, 51, 323-332, <https://doi.org/10.1016/j.envsoft.2013.10.017>, 2014.
- Vilkien , M., Ambrazaitien , D., Kar auskien , D., and Dabkevi ius, Z.: Assessment of soil organic matter mineralization under various management practices, *Acta Agriculturae Scandinavica, Section B — Soil & Plant Science*, 66, 641-646, 10.1080/09064710.2016.1162845, 2016.
- Vos, C., Jaconi, A., Jacobs, A., and Don, A.: Hot regions of labile and stable soil organic carbon in Germany – Spatial variability and driving factors, *SOIL*, 4, 153-167, 10.5194/soil-4-153-2018, 2018.
- Vos, C., Don, A., Hobley, E. U., Prietz, R., Heidkamp, A., and Freibauer, A.: Factors controlling the variation in organic carbon stocks in agricultural soils of Germany, *European Journal of Soil Science*, 70, 550-564, <https://doi.org/10.1111/ejss.12787>, 2019.
- Vrugt, J. A.: Markov chain Monte Carlo simulation using the DREAM software package: Theory, concepts, and MATLAB implementation, *Environmental Modelling & Software*, 75, 273-316, <https://doi.org/10.1016/j.envsoft.2015.08.013>, 2016.
- Wang, J., Xiong, Z., and Kuzyakov, Y.: Biochar stability in soil: meta-analysis of decomposition and priming effects, *GCB Bioenergy*, 8, 512-523, <https://doi.org/10.1111/gcbb.12266>, 2016.



- Wei, H., Guenet, B., Vicca, S., Nunan, N., Asard, H., AbdElgawad, H., Shen, W., and Janssens, I. A.: High clay content accelerates the decomposition of fresh organic matter in artificial soils, *Soil Biology and Biochemistry*, 77, 100-108, <https://doi.org/10.1016/j.soilbio.2014.06.006>, 2014.
- 1035 Wiesler, F. H.-R., Kerstin; Gäth, Stefan; George, Eckhard; Greef, Jörg M; Hölzle, Ludwig E; Holz, Falko; Hülsbergen, Kurt -Jürgen; Pfeil Rudolf; Severin, Karl; Frede, Hans-Georg; Blum, Birgit; Schenkel, Hans; Horst, Walter; Dittert, Klaus; Ebertseder, Thomas; Osterburg, Bernhard; Philipp, Werner; Pietsch, Magdalena: Anwendung von organischen Düngern und organischen Reststoffen in der Landwirtschaft, <https://doi.org/10.12767/buel.v94i1.124> 2016.
- Willmott, C. J. and Matsuura, K.: Advantages of the mean absolute error (MAE) over the root mean square error (RMSE) in assessing average model performance, *Climate research*, 30, 79-82, 2005.
- 1040 Winkhart, F., Mösl, T., Schmid, H., and Hülsbergen, K.-J.: Effects of Organic Maize Cropping Systems on Nitrogen Balances and Nitrous Oxide Emissions, 10.3390/agriculture12070907, 2022.
- Wolf, U., Fuß, R., Höppner, F., and Flessa, H.: Contribution of N<sub>2</sub>O and NH<sub>3</sub> to total greenhouse gas emission from fertilization: results from a sandy soil fertilized with nitrate and biogas digestate with and without nitrification inhibitor, *Nutrient cycling in agroecosystems*, 100, 121-134, 2014.
- 1045 Yin, X., Kersebaum, K. C., Kollas, C., Manevski, K., Baby, S., Beaudoin, N., Öztürk, I., Gaiser, T., Wu, L., and Hoffmann, M.: Performance of process-based models for simulation of grain N in crop rotations across Europe, *Agricultural Systems*, 154, 63-77, 2017.
- Zambrano-Bigiarini, M.: hydroGOF: Goodness-of-fit functions for comparison of simulated and observed hydrological time series, 2020.
- Zhang, Y., Liu, Q., Zhang, W., Wang, X., Mao, R., Tigabu, M., and Ma, X.: Linkage of aggregate formation, aggregate-associated C distribution, and microorganisms in two different-textured ultisols: A short-term incubation experiment, *Geoderma*, 394, 114979, <https://doi.org/10.1016/j.geoderma.2021.114979>, 2021.
- 1050 Zhao, F., Wu, Y., Hui, J., Sivakumar, B., Meng, X., and Liu, S.: Projected soil organic carbon loss in response to climate warming and soil water content in a loess watershed, *Carbon Balance and Management*, 16, 24, 10.1186/s13021-021-00187-2, 2021.