

# Shallow and Deep Convection Characteristics in the Greater Houston, Texas Area Using Cell Tracking Methodology

Kristofer S. Tuftedal<sup>1</sup>, Bernat Puigdomènech Treserras<sup>2</sup>, Mariko Oue<sup>1</sup>, Pavlos Kollias<sup>1,3</sup>

<sup>1</sup>Division of Atmospheric Sciences, Stony Brook University, Stony Brook, NY, USA

<sup>2</sup>Department of Atmospheric and Oceanic Sciences, McGill University, Montréal, QC, Canada

<sup>3</sup>Center for Multiscale Applied Sensing, Brookhaven National Laboratory, Upton, NY, USA

*Correspondence to:* Kristofer S. Tuftedal (*kristofer.tuftedal@stonybrook.edu*)

**Abstract.** The convective lifecycle, from initiation to maturity and dissipation, is driven by a combination of kinematic, thermodynamic, microphysical, and radiative processes that are strongly coupled and variable in time and space. Weather radars have been traditionally used to provide various convective clouds characteristics. Here, we analyzed climatological convective cell radar characteristics to obtain and assess the diurnal cycle of three convective cell types: shallow, modest deep, and vigorous deep convective cells, that formed in the Greater Houston area, using the National Weather Service radar from Houston, Texas and a multi-cell identification and tracking algorithm. The examined dataset spans over four years (2018–2021) and for the warm season months (June to September). The analysis showed the clear diurnal cycles in cell initiation (CI) consistent with the sea breeze circulation, and diurnal and normalized lifetime relationships in cell evolution parameters (e.g., maximum reflectivity, echo top height, GOES-16 Channel 13 brightness temperature, and the height of maximum reflectivity). The cell evolution is well represented by relationships between 1) the height and value of the maximum radar reflectivity, 2) the minimum GOES-16 Channel 13 brightness temperature and the maximum vertically integrated liquid, 3) the maximum reflectivity and columnar average reflectivity, and 4) echo top ascent rate and cell lifetime. The relationships presented herein help to identify the cell lifecycle stages such as early shallow convection, vigorous vertical development, anvil development, and convective core dissipation. GOES-16 Aerosol Optical Depth values are also used as a proxy for cell initiation aerosol concentrations to investigate any potential relationships between initiation location and aerosol concentration. Overall, no significant relationships were found for the three cell types investigated between initiation location and aerosol concentration, but some minor differences exist in the pre-CI aerosol optical depth for vigorous deep convective cells.

## 1 Introduction

Convection is one of the most important contributors to the Earth's climate system through its transport of heat, moisture, and momentum. These processes strongly depend on cloud evolution. While an ordinal convective cloud model is proposed, the cloud evolution may vary depending on environment, diurnal cycle, etc. (e.g. Bony et al. 2015; Fridlind et al. 2017; Ladino et al. 2017; Colin and Sherwood 2021). These variabilities may cause large uncertainties in convective parameterizations in numerical climate and cloud models. Modeling studies have attempted to answer some of these outstanding questions (Lee et al. 2008; Zhu et al. 2012; Varble et al 2014; Igel et al. 2015; Peters et al. 2019), but the general lack of high-quality observational data to compare with these modelling studies makes it difficult to assess the validity of the results presented therein. Previous modeling studies tend to focus only on comparisons with observed reflectivity, which can be a rather convoluted to use for microphysical analyses. However, recent work in simulating polarimetric variables using forward operators (Ryzhkov et al. 2011; Wolfensberger and Berne 2018; Kumjian et al. 2019) may provide better insight into the changes in storm microphysics through comparisons with observed polarimetry.

The ongoing debate in the literature about warm- and cold-phase convective invigoration (i.e., whether aerosol particles invigorate convection through latent heating caused by condensation or freezing, respectively; Sheffield et al. 2015; Fan et al. 2018; Abbott and Cronin 2021; Igel and van den Heever 2021; Grabowski and Morrison 2021) is one such example of the need for high-quality observational datasets to compare with modeling studies. While model-based studies may provide some context to the physical processes ongoing within convection, their model intercomparisons also show inconsistency among the models and microphysical parameterizations. The collection of a robust observational dataset that can be used as observational benchmark for modeling studies is challenging. Such a dataset requires a sufficient large sample size of convective clouds properties over a wide range of meteorological and aerosol conditions and convective cell centric methodologies rather than domain averaging approaches that fail to capture the lifecycle of convective clouds.

To address some of the observational shortcomings and needs, a diverse, interagency, coordinated effort took place in the greater Houston, Texas metropolitan area and surrounding region from the summer of 2021 through the summer of 2022 to collect a comprehensive dataset of isolated convective cells (Jensen et al. 2022). These efforts include the Tracking Aerosol Convection interactions Experiment (TRACER) supported by the United States

Department of Energy (DOE) Atmospheric Radiation Measurement (ARM) facility and the National Science  
55 Foundation (NSF) Experiment of Sea Breeze Convection, Aerosols, Precipitation and Environment (ESCAPE)  
campaign. The study region was selected because it is warm and humid in the summer, commonly experiences onshore  
flow and sea breeze-forced convection, has variable aerosol concentrations along the Gulf Coast. Recent studies on  
convective cells that form in coastal regions have illustrated that these cells are less influenced by the synoptic-scale  
meteorological conditions than cells that exist in only maritime or continental environments (e.g. Bergemann and  
60 Jakob 2016; Birch et al. 2016). The land-sea breeze circulation that develops in these regions has been shown to have  
a greater effect on local convection by acting as a forcing mechanism for convective initiation (CI) through an increase  
in surface convergence (Haurwitz 1947; Rotunno 1983). Study of convective cells that form under land-sea breeze  
circulation forced CI can be used to more directly attribute convective characteristics to ongoing convective processes  
rather than the synoptic-scale meteorological conditions present. This lack of dependence on the larger scale  
65 meteorology allows for more generalizable conclusions to be drawn for any convective characteristics presented  
herein.

Previous studies of long-term remote sensing data collected in Houston suggest that variability in convective  
cloud microstructure, hydrometeor properties, and electrification is correlated to variability in aerosol conditions over  
and downwind of Houston (Hu et al. 2019a, b). In these studies, NEXRAD radar observations were used to track  
70 convective cells during different cloud condensation nuclei (CCN) conditions (satellite-retrieved; Rosenfeld et al.  
2016), and to investigate how variability in cloud, precipitation and lightning characteristics related to CCN  
conditions, though neglected proper control of meteorology. Also, a recent pilot study has identified the need to collect  
observations of convective clouds at a temporal and spatial resolution higher than is possible from the existing  
operational observations (Fridlind et al. 2019). Those studies show large variabilities of convective cloud properties  
75 associated with cloud lifecycles. In addition, cloud lifecycles also vary from one individual cloud to another. These  
substantial variabilities make it prudent to analyze very large datasets worth of convective clouds to obtain more  
representative bulk statistics of these phenomena. This lack of generalizable data also hinders our ability to evaluate  
cloud model simulations.

Here, we add and expand on these previous studies using a similar methodology, but focusing on  
80 characteristics of both shallow and deep convection and utilizing an extensive convective-cell centric methodology

for the different types of isolated convection observed around the Houston metropolitan and surrounding areas. The climatologies of the observed characteristics of shallow and deep convection are derived from the National Weather Service (NWS) KHGX Weather Surveillance Radar–1988 Doppler (WSR-88D; Crum and Alberty 1993; Radar Operations Center 2022) and the Geostationary Operational Environmental Satellite-16 (GOES-16). Strict thresholding of the radar and satellite derived characteristics of tracked convective cells allows for the analysis of the behaviors of each case type (i.e., shallow, modest deep, and vigorous deep convection) distinct from one another. Sensitivity testing is also performed to determine how much the results presented herein vary with small variations in threshold selection. Section 2 outlines the data, threshold variables, and analysis methods used, section 3 presents the analyses of these convective types, and section 4 discusses the observations and results.

90

## **2 Data and Methods**

### **2.1 Domain and Data**

Our study domain was centered on the KHGX WSR-88D near Houston, Texas and extended  $\pm 125$  km to the north, east, south, and west, creating a 250 km by 250 km domain. This region was selected because the coastline with the Gulf of Mexico generates regular land-sea breeze circulations which provide a forcing mechanism for CI. This domain also provides regions of relatively “pristine” and “polluted” aerosol regimes. For example, the area to the northeast of Houston is generally downstream of pollution sources, such as the Houston metropolitan area itself and oil refineries near Houston allowing for the advection of this polluted air over this sub-region. The area to the southwest of Houston is generally upstream and made up of mostly rural land even further upstream, allowing the air here to be far less polluted. However, the general flow pattern described does not hold true for all cases (i.e., easterly flow at the surface), and are not accounted for. As such, flow patterns that deviate substantially from the general flow will introduce considerable error to any aerosol concentration relationships presented herein.

Data from KHGX and GOES-16 were collected for the period of June through September 2018–2021 where convective initiation occurred during local daytime (09:00–21:00 Central Daylight Time (CDT)). Initiation during local daytime was chosen to increase the likelihood that sea breeze propagation was a primary forcing mechanism for CI and to enable the analysis of GOES aerosol optical depth (AOD) data.

105

During the study period, the volume coverage patterns (VCPs) available to KHGX changed with the implementation of VCP 215 (15 elevation angles from 0.5 and 19.5°) and 112 (14 elevation angles from 0.5 and 19.5°) and the removal of VCPs 11, 211, 21, and 221 (Radar Operations Center 2015; Zittel 2019). Aside from these differing VCPs, different dynamic scanning techniques can adjust how the radar scans without changing the VCP (Chrisman 2009; 2013; 2014; 2016). These differing VCPs and dynamic scanning techniques have an effect on the time required to complete a single VCP. Despite these variations in rescan time, the time required to complete any of these VCPs is on the order of approximately five minutes. These different VCPs also collect slightly different sets of elevation data, which can substantially affect the uncertainty in any height or vertically integrated liquid (VIL; Greene and Clark 1972; Amburn and Wolf 1997) analyses presented herein.

Horizontal radar reflectivity factor ( $Z_H$ ) from KHGX was collected during normal weather surveillance operations and regridded to a 250 km by 250 km domain with 500 m by 500 m horizontal spacing for each case day considered, which was then used to estimate VIL, radar-derived echo top height (ETH), and radar-derived profile depth ( $H_{cell}$ ). GOES-16 Channel 13 cloud top brightness temperature data (GOESBT) were regridded to the same grid as before and analyzed such that each 5-minute image was linked to the KHGX scan time nearest to each GOESBT product time. Since KHGX and GOES-16 are not scanning at the exact same time, some error may be introduced by the difference between product times. However, since both KHGX and GOES-16 scan at approximately 5-minute intervals, the maximum difference between product times is approximately 2.5 minutes. GOES-16 AOD calculation requires cloud-free pixels to generate AOD data. In many cases, the location of cell initiation already contains some form of cloud cover (be it low-, middle-, or upper-level clouds) at the time of initiation, which inhibits AOD generation. When AOD values are generated, they are classified as low-, medium-, or high-quality returns. To ensure a large enough sample size, AOD data denoted as medium- or high-quality were temporally averaged for the 30 minutes preceding cell initiation at the location of cell initiation. We calculated VIL from gridded  $Z_H$  (with  $Z_H$  capped at 56 dBZ, as done in Greene and Clark 1972) data from KHGX using equation (1):

$$VIL (dB) = 10 * \log_{10} \left( \frac{\sum_{i=0}^{i=(i_{max}-1)} 3.44 * 10^{-6} [(Z_i + Z_{i+1})/2]^{4/7} dh}{1 \text{ kg/m}^2} \right) \quad (1)$$

where  $i_{\max}$  is the total number of sweeps for a given VCP,  $Z$  is the radar reflectivity of a given sweep ( $i$ ) in units of  $\text{mm}^6 \text{m}^{-3}$ , and  $dh$  is the vertical spacing between sweeps. VIL assumes a Marshall-Palmer drop-size distribution, which introduces uncertainty since this relationship may not be completely valid for convective events in coastal regions.

135 Other sources of uncertainty in VIL include beam spreading with increasing distance from the radar and data sparsity at high elevations within 30 km of KHGX. The uncertainty because of beam spreading with distance has been shown to be relatively small, but that VIL within 30 km of the radar tends to be underestimated for 5-minute WSR-88D VCPs (Oue et al. 2022).

## 2.2 Cell Tracking

140 KHGX VIL and GOES-16 observations were used as input to a modified version of the multi-cell identification and tracking (MCIT) algorithm (Rosenfeld 1987; Gagin et al. 1985; Hu et al. 2019a) to track all features with VIL of  $\geq -20$  dB during that period. The specifics of the MCIT algorithm can be found in section 2.3 to 2.4.2 in Hu et al. 2019a and the modifications made to the algorithm are provided in detail in Lamer et al. (2023). The MCIT algorithm ingests time series of volume scans and tracks local maxima of VIL by identifying the two cells in  
145 consecutive radar scans that have common maximum VIL (i.e. the areas of VIL have some overlap from one time to the next.) To check for overlap from timestep to the next, the pySTEPS (Pulkkinen et al. 2019a,b) is used to compute velocity fields to find the shift in images. The current image is then checked with the advected cell from the computed velocity field and checks for overlap between the actual cell and computed cell location. This tracking continues until the cell dies, splits, or merges. In addition to the KHGX data, we used GOES-16 Channel 13 cloud top brightness  
150 temperature (GOESBT) to identify cold topped and warm topped cells. The initial analysis from the modified MCIT algorithm identified 1,664,215 features with a VIL value  $\geq -20$  dB during the analysis period.

## 2.3 Cell Classification

To better characterize evolution of each cell, we employed the following cloud properties:

- 1) ETH: the height at the middle of the highest gate with detectible signal ( $Z_H > -10$  dBZ) from KHGX;
- 155 2)  $H_{\text{cell}}$ : the depth between the top of the highest gate of the radar detectible signal for a cell and the bottom of the lowest gate of the radar detectible signal (not corrected for increasing altitude with increasing range from the radar).

- 3) ERatio (Fig. 1): the ratio of the ETH to  $H_{\text{cell}}$ ; and
- 4) the tracked cell area based on VIL (Area)

160 Using these properties, GOESBT, and VIL, we classified tracked features into three categories: 1) shallow, 2) modest deep, and 3) vigorous deep convective cells. Thresholds for the classification are listed in Table 1. These thresholds were empirically derived to avoid false classification of cloud systems such as high cirrus clouds, mesoscale convective systems (MCSs), or large regions of stratiform precipitation. We also used the initial cluster fraction of the cell, which is the ratio of the area of a given cell to the area of a cluster (parent) of the cell at the beginning of the cell lifetime, equal to 1 (a value of 1 meaning the cell is discrete). The split/merged cells are removed in this study using 165 the process outlined in Lamer et al. (2023). The area threshold is used only to classify the shallow cells to ensure that these shallow features are not large regions of stratiform precipitation. This extensive thresholding removes noisy features, likely due to non-meteorological echoes (e.g., ground clutter, insects, etc.). Cells that have their “birth” or “death” occur on a domain edge are removed to avoid analyzing cases which are ongoing before or after entering or 170 leaving the study domain. Cells which pass within 15 km of KHGX are also removed to mitigate the lack of data in the “cone of silence.”

The shallow and deep convective cells selected based on these thresholds were then analyzed separately and compared with one another. Sensitivity studies were also performed on each case type by varying all thresholds (except initial cluster fraction and split/merge status) by  $\pm 5\%$  individually and simultaneously to observe any changes in the 175 distributions of certain variables for each case type (See Appendix).

## 2.4 Climatological Analyses and Statistical Analysis

Cell properties introduced in Sections 2.1 and 2.2 for all cells classified into the three categories during the four-year observation period were examined to allow for bulk analyses of cell type characteristics. Analyses used 180 herein include observations during specific times during cells’ lifetimes (such as the time of initiation), aggregates of all scans from the entire life of all cells of a given type, changes in variables over cell normalized lifetimes, diurnal changes in these variables, and spatial differences in initiation location. These analysis types enabled us to investigate how these cells changed as they grew, matured, and decayed, as well as allowed for the direct comparison of how

these case types differ from each other. To parse out any potential aerosol effects on shallow and deep CI and intensity  
185 GOES-16 AOD data were analyzed. The nonparametric Mann-Whitney U test was used to investigate statistical  
differences between regional cell initiation AOD distributions (Mann and Whitney 1947).

### 3 Results

#### 3.1 Overall Cell Characteristics

The monthly average number of cells varies little from month to month for June, July, August, and  
190 September, is highest in August, and has no significant difference between months. The storm motion could be  
important to understand storm evolution. To evaluate the primary speed and direction of cell motion, Figure 3 shows  
frequency distributions of the propagation speed and direction as a function of normalized lifetime for the three cell  
types investigated. The frequencies shown are normalized by the total number of samples at each normalized lifetime  
bin (every 0.025). All cell types tend to have storm motions ranging primarily between southerly to easterly. Shallow  
195 convective cells, overall, move the slowest out of the three cell types and have less variability in speed than modest  
and vigorous deep convective cell. Over the lifetime of these cells, shallow convective cell speed varies little and is  
much slower when compared to deep convection. Overall, for the three categories, the convective cells mainly tend to  
have storm motions spanning from southwesterly to easterly. Larger variability in storm motion is found at the later  
period of the cell lifetime for modest and vigorous deep cells, where the greatest variability is seen in the vigorous  
200 deep cells. This finding suggests that some of the vigorous deep convective cells may be supercellular in nature  
because of the large deviations from the early storm motions near the time of initiation, but cannot be confirmed  
without further analyses of detailed conditions for supercells outside of the scope of this research. The direction of  
motion of sea-breeze induced convective cells can depend on the low-level (i.e., 925 hPa) wind direction relative to  
the coastline (Die Wang, 2023, personal communication). The direction of cell motion (particularly early in the  
205 lifetimes of these cells) indicates the possibility that the sea breeze along the Gulf Coast plays a part in storm initiation  
and propagation as these directions would be consistent with the inland propagation of the sea-breeze, which  
propagates from southeast to northwest from the coastline.

#### 3.2 Location and Diurnal Frequency of Initiation



Figure 4 shows the diurnal frequency of initiation times as a function of local time. All cell types have their  
210 peak in initiation in the late morning/early afternoon hours, which then sharply decreases as the day progresses.  
Overall, all types of convection in this region preferentially initiate in the late morning/early afternoon. However,  
there is a slight difference in peak time. Shallow convection and vigorous deep convection show earlier peaks in  
initiation time – between 10:00 and 14:00 CDT – when compared to the peak for modest deep convection, which is  
shifted slightly later (between 12:00 and 15:00; Fig. 4).

215 Figure 5 shows the initiation location for shallow (Fig. 5a,d,e), modest deep (Fig. 5b,e,h), and vigorous deep  
(Fig. 5c,f,i) convection. The three types all show a preference to initiate over land along the coastline to the southwest  
and northeast of Houston (within 100 km from the coast) with a local minimum in initiation over Galveston Bay (the  
large bay to the east of Houston), suggesting that a land-sea contrast, hence sea breeze, is a key factor to initiate  
convection in this area. The inland propagation of the sea breeze can also be observed based on the cells' overall speed  
220 and direction of travel early in their lifetimes (Fig. 3). One feature of note is the obvious preference in shallow CI to  
the southwest of Houston (Fig. 5a,d,e). Another notable feature is that both moderate and vigorous deep convective  
cell types show a local maximum over the Houston metropolitan area, which is especially apparent in vigorous deep  
convective cases. This initiation maximum could be caused by the enhancement from aerosol loading and/or urban  
heating, but will need further examination in the future. There is an obvious inland shift in initiation location, that is  
225 most obvious for vigorous deep convection, from the 09:00-11:59 to 12:00-14:59 CDT period, further supporting the  
influence of the sea breeze on CI for all case types. The diurnal spatial shift in initiation location shows that the sea  
breeze is a key factor for CI along the coast. Figures 4 and 5 highlight that the early afternoon hours are the most  
preferable time for CI for all cell types. This is consistent with the previous studies (e.g., Park et al. 2020).

The domain maximum in shallow CI to the southwest of Houston is also apparent in all periods except 18:00-  
230 20:59 CDT (not shown), but modest and vigorous deep convective types do not visually show the same southwest  
initiation preference. Rather, the two deep convective types (especially vigorous deep convection) do show maximums  
in initiation over the Houston metropolitan area in the mid-to-late afternoon. We further speculate that this may be the  
result of urban heating allowing parcels to reach their convective temperatures during this time, even on days where  
CAPE is high, but CIN elsewhere in the domain is too strong to allow for other convection to initiate. However, as

235 stated previously, further research is necessary to elucidate what mechanism(s) is(are) responsible for this local initiation maximum in deep convection.

### 3.3 Diurnal Changes in Cell Characteristics

To assess diurnal cycle of convective cells, we analyzed cell hour-by-hour basis characteristics for each scan for the duration of its life over the course of the day. Figure 6 shows the diurnal trends in the distributions of GOESBT, 240 the height of cell maximum radar reflectivity ( $H_{dBZ_{max}}$ ), and the cell maximum radar reflectivity ( $dBZ_{max}$ ) for all scans of all convective cases. For shallow cells, the GOESBT tends to be constant at around 285 K over time, and the majority of the cells have  $H_{dBZ_{max}}$ . In contrast, the two deep convection types exhibit similar maximum frequencies of GOESBT before noon with a substantial shift of the maximum frequency to  $\sim 240$  K for modest deep cells and  $\sim 220$  K for the vigorous deep cells around noon, then taper off into the later evening and overnight hours. The frequency 245 peak of  $H_{dBZ_{max}}$  for the modest and vigorous convective types is found below 6 km before noon, similar to shallow cells, which then shifts to 6–10 km for modest deep cells and 9–13 km for the vigorous deep cells until  $\sim 18:00$  CDT for modest deep cells and 23:00 CDT for vigorous deep cells. ETH also shows the same change in characteristics as GOESBT and  $H_{dBZ_{max}}$  (not shown), further supporting late morning/early afternoon initiation. There is little dependency of maximum reflectivity on time.

### 250 3.4 Analysis of Bulk Cell Type and Normalized Cell Lifetime Characteristics

Even though the peak in CI has been shown to occur in the late morning/early afternoon, and peak in intensity in the early-to-mid afternoon, there may be diurnal variability and differences in duration in cell lifetime that are unaccounted for. To observe how these cells behave over the course of their respective lifetimes rather than the course of a given day, we have analyzed these cells by aggregating scans during specific periods of cell lifetimes, aggregating 255 all scans of all cells of each type, and by normalizing each by its respective lifetime. These analyses allow for direct comparisons of case types, regardless of what time the cells initiated or how long they lasted.

The growth and decay of these cells can be seen by analyzing the change in the distributions of  $dBZ_{max}$ ,  $H_{dBZ_{max}}$ , maximum VIL, GOESBT, and the average of the maximum radar reflectivity for all columns within a given cell ( $dBZ_{avg}$ ). Shallow convective cells show overall low  $dBZ_{max}$  at low  $H_{dBZ_{max}}$  over the course of entire cell lifetimes 260 (Fig. 7). Cell growth and decay is apparent, as  $H_{dBZ_{max}}$  and  $dBZ_{max}$  shift to overall higher values during the first half

of cell lifetimes and then decrease back to a distribution that looks most similar to cell initiation at the end of their lives. Modest and vigorous deep convection show clear signals of the birth (Fig. 7b,c), mature (Fig. 7e,f,h,i), and dissipation (Fig. 7k,l) phases. Early in their lifetimes, these cell types are dominated by low  $\text{dBZ}_{\text{max}}$  values at low  $H_{\text{dBZ}_{\text{max}}}$ , which both increase considerably moving into their mature phases. This trend is especially apparent in vigorous deep convection, where early in cell lifetimes, there is a clear signal of initiation and some cells that have begun to grow (based on the secondary maximum in high  $\text{dBZ}_{\text{max}}$  at high  $H_{\text{dBZ}_{\text{max}}}$  values). As these cells continue to mature, we see that  $H_{\text{dBZ}_{\text{max}}}$  remains high, but that  $\text{dBZ}_{\text{max}}$  begins to decrease. This is indicative of cells where the convective core has dissipated leaving “orphan anvils” (Hitschfeld 1960). However,  $H_{\text{dBZ}_{\text{max}}}$  appears to be slightly skewed by bright banding, as both modest and vigorous deep convection show rather unnatural looking discontinuities in their distributions around 6 km above radar level. These discontinuity features are especially apparent in Figure 8b,e.

Figure 8 supports that the feature seen in Figure 7 around 6 km above radar level to be caused by the bright band, since there are no similar discontinuities in these distributions where GOESBT is warmer. However, this is one interpretation and the discontinuities shown in Figure 7 may rather be indicative of some ongoing storm physics. As with Figure 7, Figure 8 clearly shows the initiation, growth, and decay of these cell types. Shallow convective cells remain at relatively warm GOESBT and low maximum VIL throughout their lives, whereas modest and vigorous deep convection show clear growth from warm GOESBT, low maximum VIL cells to cold GOESBT high maximum VIL cells. The rapid changes shown in Figures 7 and 8 during the early stages of vigorous deep convective shows the quick vertical evolution of these cells. Also shown as in Figure 7, is the dissipation phase, where GOESBT remains cold, but VIL drops off considerably. This further supports the idea that this is the period when the convective cores have dissipated, leaving orphan anvils behind.

As stated previously,  $\text{dBZ}_{\text{avg}}$  is the average of the column  $Z_{\text{H}}$  maximums within a given cell and  $\text{dBZ}_{\text{max}}$  is the cell maximum  $Z_{\text{H}}$  for a given time. To investigate potential anvil generation, the distributions of  $\text{dBZ}_{\text{avg}}$  versus  $\text{dBZ}_{\text{max}}$  are analyzed (Fig. 9). During the anvil generation phase, we would expect cells to be at their most intense state, meaning they would have a high value of  $\text{dBZ}_{\text{max}}$ . We would also expect an increasingly large region of low  $Z_{\text{H}}$  high aloft in the storm to form. This region of low  $Z_{\text{H}}$  will skew  $\text{dBZ}_{\text{avg}}$  lower while  $\text{dBZ}_{\text{max}}$  will remain unaffected until the storm begins to dissipate. When comparing shallow and deep convection with this information in mind, the feature, apparent in deep convection, of low  $\text{dBZ}_{\text{avg}}$  and high  $\text{dBZ}_{\text{max}}$  could be interpreted as anvil generation. As with

previous figures in this section, these anvil signatures are especially apparent in vigorous deep convective cells. Early  
290 in deep convective lifetimes, cells have not had enough time to grow to a state where anvil generation is possible (with  
the exception of the most intense cells). The vast majority of cells exist very near the one-to-one line of  $\text{dBZ}_{\text{avg}}$  to  
 $\text{dBZ}_{\text{max}}$  with only a small subset of cells moving into the high  $\text{dBZ}_{\text{max}}$  and lower  $\text{dBZ}_{\text{avg}}$  indicative of cells growing  
and beginning to produce anvils (Fig. 9c). As cells move into the mature portion of their lives, the low  $\text{dBZ}_{\text{avg}}$ /high  
 $\text{dBZ}_{\text{max}}$  region, where strong, anvil generating cells are present, becomes the dominant regime for these cells (Fig.  
295 9f,i). As cells decay, the distribution maximum shifts back down near the one-to-one line (Fig. 9l), which would be  
suggestive of orphan anvils, given the analyses provided in the previous paragraphs. One feature of note in the  
vigorous deep convective cells during the dissipation phase is the secondary maximum in the high  $\text{dBZ}_{\text{avg}}$ /high  $\text{dBZ}_{\text{max}}$ .  
This feature represents the transition from the mature stage having the large  $\text{dBZ}_{\text{max}}$  in a deeper depth within the clouds  
to the later stage having lower  $\text{dBZ}_{\text{max}}$  during the normalized lifetime 0.75 – 1.0.

300 Shallow convective cells reach their maximum ETH during approximately the first 10% of their life, appear  
to maintain this height until around 75%, and then gradually decrease in height until dissipation (Fig. 10a). Modest  
convective cells take substantially longer (~35-55% of the way through their life) to reach their maximum ETH and  
then sharply decrease in height during the last 15% (Fig. 10b). Vigorous convective cells, on the contrary, reach their  
maximum ETH quite quickly (within the first 15 to 35% of their life), maintain a tall echo top, and then gradually  
305 decrease in height to dissipation (Fig. 10c). The gradual decrease in ETH leading to dissipation is further indicative  
of convective anvils. In this case, we believe this signal further supports that we are observing convective core  
dissipation, thus leaving orphan anvils behind. Both  $H_{\text{dBZ}_{\text{max}}}$  and  $\text{dBZ}_{\text{max}}$  show similar trends to ETH for all case types.  
Shallow convective cells reach their  $\text{dBZ}_{\text{max}}$  and  $H_{\text{dBZ}_{\text{max}}}$  early in the cells' normalized lifetime (within the first 20-  
30%), maintain, and then gradually decay to dissipation (Fig. 10d,g). Modest convection shows a gradual increase in  
310  $H_{\text{dBZ}_{\text{max}}}$  that looks nearly identical to its maximum ETH. It appears that these cells reach their  $H_{\text{dBZ}_{\text{max}}}$  anywhere from  
35 to 65% of the way through their lifetimes. However,  $\text{dBZ}_{\text{max}}$  is reached much earlier (within the first 20% of their  
life) than  $H_{\text{dBZ}_{\text{max}}}$  (Fig. 10e,h). Vigorous deep convection reaches its  $H_{\text{dBZ}_{\text{max}}}$  much sooner in its normalized lifetime  
(between 10 and 25% of their normalized lives), like ETH (Fig. 10f). As with modest deep convection, vigorous deep  
convection also reaches its  $\text{dBZ}_{\text{max}}$  early in its life (within the first 20% of their normalized life) and maintains these  
315 values until about halfway through their lives, when dissipation begins (Fig. 10i). One feature that is seen in  $\text{dBZ}_{\text{max}}$   
and  $H_{\text{dBZ}_{\text{max}}}$  for both modest and vigorous deep convection is the apparent bimodality at the later of these cells'

lifetimes ( $> \sim 0.4$ ), where it appears that some cells maintain high values of  $\text{dBZ}_{\text{max}}$  and VIL (not shown) at the higher altitudes ( $\sim 9$  km) all the way up to dissipation (Fig. 10f,i). This is particularly apparent in the modest deep cells. This bimodality suggests that some of the modest deep cells with lower ETHs were dominated by warm phase precipitation processes similar to the shallow cells, as  $H_{\text{dBZ}_{\text{max}}}$  was found below 6 km through the lifetime. This reasoning would explain why  $\text{dBZ}_{\text{max}}$  and VIL remain so high all the way up to the end of their life, as they are deemed “dissipated” when leaving the domain, even though these cells may still persist for some time.

Cell echo base height ( $H_{\text{EBase}}$ ) was determined using the maximum ETH minus the radar-derived echo profile depth to estimate the precipitation base of these cells. For shallow convection,  $H_{\text{EBase}}$  remains relatively flat for the duration of these cells, whereas it shows an increase in height during dissipation for modest and vigorous deep convective cells (Fig. 10j,k,l). This signature is especially apparent in the vigorous deep convective cells. This increase in  $H_{\text{EBase}}$ , in tandem with the minimal decrease in maximum ETH during dissipation of both types of deep convection (especially vigorous deep convection), reinforces the idea that we are observing the generation of orphan anvils. The bimodality here may also partially be caused by some cells leaving the domain during their mature phase, prior to dissipation despite the mitigation techniques we applied, but this is less certain than for the reasoning given for  $\text{dBZ}_{\text{max}}$ .

In an attempt to quantify vertical motion within vigorous deep convective cells, we approximated this quantity three different ways. We used the maximum ETH, and  $H_{\text{dBZ}_{\text{max}}}$  during each scan, and GOESBT at the time closest to radar scan time with the time between samples to calculate maximum ETH and  $H_{\text{dBZ}_{\text{max}}}$  “ascent rates,” as well as GOESBT “cloud top lapse rates.” As shown in Figure 11, there are maximum ETH and  $H_{\text{dBZ}_{\text{max}}}$  ascent signatures early and descent signatures late in the lives of these cells for all case types. GOESBT shows cooling during the same period where ascent is seen and warming during the same period where descent is seen in maximum ETH and  $H_{\text{dBZ}_{\text{max}}}$ . The near identical timing and structure of these derived ascent rates suggest that these values may be good proxies for updraft intensity during the early parts of these cell. However, the later portions of these cells’ lives are dominated by the tops or high upper portions of these features (as shown in Figure 11) and most likely do not represent downdraft intensity during dissipation. We suspect that, in vigorous deep convective cases, we may be seeing orphan anvils falling out as virga since the descent signature occurs quite late in these cells’ normalized lifetimes after their  $H_{\text{EBase}}$  increases considerably. Maximum ETH and  $H_{\text{dBZ}_{\text{max}}}$  descent rates and GOESBT warming rates match well late in these cells’ lives, which further supports the idea that we may be observing anvil fall-out. Further analysis is

necessary to validate these results. There is a hint of this signature as well for modest deep convective cells, but it is  
345 not as obvious. To further assess the quality of these approximations, analyses outside the scope of this study will be  
needed. One caveat to consider with these ascent rate results is the increasing uncertainty in ETH and  $H_{dBZ_{max}}$  with  
increasing range from KHGX. This uncertainty may account for the large spread always shown since we have large  
sample sizes for all case types. The large uncertainty for cells in the farthest portions of our domain muddle these  
results considerably. However, since the same signal is seen when looking at the GOESBT cloud top lapse rates, more  
350 credence is led to the above analyses.

### 3.5 Aerosol Effect Analysis

One potential ingredient that may influence CI and has been a key point of debate in recent literature is  
aerosol loading. Aerosol particles in this region range from natural particles, such as sea spray, to industrial pollutants  
355 from oil refineries and urban sources. However, no such dataset exists which tracks individual aerosol species on  
the spatiotemporal scale needed for direct comparison with MCIT output. As such, we used temporal averages of  
GOES-16 AOD at the locations of cell initiation for the 30 minutes prior to MCIT detected initiation. We analyzed  
these data for regions to the southwest and northeast of Houston. The analyses of AOD for shallow and modest deep  
(Fig. 12a,b) cells at the locations of their initiation show essentially identical distributions from southwest to northeast;  
360 the primary difference being that both shallow and modest deep cells exhibit a longer tail toward higher AOD values  
to the southwest. Median (mean) AOD for shallow convective cell initiation locations were 0.338 (0.400) to the  
southwest and 0.335 (0.366) to the northeast, and 0.366 (0.453) to the southwest and 0.346 (0.384) to the northeast  
for modest deep convection. Considering the substantial difference in initiation location for shallow convection and  
nearly identical initiation AOD distributions, this result can be interpreted that bulk AOD does not play an important  
365 role in controlling cell initiation. However, this does not mean that aerosol particles play no role as a control on cell  
initiation. Rather, it may be specific species of aerosol particles that are more or less important to these processes.  
Vigorous deep convective cells, however, do show differing distributions from southwest to northeast (Fig. 19c). As  
with shallow and modest deep convection, vigorous deep convection also exhibits a longer tail extending to higher  
values of AOD to the southwest. The initiation locations for vigorous deep convective cells tend to have marginally  
370 higher AOD values to the southwest. The median (mean) of the AOD distributions for vigorous deep convective cell

initiation locations was 0.428 (0.513) to the southwest and 0.373 (0.396) to the northeast. However, these differences in distributions are not statistically significant. From the previous area-normalized cell initiation, we speculate that the marginally higher values of AOD and rates of cell initiation to the southwest in vigorous deep convection suggest that aerosol loading may indeed factor into vigorous deep CI, but that more marginal convective cells are either more dependent on specific species of aerosol particles rather than overall aerosol loading or are not as affected by aerosol loading overall.

#### 4 Summary and Discussion

The climatological characteristics of convective cell evolution and their diurnal cycles were analyzed using the NWS KHGX WSR-88D from Houston, Texas for the warm season months (June to September) from 2018 and 2021 and a modified version of the MCIT algorithm. In total, this study analyzed 48,773 convective cells (35,974 shallow cells for 142,921 volume scans, 7,930 modest deep cells for 92,770 volume scans, and 4,869 vigorous deep cells for 95,219 volume scans). Analysis of these case types together allowed for the direct comparison of cell characteristics and the environments in which they form. The key findings from this study are:

1. CI for all cell types occurs most frequently in the late morning/early afternoon over land, consistent with the inland incursion of the sea breeze front.
2. There is a spatial variability in CI for shallow, modest, and vigorous deep convective cells, suggesting some effects of aerosol loading and/or urban heating. This is particularly clear for shallow cells. Higher AOD values tend to be present to the southwest of Houston, which is most easily observed when looking at the vigorous deep CI AOD distributions. While these results are not statistically significant, they suggest that aerosol loading may have some effect on deep CI. The initiation biases do not appear to be related to overall aerosol loading based on the pre-CI AOD analysis with the exception of vigorous deep cells. Further analysis using high spatiotemporal aerosol and urban heat data will be needed.
3. The modest and vigorous deep convective cells particularly deepen in the afternoon/evening (12:00-21:00 CDT) as the frequency peak of their heights of maximum reflectivity increase to 11 km and that of the brightness temperature decrease to 220 K. The shallow cells do not have clear diurnal variability in those parameters.

4. The cell evolution is well represented by relationships between the following cell properties:
- 1) The height and value of the maximum radar reflectivity: The developing stage (cell lifetime normalized by cell duration  $< 0.75$ ) is well represented by an exponential curve as the  $H_{dBZ_{max}}$  gradually increases from around 2 to 4 km with a maximum reflectivity of  $\sim 50$  dBZ, which then dramatically increases to a height of 12 km for  $dBZ_{max}$  values of 50-60 dBZ. These coincident increases in  $dBZ_{max}$  and  $H_{dBZ_{max}}$  occur in the early stage of cell lifetime (normalized lifetime  $< 0.5$ ). The dissipation stage is represented by a wide distribution of the maximum reflectivity at a high altitude ( $\sim 10$  km) suggesting anvil development and convective core dissipation.
  - 2) The brightness temperature (hence echo top height) and the maximum VIL: The developing stage (normalized lifetime  $< 0.75$ ) is well represented by an exponential curve as the brightness temperature gradually decreases from 290 to 260 K from the maximum VIL until 10 dB then dramatically decreases to 210 K for  $VIL > 10$  dB.
  - 3) The maximum reflectivity and columnar average reflectivity: As the cells begin to develop for all case types, these variables remain near the one-to-one line. However, as deep convective cells (especially vigorous deep convective cells) reach the middle phases of their lifecycles ( $0.25 < \text{normalized lifetime} \leq 0.75$ ), an obvious extension of high  $dBZ_{max}$ /low  $dBZ_{avg}$  becomes apparent. The extension of the distribution during this period further supports the development of anvils. In the remaining 25% of vigorous deep convective cell lifetimes, the distribution of  $dBZ_{max}/dBZ_{avg}$  shifts to low values for both, suggesting convective core dissipation leaving only orphan anvils behind.
5. The ETH ascent rate is slightly more positive (5 m/s) in the early stage (normalized lifetime  $< 0.4$ ) and negative for the later stage (normalized lifetime  $> 0.8$ ) for deep convective cells. Early in these cells' lives, the cell tops grow vertically as they intensify and the rates at which they ascent should be close to the actual updraft intensity. Late in the cells' lives,  $H_{dBZ_{max}}$  remains high aloft as it appears that  $H_{dBZ_{max}}$  remains in the anvil portion of the storm after it reaches maturity. These descent rates are likely to be representative of orphan anvils falling out as virga.



Based on the findings in this study, the analysis techniques presented can identify individual features within  
425 convective cells. Further parsing of these data may allow for the tracking of individual features within cells, such as  
tracking cores and anvils separately and being able to analyze their behaviors over their lives. Polarimetric variables  
can also be added, as in Hu et al. (2019a,b), for automated tracking of features such as differential reflectivity columns  
and arcs. The convective cells considered in this study only constitute about 3% of the features tracked during the  
climatology period and were selected based on the empirically derived thresholds in Table 1. Some of the features we  
430 excluded by using these thresholds are non-meteorological in nature, but others include high clouds and large  
precipitation shields. Different empirically derived thresholds can be developed to isolate these and other features  
from the full dataset and used to create large climatologies of these features of interest. The analysis techniques  
presented in this study can also be applied to the cloud resolving model simulations using radar simulator and cell  
tracking techniques (e.g. Oue et al. 2022). Analyses of this type will aid in better evaluating simulation results to  
435 understand isolated convective cell formation and evolution mechanisms including effects of environmental factors  
such as aerosols. Some such formation mechanisms have been discussed here, such as aerosol loading and urban  
heating, but have not been explored in-depth, but may be of considerable importance in determining what processes  
are most important to convective growth, maintenance, and decay.

#### 440 **Appendix**

To assess the sensitivity of the results in this study, the thresholds were varied individually and  
simultaneously by  $\pm 5\%$  from the values presented in Table 1. The number of cells selected in each sensitivity test  
were recorded and plots were qualitatively analyzed to investigate substantial visual differences as these variables  
were adjusted. Table A1 shows the number of cells selected for each variable adjusted. Figure A1 shows the visual  
445 differences of the distributions based on changes in the most sensitive thresholds.

The number of shallow convective cases varied more for lifetime minimum GOESBT than any other variable.  
The number of cases increased by 16.80% for a five percent reduction in the GOESBT threshold (from 268.0 K to  
254.6 K) and decreased by 48.99% for a five percent increase in the same threshold (from 268.0 K to 281.4 K). The  
visual differences in the distributions of cases are shown in Figure A1a,b,c. As shown, the visual differences in the  
450 distributions are caused by the addition or removal of colder topped shallow convection (Fig. A1c). The overall shape

of the distribution does not change otherwise. Increasing GOESBT only removes the more “intense” shallow convective cells that have colder GOESBT values during their lifetimes.

Like the shallow convective cases, the number of modest deep convection cases varied most with changes in lifetime minimum GOESBT. There was a 27.25% reduction in cases with a five percent decrease in GOESBT (250.0 K to 237.5 K). This removes the slightly warmer GOESBT cloud tops, leaving an upper bound of more intense modest deep convective cases (Fig. A1e). There was also a 24.40% increase in cases with a five percent increase in lifetime minimum GOESBT (250.0 K to 262.5 K). This clearly shows the addition of warmer, lower intensity convective cells (Fig. A1f). The modest convective cells also varied by more than 10% for the upper bound of the lifetime maximum ETH, but there were no visible changes in distributions of these variable, only a reduction/increase in the number of cases (Fig. A1g,h,i).

Vigorous convection varied most by lifetime maximum ETH. The number of cases increased by 28.94% for a five percent decrease in maximum ETH (12 km to 11.4 km) and decreased by 26.95% for a five percent increase in maximum ETH (12 km to 12.6 km). The variability in maximum ETH is shown in Figure A1j,k,l and appears to behave similarly to how the sensitivity in maximum ETH affects modest deep convection. Adjusting this threshold, only appears to reduce or increase the number of cases and not change the shape of the distribution.

### **Author Contribution**

Kristofer S. Tuftedal performed the threshold selection processes, all analyses presented herein, and the bulk of manuscript preparation. Bernat Puigdomènech Treserras wrote the script for the MCIT algorithm and generated the dataset analyzed in this paper. Mariko Oue provided assistance in the interpretation of results and assistance in preparing this manuscript. Pavlos Kollias provided assistance in the interpretation of results, suggestions for analysis techniques, and assistance as a Ph.D. advisor to Kristofer S. Tuftedal.

### **Code Availability**

The MCIT tracking algorithm and analysis codes are available upon request.

## Data Availability

The datasets generated for this study are available upon request. All other datasets used (WSR-88D KHGX and GOES-16) in this study are freely available through the National Centers for Environmental Information.

480

## Competing Interests

The authors declare that they have no conflict of interest.

## Acknowledgements

485 This study was supported by NSF Grant AGS-2019968 (Kollias and Oue) and the U.S. Department of Energy DE-SC0021160 (Oue). The authors would like to thank the four anonymous reviewers and the editor, Zhanqing Li, whose comments helped improve the content and clarity of this manuscript.

## References

490 Abbott, T. H., and T. W. Cronin, 2021: Aerosol invigoration of atmospheric convection through increases in humidity. *Sci.*, **371**, 83–85, doi:10.1126/science.abc5181.

Amburn, S. A., and P. L. Wolf, 1997: VIL Density as a hail indicator. *Wea. Forecasting*, **12**, 473–478, doi:10.1175/1520-0434(1997)012<0473:VDAAH1>2.0.CO;2.

495

Bergemann, M., and C. Jakob, 2016: How important is tropospheric humidity for coastal rainfall in the tropics? *Geophys. Res. Lett.*, **43**, 5860–5868, doi:10.1002/2016GL069255.

Birch, C. E., S. Webster, S. C. Peatman, D. J. Parker, A. J. Matthews, Y. Li, and M. E. E. Hassim, 2016: Scale  
500 interactions between the MJO and the western maritime continent. *J. Climate*, **29**, 2471–2492, doi:10.1175/JCLI-D-  
15-0557.1.

Bony, S., B. Stevens, D. M. W. Frierson, C. Jakob, M. Kageyama, R. Pincus, T. G. Shepherd, S. C. Sherwood, A. P.  
Siebesma, A. H. Sobel, M. Watanabe, and M. J. Webb, 2015: Clouds, circulation, and climate sensitivity. *Nature*  
505 *Geoscience*, **8**, 261–268, doi:10.1038/ngeo2398.

Chrisman, J. N., 2009: Automated Volume Scan Evaluation and Termination (AVSET)—A simple technique to  
achieve faster volume scan updates. 34th Conf. on Radar Meteorology, Williamsburg, VA, Amer. Meteor. Soc., P4.4,  
[https://ams.confex.com/ams/34Radar/techprogram/paper\\_155324.htm](https://ams.confex.com/ams/34Radar/techprogram/paper_155324.htm).

510

Chrisman, J. N., 2013: Dynamic scanning. NEXRAD Now, 22, 1–3,  
<https://www.roc.noaa.gov/WSR88D/PublicDocs/NNOW/NNow22c.pdf>.

Chrisman, J. N., 2014: Multiple elevation scan option for SAILS (MESO-SAILS)—The next step in dynamic scanning  
515 for the WSR-88D. Radar Operations Center, 27 pp.,  
[https://www.roc.noaa.gov/wsr88d/PublicDocs/NewTechnology/MESO-SAILS\\_Description\\_Briefing\\_Jan\\_2014.pdf](https://www.roc.noaa.gov/wsr88d/PublicDocs/NewTechnology/MESO-SAILS_Description_Briefing_Jan_2014.pdf).

Chrisman, J. N., 2016: Mid-volume rescan of low-level elevations (MRLE): A new approach to enhance sampling of  
quasi-linear convective systems (QLCSs). New Radar Technologies Web Page, NOAA/NWS/Radar Operations  
520 Center, 21 pp.,  
[https://www.roc.noaa.gov/WSR88D/PublicDocs/NewTechnology/DQ\\_QLCS\\_MRLE\\_June\\_2016.pdf](https://www.roc.noaa.gov/WSR88D/PublicDocs/NewTechnology/DQ_QLCS_MRLE_June_2016.pdf).

Colin, M., and S. C. Sherwood, 2021: Atmospheric convection as an unstable predator-prey process with memory. *J. Atmos. Sci.*, **78**, 3781–3797, doi:10.1175/JAS-D-20-0337.1.

525

Crum, T. D. and R. L. Alberty, 1993: The WSR-88D and the WSR-88D operational support facility. *Bull. Amer. Meteor. Soc.*, **74**, 1669–1688, doi:10.1175/1520-0477(1993)074<1669:TWATWO>2.0.CO;2.

Fan, J., D. Rosenfeld, Y. Zhang, S. E. Giangrande, Z. Li, L. A. T. Machado, S. T. Martin, Y. Yang, J. Wang, P. Artaxo, H. M. J. Barbosa, R. C. Braga, J. M. Comstock, Z. Feng, W. Gao, H. B. Gomes, F. Mei, C. Pöhlker, U. Pöschl, and R. A. F. De Souza, 2018: Substantial convection and precipitation enhancements by ultrafine aerosol particles. *Sci.*, **359**, 411–418, doi:10.1126/science.aan8461.

Fridlind, A. M., X. Li, D. Wu, M. van Lier-Walqui, A. S. Ackerman, W. Tao, G. M. McFarquhar, W. Wu, X. Dong, J. Wang, A. Ryzhkov, P. Zhang, M. R. Poellot, A. Neumann, and J. M. Tomlinson, 2017: Derivation of aerosol profiles for MC3E convection studies and use in simulations of the 20 May squall line case. *Atmos. Chem. Phys.*, **17**, 5947–5972, doi:10.5194/acp-17-5947-2017.

Fridlind, A. M., M. van Lier-Walqui, S. Collis, S. E. Giangrande, R. C. Jackson, X. Li, T. Matsui, R. Orville, M. H. Picel, D. Rosenfeld, A. Ryzhkov, R. Weitz, and P. Zhang, 2019: Use of polarimetric radar measurements to constrain simulated convective cell evolution: a pilot study with Lagrangian tracking. *Atmos. Meas. Tech.*, **12**, 2979–3000, doi:10.5194/amt-12-2979-2019.

Gagin, A., D. Rosenfeld and R. E. Lopez, 1985: The relationship between height and precipitation characteristics of summertime convective cells in south Florida. *J. Atmos. Sci.*, **42**, 84–94.

545

Grabowski, W. W., and H. Morrison, 2021: Supersaturation, buoyancy, and deep convection dynamics. *Atmos. Chem. Phys.*, **21**, 13997–14018, doi:10.5194/acp-21-13997-2021.

550 Greene, D. R., and R. A. Clark, 1972: Vertically Integrated Liquid Water – A new analysis tool. *Mon. Wea. Rev.*, **100**, 548–552, doi:10.1175/1520-0493(1972)100<0548:VILWNA>2.3.CO;2.

Haurwitz, B., 1947: Comments on the sea-breeze circulation. *J. Atmos. Sci.*, **4**, 1–8, doi:10.1175/1520-0469(1947)004<0001:COTSBC>2.0.CO;2.

555

Hitschfeld, W., 1960: The motion and erosion of convective storms in severe vertical wind shear. *J. Atmos. Sci.*, **17**, 270–282, doi:10.1175/1520-0469(1960)017<0270:TMAEOC>2.0.CO;2.

Hu, J., D. Rosenfeld, D. Zrníc, E. Williams, P. Zhang, J. C. Snyder, A. Ryzhkov, E. Hashimshoni, R. Zhang, and R.  
560 Weitz, 2019a: Tracking and characterization of convective cells through their maturation into stratiform storm elements using polarimetric radar and lightning detection. *J. Atmos. Res.*, **226**, 192–207, doi:10.1016/j.atmosres.2019.04.015.

Hu, J., D. Rosenfeld, A. Ryzhkov, D. Zrníc, E. Williams, P. Zhang, J. C. Snyder, R. Zhang, and R. Weitz, 2019b:  
565 Polarimetric radar convective cell tracking reveals large sensitivity to cloud precipitation and electrification properties of CCN. *J. Geophys. Res. Atmos.*, **124**, 12194–12205, doi:10.1029/2019jd030857.

Igel, A. L., M. R. Igel, and S. C. van den Heever, 2015: Make it a double? Sobering results from simulations using single-moment microphysics schemes. *J. Atmos. Sci.*, **72**, 910–925, doi:10.1175/JAS-D-14-0107.1.

570

Igel, A. L., and S. C. van den Heever, 2021: Invigoration or enervation of convective clouds by aerosols? *Geophys. Res. Lett.*, **48**, e2021GL093804, doi:10.1029/2021GL093804.

575 Jensen, M. P., J. H. Flynn, L. M. Judd, P. Kollias, C. Kuang, G. Mcfarquhar, R. Nadkarni, H. Powers, and J. Sullivan,  
2022: A succession of cloud, precipitation, aerosol, and air quality field experiments in the coastal urban environment.  
*Bull. Amer. Meteor. Soc.*, **103**, 103–105, doi:10.1175/BAMS-D-21-0104.1.

Kumjian, M. R., C. P. Martinkus, O. P. Prat, S. Collis, M. van Lier-Walqui, and H. C. Morrison, 2019: A moment-  
based polarimetric radar forward operator for rain microphysics. *J. Appl. Meteor. Climatol.*, **58**, 113–130,  
580 doi:10.1175/JAMC-D-18-0121.1.

Ladino, L. A., A. Korolev, I. Heckman, M. Wolde, A. M. Fridlind, and A. S. Ackerman, 2017: On the role of ice-  
nucleating aerosol in the formation of ice particles in tropical mesoscale convective systems. *Geophys. Res. Lett.*, **44**,  
1574–1582, doi:10.1002/2016GL072455.

585

Lamer, K., P. Kollias, E. P. Luke, B. P. Treserras, M. Oue, B. Dolan, 2023: Multisensor agile adaptive sampling  
(MAAS): a methodology to collect radar observations of convective cell life cycle. *J. Atmos. Oceanic Technol.*,  
submitted.

590 Lee, M., S. D. Schubert, M. J. Suarez, J. E. Schemm, H. Pan, J. Han, and S. Yoo, 2008: Role of convection triggers  
in the simulations of the diurnal cycle of precipitation over the United States Great Plains in a general circulation  
model. *J. Geophys. Res. Atmos.*, **113**, D02111, doi:10.1029/2007JD008984.

Mann, H. B., and D. R. Whitney, 1947: On a test of whether one of two random variables is stochastically larger than  
595 the other. *Ann. Math. Stat.*, **18**, 50–60, doi:10.1214/aoms/1177730491.

Oue, M., S. M. Saleeby, P. J. Marinescu, P. Kollias, and S. C. van den Heever, 2022: Optimizing radar scan strategies  
for tracking isolated deep convection using observing system simulation experiments. *Atmos. Meas. Tech.*, **15**, 4931–  
4950, doi:10.5194/amt-15-4931-2022.

600

Park, J. M., S. C. van den Heever, A. L. Igel, L. D. Grant, J. S. Johnson, S. M. Saleeby, S. D. Miller, and J. S. Reid,  
2020: Environmental controls on tropical sea breeze convection and resulting aerosol redistribution. *J. Geophys. Res.*  
*Atmos.*, **125**, e2019JD031699, doi:10.1029/2019JD031699.

605 Peters, K., C. Hohenegger, and D. Klocke, 2019: Different representation of mesoscale convective systems in  
convection-permitting and convection-parameterizing NWP models and its implications for large-scale forecast  
evolution. *Atmos.*, **10**, 503, doi:10.3390/atmos10090503.

Pulkkinen, S., D. Nerini, A. Perez Hortal, C. Velasco-Forero, U. Germann, A. Seed, and L. Foresti, 2019a: Pysteps:  
610 an open-source Python library for probabilistic precipitation nowcasting (v1.0). *Geosci. Model Dev.*, **12**, 4185–4219.  
doi:10.5194/gmd-12-4185-2019.

Pulkkinen, S., D. Nerini, A. Perez Hortal, C. Velasco-Forero, U. Germann, A. Seed, and L. Foresti, 2019b: Pysteps -  
a community-driven open-source library for precipitation nowcasting. *Poster presented at the 3rd European*  
615 *Nowcasting Conference*, Madrid, ES. doi: 10.13140/RG.2.2.31368.67840.



Radar Operations Center, 2015: WSR-88D Volume Coverage Pattern (VCP) improvement initiatives. New Radar Technologies Web Page, NOAA/NWS/Radar Operations Center, 8pp., [https://www.roc.noaa.gov/WSR88D/PublicDocs/NewTechnology/New\\_VCP\\_Paradigm\\_Public\\_Oct\\_2015.pdf](https://www.roc.noaa.gov/WSR88D/PublicDocs/NewTechnology/New_VCP_Paradigm_Public_Oct_2015.pdf).

620

Radar Operations Center, 2022: NEXRAD technical information. NEXRAD Technical Information Web Page, NOAA/NWS/Radar Operations Center, <https://www.roc.noaa.gov/WSR88D/Engineering/NEXRADTechInfo.aspx>.

Rosenfeld, D., 1987: Objective method for analysis and tracking of convective cells as seen by radar. *J. Atmos. Oceanic Technol.*, **4**, 422–434.

625

Rosenfeld, D., Y. Zheng, E. Hashimshoni, M. L. Pöhlker, A. Jefferson, C. Pöhlker, X. Yu, Y. Zhu, G. Liu, Z. Yue, B. Fischman, Z. Li, D. Giguzin, T. Goren, P. Artaxo, H. M. J. Barbosa, U. Pöschl, and M. O. Andreae, 2016: Satellite retrieval of cloud condensation nuclei concentrations by using clouds as CCN chambers. *Proc. Natl. Acad. Sci. USA*, **113**, 5828–5834, doi:10.1073/pnas.1514044113.

630

Rotunno, R., 1983: On the linear theory of the land and sea breeze. *J. Atmos. Sci.*, **40**, 1999–2009, doi:10.1175/1520-0469(1983)040<1999:OTLTOT>2.0.CO;2.

Ryzhkov, A., M. Pinsky, A. Pokrovsky, and A. Khain, 2011: Polarimetric Radar Observation Operator for a Cloud Model with Spectral Microphysics. *J. Appl. Meteor. Climatol.*, **50**, 873–894, doi:10.1175/2010JAMC2363.1.

635

Sheffield, A. M., S. M. Saleeby, and S. C. van den Heever, 2015: Aerosol-induced mechanisms for cumulus congestus growth. *J. Geophys. Res. Atmos.*, **120**, 8941–8952, doi:10.1002/2015JD023743.

640

Varble, A., E. J. Zipser, A. M. Fridlind, P. Zhu, A. S. Ackerman, J. Chaboureau, S. Collis, J. Fan, A. Hill, and B. Shipway, 2014: Evaluation of cloud-resolving and limited area model intercomparison simulations using TWP-ICE observations: 1. Deep convective updraft properties. *J. Geophys. Res. Atmos.*, **119**, 13891–13918, doi:10.1002/2013JD021371.

Wolfensberger, D. and A. Berne, 2018.: From model to radar variables: a new forward polarimetric radar operator for COSMO, *Atmos. Meas. Tech.*, **11**, 3883–3916, doi:10.5194/amt-11-3883-2018.

650 Zhu, P., J. Dudhia, P. R. Field, K. Wapler, A. Fridlind, A. Varble, E. Zipser, J. Petch, M. Chen, and Z. Zhu, 2012: A limited area model (LAM) intercomparison study of a TWP-ICE active monsoon mesoscale convective event. *J. Geophys. Res. Atmos.*, **117**, D11208, doi:10.1029/2011JD016447.


Zittel, W. D., 2019: Theory and concept of operations for multi-PRF dealiasing algorithm's VCP 112. New Radar Technologies Web Page, NOAA/NWS/Radar Operations Center, 13 pp., [https://www.roc.noaa.gov/WSR88D/PublicDocs/NewTechnology/Theory\\_ConOps\\_VCP112.pdf](https://www.roc.noaa.gov/WSR88D/PublicDocs/NewTechnology/Theory_ConOps_VCP112.pdf).

## Tables


	Shallow Convection	Modest Deep Conv.	Vigorous Deep Conv.
Initial Cluster Fraction	1	1	1
Lifetime Min. GOESBT	$\geq 268 \text{ K}$	$\leq 250 \text{ K}$	$\leq 250 \text{ K}$
Lifetime Max. VIL	$\geq -10 \text{ dB}$	$\geq 0 \text{ dB}$	$\geq 0 \text{ dB}$
Lifetime Max. ETH	$< 8 \text{ km}$	$8 \leq \text{ETH} < 12 \text{ km}$	$\geq 12 \text{ km}$
Lifetime Max. ERatio	$\geq 0.60$	$\geq 0.75$	$\geq 0.75$
Lifetime Max. Area	$\leq 30 \text{ km}^2$	N/A	N/A
Splits/Merges Removed	Yes	Yes	Yes
# of cells	35974	7930	4869
# of radar scans	142921	92770	95219

660 Table 1: A table of the thresholds used to isolate shallow, modest deep, and vigorous deep convection from all tracked features from the modified version of the MCIT algorithm.

## Figures



665 Figure 1: A visual illustration of  $ERatio$ ,  $H_{cell}$ , and  $H_{EBase}$ . This image is meant to provide a visual context for these three variables. The red, yellow, and green shaded areas represent arbitrary high, medium, and low reflectivity contours respectively.  $ERatio$  and  $H_{EBase}$  are calculated using the maximum  $ETH$  and  $H_{cell}$ , which are radar-derived quantities and will always be less than the actual height of the storm top and physical depth of a given cell.



670 Figure 2: Bar graphs showing the monthly average cell count for shallow (green), modest deep (yellow), and vigorous deep (red) convection. The vertical black lines at the top of each bar denote  $\pm$  one standard deviation.






Figure 3: The normalized lifetime distributions of the bin count normalized cell motion speed (a, b, c) and cell motion direction (d, e, f) for all shallow (a, d), modest deep (b, e), and vigorous deep (c, f) convective cells.




675

Figure 4: The frequency of initiation based on local time of day for (green) shallow convection, (yellow) modest deep convection, and (red) vigorous deep convection.




680 Figure 5: Maps showing the spatial distributions of initiation locations normalized by the number of cells that initiated during that period for the 3-hour periods 09:00 to 11:59 (a, b, c), 12:00 to 14:59 (d, e, f), and 15:00 to 17:59 (g, h, i), local time for shallow (a, d, g), modest deep (b, e, h), and vigorous deep (c, f, i) convection. The black (red) star in each plot denotes the location of KHGX (downtown Houston).





685 Figure 6: The time bin normalized distributions of GOESBT (a, b, c),  $H_{dBZ_{max}}$  (d, e, f), and  $dBZ_{max}$  (g, h, i) over the course of a day for all scans over the lifetimes of shallow (a, d, g), modest deep (b, e, h), and vigorous deep (c, f, i) convection.



690 Figure 7: The distributions normalized by the number of scans considered for each cell lifetime segment of  $\text{dBZ}_{\text{max}}$  versus  $H_{\text{dBZmax}}$  for the first 25% (a, b, c), the second 25% (b, e, f), the third 25% (g, h, i), and the final 25% of cell lifetimes (j, k, l) for shallow (a, d, g, j), modest deep (b, e, h, i), and vigorous deep (c, f, i, l) convection.





Figure 8: As in Fig. 7, but for maximum VIL versus GOESBT.



695 Figure 9: As in Fig. 7, but for  $\text{dBZ}_{\text{avg}}$  versus  $\text{dBZ}_{\text{max}}$ . The dashed line shows the one-to-one value of  $\text{dBZ}_{\text{avg}}$  and  $\text{dBZ}_{\text{max}}$ .




Figure 10: The bin normalized distributions for normalized lifetime evolutions of ETH (a, b, c),  $H_{dBZ_{max}}$  (d, e, f),  $dBZ_{max}$  (g, h, i), and  $H_{EBase}$  (j, k, l) for shallow (a, d, g, j), modest deep (b, e, h, k), and vigorous deep (c, f, i, l) convection.

700




Figure 11: The bin normalized ETH (a, b, c), GOESBT (d, e, f), and  $H_{dBZ_{max}}$  (g, h, i) based ascent rates for shallow (a, d, g), modest deep (b, e, h), and vigorous deep (c, f, i) convection. The dashed lines denote the zero line, the solid lines denote the median values per bin for all normalized lifetime bins, and the dotted lines above and below the median lines denote the 90th and 10th percentile values per bin for all normalized lifetime bins.

705




Figure 12: Violin plots depicting the distributions of the average of the 30-minute AOD values for the period prior to cell initiation to the southwest (red) and northeast (blue) of Houston for (a) shallow, (b) modest deep, and (c) vigorous deep convective cells. The number n under each violin plot corresponds to the number of cells considered for each distribution.

710

## Appendix Tables and Figures

Shallow Convection				
	-5%		+5%	
	n	% change	n	% change
All Thresholds Simultaneously	41683	<i>15.87</i>	17857	<b>-50.36</b>
Lifetime Max. Area	34996	-2.72	36829	2.38
Lifetime Min. GOESBT	42016	<i>16.80</i>	18351	<b>-48.99</b>
Lifetime Max. ERatio	36025	0.14	35888	-0.24
Lifetime Max. ETH	35823	-0.42	36068	0.26
Lifetime Max. VIL	36903	2.58	35059	-2.54
Modest Deep Convection				
	-5%		+5%	
	n	% change	n	% change
All Thresholds Simultaneously	4991	<b>-37.06</b>	10168	<b>28.22</b>
Lifetime Min. GOESBT	5769	<b>-27.25</b>	9865	<b>24.40</b>
Lifetime Max. ERatio	7951	0.26	7901	-0.37
Lifetime Max. ETH (Lower Bound)	8563	7.98	7307	-7.86
Lifetime Max. ETH (Upper Bound)	6522	<i>-17.76</i>	9242	<i>16.54</i>
Lifetime Max. VIL	7987	0.72	7879	-0.64
Vigorous Deep Convection				
	-5%		+5%	
	n	% change	n	% change
All Thresholds Simultaneously	5933	<b>21.85</b>	3599	<b>-26.08</b>
Lifetime Min. GOESBT	4684	-3.80	4975	2.18
Lifetime Max. ERatio	4873	0.08	4863	-0.12
Lifetime Max. ETH	6278	<b>28.94</b>	3557	<b>-26.95</b>
Lifetime Max. VIL	4877	0.16	4862	-0.14

715 Table A1: A table showing the sensitivity of cell selection based on  $\pm 5\%$  adjustments of a given variable. The number n represents the number of cells selected after the threshold adjustments were applied. Values for percent changes between  $\pm 10\%$  and  $\pm 20\%$  and  $> \pm 20\%$  are denoted by *italicized* and **bolded** values, respectively.






Figure A1: The lifetime distributions of GOESBT versus maximum VIL to illustrate threshold sensitivity for the most highly sensitive variables shown in Table A1. The baseline (no change) distributions are shown in (a, d, g, j), 5% decrease is shown in (b, e, h, k) and 5% increase is shown in (c, f, i, l) for shallow convection GOESBT sensitivity (a,b,c), modest deep convection GOESBT sensitivity (d, e, f), modest deep convection upper bound ERatio sensitivity (g, h, i) and vigorous deep convection ETH sensitivity (j, k, l).

720