

pH-Dependence of Brown Carbon Optical Properties in Cloud Water

Christopher J. Hennigan¹, Michael McKee¹, Vikram Pratap¹, Bryanna Boegner¹, Jasper Reno¹, Lucia Garcia¹, Madison McLaren¹, Sara M. Lance²

5

¹Department of Chemical, Biochemical and Environmental Engineering, University of Maryland, Baltimore County, Baltimore, 21250, USA

²Atmospheric Sciences Research Center (ASRC), University at Albany, Albany, 12226, USA

10 *Correspondence to:* Christopher J. Hennigan (hennigan@umbc.edu)

Abstract. Light-absorbing organic species present in aerosols, collectively called brown carbon (BrC), contribute important but highly uncertain effects on climate. Clouds likely represent a significant medium for secondary BrC production and for bleaching reactions, though the relative importance of formation and loss processes in clouds is unknown at present. The acidity (or pH) of atmospheric particles and clouds affects the optical properties of BrC and bleaching rates. Given the wide variability of pH in the atmosphere (pH in particles and clouds ranges from -1 to 8), the optical properties of BrC and its bleaching behavior are expected to vary significantly, and the link between pH and BrC is yet another uncertainty in attempts to constrain its climate forcing effects. In this work, we characterize the pH-dependence of BrC optical properties – including light absorption at 365 nm (Abs_{365}), mass absorption coefficient (MAC_{365}), and the absorption Ångström exponent (AAE) – in bulk cloud water sampled from the summit of Whiteface Mountain, NY. In all samples ($n = 17$), Abs_{365} and MAC_{365} increased linearly with increasing pH, highlighting the importance of reporting pH in studies of BrC in aqueous media. There was strong variability in the sensitivity of Abs_{365} to pH, with normalized slopes that ranged from 5.1% to 17.2% per pH unit. The normalized slope decreased strongly with increasing cloud water $[K^+]$, suggesting that the non-biomass burning BrC has optical properties that are more sensitive to pH than BrC associated with biomass burning. AAE also showed a distinct pH-dependence, as it was relatively flat between pH 1.5 – 5, then decreased significantly above pH 5. The cloud water composition was used to inform thermodynamic predictions of aerosol pH upwind/downwind of Whiteface Mountain and the subsequent changes in BrC optical properties. Overall, these results show that,

30 in addition to secondary BrC production, photobleaching, and the altitudinal distribution, the climate forcing of BrC is quite strongly affected by its pH-dependent absorption.

1 Introduction

Light-absorbing organic compounds, or chromophores, in particulate matter are collectively referred to as brown carbon (BrC). BrC exhibits a strong spectral dependence whereby the absorption efficiency
35 increases as wavelength decreases (Lack and Cappa, 2010; Kirchstetter et al., 2004). The light absorbing properties of BrC – both the absorption efficiency at a given wavelength and the wavelength-dependence – vary considerably in the atmosphere (Saleh, 2020). This variability in optical properties is due to the multitude of different organic compounds that contribute to BrC (Laskin et al., 2015). Regionally, the radiative forcing of BrC can represent an important component in the direct effect of aerosols on the
40 radiative balance (Zhang et al., 2017). However, the global climate forcing attributed to BrC is largely unconstrained at present due to the chemical complexity and widely variable optical properties; global climate models predict a net warming effect from BrC but estimates of the direct radiative forcing vary by almost a factor of 20 (from $+0.03 \text{ W m}^{-2}$ up to $+0.57 \text{ W m}^{-2}$) (Saleh, 2020).

45 BrC is different from other absorbing aerosols, dust and black carbon (BC), because it has prominent primary and secondary sources (Laskin et al., 2015). Like BC and dust, BrC is removed from the atmosphere through wet and dry deposition; however, BrC also undergoes chemical losses initiated by oxidants and direct photolysis (collectively termed bleaching) that can rapidly diminish its light
50 absorbing properties (Hems and Abbatt, 2018). Multiphase atmospheric processes, including those in clouds, play a key role in the life cycle of BrC, as they can facilitate production or loss, depending on the conditions (Hems and Abbatt, 2018; Laskin et al., 2015; Schnitzler and Abbatt, 2018; Zhao et al., 2015; Lee et al., 2014; Yu et al., 2016; Lin et al., 2015). Many laboratory studies have investigated BrC formation and loss using cloud water mimics (e.g., Powelson et al., 2014; De Haan et al., 2018) but few using real atmospheric cloud water samples. This represents a key knowledge gap in understanding the
55 temporal and spatial distribution of BrC and its effects on climate.

The acidity (or pH) of aerosols and clouds has tremendous importance for numerous atmospheric processes and for associated environmental effects (Pye et al., 2020). This includes an effect of pH on the radiative forcing of BrC compounds. pH affects the light-absorbing properties of many organic
60 compounds in aqueous media (Baes and Bloom, 1990). Aerosol samples collected in the southeastern U.S. showed pronounced pH-dependent absorbance spectra (Phillips et al., 2017). This was observed for both background ambient samples and those that had been influenced by biomass burning emissions. The increase in absorbance with increasing pH suggests that carboxylic acids and phenols contributed significantly to BrC in this study (Phillips et al., 2017). A similar study found that the aqueous extracts
65 of biomass burning aerosols exhibited a strong spectral dependence with pH (Cai et al., 2018). Further, pH had an important effect on the evolution of BrC bleaching in photolysis experiments conducted with the aqueous aerosol extracts (Cai et al., 2018). This is expected because many aqueous reaction rates are pH-dependent (Pye et al., 2020; Tilgner et al., 2021).

70 The pH of atmospheric particles, cloud, and fog droplets spans a wide dynamic range, from approximately -1 in highly acidic particles up to 8 in clouds containing alkaline components (Shah et al., 2020; Pye et al., 2020). Therefore, the optical properties of BrC and its bleaching behavior are expected to vary significantly in the atmosphere, as well. The pH-dependence of BrC absorbance and bleaching may be especially dynamic in cloud cycles because the pH of CCN particles can increase by 3-4 pH units when
75 they activate and form cloud droplets or decrease by the same amount when they transition from cloud droplets back to aqueous particles (Rusumdar et al., 2020). This suggests that the light absorbing properties of chromophores may be dramatically different in clouds compared to the properties of the same compounds in aqueous haze particles. In addition to in-cloud BrC transformations due to secondary formation and bleaching, changes in BrC absorbance with pH need to be characterized in order to
80 understand the impact of clouds on BrC.

The purpose of this study is to characterize the pH-dependence of BrC optical properties in cloud water samples collected at Whiteface Mountain, NY (WFM). The samples were collected over two summers and have diverse chemical composition and back trajectories, suggesting a variety of source influences

85 and airmass ageing. The analysis of BrC optical properties is paired with aerosol thermodynamic equilibrium modeling and cloud water composition measurements to calculate the aerosol liquid water content and pH of aerosols upwind of WFM. Using the measured pH-dependence of BrC optical properties, the modeled aerosol pH upwind of WFM informs changes to BrC radiative forcing in the atmosphere.

90 **2 Materials and Methods**

Cloud water samples were collected at the summit of Whiteface Mountain, NY (latitude N44°21'58" and longitude W 73°54'10", 1483 m a.s.l.) during the summers of 2018 and 2019. The site has been used continuously for atmospheric chemistry and cloud research for more than three decades (Mohnen and Kadlec, 1989). Procedures for cloud water collection and chemical analysis have been described in
95 detail before (Lance et al., 2020; Schwab et al., 2016; Lawrence et al., 2023). Briefly, cloud water collection occurred via a passive Mohnen omni-directional sampler when non-precipitating liquid clouds were present under select meteorological conditions. The collected cloud water was refrigerated (4°C), and analyzed for inorganic ionic species, organic acids, pH, conductivity, and total organic carbon (TOC) content. A subset of the samples was syringe filtered (0.45 µm) prior to analysis and thus, the analyzed
100 organics represents WSOC instead of TOC (Lance et al., 2020). The remaining sample volume was frozen (-20°C) for further analysis. Some of these frozen samples were packed in dry ice and shipped overnight to the University of Maryland, Baltimore County for BrC analysis.

In this study, BrC is operationally defined as the water-soluble organic carbon compounds that absorb
105 light in the 300 – 500 nm wavelength range. Many chromophores are found in the atmosphere that are insoluble or are soluble in other solvents (e.g., methanol) (Jiang et al., 2022; Zeng et al., 2020; Zhang et al., 2013). For the cloud water samples that were filtered, and since no additional organics were added, our analyses neglect any chromophores that are insoluble in water or soluble in other solvents (e.g., methanol), which can be 40-50% of BrC globally (Zeng et al., 2020). Absorbance spectra of cloud water
110 samples at each pH were recorded based on the method described previously (Pratap et al., 2020). A liquid waveguide capillary cell (LWCC, World Precision Instruments) with 50 cm path length was

coupled via fiber optic cables to a light source (200 – 1600 nm, Ocean Insight DH-mini) and two spectrometers, one to record absorbance spectra (FLAME-S, Ocean Optics) and one to continuously monitor the light source stability (STS-VIS, Ocean Insight). An automated syringe pump (model C3000, TriContinent Scientific) delivered the pH-adjusted cloud water sample to the LWCC, which was thoroughly rinsed with DI water ($>18.2 \text{ M}\Omega \cdot \text{cm}$) in between each sample. Absorbance spectra from 300 – 500 nm were recorded every 3 seconds and averaged over 4 minutes. Mass absorption coefficients at 365 nm ($\text{MAC}_{365}, \text{m}^2 \text{g}^{-1}$) were calculated according to Saleh (2020). The absorbance spectra from 300 – 500 nm were fit with an exponential function (Igor Pro, WaveMetrics) to calculate the absorption 120 Ångström exponent at each pH (AAE_{pH}).

The optical properties of each cloud water sample were measured at pH from 11 to 1.5. Following the approach of Phillips et al. (2017), sample pH was cycled from 2 to 11 to avoid potential hysteresis effects prior to the optical measurements. The cloud water sample was adjusted to pH 11 using 1.0 M NaOH (Fisher Scientific). Downward pH adjustments of ~ 1 pH unit were made through additions of 0.1 M and 3.0 M HCl (Fisher Scientific). pH was measured using an Orion Star A211 meter with Ross Ultra pH Electrode (Orion 8103BNUWP). The actual pH after NaOH and HCl additions was variable and depended on the cloud water composition (Table 1). Therefore, the actual pH at each step was recorded and used for the analyses. After the pH stabilized, approximately 1 mL of the cloud water sample was injected into the LWCC to fill the LWCC sample loop and tubing void volumes. The sample was injected into the LWCC within minutes of the pH adjustment, minimizing the time for any acid-catalyzed reactions of BrC chromophores to occur. The optical measurements accounted for the small diluting effects of the NaOH and HCl additions. Initially, these pH titrations and optical analyses were performed in duplicate on each cloud water sample; however, a high degree of repeatability was observed (Figure 1), so subsequent samples were only analyzed once to conserve the limited sample volumes. QA/QC on the experimental procedure was run using the full range of pH adjustments to DI water. The DI water at all pH levels showed absorbance values approximately 2 orders of magnitude lower than the cloud water samples and often below the system limit of detection: No background adjustments were made to the cloud sample measurements. Cloud water samples were also compared to Suwannee River Natural

140 Organic Matter (SR-NOM, International Humic Substances Society), a well-characterized material that has similarities to BrC sampled in the atmosphere (Green et al., 2015; Graber and Rudich, 2006).

Back trajectories for each cloud water sample were calculated using the Hybrid Single-Particle Lagrangian Integrated Trajectory model (HYSPLIT, (Stein et al., 2015)). Following the approach of
145 Lawrence et al. (2021), 6-day back trajectories were initiated at each hour of cloud water sampling using meteorology from the North American Regional Reanalysis (NARR) 32 km x 32 km. Output from the HYSPLIT model included the air mass location and altitude, as well as meteorological parameters. The temperature (T) and relative humidity (RH) output from HYSPLIT were used as inputs to the ISORROPIA-II aerosol thermodynamic equilibrium model (Fountoukis and Nenes, 2007). The inorganic ionic
150 composition of the cloud water samples was used for the ISORROPIA-II model input, with the exception that Ca^{2+} and Mg^{2+} concentrations were excluded because a decadal analysis of WFM cloud composition revealed that these species likely derive predominantly from coarse particles (Lawrence et al., 2023). To provide an estimate of fine aerosol pH along the back trajectory, ISORROPIA-II was run in forward mode with solids formation suppressed (i.e., metastable) according to Pye et al. (2020). We note that this
155 approach assumes a constant chemical composition along the entire 6-day back trajectory, which is clearly not realistic. However, this exercise shows pH variations along the back trajectory that may arise solely from the changes in T and RH as air mass altitude and position changes. In many locations, T and RH exert a greater influence on fine aerosol pH than composition, like sulfate and ammonium (Battaglia et al., 2017; Zheng et al., 2020; Tao and Murphy, 2021). Therefore, this exercise, while not accurate in
160 predicting the actual fine aerosol pH along the back trajectory, does reflect the magnitude of pH variability that may be expected as air masses were transported to WFM and how they compared to the cloud water pH.

3 Results

An overview of the cloud water samples analysed for this study is presented in **Table 1**. The cloud liquid
165 water content ranged from 0.26 g m^{-3} to 0.87 g m^{-3} (median 0.53 g m^{-3}) and the temperature ranged from $7.2 \text{ }^\circ\text{C}$ to $21.0 \text{ }^\circ\text{C}$ (median $17.2 \text{ }^\circ\text{C}$). Cloud water composition also showed quite a bit of variability: pH

ranged from 4.27 to 6.41 (median 4.70). TOC concentrations in the cloud water varied by more than an order of magnitude, from 106 μM to 1770 μM (median 505 μM). Amongst the inorganic ionic species measured, NH_4^+ had the highest median concentration (58.3 μM), followed by NO_3^- (32.5 μM), SO_4^{2-} (22.0 μM), and Cl^- (19.0 μM). In a prior study at WFM, Cook et al. (2017) found that K^+ concentrations exceeded 0.04 mg L^{-1} (1 μM) in cloud water samples influenced by fire emissions. In the present study, 12 of the 17 samples had K^+ concentrations above 0.04 mg L^{-1} , suggesting fire emissions had an important impact on cloud water composition. This is supported by the linear correlation between K^+ and TOC concentrations ($R^2 = 0.62$) in our samples and prior studies at WFM observing frequent influence of fire emissions on cloud water samples during the summer (Lance et al., 2020; Lee et al., 2022).

The distribution of MAC_{365} values at pH 4 is shown in **Figure 2**. MAC_{365} values at pH 4 ranged from 0.12 $\text{m}^2 \text{g}^{-1}$ up to 0.68 $\text{m}^2 \text{g}^{-1}$ (median of 0.34 $\text{m}^2 \text{g}^{-1}$). The median MAC_{365} value for WFM cloud water samples was similar to the value observed for Suwannee River NOM (0.30, green hatched bar in Fig. 2). A key feature of the BrC optical properties in our samples was the strong pH-dependence of absorption, as illustrated in Fig. 1 for duplicate analyses of sample 1922801. All samples showed a positive linear relationship, as absorption (and MAC) increased with increasing pH. This result is consistent with observations of BrC in ambient aerosols sampled in Georgia, including under conditions of biomass burning influence (Phillips et al., 2017). Based on the results of Schendorf et al. (2019), we expect the observed pH-dependence to be reversible, although this was not verified experimentally.

Although a positive linear relationship between MAC_{365} and pH was observed in all samples, there was considerable variation in the slope of this relationship for different cloud water samples (**Figure 3** and **Table 2**). Each sample's absorbance at 365 nm (Abs_{365}) at each pH was normalised to the sample Abs_{365} at pH 1.5. The slope of the normalized Abs_{365} vs. pH is a measure of how much BrC absorption changes for each pH unit change. The lowest relative slope was observed for sample 1822702 (slope of 0.051), which indicates the Abs_{365} varied by 5.1% per pH unit. Thus, a change of 6 pH units would only change the BrC absorption at 365 nm for this sample by $\sim 30\%$. The highest relative slope was observed for sample 1920702 (17.2% change in Abs_{365} per pH unit), where a 6 pH unit change would result in greater

195 than a factor of two change for the BrC absorption at 365 nm. The results in **Figure 3** are contrasted with results from Phillips et al. (2017), where slopes of 8% and 13% per pH unit were observed for BrC sampled in ambient aerosols uninfluenced and significantly influenced by biomass burning, respectively (solid red and green lines in **Figure 3**). The sensitivity of Abs_{365} to pH observed in WFM samples was similar to that of SR-NOM (thick dashed black line in **Figure 3**), which had a slope of 10.7% per pH unit.
200 The greater variability observed in our study may be due to the ageing of air masses sampled at WFM and more diverse source influences, which is discussed in detail below.

Variability in the response of BrC absorption to changes in pH was likely influenced by BrC sources. **Figure 4** shows the relationship between the normalized slope from Fig. 3 and the cloud water K^+
205 concentration. There was a clear decreasing trend in the sensitivity of BrC absorption to pH as $[K^+]$ increased. As discussed above, $[K^+]$ in WFM cloud water samples was used to identify biomass burning influence (Cook et al., 2017). The results in **Figure 4** suggest that aged biomass burning samples have less sensitivity to pH than ambient samples uninfluenced (or only lightly influenced) by biomass burning. This contrasts the results of a prior study that observed samples heavily influenced by fresh biomass
210 burning emissions showed higher sensitivity of BrC absorption to pH (Phillips et al., 2017). The apparent discrepancy between the results in Fig. 4 and those in Phillips et al. (2017) is likely due to atmospheric ageing. Phillip et al. (2017) sampled under conditions when large fires were burning relatively close to the sampling site, suggesting that the emissions were fresh. By contrast, fires burning in the western U.S. and Canada undergo transport of days (typically 3 – 7 days) before sampling at WFM (Cook et al., 2017;
215 Lance et al., 2020). Changes in BrC optical properties as biomass burning emissions age in the atmosphere are well documented (Saleh et al., 2013; Laskin et al., 2015; Forrister et al., 2015). Photobleaching and secondary BrC production alter the MAC and AAE of primary biomass burning emissions, though the timescale for these changes varies widely in the atmosphere and remains a major uncertainty in better representing BrC in global models (Saleh, 2020). The present results suggest that
220 atmospheric ageing may reduce the sensitivity of biomass burning BrC optical properties to pH, though future studies are needed to substantiate this finding through investigations that provide for controlled ageing conditions.

Similar to the MAC_{365} , the absorption Ångström exponent showed wide variability across the cloud water
225 samples (**Figure 5** and **Table 2**). The AAE_5 ranged from 4.63 (sample 1822702) to 7.09 (sample
1818204). AAE_5 values of cloud water samples were systematically higher than the SR-NOM, which
had an AAE_5 of 3.39 (green hatched bar in **Figure 5**). AAE_5 values indicate three samples (1822702,
1920004, and 1817601) were ‘moderately absorptive’ ($2.5 < AAE < 5$), while the remainder were ‘weakly
absorptive’ ($5 < AAE < 8$) according to the optical bins defined by Saleh (2020). The AAE values also
230 showed a distinct pH dependence. **Figure 6** shows the mean AAE values for WFM cloud water samples
as a function of pH. For pH 1.5 – 5, there was not much variation in AAE with pH; however, AAE
decreased significantly above pH 7. The results in Figure 6 are consistent with the results of Qin et al.
(2022b), who characterized water-soluble BrC in Beijing and observed a non-linear relationship between
AAE and pH, with significant reductions in AAE at $pH > 6$. This is likely due to chromophores with pKa
235 values above 6. For weakly acidic and moderately basic BrC chromophores, changes in pH below pH 6
impart only minor changes to the acid-base equilibrium and thus the spectral dependence on pH. Above
pH 6, BrC speciation becomes more sensitive to pH variations, resulting in changes to AAE in this range.

The optical properties of BrC are highly variable, reflecting the chemical diversity of atmospheric organic
240 compounds that absorb light (Laskin et al., 2015). In general, as BrC becomes more highly oxidized (and
thus water-soluble), BrC absorption efficiency weakens and AAE increases (Saleh, 2020). The optical
properties of BrC in WFM samples were broadly consistent with the findings in other locations. For
example, the AAE values of WSOC sampled in cloud water at Mt. Tianjing, China ranged from 5.37 –
6.31 during a study where cloud water composition was strongly influenced by biomass burning (blue
245 hatched bar in **Figure 5**, (Guo et al., 2022)). Water-soluble BrC extracted from $PM_{2.5}$ samples exhibited
average AAE values of 6.6 – 6.8 in Beijing and 4.8 – 7.3 in other cities in China (Qin et al., 2022a; Wu
et al., 2020), while water-soluble $PM_{2.5}$ extracts at various locations in India had AAE values of 5.1 – 5.3
(Kirillova et al., 2014; Srinivas et al., 2016). In the U.S., water-soluble organic carbon had mean AAE
values of 7.6 in Los Angeles (Zhang et al., 2013) and 6.0 – 8.3 in Atlanta, depending on the time of year
250 (Hecobian et al., 2010). In the Atmospheric Tomography Missions (ATom-2, ATom-3, and ATom-4)

aircraft studies probing global aerosol compositions, water-soluble BrC exhibited mean AAE values of 4.1 – 6.5 (red patterned bars in **Figure 5**; (Zeng et al., 2020)). There was not an apparent trend in the geographic distribution of AAE, though there was an altitudinal dependence, with lowest AAE values observed at the highest (10 – 13 km) and lowest altitudes (< 1 km) (Zeng et al., 2020).

255

A comparison of the present results to other studies brings up a critical point: Most of the prior studies did not report the pH of the aqueous extracts or the pH-dependence of absorption or AAE. The results in Figures 1, 3, and 6 show that the optical properties of atmospheric water-soluble BrC depend strongly on pH. Therefore, measurements of BrC in aqueous environments need to include and report pH in order to facilitate interstudy comparisons and to assess the climate forcing effects of BrC. **Further, optical properties of water-soluble BrC in aqueous environments should be measured at two pH levels, when feasible, to enable translation to other conditions in the atmosphere.**

In addition to the changes in Abs_{365} (and MAC_{365}) and AAE with pH, the relative absorption changed with wavelength at each pH, as well. **Figure 7** shows the absorption at a given pH relative to the absorption at pH 1.5 averaged over all cloud water samples. There was a clear wavelength-dependence of the absorption enhancements with increasing pH. At $pH < 6$, there was a minor increase in absorption from 300 nm up to 340 nm. The mean absorbance spectra below pH 6 were relatively flat from 340 nm up to 400 nm, declined gradually to a minimum around 450 nm, then gradually increased up to 500 nm. Above pH 7, mean absorbance spectra showed a steady increase from 300 nm – 380 nm, peak absorption relative to pH 1.5 at 380 – 400 nm, followed by a similar decline and minimum at ~450 nm. Comparison to other studies reveals both similarities and differences in the wavelength-dependent variations with pH shown in **Figure 7**. For example, in a study characterizing the effects of pH on water-soluble BrC aerosols in Beijing, there was a similar wavelength-dependence to the absorption enhancement we observed in **Figure 7** (Qin et al., 2022a); however, the magnitude of the absorption enhancement at 400 nm relative to pH 1.5 was higher in WFM cloud water at each pH level.

4 Discussion

Our results have important implications for understanding BrC in the atmosphere. It is imperative that studies reporting the optical properties of water-soluble BrC – whether in aerosol extracts, cloud, or fog water – also report the pH at which the measurements were conducted. The *in situ* aerosol pH (ranging from -1 to 8) may be dramatically different from the pH of dilute aqueous extract solutions (most likely in the range of pH 4 to 6). This will facilitate more accurate comparisons between studies. Cloud cycling, with the accompanying 2 – 3 pH unit step changes, results in BrC absorption efficiency changes of 10% – 50% for the WFM samples. Particle composition and changes in meteorology also change aerosol pH, such that BrC compounds may experience variability of 5 - 6 pH units over their atmospheric lifetime. Our results suggest that these pH changes dramatically change the climate forcing effects of BrC. Aerosols are usually more acidic than clouds, so BrC exhibits stronger absorption in clouds than in aerosols. The activated fraction of BrC has, to our knowledge, never before been explored but also affects the radiative forcing of BrC in aerosols and clouds. Therefore, in addition to factors that have been considered, such as the altitudinal distribution of BrC (Zhang et al., 2017), accurately constraining the climate forcing effects of BrC requires accounting for the sensitivity to pH.

The results in this study suggest that aromatic carboxylic acids and phenolic compounds, including nitrophenols, are primarily responsible for the observed pH-dependent optical properties. Phenols and aromatic carboxylic acids are major contributors to atmospheric BrC (Laskin et al., 2015). Prior studies have shown that aromatic carboxylic acids contribute most to the pH-dependence below pH 7 while phenols are most responsible above pH 7 (Schendorf et al., 2019; Qin et al., 2022c). Our results also suggest that phenols are primarily responsible for the observed dependence of AAE on pH (Figure 6). AAE only varied with pH above pH 7, consistent with phenolic pKa values that are typically in the 7 – 10 range. The wavelength-dependent enhancements with increasing pH shown in Figure 7 also point to the influence of phenols because the shape and magnitude of the enhancements become much more prominent above pH 7.

305 The variability in BrC absorption due to pH is illustrated in **Figure 8**, which shows back trajectories corresponding to each cloud water sample analysed in this study. The 6-day back trajectories are coloured by the Abs_{365} along the trajectory normalized to the Abs_{365} for cloud water conditions. The relative Abs_{365} changes in Figure 8 are based upon the changes in pH along the back trajectories as well as the sensitivity of absorption to pH for each sample (normalized slope in Table 2). For all samples, the maximum Abs_{365} occurs in-cloud because the pH is systematically higher than in particles due to the effect of dilution. The predicted changes range from minor to major, depending on the sample. The mean normalized Abs_{365} among all trajectories was 0.637 (median 0.661), indicating a $\sim 1/3$ reduction in BrC absorption in the aerosols relative to the cloud water. As described above, aerosol pH modelled along the back trajectory was computed using the cloud water composition so it only accounts for changes in T and RH. Although these factors have a major effect on pH (Tao and Murphy, 2019), the actual pH along the back trajectories will differ due to changes in aerosol composition. Abs_{365} will further change if secondary BrC production or bleaching occurs during transport to WFM. Therefore, the results in **Figure 8** serve only as a guide to illustrate the magnitude of BrC absorbance changes that may occur in the atmosphere due to changes in pH associated with changes in T and RH. **Figure 8** also illustrates the diversity of sources influencing the cloud water samples, consistent with the variable cloud water composition discussed above. The trajectories collectively show marine, clean arctic, biogenic, and polluted continental air mass origins. We attempted to correlate BrC optical properties with the trajectory analyses; however, this was inconclusive because of the small number of trajectories originating in some regions.

325 These results also inform measurements of BrC that are not conducted in aqueous matrices. For example, experimental approaches such as cavity ringdown spectroscopy (CRDS), photoacoustic spectrometry (PAS), and aethalometry are frequently used to measure total BrC, not just the water-soluble fraction (Liu et al., 2015). Non-filter based approaches, including CRDS and PAS, typically dry the air sample before measurement (Washenfelder et al., 2013; Lack et al., 2012). It is unclear how the optical properties of chromophoric WSOC change as the particles transition from ambient conditions, where they often contain liquid water, to the dry environment within the instrument. Our results suggest that the water-soluble BrC compounds that exhibit a pH dependence will also exhibit different absorbance behaviours

transitioning from an aqueous to non-aqueous phase state, though this topic should be explored in detail in the future.

335

BrC compounds are likely to encounter liquid water during much of their time in the atmosphere and water-soluble chromophores can dissolve or partially dissolve in the aqueous fraction (Zhang et al., 2012; Pye et al., 2018). Therefore, the optical properties of BrC are not static, but evolve in space and time. Secondary BrC formation and photobleaching contribute to these dynamic changes. Our results show that changing pH, which can be due to changing inorganic aerosol composition, changing T and RH, also contribute to changing BrC optical properties that may be as - or more - impactful as these other processes.

Our results indicate that pH is a key parameter that may dramatically change the radiative forcing of water-soluble BrC relative to insoluble BrC. Water-soluble BrC makes a variable contribution to the overall light absorption. Guo et al. (2022) found that water-soluble BrC contributed approximately half of the Abs_{370} in cloud droplet residuals sampled at Mt. Tianjing, China, and Zeng et al. (2020) estimated that the water-soluble fraction of BrC was approximately half of the total BrC in the ATom aircraft studies characterizing the global distribution of aerosol composition and concentration. However, our results show that changes in aerosol and cloud water pH will clearly change the climate forcing of water-soluble BrC relative to the total. Changes in composition, aerosol and cloud processes (e.g., cloud activation or drying), and meteorology (temperature and RH) strongly affect pH (Pye et al., 2020). Therefore, the contribution of water-soluble BrC to total BrC radiative forcing will change in time and space, as well.

The WFM cloud water samples showed similarities with Suwannee River NOM. Specifically, the MAC at pH 4 (Fig. 2) and the normalized slope of Abs_{365} vs. pH (Fig. 3) for SR-NOM were both very close to the median WFM cloud water samples. More broadly, Schendorf et al. (2019) characterized the pH effects on optical properties of five different humic substances isolated from aquatic and terrestrial systems and found similar behaviour to our cloud water samples, as absorption increased with increasing pH, though the wavelength-dependence and spectral characteristics differed between different samples. Our WFM cloud water samples were most similar to their analysis of Suwannee River fulvic and humic

360

acids (Schendorf et al., 2019), materials that contain hydrophobic organic acids isolated from the SR-NOM we analysed (<https://humic-substances.org/>, last accessed 24-Feb-2023). It is now well known that chemical and physical characteristics of atmospheric organic matter – especially the light-absorbing fraction – bear similarities to organic material found in aquatic and terrestrial systems (Kalberer et al., 2004; Graber and Rudich, 2006). The present results build upon this body of work and support the use of these materials, which are abundant and commercially available, as surrogates for further studies into the pH-dependent optical properties of atmospheric BrC.

Turnock et al. (2019) analysed changes in aerosol radiative forcing due to cloud water pH. Their study considered the effects of pH on the aqueous oxidation of SO₂ in clouds and how this changed global particle size distributions. Ultimately, their model predicted stronger aerosol radiative forcing at higher cloud water pH because of enhanced SO₂ oxidation to sulphate in clouds and the resulting reduction in gas-phase production of H₂SO₄ (Turnock et al., 2019). Our results show that changes in cloud water pH also affect the climate forcing of aerosols through effects on BrC absorption. Increasing pH increased the absorption efficiency of BrC for all cloud water samples analysed. Because BrC absorption is sensitive to pH, photolysis rates (i.e., photobleaching) are sensitive to pH, as well. For example, Zhao et al. (2015) observed a factor of ~2 increase in the photolysis rate of a nitrophenol (4-nitrocatechol) going from pH 3 to pH 5. Another study observed an increase in the photolysis rates of nitrophenols (guaiacol, catechol, and 5-nitroguaiacol) with decreasing pH (Yang et al., 2023). Therefore, cloud and aerosol pH affect the lifetime of BrC compounds, which further affects their climate forcing.

Finally, our measurements were conducted on liquid cloud samples, so these results are limited to aqueous environments. Aerosol liquid water, while often abundant, is also highly variable in the atmosphere (Nguyen et al., 2016). Aqueous aerosols contain orders of magnitude less water than cloud droplets, so organic solutes rapidly transition between highly concentrated and dilute environments during cloud cycling. Organic compounds in atmospheric particles span a very wide range of water solubilities, including compounds measured in the water-soluble fraction (i.e., as WSOC) (Psichoudaki and Pandis, 2013). Changes in liquid water content cause organic aerosols to undergo various phase transitions,

including liquid-liquid phase separation and transition to highly viscous “glassy” states (Reid et al., 2018).
390 Acidity can be defined in non-aqueous media, though the pH scale as a measure of solution acidity does
not directly transfer between different solvents (Himmel et al., 2018). Therefore, our results do not inform
the optical properties of BrC in non-aqueous environments; however, future studies should address this
question. These results reflect the optical properties of organic compounds dissolved in cloud water at
their time of sampling under the environmental conditions given in Table 1; a different distribution of
395 solubility would likely change the measured optical properties, as well.

Data Availability

All data presented in this work are published in Tables 1 and 2. Any data not published, e.g., raw spectra,
are available upon request.

400

Author Contributions

C. Hennigan: Conceptualization, Methodology, Resources, Funding acquisition, Supervision, Writing –
Original Draft, Writing – Review & Editing. M. McKee: Methodology, Validation, Formal analysis,
Writing – Review & Editing, Visualization. V. Pratap: Methodology, Validation, Formal analysis,
405 Writing – Review & Editing, Visualization. B. Boegner: Methodology, Validation, Formal analysis,
Writing – Review & Editing, Visualization. J. Reno: Methodology, Validation, Formal analysis, Writing
– Review & Editing, Visualization. L. Garcia: Methodology, Validation, Formal analysis, Writing –
Review & Editing, Visualization. M. McLaren: Methodology, Validation, Formal analysis, Writing –
Review & Editing, Visualization. S. Lance: Methodology, Resources, Writing – Original Draft, Writing
410 – Review & Editing.

Competing Interests

The authors declare no competing financial interests.

Acknowledgements

415

We acknowledge Paul Casson, Richard Brandt, and Eric Hebert for helping set up and service the cloud water collection system. We acknowledge Dan Kelting and Liz Yerger, Paul Smith's College, for conducting the chemical analysis of the cloud water samples.

420 **Financial Support**

This work was supported by the U.S. Department of Energy, Office of Biological & Environmental Research (BER) through grant DE-SC0022049. Cloud water and meteorological measurements reported in this paper were supported by the New York State Energy Research and Development Authority (NYSERDA) Contract 124461. NYSERDA has not reviewed the information contained herein, and the
425 opinions expressed in this report do not necessarily reflect those of NYSERDA or the State of New York.

Table 1: Overview of the cloud water composition for samples collected from the summit of Whiteface Mountain and analysed for this study.

Sample ID	Date	Cloud sampling duration (hr)	Cloud LWC (g/m ³)	Temp. (°C)	pH	TOC (µmol/L)
1817601	25-Jun-2018	4.7	0.51	7.2		439.3
1818204*	1-Jul-2018	10.9	0.66	19.0	4.51	988.3
1818205*	1-Jul-2018	3.8	0.33	19.0	4.57	1770
1820401	23-Jul-2018	4.95	0.40	17.7	4.60	206.1
1820702	26-Jul-2018	6.3	0.75	15.2	4.64	106
1821005	29-Jul-2018	11.1	0.87	11.5	4.88	216
1821301	1-Aug-2018	7.4	0.36	15.2	4.27	498.2
1822701	15-Aug-2018	2.1	0.55	17.2	4.62	869.2
1822702	15-Aug-2018	2.1	0.55	17.2		732.1
1822901	17-Aug-2018	5.8	0.72	15.8	4.76	311.4
1823805	26-Aug-2018	7.4	0.78	13.8	5.35	801.8
1920004*	19-Jul-2019	11.3	0.57	18.9	5.18	512.3
1921102*	30-Jul-2019	1.9	0.27	21.0	6.41	1323.3
1922202*	10-Aug-2019	10.8	0.50	15.0	5.38	344.6
1922801*	16-Aug-2019	6.8	0.51	18.3	4.28	540
1926801	25-Sept-2019	2.9	0.26	16.7	5.18	385

430

Sample ID	[Na ⁺] (µM)	[K ⁺] (µM)	[NH ₄ ⁺] (µM)	[Ca ²⁺] (µM)	[Mg ²⁺] (µM)	[Cl ⁻] (µM)	[NO ₃ ⁻] (µM)	[SO ₄ ²⁻] (µM)
1817601	0.9	1.0	20.0	1.0	0.4	25.7	11.3	14.9
1818204*	5.7	2.0	510.0	23.0	5.3	46.8	93.5	67.0
1818205*	10.0	3.1	908.9	28.2	8.2	57.3	125.0	102.4
1820401	31.3	1.3	14.4	3.5	2.5	28.5	21.3	10.5
1820702		0.8	20.0	2.0	0.8	13.8	24.4	9.9
1821005	1.3	0.5	10.6	3.2	1.2	22.6	6.1	6.7
1821301	13.9	1.5	52.2	7.5	2.1	22.3		21.8
1822701	7.0	3.1	141.1	21.5	4.9	18.9	40.5	22.3
1822702	0.4	2.6	80.0	19.2	4.9	19.2	35.6	21.4


1822901	1.3	2.3	60.0	13.0	3.7	0.8	37.7	25.2
1823805	13.0	2.6	141.7	39.2	13.6	3.7	69.7	32.8
1920004*	2.6	1.8	38.9	11.5	2.5	24.3	25.8	13.1
1921102*	3.0	4.6	248.3	94.6	11.9	1.7	158.5	85.7
1922202*	2.6	1.8	56.7	10.2	2.5	9.9	29.4	22.9
1922801*	2.6	2.0	96.7	10.7	4.1	9.9	90.6	30.3
1926801			9.4			5.4	10.6	7.9

*Denotes samples that were syringe-filtered (0.45 μm); thus the TOC reported is actually WSOC for these samples.


Table 2: Summary of key optical properties for cloud water samples.

Sample ID	MAC ₃₆₅ (m ² g ⁻¹)	*Normalized slope Abs ₃₆₅ vs. pH	AAE ₂	AAE ₅	AAE ₉
1817601	0.220	0.171	5.49	4.96	4.53
1818204	0.121	0.066	7.59	7.09	4.43
1818205	0.123	0.070	6.86	6.50	4.81
1820401	0.473	0.146	5.66	6.16	6.22
1820702	0.678	0.172	-	-	-
1821005	0.486	0.109	5.94	6.41	5.01
1821301	0.122	0.154	6.20	5.79	6.08
1822701	0.575	0.058	6.23	6.00	5.19
1822702	0.427	0.051	4.53	4.63	4.77
1822901	0.678	0.093	6.13	5.76	5.08
1823805	0.377	0.054	5.84	5.32	4.13
1920004	0.527	0.087	4.83	4.80	5.09
1921102	0.145	0.055	6.56	6.45	5.63
1922202	0.297	0.083	6.15	6.02	5.11
1922801	0.209	0.163	6.70	6.43	6.25
1926801	0.122	0.139	6.07	5.66	5.44

435 *Slope determined as Abs₃₆₅(pH)/Abs₃₆₅(pH 1.5) vs. pH for each sample (see Fig. 3)




440 **Figure 1:** Duplicate measurements of MAC₃₆₅ as a function of pH for cloud water sample #1922801. The slope, intercept, and coefficient of variation (R²) showed excellent repeatability for the analyses run for the same sample on different days. WSOC for this sample was 540 μM, close to the median value of 505 μM observed for this study.



445

Figure 2: Average mass absorption coefficient measured at 365 nm (MAC₃₆₅) and pH 4 for cloud water samples (solid blue bars) and Suwannee River natural organic matter (hatched green bar).



450

Figure 3: Normalized absorbance at 365 nm ($Abs_{365}(pH)/Abs_{365}(pH\ 1.5)$) versus pH for all cloud water samples analyzed in this study. Lines of best fit (least squares linear regression) for each cloud water sample are shown as the thin dotted lines. Comparison to BrC sampled in ambient aerosols both influenced and uninfluenced by biomass burning are shown with the solid green and red lines, respectively, both from Phillips et al. (2017). Mean behaviour of Suwannee River NOM shown with the thick dashed black line.

455




Figure 4: Relationship between the normalized slope of Abs₃₆₅ vs. pH (from Table 2) and the cloud
460 water K⁺ concentration.




Figure 5: Distribution of absorption Ångström exponent values at pH 5 (AAE₅) for cloud water samples (solid bars), Suwannee River Natural Organic Matter (green hatched bar), cloud water samples from Mt. Tianjing, China (blue hatched bar, mean value $\pm 1\sigma$ reported from Guo et al., 2022 for WSOC), and mean values from the ATom aircraft missions (red patterned bars, from Zeng et al., GRL, 2020). Note, the pH corresponding to the AAE values from Mt. Tianjing and ATom missions were not specified.




Figure 6: Mean absorption Ångström exponent value for all cloud water samples as a function of pH. Error bars represent $\pm 1\sigma$.

475





Figure 7: Mean absorption at a given pH relative to the absorption at pH 1.5 as a function of wavelength
 480 for all cloud water samples.



485 **Figure 8:** Map showing 6-day back trajectories for cloud water samples. Trajectories are coloured according to the relative BrC absorption at 365 nm (Abs_{365}) based upon the slopes (Table 2 and Fig. 3) and aerosol pH modelled along the back trajectory by ISORROPIA-II. The thermodynamic calculations assume constant aerosol composition along the back trajectory and only account for differences in pH based on T and RH.

References

- Baes, A. U. and Bloom, P. R.: FULVIC-ACID ULTRAVIOLET-VISIBLE SPECTRA - INFLUENCE OF SOLVENT AND PH, *Soil Science Society of America Journal*, 54, 1248-1254, 10.2136/sssaj1990.03615995005400050008x, 1990.
- 495 Battaglia, M. A., Douglas, S., and Hennigan, C. J.: Effect of the Urban Heat Island on Aerosol pH, *Environmental Science & Technology*, 51, 13095-13103, 10.1021/acs.est.7b02786, 2017.
- Cai, J., Zhi, G. R., Yu, Z. Q., Nie, P., Gligorovski, S., Zhang, Y. Z., Zhu, L. K., Guo, X. X., Li, P., He, T., He, Y. J., Sun, J. Z., and Zhang, Y.: Spectral changes induced by pH variation of aqueous extracts derived from biomass burning aerosols: Under dark and in presence of simulated sunlight irradiation, *Atmospheric Environment*, 185, 1-6, 10.1016/j.atmosenv.2018.04.037, 2018.
- 500 Cook, R. D., Lin, Y. H., Peng, Z. Y., Boone, E., Chu, R. K., Dukett, J. E., Gansch, M. J., Zhang, W. L., Tolic, N., Laskin, A., and Pratt, K. A.: Biogenic, urban, and wildfire influences on the molecular composition of dissolved organic compounds in cloud water, *Atmospheric Chemistry and Physics*, 17, 15167-15180, 10.5194/acp-17-15167-2017, 2017.
- 505 De Haan, D. O., Tapavicza, E., Riva, M., Cui, T. Q., Surratt, J. D., Smith, A. C., Jordan, M. C., Nilakantan, S., Almodovar, M., Stewart, T. N., de Loera, A., De Haan, A. C., Cazaunau, M., Gratien, A., Pangui, E., and Doussin, J. F.: Nitrogen-Containing, Light-Absorbing Oligomers Produced in Aerosol Particles Exposed to Methylglyoxal, Photolysis, and Cloud Cycling, *Environmental Science & Technology*, 52, 4061-4071, 10.1021/acs.est.7b06105, 2018.
- 510 Forrister, H., Liu, J., Scheuer, E., Dibb, J., Ziemba, L., Thornhill, K. L., Anderson, B., Diskin, G., Perring, A. E., Schwarz, J. P., Campuzano-Jost, P., Day, D. A., Palm, B. B., Jimenez, J. L., Nenes, A., and Weber, R. J.: Evolution of brown carbon in wildfire plumes, *Geophysical Research Letters*, 42, 4623-4630, 10.1002/2015gl063897, 2015.
- Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for K⁺-Ca²⁺-Mg²⁺-Nh(4)⁽⁺⁾-Na⁺-SO₄²⁻-NO₃⁻-Cl⁻-H₂O aerosols, *Atmospheric Chemistry and Physics*, 7, 4639-4659, 10.5194/acp-7-4639-2007, 2007.
- 515 Graber, E. R. and Rudich, Y.: Atmospheric HULIS: How humic-like are they? A comprehensive and critical review, *Atmospheric Chemistry and Physics*, 6, 729-753, DOI 10.5194/acp-6-729-2006, 2006.

- Green, N. W., McInnis, D., Hertkorn, N., Maurice, P. A., and Perdue, E. M.: Suwannee River Natural Organic
520 Matter: Isolation of the 2R101N Reference Sample by Reverse Osmosis, *Environ Eng Sci*, 32, 38-44,
10.1089/ees.2014.0284, 2015.
- Guo, Z., Yang, Y., Hu, X., Peng, X., Fu, Y., Sun, W., Zhang, G., Chen, D., Bi, X., Wang, X., and Peng, P.: The
optical properties and in-situ observational evidence for the formation of brown carbon in clouds, *Atmos.*
Chem. Phys., 22, 4827-4839, 10.5194/acp-22-4827-2022, 2022.
- 525 Hecobian, A., Zhang, X., Zheng, M., Frank, N., Edgerton, E. S., and Weber, R. J.: Water-Soluble Organic Aerosol
material and the light-absorption characteristics of aqueous extracts measured over the Southeastern United
States, *Atmospheric Chemistry and Physics*, 10, 5965-5977, 10.5194/acp-10-5965-2010, 2010.
- Hems, R. F. and Abbatt, J. P. D.: Aqueous Phase Photo-oxidation of Brown Carbon Nitrophenols: Reaction
Kinetics, Mechanism, and Evolution of Light Absorption, *Acs Earth and Space Chemistry*, 2, 225-234,
530 10.1021/acsearthspacechem.7b00123, 2018.
- Himmel, D., Radtke, V., Butschke, B., and Krossing, I.: Basic Remarks on Acidity, *Angewandte Chemie*
International Edition, 57, 4386-4411, <https://doi.org/10.1002/anie.201709057>, 2018.
- Jiang, X., Liu, D., Li, Q., Tian, P., Wu, Y., Li, S., Hu, K., Ding, S., Bi, K., Li, R., Huang, M., Ding, D., Chen, Q.,
Kong, S., Li, W., Pang, Y., and He, D.: Connecting the Light Absorption of Atmospheric Organic Aerosols
535 with Oxidation State and Polarity, *Environmental Science & Technology*, 56, 12873-12885,
10.1021/acs.est.2c02202, 2022.
- Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R., Weingartner, E.,
Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as major components of
atmospheric organic aerosols, *Science*, 303, 1659-1662, DOI 10.1126/science.1092185, 2004.
- 540 Kirchstetter, T. W., Novakov, T., and Hobbs, P. V.: Evidence that the spectral dependence of light absorption by
aerosols is affected by organic carbon, *Journal of Geophysical Research-Atmospheres*, 109,
10.1029/2004jd004999, 2004.
- Kirillova, E. N., Andersson, A., Tiwari, S., Srivastava, A. K., Bisht, D. S., and Gustafsson, O.: Water-soluble
organic carbon aerosols during a full New Delhi winter: Isotope-based source apportionment and optical
545 properties, *Journal of Geophysical Research-Atmospheres*, 119, 3476-3485, 10.1002/2013jd020041, 2014.
- Lack, D. A. and Cappa, C. D.: Impact of brown and clear carbon on light absorption enhancement, single scatter
albedo and absorption wavelength dependence of black carbon, *Atmospheric Chemistry and Physics*, 10,
4207-4220, 10.5194/acp-10-4207-2010, 2010.

- Lack, D. A., Richardson, M. S., Law, D., Langridge, J. M., Cappa, C. D., McLaughlin, R. J., and Murphy, D. M.,
550 Aircraft Instrument for Comprehensive Characterization of Aerosol Optical Properties, Part 2: Black and
Brown Carbon Absorption and Absorption Enhancement Measured with Photo Acoustic Spectroscopy,
Aerosol Science and Technology, 46 (5), DOI:10.1080/02786826.2011.645955, 2012.
- Lance, S., Zhang, J., Schwab, J. J., Casson, P., Brandt, R. E., Fitzjarrald, D. R., Schwab, M. J., Sicker, J., Lu, C.
H., Chen, S. P., Yun, J., Freedman, J. M., Shrestha, B., Min, Q. L., Beauharnois, M., Crandall, B., Joseph,
555 E., Brewer, M. J., Minder, J. R., Orłowski, D., Christiansen, A., Carlton, A. G., and Barth, M. C.: Overview
of the CPOC Pilot Study at Whiteface Mountain, NY Cloud Processing of Organics within Clouds (CPOC),
Bulletin of the American Meteorological Society, 101, E1820-E1841, 10.1175/bams-d-19-0022.1, 2020.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, Chemical Reviews, 115,
4335-4382, 10.1021/cr5006167, 2015.
- 560 Lawrence, C. E., Lance, S., Kelting, G., and Yererger, E.: Investigating Characteristic Air Masses Affecting Organic
and Inorganic Cloud Water Composition at Whiteface Mountain Using HYSPLIT and Cluster Analysis,
American Meteorological Society, Virtual,
- Lawrence, C. E., Casson, P., Brandt, R., Schwab, J. J., Dukett, J. E., Snyder, P., Yererger, E., Kelting, D.,
VandenBoer, T. C., and Lance, S.: Long-term monitoring of cloud water chemistry at Whiteface Mountain:
565 the emergence of a new chemical regime, Atmos. Chem. Phys., 23, 1619-1639, 10.5194/acp-23-1619-
2023, 2023.
- Lee, H. J., Aiona, P. K., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Effect of Solar Radiation on the Optical
Properties and Molecular Composition of Laboratory Proxies of Atmospheric Brown Carbon,
Environmental Science & Technology, 48, 10217-10226, 10.1021/es502515r, 2014.
- 570 Lee, J. Y., Peterson, P. K., Vear, L. R., Cook, R. D., Sullivan, A. P., Smith, E., Hawkins, L. N., Olson, N. E., Hems,
R., Snyder, P. K., and Pratt, K. A.: Wildfire Smoke Influence on Cloud Water Chemical Composition at
Whiteface Mountain, New York, Journal of Geophysical Research: Atmospheres, 127, e2022JD037177,
<https://doi.org/10.1029/2022JD037177>, 2022.
- Lin, P., Laskin, J., Nizkorodov, S. A., and Laskin, A.: Revealing Brown Carbon Chromophores Produced in
575 Reactions of Methylglyoxal with Ammonium Sulfate, Environmental Science & Technology, 49, 14257-
14266, 10.1021/acs.est.5b03608, 2015.
- Liu, S., Aiken, A. C., Gorkowski, K., Dubey, M. K., Cappa, C. D., Williams, L. R., Herndon, S. C., Massoli, P.,
Fortner, E. C., Chhabra, P. S., Brooks, W. A., Onasch, T. B., Jayne, J. T., Worsnop, D. R., China, S.,
Sharma, N., Mazzoleni, C., Xu, L., Ng, N. L., Liu, D., Allan, J. D., Lee, J. D., Fleming, Z. L., Mohr, C.,

- 580 Zotter, P., Szidat, S., and Prévôt, A. S. H.: Enhanced light absorption by mixed source black and brown carbon particles in UK winter, *Nature Communications*, 6, 8435, 10.1038/ncomms9435, 2015.
- Mohnen, V. A. and Kadlecek, J. A.: Cloud chemistry research at Whiteface Mountain, *Tellus B*, 41, 79-91, 10.1111/j.1600-0889.1989.tb00127.x, 1989.
- Nguyen, T. K. V., Ghate, V. P., and Carlton, A. G.: Reconciling satellite aerosol optical thickness and surface fine
585 particle mass through aerosol liquid water, *Geophysical Research Letters*, 43, 11,903-911,912, <https://doi.org/10.1002/2016GL070994>, 2016.
- Phillips, S. M., Bellcross, A. D., and Smith, G. D.: Light Absorption by Brown Carbon in the Southeastern United States is pH-dependent, *Environmental Science & Technology*, 51, 6782-6790, 10.1021/acs.est.7b01116, 2017.
- 590 Powelson, M. H., Espelien, B. M., Hawkins, L. N., Galloway, M. M., and De Haan, D. O.: Brown Carbon Formation by Aqueous-Phase Carbonyl Compound Reactions with Amines and Ammonium Sulfate, *Environmental Science & Technology*, 48, 985-993, 10.1021/es4038325, 2014.
- Pratap, V., Battaglia, M. A., Carlton, A. G., and Hennigan, C. J.: No evidence for brown carbon formation in ambient particles undergoing atmospherically relevant drying, *Environmental Science-Processes &*
595 *Impacts*, 22, 442-450, 10.1039/c9em00457b, 2020.
- Psichoudaki, M. and Pandis, S. N.: Atmospheric Aerosol Water-Soluble Organic Carbon Measurement: A Theoretical Analysis, *Environmental Science & Technology*, 47, 9791-9798, 10.1021/es402270y, 2013.
- Pye, H. O. T., Zuend, A., Fry, J. L., Isaacman-VanWertz, G., Capps, S. L., Appel, K. W., Foroutan, H., Xu, L., Ng, N. L., and Goldstein, A. H.: Coupling of organic and inorganic aerosol systems and the effect on gas-
600 particle partitioning in the southeastern US, *Atmos. Chem. Phys.*, 18, 357-370, 10.5194/acp-18-357-2018, 2018.
- Pye, H. O. T., Nenes, A., Alexander, B., Ault, A. P., Barth, M. C., Clegg, S. L., Collett, J. L., Fahey, K. M., Hennigan, C. J., Herrmann, H., Kanakidou, M., Kelly, J. T., Ku, I. T., McNeill, V. F., Riemer, N., Schaefer, T., Shi, G. L., Tilgner, A., Walker, J. T., Wang, T., Weber, R., Xing, J., Zaveri, R. A., and Zuend, A.: The
605 acidity of atmospheric particles and clouds, *Atmospheric Chemistry and Physics*, 20, 4809-4888, 10.5194/acp-20-4809-2020, 2020.
- Qin, J., Zhang, L., Qin, Y., Shi, S., Li, J., Gao, Y., Tan, J., and Wang, X.: pH-Dependent Chemical Transformations of Humic-Like Substances and Further Cognitions Revealed by Optical Methods, *Environmental Science & Technology*, 56, 7578-7587, 10.1021/acs.est.1c07729, 2022a.

- 610 Qin, Y., Qin, J., Wang, X., Xiao, K., Qi, T., Gao, Y., Zhou, X., Shi, S., Li, J., Gao, J., Zhang, Z., Tan, J., Zhang, Y., and Chen, R.: Measurement report: Investigation of pH- and particle-size-dependent chemical and optical properties of water-soluble organic carbon: implications for its sources and aging processes, *Atmos. Chem. Phys.*, 22, 13845-13859, doi.10.5194/acp-22-13845-2022, 2022b.
- Qin, Y., Qin, J., Zhou, X., Yang, Y., Chen, R., Tan, J., Xiao, K., and Wang, X., Effects of pH on light absorption
615 properties of water-soluble organic compounds in particulate matter emitted from typical emission sources, *Journal of Hazardous Materials*, 424, 127688, doi.org/10.1016/j.jhazmat.2021.127688, 2022c.
- Reid, J. P., Bertram, A. K., Topping, D. O., Laskin, A., Martin, S. T., Petters, M. D., Pope, F. D., and Rovelli, G.: The viscosity of atmospherically relevant organic particles, *Nature Communications*, 9, 956, 10.1038/s41467-018-03027-z, 2018.
- 620 Rusumdar, A. J., Tilgner, A., Wolke, R., and Herrmann, H.: Treatment of non-ideality in the SPACCIM multiphase model – Part 2: Impacts on the multiphase chemical processing in deliquesced aerosol particles, *Atmos. Chem. Phys.*, 20, 10351-10377, 10.5194/acp-20-10351-2020, 2020.
- Saleh, R.: From Measurements to Models: Toward Accurate Representation of Brown Carbon in Climate Calculations, *Current Pollution Reports*, 6, 90-104, 10.1007/s40726-020-00139-3, 2020.
- 625 Saleh, R., Hennigan, C. J., McMeeking, G. R., Chuang, W. K., Robinson, E. S., Coe, H., Donahue, N. M., and Robinson, A. L.: Absorptivity of brown carbon in fresh and photo-chemically aged biomass-burning emissions, *Atmospheric Chemistry and Physics*, 13, 7683-7693, 10.5194/acp-13-7683-2013, 2013.
- Schendorf, T. M., Del Vecchio, R., Bianca, M., and Blough, N. V.: Combined Effects of pH and Borohydride Reduction on Optical Properties of Humic Substances (HS): A Comparison of Optical Models,
630 *Environmental Science & Technology*, 53, 6310-6319, 10.1021/acs.est.9b01516, 2019.
- Schnitzler, E. G. and Abbatt, J. P. D.: Heterogeneous OH oxidation of secondary brown carbon aerosol, *Atmospheric Chemistry and Physics*, 18, 14539-14553, 10.5194/acp-18-14539-2018, 2018.
- Schwab, J. J., Wolfe, D., Casson, P., Brandt, R., Demerjian, K. L., Husain, L., Dutkiewicz, V. A., Civerolo, K. L., and Rattigan, O. V.: Atmospheric Science Research at Whiteface Mountain, NY: Site Description and
635 History, *Aerosol and Air Quality Research*, 16, 827-840, 10.4209/aaqr.2015.05.0343, 2016.
- Shah, V., Jacob, D. J., Moch, J. M., Wang, X., and Zhai, S.: Global modeling of cloud water acidity, precipitation acidity, and acid inputs to ecosystems, *Atmos. Chem. Phys.*, 20, 12223-12245, 10.5194/acp-20-12223-2020, 2020.

- 640 Srinivas, B., Rastogi, N., Sarin, M. M., Singh, A., and Singh, D.: Mass absorption efficiency of light absorbing organic aerosols from source region of paddy-residue burning emissions in the Indo-Gangetic Plain, *Atmospheric Environment*, 125, 360-370, <https://doi.org/10.1016/j.atmosenv.2015.07.017>, 2016.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059-2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- 645 Tao, Y. and Murphy, J. G.: The sensitivity of PM_{2.5} acidity to meteorological parameters and chemical composition changes: 10-year records from six Canadian monitoring sites, *Atmospheric Chemistry and Physics*, 19, 9309-9320, 10.5194/acp-19-9309-2019, 2019.
- Tao, Y. and Murphy, J. G.: Simple Framework to Quantify the Contributions from Different Factors Influencing Aerosol pH Based on NH_x Phase-Partitioning Equilibrium, *Environmental Science & Technology*, 55, 650 10310-10319, 10.1021/acs.est.1c03103, 2021.
- Tilgner, A., Schaefer, T., Alexander, B., Barth, M., Collett Jr, J. L., Fahey, K. M., Nenes, A., Pye, H. O. T., Herrmann, H., and McNeill, V. F.: Acidity and the multiphase chemistry of atmospheric aqueous particles and clouds, *Atmos. Chem. Phys.*, 21, 13483-13536, 10.5194/acp-21-13483-2021, 2021.
- Turnock, S. T., Mann, G. W., Woodhouse, M. T., Dalvi, M., O'Connor, F. M., Carslaw, K. S., and Spracklen, D. 655 V.: The Impact of Changes in Cloud Water pH on Aerosol Radiative Forcing, *Geophysical Research Letters*, 46, 4039-4048, 10.1029/2019gl082067, 2019.
- Washenfelder, R. A., Flores, J. M., Brock, C. A., Brown, S. S., and Rudich, Y.: Broadband measurements of aerosol extinction in the ultraviolet spectral region, *Atmos. Meas. Tech.*, 6, 861-877, 10.5194/amt-6-861-2013, 2013.
- 660 Wu, C., Wang, G., Li, J., Li, J., Cao, C., Ge, S., Xie, Y., Chen, J., Li, X., Xue, G., Wang, X., Zhao, Z., and Cao, F.: The characteristics of atmospheric brown carbon in Xi'an, inland China: sources, size distributions and optical properties, *Atmos. Chem. Phys.*, 20, 2017-2030, 10.5194/acp-20-2017-2020, 2020.
- Yang, J., Au, W. C., Law, H., Leung, C. H., Lam, C. H., and Nah, T.: pH affects the aqueous-phase nitrate-mediated photooxidation of phenolic compounds: implications for brown carbon formation and evolution, 665 *Environmental Science: Processes & Impacts*, 25, 176-189, 10.1039/D2EM00004K, 2023.
- Yu, L., Smith, J., Laskin, A., George, K. M., Anastasio, C., Laskin, J., Dillner, A. M., and Zhang, Q.: Molecular transformations of phenolic SOA during photochemical aging in the aqueous phase: competition among oligomerization, functionalization, and fragmentation, *Atmospheric Chemistry and Physics*, 16, 4511-4527, 10.5194/acp-16-4511-2016, 2016.

- 670 Zeng, L., Zhang, A., Wang, Y., Wagner, N. L., Katich, J. M., Schwarz, J. P., Schill, G. P., Brock, C., Froyd, K. D.,
Murphy, D. M., Williamson, C. J., Kupc, A., Scheuer, E., Dibb, J., and Weber, R. J.: Global Measurements
of Brown Carbon and Estimated Direct Radiative Effects, *Geophysical Research Letters*, 47,
e2020GL088747, <https://doi.org/10.1029/2020GL088747>, 2020.
- Zhang, K., O'Donnell, D., Kazil, J., Stier, P., Kinne, S., Lohmann, U., Ferrachat, S., Croft, B., Quaas, J., Wan, H.,
675 Rast, S., and Feichter, J.: The global aerosol-climate model ECHAM-HAM, version 2: sensitivity to
improvements in process representations, *Atmos. Chem. Phys.*, 12, 8911-8949, 10.5194/acp-12-8911-
2012, 2012.
- Zhang, X. L., Lin, Y. H., Surratt, J. D., and Weber, R. J.: Sources, Composition and Absorption Angstrom Exponent
of Light-absorbing Organic Components in Aerosol Extracts from the Los Angeles Basin, *Environmental*
680 *Science & Technology*, 47, 3685-3693, 10.1021/es305047b, 2013.
- Zhang, Y. Z., Forrister, H., Liu, J. M., Dibb, J., Anderson, B., Schwarz, J. P., Perring, A. E., Jimenez, J. L.,
Campuzano-Jost, P., Wang, Y. H., Nenes, A., and Weber, R. J.: Top-of-atmosphere radiative forcing
affected by brown carbon in the upper troposphere, *Nature Geoscience*, 10, 486+, 10.1038/ngeo2960,
2017.
- 685 Zhao, R., Lee, A. K. Y., Huang, L., Li, X., Yang, F., and Abbatt, J. P. D.: Photochemical processing of aqueous
atmospheric brown carbon, *Atmospheric Chemistry and Physics*, 15, 6087-6100, 10.5194/acp-15-6087-
2015, 2015.
- Zheng, G. J., Su, H., Wang, S. W., Andreae, M. O., Poschl, U., and Cheng, Y. F.: Multiphase buffer theory explains
690 contrasts in atmospheric aerosol acidity, *Science*, 369, 1374+, 10.1126/science.aba3719, 2020.