



Seasonal variations in photooxidant formation and light absorption in aqueous extracts of ambient particles

Lan Ma^{1a}, Reed Worland^{1b}, Laura Heinlein¹, Chrystal Guzman^{1c}, Wenqing Jiang², Christopher Niedek², Keith J. Bein³, Qi Zhang², Cort Anastasio¹

5 ¹Department of Land, Air and Water Resources, University of California, Davis, One Shields Avenue, Davis, CA 95616-8627, USA

²Department of Environmental Toxicology, University of California, Davis, One Shields Avenue, Davis, CA 95616-8627, USA

³Center for Health and the Environment, University of California, Davis, One Shields Avenue, Davis, CA 95616-8627, USA

^aNow at: SGS-CSTC Standards Technical Services Co.,Ltd. Hangzhou Branch, Hangzhou, Zhejiang Province, 310052, China

10 ^bNow at: Department of Chemistry, University of Washington, WA, USA

^cNow at: Department of Pharmacology, University of Washington, WA, USA

Correspondence to: Cort Anastasio (canastasio@ucdavis.edu)

Abstract. Atmospheric waters – including fog/cloud drops and aerosol liquid water – are important sites for the transformations of atmospheric species, largely through reactions with photoformed oxidants such as hydroxyl radical ($\bullet\text{OH}$), singlet molecular oxygen ($^1\text{O}_2^*$), and oxidizing triplet excited states of organic matter ($^3\text{C}^*$). Despite this, there are few measurements of these photooxidants, especially in extracts of ambient particles, and very little information about how oxidant levels vary with season or particle type. To address this gap, we collected ambient $\text{PM}_{2.5}$ from Davis, California over the course of a year and measured photooxidant concentrations in dilute aqueous extracts of the particles. We categorized samples into four groups: Winter & Spring (Win-Spr), Summer & Fall (Sum-Fall) without wildfire influence, fresh biomass burning (FBB), and aged biomass burning (ABB). FBB contains significant amounts of brown carbon (BrC) from wildfires, and the highest mass absorption coefficients (MAC) normalized by dissolved organic carbon, with an average ($\pm 1 \sigma$) value of $3.3 (\pm 0.4) \text{ m}^2 (\text{g C})^{-1}$ at 300 nm. Win-Spr and ABB have similar MAC averages, $1.9 (\pm 0.4)$ and $1.5 (\pm 0.3) \text{ m}^2 (\text{g C})^{-1}$, respectively, while Sum-Fall has the lowest MAC_{DOC} ($0.65 (\pm 0.19) \text{ m}^2 (\text{g C})^{-1}$). $\bullet\text{OH}$ concentrations in extracts range from $(0.2\text{-}4.7) \times 10^{-15} \text{ M}$ and generally increase with concentration of dissolved organic carbon (DOC), although this might be because DOC is a proxy for extract concentration. The average quantum yield for $\bullet\text{OH}$ formation (Φ_{OH}) across all sample types is $3.7 (\pm 2.4) \%$, with no statistical difference among sample types. $^1\text{O}_2^*$ concentrations have a range of $(0.7\text{-}45) \times 10^{-13} \text{ M}$, exhibiting a good linearity with DOC that is independent of sample type ($R^2 = 0.93$). Fresh BB samples have the highest $[^1\text{O}_2^*]$ but the lowest average $\Phi_{1\text{O}_2^*}$, while Sum-Fall samples are the opposite. $\Phi_{1\text{O}_2^*}$ is negatively correlated with MAC_{DOC} , indicating that less light-absorbing samples form $^1\text{O}_2^*$ more efficiently. We quantified $^3\text{C}^*$ concentrations with two triplet probes: syringol (SYR), which captures both strongly and weakly oxidizing triplets, and (phenylthio)acetic acid (PTA), which is only sensitive to strongly oxidizing triplets. Concentrations of $^3\text{C}^*$ are in the range of $(0.03 - 7.9) \times 10^{-13} \text{ M}$ and linearly increase with DOC ($R^2 = 0.85$ for SYR and $R^2 = 0.80$ for PTA); this relationship for $[^3\text{C}^*]_{\text{SYR}}$ is independent of sample type. The average ratio of $[^3\text{C}^*]_{\text{PTA}}/[^3\text{C}^*]_{\text{SYR}}$ is $0.58 (\pm 0.38)$, indicating that roughly 60% of oxidizing triplets are strongly oxidizing. Win-Spr samples have the highest fraction of strongly oxidizing $^3\text{C}^*$, with an average of $86 (\pm 43)\%$. $\Phi_{3\text{C}^*,\text{SYR}}$ is in the range of $(0.6\text{-}8.8) \%$, with an average value, $3.3 (\pm 1.9)\%$, two times higher than $\Phi_{3\text{C}^*,\text{PTA}}$. FBB has the lowest average $\Phi_{3\text{C}^*}$, while the aging process tends to enhance $\Phi_{3\text{C}^*}$, as well as $\Phi_{1\text{O}_2^*}$.

To estimate photooxidant concentrations in particle water, we extrapolate the photooxidant kinetics in our dilute particle extracts to aerosol liquid water (ALW) conditions of $1 \mu\text{g PM}/\mu\text{g H}_2\text{O}$ for each sample type. The estimated ALW $\bullet\text{OH}$ concentration is $7 \times 10^{-15} \text{ M}$ when including mass transport of gas-phase $\bullet\text{OH}$ to the particles. $^1\text{O}_2^*$ and $^3\text{C}^*$ concentrations in ALW have ranges of



40 $(0.6 - 7) \times 10^{-12}$ M and $(0.08 - 1) \times 10^{-12}$ M, respectively. In the Win-Spr and Sum-Fall samples, photooxidant concentrations increase significantly from lab particle extracts to ALW, while the changes for the FBB and ABB samples are minor. The small increases in $^1\text{O}_2^*$ and $^3\text{C}^*$ from extract to ALW for the biomass burning particles are likely due to the high amounts of organic compounds in the extracts, which lead to strong quenching of these oxidants even under our dilute conditions. Compared to the photooxidant concentration estimates in Kaur et al. (2019), our updated ALW estimates show higher $\bullet\text{OH}$ (by roughly a factor of
45 10), higher $^3\text{C}^*$ (by factors of 1-5) and lower $^1\text{O}_2^*$ concentrations (by factors of 20-100). Our results indicate that $^3\text{C}^*$ and $^1\text{O}_2^*$ in ALW dominate the processing of organic compounds that react quickly with these oxidants (such as phenols and furans, respectively), while $\bullet\text{OH}$ is more important for less reactive organics.

1 Introduction

Atmospheric waters, including fog/cloud drops and liquid water on aerosol particles, are important media for photochemical
50 transformations of chemical species (Herrmann et al., 2010, 2015). These include formation of aqueous secondary organic aerosol (aqSOA), formation and photobleaching of brown carbon (BrC), oxidation of reduced sulfur, and aerosol aging (Ervens, 2018; Ervens et al., 2011; Gilardoni et al., 2016; Laskin et al., 2015; McNeill, 2015; Seinfeld and Pandis, 2016; Wang et al., 2016; Zhao et al., 2015). Many of these processes are driven by photochemically generated oxidants, including hydroxyl radical ($\bullet\text{OH}$), triplet excited states of organic matter ($^3\text{C}^*$), and singlet molecular oxygen ($^1\text{O}_2^*$) (Ervens et al., 2014; Finlayson-Pitts and Pitts, 2000;
55 He et al., 2013; Herrmann, 2003; Kaur et al., 2019; Lim et al., 2010).

Hydroxyl radical ($\bullet\text{OH}$), the best studied aqueous oxidant in the atmosphere, is highly reactive with most reduced species but has a relatively low abundance compared to $^3\text{C}^*$ and $^1\text{O}_2^*$. Concentrations of $\bullet\text{OH}$ in fog and cloud waters, as well as aqueous extracts of ambient particles and lab-generated secondary organic aerosol, are typically 10^{-17} to 10^{-15} M (Anastasio and McGregor, 2001;
60 Arakaki et al., 2013; Dorfman and Adams, 1973; Kaur and Anastasio, 2017; Kaur et al., 2019; Manfrin et al., 2019; Tilgner and Herrmann, 2018). Sources of $\bullet\text{OH}$ in the aqueous phase include mass transfer from the gas phase, Fenton or Fenton-like reactions of reduced metals with hydrogen peroxide, and photolysis of nitrate, nitrite, iron complexes, hydrogen peroxide, and organic hydroperoxides (Badali et al., 2015; Herrmann et al., 2010; Tilgner and Herrmann, 2018; Tong et al., 2016). Additionally, organic compounds in atmospheric waters can affect $\bullet\text{OH}$ production. For example, the interaction of humic-like substances (HULIS) or
65 SOA with Fe(II) can enhance or suppress $\bullet\text{OH}$ formation (Baba et al., 2015; Gonzalez et al., 2017; Hems and Abbatt, 2018; Tong et al., 2016; Zuo and Hoigne, 1992). This suggests that seasonal variations in particle composition (e.g., SOA and Fe) can affect $\bullet\text{OH}$ kinetics, as reported recently for $\bullet\text{OH}$ photoproduction in extracts of particulate matter (PM) from Colorado: winter $\bullet\text{OH}$ originated from nitrate photolysis, while summer $\bullet\text{OH}$ was more linked to soluble iron (Leresche et al., 2021). But little is known about how $\bullet\text{OH}$ concentrations in particles vary with season or among particle types.

70

Triplet excited states ($^3\text{CDOM}^*$) are formed when organic chromophores (i.e., brown carbon (BrC)) absorb sunlight and are promoted to a higher energy state (McNeill and Canonica, 2016). Oxidizing triplets ($^3\text{C}^*$), i.e., the subset of triplets that have high reduction potentials, are effective oxidants, reacting with phenols and biogenic volatile compounds to form SOA and BrC, and oxidizing bisulfite to sulfate (González Palacios et al., 2016; Monge et al., 2012; Rossignol et al., 2014; Smith et al., 2014; Wang
75 et al., 2020b; Yu et al., 2014). For compounds (like phenols) that react rapidly with triplets, $^3\text{C}^*$ can be as important an oxidant as $\bullet\text{OH}$ in cloud and fog drops, where oxidizing triplet concentrations are 10^{-15} - 10^{-13} M (Kaur and Anastasio, 2018; Kaur et al., 2019; Ma et al., 2021; Smith et al., 2015). Moreover, triplet concentrations are estimated to be enhanced by one or two orders of



magnitude in aerosol liquid water (Kaur et al., 2019; Ma et al., 2023a). The ability of dissolved organic matter (DOM) to form $^3\text{C}^*$ depends on its composition. In surface waters, quantum yields of $^3\text{C}^*$ are higher for organic compounds with lower average
80 molecular weights and lower aromaticity (Berg et al., 2019; Maizel and Remucal, 2017; McCabe and Arnold, 2017, 2018; Mckay et al., 2017). However, little is known about how $^3\text{C}^*$ formation in atmospheric waters depends on BrC characteristics or season.

The final oxidant we consider, singlet molecular oxygen ($^1\text{O}_2^*$), is formed when triplet excited states transfer energy to dissolved molecular oxygen. $^1\text{O}_2^*$ reacts rapidly with electron-rich compounds such as furans, polycyclic aromatic hydrocarbons, some
85 amino acids, and substituted alkenes (Gollnick and Griesbeck, 1985; McGregor and Anastasio, 2001; Richards-Henderson et al., 2015; Wilkinson et al., 1995; Zeinali et al., 2019). $^1\text{O}_2^*$ concentrations in fog and cloud waters and aqueous particle extracts are the highest of the three oxidants, in the range of 10^{-14} - 10^{-12} M (Bogler et al., 2022; Kaur and Anastasio, 2017; Kaur et al., 2019; Leresche et al., 2021; Manfrin et al., 2019). Dissolved black carbon also can produce $^1\text{O}_2^*$, resulting in concentrations on the order of 10^{-12} M (Li et al., 2019). Though $^1\text{O}_2^*$ is not as reactive as $^3\text{C}^*$ and $\bullet\text{OH}$, its concentration increases by orders of magnitude
90 when moving from dilute cloud/fog conditions towards the more concentrated conditions of aerosol liquid water (Kaur et al., 2019; Ma et al., 2023a). Since $^1\text{O}_2^*$ is born from $^3\text{CDOM}^*$, these two oxidants are tightly linked. For example, in surface waters the quantum yield of $^1\text{O}_2^*$ ($\Phi_{1\text{O}_2^*}$) is also higher in samples with lower average molecular weight DOM, as seen for $^3\text{C}^*$ (Berg et al., 2019; Maizel and Remucal, 2017; Ossola et al., 2021; Wang et al., 2020a). Some studies on the seasonal trend of $\Phi_{1\text{O}_2^*}$ in surface waters hypothesized that summer samples where photodegradation is more rapid have higher $\Phi_{1\text{O}_2^*}$ based on DOM
95 photodegradation increasing $^1\text{O}_2^*$ quantum yields (McCabe and Arnold, 2016; Ossola et al., 2021; Sharpless et al., 2014). However, there are differences in singlet oxygen generation and concentrations between surface and atmospheric waters. For example, while ozonation and photodegradation of DOM enhances $\Phi_{1\text{O}_2^*}$ in surface waters, photodegradation of aqueous particle extracts has no significant effect on $\Phi_{1\text{O}_2^*}$ (Leresche et al., 2019, 2021; Sharpless et al., 2014). In addition, we know very little about the seasonality of $^1\text{O}_2^*$ concentrations in particles or how this oxidant varies between particle types.

100

Although $\bullet\text{OH}$, $^3\text{C}^*$, and $^1\text{O}_2^*$ are important in the transformation of atmospheric species, there are few measurements of these photooxidants in atmospheric condensed phases, especially in extracts of ambient particles. In addition, very little is known about seasonal variations in these oxidant concentrations and kinetics. To address this gap, we collected $\text{PM}_{2.5}$ from November 2019 to October 2020 in Davis CA, extracted them in water, and measured light absorption and photooxidant formation. This period
105 included four main types of samples: winter samples influenced by residential wood combustion and high humidity, summer samples impacted by nearby wildfires (i.e., fresh biomass burning (BB) particles), summer samples impacted by more distant wildfires (i.e., aged BB particles), and spring/summer samples with little to no biomass burning. We measured photooxidant concentrations ($\bullet\text{OH}$, $^1\text{O}_2^*$, $^3\text{C}^*$) in water extracts of the particles, and investigated how photooxidant formation depends on particle type, optical properties, and biomass burning influence. Finally, we extrapolated our dilute extract results to predict photooxidant
110 concentrations in aerosol liquid water (ALW) and assessed the importance of photooxidants in processing particulate organic compounds.

2 Experimental methods

2.1 Chemicals

Furfuryl alcohol (FFA, 98%), benzoic acid (BA, $\geq 99.5\%$), *p*-hydroxybenzoic acid (*p*-HBA, 99%), (phenylthio)acetic acid (PTA, 96%), syringol (SYR, 99%), 3,4-dimethoxybenzaldehyde (DMB, 99%), and deuterium oxide (D_2O , 99.9% D-atom) were received
115



from Millipore Sigma. All chemical solutions and particulate matter extracts were prepared using air-saturated ultrapure water (Milli-Q water) from a Milli-Q Advantage A10 system (Millipore; $\geq 18.2 \text{ M}\Omega \text{ cm}$) that was pretreated with a Barnstead activated carbon cartridge.

2.2 Particle collection and extraction

120 More detailed descriptions of sampling and extraction procedures are provided in Ma et al. (2023a) and are only briefly discussed here. Fine particle ($\text{PM}_{2.5}$) sampling was conducted from November 2019 to October 2020 on the roof of Ghauri Hall on the University of California, Davis campus. Winter in Davis is humid and sometimes foggy, and the air quality is often impacted by residential wood combustion, while Davis in summer is hot and dry. During the summer of 2020, several severe wildfires occurred in Northern California and Oregon, including the largest wildfires in the recorded history of California: the August complex (size: 125 4179 km^2), LNU Lightning complex (1605 km^2), and SCU lightning complex (1470 km^2) (<https://www.fire.ca.gov/incidents/2020>; last access: 15 July 2022). These fires caused extremely heavy air pollution in Davis with daily $\text{PM}_{2.5}$ concentrations sometimes exceeding $80 \mu\text{g m}^{-3}$ (<https://www.arb.ca.gov/aqmis2/aqmis2.php>, last access: 20 June 2022). Particles were collected with a high-volume sampler containing a PM_{10} inlet (Graseby Andersen) and two offset, slotted impactor plates (Tisch Environmental, Inc., 230 series) to remove particles larger than $2.5 \mu\text{m}$. $\text{PM}_{2.5}$ was collected onto pre-cleaned Teflon-coated borosilicate glass microfiber 130 filters (Pall Corporation, EmFab™ filters, 8 in. \times 10 in.) and stored at $-20 \text{ }^\circ\text{C}$ immediately after collection. The sampling duration was either 24 hr or up to a week (Table S1). The sampling campaign was paused from March to June 2020 because of COVID-related restrictions on campus activities.

To prepare particulate matter extracts (PMEs), filters were cut into $2 \text{ cm} \times 2 \text{ cm}$ squares, and then extracted with 1.0 mL Milli-Q 135 water by shaking for 4 h in the dark. The extracts from the same filter were combined, filtered ($0.22 \mu\text{m}$ PTFE; Pall), and adjusted to pH 4.2 by sulfuric acid to mimic the acidity of winter particle water in the Central Valley of California (Parworth et al., 2017). The acidity of extracts was measured by a pH microelectrode (MI-414 series, protected tip; Microelectrodes, Inc.). PMEs were flash-frozen in liquid nitrogen immediately after preparation and were later thawed on the day of the experiment. Particle mass extracted was determined by weighing filter squares before and after extraction with a microbalance (M2P, Sartorius); the extracted 140 mass is an upper bound because we cannot account for insoluble material that is extracted from the square but removed by subsequent filtration. UV-Vis spectra of PMEs were measured with a Shimadzu UV-2501PC spectrophotometer. Dissolved organic carbon (DOC) and major ions were measured by a total organic carbon analyzer (TOC-VCPH, Shimadzu) and ion chromatographs (881 Compact IC Pro, Metrohm) equipped with conductivity detectors, respectively. PME sample information is provided in Table S1, while DOC and ion concentrations are in Table S2.

145 2.3 Sample illumination and chemical analysis

Illumination experiments were conducted using light from a 1000 W xenon arc lamp that was passed through optical filters to simulate tropospheric sunlight (Kaur and Anastasio, 2017). 1.0 mL of extract at pH 4.2 was spiked with a photooxidant probe and illuminated in a silicone-plugged GE 021 quartz tube (5 mm inner diameter, 1.0 mL volume) at $20 \text{ }^\circ\text{C}$. Dark control samples were wrapped in aluminum foil and kept in the same photoreactor chamber. During illumination, aliquots were removed from the 150 illuminated and dark tubes periodically to measure probe concentrations with high-performance liquid chromatography (HPLC, Shimadzu LC-20AB pump, Thermo Scientific Accucore XL C18 column ($50 \times 3 \text{ mm}$, $4 \mu\text{m}$ bead), and Shimadzu-M20A UV-Vis detector). The photon flux in an identical quartz tube was determined on each experiment day by measuring the photolysis rate constant of a $10 \mu\text{M}$ 2-nitrobenzaldehyde (2NB) solution (Galbavy et al., 2010).



2.4 Photooxidant measurements

155 Photooxidant methods are detailed in past papers (Anastasio and McGregor, 2001; Kaur and Anastasio, 2017; Ma et al., 2023a) and are only briefly described here. The uncertainty on an individual oxidant concentration is 1 standard error, determined by propagating the errors of the individual parameters required to calculate the concentration. Uncertainties on average values are 1 standard deviation, calculated from the spread of the individual values.

2.4.1 Hydroxyl radical ($\bullet\text{OH}$)

160 $\bullet\text{OH}$ concentration was quantified using 10 μM benzoic acid (BA) as a probe and simultaneously monitoring the rates of probe decay and product (*p*-hydroxybenzoic acid, *p*-HBA) formation. For dilute samples ($\text{DOC} < 15 \text{ mg C L}^{-1}$), 2 μM BA was used in order not to perturb the natural $\bullet\text{OH}$ sink in PME. Aliquots were taken during illumination to measure BA and *p*-HBA concentrations. From the BA probe loss, a linear regression of $\ln([\text{BA}]_t/[\text{BA}]_0)$ versus illumination time (t) was fitted, where $[\text{BA}]_0$ is the concentration at time zero. The negative value of the regression slope is the BA pseudo-first order decay rate constant (k'_{BA}).

165 The $\bullet\text{OH}$ concentration was then determined using:

$$[\bullet\text{OH}]_{\text{exp}} = \left[\frac{k'_{\text{BA}}}{k_{\text{BA}+\bullet\text{OH}}} \right] \quad (1)$$

where $k_{\text{BA}+\bullet\text{OH}}$ is the second-order rate constant of BA reacting with $\bullet\text{OH}$ at pH 4.2 ($5.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) (Ashton et al., 1995; Wander et al., 1968). Next, $[\bullet\text{OH}]_{\text{exp}}$ was normalized to sunlight conditions at midday on the winter solstice at Davis (solar zenith = 62° ; $j_{2\text{NB},\text{win}} = 0.0070 \text{ s}^{-1}$) (Galbavy et al., 2010) and corrected for internal light screening due to absorption by chromophores in PME:

$$170 \quad [\bullet\text{OH}]_{\text{win}} = \left[\frac{[\bullet\text{OH}]_{\text{exp}}}{S_\lambda \times j_{2\text{NB},\text{exp}}} \right] \times j_{2\text{NB},\text{win}} \quad (2)$$

where S_λ is the internal light screening factor in an individual sample (Table S1), and $j_{2\text{NB},\text{exp}}$ is the photolysis rate constant of 2NB measured on the experiment day.

We also determined the $\bullet\text{OH}$ concentration in each sample from *p*-HBA formation. The initial formation rate of *p*-HBA was determined from the regression between *p*-HBA concentration and illumination time, either using a linear regression or a three-parameter exponential fit:

$$[p\text{-HBA}]_t = [p\text{-HBA}]_0 + a(1 - e^{-bt}) \quad (3)$$

where $[p\text{-HBA}]_t$ and $[p\text{-HBA}]_0$ are the measured concentrations at illumination times t and zero, respectively, and a and b are regression fit parameters. With this fitting, the initial formation rate of *p*-HBA, R_p , is calculated with:

$$180 \quad R_p = a \times b \quad (4)$$

and then the $\bullet\text{OH}$ concentration was calculated using:

$$[\bullet\text{OH}]_{\text{exp}} = \frac{R_p}{[\text{BA}]_0 \times k_{\text{BA}+\bullet\text{OH}} \times Y_{p\text{-HBA}}} \quad (5)$$

where $Y_{p\text{-HBA}}$ (0.18) is the yield of *p*-HBA from the reaction of BA with $\bullet\text{OH}$ (Anastasio and McGregor, 2001). $\bullet\text{OH}$ concentrations were normalized by $j_{2\text{NB}}$ and light screening factor using Eq.2. In some samples, BA decay and *p*-HBA formation were faster at the beginning of illumination and then slowed (e.g., Fig. S1), indicating an initially higher $\bullet\text{OH}$ concentration compared to later times, as seen previously (Paulson et al., 2019). For each sample we generally used all data points for the regressions of BA and *p*-HBA and then determined the reported $[\bullet\text{OH}]$ as the average of the BA and *p*-HBA results (Table S3).



2.4.2 Singlet molecular oxygen ($^1\text{O}_2^*$)

To determine $^1\text{O}_2^*$ concentrations, FFA was used as a probe and deuterium oxide (D_2O) was used as a diagnostic tool (Anastasio and McGregor, 2001) because $^1\text{O}_2^*$ decays more rapidly in H_2O than D_2O . Therefore, the difference of FFA decay rates in H_2O and D_2O is attributed to $^1\text{O}_2^*$ (instead of other oxidants). For each sample, 1.0 mL of PME was divided into two 0.5 mL aliquots, with one diluted with 0.5 mL H_2O and the other 0.5 mL D_2O . 10 μM FFA was spiked into both solutions and pseudo-first order rate constants of FFA loss during illumination were determined ($k_{\text{exp,H}_2\text{O}}$ and $k_{\text{exp,D}_2\text{O}}$). The difference between the FFA first-order rate constants was used to calculate the steady-state $^1\text{O}_2^*$ concentration (Anastasio and McGregor, 2001). This experimental $^1\text{O}_2^*$ concentration was normalized by photon flux and light screening factors of PME using an analog of Eq. 2 to determine $^1\text{O}_2^*$ winter-solstice values (Table S4).

2.4.3 Oxidizing triplet excited states of organic matter ($^3\text{C}^*$)

Oxidizing triplets were measured with two probes, syringol (SYR) and (phenylthio)acetic acid (PTA). SYR reacts rapidly with all oxidizing triplets, but its decay by $^3\text{C}^*$ can be inhibited by high concentrations of dissolved organic matter (DOM) (Ma et al., 2023a, 2023b; Maizel and Remucal, 2017; McCabe and Arnold, 2017). In contrast, PTA is more resistant to this inhibition, but it can only capture strongly oxidizing triplets (Ma et al., 2023b). To determine $^3\text{C}^*$ concentrations, two 1.0 mL aliquots of PME were spiked with 10 μM of either SYR or PTA, and then illuminated to determine the pseudo-first order rate constant for loss of each probe ($k'_{\text{P,exp}}$). We then removed the contributions of direct photodegradation, $\bullet\text{OH}$, and $^1\text{O}_2^*$ to triplet probe decay (Ma et al., 2023a). Since $^3\text{C}^*$ is a complex mixture of triplets with a wide range of reactivities, there is no exact value for the second-order rate constant of $^3\text{C}^*$ in PME reacting with probes. Our past work indicated that $^3\text{C}^*$ in Davis winter PM have a similar average reactivity to the triplet state of DMB (Kaur and Anastasio, 2018; Kaur et al., 2019), which is a component of BB BrC (Fleming et al., 2020; Schauer et al., 2001). However, it is possible that this model compound is more reactive than natural oxidizing triplets, which would lead to an underestimate of $^3\text{C}^*$ (Ma et al., 2023b). We quantified the inhibition effect of DOM on the decay of SYR and PTA by measuring inhibition factors of each probe ($IF_{\text{P,corr}}$) in each sample, and used them to correct $^3\text{C}^*$ concentrations (Canonica and Laubscher, 2008; Ma et al., 2023b; McCabe and Arnold, 2017; Wenk et al., 2011). Details about determining inhibition factors and correcting $^3\text{C}^*$ concentrations are provided in Supplemental Information Section S1. $^3\text{C}^*$ concentrations in PME during each experiment were calculated with:

$$[^3\text{C}^*]_{\text{P,exp}} = \frac{k'_{\text{P},3\text{C}^*}}{k_{\text{P}+3\text{DMB}^*} \times IF_{\text{P,corr}}} \quad (6)$$

where $k_{\text{P}+3\text{DMB}^*}$ is the second-order rate constant of probe with $^3\text{DMB}^*$ (Table S5). These values were converted to $^3\text{C}^*$ concentrations expected on midday of the winter solstice in Davis (after correction for internal light screening) using an equation analogous to Eq. 2; these are the concentrations reported in the main text. Details of $^3\text{C}^*$ measurements by SYR and PTA are in Tables S7 and S8, respectively.

2.4.4 Extrapolating extract results to aerosol liquid water conditions

Photooxidant concentrations in PM extracts represent dilute conditions similar to cloud/fog waters, while our goal is to estimate photooxidant concentrations in aerosol liquid water, which is orders of magnitude more concentrated. To predict photooxidant concentrations in ALW, we quantified photooxidant kinetics (i.e., oxidant formation rates and loss rate constants) for each sample type as a function of particle mass concentration and then extrapolated to ALW conditions (Kaur et al., 2019; Ma et al., 2023a). Details about the extrapolations are provided in Section S4.




225 3. Results and Discussion

3.1 General extract characteristics

To investigate the seasonal variation of photooxidant formation, we studied 18 PM_{2.5} samples across a year of sampling. Samples were from all seasons, but there was only one Spring sample because of COVID restrictions from March through June of 2020 (Fig. 1 and Table S1). Most particle samples were collected for 24 h, while four of the winter samples were collected for seven
230 days to obtain more particle mass. Winters were marked by residential wood burning and high relative humidities, while the summer samples represented both periods influenced by fresh and aged biomass burning (from wildfires) and clean conditions. From August to October 2020, Davis periodically experienced severe air pollution caused by wildfires in California and Oregon. Section S2 of the supplement provides satellite images with fire points detected by the NASA Visible Infrared Imaging Radiometer Suite (VIIRS) and 24-h back trajectories estimated by the Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPPLIT) on
235 the day of sampling for wildfire periods (Rolph et al., 2017; Stein et al., 2015). Based on the satellite images and back trajectories, smoke plumes were transported from their sources to Davis in as short as 1~2 h or as long as 12~24 h or more.

Figure 1 shows the average PM_{2.5} concentration during each extract sampling period. We categorized the 18 samples into four groups based on sampling date and positive matrix factorization (PMF) results obtained using UV/Vis absorption spectra and
240 aerosol mass spectrometer chemical characterization (Jiang et al., 2023). The first group is termed Winter & Spring samples (Win-Spr), which were collected from November 2019 to March 2020 and have an average PM_{2.5} concentration of 9.9 (± 1.5) µg m⁻³ (Table S1). Three samples collected in July, August, and October without wildfire influence are classified as Summer & Fall samples (Sum-Fall), with an average PM_{2.5} of 7.4 (± 0.4) µg m⁻³. The seven wildfire-influenced samples collected from August to October are classified as fresh biomass burning (FBB) or aged biomass burning (ABB), with average PM_{2.5} values of 55 (± 10)
245 and 24 (± 8) µg m⁻³, respectively. The PMF results indicate that FBB samples are dominated by biomass-burning organic aerosol factors characterized by elevated levels of levoglucosan (m/z 60) signature ions in the AMS mass spectra (Alfarra et al., 2007). ABB samples were also collected during the wildfire-influenced period, but they are dominated by an oxidized organic aerosol factor with high O/C ratio and little levoglucosan (Jiang et al., 2023).

250 Our PM extracts are much more dilute than aerosol liquid water in the ambient atmosphere, a result of physical limitations on the amount of water we need to extract and study particle photochemistry. Particle mass/liquid water mass ratios of our extracts were in the range (0.7 – 4.1) × 10⁻⁴ µg PM/µg H₂O for one-day samples (Fig. S9) and correlated well with the ambient PM_{2.5} concentrations (Table S1). The seven-day winter samples had higher particle mass/water mass ratios, up to 9.1 × 10⁻⁴ µg PM/µg H₂O. Based on the PM mass concentrations, our particle extracts are similar to dilute atmospheric waters such as cloud and fog
255 drops (10⁻⁵ - 10⁻³ µg PM/µg H₂O), instead of concentrated particle liquid water (roughly 1 µg PM/µg H₂O) (Nguyen et al., 2016; Seinfeld and Pandis, 2016).



260 **Figure 1. Average PM_{2.5} concentrations (circles) during each sampling period and DOC-normalized mass absorption coefficients at 300 nm (x) in particle extracts for Winter & Spring samples (blue), Summer & Fall samples (green), Fresh biomass burning (red), and Aged biomass burning (yellow).**

Dissolved organic carbon concentrations in the extracts range from 5 to 192 mg C L⁻¹ (Table S1). The ratio of organic carbon (OC) mass to total extracted PM mass is high in the wildfire samples, with average values of 31 (± 6) % and 26 (± 6) % for FBB and ABB, respectively. These fractions are lower than values for BB particles in other studies (43-59%) (Schauer et al., 2001; Vicente et al., 2013), probably because we used water as the extraction solvent, thereby missing water insoluble organics. The OC/PM fractions for Win-Spr and Sum-Fall samples are similar to each other, with values of 16 (± 5) % and 11 (± 3) %, respectively. Win-Spr PMEs have high concentrations of nitrate (NO₃⁻), 84-3300 µM (Table S2), which contributed up to 33 % of PM mass. PMEs in the other three groups have nitrate concentrations from 25 to 300 µM, which are less than 10 % of PM mass. Win-Spr samples also have the highest ammonium concentrations, 168-4900 µM, followed by wildfire-influenced samples (46-803 µM), and Sum-Fall samples (< 100 µM). Potassium, a marker of biomass burning (Silva et al., 1999), has its highest concentrations in winter and wildfire samples with a range of 62-220 µM. The Sum-Fall samples have the highest fraction of sodium with an average of 11%, suggesting the influence of sea salt (Parworth et al., 2017). We employed three field blanks in this study at the beginning, middle, and end of the sampling campaign. In field blanks, ions and DOC concentrations are less than 10% of their concentrations in most PME samples, though FB1 was contaminated by the filling solution of a pH electrode, resulting in extremely high chloride concentrations (Table S2).

275 3.2 Light absorption in particle extracts

DOC-normalized mass absorption coefficients at 300 nm (MAC_{DOC,300}) are shown in Figure 1. For wildfire samples, MAC is correlated with the PM_{2.5} concentration, which probably reflects the dominant influence of BB emissions on both PM levels and light absorbance since FBB has the highest MAC among sample types, with an average of 3.3 (± 0.4) (g C)⁻¹. This is expected because fresh biomass burning organic aerosols (BBOA) contain abundant amounts of highly light-absorbing products, including substituted aromatics with high unsaturation and nitroaromatics (Budisulistiorini et al., 2017; Claeys et al., 2012; Fleming et al., 2020; Hettiyadura et al., 2021; Lin et al., 2016, 2017). The average MAC for FBB at 365 nm is 1.2 (± 0.4) m² (g C)⁻¹, similar to past values determined in water extracts of biomass burning particles (0.9 – 1.4 m² (g C)⁻¹) (Du et al., 2014; Fan et al., 2018; Park and Yu, 2016). At 300 nm, the average MAC of ABB is 1.5 (± 0.3) m² (g C)⁻¹, half the value of FBB, likely because of



photobleaching of brown carbon during aging (Hems and Abbatt, 2018; Hems et al., 2021; Laskin et al., 2015; Wong et al., 2017; Zhao et al., 2015). Win-Spr has an average $MAC_{DOC,300}$ ($1.9 (\pm 0.4) \text{ m}^2 (\text{g C})^{-1}$) that is three times higher than that of Sum-Fall ($0.65 (\pm 0.19) \text{ m}^2 (\text{g C})^{-1}$), though they have similar $PM_{2.5}$ concentrations. This indicates that winter wood combustion can significantly enhance light absorption by particles. Our winter MAC value is similar to the average value ($2.2 (\pm 0.7) \text{ m}^2 (\text{g C})^{-1}$) determined in previous water extracts of Davis winter particles (Kaur et al., 2019).

We also calculated the average MAC_{DOC} for each sample type in the wavelength range of 300-600 nm, as shown in Figure 2. Fresh wildfire samples have the highest MAC values across the wavelength range and the lowest absorption Ångström exponent (AAE, 300 – 450 nm), which is $7.3 (\pm 0.2)$. ABB shows slightly lower MAC values than Win-Spr. This might be explained by faster rates of aging and photobleaching during summer as well as higher amounts of less absorbing SOA. AAE values of ABB and Win-Spr are similar, $7.7 (\pm 0.3)$ and $7.9 (\pm 0.3)$, respectively, and are comparable to previously reported values of water-soluble organic carbon from biomass burning (Du et al., 2014; Hecobian et al., 2010; Lin et al., 2017). Sum-Fall has the lowest MAC but the highest AAE ($9.1 (\pm 0.5)$).




Figure 2. Average DOC-normalized mass absorption coefficients for Fresh wildfire samples (red), Winter & Spring samples (blue), Aged wildfire samples (orange), and Summer & Fall samples (green). Each shaded area represents ± 1 standard deviation.


300

An optical property frequently used to characterize surface water DOM is E_2/E_3 , which is the ratio of absorbance at 250 nm to that at 365 nm. In surface waters, this ratio is an indicator of the molecular weight of dissolved organic matter, with low E_2/E_3 representing high molecular-weight DOM (Ossola et al., 2021). E_2/E_3 in our PMEs ranges from 4.2 to 17 and is related to MAC values: as shown in Fig. 3, MAC decreases with increasing E_2/E_3 , i.e., absorbance decreases as DOM molecular weight decreases.

FBB has the lowest average E_2/E_3 ($5.8 (\pm 1.5)$) of our sample types, including ABB ($12.5 (\pm 2.3)$), which suggests that organic molecules in fresh BB are fragmented during aging. This is consistent with the observation that high-molecular weight compounds are less abundant in aged BBOA (Farley et al., 2022). Therefore, E_2/E_3 may be an easy and effective indicator to differentiate fresh



and aged samples. E_2/E_3 ratios for the Win-Spr samples are intermediate between the summer fresh and aged BB samples, again suggesting these biomass-burning influenced winter samples are less aged than ABB.



310

Figure 3. Mass absorption coefficients of dissolved organic carbon at 300 nm (circles) and 365 nm (triangles) as a function of E_2/E_3 for each sample type. Solid lines represent linear regressions.

Since the light absorption of methanol extracts of particles are usually greater than those of water extracts (Liu et al., 2013; Zhang et al., 2013), we also examined the absorbance of a FBB filter extracted with different solvents (water, methanol, and hexane). As shown in Fig. S10, the absorbance of the methanol (MeOH) extract is more than twice as high as the water extract, and five times higher than the hexane extract, indicating this FBB contains a high fraction of water-insoluble brown carbon. We also did a sequential extraction with this FBB sample and with a Win-Spr sample, with 1st, 2nd, and 3rd extraction solvents of water, methanol, and hexane, respectively. The UV-Vis spectra and PM mass extracted for each solvent extraction are shown in Fig. S11. For the Win-Spr and FBB samples, the PM mass recovered by the second extraction (in methanol) are only 20% and 56% of the mass by the first extraction (in water), respectively, but the MeOH extract absorbance at 365 nm is similar or even greater than the water extract. This is consistent with a previous study of sequential extraction with US western wildfire samples (Zeng et al., 2022), which found that water-insoluble brown carbon (e.g., polycyclic aromatic hydrocarbons) is highly light-absorbing, despite accounting for little of the PM mass. The high light absorption in methanol extracts suggests that the water-insoluble chromophores have high potential to produce photooxidants. However, since the oxidant probes we use were developed for aqueous, and not organic, solutions we did not study photooxidant generation in methanol or hexane extracts.

3.3 Photooxidant concentrations

3.3.1 Normalization by sample duration

While most of our PM samples were collected for 1 day, we also collected four samples for 7 days, which resulted in extracts that were more concentrated and that had higher oxidant concentrations. To properly compare these longer samples with the rest, we normalized photooxidant concentrations in the 7-day samples to what would be expected for a 24-h sample. For $^1O_2^*$ and $^3C^*$, the production rate is proportional to the brown carbon mass (Faust and Allen, 1992; Kaur et al., 2019) and so we normalized their concentrations by dividing by the duration of sampling (i.e., number of sampling days). The case for hydroxyl radical is more complicated, since past work has found that the $\bullet OH$ concentration can be independent of extract concentration (Arakaki et al.,


330



2013; Kaur et al., 2019), but unnormalized $\bullet\text{OH}$ concentrations in our 7-day samples are clearly higher than in the adjacent 24-h samples (Fig. S12). If we normalize $\bullet\text{OH}$ using the same method as for $^1\text{O}_2^*$ and $^3\text{C}^*$ (i.e., by the duration of sampling), the resulting $\bullet\text{OH}$ concentrations are lower than the adjacent 24 h samples (Fig. S12). To obtain more reasonable estimates for $[\bullet\text{OH}]$ in the 7-day samples, we fitted the plot of $\bullet\text{OH}$ concentration versus particle mass/water mass ratio for Win-Spr samples with a linear regression (Fig. S13), and then used the regression to estimate $\bullet\text{OH}$ concentrations in the 7-day samples using the time-normalized particle mass/water mass ratio values (i.e., measured particle mass/water ratio divided by 7).

340 3.3.2 Hydroxyl radical ($\bullet\text{OH}$)

As shown in Fig. 4a, normalized $\bullet\text{OH}$ concentrations have a range of $(0.2\text{-}3.2) \times 10^{-15}$ M. The values are similar to those in illuminated particle extracts from Davis and Colorado (Kaur et al., 2019; Leresche et al., 2021), but much higher than those in illuminated extracts of lab SOA and PM_{10} from Switzerland $((2.2\text{-}4.9) \times 10^{-17}$ M) that had low DOC (5 mg C L^{-1}) (Manfrin et al., 2019). Among our four sample types, fresh biomass burning samples have the highest average $[\bullet\text{OH}]$, $2.5 (\pm 0.3) \times 10^{-15}$ M, while aged BB particles have a similar average concentration that is statistically indistinguishable, $1.7 (\pm 1.4) \times 10^{-15}$ M. This is parallel to a previous finding that BBOA, compared to other types of organic aerosols, has the highest oxidative potential as measured by the DTT assay and this potential decreases with simulated atmospheric aging (Verma et al., 2015; Wong et al., 2019). Win-Spr has a similar average $[\bullet\text{OH}]$, $1.5 (\pm 0.3) \times 10^{-15}$ M, while Sum-Fall is the lowest at $0.4 (\pm 0.3) \times 10^{-15}$ M. Our winter values are roughly three to four times higher than average values in previous Davis winter particle extracts and fog waters ($0.51 (\pm 0.24) \times 10^{-15}$ M and $0.42 (\pm 0.07) \times 10^{-15}$ M, respectively) (Kaur and Anastasio, 2017; Kaur et al., 2019). While nitrate and nitrite can be important sources of $\bullet\text{OH}$ in atmospheric waters (Anastasio and McGregor, 2001; Kaur and Anastasio, 2017; Kaur et al., 2019; Leresche et al., 2021), these species account for less than 10% of $\bullet\text{OH}$ in most of our current samples (Table S3). In our kinetic experiments, in 6 of our 18 samples (5 winter samples and 1 wildfire sample) BA decayed faster at the beginning of irradiation but was slower at later times, with a rate difference up to a factor of 3.4 (Fig. S1). This indicates $[\bullet\text{OH}]$ in some samples is higher during the initial stage of irradiation, possibly because a portion of the compounds that produce $\bullet\text{OH}$ are labile and undergo rapid decomposition. A similar effect was seen in biomass burning aerosols from Fresno CA, where a burst of $\bullet\text{OH}$ was observed within the first few minutes of irradiation and was hypothesized to be due to the decomposition of peroxides through photo-Fenton reactions (Paulson et al., 2019).




360 **Figure 4.** Steady-state concentrations of (a) hydroxyl radical, (b) singlet molecular oxygen, and (c) oxidizing triplet excited states of organics determined by syringol (circles) and (phenylthio)acetic acid (crosses) in particle extracts. Concentrations are all normalized by sampling duration and to midday winter solstice sunlight in Davis to highlight seasonal differences in particle reactivity; the equivalent plots with concentrations calculated for the midday sunlight of each sample collection period is shown in Figure S14.

365 Figure. 5a shows $\bullet\text{OH}$ concentration as a function of dissolved organic carbon for the four sample types. For comparison, we also include data from Kaur et al. (2019), who measured photooxidant concentrations in Davis winter particle extracts. Though samples in Kaur et al. (2019) have similar values of DOC as our 24-h Win-Spr samples, their $[\bullet\text{OH}]$ is 5 times lower and independent of DOC. While $\bullet\text{OH}$ appears to increase with DOC (Fig. 5a), the data is noisy and the linear correlation is weak ($R^2 = 0.40$). A previous study on Minnesota surface waters observed a logarithmic relationship between $[\bullet\text{OH}]$ and absorbance coefficient at 440 nm (Chen et al., 2020), which in turn was correlated to DOC. They speculated this is because the dominant $\bullet\text{OH}$ sink changes from



370 bicarbonate/carbonate to DOC with increasing DOC levels, but bicarbonate/carbonate are negligible sinks in our extracts since they are acidic (pH 4.2). [$\bullet\text{OH}$] in FBB is independent of DOC, but the three ABB samples show $\bullet\text{OH}$ increasing with DOC. We also found that [$\bullet\text{OH}$] increases with DOC in a dilution series of summer wildfire PM and hypothesized that $\bullet\text{OH}$ production is a bimolecular reaction (primarily $\text{Fe(II)} + \text{HOOH}$) that increases as the square of PM mass concentration (Ma et al., 2023a). This might also explain our current ABB results.



375

Figure 5. Steady-state concentrations of (a) hydroxyl radical, (b) singlet molecular oxygen, and oxidizing triplet excited states of organic matter determined by (c) syringol and (d) (phenylthio)acetic acid as a function of dissolved organic matter for each sample type (solid circles). Previous measurements made in Davis winter particle extracts are in open circles (Kaur et al., 2019). Solid black lines are linear regressions between oxidant concentrations in this work and DOC. The blue dashed line in panel (d) is the linear regression of the Win-Spr samples. Error bars represent standard error propagated from linear regression and error in rate constants. Oxidant concentration values are not normalized by the sampling duration.

380

3.3.2 Singlet molecular oxygen ($^1\text{O}_2^*$)

Winter-solstice-sunlight normalized $^1\text{O}_2^*$ has a concentration range of $(0.7\text{--}32) \times 10^{-13}$ M (Fig. 4b) and correlates well with ambient $\text{PM}_{2.5}$ concentration (Fig. S15). These concentrations are similar to the wide range of previously reported values in particle extracts, $(0.6\text{--}22) \times 10^{-13}$ M (Bogler et al., 2022; Kaur et al., 2019; Leresche et al., 2021), but are roughly 100 times higher than concentrations in illuminated extracts of biogenic and anthropogenic SOA, $(0.8\text{--}45) \times 10^{-15}$ M (Manfrin et al., 2019). Our higher $^1\text{O}_2^*$ concentrations are only partially explained by our 1 – 40 times higher DOC concentrations; the remaining difference is likely due to greater light absorption by our samples. Our values are also similar to [$^1\text{O}_2^*$] in solutions of dissolved soot illuminated with simulated sunlight, $(0.6\text{--}65) \times 10^{-13}$ M (Li et al., 2019), even though their samples absorbed very little light. Among our samples, Fresh BB has the highest average [$^1\text{O}_2^*$], followed by ABB, with values of $29 (\pm 7) \times 10^{-13}$ M and $7.3 (\pm 0.4) \times 10^{-13}$ M, respectively. Leresche et al. (2021) found that [$^1\text{O}_2^*$] decreased by a factor of two in particle extracts after sunlight irradiation, which is consistent with our observation that aged particle extracts have lower [$^1\text{O}_2^*$]. Win-Spr and Sum-Fall samples have average [$^1\text{O}_2^*$] values of

390



3.8 (\pm 1.6) $\times 10^{-13}$ M and 1.1 (\pm 0.6) $\times 10^{-13}$ M, respectively. The higher Win-Spr concentrations are probably because of the
395 influence of biomass burning.

As shown in Fig. 5b, $^1\text{O}_2^*$ concentrations linearly increase with DOC ($R^2 = 0.93$), consistent with our understanding that organic matter is the primary source of $^1\text{O}_2^*$ (Bogler et al., 2022; Kaur and Anastasio, 2017; Kaur et al., 2019; Ossola et al., 2021). Moreover, all four types of samples share the same slope, suggesting the relationship between [$^1\text{O}_2^*$] and DOC is independent of
400 particle type or chemical composition, which is somewhat surprising given the large differences in DOC-normalized light absorption for the different samples types (Fig. 2). When plotting [$^1\text{O}_2^*$] as a function of absorbance at 300 and 365 nm (Figs. S16b and S17b, respectively), we do observe differences among sample types. In these plots, Win-Spr samples present a steeper slope (as do samples from Kaur et al. (2019)) compared to wildfire samples, consistent with our previous work (Ma et al., 2023a). The $^1\text{O}_2^*$ concentrations in previous Davis winter particle extracts (Kaur et al., 2019) also follow the linear regression of this work.
405 While this suggests DOC is a robust descriptor for $^1\text{O}_2^*$ concentrations, most of our particle samples were influenced by biomass burning. Other particle types - such as anthropogenic SOA, biogenic SOA, and emissions from fossil fuel combustion appear to have different relationships between $^1\text{O}_2^*$ and DOC, as suggested by results from Manfrin et al. (2019), Jiang et al. (2023), and Bogler et al. (2022).

3.3.3 Oxidizing triplet excited states of brown carbon ($^3\text{C}^*$)

410 We used two probes – syringol (SYR) and (phenylthio)acetic acid (PTA) – to quantify oxidizing triplet excited states. SYR reacts rapidly with both strongly and weakly oxidizing triplets, while PTA is only reactive with strongly oxidizing triplets (Ma et al., 2023b). However, syringol has a disadvantage that its decay by $^3\text{C}^*$ can be inhibited by dissolved organic matter, while PTA is largely resistant to inhibition (Ma et al., 2023b; Maizel and Remucal, 2017; McCabe and Arnold, 2017; Wenk et al., 2011). As shown in Fig. 4c, winter-solstice-normalized (and inhibition-corrected) $^3\text{C}^*$ concentrations have a range of (0.13 – 6.9) $\times 10^{-13}$ M
415 as determined by SYR and (0.03 – 1.9) $\times 10^{-13}$ M by PTA. The $^3\text{C}^*$ concentration follows $\text{PM}_{2.5}$ concentration well, with low values during non-wildfire periods and very high values during wildfire-influenced periods (Fig. S15). For nearly all samples, [$^3\text{C}^*$]_{SYR} is higher than [$^3\text{C}^*$]_{PTA}. As seen for $^1\text{O}_2^*$, FBB has the highest average [$^3\text{C}^*$], 4.8 (\pm 1.4) $\times 10^{-13}$ M from SYR and 1.8 (\pm 1.6) $\times 10^{-13}$ M from PTA, due to the high organic amounts in these samples. Relative to the FBB average, the FBB, ABB, Win-Spr, and Sum-Fall samples have triplet concentration ratios of 1 : 0.32 : 0.12 : 0.04 as determined by SYR and 1 : 0.32 : 0.21 : 0.03 as determined
420 by PTA. These ratios are similar to the ratio of average DOC concentrations, which is 1 : 0.45 : 0.15 : 0.08, indicating DOC is the main driver of $^3\text{C}^*$ concentration differences among sample types. This relationship is complicated at high DOC where dissolved organics can be the dominant triplet sink (up to roughly 60% of the total sink), larger than the contribution from dissolved oxygen.

Figure 5c shows the correlation between [$^3\text{C}^*$]_{SYR} and DOC for our samples, along with data from Kaur et al. (2019). [$^3\text{C}^*$]_{SYR}
425 linearly increases with DOC ($R^2 = 0.83$) independent of sample type, likely because SYR reacts rapidly with a wide range of oxidizing triplets (Kaur and Anastasio, 2018). However, Figs. S16c and S17c show some differences between sample types in the relationship between [$^3\text{C}^*$]_{SYR} and absorbance at 300 or 365 nm, with Win-Spr samples having a steeper slope. However, the trend of FBB samples is hard to discern, in part because of the limited number of samples available (only four). As shown in Fig. 5d, [$^3\text{C}^*$]_{PTA} also linearly increases with DOC, though the correlation is not as good as those for [$^1\text{O}_2^*$] or [$^3\text{C}^*$]_{SYR}. Win-Spr samples
430 present a slightly higher slope than wildfire samples (FBB and ABB); oddly, [$^3\text{C}^*$]_{PTA} is nearly independent of DOC within either biomass burning group. The steeper slope of [$^3\text{C}^*$]_{PTA} with DOC for the Win-Spr samples suggests these samples contain a higher



fraction of highly oxidizing $^3\text{C}^*$ than the wildfire samples. This difference in slopes is particularly noticeable in Figs. S16d and S17d, where $[\text{}^3\text{C}^*]_{\text{PTA}}$ is plotted against absorbance at 300 or 365 nm.

435 Since PTA only captures $^3\text{C}^*$ that have high reduction potentials, while SYR reacts rapidly with both strongly and weakly oxidizing triplets, the ratio $[\text{}^3\text{C}^*]_{\text{PTA}}/[\text{}^3\text{C}^*]_{\text{SYR}}$ provides an estimate of the fraction of oxidizing $^3\text{C}^*$ that are strong oxidants. As shown in Fig. 6, the ratio $[\text{}^3\text{C}^*]_{\text{PTA}}/[\text{}^3\text{C}^*]_{\text{SYR}}$ ranges from 0.27 (± 0.10) to 1.7 (± 0.7) with an average value of 0.58 (± 0.38), indicating roughly 60% of oxidizing triplets are strong oxidants. The Win-Spr samples have an average ratio of 0.86 (± 0.43), significantly higher than the rest of the samples (0.37 ± 0.07), indicating that they produce a higher fraction of strongly oxidizing $^3\text{C}^*$. Precursors for
440 more oxidizing triplets include quinones, aromatic ketones and aromatic aldehydes, while weakly oxidizing triplet precursors include polycyclic aromatic compounds (McNeill and Canonica, 2016).




Figure 6. The ratio of oxidizing triplet excited state concentrations determined by PTA to those determined by SYR as a function of DOC for each sample type.

445 We can also gain some insight into extract compositions from the inhibition factors (*IF*) (Section S1) for SYR and PTA in each sample. An *IF* of 1 represents no inhibition of probe decay by the sample, while an *IF* of 0 indicates that the triplet-mediated decay of probe is completely reversed by DOM in the sample (Cannonica and Laubscher, 2008; Ma et al., 2023b). Among our samples, *IF* for SYR ($IF_{\text{SYR,corr}}$) ranges from 1.2 to 0.21, with an average value of 0.64 (± 0.29) (Table S6 and Fig. S18). This indicates that SYR decay by $^3\text{C}^*$ in PME can be heavily inhibited, suggesting that our PMEs contain abundant antioxidants such as phenolic
450 or aniline moieties (Wenk and Canonica, 2012; Wenk et al., 2011). As shown in Fig. S18b, $IF_{\text{SYR,corr}}$ generally decreases with increasing DOC, consistent with previous surface water studies (Cannonica and Laubscher, 2008; McCabe and Arnold, 2017). We fit $IF_{\text{SYR,corr}}^{-1}$ versus DOC using a linear regression with all samples (Ma et al., 2023b; Wenk et al., 2011) as shown in Fig S18b. The fitted slope is 0.015 L mg C⁻¹; the inverse of this slope, 67 (± 13) mg C L⁻¹, represents the DOC concentration that causes $IF_{\text{SYR,corr}}$ to equal 0.5. All of the sample groups essentially fit on the same line. The *IF* for PTA ($IF_{\text{PTA,corr}}$) ranges from 1.5 to 0.6,
455 with an average value of 1.1 (± 0.2), demonstrating its better resistance to inhibition (Fig. S18c). We also measured the inhibition factor of furfuryl alcohol (IF_{FFA}) as the indicator of the ability of DOM in PME to quench $^3\text{C}^*$ (Fig. S18a). IF_{FFA} decreases with increasing DOC, ranging from 1.4 (i.e., no quenching of triplets by PME DOM) to 0.5 (i.e., DOM is reducing the triplet concentration to 50 % of its non-quenched value). From the linear fit between IF_{FFA}^{-1} and DOC, we obtain a second-order rate



constant of DOM quenching ${}^3\text{DMB}^*$ (Ma et al., 2023b; Wenk et al., 2011, 2013) of $2.7 (\pm 0.7) \times 10^7 \text{ L (mol-C)}^{-1} \text{ s}^{-1}$. This value is
460 somewhat lower than rate constants of DOM quenching oxidizing ${}^3\text{C}^*$ in Davis particle extracts ($(5.7 - 12) \times 10^7 \text{ L (mol C)}^{-1} \text{ s}^{-1}$)
(Ma et al., 2023a) but in the range of values for DOM quenching ${}^3\text{C}^*$ in surface waters. $(1.3-7.9) \times 10^7 \text{ L (mol C)}^{-1} \text{ s}^{-1}$ (Wenk et al.,
2013).

3.3.5 Normalization by photon flux

Photooxidant concentrations in Figures 4 and 5 are all normalized to the same actinic flux condition (i.e., solar noon on the winter
465 solstice in Davis CA, $j_{2\text{NB}} = 0.007 \text{ s}^{-1}$) to highlight seasonal differences in particle reactivity. However, photon fluxes vary
throughout the year, which will affect the rate of photooxidant formation and accompanying concentration. To account for this
effect, we calculated midday $j_{2\text{NB}}$ values as a function of date during our sampling campaign, as shown in Fig. S19 and described
in Section S3. The estimated $j_{2\text{NB}}$ value at midday of the summer solstice is 0.013 s^{-1} , which is nearly twice the value during winter.
Next, we estimated midday $j_{2\text{NB}}$ values for each sampling day and normalized photooxidant concentrations to the corresponding
470 sunlight condition. Figure S14 shows the equivalent plot of Figure 3 after photon flux normalization, which increased oxidant
concentrations by factors ranging from 1.0 to 1.9. The average normalization factors for FBB and Sum-Fall samples are 1.7, while
ABB and Win-Spr have average factors of 1.5 and 1.2, respectively. These $j_{2\text{NB}}$ values do not account for optical confinement of
sunlight within particles; recent work suggests that this will enhance in-particle actinic fluxes by approximately a factor of two
(Corral Arroyo et al., 2022), which would cause a proportional increase in oxidant concentrations. At this point we do not have
475 enough information to understand how seasonal variations in temperature might affect oxidant concentrations, so we have not
attempted to factor this into our analysis.

3.4 Quantum yields for photooxidants

3.4.1 Hydroxyl radical


480 To investigate how sample type affects the efficiency of photooxidant formation, we determined apparent quantum yields of
photooxidant formation (Φ_{Ox}), i.e., the fraction of absorbed photons that result in formation of a particular photooxidant:

$$\Phi_{\text{Ox}} = \frac{P_{\text{Ox}}}{R_{\text{abs}}} \quad (7)$$

where P_{Ox} is the oxidant production rate and R_{abs} is the rate of sunlight absorption by the sample between 300 and 450 nm (Kaur
et al., 2019). We calculate the production rate of $\bullet\text{OH}$, P_{OH} , by assuming it is equal to the $\bullet\text{OH}$ consumption rate since hydroxyl
485 radical (and the other photooxidants) are at steady state. Thus, P_{OH} is equal to the product of $[\bullet\text{OH}]$ and the first-order rate constant
of $\bullet\text{OH}$ loss by natural sinks (k'_{OH}). To estimate k'_{OH} , we assume that organic matter is the dominant sink for $\bullet\text{OH}$ (Kaur et al.,
2019) and that k'_{OH} is the product of DOC concentration and the second-order rate constant of DOC with $\bullet\text{OH}$ ($k_{\text{DOC}+\text{OH}}$). For
 $k_{\text{DOC}+\text{OH}}$, we used the average value measured in Davis winter and summer wildfire particle extracts (Ma et al., 2023a), which is
2.7 (± 0.4) $\times 10^8 \text{ L (mol-C)}^{-1} \text{ s}^{-1}$. This value is slightly lower than that determined by Arakaki et al. (2013) for a broad range of
490 atmospheric waters (3.8 (± 1.9) $\times 10^8 \text{ L (mol-C)}^{-1} \text{ s}^{-1}$) and the one from Leresche et al. (2021) for Colorado PM extracts (4.9 (± 2.3)
 $\times 10^8 \text{ L (mol-C)}^{-1} \text{ s}^{-1}$), but none of these are statistically different. In our samples, the resulting calculated k'_{OH} is in the range (0.11
– 4.3) $\times 10^6 \text{ s}^{-1}$ (Table S3), yielding P_{OH} in the range of (0.04-14) $\times 10^{-9} \text{ M s}^{-1}$, similar to past measured and modeled values for
fog/cloud waters and particle extracts (Arakaki et al., 2013; Leresche et al., 2021; Tilgner and Herrmann, 2018).



495 Our calculated apparent quantum yields of $\bullet\text{OH}$ are shown in Fig. 7a, along with past Davis winter PME samples from Kaur et al. (2019). Φ_{OH} ranges from 0.01 % to 0.10 % in our samples, which are generally higher than values from Kaur et al. (2019) and from PM_{10} and lab SOA water extracts (Manfrin et al., 2019). As expected, Φ_{OH} appears independent of DOC. Average $\bullet\text{OH}$ quantum yields for Win-Spr, Sum-Fall, FBB, and ABB are $0.044 (\pm 0.022)$ %, $0.028 (\pm 0.010)$ %, $0.021 (\pm 0.005)$ %, and $0.049 (\pm 0.050)$ %, respectively. These averages are not statistically different ($p > 0.05$).



500

Figure 7. Apparent quantum yields of (a) hydroxyl radical, (b) singlet molecular oxygen, and oxidizing triplets determined by (c) syringol and (d) (phenylthio)acetic acid as a function of dissolved organic matter for each sample type (solid circles). Previous measurements made in Davis winter particle extracts are in open circles (Kaur et al., 2019).

3.4.2 Singlet molecular oxygen

505 To calculate the apparent quantum yields of $^1\text{O}_2^*$ ($\Phi_{1\text{O}_2^*}$), we assume that H_2O is the dominant sink for $^1\text{O}_2^*$ in our PM extracts. This is a reasonable assumption since the first-order rate constants for $^1\text{O}_2^*$ loss via DOC are $(0.04 - 2) \times 10^3 \text{ s}^{-1}$ in our samples (based on an estimated $^1\text{O}_2^* + \text{DOC}$ rate constant of $1 \times 10^5 \text{ L (mol-C)}^{-1} \text{ s}^{-1}$ (Ma et al., 2023a)), while the rate constant for $^1\text{O}_2^*$ loss by water is $2.2 \times 10^5 \text{ s}^{-1}$ (Bilski et al., 1997). Therefore, we calculated the production rate of $^1\text{O}_2^*$ ($P_{1\text{O}_2^*}$) by multiplying the rate of $^1\text{O}_2^*$ loss by water ($k'_{\text{H}_2\text{O}}$) by $[^1\text{O}_2^*]$. As shown in Fig. 7b, $\Phi_{1\text{O}_2^*}$ ranges from 1.7% to 8.4%, comparable to values from

510 Kaur et al. (2019), which are shown as open circles in the figure, as well as from SOA and ambient particle extracts in other studies (0.1 – 4.5 %) (Bogler et al., 2022; Leresche et al., 2021; Manfrin et al., 2019). But our $\Phi_{1\text{O}_2^*}$ values are significantly lower than those in dissolved soot extracts (33%) (Li et al., 2019) under 377 nm irradiation; we do not expect significant black carbon in our extracts since they were filtered. Sum-Fall has the highest average $\Phi_{1\text{O}_2^*}$, $7.9 (\pm 0.4)$ %, which is significantly different from the others, while Win-Spr and ABB have similar average values, $4.0 (\pm 1.1)$ and $3.9 (\pm 1.0)$, respectively, and FBB shows the lowest

515 average $\Phi_{1\text{O}_2^*}$ of $2.2 (\pm 0.5)$ %. The higher quantum yield for aged biomass burning PM compared to fresh BB PM is broadly consistent with the enhancement in $\Phi_{1\text{O}_2^*}$ resulting from ozonation of surface water DOM (Leresche et al., 2019). The difference



among sample types is more pronounced when $\Phi_{1O_2^*}$ is plotted as a function of MAC. As shown in Figs. S20b and S21b, $\Phi_{1O_2^*}$ decreases with absorbance at 300 or 365 nm, indicating that less light-absorbing brown carbon (e.g., Sum-Fall) more efficiently produces $^1O_2^*$ compared to high-MAC samples (e.g., FBB). In surface waters, $\Phi_{1O_2^*}$ is positively correlated with E_2/E_3 , i.e., the $^1O_2^*$ quantum yield increases for DOM with lower average molecular weight molecules (Berg et al., 2019; Ossola et al., 2021). We find a similar linear relationship in our samples, with an R^2 of 0.54 (Fig. S22). The fresh BB extract has low E_2/E_3 (and low $\Phi_{1O_2^*}$), suggesting that it contains more high-molecular-weight compounds that absorb significant amounts of light but inefficiently produce $^1O_2^*$. It has been suggested that DOM with a high lignin content (as expected for BB PM) can have a high degree of charge transfer interactions, which results in low $\Phi_{1O_2^*}$ (Ossola et al., 2021). Despite the relatively inefficient production of singlet oxygen by the fresh BB extracts, these samples have some of the highest $^1O_2^*$ concentrations (Fig. 4), a result of their very strong light absorption (Fig. 2).

3.4.3 Oxidizing triplet excited states

To calculate the production rate of $^3C^*$, we first need to estimate the $^3C^*$ sink, which is dominated by dissolved oxygen at low DOC but by organic matter as DOC increases. We estimated average second-order rate constants for DOC reacting with and physically quenching $^3C^*$ ($k_{rxn+Q,3C^*}$) in our samples by fitting $[^3C^*]$ as a function of DOC with a hyperbolic regression (Fig. S23). Values of $k_{rxn+Q,3C^*}$, calculated from one of the regression fitting parameters (Kaur et al., 2019), are $7.2 (\pm 2.2) \times 10^7 \text{ L (mol-C)}^{-1} \text{ s}^{-1}$ for $^3C^*$ determined by SYR and $7.4 (\pm 2.5) \times 10^7 \text{ L (mol-C)}^{-1} \text{ s}^{-1}$ for $^3C^*$ by PTA. Since the production rate of $^3C^*$ (P_{3C^*}) is equal to its loss rate, we calculate the former with:

$$P_{3C^*} = (k_{rxn+Q,3C^*}[DOC] + k_{3C^*+O_2}[O_2]) \times [^3C^*] \quad (8)$$

where $k_{3C^*+O_2}$ is the second-order rate constant of dissolved oxygen reacting with $^3C^*$ ($2.8 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) (Kaur et al., 2019) and $[O_2]$ is the dissolved oxygen concentration, 280 μM at 20 °C for an air-saturated solution (U.S. Geological Survey, 2020). The apparent quantum yield of $^3C^*$ is then calculated using P_{3C^*} divided by the rate of light absorption (Eq. 7).

Figures 7c and 7d show quantum yields of $^3C^*$ determined by SYR ($\Phi_{3C^*,SYR}$) and PTA ($\Phi_{3C^*,PTA}$). $\Phi_{3C^*,SYR}$ has a range of (0.9-8.8) % and an average value of $3.5 (\pm 1.8) \%$. Our values are similar to Φ_{3C^*} in past Davis winter PM extracts (as shown by the open circles in the figures), as well as fog waters and surface waters, which are in the range (0.3-14) % (Kaur and Anastasio, 2018; McCabe and Arnold, 2018). We do not observe significant differences in $\Phi_{3C^*,SYR}$ among sample types (Fig. S24), consistent with the similarities among sample types in the relationship of $[^3C^*]_{SYR}$ versus DOC (Fig. 4). $\Phi_{3C^*,PTA}$ has a range of (0.6-3.4) %, with an average value of $1.7 (\pm 0.7) \%$, half of the average $\Phi_{3C^*,SYR}$. Win-Spr has the highest average $\Phi_{3C^*,PTA}$, $2.1 (\pm 0.7) \%$, while FBB has the lowest, $0.96 (\pm 0.39) \%$, but they are not statistically different. Though $^3C^*$ is the precursor of $^1O_2^*$, Φ_{3C^*} does not correlate well with MAC, unlike $\Phi_{1O_2^*}$ (Figs. S20 and S21), probably because we are measuring only the oxidizing portion of the triplet pool. In surface waters, Φ_{3C^*} often increases with E_2/E_3 , similar to $\Phi_{1O_2^*}$ (Berg et al., 2019; Maizel and Remucal, 2017; McCabe and Arnold, 2017), but we do not see this triplet behavior in our samples (Fig. S25) even though we do for $^1O_2^*$ (Fig. S22).

We next use our quantum yields to estimate the fraction of the total triplet pool that can oxidize SYR or PTA. Since almost all triplets can transfer energy to dissolved oxygen to make $^1O_2^*$, we estimate the quantum yield of total $^3C^*$ as $\Phi_{1O_2^*}/f_{\Delta}$, where f_{Δ} is the fraction of $^3C^*$ interaction with dissolved oxygen that forms $^1O_2^*$, estimated as 0.53 (Kaur and Anastasio, 2018; McNeill and Canonica, 2016). Therefore, the fraction of triplets that are oxidizing can be calculated as $\Phi_{3C^*}/(\Phi_{1O_2^*}/f_{\Delta})$, with values shown in Fig. S26. For $^3C^*$ determined by SYR, the fraction ranges from 0.14 to 0.81, with an average of $0.47 (\pm 0.20)$ and no statistical difference among the four sample types. This average value is similar to those determined in fog waters (0.55 ± 0.44) as well as in



previous Davis winter particle extracts (0.31 ± 0.11) (Kaur and Anastasio, 2018; Kaur et al., 2019), indicating that roughly half of the triplets in Davis PM and fog samples are oxidizing. For strongly oxidizing triplets determined by PTA, the fraction ranges from 0.07 to 0.45, with an average of 0.24 (± 0.09); this is half the SYR value, suggesting that approximately half of oxidizing $^3\text{C}^*$ possesses a high reduction potential, consistent with the results of Fig. 6. For $^3\text{C}^*$ determined by PTA, Sum-Fall has a statistically lower average value, 0.11 (± 0.05), compared to Win-Spr (0.29 ± 0.09), FBB (0.22 ± 0.04), and ABB (0.23 ± 0.06). This is reasonable because Sum-Fall samples were not significantly influenced by biomass burning, leading to a lower aromatic content and more weakly oxidizing triplets (McNeill and Canonica, 2016).

3.4.4. Quantum yields in aerosol liquid water

We calculated the quantum yields above for the relatively dilute conditions of our particle extracts, but these results are not necessarily applicable to the more concentrated conditions of aerosol liquid water. This is because the formation rate of each oxidant (P_{Ox}) is not necessarily proportional to the concentration factor of the sample, while the light absorption should be proportional; based on Eq. 7, if these factors do not vary in the same way as samples get more concentrated, the quantum yield will vary with concentration. As described by Ma et al. (2023a), as we move from dilute extracts to concentrated particle water $P_{^3\text{C}^*}$ appears to increase linearly with concentration factor, $P_{^1\text{O}_2^*}$ does not, and P_{OH} only does sometimes. This suggests that triplet quantum yields in ALW will be similar to those determined in PME, but that yields for singlet oxygen and hydroxyl radical can be lower in ALW compared to in PME. In each case, care needs to be taken when applying the extract quantum yields from above to more concentrated conditions.

3.5 Extrapolation of photooxidant concentrations to aerosol liquid water (ALW) conditions

Particle mass/water mass ratios in our PM extracts range from 10^{-5} to 10^{-3} $\mu\text{g PM}/\mu\text{g H}_2\text{O}$ (Table S1), which are typical for dilute hydrometeors like cloud and fog drops (Hess et al., 1998; Nguyen et al., 2016; Parworth et al., 2017). While the results in dilute extracts are interesting and applicable to cloud and fog chemistry, our goal is to understand photooxidant concentrations for each sample type in aerosol liquid water, which is orders of magnitude more concentrated (typically near $1 \mu\text{g PM}/\mu\text{g H}_2\text{O}$). Due to the very limited water content of particles, we cannot study this condition directly using our current probe techniques. Instead, our approach has been to quantify photooxidant kinetics (i.e., formation rates and loss rate constants) in a single PM sample as a function of particle dilution and then extrapolate to ALW conditions (Kaur et al., 2019; Ma et al., 2023a). We do this with our current samples by applying parameters obtained from our recent dilution study of a winter (WIN) and a summer (SUM) $\text{PM}_{2.5}$ sample (Ma et al., 2023a). Details about the extrapolations and accompanying parameters are provided in Section S4 and Table S10. Moreover, we take the influence of actinic flux on sample types into consideration by using the average midday $j_{2\text{NB}}$ value for each sample type to normalize photooxidant concentrations to that sunlight condition.


We calculate $[\bullet\text{OH}]$ in ALW using the average P_{OH} and k'_{OH} values that were determined from the Davis winter and summer particle extracts in our previous study (Ma et al., 2023a). We do not consider the effect of sample type because we do not observe significant differences in the relationship of $[\bullet\text{OH}]$ versus DOC among our four sample types (Fig. 5a). As shown in Fig. S27, the predicted $\bullet\text{OH}$ concentration is relatively constant across drop to particle conditions, with a range of $(6 - 9) \times 10^{-15}$ M. The predicted $[\bullet\text{OH}]$ in dilute condition is higher than our measured values because we include $\bullet\text{OH}$ from the gas phase in our calculation (Kaur et al., 2019). As shown in Figure 8, $[\bullet\text{OH}]$ at $1 \mu\text{g PM}/\mu\text{g H}_2\text{O}$ has a range of $(8.8 - 13) \times 10^{-15}$ M, of which the difference among



sample types is driven by the seasonal variation in actinic flux. Our $\bullet\text{OH}$ concentrations are around 10 times higher than the previous ALW value predicted by Kaur et al. (2019).

595

We next consider singlet oxygen. As shown in Fig. S29, $[^1\text{O}_2^*]$ for each sample type increases with particle mass/water mass ratio under dilute conditions, peaks near $0.01 - 0.1 \mu\text{g PM}/\mu\text{g H}_2\text{O}$, and then decreases under more concentrated conditions. At $1 \mu\text{g PM}/\mu\text{g H}_2\text{O}$, Win-Spr has the highest $[^1\text{O}_2^*]$ ($8 \times 10^{-12} \text{ M}$), followed by Sum-Fall ($3 \times 10^{-12} \text{ M}$), FBB ($2 \times 10^{-12} \text{ M}$), and ABB ($1 \times 10^{-12} \text{ M}$) (Fig. 8). Win-Spr is characterized by its high $^1\text{O}_2^*$ quantum yield, second highest light absorption, and low rate of DOC quenching for both $^3\text{C}^*$ and $^1\text{O}_2^*$. In contrast, FBB and ABB have more brown carbon (and therefore greater sources of $^1\text{O}_2^*$) but high DOC, which leads to greater sinks for triplets and singlet oxygen. Moreover, DOC in FBB and ABB quenches $^3\text{C}^*$ more efficiently than that in Win-Spr (i.e., the BB samples have higher values of $k_{3\text{C}^*+\text{DOC}}$). Therefore, their $[^1\text{O}_2^*]$ in ALW are similar to, or even lower than, $[^1\text{O}_2^*]$ measured in FBB and ABB extracts, while the ALW singlet oxygen concentrations for Win-Spr and Sum-Fall are nearly 20 times higher than their corresponding averages in extracts. Our estimated $[^1\text{O}_2^*]$ in ALW is 20 – 200 times lower than the value derived by Kaur et al. (2019), $1.6 \times 10^{-10} \text{ M}$, for Davis winter particle water. This is primarily because we account for DOC suppressing $^3\text{C}^*$ concentrations, and therefore lowering the rate of $^1\text{O}_2^*$ production at high DOC values; this was not done in the previous work.



610 **Figure 8.** Predicted photooxidant concentrations for each sample type under aerosol liquid water conditions ($1 \mu\text{g PM}/\mu\text{g H}_2\text{O}$), normalized to the average midday actinic flux for each sample type. Previous extrapolations made from Davis winter particle extracts are in open bars, where photooxidant concentrations are normalized to Davis winter solstice sunlight and $^3\text{C}^*$ is the lower-bound estimate (Kaur et al., 2019). Photooxidant concentrations all normalized to Davis winter solstice sunlight are in Figure S31.


Our final ALW predictions are for oxidizing triplets. $[^3\text{C}^*]$ for all sample types increases with particle mass concentration under dilute conditions, but then reaches a plateau as solutions become more concentrated and DOC becomes the dominant sink for triplets (Fig. S30). As shown in Fig. 8, $[^3\text{C}^*]_{\text{SYR}}$ and $[^3\text{C}^*]_{\text{PTA}}$ at $1 \mu\text{g PM}/\mu\text{g H}_2\text{O}$ have a range of $(0.4 - 13) \times 10^{-12} \text{ M}$ and $(0.1 - 10) \times 10^{-12} \text{ M}$, respectively, with Win-Spr and Sum-Fall having the maximum and minimum values, respectively. Sum-Fall samples might contain a lower fraction of carbonyl or ketone compounds compared to other sample types, leading to lower production of oxidizing $^3\text{C}^*$ (McNeill and Canonica, 2016). Compared to our average measured $^3\text{C}^*$ concentration in the PM extracts, $[^3\text{C}^*]$ in ALW for Win-Spr and Sum-Fall samples increases by a factor of approximately 20, while ALW concentrations for FBB and ABB

620



are only around 2 times higher than their extract values. Our predicted $[^3\text{C}^*]_{\text{SYR}}$ is 2 – 5 times higher than the lower-bound estimate of Kaur et al. (2019) (Fig. 8).

625 Finally, to understand how photooxidants affect the fate of organic compounds in ALW, we revisit the Kaur et al. (2019) estimates for the lifetimes and fates of five model organic compounds: (1) syringol, (2) methyl jasmonate, (3) tyrosine, (4) 1,2,4-butanetriol, and (5) 3-hydroxy-2,5-bis(hydroxymethyl)furan. To estimate the fate of each compound, we assume equilibrium gas-aqueous partitioning in an aerosol with an ALW of $20 \mu\text{g m}^{-3}$ and consider reactions with two gas-phase oxidants ($\bullet\text{OH}$, O_3) and four aqueous-phase oxidants ($\bullet\text{OH}$, O_3 , $^3\text{C}^*$, $^1\text{O}_2^*$). In our calculations, we employed rate constants and Henry's law constants (K_{H}) from Kaur et al. (2019) and used our predicted ALW photooxidant concentrations in Win-Spr ($[\bullet\text{OH}] = 7 \times 10^{-15} \text{ M}$, $[^1\text{O}_2^*] = 7 \times$
630 10^{-12} M , $[^3\text{C}^*]_{\text{SYR}} = 1 \times 10^{-12} \text{ M}$, normalized to Davis winter solstice). More details about the calculations are in Kaur et al. (2019). As shown in Figure 9, compounds (1) and (2), which have low K_{H} values, partition negligibly to the aqueous phase and so gas-phase reactions dominate their fates, with overall lifetimes of 2-3 h; these results are the same for both the aqueous oxidant concentrations of Kaur et al. (2019) and those determined in this work (i.e., Figure 8). For compounds (3), (4), and (5), which have high K_{H} values, 30 – 100% of the species are present in the aqueous phase of the aerosol. With photooxidant concentrations predicted by Kaur et al. (2019), organic lifetimes range from 0.04 to 20 h and $^1\text{O}_2^*$ is the major sink. However, in this work we predict higher $\bullet\text{OH}$ and $^3\text{C}^*$ concentrations but significantly lower $^1\text{O}_2^*$ in ALW (Figure 8). The lower $^1\text{O}_2^*$ leads to lifetimes of compounds (3) and (5) increasing by factors of 6 and 17, respectively. $^3\text{C}^*$ becomes the dominant oxidant for the phenolic amino acid, compound (3), but singlet oxygen is still the dominant sink for the substituted furan, compound (5). With the new oxidant concentrations, the lifetime of the aliphatic alcohol, compound (4), decreases by a factor of almost 3 due to the enhanced $\bullet\text{OH}$
640 concentration and singlet oxygen is much less important. Overall, results with the new oxidant concentrations show some significant shifts in the lifetimes of the three highly soluble organics as well as in the contributions of individual oxidants. But our new results still indicate that $^3\text{C}^*$ and $^1\text{O}_2^*$ dominate the particle processing for highly soluble organic compounds with which they react quickly while $\bullet\text{OH}$ dominates for aqueous organics that react slowly with the other two oxidants. Based on our Win-Spr oxidant concentrations (Figure 8), for an organic compound that has an $\bullet\text{OH}$ rate constant of $1 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$, singlet oxygen will be the dominant oxidant if its rate constant with the organic is larger than roughly $1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$, while oxidizing triplets will dominate if their rate constant is larger than approximately $1 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$,



650 **Figure 9.** Fates of five model organic compounds – (1) syringol, (2) methyl jasmonate, (3) tyrosine, (4) 1,2,4-butanetriol, and (5) 3-hydroxy-2,5-bis(hydroxymethyl)furan in an aerosol ($20 \mu\text{g dry PM}/\text{m}^3\text{-air}$) containing equal amounts of PM and liquid water (i.e., $1 \mu\text{g PM}/\mu\text{g H}_2\text{O}$). Results to the left of the dashed lines are calculated using estimated photooxidant concentrations from Kaur et al. (2019), while those to the right are calculated using oxidant concentrations for Win-Spr from this work. Panel (a) shows the overall lifetime (blue columns, left y-axis) and the fraction in the aqueous phase (red circles) for each organic. Panel (b) shows the fraction of organic lost due to each oxidant.

655 **4. Conclusions, Implications, and Uncertainties**

In this work, we measured concentrations of three photooxidants – hydroxyl radical, singlet molecular oxygen, and oxidizing triplet excited states of brown carbon – in particle extracts. Our extracts have particle mass/liquid water mass ratios in the range of $(0.7\text{--}9.1) \times 10^{-4} \mu\text{g PM}/\mu\text{g H}_2\text{O}$, which are close to fog/cloud water conditions but much more dilute than aerosol liquid water. We categorized samples into four types based on sampling dates and chemical characterization: Winter & Spring (Win-Spr), Summer & Fall (Sum-Fall) without wildfire influence, fresh biomass burning (FBB), and aged biomass burning (ABB). FBB contains the highest amounts of BrC, leading to the highest average mass absorption coefficients normalized by dissolved organic carbon, e.g., $3.3 (\pm 0.4) \text{ m}^2 (\text{g C})^{-1}$ at 300 nm. Win-Spr and ABB have similar MACs at this wavelength ($1.9 (\pm 0.4) \text{ m}^2 (\text{g C})^{-1}$ and $1.5 (\pm 0.3) \text{ m}^2 (\text{g C})^{-1}$, respectively), while Sum-Fall has the lowest MAC_{DOC} ($0.65 (\pm 0.19) \text{ m}^2 (\text{g C})^{-1}$).



665 Photooxidant concentrations in the particle extracts are in the range $(0.2-4.7) \times 10^{-15}$ M for $\bullet\text{OH}$, $(0.07-4.5) \times 10^{-12}$ M for $^1\text{O}_2^*$, and $(0.03 - 7.9) \times 10^{-13}$ M for $^3\text{C}^*$, respectively. All oxidant concentrations generally increase with the concentration of dissolved organic carbon (DOC), which ranged from 5 to 192 mg C L⁻¹. $^1\text{O}_2^*$ concentrations exhibit good linearity with DOC with all sample types falling roughly on the same line. Fresh BB extracts have the highest [$^1\text{O}_2^*$] but the lowest average quantum yield ($\Phi_{1\text{O}_2^*}$), while Sum-Fall samples are the opposite. $\Phi_{1\text{O}_2^*}$ is negatively correlated with MAC_{DOC}, indicating that less light-absorbing samples form $^1\text{O}_2^*$ more efficiently. Triplet concentrations determined by both probes linearly increase with DOC, and this relationship for [$^3\text{C}^*$]_{SYR} is independent of sample type. We find that approximately half of the total triplets are oxidizing based on SYR loss, while roughly half of the oxidizing triplets are strongly oxidizing based on PTA loss. FBB has the lowest average $\Phi_{3\text{C}^*}$, while atmospheric aging appears to enhance $\Phi_{3\text{C}^*}$, as well as $\Phi_{1\text{O}_2^*}$, based on the higher quantum yields for ABB samples.

675 Based on our results in dilute PM extracts (as well as past work), light absorption by brown carbon produces significant amounts of photooxidants in particles. To estimate the corresponding photooxidant concentrations, we extrapolate measured photooxidant kinetics in our particle extracts to an aerosol liquid water condition (1 μg PM/μg H₂O). Estimated concentrations of $^1\text{O}_2^*$, $^3\text{C}^*$, and $\bullet\text{OH}$ in ALW are on the order of 10^{-12} - 10^{-11} , 10^{-13} - 10^{-12} and 10^{-14} M with the ratio of $^1\text{O}_2^*$: $^3\text{C}^*$: $\bullet\text{OH}$ of (900 - 90) : (150 - 10) : 1. The corresponding ratio in our particle extracts is (40 - 5) : (1 - 10) : 1. For Win-Spr and Sum-Fall samples, singlet oxygen and oxidizing triplet concentrations increase significantly in ALW compared to in dilute extracts, while the changes in FBB and ABB are minor, likely due to the high DOC in the extracts, which cause strong quenching of $^1\text{O}_2^*$ and $^3\text{C}^*$. Compared to the predicted photooxidant concentrations in Davis winter particle water by Kaur et al. (2019), our Win-Spr predictions for [$\bullet\text{OH}$] and [$^3\text{C}^*$] are nearly 10 and 5 times higher, respectively, but our ALW value for [$^1\text{O}_2^*$] is 20 times lower. Based on our estimated ALW concentrations, lifetimes of organic compounds with high Henry's law constants in ALW can be significantly shortened compared to foggy conditions (Kaur et al., 2019), due to enhanced $^3\text{C}^*$ and $^1\text{O}_2^*$ concentrations in particle water.

While oxidant concentrations are required to calculate the lifetimes of individual organic species in ALW, the formation rate of a photooxidant provides insight into the overall significance of that oxidant as a sink for organics. Since organic compounds appear to be the major sink for all three photooxidants in ALW, the formation rate of an oxidant is approximately equal to the rate of DOM processing by that oxidant, although organics can also physically quench a triplet without transforming the organic (Grebil et al., 2011; Ma et al., 2021; Smith et al., 2014). Based on our extrapolations, the ratio of formation rates in ALW for $^1\text{O}_2^*$, $^3\text{C}^*$, and $\bullet\text{OH}$ (including mass transfer from the gas phase) is 1: 100: 4, taking Win-Spr as an example. Since the triplet formation rate is much higher than those of $\bullet\text{OH}$ or $^1\text{O}_2^*$, our results indicate that $^3\text{C}^*$ might be more important for the overall oxidation of organic compounds compared to the other two oxidants. However, the picture for any specific organic compound depends on its rate constants with each oxidant. For example, $\bullet\text{OH}$ will be relatively more important for organics that are less reactive with $^3\text{C}^*$ and $^1\text{O}_2^*$.

There are important uncertainties in the oxidant concentrations reported in our work. Foremost, predicting photooxidant concentrations from our dilute extracts to ALW conditions is highly uncertain as it requires extrapolating over a concentration difference of approximately a factor of 1000. While our current extracts have more DOC than those in our past work (Kaur et al., 2019), allowing us to get closer to ALW chemistry, we are still orders of magnitude more dilute. Despite this improvement, additional approaches – such as chamber and flow tube studies – are needed to measure photooxidants and their chemical impacts under conditions more similar to ambient aerosols. The oxidizing triplet concentrations are less certain than those of the other two



705 oxidants, both because we use an individual triplet (³DMB*) to model the wide range of natural triplet reactivities, but also because
of uncertainties in correcting the inhibition of syringol oxidation by particle components. Another uncertainty with our current
(and past) results is that we are missing the water-insoluble chromophores from particles. Consistent with past results from other
groups, we find significant amounts of highly light-absorbing water-insoluble brown carbon in our particle samples, suggesting
that by using aqueous extracts we are underestimating the concentrations and significance of photooxidants in ambient particles.
This issue should be addressed in future photochemistry studies.

710

Data availability

All data are available upon request.

Author contribution

715 CA and LM developed the research goals and designed the experiments. KB lent and set up the sampler, while
LM and CG collected samples. LM, RW, and LH performed the photochemistry experiments while WJ and CN analyzed OC and
ions, respectively. LM analyzed the data and prepared the manuscript with contributions from all co-authors. CA reviewed, wrote
portions of, and edited the manuscript. CA and QZ provided supervision and oversight during the experiments and writing.

Competing interests

The authors declare that they have no conflict of interest.

720 Acknowledgement

We gratefully acknowledge the following agencies for their publicly available data: the California Air Resources Board for PM_{2.5}
data, NASA's Land, Atmosphere Near real-time Capability for EOS (LANCE) system (<https://earthdata.nasa.gov/lance>), part of
the Earth Observing System Data and Information System (EOSDIS) for wildfire and smoke images, and the NOAA Air Resources
Laboratory (ARL) for the HYSPLIT model and READY website (<https://www.ready.noaa.gov>).

725 Financial support

This research has been supported by the National Science Foundation (AGS-1649212 and AGS-2220307); the California
Agricultural Experiment Station (Projects CA-D-LAW-6403-RR and CA-D-ETX-2102-H); and the University of California, Davis
(Donald G. Crosby Graduate Fellowships in Environmental Chemistry and Jastro Shields Research Awards).

730



References

- 735 Alfarrá, M. R., Prevot, A. S. H., Szidat, S., Sandradewi, J., Weimer, S., Lanz, V. A., Schreiber, D., Mohr, M. and Baltensperger, U.: Identification of the Mass Spectral Signature of Organic Aerosols from Wood Burning Emissions, *Environ. Sci. Technol.*, 41(16), 5770–5777, doi:10.1021/es062289b, 2007.
- Anastasio, C. and McGregor, K. G.: Chemistry of fog waters in California’s Central Valley: 1. In situ photoformation of hydroxyl radical and singlet molecular oxygen, *Atmos. Environ.*, 35(6), 1079–1089, doi:10.1016/S1352-2310(00)00281-8, 2001.
- 740 Arakaki, T., Anastasio, C., Kuroki, Y., Nakajima, H., Okada, K., Kotani, Y., Handa, D., Azechi, S., Kimura, T., Tshako, A. and Miyagi, Y.: A general scavenging rate constant for reaction of hydroxyl radical with organic carbon in atmospheric waters., *Environ. Sci. Technol.*, 47(15), 8196–8203, doi:10.1021/es401927b, 2013.
- Ashton, L., Buxton, G. V. and Stuart, C. R.: Temperature dependence of the rate of reaction of OH with some aromatic compounds in aqueous solution. Evidence for the formation of a π -complex intermediate?, *J. Chem. Soc., Faraday Trans.*, 91(11), 1631–
- 745 1633, doi:10.1039/FT9959101631, 1995.
- Baba, Y., Yatagai, T., Harada, T. and Kawase, Y.: Hydroxyl radical generation in the photo-Fenton process: Effects of carboxylic acids on iron redox cycling, *Chemical Engineering Journal*, 277, 229–241, doi:10.1016/j.cej.2015.04.103, 2015.
- Badali, K. M., Zhou, S., Aljawhary, D., Antiñolo, M., Chen, W. J., Lok, A., Mungall, E., Wong, J. P. S., Zhao, R. and Abbatt, J. P. D.: Formation of hydroxyl radicals from photolysis of secondary organic aerosol material, *Atmos. Chem. Phys.*, 15(14),
- 750 7831–7840, doi:10.5194/acp-15-7831-2015, 2015.
- Berg, S. M., Whiting, Q. T., Herrli, J. A., Winkels, R., Wammer, K. H. and Remucal, C. K.: The role of dissolved organic matter composition in determining photochemical reactivity at the molecular level., *Environ. Sci. Technol.*, 53(20), 11725–11734, doi:10.1021/acs.est.9b03007, 2019.
- Bilski, P., Holt, R. N. and Chignell, C. F.: Properties of singlet molecular oxygen in binary solvent mixtures of different polarity and proticity, *J. Photochem. Photobiol. A*, 109(3), 243–249, doi:10.1016/S1010-6030(97)00147-0, 1997.
- Bogler, S., Daellenbach, K. R., Bell, D. M., Prévôt, A. S. H., El Haddad, I. and Borduas-Dedekind, N.: Singlet oxygen seasonality in aqueous PM10 is driven by biomass burning and anthropogenic secondary organic aerosol, *Environ. Sci. Technol.*, doi:10.1021/acs.est.2c04554, 2022.
- Budisulistiorini, S. H., Riva, M., Williams, M., Chen, J., Itoh, M., Surratt, J. D. and Kuwata, M.: Light-Absorbing Brown Carbon
- 760 Aerosol Constituents from Combustion of Indonesian Peat and Biomass., *Environ. Sci. Technol.*, 51(8), 4415–4423, doi:10.1021/acs.est.7b00397, 2017.
- Canonica, S. and Laubscher, H.-U.: Inhibitory effect of dissolved organic matter on triplet-induced oxidation of aquatic contaminants, *Photochem. Photobiol. Sci.*, 7(5), 547, doi:10.1039/b719982a, 2008.
- Chen, Y., Hozalski, R. M., Olmanson, L. G., Page, B. P., Finlay, J. C., Brezonik, P. L. and Arnold, W. A.: Prediction of
- 765 photochemically produced reactive intermediates in surface waters via satellite remote sensing., *Environ. Sci. Technol.*, 54(11), 6671–6681, doi:10.1021/acs.est.0c00344, 2020.
- Claeys, M., Vermeylen, R., Yasmeen, F., Gómez-González, Y., Chi, X., Maenhaut, W., Mészáros, T. and Salma, I.: Chemical characterisation of humic-like substances from urban, rural and tropical biomass burning environments using liquid chromatography with UV/vis photodiode array detection and electrospray ionisation mass spectrometry, *Environ. Chem.*, 9(3),
- 770 273, doi:10.1071/EN11163, 2012.
- Corral Arroyo, P., David, G., Alpert, P. A., Parmentier, E. A., Ammann, M. and Signorell, R.: Amplification of light within aerosol particles accelerates in-particle photochemistry., *Science*, 376(6590), 293–296, doi:10.1126/science.abm7915, 2022.



- Dorfman, L. M. and Adams, G. E.: Reactivity of the hydroxyl radical in aqueous solutions, 0 ed., National Bureau of Standards, Gaithersburg, MD., 1973.
- 775 Du, Z., He, K., Cheng, Y., Duan, F., Ma, Y., Liu, J., Zhang, X., Zheng, M. and Weber, R.: A yearlong study of water-soluble organic carbon in Beijing II: Light absorption properties, *Atmos. Environ.*, 89, 235–241, doi:10.1016/j.atmosenv.2014.02.022, 2014.
- Ervens, B.: Progress and problems in modeling chemical processing in cloud droplets and wet aerosol particles, in *Multiphase environmental chemistry in the atmosphere*, vol. 1299, edited by S. W. Hunt, A. Laskin, and S. A. Nizkorodov, pp. 327–345, 780 American Chemical Society, Washington, DC., 2018.
- Ervens, B., Sorooshian, A., Lim, Y. B. and Turpin, B. J.: Key parameters controlling OH-initiated formation of secondary organic aerosol in the aqueous phase (aqSOA), *J. Geophys. Res. Atmos.*, 119(7), 3997–4016, doi:10.1002/2013JD021021, 2014.
- Ervens, B., Turpin, B. J. and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmos. Chem. Phys.*, 11(21), 11069–11102, doi:10.5194/acp-11-11069-2011, 785 2011.
- Fan, X., Li, M., Cao, T., Cheng, C., Li, F., Xie, Y., Wei, S., Song, J. and Peng, P.: Optical properties and oxidative potential of water- and alkaline-soluble brown carbon in smoke particles emitted from laboratory simulated biomass burning, *Atmos. Environ.*, 194, 48–57, doi:10.1016/j.atmosenv.2018.09.025, 2018.
- Farley, R., Bernays, N., Jaffe, D. A., Ketcherside, D., Hu, L., Zhou, S., Collier, S. and Zhang, Q.: Persistent Influence of Wildfire 790 Emissions in the Western United States and Characteristics of Aged Biomass Burning Organic Aerosols under Clean Air Conditions., *Environ. Sci. Technol.*, 56(6), 3645–3657, doi:10.1021/acs.est.1c07301, 2022.
- Faust, B. C. and Allen, J. M.: Aqueous-phase photochemical sources of peroxy radicals and singlet molecular oxygen in clouds and fog, *J. Geophys. Res.*, 97(D12), 12913, doi:10.1029/92JD00843, 1992.
- Finlayson-Pitts, B. J. and Pitts, J. N.: *Chemistry of the upper and lower atmosphere:: theory, experiments, and applications*, 795 Elsevier., 2000.
- Fleming, L. T., Lin, P., Roberts, J. M., Selimovic, V., Yokelson, R., Laskin, J., Laskin, A. and Nizkorodov, S. A.: Molecular composition and photochemical lifetimes of brown carbon chromophores in biomass burning organic aerosol, *Atmos. Chem. Phys.*, 20(2), 1105–1129, doi:10.5194/acp-20-1105-2020, 2020.
- Galbavy, E. S., Ram, K. and Anastasio, C.: 2-Nitrobenzaldehyde as a chemical actinometer for solution and ice photochemistry, 800 *J. Photochem. Photobiol. A*, 209(2-3), 186–192, doi:10.1016/j.jphotochem.2009.11.013, 2010.
- Gilardoni, S., Massoli, P., Paglione, M., Giulianelli, L., Carbone, C., Rinaldi, M., Decesari, S., Sandrini, S., Costabile, F., Gobbi, G. P., Pietrogrande, M. C., Visentin, M., Scotto, F., Fuzzi, S. and Facchini, M. C.: Direct observation of aqueous secondary organic aerosol from biomass-burning emissions., *Proc. Natl. Acad. Sci. USA*, 113(36), 10013–10018, doi:10.1073/pnas.1602212113, 2016.
- 805 Gollnick, K. and Griesbeck, A.: ChemInform Abstract: Singlet Oxygen Photooxygenation of Furans. Isolation and Reactions of [4 +2]-Cycloaddition Products (Unsaturated sec.-Ozonides), *Tetrahedron*, 41(11), 2057–2068, doi:10.1002/chin.198540196, 1985.
- Gonzalez, D. H., Cala, C. K., Peng, Q. and Paulson, S. E.: HULIS Enhancement of Hydroxyl Radical Formation from Fe(II): Kinetics of Fulvic Acid-Fe(II) Complexes in the Presence of Lung Antioxidants., *Environ. Sci. Technol.*, 51(13), 7676–7685, 810 doi:10.1021/acs.est.7b01299, 2017.



- González Palacios, L., Corral Arroyo, P., Aregahegn, K. Z., Steimer, S. S., Bartels-Rausch, T., Nozière, B., George, C., Ammann, M. and Volkamer, R.: Heterogeneous photochemistry of imidazole-2-carboxaldehyde: HO₂ radical formation and aerosol growth, *Atmos. Chem. Phys.*, 16(18), 11823–11836, doi:10.5194/acp-16-11823-2016, 2016.
- 815 Gebel, J. E., Pignatello, J. J. and Mitch, W. A.: Sorbic acid as a quantitative probe for the formation, scavenging and steady-state concentrations of the triplet-excited state of organic compounds., *Water Res.*, 45(19), 6535–6544, doi:10.1016/j.watres.2011.09.048, 2011.
- He, C., Liu, J., Carlton, A. G., Fan, S., Horowitz, L. W., Levy II, H. and Tao, S.: Evaluation of factors controlling global secondary organic aerosol production from cloud processes, *Atmos. Chem. Phys.*, 13(4), 1913–1926, doi:10.5194/acp-13-1913-2013, 2013.
- 820 Hecobian, A., Zhang, X., Zheng, M., Frank, N., Edgerton, E. S. and Weber, R. J.: Water-Soluble Organic Aerosol material and the light-absorption characteristics of aqueous extracts measured over the Southeastern United States, *Atmos. Chem. Phys.*, 10(13), 5965–5977, doi:10.5194/acp-10-5965-2010, 2010.
- Hems, R. F. and Abbatt, J. P. D.: Aqueous Phase Photo-oxidation of Brown Carbon Nitrophenols: Reaction Kinetics, Mechanism, and Evolution of Light Absorption, *ACS Earth Space Chem.*, 2(3), 225–234, doi:10.1021/acsearthspacechem.7b00123, 2018.
- 825 Hems, R. F., Schnitzler, E. G., Liu-Kang, C., Cappa, C. D. and Abbatt, J. P. D.: Aging of atmospheric brown carbon aerosol, *ACS Earth Space Chem.*, doi:10.1021/acsearthspacechem.0c00346, 2021.
- Herrmann, H.: Kinetics of aqueous phase reactions relevant for atmospheric chemistry., *Chem. Rev.*, 103(12), 4691–4716, doi:10.1021/cr020658q, 2003.
- Herrmann, H., Hoffmann, D., Schaefer, T., Brüner, P. and Tilgner, A.: Tropospheric aqueous-phase free-radical chemistry: radical 830 sources, spectra, reaction kinetics and prediction tools., *ChemPhysChem*, 11(18), 3796–3822, doi:10.1002/cphc.201000533, 2010.
- Herrmann, H., Schaefer, T., Tilgner, A., Styler, S. A., Weller, C., Teich, M. and Otto, T.: Tropospheric aqueous-phase chemistry: kinetics, mechanisms, and its coupling to a changing gas phase., *Chem. Rev.*, 115(10), 4259–4334, doi:10.1021/cr500447k, 2015.
- 835 Hess, M., Koepke, P. and Schult, I.: Optical properties of aerosols and clouds: the software package OPAC, *Bull. Amer. Meteor. Soc.*, 79(5), 831–844, doi:10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2, 1998.
- Hettiyadura, A. P. S., Garcia, V., Li, C., West, C. P., Tomlin, J., He, Q., Rudich, Y. and Laskin, A.: Chemical composition and molecular-specific optical properties of atmospheric brown carbon associated with biomass burning, *Environ. Sci. Technol.*, 55(4), 2511–2521, doi:10.1021/acs.est.0c05883, 2021.
- 840 Jiang, W., Ma, L., Niedek, C., Anastasio, C. and Zhang, Q.: Chemical and light absorption properties of water-soluble organic aerosols in Northern California and photooxidant production by brown carbon components, *ACS Earth and Space Chemistry*, in press, 2023.
- Kaur, R. and Anastasio, C.: Light absorption and the photoformation of hydroxyl radical and singlet oxygen in fog waters, *Atmos. Environ.*, 164, 387–397, doi:10.1016/j.atmosenv.2017.06.006, 2017.
- 845 Kaur, R. and Anastasio, C.: First measurements of organic triplet excited states in atmospheric waters., *Environ. Sci. Technol.*, 52(9), 5218–5226, doi:10.1021/acs.est.7b06699, 2018.
- Kaur, R., Labins, J. R., Helbock, S. S., Jiang, W., Bein, K. J., Zhang, Q. and Anastasio, C.: Photooxidants from brown carbon and other chromophores in illuminated particle extracts, *Atmos. Chem. Phys.*, 19(9), 6579–6594, doi:10.5194/acp-19-6579-2019, 2019.



- 850 Laskin, A., Laskin, J. and Nizkorodov, S. A.: Chemistry of atmospheric brown carbon., *Chem. Rev.*, 115(10), 4335–4382, doi:10.1021/cr5006167, 2015.
- Leresche, F., McKay, G., Kurtz, T., von Gunten, U., Canonica, S. and Rosario-Ortiz, F. L.: Effects of ozone on the photochemical and photophysical properties of dissolved organic matter., *Environ. Sci. Technol.*, 53(10), 5622–5632, doi:10.1021/acs.est.8b06410, 2019.
- 855 Leresche, F., Salazar, J. R., Pfothner, D. J., Hannigan, M. P., Majestic, B. J. and Rosario-Ortiz, F. L.: Photochemical aging of atmospheric particulate matter in the aqueous phase., *Environ. Sci. Technol.*, 55(19), 13152–13163, doi:10.1021/acs.est.1c00978, 2021.
- Li, M., Bao, F., Zhang, Y., Sheng, H., Chen, C. and Zhao, J.: Photochemical aging of soot in the aqueous phase: release of dissolved black carbon and the formation of IO₂, *Environ. Sci. Technol.*, 53(21), 12311–12319, doi:10.1021/acs.est.9b02773, 2019.
- 860 Lim, Y. B., Tan, Y., Perri, M. J., Seitzinger, S. P. and Turpin, B. J.: Aqueous chemistry and its role in secondary organic aerosol (SOA) formation, *Atmospheric Chemistry & Physics Discussions*, 10(6), 2010.
- Lin, P., Aiona, P. K., Li, Y., Shiraiwa, M., Laskin, J., Nizkorodov, S. A. and Laskin, A.: Molecular characterization of brown carbon in biomass burning aerosol particles., *Environ. Sci. Technol.*, 50(21), 11815–11824, doi:10.1021/acs.est.6b03024, 2016.
- Lin, P., Bluvshstein, N., Rudich, Y., Nizkorodov, S. A., Laskin, J. and Laskin, A.: Molecular Chemistry of Atmospheric Brown Carbon Inferred from a Nationwide Biomass Burning Event., *Environ. Sci. Technol.*, 51(20), 11561–11570, doi:10.1021/acs.est.7b02276, 2017.
- 865 Liu, J., Bergin, M., Guo, H., King, L., Kotra, N., Edgerton, E. and Weber, R. J.: Size-resolved measurements of brown carbon in water and methanol extracts and estimates of their contribution to ambient fine-particle light absorption, *Atmos. Chem. Phys.*, 13(24), 12389–12404, doi:10.5194/acp-13-12389-2013, 2013.
- 870 Ma, L., Guzman, C., Niedek, C., Tran, T., Zhang, Q. and Anastasio, C.: Kinetics and mass yields of aqueous secondary organic aerosol from highly substituted phenols reacting with a triplet excited state, *Environ. Sci. Technol.*, 55(9), 5772–5781, doi:10.1021/acs.est.1c00575, 2021.
- Ma, L., Worland, R., Jiang, W., Niedek, C., Guzman, C., Bein, K. J., Zhang, Q. and Anastasio, C.: Predicting photooxidant concentrations in aerosol liquid water based on laboratory extracts of ambient particles, *EGUsphere [preprint]*, doi:10.5194/egusphere-2023-566, 2023a.
- 875 Ma, L., Worland, R., Tran, T. and Anastasio, C.: Evaluation of probes to measure oxidizing organic triplet excited states in aerosol liquid water, *Environ. Sci. Technol.*, 57(15), 6052–6062, doi:10.1021/acs.est.2c09672, 2023b.
- Maizel, A. C. and Remucal, C. K.: The effect of probe choice and solution conditions on the apparent photoreactivity of dissolved organic matter., *Environ. Sci. Process. Impacts*, 19(8), 1040–1050, doi:10.1039/c7em00235a, 2017.
- 880 Manfrin, A., Nizkorodov, S. A., Malecha, K. T., Getzinger, G. J., McNeill, K. and Borduas-Dedekind, N.: Reactive oxygen species production from secondary organic aerosols: the importance of singlet oxygen, *Environ. Sci. Technol.*, 53(15), 8553–8562, doi:10.1021/acs.est.9b01609, 2019.
- McCabe, A. J. and Arnold, W. A.: Seasonal and spatial variabilities in the water chemistry of prairie pothole wetlands influence the photoproduction of reactive intermediates., *Chemosphere*, 155, 640–647, doi:10.1016/j.chemosphere.2016.04.078, 2016.
- 885 McCabe, A. J. and Arnold, W. A.: Reactivity of triplet excited states of dissolved natural organic matter in stormflow from mixed-use watersheds., *Environ. Sci. Technol.*, 51(17), 9718–9728, doi:10.1021/acs.est.7b01914, 2017.
- McCabe, A. J. and Arnold, W. A.: Multiple linear regression models to predict the formation efficiency of triplet excited states of dissolved organic matter in temperate wetlands, *Limnol. Oceanogr.*, 63(5), 1992–2014, doi:10.1002/lno.10820, 2018.



- McGregor, K. G. and Anastasio, C.: Chemistry of fog waters in California's Central Valley: 2. Photochemical transformations of
890 amino acids and alkyl amines, *Atmos. Environ.*, 35(6), 1091–1104, doi:10.1016/S1352-2310(00)00282-X, 2001.
- Mckay, G., Huang, W., Romera-Castillo, C., Crouch, J. E., Rosario-Ortiz, F. L. and Jaffé, R.: Predicting Reactive Intermediate
Quantum Yields from Dissolved Organic Matter Photolysis Using Optical Properties and Antioxidant Capacity., *Environ. Sci.
Technol.*, 51(10), 5404–5413, doi:10.1021/acs.est.6b06372, 2017.
- McNeill, K. and Canonica, S.: Triplet state dissolved organic matter in aquatic photochemistry: reaction mechanisms, substrate
895 scope, and photophysical properties., *Environ. Sci. Process. Impacts*, 18(11), 1381–1399, doi:10.1039/c6em00408c, 2016.
- McNeill, V. F.: Aqueous organic chemistry in the atmosphere: sources and chemical processing of organic aerosols., *Environ. Sci.
Technol.*, 49(3), 1237–1244, doi:10.1021/es5043707, 2015.
- Monge, M. E., Rosenørn, T., Favez, O., Müller, M., Adler, G., Abo Riziq, A., Rudich, Y., Herrmann, H., George, C. and D'Anna,
B.: Alternative pathway for atmospheric particles growth., *Proc. Natl. Acad. Sci. USA*, 109(18), 6840–6844,
900 doi:10.1073/pnas.1120593109, 2012.
- Nguyen, T. K. V., Zhang, Q., Jimenez, J. L., Pike, M. and Carlton, A. G.: Liquid water: ubiquitous contributor to aerosol mass,
Environ. Sci. Technol. Lett., 3(7), 257–263, doi:10.1021/acs.estlett.6b00167, 2016.
- Ossola, R., Jönsson, O. M., Moor, K. and McNeill, K.: Singlet oxygen quantum yields in environmental waters., *Chem. Rev.*,
121(7), 4100–4146, doi:10.1021/acs.chemrev.0c00781, 2021.
- 905 Park, S. S. and Yu, J.: Chemical and light absorption properties of humic-like substances from biomass burning emissions under
controlled combustion experiments, *Atmos. Environ.*, 136, 114–122, doi:10.1016/j.atmosenv.2016.04.022, 2016.
- Parworth, C. L., Young, D. E., Kim, H., Zhang, X., Cappa, C. D., Collier, S. and Zhang, Q.: Wintertime water-soluble aerosol
composition and particle water content in Fresno, California, *J. Geophys. Res. Atmos.*, 122(5), 3155–3170,
doi:10.1002/2016JD026173, 2017.
- 910 Paulson, S. E., Gallimore, P. J., Kuang, X. M., Chen, J. R., Kalberer, M. and Gonzalez, D. H.: A light-driven burst of hydroxyl
radicals dominates oxidation chemistry in newly activated cloud droplets., *Sci. Adv.*, 5(5), eaav7689,
doi:10.1126/sciadv.aav7689, 2019.
- Richards-Henderson, N. K., Pham, A. T., Kirk, B. B. and Anastasio, C.: Secondary organic aerosol from aqueous reactions of
green leaf volatiles with organic triplet excited states and singlet molecular oxygen., *Environ. Sci. Technol.*, 49(1), 268–276,
915 doi:10.1021/es503656m, 2015.
- Rolph, G., Stein, A. and Stunder, B.: Real-time Environmental Applications and Display sYstem: READY, *Environ. Model.
Softw.*, 95, 210–228, doi:10.1016/j.envsoft.2017.06.025, 2017.
- Rossignol, S., Aregahegn, K. Z., Tinel, L., Fine, L., Nozière, B. and George, C.: Glyoxal induced atmospheric photosensitized
chemistry leading to organic aerosol growth., *Environ. Sci. Technol.*, 48(6), 3218–3227, doi:10.1021/es405581g, 2014.
- 920 Schauer, J. J., Kleeman, M. J., Cass, G. R. and Simoneit, B. R.: Measurement of emissions from air pollution sources. 3. C1-C29
organic compounds from fireplace combustion of wood., *Environ. Sci. Technol.*, 35(9), 1716–1728, doi:10.1021/es001331e,
2001.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry And Physics: From Air Pollution To Climate Change*, 3rd ed., Wiley,
Hoboken, New Jersey., 2016.
- 925 Sharpless, C. M., Aeschbacher, M., Page, S. E., Wenk, J., Sander, M. and McNeill, K.: Photooxidation-induced changes in optical,
electrochemical, and photochemical properties of humic substances., *Environ. Sci. Technol.*, 48(5), 2688–2696,
doi:10.1021/es403925g, 2014.



- Silva, P. J., Liu, D.-Y., Noble, C. A. and Prather, K. A.: Size and chemical characterization of individual particles resulting from biomass burning of local Southern California species, *Environ. Sci. Technol.*, 33(18), 3068–3076, doi:10.1021/es980544p, 930 1999.
- Smith, J. D., Kinney, H. and Anastasio, C.: Aqueous benzene-diols react with an organic triplet excited state and hydroxyl radical to form secondary organic aerosol., *Phys. Chem. Chem. Phys.*, 17(15), 10227–10237, doi:10.1039/c4cp06095d, 2015.
- Smith, J. D., Sio, V., Yu, L., Zhang, Q. and Anastasio, C.: Secondary organic aerosol production from aqueous reactions of atmospheric phenols with an organic triplet excited state., *Environ. Sci. Technol.*, 48(2), 1049–1057, doi:10.1021/es4045715, 935 2014.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D. and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bull. Amer. Meteor. Soc.*, 96(12), 2059–2077, doi:10.1175/BAMS-D-14-00110.1, 2015.
- Tilgner, A. and Herrmann, H.: Tropospheric aqueous-phase OH oxidation chemistry: current understanding, uptake of highly oxidized organics and its effects, in *Multiphase environmental chemistry in the atmosphere*, vol. 1299, edited by S. W. Hunt, 940 A. Laskin, and S. A. Nizkorodov, pp. 49–85, American Chemical Society, Washington, DC., 2018.
- Tong, H., Arangio, A. M., Lakey, P. S. J., Berkemeier, T., Liu, F., Kampf, C. J., Brune, W. H., Pöschl, U. and Shiraiwa, M.: Hydroxyl radicals from secondary organic aerosol decomposition in water, *Atmos. Chem. Phys.*, 16(3), 1761–1771, doi:10.5194/acp-16-1761-2016, 2016.
- U.S. Geological Survey: Dissolved oxygen: U.S. Geological Survey Techniques and Methods, book 9, chap. A6.2, 33 p., 945 <https://doi.org/10.3133/tm9A6.2>, 2020.
- Verma, V., Fang, T., Xu, L., Peltier, R. E., Russell, A. G., Ng, N. L. and Weber, R. J.: Organic aerosols associated with the generation of reactive oxygen species (ROS) by water-soluble PM_{2.5}, *Environ. Sci. Technol.*, 49(7), 4646–4656, doi:10.1021/es505577w, 2015.
- Vicente, A., Alves, C., Calvo, A. I., Fernandes, A. P., Nunes, T., Monteiro, C., Almeida, S. M. and Pio, C.: Emission factors and detailed chemical composition of smoke particles from the 2010 wildfire season, *Atmos. Environ.*, 71, 295–303, doi:10.1016/j.atmosenv.2013.01.062, 2013.
- Wander, R., Neta, P. and Dorfman, L. M.: Pulse radiolysis studies. XII. Kinetics and spectra of the cyclohexadienyl radicals in aqueous benzoic acid solution, *J. Phys. Chem.*, 72(8), 2946–2949, doi:10.1021/j100854a044, 1968.
- Wang, G., Zhang, R., Gomez, M. E., Yang, L., Levy Zamora, M., Hu, M., Lin, Y., Peng, J., Guo, S., Meng, J., Li, J., Cheng, C., 955 Hu, T., Ren, Y., Wang, Y., Gao, J., Cao, J., An, Z., Zhou, W., Li, G., Wang, J., Tian, P., Marrero-Ortiz, W., Secrest, J., Du, Z., Zheng, J., Shang, D., Zeng, L., Shao, M., Wang, W., Huang, Y., Wang, Y., Zhu, Y., Li, Y., Hu, J., Pan, B., Cai, L., Cheng, Y., Ji, Y., Zhang, F., Rosenfeld, D., Liss, P. S., Duce, R. A., Kolb, C. E. and Molina, M. J.: Persistent sulfate formation from London Fog to Chinese haze., *Proc. Natl. Acad. Sci. USA*, 113(48), 13630–13635, doi:10.1073/pnas.1616540113, 2016.
- Wang, H., Zhou, H., Ma, J., Nie, J., Yan, S. and Song, W.: Triplet photochemistry of dissolved black carbon and its effects on the photochemical formation of reactive oxygen species., *Environ. Sci. Technol.*, 54(8), 4903–4911, doi:10.1021/acs.est.0c00061, 960 2020a.
- Wang, X., Gemayel, R., Hayeck, N., Perrier, S., Charbonnel, N., Xu, C., Chen, H., Zhu, C., Zhang, L., Wang, L., Nizkorodov, S. A., Wang, X., Wang, Z., Wang, T., Mellouki, A., Riva, M., Chen, J. and George, C.: Atmospheric photosensitization: A new pathway for sulfate formation., *Environ. Sci. Technol.*, 54(6), 3114–3120, doi:10.1021/acs.est.9b06347, 2020b.
- Wenk, J. and Canonica, S.: Phenolic antioxidants inhibit the triplet-induced transformation of anilines and sulfonamide antibiotics in aqueous solution, *Environ. Sci. Technol.*, 46(10), 5455–5462, doi:10.1021/es300485u, 2012.



- Wenk, J., Eustis, S. N., McNeill, K. and Canonica, S.: Quenching of excited triplet states by dissolved natural organic matter., *Environ. Sci. Technol.*, 47(22), 12802–12810, doi:10.1021/es402668h, 2013.
- Wenk, J., von Gunten, U. and Canonica, S.: Effect of dissolved organic matter on the transformation of contaminants induced by
970 excited triplet states and the hydroxyl radical., *Environ. Sci. Technol.*, 45(4), 1334–1340, doi:10.1021/es102212t, 2011.
- Wilkinson, F., Helman, W. P. and Ross, A. B.: Rate constants for the decay and reactions of the lowest electronically excited singlet state of molecular oxygen in solution. an expanded and revised compilation, *J. Phys. Chem. Ref. Data*, 24(2), 663–677, doi:10.1063/1.555965, 1995.
- Wong, J. P. S., Nenes, A. and Weber, R. J.: Changes in light absorptivity of molecular weight separated brown carbon due to
975 photolytic aging., *Environ. Sci. Technol.*, 51(15), 8414–8421, doi:10.1021/acs.est.7b01739, 2017.
- Wong, J. P. S., Tsagkaraki, M., Tsiodra, I., Mihalopoulos, N., Violaki, K., Kanakidou, M., Sciare, J., Nenes, A. and Weber, R. J.: Effects of atmospheric processing on the oxidative potential of biomass burning organic aerosols., *Environ. Sci. Technol.*, 53(12), 6747–6756, doi:10.1021/acs.est.9b01034, 2019.
- Yu, L., Smith, J., Laskin, A., Anastasio, C., Laskin, J. and Zhang, Q.: Chemical characterization of SOA formed from aqueous-
980 phase reactions of phenols with the triplet excited state of carbonyl and hydroxyl radical, *Atmos. Chem. Phys.*, 14(24), 13801–13816, doi:10.5194/acp-14-13801-2014, 2014.
- Zeinali, N., Oluwoye, I., Altarawneh, M. K., Almatarneh, M. H. and Dlugogorski, B. Z.: Probing the Reactivity of Singlet Oxygen with Cyclic Monoterpenes., *ACS Omega*, 4(9), 14040–14048, doi:10.1021/acsomega.9b01825, 2019.
- Zeng, L., Dibb, J., Scheuer, E., Katich, J. M., Schwarz, J. P., Bourgeois, I., Peischl, J., Ryerson, T., Warneke, C., Perring, A. E.,
985 Diskin, G. S., DiGangi, J. P., Nowak, J. B., Moore, R. H., Wiggins, E. B., Pagonis, D., Guo, H., Campuzano-Jost, P., Jimenez, J. L., Xu, L. and Weber, R. J.: Characteristics and evolution of brown carbon in western United States wildfires, *Atmos. Chem. Phys.*, 22(12), 8009–8036, doi:10.5194/acp-22-8009-2022, 2022.
- Zhang, X., Lin, Y.-H., Surratt, J. D. and Weber, R. J.: Sources, Composition and Absorption Ångström Exponent of Light-absorbing Organic Components in Aerosol Extracts from the Los Angeles Basin, *Environ. Sci. Technol.*, 47(8), 3685–3693,
990 doi:10.1021/es305047b, 2013.
- Zhao, R., Lee, A. K. Y., Huang, L., Li, X., Yang, F. and Abbatt, J. P. D.: Photochemical processing of aqueous atmospheric brown carbon, *Atmos. Chem. Phys.*, 15(11), 6087–6100, doi:10.5194/acp-15-6087-2015, 2015.
- Zuo, Y. and Hoigne, J.: Formation of hydrogen peroxide and depletion of oxalic acid in atmospheric water by photolysis of iron(III)-oxalato complexes, *Environ. Sci. Technol.*, 26(5), 1014–1022, doi:10.1021/es00029a022, 1992.
- 995