

1 **A Major Midlatitude Hurricane in the Little Ice Age**

2 **John Dickie^{1,2} and Grant Wach¹**

3 ¹Basin and Reservoir Lab, Department of Earth and Environmental Sciences

4 Dalhousie University, Halifax, Canada B3H 4R2

5 ²Corresponding Author

6 Contacts: john.dickie@dal.ca; grant.wach@dal.ca


7 **Abstract**

8 An unusually severe hurricane (Louisbourg Storm) struck Nova Scotia Canada in 1757. Historic
9 records describing storm conditions as well as damage to ships and coastal fortifications indicate
10 an intensity beyond any modern (post-1851) Atlantic cyclones striking the same region, yet this
11 storm struck during a cold climate period known as the Little Ice Age (LIA). Its track and timing
12 coincided with a British naval blockade of a French fleet at Fortress Louisbourg during the Seven
13 Years' War (1756-1763). This provides a unique opportunity to explore growing scientific
14 evidence of heightened storminess in the North Atlantic despite a colder climate expected to
15 suppress hurricane intensification but which research is increasingly showing to have supported
16 North Atlantic storms of exceptional strength. Weather attributes extracted from the logs of
17 naval vessels scattered by the Louisbourg Storm provided multiple hourly observations recorded
18 at different locations. Wave height and wind force estimates at ship locations were compared to
19 extreme storm surge heights calculated for Louisbourg Harbour and a shipwreck site south of
20 Fortress Louisbourg. Comparing these metrics to those of modern analogs that crossed the same
21 bathymetry reflects landfall intensity consistent with a powerful major hurricane. Historical
22 records show this storm originated as a tropical cyclone at the height of hurricane season and
23 intensified into the northern midlatitudes along the Gulf Stream. Its intensity at landfall is

24 consistent with established seasonal climatological models where highly baroclinic westerlies
25 driven by autumn continental cooling encounter intensifying north-tracking tropical cyclones
26 fueled by sea surface temperatures that peak in autumn. Stronger seasonal contrasts from earlier
27 and colder continental westerlies in the Little Ice Age (LIA) may have triggered explosive
28 extratropical transition from a large hurricane resulting in a more severe strike. It suggests that
29 tropical cyclones lasting days to weeks and the conditions that generate them are likely masked
30 by cooler historic mean-annual to multi-decadal LIA climate reconstructions.

31 1.0 Introduction

32 On September 25, 1757, a powerful hurricane struck the coast of Cape Breton Island,
33 Nova Scotia, Canada (Fig. 1). There would have had no record of the ‘Louisbourg Storm’ had it
34 not coincided with a British naval blockade of France’s Fortress Louisbourg during the Seven
35 Years’ War (1756-1763). Three French naval squadrons at Louisbourg and the blockading



36

37 **Figure 1.** Study location in Nova Scotia, Canada. Arrow length and orientation represents the
38 distance and direction traveled by the British fleet on September 21-26, 1757. September 25 and
39 26 show the path of the *Invincible* south of the wider dispersal of the British fleet after being
40 scattered by the storm (dotted oval). The storm's location off New England is estimated (off
41 map). The estimated storm track (dashed line) shows eye locations for the dates shown. Inset
42 shows the study area relative to the North Atlantic and the hurricane track based on historic
43 records showing its progressive northward translation seaward of (1) Florida (no date), (2) North
44 Carolina (September 23), (3) New England (September 24) and (4) Cape Breton Canada
45 (September 25-26). Fort Cumberland is 70 km toward 293° azimuth.

46 British fleet placed 49 sailing battleships and other warships (Supplemental Tables S1, S2) in the
47 path of a storm descriptions of damage to ships and coastal infrastructure, severe flooding from
48 rainfall and extreme storm surge suggest was more intense than any landfalling storm in
49 Canadian waters since modern records began in 1851 (Landsea et al., 2004, Finck, 2015). This
50 suggests it had the intensity of a major hurricane at landfall (Category 3+ on the Saffir-Simpson
51 Hurricane Wind Scale) yet it struck during the colder climate of the 'Little Ice Age' (LIA;
52 c1300-1850).

53 Hurricanes are fueled by sea surface temperatures (SSTs) over 28°C. They rapidly lose
54 energy as they move north over cooler midlatitude waters, and many tropical cyclones undergo
55 extratropical transition which releases tropical energy at increasingly higher latitudes later in
56 hurricane season (Hart and Evans, 2001). Modern tropical cyclone intensity is characterized in
57 real time with instruments carried by aircraft, satellites and at ground stations. In contrast, pre-
58 industrial metrics must be derived from historical observational records. Subjective interpretation
59 and geographic bias can make them less reliable than instrumental data (e.g., Jones and Mann

Commented [JD1]: period missing

60 2004), yet they offer a temporal resolution unavailable in natural climate archives, and they
61 straddle the end of the LIA and the rise of modern anthropogenic emissions. Oliver and Kington
62 (1970) and Lamb (1982) first explored their suitability for weather research. Naval logbooks
63 were subsequently found to be a superior source of historical weather data given that hourly ship
64 observations were systematically recorded in real time with a consistent terminology. Logbook
65 data have been compiled to assess historical atmospheric circulation patterns (e.g., Garcia et al.,
66 2001, Garcia-Herrera et al., 2005a, Wheeler et al., 2010, Barriopedro et al., 2014). CLIWOC, the
67 Climatological Database for the World's Oceans, was compiled from historical British, French,
68 Dutch and Spanish naval logbooks. It established a common historical wind force terminology to
69 document ocean surface atmospheric circulation patterns between 1750 and 1850 (Garcia-
70 Herrera et al., 2005b).

71 To date, pooled historical naval records have been used to identify longer-term regional
72 circulation patterns and extend the multidecadal climate signal into the industrial period (e.g.,
73 Garcia-Herrera et al., 2005a, 2005b, Wheeler et al., 2010, Barriopedro et al., 2014). In contrast,
74 this study takes advantage of an unusual concentration of warships in the path of a single
75 hurricane to characterize its intensity. It seems counterintuitive that the colder LIA climate would
76 generate more powerful midlatitude Atlantic cyclones than in the modern era, yet historical
77 records show the LIA to be generally 'stormier' with unusually powerful midlatitude hurricanes
78 despite conditions that dampen hurricane energy. Donnelly et al.'s (2001) historic storm
79 reconstruction from Mattapoissett Pond, Massachusetts, and Oliva et al.'s (2018) historic storm
80 reconstruction from Robinson Lake, Nova Scotia, are among a growing number of proxy studies
81 showing that major Atlantic cyclones struck the northeastern seaboard of North America in the
82 LIA. Since winter extratropical cyclones known as Nor'easters cannot be differentiated from

83 Atlantic tropical cyclones and their extratropical derivatives from proxy data alone, historical
84 records can constrain the timing of midlatitude hurricanes and tropical storms.

85 This study utilizes a unique historical data set to characterize the intensity of the
86 Louisbourg Storm using spatial and temporal weather metrics extracted from ship logbooks of
87 both the English and French fleets, British Admiralty records and official documents of both
88 nations, and compares the derived storm metrics to those of modern systems that tracked across
89 the same bathymetry. Characterizing its intensity tests historical descriptions of an unusually
90 severe storm and may help establish a more thorough understanding of LIA hurricane
91 climatology.

92 **2.0 Methodology**

93 *2.1 Historical Records*

94 Eighteenth century navigation and weather data were entered hourly in the daily logs of
95 naval vessels, resulting in reliable records suitable for historical climate research. A noon
96 sighting of the sun fixed latitude and marked the start of the sea day. Britain adopted the
97 Gregorian calendar in 1752 so dates in logs used for this study did not require correction. In 1757
98 a local meridian was used to determine longitude, deduced from logs to have been based on
99 Louisbourg Lighthouse (Fig. 2).

100 Historical British Admiralty Correspondence and Papers (ADM1/481, 1488, 2294)
101 covering storm damage to British vessels on the ‘Halifax Station’ in 1757 and Fleet Lists
102 (ADM8/31, 32) are preserved at the National Archives at Kew (UK), as are Royal Navy Master’s
103 (ADM 51/409, 633,1075) and Captain’s (ADM 52/578,819,1064) logbooks. Lieutenant’s logs
104 (ADM51) kept at the National Maritime Museum, Greenwich, were often incorporated into
105 Captain’s logs with addenda. Master’s and Captain’s logs of the Royal Navy warships *Invincible*,

106 *Windsor, Sunderland, Eagle, Terrible, Grafton, Newark, and Captain*, plus ancillary official
107 correspondence, were used in this study. The first author reviewed all logs and found them to be
108 consistent in content and format, then he copied letters and logbook entries written in cursive in
109 multiple handwriting styles to a more readable format (Supplemental Fig. S1). These were
110 interpreted, compiled into a time sequence and cross referenced. Logs from French warships
111 *Fleur de Lys, l'Abenaquise, Tonnant, l'Inflexible* and *Dauphin Royal* translated from French
112 describe conditions in Louisbourg Harbour (McLennan, 1918). Wind directions from gimballed
113 ships' compasses reference magnetic north. Bearings and wind directions used the 32 points of
114 the compass (Smyth, 1867, Blake and Lawrence, 1999) and were translated to azimuths. The
115 logs of British ships at sea and French ships moored in Louisbourg Harbour contained: (1) dates
116 and times, (2) position, (3) bearing, (4) wind direction, (5) wind speed terms that evolved into the
117 Beaufort Wind Scale (e.g., Garcia-Herrera et al., 2005a, 2005b, Wheeler, 2005, Wheeler et al.,
118 2010), and (6) descriptions of sea state.

119 2.2 *Climate Context*

120 Major atmospheric circulation patterns that influence Atlantic tropical cyclone behaviour,
121 specifically the El Nino Southern Oscillation (ENSO) and North Atlantic Oscillation (NAO),
122 have been reconstructed for the historical period (e.g., Gurgis and Fowler, 2009, Trouet et al.,
123 2012). These trends provide an overarching context since La Nina years create conditions
124 conducive to driving hurricanes in the Atlantic, and a negative NAO allows Atlantic tropical
125 cyclones to enter the Atlantic and potentially reach the midlatitude eastern seaboard.
126 Atmospheric circulation patterns for 1757 were studied to assess overarching conditions
127 conducive to Atlantic hurricane generation.

128 2.3 *Wind Speed*

Commented [JD2]: clarify figure is supplemental

129 Wheeler and Wilkinson's (2004) analysis of the derivation of the Beaufort scale shows
 130 terms that vary little from the logbook terms used in this study. A similar approach has been
 131 adopted here with adjectives describing primary nomenclature. A 'gale' (Beaufort Force 8) was
 132 originally between a breeze (Force 2) and a violent storm (Force 11) and established a
 133 benchmark (Table 1). A 'near gale,' its diminutive (Smyth, 1867) corresponds to a 'moderate
 134 gale.' Wheeler et al. (2010) categorized 'strong gale,' 'hard gale,' 'blew hard' and 'storm' as
 135 stronger than 'fresh gale.' Adjectives 'stiff' and 'fresh' indicate winds stronger than a gale
 136 (Force 9) while a 'severe' or 'hard' gale reflects a 'storm' (Force 10). 'Excessive' and 'extreme'
 137 hard gale, necessarily stronger than a 'hard gale' would then correspond to 'violent storm' (Force
 138 11) which does not appear in the logs used here. 'Hurricane' (Force 12) is mentioned in both
 139 French and British records. 'Squall' is a historical term for an increase in wind speed sustained
 140 above threshold for at least one minute. The National Oceans and Atmospheric Administration
 141 (NOAA) defines it as a sudden increase by at least 16 knots (33 ~~kph~~km h⁻¹) and sustained at over
 142 22 knots (41 ~~km h-1~~) for one minute. Environment and Climate Change Canada (ECCC) defines

Commented [JD3]: change kph to kmh⁻¹

Deleted: kph

Logbook Term	Beaufort Scale	Rating	Wind (kph)
Hurricane	Hurricane	12	118+
Excessive / Extreme Hard Gale	Violent storm	11	103-117
Severe / Hard Gale	Storm	10	89-102
Strong / Stiff Gale	Strong Gale	9	75-88
Gale	Gale	8	62-74
Moderate Gale	Near Gale	7	50-61
Strong / Stiff Breeze	Strong Breeze	6	39-49

Logbook Term	Beaufort Scale	Rating	Wind (km h ⁻¹)
Hurricane	Hurricane	12	118+
Excessive / Extreme Hard Gale	Violent storm	11	103-117
Severe / Hard Gale	Storm	10	89-102
Strong / Stiff Gale	Strong Gale	9	75-88
Gale	Gale	8	62-74
Moderate Gale	Near Gale	7	50-61
Strong / Stiff Breeze	Strong Breeze	6	39-49

Logbook Term
Hurricane
Excessive / Extreme Hard Gale
Severe / Hard Gale
Strong / Stiff Gale
Gale
Moderate Gale
Strong / Stiff Breeze

Deleted:

144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160

Table 1. Logbook Beaufort Terms and Associated Windspeeds (kph km h⁻¹).
squalls as increases of 34 knots (63 kph km h⁻¹) or more above prevailing winds sustained for
over a minute. The World Meteorological Organization (WMO) uses 8 m/s m s⁻¹ and 11 m/s m s⁻¹
(29 and 40 kph km h⁻¹) above threshold for over one minute while the American
Meteorological Association (AMA) notes squalls are of ‘several minutes’ duration. In
considering these definitions ‘squall’ is taken to be a sudden increase in wind speed of 40-60 kph
km h⁻¹ above threshold and sustained for at least one minute. We interpret ‘hard’ squalls as the
upper end of the spectrum by applying the same historical adjectives used to create the historic
Beaufort scale (Wheeler and Wilkinson, 2004). Heavy rains accompanying squalls noted in the
logs appear to be consistent with descriptions of hurricane spiral bands.

In this study the Beaufort Wind Force Scale is used to describe wind speeds from gale to
hurricane force (63-118 kph km h⁻¹). The Saffir-Simpson Hurricane Wind Scale describes
hurricane winds greater than 118 kph km h⁻¹ with peak wind speeds averaged over one minute
defining hurricane intensity Categories 1-5. A major hurricane is Category 3 (178-208 kph km h⁻¹)
or stronger. Wind speeds derived from log entries were plotted from the first southeasterlies
noted off Nova Scotia on September 22, 1757, to diminishing westerlies at the storm’s end on

Commented [JD4]: change kph to km h⁻¹

Commented [JD5]: change kph to km h⁻¹

Commented [JD6]: change kph to km h⁻¹

Commented [JD7]: change kph to km h⁻¹

Commented [JD8]: change kph to km h⁻¹

162 September 26. Ephemeral squalls of 1 min duration above threshold winds provide an estimate
163 of total wind speed sustained for one minute or longer. Wind speeds at mid-mast height above
164 the deck plus freeboard (distance from the waterline to the upper deck) approximate the 10 m
165 height above ground level for modern hurricane wind speed measurements.

166 Eighteenth century navies knew hurricanes commonly encountered in the Caribbean
167 sometimes reached North America's eastern seaboard. Since no real time wind force
168 measurement existed in 1757, to measure and categorize hurricane intensity, this study has
169 adopted Virot et al.'s (2016) engineering analysis of critical hurricane wind speeds that break
170 trees as a model for estimating threshold wind speeds needed to break ships' masts. *Invincible's*
171 log indicates it maintained course relative to prevailing storm winds. This placed the vessel
172 oblique to wave crests which minimized pitch and yaw, and held masts within a stable plane of
173 reference against which wind applied a sustained force. In addition, large vessels (74-gun third
174 rates) with up to nine feet of flooding in the hold would have a lower center of mass that would
175 have affected its righting moment and minimized directional variance in the wind force striking
176 the masts. Rigging designed to stabilize the masts and transfer wind energy through the sails
177 would likely have required a higher sustained wind force to achieve failure.

178 *2.4 Wind Direction*

179 Wind direction was measured using the ship's magnetic compass and entered in the
180 ships' logs as 'points of the compass.' These entries were translated to azimuths. Compass
181 directions are relative to magnetic north and not corrected for declination given the small study
182 area and short time frame. Eighteenth century navigation was inaccurate but this study benefits
183 from (1) log entries of the fleet relying on smaller vessels sent inshore to establish distance from
184 coastal landmarks, and (2) during the storm ships were driven sufficiently close to land that their

185 positioning entries were based on triangulation using landmarks which greatly improves
186 accuracy. Experienced navigators were also able to correct for ship motion in their readings
187 while the ship's position was typically determined by a Lieutenant plus one or more midshipmen
188 and the sailing master's mate.

189 *2.5 Wave Height*

190 Wave height was estimated based on descriptions compared to ship dimensions and is the
191 last accurate metric. Historic references to ship structure in Imperial Units have been converted
192 to metric. This includes the distance from the keel to the upper deck and freeboard from the
193 waterline to the upper deck. The depth of water needed to spill over the bow to flood the upper
194 deck and tear away large ship's boats tethered to the deck is estimated. References such as
195 sailors being swept off spars 24 m (80') above the waterline offers an estimate of peak wave
196 heights. Warships were designed for stability as floating gun platforms and to return to an 'even
197 keel' as quickly as possible after firing. Wave descriptions in Louisbourg Harbour are the least
198 reliable since they include storm surge.

199 *2.6 Surge*

200 Surge is a rise in sea level due to atmospheric pressure and storm winds and is
201 proportional to a tropical cyclone's intensity and translation rate. Coastal surge is a reasonable
202 estimate of storm intensity and can serve as a test of intensity derived from wind data. The surge
203 height of modern analogs that struck Nova Scotia after tracking across the Scotian Shelf and
204 whose intensity has been characterized with metrics derived using modern meteorological
205 methods provides a reliable benchmark for comparison to surge calculated for the 1757 storm. In
206 this study, storm surge at known locations and elevations above sea level were described at (1)
207 Battery de la Grave at Fortress Louisbourg and (2) the historic town within the Fortress (Fig. 2),

Deleted: ' (24 m

209 and (3) St. Esprit (Fig. 1) where the British warship *HMS Tilbury* was stranded in water depths it
210 could not normally navigate given its displacement. All surge calculations were then corrected
211 for (1) relative sea level (RSL) rise since 1757, and (2) a mid-tide RSL datum used by Google
212 Earth versus a lowest low water (tide) datum used by the Canadian Hydrographic Service for a
213 (draft) navigation chart used for the *Tilbury* wreck site. In addition, French records noting the
214 tidal change at Louisbourg allowed for the timing of the tidal cycle to be backed out to determine
215 storm surge versus storm tide.

216 *Tilbury's* wreck site offered a chance to estimate surge at a second location 45 km
217 southwest of Louisbourg. *Tilbury's* identity was confirmed in 1986 with the discovery of the
218 ship's bell, most of its guns, anchors and artifacts (Storm, 2002). Locating the wreck to confirm
219 its water depth required creating a digital bathymetric chart needed to guide a marine
220 magnetometer survey leading to site confirmation by divers.

221 **3.0 Little Ice Age Storminess**

222 Matthes (1939) named the LIA to explain European glacier expansion during a
223 historically colder climate period. Heightened climate variability saw deeply cold winters and
224 cooler mean annual temperatures primarily in the northern hemisphere (e.g., Kreutz et al., 1997,
225 Mann, 2002, Jones and Mann, 2004). It may have been triggered by late 13th Century volcanic
226 eruptions and a cooling feedback process sustained by Arctic sea-ice expansion (Miller et al.,
227 2012). North Atlantic mean annual SSTs were 1-2°C cooler than today (e.g., Keigwin, 1996,
228 Winter et al., 2000, Richey et al., 2009, Saenger et al., 2009, Cronin et al., 2010, Bertler et al.,
229 2011, Mazzarella and Scaffeta, 2018, Gebbie, 2019). The Maunder Minimum, the coldest part of
230 the LIA, (MM; 1645-1715) saw greater 'storminess' during polar air breakouts from Europe
231 correlating to more frequent easterly gales in the English Channel and Approaches in 1685-1750

232 (Wheeler et al. 2010). Concentrated storm horizons in coastal dunes across western Europe and
233 in Brittany and on France's Mediterranean coast correlate to the coldest part of the LIA
234 (Dezileau et al., 2011, Van Vliet-Lanoe et al., 2014, Sicre et al., 2016, Jackson et al. 2019).
235 Dezileau et al. (2011) attributed LIA storminess to cold-enhanced lower tropospheric
236 baroclinicity modifying prevailing westerlies. In the northwest Atlantic, Donnelly et al. (2015)
237 described major hurricane deposits in New England coastal sediments dating to 1635, 1638 and
238 1815. Ludlum's (1963) compilation of historical northwest Atlantic hurricanes and tropical
239 storms includes the LIA's major 'Independence Hurricane' that struck New England on August
240 29, 1775 and the 'Newfoundland Hurricane' of September 9, 1775, a storm that left 4000 dead to
241 become Canada's deadliest hurricane (Ludlum, 1963, Ruffman, 1996). Lamb's (1991)
242 exhaustive survey of British and European storms includes the Great Storm that devastated the
243 British Isles on November 26, 1703. It was an extratropical cyclone equal to a Category 2
244 hurricane yet Wheeler (2003) notes a far more powerful Atlantic storm on December 1-12, 1792,
245 also late in Atlantic hurricane season. Both were anomalous for a colder climate period.

246 The Scotian Shelf on Canada's Atlantic seaboard (Fig. 1) is dominated by the cold, south-
247 flowing, low-salinity Labrador Current. It originates in the Davis Strait of the Canadian Arctic
248 and hugs the coast to the start of the midlatitudes at Cape Hatteras, North Carolina where it
249 meets, mixes with, and redirects seaward the tropical, north-flowing more saline Gulf Stream.
250 The Labrador Current plays a critical role in hurricane extratropical transition by providing a
251 coastal buffer of cooler sea surface temperatures that effectively cut off the tropical energy of the
252 Gulf Stream (Hart and Evans, 2001). Summer and fall bring warm water eddies from the Gulf
253 Stream and warmer coastal SST. Sediment cores from the Emerald Basin off Nova Scotia show
254 1600 years of cold Labrador Current temperatures and a sudden and sustained warming around

255 1850 that has continued into the present (Keigwin et al., 2003) and coincides with the end of the
256 LIA. Storm compilations by Landsea et al. (2004) and Chenowith (2006) show a progressive
257 increase in the number of historical Atlantic tropical cyclones from 1700 and a sharp increase in
258 the number and percentage reaching New England and eastern Canada beginning around 1850.
259 Vecchi and Knutson (2008) in a study of data from the start of instrumental data collection in
260 1880 show a strong correlation between mean annual SST and storm frequency.

261 Historical records offer seasonal weather detail not captured by annual to multidecadal
262 proxy trends. Anomalous midlatitude coastal sea surface temperatures (SSTs) over days to
263 weeks, conditions that fuel tropical cyclones, are therefore not likely to appear in annualized data
264 weighted by colder, sustained LIA winters. Northern and Arctic temperature reconstructions for
265 temperate North America show cooler mean temperatures over the whole of the LIA (e.g.,
266 Jacoby and D'Arrigo, 1989 and Trouet et al., 2013). Trouet et al., (2013) demonstrate a multi-
267 decadal warming to cooling trend peaking in the mid-eighteenth century.

268 Lieutenant John Knox recorded unusually high temperatures in Halifax Harbour on July
269 20, 1757, which fellow officers found hotter than Gibraltar and the Mediterranean (Knox, 1769).
270 This coincided with a heat wave in Britain and southwest Europe from July into early August
271 1757 that set temperature records that stood for over 250 years (The London Chronicle, July 23-
272 26, 1757; London Magazine, November 1758 p. 563-4). London on July 16-26 had an average
273 high of 41.2C (Nature Notes, 24 August 1882, p. 415). This does not assume weather conditions
274 in Europe fueled a hurricane tracking into Atlantic Canada, but demonstrates that unusually hot
275 temperatures across the northern hemisphere capable of warming midlatitude SSTs that intensify
276 midlatitude hurricanes existed in the summer of 1757.

277 The 1757 hurricane noted by Poey (1855) and Ludlum (1963) was confirmed as a
278 hurricane, storm 73 in Table IV in Chenowith's (2006) re-assessment compilation. It was first
279 seen off Florida and followed the coastline past Cape Hatteras to New England on September 22-
280 24 (Ludlum, 1963). Benjamin Franklin's observations of this specific storm led him to conclude
281 that hurricanes "are produced by currents of cold winds rushing from the north along the Atlantic
282 coast and mingling with the warm winds produced by the gulf-stream" (Warden, 1819). It struck
283 the British frigate *HMS Winchelsea* on September 23 to 24 at 36°45'N 70° 54'W (off North
284 Carolina over the Gulf Stream). The log notes gale force east then east-southeast and south winds
285 between 10 p.m. and 5 a.m. on September 23 which, 15 minutes later, veered violently to the
286 northeast and then northwest at 'near hurricane' intensity. It split the main sail and broke the
287 main mast and was accompanied by a 'great sea' (ADM 52/1105).

288 The storm passed New England on September 24 (Boston Herald, Oct. 17, 1757, Ludlum
289 1963) and struck Nova Scotia as the Louisbourg Storm on September 25, 1757. Its arrival at Fort
290 Cumberland on the Nova Scotia border 200 km inland late September 22 included 'violent rain'
291 and 'constant heavy rain' into the 23rd. Knox's journal on the 27th describes September 24-26
292 with ... 'I never saw such storms of wind and rain as we have had for some days past...'
293 followed by 'windy, showery and very cold' weather on the 27-28th and 'dry, cold windy
294 weather' on the 29th, followed by frost and snow across Nova Scotia by mid-October (Knox
295 1769).

296 **4.0 Historical Context**

297 The Seven Years' War (1756-1763) arose from unresolved issues following the Treaty
298 of Aix-la-Chappelle that ended the War of the Austian Succession (1740-1748). It began as a
299 European conflict between Great Britain and allies and France and its allies, but soon extended to

300 the colonial interests of both nations in North America and India. It resulted in significant losses
301 for France including the loss of New France, now Canada, to Great Britain (Syrett, 2008).
302 Britain's overwhelming success in gaining territory at France's expense during the war led
303 France to subsequently support the secession of the American colonies in 1775.

304 Great Britain's 'Grand Plan' for the North American campaign began with John
305 Campbell, the 4th Earl of Loudoun, being appointed Commander-in-Chief of the British military
306 in North America. His adversary was Louis-Joseph de Montcalm-Grozon, Marquis de Montcalm
307 de Saint-Veran, commander of French forces in North America. To attack Montcalm at Quebec
308 without leaving a powerful French fortress at his rear, Loudoun needed to first seize Fortress
309 Louisbourg in Nova Scotia. On May 22 to 25, 1757, troops boarded 134 transport ships in New
310 York to rendezvous at Halifax with a fleet departing Britain under Vice Admiral Frances
311 Holbourne. Pitt's brief removal as Prime Minister delayed the fleet but his return to power with a
312 coalition government saw it depart Cork, Ireland, on May 8, 1757. The delay allowed France to
313 reinforce Louisbourg with three naval squadrons ahead of the British arrival. On May 23 five
314 French battleships and a frigate under Chevalier Joseph de Beaufremont arrived from the West
315 Indies, followed on June 15 by four battleships and two frigates under Joseph Francois de Noble
316 du Revest from Toulon. On June 20 nine battleships and two frigates under Vice Admiral
317 Emmanuel-Auguste de Cahideuc (Comte Dubois de la Motte) arrived from Brest. 4000 French
318 troops bolstered a garrison of 3200 plus 300 Acadians and Mi'kmaq warriors (McLennan, 1918,
319 Stoetzel, 2008). Holbourne's arrival at Halifax on June 30 bolstered Loudoun's force to create an
320 army of 12 000. *HMS Gosport* arrived on August 5 with letters intercepted from a French
321 schooner captured off Newfoundland detailing Louisbourg's reinforcement. It rendered the

Formatted: Font: Not Italic

322 attack on the fortress untenable. Loudoun returned to New York and on September 11, 1757

323 Holbourne sailed his fleet north to blockade Louisbourg (Fig 1).

324 **5.0 The Louisbourg Storm**

325 The British fleet cruised off the coast of Cape Breton Nova Scotia (Fig. 1) to lure the
 326 French fleet out of Louisbourg Harbour to do battle. On September 21, the British 80-gun
 327 flagship *Newark* noted fresh westerly gales followed by fair weather and light breezes then calm
 328 with fog on the 22nd. That day an officer on the French 28-gun frigate *Fleur de Lys* saw a low
 329 mist enter Louisbourg Harbour. The mist was also seen at sea by the British *Invincible* until it
 330 dissipated under a rising southeast breeze. Britain's *Newark* and France's *Fleur de Lys* recorded
 331 that the breeze veered to the southeast and intensified to moderate gales on September 22. The
 332 *Invincible* recorded strengthening easterlies September 22-26 from otherwise prevailing
 333 westerlies through the second half of September (Table 2).


Commented [JD9]: Insert new Table 2 with no coloured cells (replaced by shaded B&W)

SEPT 16			SEPT 17			SEPT 18		
SW	SW	WSW	SW	W	NNW	NNW	NNW	NNW
225	225	247.5	225	270	337.5	337.5	337.5	337.5
SEPT 19			SEPT 20			SEPT 21		
NNW	NE	WNW	WSW	WSW	W	W	W	NNW
337.5	45	292.5	247.5	247.5	270	270	270	337.5
SEPT 22			SEPT 23			SEPT 24		
SE	SSE	SEBS	SE	SE	SEBS	SEBS	SEBS	EBS
135	157.5	146.25	135	135	146.25	146.25	146.25	101.25
SEPT 25			SEPT 26			SEPT 27		
EBS	SW	W	W	W	NW	SWBW	SEBS	WBS
101.25	225	270	270	270	315	236.25	146.25	258.75

334
 335 **Table 2.** Prevailing Winds (*HMS Invincible* logbook)

336 Prevailing wind direction measured for each of three successive 8-hour watches per day and
337 azimuth equivalent on the *Invincible*. Storm winds, arriving September 22, 1757, off Cape
338 Breton, are shaded and in italics; two watches with easterlies not associated with the storm are
339 shaded only. Mean 250.5 (WSW) prevailing wind direction six days before and five days
340 following storm (continued westerly on 28 and 29). Mean 135 (SE) wind direction during storm.
341 Ships off St. Esprit on September 25 saw prevailing southeasterly winds last until September 26.
342 Ships south of St. Esprit including *Invincible*, *Sunderland* and *Windsor* faced southwesterly
343 winds on September 25. ‘B’ stands for ‘by,’ a historical modifier defining a point of the compass
344 (e.g., SWBW means southwest by west which is 11.25° west of southwest or 236.25° azimuth).

345 French naval officers, expecting a storm due to the southeast winds, moored the French
346 fleet in two lines off Royal Battery (Fig. 2) with four 2-ton anchors set from the bow of each ship
347 with four 20 cm diameter anchor cables. The southeast winds led the British ships at sea to
348 secure masts and naval guns, weighing as much as 3 tons apiece, anticipating a storm. On the
349 24th *Invincible* and *Newark* reported increasing cloud, haze and rain beginning under southeast
350 gales.



351

352 **Figure 2.** Louisbourg Harbour showing the French fleet anchorage, Louisbourg Lighthouse,
 353 Royal Battery, Battery de la Grave Guardhouse, and the southeast seawall overlain on chart
 354 image © Canadian Hydrographic Service (2011) Chart Guyon Island to Flint Island 1:37,866
 355 [Issued 2022-11-26]. Shoals (shaded) relative to ship hull displacements of 5.8-7.0 m (19 to 23')
 356 give a general sense of the scale of waves and surge needed to throw battleships on shore and
 357 destroy the southeast facing seawall.

358 On September 25 fresh southeast gales rose to excessive hard gales with very heavy rain.
 359 The British *Windsor* noted heavy rain and mist and intensifying strong gales with hard squalls.
 360 At 7 p.m. *Sunderland* faced very hard gales that rose to extreme hard gales by 10 p.m. At 12
 361 a.m. *Invincible* faced strong gales, torrential rains and a 'great sea.' At 2 a.m. on the 25th
 362 *Invincible* noted an 'excessive hard gale' and 'a hurricane of wind' and mountainous waves.
 363 Topsails used to control ships in severe weather were 'blown to rags.' *Sunderland's* main
 364 staysail was torn away. Waves 'made a free passage over...' the 70-gun *Devonshire* and

Commented [JD10]: Figure with colour cells changed to Black and White

365 smashed in *Lightning's* stern. The wind tore away the 8-gun *Cruiser* sloop's mizzen mast and
366 three sailors were swept overboard. *Cruiser* was 'very near foundering having been underwater
367 several times' and jettisoned its guns to stay afloat.

368 *Windsor's* log records extreme gales with severe squalls, heavy rain and a great sea.
369 Canvas tarpaulins stripped off deck gratings by the wind allowed waves and rain to flood the
370 ships which soon had up to 2.5 m (9') of water in the holds despite the pumps in full operation.
371 *Windsor* and *Sunderland* sailed S across SSW winds. *Grafton's* three-ton 7 m (30') rudder was
372 torn off the ship. *Invincible's* rudder, also torn free, was only saved by its preventer chains.
373 Sails on all the British ships at sea were torn away by the wind. Captain Bently later reported
374 that *Invincible's* hull planking had opened and strain on the hull broke iron reinforcing brackets
375 and bolts, allowing the entire gun deck and its tens of tons of heavy naval guns to drop several
376 inches (Captain's Letters, ADM 1/1488). *Sunderland's* foretopmast, reinforced by ten 5 cm (2")
377 rope shrouds plus stays, was torn off the ship and it disappeared into the night with two sailors.
378 *Invincible* was thrown onto her 'beam ends' (side), forcing it to heave overboard ten 12-pounder
379 upper deck guns and carriages, roughly twenty tons, to right the ship. *Invincible's* main yard was
380 ordered taken down but before it could be done the wind broke off the 38" (1 m) diameter
381 mainmast 20' (6 m) above the deck. The falling mast tore down the foretopmast and mizzen mast
382 and crushed the starboard gunwale. The wreckage pulled the ship onto its side and swept sailors
383 John Guttredge and Samuel Kirby into the sea. *Invincible's* sailors cut the tangled mass free
384 before it sank the ship.

385 At Louisbourg, the French military officer at La Grave Battery (Fig. 2) led his troops to
386 safety after the sea rose steadily above their knees (Chevalier de Johnstone, 1758). Offshore, the
387 British 14-gun *Ferret* sloop under Francis Upton and a crew of 104, 125 was lost with all hands.

Deleted: 125-was

Formatted: Highlight

389 Around 6 a.m. *Invincible* noted five British ships dangerously close to shore. *Eagle* was blown
390 onto its beam ends and jettisoned ten upper deck guns and cut down its mizzen mast to right the
391 ship. *Captain's* foretopmast was torn away and took its two topmen. *Lightning* found it was
392 drifting toward offshore breakers less than 200 m away. Captain Faulkner ordered *Windsor's*
393 guns jettisoned. He noted *Invincible* had lost all but its lower foremast and bowsprit. *Sunderland*
394 was swept by 'a very heavy large sea' that 'passed freely over us.' Barges lashed to the decks of
395 *Windsor* and *Invincible* were smashed and swept overboard. *Sunderland* cut down its main
396 topmast and threw guns overboard to right the ship. The wind snapped its 61 cm (24") diameter
397 mizzen mast as it drifted toward the offshore breakers. Anchors did not slow its drift so the
398 mainmast was cut down. *Sunderland* stopped close to the breakers and less than a kilometer from
399 shore (Fig. 3). The 74-gun *Terrible* also stopped its drift almost at the breakers. *Eagle's*
400 foretopmast was cut down to lessen the strain on the ship. It sailed southward narrowly missing
401 the breakers (Fig. 3). *Newark* regained control after cutting the anchor cable and heaving guns
402 overboard and barely cleared the line of breakers. Dawn revealed a signal flag had been raised
403 by the French fishing village of St. Esprit to give the crews of the British ships hope (Knox
404 Bristol Journal, November 12, 1757).

405 At Louisbourg the French fleet was pummeled by severe winds and waves. The 70-gun
406 French battleship *Dauphin Royale* fired a gun in distress when its anchor cables snapped under
407 the strain. *Dauphin Royale* collided with the 80-gun *Tonnant*, destroying its bowsprit, figurehead
408 and cutwater, and damaged *Tonnant's* rudder and poop deck. The two ships crossed
409 *l'Abenaquise's* anchor cables and the three entangled ships were heaved on shore at Royal
410 Battery (Fig. 2) along with 25 merchant ships, 50 schooners and 80 small vessels, many high and
411 dry and with many sailors drowned (McLennan, 1918).

412 At sea, by 10 a.m. the British fleet was dangerously close to the breakers off St. Esprit.
413 Many sailors were certain they were doomed (Knox Bristol Journal, November 12, 1757).
414 *Grafton* struck a rock but floated free and managed to set an anchor. *Windsor* and *Eagle* had
415 been able to sail south of the main British fleet off St. Esprit. *Eagle*'s Captain Palliser saw what
416 he judged to be *Nottingham* or *Tilbury* near shore, within the breakers, its bow facing shore with
417 its fore and mizzen masts gone. He also recorded that it was afloat and attempting to wear (turn)
418 but lost sight of it in heavy rain.

419 Waves tore down sections of the French Fortress Louisbourg's massive southeast facing
420 stone seawalls. Locals brought news of lakes 10 km inland being reached by the sea. Seawater
421 rose to flood the streets of the Town of Louisbourg, 'something never before seen' (Chevalier de
422 Johnstone, 1758). Eventually the beached French battleship *Tonnant* 'floated with the tide' as
423 the wind veered south and then west at 11 a.m.

424 At sea the British warship *Windsor* noted the wind turned to blow from the west at 11:30
425 a.m. but had strengthened. *Eagle* recorded that the squalls had lessened by noon. On the
426 *Sunderland* massive waves swept sailor George Lancey from the fore yard 24 m (80') above the
427 keel. By 3 p.m. waves at Louisbourg fell enough that *l'Inflexible* was able to send sailors to assist
428 other ships. French captains petitioned 74-year-old Admiral Dubois de la Motte to attack the
429 stricken British ships off their coast but his orders to defend Louisbourg had been met and he
430 kept his ships in port. James Johnstone, a Scot serving as a French officer, felt that five French
431 warships if they had ventured to sea could have captured the entire British fleet (Chevalier de
432 Johnstone, 1758). This sentiment was subsequently shared by Lady Anson, daughter of a
433 confidante of Lord Newcastle with whom Pitt had formed his coalition government, in an
434 October 31, 1757 letter to the First Lord of the Admiralty, her husband George Anson (Anson,

435 1757). On September 27th a boat arrived at Louisbourg from St. Esprit with news that the British
 436 warship *Tilbury* had wrecked there with over 120 lost. Four schooners with 160 French troops

TIME	BRITISH AT SEA	WINDS	DESCRIPTION	FRENCH IN PORT	WINDS	DESCRIPTION
7 p.m.	Sunderland	SSE	Very hard gales and hard squalls	Fleet	SE	Moored in Louisbourg Harbour
10 p.m.	Sunderland	SSE	Extreme hard gales			
10 p.m.	Windsor	SSW	Very heavy rain, intensifying strong gales Hard squalls			
12 a.m.	Invincible	SW	Strong gales; great sea, torrential rain			
2-4 a.m.	Invincible	SW	Excessive hard gale, hurricane of wind seas like mountains,	La Grave Battery	SE	Sea level rises 3.4 m (11')
2-4 a.m.	Sunderland	SSW	topsails and staysails blown to rags	Dauphin Royale		Dauphin Royale collides with Tonnant
2-4 a.m.	Devonshire	SE	Waves swept over the ship	Tonnant		Dauphin Royale and Tonnant driven across
2-4 a.m.	Lightning	SE	Waves overrun and destroy stern gallery	l'Abenaquise		l'Abenaquise anchor cable and the three entangled ships are thrown ashore at
2-4 a.m.	Cruiser	SE	Waves sweep over the ship			Royal Battery
			Guns jettisoned to avoid sinking	Royal Battery		
			Mizzen mast torn off ship by wind	Merchant ships		25 merchant ships thrown on shore
2-4 a.m.	Windsor	SSW	Severe squalls, heavy rain, great sea	Schooners		50 schooners thrown on shore
2-4 a.m.	fleet		Flooding by rain and waves	Small vessels		80 small vessels thrown on shore
	Grafton	SSE	Rudder torn off ship			
2-4 a.m.	Invincible	SW	Rudder torn off ship	SE facing sea wall		Waves tear down fortress stone seawalls
		SW	Hull planking sprung, hold flooding	Lakes in region		Lakes 10 km inland flooded by the sea
		SW	Gun deck brackets/bolts snapped	Louisbourg		Seawater floods the Town of Louisbourg requiring at least 4.4-6.4 m (14.4-21') surge
2-4 a.m.	Sunderland	SW	Foretopmast torn off ship			
	Invincible	SW	Driven onto its side by wind force			
		SW	Ten upper deck guns jettisoned			
		SW	Main mast snapped off which tears down foretopmast and mizzen mast			
		SW	Ship hauled onto its side by wreckage			
2-4 a.m. ?	Ferret	SE?	Ship swallowed by the sea with all hands			
4-6 a.m.	Invincible	SW	Near shore, sees five ships close to shore			
4-6 a.m.	Eagle	SE	Driven onto its side by wind force			
			Jettisons guns and cuts down mizzen			
4-6 a.m.	Captain	SE	Foretopmast torn off ship			
4-6 a.m.	Lightning	SE	Near offshore breakers 200 m away			
4-6 a.m.	Windsor	SSW	Jettisons guns to stay afloat			
4-6 a.m.	Sunderland	SSW	Swept by waves			
			Barge torn off the upper deck by waves			
4-6 a.m.	Windsor	SSW	Barge torn off the upper deck by waves			
	Sunderland	SSW	Driven onto its side by wind force			
		SW	Jettisons guns to stay afloat			
		SW	Mizzen mast torn off ship by wind			
		SW	Anchors at breakers 1 km from shore			
6-8 a.m.	Terrible	SE	Anchors at breakers			
	Newark	SE	Clears breakers			
10 a.m.	Grafton	SE	Strikes rock near St. Esprit			
	Eagle	SE	Notes <i>Tilbury</i> near shore at St. Esprit			
	<i>Tilbury</i>	SE	Aground at St. Esprit			
	fleet	SE	Most ships dangerously close to shore			
11 a.m.	Windsor	W	Winds shifted to westerlies			
12 p.m.	Eagle	W	Squalls lessening in strength			
3 p.m.	Invincible	W to NW	ship under jury rig drifting seaward	Inflexible	W	Waves reduced enough to assist other ships

437
 438 **Table 3.** Timeline of Louisbourg Storm (September 25)

439 Timeline of storm impacts on the British fleet at sea increasingly scattered by the storm and the
 440 French fleet moored in Louisbourg Harbour. Relative ship locations, south to north, are blue,
 441 orange, green and grey. British ships were relatively static (drifting, sailing under reefed sails or

442 at anchor) but *Invincible* sailed across storm winds to end up south of *Windsor* and *Sunderland*.
443 It is not known when *Ferret* sank but it had been sent ahead of the fleet prior to the storm to
444 undertake reconnaissance of the French fleet at Louisbourg.



445
446 **Figure 3.** Location of British ships estimated for 8 a.m. September 25 (sea day). The fleet sailed
447 in close formation until scattered by the hurricane south of Louisbourg (Fig. 1). Named ship
448 locations reflect best estimates of ship positions based on logbook references to sightings and
449 estimated distances and bearings to the coastline, known islands, Louisbourg, the breakers at St.
450 Esprit and other ships. The displacement of *Invincible* is based on the ship's logbook entries for
451 September 24 where the ship's position was fixed at noon with sextants to establish latitude with
452 the sun highest in the sky marking the start of the sea day. The entry 45°36'N 0°12'E, correcting
453 for 12' east longitude relative to Louisbourg Lighthouse as the zero meridian, corrects
454 *Invincible's* position to 45°36'N 59°45' W. *Invincible's* position on September 26 based on a
455 bearing of NBE (11.25 azimuth) and 4 miles (5 km) from 'Peddigrah,' a phonetic spelling for
456 'Petit de Grat' gives 45°23'51" N 60°58'55" W. *Sunderland* halted its drift one km from shore

Formatted: Font: 12 pt, Highlight

457 when the anchor finally held. The southwesterly winds encountered by *Invincible*, *Sunderland*
 458 and *Windsor* reflect the southernmost vessels sailing southwest into a northeast tracking storm. A
 459 displacement of 97.25 km toward 257.43 azimuth when the bearing was taken at 11 a.m. when
 460 the wind shifted to westerly (September 25; sea day), giving an average speed of 2.07 ~~km h⁻¹~~ km h⁻¹
 461 over 47 hours. Plotting the hourly displacement allowed the position of the ship to be estimated
 462 for noon, September 25, at the height of the storm at 3 a.m. and when the ship was dismantled at 6
 463 a.m., and at 8 a.m. when the positions of multiple ships could be estimated when *Windsor*,
 464 *Sunderland* and *Invincible* were under southwesterly winds while the rest of the British ships
 465 were still recording south-southeasterly winds and when the British ship positions had been
 466 stabilized by anchoring or limiting their rate of drift. Logbook records: *Orford* 6 km from the
 467 coastline running northwest to north; *Windsor* 3 km from the breakers; *Terrible* 1.3 km from
 468 breakers and 3 km from the land to the west-northwest; *Lightning* 200 m off the breakers before
 469 halting its drift; *Sunderland* 1 km from shore after sailing SE, across SW winds; *Tilbury*
 470 shoreward of the breakers; *Newark* near breakers with *Northumberland* and *Kingston* and
 471 *Windsor*; *Eagle* south of breakers until 11:30 when the breakers were 3 km to their lee. Image ©
 472 Google Earth Pro 7.3.6.9345 (2022) Cape Breton, Nova Scotia Canada. Image date 12/13/2015
 473 45°33'51.38" N 60°13'56.57" W Eye alt 132.12 km TerraMetrics © 2023 MaxarTechnologies ©
 474 2023.

475 were unable to counter the heavy seas so they marched to the site across land flooded by the
 476 torrential rain. Mi'kmaq warriors gained the wreck first but informed the shipwrecked British
 477 they would not be harmed since the storm had brought them to their lands (Moreau St. Mery in
 478 McLennan, 1918).

479 6.0 Deriving Storm Metrics

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Highlight

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt, Highlight

Formatted: Highlight

Formatted: Font: 12 pt

480 Storm intensity is reflected in key metrics including wind speed and direction, wave
481 height and surge which is driven by a rise in sea level due to atmospheric pressure and sustained
482 storm winds and is proportional to a cyclone's intensity, translation rate and the bathymetric
483 gradient of the continental shelf.

484 6.1 Estimating Storm Wind Speed

485 The wind speed required to break *Invincible's* main mast, and other ships' mizzen masts
486 and topmasts is estimated based on the engineering model of Viro et al. (2016) who determined
487 the critical wind force needed to break trees of average integrity is 151 kph km h^{-1} irrespective of
488 species with a +9% factor for large diameter trees. This is relevant since masts in 1757 were
489 made from single trees. 165 kph km h^{-1} assumes structural defects due to longer tree life offset
490 the structural advantage of size, yet masts were chosen for their lack of defects. Fir and pine trees
491 of superior structural integrity were selectively harvested for Royal Navy masts into the 1770's
492 from North America, Great Britain and the Baltic (Lavery, 1984). Masts were also not free-
493 standing (like trees) but reinforced by rigging to effectively transfer wind energy from the sails
494 to the hull. *Invincible's* masts were secured by sixteen 5 cm (2") hemp shrouds per side, each
495 tensioned with paired deadeye blocks, the lower block in an iron band bolted to the ship's frame.
496 Its 1 m (38") diameter lower mainmast stepped against the ship's keelson rose 35.7 m (117')
497 through two decks. Above it stood a 21.3 m (70') 51 cm (20") diameter topmast and above that
498 the 10.7 m (35') 28 cm (11") diameter topgallant mast (Lavery, 1984, 1988). To achieve the
499 critical wind speed of 165 kph km h^{-1} , taken as a minimum due to the factors noted, *Invincible's*
500 motion must be considered.

501 *Invincible* sailed SW under SE winds, but gradually encountered SW winds. *Sunderland*
502 and *Windsor* sailed south across SSW winds while most ships of the British fleet to their north

Commented [JD11]: change kph to km h^{-1}

Commented [JD12]: This is not critical but helps understand the displacement of the ship between September 24-26



503 near St. Esprit faced SSE winds. *Invincible* was among the southernmost ships (Fig. 1). It sailed
 504 SW½W (230°) against EbS (101°) winds on September 24. During the storm its displacement
 505 was 98 km toward 256.7° (22.5 km S; 96 km W). 6 km SE (135°) of Chedabucto Bay it faced W
 506 (270°) winds.

507 Ephemeral squalls of 40-60 **kph km h⁻¹** added to sustained winds of **165 km h⁻¹** kph
 508 suggests peak winds might have reached 205-225 **kph km h⁻¹** around 6 a.m. when *Invincible*'s
 509 mast broke. *Sunderland*'s foretopmast broke at 7 a.m. and the mizzen mast broke at 9:30 a.m.
 510 While it is an imperfect solution, it does not consider the inherently superior structural integrity
 511 of masts plus their reinforcement by rigging, which requires only an additional strength factor to
 512 withstand an additional sustained 13 **kph km h⁻¹** to meet major hurricane threshold (178 **kph km**
 513 **h⁻¹**) without considering squalls.

Commented [JD13]: change kph to kmh⁻¹

Commented [JD14]: change kph to kmh⁻¹

Commented [JD15]: corrected to kmh⁻¹



Deleted:

514 **Figure 4.** Hurricane wind evolution with time. The time sequence shows the arrival of southeast
 515 winds (Beaufort Scale) intensifying to hurricane winds (118 **kph km h⁻¹**) peaking to sustained
 516 winds (Beaufort Scale) intensifying to hurricane winds (118 **kph km h⁻¹**) peaking to sustained

518 165 km h^{-1} critical wind force with increasing squall frequency, followed by a rapid decline
 519 to gale force westerlies. The horizontal axis is divided into days (noon) and 2-hour intervals. The
 520 vertical scale is wind speed in km h^{-1} . A best fit curve [1] is typical of windspeeds as a
 521 hurricane passes a fixed point. A best fit curve for squall frequency [2] in ships' logs adds
 522 ephemeral wind speed to sustained winds. 165 km h^{-1} is considered the minimum critical
 523 wind force considering the superior materials integrity of masts and their reinforcement with
 524 rigging. Peak winds lasted 9 hours while hurricane force winds impacting the fleet lasted 15
 525 hours. Wind directions represent, north to south, winds affecting: French ships at Louisbourg,
 526 British ships near St. Esprit, *Windsor* and *Sunderland* south of St. Esprit, and *Invincible* closest
 527 to the eye (Fig 1). Southernmost (blue) through southern (orange), off St. Esprit (green) and
 528 Louisbourg (grey) show the general distribution of ships (*see* Table 3). *Invincible* sailed past
 529 *Windsor* and *Sunderland* during the storm.





Figure 5. Estimate of wind force at *Invincible* under threshold winds. *Invincible*, maintaining its bearing of SW½W of September 24 sailed into winds that progressively became SW (at the ship)



534 as the hurricane tracked northeast. Square-rigged ships cannot sail closer than 35° degrees into
535 the wind. This reduced the wind speed acting on the masts by a minor amount, suggesting that
536 squalls whose frequency corresponds to the frequency of ship damage (Fig. 4) were needed to
537 overcome the reinforcing factors of superior mast structural integrity and rigging to achieve
538 critical force. Not to scale.

539 Anticlockwise wind vectors at ship locations are tangential to concentric cyclonic wind bands.
540 Normal lines drawn to these vectors converge to identify the location of the eye. Interestingly
541 they lack the asymmetry diagnostic of extratropical cyclone wind fields (Fig. 8). This process,
542 repeated to plot the eye location on September 26, 1757, indicates the storm crossed Cape Breton
543 and entered the Gulf of St. Lawrence. Even if the wind field began to collapse, the location of the
544 storm center suggests the system may have slowed while passing over Cape Breton Island.

545 6.2 Estimating Storm Wave Height

546 *Sunderland's* and *Devonshire's* upper decks were submerged after waves broke over the
547 forecastle. The 12.2 m (40') distance from the keel to the upper deck plus an estimated 3-6 m
548 (15-20') to break over the forecastle and tear away ship's boats lashed to the deck requires a
549 wave height of about 18 m (60') (Lavery 1983). *Lightning's* stern gallery 20-15 m (40-50')
550 above the keel was destroyed by waves striking the ship from astern, also requiring waves of
551 about 12.2 m (60'). A sailor swept out of *Sunderland's* fore yard by a wave necessitates a wave
552 of about 25-30 m (80-90'). While carrying considerable uncertainty, these examples provide
553 estimates of significant and maximum wave heights. Waves sufficiently large to tear down stone
554 seawall rampart of Fortress Louisbourg are consistent with these estimates, as are waves capable
555 of reaching inland lakes. Descriptions of the sea state in Louisbourg Harbour by French naval
556 officers resulting in extensive damage to ships and boats suggests waves much larger than any

Deleted: 40-50'

558 recorded in modern times even though wave energy from the southeast would have been partly
559 attenuated by shoals (Fig. 2).

560 On September 26-28, 1818, the American frigate *USS Macedonian* met a hurricane off
561 Bermuda (35°N 53°W) and suffered damage nearly identical to *HMS Invincible* in 1757 from
562 waves of 12 m (40') (Saegesser, 1970). The dates appear to coincide with Chenowith's (2006)
563 'Final Storm Number 253' listed as a hurricane in Chenowith's Table IV. Damage to the ship
564 closely parallels that described for the 1757 hurricane except that line of battle ships had a much
565 heavier construction than a frigate. Saegesser (1970) provides a detailed account from the ship's
566 log and ancillary damage reports, and notes that in the same storm the Dutch brig *De Hoope* lost
567 all topmasts and spars, the brig *Ann* from Nova Scotia was abandoned at sea, the brig *Mary* from
568 Bristol was overturned, the ship *Catherine Dawes* from Philadelphia sank and a Baltimore
569 schooner and a Nantucket whaler were both dismasted. *Invincible's* substantially more robust
570 build than the frigate *Macedonian* implies larger, more powerful waves caused its damage.

571 6.3 Estimating Surge Height

572 6.3.1 Surge at Louisbourg Harbour

573 A Parks Canada coastal erosion study at Fortress Louisbourg National Historic Site
574 revealed iron mooring rings set in the remains of a seawall. Modern high tide compared to these
575 rings established historical high tide 0.90 m (3') of sea level rise since 1757 (Duggan, 2010). La
576 Grave Battery (Fig. 2) is 2.0 m (6.6') above sea level (asl; Google Earth mid-tide datum), so sea
577 level rise plus flooding to sentries' knees (0.5 m) yields a 3.4 m (11') mid-storm surge. Historic
578 buildings along the waterfront (Fig. 2; 45°53'33.57" N 59°59'07.89" W) are 5 m (16.4') asl
579 while the first street, Rue Royale, is 7 m (22.9') asl. Seawater flooding the town streets at the
580 lowest levels and adjusted for sea level rise indicates 5.9 m (19.4') to 7.9 m (21.4') of surge.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Italic, Highlight

Formatted: Font: Italic, Highlight

Formatted: Font: Italic, Highlight


Formatted: Font: Italic, Highlight

581 *Tonnant* ‘floated with the tide’ when the wind veered south at 11 a.m. on September 26 (*Fleur de*
582 *Lys* log in McLennan, 1918). Louisbourg’s 12-hour tidal cycle and assuming low tide around 10
583 a.m. gives a high tide at 4 a.m. coinciding with storm landfall and creating a storm tide (Fig. 4).
584 Backing out the 1.5 m (5’) tidal range gives a 4.4-6.4 m (14.4-21’) peak surge, consistent with
585 the earlier surge of 3.4 m (11’) at La Grave.

586 6.3.2 Surge at St. Esprit (*Tilbury Wreck*)

587 *HMS Tilbury* was a 58-gun square-rigged warship lost on the coast in the storm. *Eagle*’s
588 captain saw either *Tilbury* or *Nottingham* shoreward of the breakers near St. Esprit, 45 km south
589 of Louisbourg. It was deduced to have been *Tilbury* since *Nottingham* survived the storm with a
590 different array of masts than seen on this ship. ‘Wreck’ appears on a 1776 chart (Fig. 6). Storm
591 (2002) used Zinck’s (1975) image of an 18th Century 6-pounder British naval gun at ‘Tilbury
592 Rocks’ to view *Tilbury*’s wreckage in 4 m (15’) from a boat in 1969.

Commented [JD16]: Italics




593

594 **Figure 6.** Excerpt from a historic chart of Cape Breton Island showing the general St. Esprit
595 study area and *HMS Tilbury* wreck site, from Mowat (1776), depicted in Fig. 7a, b. The faint
596 dotted line right of Barnsley Lake, named for *Tilbury's* captain, marks a parish boundary.


Commented [JD17]: Italics

597 The historic navigation chart (Fig. 6) showed parish boundaries marked by fieldstone
598 walls of historic St. Esprit (Fig. 7a, b) which helped identify the line of offshore breakers
599 described in British naval logs. A draft hydrographic chart (Hanson, 1954) was digitized and
600 gridded with missing data interpolated. Paired depths and locations were entered in a spreadsheet
601 and a grid-plot of local bathymetry supported a marine proton magnetometer survey of Tilbury
602 Reef isobaths following best practices for submerged archaeological sites (Cornwall Council
603 Report 2010-R012). Dipole targets investigated by divers led to locating a mid-18th Century 6-

604 pounder British naval gun *in situ* in 3 m (10') which was 2.1 m (7') in 1757, near the site of the
605 6-pounder on shore, both interpreted to be from *Tilbury's* forecastle. In 1757 *Tilbury* was
606 observed at the time as 'bow in' near shore, landward of the breakers and 'attempting to wear'
607 (turn). It was in water sufficiently deep for its 18' displacement as it was, at the time, afloat and
608 under sail. Adding in the hydrographic survey datum offset of 0.6 m (2') between lowest low tide
609 at St. Esprit and the Google Earth WGS84 (World Geodetic Standard 1984) mid-tide datum for
610 Louisbourg suggests a minimum 4.0 m (13') surge at St. Esprit. Post-storm relaxation flow
611 stranded the *Tilbury* (Fig. 7b) allowing native warriors to reach it.



612



613

614 **Figure 7a.** Location of the *Tilbury* shipwreck. Inset map X – X' (45°38'31.21" N 60°27'41.99"

615 W to 45°38'31.61" N 60°26'05.28" W) corresponds to Fig. 6b [7b]. Dashed line is bedrock reef

616 (breakers). Image © Google Earth Pro 7.3.6.9345 (2022) St. Esprit, Nova Scotia Canada.

617 45°38'31.54" N 60°27'37.76" W Eye alt 4.50 km TerraMetrics © 2023 MaxarTechnologies ©

618 2023.

619 **Figure 7b.** Bathymetry of *Tilbury* wreck site at lowest low water adjusted for 1757 relative sea


620 level (solid line) and minimum surge (dashed line) needed to float *Tilbury*. Coastal retreat of 27

621 m (90°) calculated from historic sea level gives the 1757 shoreline. Topographic and bathymetric

622 data were kept in Imperial units (feet) for comparison to *Tilbury*'s displacement. X and X' of this

623 block diagram correspond to the same GPS positions on the areal chart in Fig. 7a.

Commented [JD18]: correct figure references



624
 625 **Figure 8. Eye location and estimated translation speed.** Plots of wind vectors on September
 626 25 (8 a.m.) at: (1) *Invincible*, (2) *Windsor* and *Sunderland*, (3) *Newark* and most of the British
 627 fleet, and (4) French ships at Louisbourg Harbour. Normal lines (dashed blue lines) taken to
 628 wind vectors cluster at the eye.

629 **7.0 Modern Analogs**

630 On September 29, 2003, Hurricane Juan struck Nova Scotia with peak winds of 165 **kph**
 631 **km h⁻¹** (Category 2), a significant wave height of 10 m (32'), a maximum wave height of 19.9 m
 632 (65') and a surge at landfall near Halifax of 1.5 m (4.9') (Lixion, 2003). On January 20-22, 2000,
 633 an extratropical meteorological 'superbomb' that developed off Cape Hatteras struck Nova
 634 Scotia with peak winds of 25-30 **m/s m s⁻¹** (90-108 **kph km h⁻¹**), a significant wave height of 12
 635 m (39'), a peak wave height of 19 m (62') to 23 m (77') at drilling rigs near Sable Island (JD
 636 pers. obs.) and a 1.4 m (4.6') surge at landfall near St. Esprit (Lalbeharry et al., 2009). Both
 637 cyclones produced similar sea states and surge which can be compared to the Louisbourg Storm.

638 On September 24, 2022, Category 3 Hurricane Fiona began extratropical transition as it crossed
 639 the Scotian shelf. A cold trough over Nova Scotia directed its landfall to the Canso Peninsula.
 640 Winds of 140 km h^{-1} kph in Nova Scotia reached 177 kph km h^{-1} in Newfoundland and Labrador.
 641 Significant and maximum wave heights were 17 m (56') and 30 m (98') and surge reached 2.4 m
 642 (8').

643 In 1969 Category 5 Hurricane Camille generated a 7.3 m (24') storm tide from 1.8-3.0 m
 644 (6-10') surge (U.S. Department of Commerce Environmental Science Services Administration
 645 1969) while Category 5 Katrina in 2005 produced a storm tide of 8.2 m (27') (Knabb et al.,
 646 2023). Hurricane Laura (Category 4) in 2020 had a peak 5.2 m (17.2') surge (Pasch et al., 2021)
 647 and a 2.7-4.0 m (9-13') spanning 130 km from Beaumont to Lake Arthur, Texas. In 2018
 648 Hurricane Dorian (Cat 5) slowed to 2 kph km h^{-1} over the Bahamas creating an 8.5 m (28') surge
 649 (Avila et al., 2020). Surge from these major hurricanes cannot be readily compared to storm
 650 strikes in Nova Scotia due to different coastal bathymetry but they allow a general comparative
 651 benchmark.

652 Hurricane Juan's translation speed before landfall was 1-5 m/s m s^{-1} (4-18 kph km h^{-1}). If
 653 the Louisbourg Storm slowed slightly as it approached Nova Scotia it may have enhanced surge
 654 height, similar to Dorian's impact on the Bahamas as it slowed which may explain the
 655 exceptional surge height at Louisbourg. The key metrics of wind speed, wave height and surge
 656 are summarized in Table 4.

Storm	Year	Date	Peak Wind (km h^{-1})	Significant Wave Height (m)	Peak Wave Height (m)	Surge (m)
Louisbourg	1757	25-Sep	205 - 225	12+	25-30	4.4 - 6.4
Unnamed	2000	22-Jan	90 - 108	12	19	1.4
Juan	2003	27-Sep	160 - 165	10	20	1.5
Fiona	2022	24-Sep	155 - 179	17	30	2.4

Commented [JD19]: changed table

Deleted:

Storm	Year	Date
Louisbourg	1757	25-Sep
Unnamed	2000	22-Jan
Juan	2003	27-Sep
Fiona	2022	24-Sep

657

659 **Table 4. Louisbourg Storm Comparison to Modern Nova Scotia Landfalling Storms.** The
660 Louisbourg Storm, a winter extratropical storm in 2000, Juan (Category 2 hurricane at landfall),
661 and Fiona, an extratropical cyclone that transitioned from a Category 3 hurricane over the
662 Scotian Shelf crossed the same coastal bathymetry with similar translation rates to strike Nova
663 Scotia. Sustained winds for the Louisbourg Storm exceeded 165 kph km h^{-1} based on the critical
664 force needed to break main and mizzen masts and break away and carry off topmasts and may
665 have reached 225 kph km h^{-1} with squalls.

666 **8.0 Discussion**

667 Metrics derived from historical data captured during the Louisbourg Storm of 1757
668 indicate its intensity surpassed any modern (post-1851) Atlantic cyclones striking the same
669 region. Historical records show the Louisbourg Storm originated in the tropics to pass Florida,
670 the Carolinas and New England to strike Nova Scotia on September 25, 1757. It developed at
671 the height of hurricane season under an optimal NAO (strongly negative) index and ENSO
672 conditions (La Nina) for Atlantic hurricanes to form and track up the Atlantic coast of North
673 America into the northern midlatitudes. The NAO index tends to decrease as the season
674 progresses (Hart and Evans, 2001) and may have helped the hurricane remain over the Gulf
675 Stream and intensify into higher latitudes. Its devastating impact on the British and French fleets
676 and coastal infrastructure was due to an unusually violent release of energy over coastal waters.
677 A UK and European heat wave in Europe in 1757, extreme even by modern standards, shows
678 seasonal temperature variability could contribute to warmer SSTs and fuel tropical cyclones in
679 the LIA. A strong correlation between SST and tropical cyclone frequency (Vecchi and Knutson,
680 2008) suggests that the LIA's cooler SSTs could see fewer storms per year. Mean-annual
681 temperature data limited by temporal resolution limitations likely mask peak temperatures that

682 must have existed over smaller areas for shorter periods since historical records (e.g., Chenowith
683 2006) clearly show tropical cyclones developed even during the coldest part of the LIA. A
684 multidecadal warming-cooling trend in temperate North America peaking in the mid-1700's
685 (Trouet et al. 2013) shows shorter-cycle warming within a cooler mean LIA. It suggests that the
686 peak latitudes reached by midlatitude hurricane patterns should be compared to multi-decadal
687 temperature cycles.

688 The large number of British warships scattered along Cape Breton's coast by the
689 Louisbourg Storm provided a spatial resolution of wind vectors not normally available in storm
690 reconstructions. It was partly facilitated by ships sailing across storm winds to avoid being
691 driven ashore. The proximity of many British ships to shore (Fig. 3) and the severe surge and
692 wave action at Louisbourg led many contemporary naval authorities of both nations to fear the
693 catastrophic loss of the British and French fleets and almost 21 000 sailors. Only the reversal of
694 wind direction at the last minute as the eye of the storm passed prevented a disaster.

695 Wind speed is the key metric used in the Saffir Simpson scale to characterize the intensity
696 of modern cyclones. Engineering models are a standard method of determining the force
697 required to trigger structural failure in materials. Trees lacking defects that negate size advantage
698 were preferentially selected for masts and so likely required higher wind speeds for structural
699 failure. Rigging not only reinforced masts but redirected wind energy to the hull. Both factors
700 imply that the wind speed estimate of 165 kph km h^{-1} is an underestimate while the 178 kph km h^{-1}
701 (Cat 3) major hurricane threshold requires that increased strength factor to only be equivalent to
702 13 kph km h^{-1} . Extreme winds are reflected in topmasts (along with shrouds and stays) not only
703 being torn off two British ships but being carried off (with sailors) instead of falling to the deck.
704 British ship positions were triangulated against known coastal landmarks, including the offshore

705 breakers at St. Esprit, and each other. This provided greater accuracy in wind vectors for the
706 period 8-10 a.m. Superimposing *Invincible's* location and the wind vectors that identify the eye
707 location at the height of the storm suggests severe damage was a consequence of proximity to the
708 eye which is the location of a cyclone's strongest winds (Figs. 1, 4, 8). Peak damage and squalls
709 above hurricane winds lasted 9 hours and hurricane force winds noted by the British ships lasted
710 more than 15 hours as the center of the storm passed the coast (Fig. 4). In comparison, Hurricane
711 Juan crossed Nova Scotia in only 3 hours while Fiona crossed the province in under 6 hours (Fig.
712 8). The Louisbourg Storm may have slowed approaching Nova Scotia. Rough estimates of the
713 storm position off North Carolina, New England and Nova Scotia suggest a translation speed of
714 33 kph km h^{-1} between the Carolinas and New England in 24 hours, and 19 kph km h^{-1} based on
715 42 hours to cross 800 km to land at Chedabucto Bay (Fig. 8) by 8 a.m. on September 25,
716 crossing the remaining 113 km in 4 hours yielding an estimate of 28 kph km h^{-1} . There is
717 significant uncertainty associated with these estimates, but if the hurricane slowed between New
718 England and Nova Scotia, its location over the Labrador Current while encountering prevailing
719 westerlies (Table 2) may have created a strong temperature gradient known to trigger
720 extratropical transition (Hart and Evans 2001) where stronger gradients drive more rapid
721 intensification and greater destructive power (e.g., Day and Hodges, 2018, Studholme et al.,
722 2022, Cheung and Chu, 2023). It can therefore be argued that while modern SST warming
723 driving steeper temperature gradients will result in more powerful storms, a similar increase in
724 baroclinic instability from steeper temperature gradients driven by colder continental autumn
725 circulation during the LIA interacted with an intensifying tropical cyclone. The hurricane was
726 fueled by SSTs that peak at their most northern latitudes at the height of Atlantic hurricane
727 season in late September and early October, consistent with the extratropical climatology of Hart

Commented [JD20]: extremely long sentence

728 and Evans (2001) and records of prevailing westerlies (Table 2) which were recorded as
729 extremely cold following the storm. Wind plots also show that the southernmost ships of the
730 British fleet faced southwest winds from the lower right quadrant of the hurricane. British ships
731 to the northeast near St. Esprit faced southeast winds. The French fleet in Louisbourg Harbour
732 also faced southeast winds and an anomalously high storm surge which allowed massive waves
733 to drive ships on shore while the surrounding region was flooded by torrential rains, all
734 consistent with the front right quadrant of the hurricane where the most severe impacts are felt.
735 There was no suggestion that the air of the storm was cold, but westerlies following the storm
736 were described at Fort Cumberland as very cold and dry.

737 Modern analogs show strong similarities in significant and maximum wave height, but
738 interpreted wind speeds for the Louisbourg storm are greater than those of Category 2 hurricane
739 Juan, a winter extratropical ‘superbomb’ in 2000, and the extratropical cyclone Fiona in 2022.
740 Surge measured at three locations is consistent with the scale of surge from major hurricanes in
741 the Gulf of Mexico and Caribbean. The 1757 surge greatly exceeds that of modern analogs that
742 crossed the same bathymetry with similar translation speeds. This consistent basis of comparison
743 of surge height, closely linked to storm intensity, shows the Louisbourg Storm had an intensity
744 far beyond a Category 2 system and was equal to a major hurricane. Surge calculated
745 independently for the lowest streets of the historic town of Louisbourg, Battery de la Grave and
746 the *Tilbury* wreck at St. Esprit were also consistent. Even accommodating the tidal range at
747 Louisbourg, the French battleship *Le Tonnant* drawing 25’ being beached requires an exceptional
748 surge. Unlike the modern analogs, storm surge at Louisbourg was one hundred kms from landfall
749 (Fig. 8).

750 The climatology of tropical cyclones on North America's eastern seaboard renders the
751 simple attribution of 'tropical' vs. 'extratropical' problematic. It is unlikely that a fully tropical
752 system with wind speeds equal to a Category 4 hurricane struck Nova Scotia. Hart and Evans'
753 (2001) climatology for North Atlantic extratropical transition of tropical cyclones showed that
754 expansion of baroclinic conditions known to trigger transition as cooling autumn continental
755 temperatures expanding under prevailing westerlies encounter north-trending tropical cyclones
756 that tend to reach the highest latitudes by October when SSTs peak. Cheung and Chu (2023)
757 modeled different concentrations of **CO₂** **CO₂** as a forcing mechanism behind future global
758 warming. Their model outputs showed that more destructive extratropical cyclones originating in
759 the tropics as tropical cyclones become more frequent in response to warming. The key factors in
760 storm destructive energy is increased wind speed and the expansion of the wind field during
761 extratropical transition. This supports the climatology of Hart and Evans (2001) who described
762 the collapse of the symmetric tropical wind field into an asymmetric extratropical storm during
763 transition, and the tendency for tropical cyclones formed below 20° **degrees** north latitude to
764 maintain their tropical integrity into higher latitudes where they have a higher probability of
765 post-transition intensification. The National Hurricane Center (NHC) uses sea surface
766 temperatures plus storm asymmetry in satellite images to gauge the degree of transition. Hart and
767 Evans (2001) also found that 'the NHC declaration (of extratropical transition) typically occurs
768 early in the 1 to 2-day period ... when the storm is just beginning to lose its tropical
769 characteristics.' This is not easy to assess for the Louisbourg Storm whose energy release may
770 have occurred over a much shorter period. The eye symmetry at landfall on September 25 is
771 based on the convergence of normal lines to vectors at ship locations (Fig. 8) suggesting it may
772 have had largely tropical characteristics at landfall. It leads to the question at what point was it

Commented [JD21]: subscript

773 'tropical' (hurricane) vs. 'extratropical' given the NHC's 1 to 2-day range. The storm's unusually
774 large size is indicated by its winds first being recorded on September 22 by both the British and
775 French fleets at Cape Breton on the same day it struck the British frigate *Winchelsea* off North
776 Carolina, 1350 km to the southwest. This may have enabled it to continue to draw tropical
777 energy from the Gulf Stream as it neared the Nova Scotia coastline. Hart and Evans's (2001)
778 extratropical climatology shows that in some cases tropical cyclones can continue to intensify
779 north of strongly baroclinic conditions that trigger transition, resulting in an explosive release of
780 energy and post-transition intensification. Their analysis of past Atlantic hurricanes shows that
781 the region most conducive to post-transition intensification in the North Atlantic basin lies
782 immediately south of Cape Breton, Nova Scotia, which covers the track of the Louisbourg Storm
783 in 1757.

784 Multidecadal climate trends for temperate North America show eighteenth century
785 warming peaking mid century followed by cooling within a cooler mean temperature associated
786 with the LIA (Trouet et al., 2013). This supports the early argument by Mann (2002) who argued
787 that the LIA was a period of natural climate variability which is indicated by relatively warmer
788 summers offset by colder winters to provide cooler mean and multidecadal LIA temperature
789 trends. Tropical cyclones continued to transfer equatorial heat northward into the midlatitudes
790 where they likely encountered colder LIA continental temperatures earlier in hurricane season,
791 driving a sharper temperature contrast and greater baroclinic instability resulting in a more
792 catastrophic energy release during extratropical transition. Oliva et al. (2017) noted the
793 importance of various proxies to study historical Atlantic hurricanes given the importance of
794 understanding their frequency and intensity as a benchmark against future storms. One area on
795 the eastern seaboard of North America showing a notable data gap is Nova Scotia (Oliva et al.,

796 2017). Not only has the population of the northeastern United States and Atlantic Canada grown
797 since 1757, but coastal waters experienced massive shipping growth between North America and
798 Europe. In addition, sea level rise since 1757 and projected rise increases storm surge risk to
799 coastlines under more powerful storms. Hart and Evans (2001) identified this region as having
800 the highest probability of post-transition intensification. Heightened temperature gradients into
801 fall driven by warmer SSTs would not only fuel more powerful tropical cyclones reaching higher
802 latitudes, but more intense extratropical cyclones as well.

803 **9.0 Conclusions**

804 In 1757 a cold air mass met a hurricane that tracked north along the Gulf Stream from the
805 coast of Florida. The resulting explosive release of energy was likely due to extratropical
806 transition driven by the heightened temperature gradient between colder continental and tropical
807 maritime circulation during the LIA, giving the Louisbourg Storm its destructive power. This
808 increase in energy requires only an incremental change in the accepted climatology of Atlantic
809 cyclone extratropical transition. The duration of hurricane force winds (15 hours) over the coast
810 may have been enhanced by the storm's large diameter, possibly a result of transition. The storm
811 drove an unusually high surge at high tide. Warmer SSTs under anthropogenic forcing creating
812 steeper autumn coastal temperature gradients could fuel future midlatitude tropical and
813 extratropical cyclones of increasing destructive power.

814 **Acknowledgements**

815 The authors would like to thank William Pretel and Antoine LaChance for constructive review
816 comments on the manuscript. Research assistance was provided by Cambria Huff (Dalhousie),
817 John Allison (UK), the National Archives (UK) and the Public Archives of Nova Scotia. All
818 figures were drafted by JD. Tony Sampson, commercial diver, offshore survival instructor and

819 owner/operator of Salty Dog Sea Tours, and Steve Jennex, Dave Murphy, Steve Dugas and Dana
820 Sheppard of Zodiac Divers coordinated marine operations and underwater exploration.

821 **Funding**

822 The authors declare that no funds, grants, or other supports were received during the preparation
823 of this manuscript and that they have no financial or proprietary interests in any material
824 discussed in this article.

825 **References**

826 Anson, Lady. Letter of October 31, 1757 from Lady Anson to George Anson, First Lord of the
827 Admiralty, British Museum Collections, London UK, Add MSS 35,376 f. 145, 1757.

828 ADM 1/481 Letters from Commanders in Chief North America 1755-1760. (Charles Holmes)
829 The State and Condition of His Majesty's Ships and Sloops under my Command at New
830 York between 3rd of May 1757 and 9th following, The National Archives, UK, 1757.
831 <https://discovery.nationalarchives.gov.uk/details/r/C4771564> [records copied in 2001]

832 ADM 1/481 Letters from Commanders in Chief North America 1755-1760. (Frances Holbourne)
833 Newark at sea 28 September, The National Archives, UK, 1757.

834 <https://discovery.nationalarchives.gov.uk/details/r/C4771564> [records copied in 2001]

835 [Letter to the Admiralty outlining his squadron's inability to continue operations and the
836 need to refit]

837 ADM 1/481 Letters from Commanders in Chief North America 1755-1760. (Frances Holbourne)
838 Newark at sea 28 September, The National Archives, UK, 1757

839 <https://discovery.nationalarchives.gov.uk/details/r/C4771564> [records copied in 2001]

840 [list of damage to ships sustained in the gale]

841 ADM 1/481 Letters from Commanders in Chief North America 1755-1760 (Frances Holbourne)
842 Newark at Sea 30 September, The National Archives, UK, 1757.
843 <https://discovery.nationalarchives.gov.uk/details/r/C4771564> [records copied in 2001]

844 ADM 1/481 Letters from Commanders in Chief North America 1755-1760. Newark at Halifax
845 14 October, The National Archives, UK, 1757.
846 <https://discovery.nationalarchives.gov.uk/details/r/C4771564> [records copied in 2001] [A
847 letter from Frances Holbourne to the Admiralty outlining the state of the squadron and
848 the enemy's ships at Louisbourg]

849 ADM 1/1488 Captain's Letters 1757 (Bently, Jonathon). An account of the damages received on
850 board His Majesty's Ship Invincible in the hurricane on the 25th September, The
851 National Archives, UK, 1757
852 <https://discovery.nationalarchives.gov.uk/details/r/C4772571> [records copied in 2001]

853 ADM 1/2294 Captain's Letters 1757 (Palliser, Hugh). Sunday 25th September 1757 at 2 a.m. An
854 account of the Eagle's situation and of the damages she received in the late gale of wind.
855 The National Archives, UK, 1757.
856 <https://discovery.nationalarchives.gov.uk/details/r/C4773376> [records copied in 2001]

857 ADM 1/2294 Captain's Letters 1757 (Palliser, Hugh). Eagle at sea 30 September, 1757. Account
858 of the Condition of His Majesty's Ship Eagle, The National Archives, UK, 1757.
859 <https://discovery.nationalarchives.gov.uk/details/r/C4771564> [records copied in 2001]

860 ADM 8/31 Admiralty List Books 1756-1757 Halifax Station, The National Archives, UK, 1757.
861 <https://discovery.nationalarchives.gov.uk/details/r/C537622> [records copied in 2001]

862 ADM 8/32 Admiralty List Books 1757-1758 Halifax Station, The National Archives, UK, 1757.
863 <https://discovery.nationalarchives.gov.uk/details/r/C537623> [records copied in 2001]

864 ADM 51/409 Captain's Log HMS Grafton (1755 Feb 7–1764 Jun 24), The National Archives,
865 UK, 1757. <https://discovery.nationalarchives.gov.uk/details/r/C4460516> [records copied
866 in 2001]

867 ADM 51/471 Captain's Log HMS Invincible (1756 Aug 7–1758 Mar 6), The National Archives,
868 UK, 1757. <https://discovery.nationalarchives.gov.uk/details/r/C4460776> [records copied
869 in 2001]

870 ADM 51/633 Captain's Log HMS Newark (1755 Jul 31–1760 Apr 1), The National Archives,
871 UK, 1757. <https://discovery.nationalarchives.gov.uk/details/r/C4462295> [records copied
872 in 2001]

873 ADM 51/921 Captain's Log HMS Sunderland (1756 Nov 15–1759 Feb 23), The National
874 Archives, UK, 1757. <https://discovery.nationalarchives.gov.uk/details/r/C4462272>
875 [records copied in 2001]

876 ADM 51/1075 Captain's Log HMS Windsor (1755 Jun 12–1759 May 20), The National
877 Archives, UK, 1757. <https://discovery.nationalarchives.gov.uk/details/r/C4462763>
878 [records copied in 2001]

879 ADM 52/578 Master's Log HMS Eagle (1757 Apr 28–1759 Mar 3), The National Archives, UK,
880 1757. <https://discovery.nationalarchives.gov.uk/details/r/C2531251> [records copied in
881 2001]

882 ADM 52/819 Master's Log HMS Captain 1756 May 21–1760 Feb 21), The National Archives,
883 UK, 1757. <https://discovery.nationalarchives.gov.uk/details/r/C2530393> [records copied
884 in 2001]

885 ADM 52/1105 Master's Log HMS Winchelsea (1755 Dec 29–1757 Dec 31), The National
886 Archives, UK, 1757. <https://discovery.nationalarchives.gov.uk/details/r/C253444> [records
887 copied in 2001]

888 ADM 52/1064 Master's Log HMS Terrible (1756 Feb 22–1758 Sep 30), The National Archives,
889 UK, 1757. <https://discovery.nationalarchives.gov.uk/details/r/C2534093> [records copied
890 in 2001]

891 Barriopedro, D., Gallego, D., Alvarez-Castro, C., Garcia-Herrera, R., Wheeler, D., Pena-Ortiz,
892 C., Barbosa, S.: Witnessing North Atlantic westerlies variability from ships' logbooks
893 (1685-2008). *Climate Dynamics*. Vol. 43, 939-955. DOI 10.1007/s00382-013-1957-8,
894 2014.

895 Bertler, N., Mayewski, P., Carter, L.: Cold conditions in Antarctica during the Little Ice Age-
896 Implications for abrupt climate change mechanisms. *Earth and Planetary Science*
897 *Letters*, Vol. 308(1-2), 41-51, 2011.

898 Blake, N., and Lawrence, R.: *The Illustrated Companion to Nelson's Navy*. Great Britain:
899 Chatham Publishing. p. 144, 1999.

900 Boston Herald, Oct. 17, 1757 [account of hurricane]

901 British High Commission, Ottawa, Canada Note 26-06: Letter advising the Minister of Foreign
902 Affairs of the British Government's position respecting the sovereign protection of the
903 HMS Fantome and HMS Tilbury shipwrecks, 2006.

904 Canadian Hydrographic Survey Chart: Guyon Island to Flint Island (2011) 1:37,866 [Issue Date
905 2022-11-26] 2022.

906 Chenowith, M.: Reassessment of Historical Atlantic Basin Tropical Cyclone Activity, 1700–
907 1855. *Climatic Change* Vol. 76, 169-240, 2006.

908 Cheung, H., and Chu, J. Global increase in destructive potential of extratropical transition events
909 in response to greenhouse warming. *Climate and Atmospheric Science*. Vol. 6. 2023. 10
910 pp. <https://doi.org/10.1038/s41612-023-00470-8> [Accessed 2023/11/04]

911 Corbett, J.: *England in the Seven Years' War: A Study in Combined Strategy*. 2 Vols. London:
912 Longmans, Green and Company. 407 pp, 1907.

913 Cornwall Council: *Developing Magnetometer Techniques to Identify Submerged Archaeological*
914 *Sites*. Cornwall Council Report 2010-R01. 221 pp, 2010.

915 Cronin, T., Hayo, K., Thunell, R., Dwyer, G., Saenger, C., Willard, D.: The medieval climate
916 anomaly and Little Ice Age in Chesapeake Bay and the North Atlantic Ocean.
917 *Palaeogeography, Palaeoclimatology and Palaeoecology*. Vol. 297, 299-310, 2010.

918 Day, J. and Hodges, K. Growing land-sea temperature contrast and the intensification of Arctic
919 cyclones. *Geophysical Research Letters*. Vol. 45. p. 3673-3681. 2018.
920 <https://doi.org/10.1029/2018GL077587> [Accessed 2023/11/04]

921 Dezileau, L., Sabatier, P., Blanchemanche, P., Joly, B., Swingedouw, D., Cassou, C., Castaings,
922 J., Martinez, P., Von Grafenstein, U.: Intense storm activity during the Little Ice Age on
923 the French Mediterranean coast. *Palaeogeography, Palaeoclimatology, Palaeoecology*.
924 Vol. 299, 289–297, 2011.

925 Duggan, R.: Coastal Heritage Planning at the Fortress of Louisbourg – Planning it Out *in*
926 Archaeology in Nova Scotia: 2010 News. Halifax: Nova Scotia Museum Collections
927 Unit, 1-8, 2010.

928 Donnelly, J., Bryant, S., Butler, J. and Dowling, J.: 700 yr. sedimentary record of intense
929 hurricane landfalls in Southern New England. Geological Society of America Bulletin
930 Vol. 113, 714-727, 2001.

931 Finck, P.: A Geological and Coastal Vulnerability Analysis of Point Michaud Provincial Park,
932 Richmond County, Nova Scotia. Nova Scotia Natural Resources Open File Report ME
933 2015-003. 25 pp. 2015.

934 Garcia, R., Diaz, R., Herrera, G., Eischeid, M., Prieto, E., Hernandez, L. Gimeno, F., Duran, R.
935 and Bascary, A.: Atmospheric circulation changes in the tropical Pacific inferred from the
936 voyages of the Manila Galleons in the sixteenth-eighteenth centuries, Bulletin of the
937 American Meteorological Society. Vol. 82, 2435–2455, 2001.

938 Garcia-Herrera, R., Wilkenson, C. Koek, F., Prieto, M., Jones, P. and Koek, F.: Description and
939 general background to ships' logbooks as a source of climatic data. Climatic Change.
940 Vol. 73, 13-36, 2005a.

941 García-Herrera R., Können G., Wheeler, D., Prieto, M., Jones, P., Koek, F.: CLIWOC: a
942 climatological database for the world's oceans 1750- 1854. Climatic Change. Vol. 73, 1–
943 12, 2005b.

944 Gebbie, G.: Atlantic warming since the Little Ice Age. Oceanography. Vol. 32, 220–230, 2019.

945 Gurgis, J. and Fowler, A.: A history of ENSO events since A.D. 1525: implications for future
946 climate change. Climatic Change. Vol. 92, 343-387, 2009.

947 Hanson, R.: St. Peter's Island to Kelpy Cove, Southeast Coast, Cape Breton Unpublished 1" =
948 3000' field sheet. Canadian Hydrographic Survey, 1954.

949 Hart, R. and Evans, J.: A climatology of extratropical transition of Atlantic tropical cyclones.
950 Journal of Climate. Vol. 14, 546-564, 2001.

951 Jackson, D., Costas, S., Guisado-Pintado, E.: Large-scale transgressive coastal dune behaviour in
952 Europe during the Little Ice Age. Global and Planetary Change. Vol. 175, 82-91, 2019.

953 Johnstone, (James) Chevalier: The campaign of Louisbourg, 1750-'58 [microform]: a short
954 account of what passed at Cape Breton, from the beginning of the last war (1750) until
955 the taking of Louisbourg, by the English, in the year of Our Lord, 1758. Memoirs of the
956 Chevalier de Johnstone Vol. 3 of 3. 33 pp, [Translated in 1871 by Charles Winchester],
957 1758.

958 Jones, P. and Mann, M.: Climate over past millennia. American Geophysical Union.
959 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2003RG000143>. 42 pp, 2004.

960 Keigwin, L.: The Little Ice Age and medieval warming period in the Sargasso Sea. Science, Vol.
961 274, 1503-1508, 1996.

962 Keigwin, L., Sachs, J., and Rosenthal, Y.: A 1600-year history of the Labrador Current off Nova
963 Scotia. *Climate Dynamics*. Vol. 21, 53–62, 2003.

964 Knabb, R. Rhone, J. and Brown, D.: Tropical cyclone report – Hurricane Katrina 23-30 August
965 2005. National Hurricane Center Tropical Cyclone Report. 43 pp,
966 https://nhc.noaa.gov/data/tcr/AL122005_katrina.pdf, 2023. [accessed 2023-09-30]

967 Knox Bristol Journal, November 12, 1757. [hurricane survivor account]

968 Knox, J. (Captain): Historical journal of the campaigns in North America for the years 1757,
969 1758, 1759 and 1760. Volume 1 of 3. London: W. Johnston (Ludgate Street), and Dodsly,
970 J. (Pall Mall), 49, 1769.

971 Kreutz, K., Mayewski, P., Meeker, L., Twickler, M., Whitlow, S., and Pittalwa, I.: Bipolar
972 changes in atmospheric circulation during the Little Ice Age. *Science*. Vol. 277 (5330),
973 1294-1296, 1997.

974 Lalbeharry, R., Bigio, R., Thomas, B. and Wilson.: Numerical simulation of extreme waves
975 during the storm of 20-22 January 2000 using winds generated by the CMC weather
976 prediction model. *Atmosphere-Ocean*. Vol. 47, 99-122, 2009.

977 Lamb, H.: *Climate, history, and the modern world*. Methuen: New York. 387 pp, 1982.

978 Landsea, C. Anderson, C., Charles, N., Dunion, J., Clark, G., Fernandez-Partag´as, J.,
979 Hungerford, P., Neumann, C., and Zimmer, M.: The Atlantic hurricane database re-
980 analysis project: Documentation for the 1851–1910 alterations and additions to the
981 HURDAT database, *in* Murnane, R. J. and Liu, K.-B. (eds.), *Hurricanes and Typhoons:
982 Past, Present, and Future*, Columbia University Press, 177–221, 2004.

983 Lavery, B.: *The Ship of the Line; Volume 1 – Development of the Battlefleet 1650-1850*.
984 London: Conway Maritime Press. 224 pp, 1983.

985 Lavery, B.: *The Ship of the Line; Volume 2 – Design, Construction and Fittings*. London:
986 Conway Maritime Press. 191 pp, 1984.

987 Lavery, B.: *The Royal Navy’s First Invincible*. Portsmouth, United Kingdom: Invincible
988 Conservations Limited – Burgess and Son (Abingdon) Limited. 119 pp, 1988.

989 Lixion, A.: National Hurricane Center Tropical Cyclone Report – Hurricane Juan. National
990 Hurricane Center Report – Hurricane Juan. [https:// www.nhc/noaa.gov/data/tcr/
991 AL152003_Juan.pdf](https://www.nhc.noaa.gov/data/tcr/AL152003_Juan.pdf) 11 pp, 2003 (Revised 2012) [accessed 2023-9-30]
992 Lixion, A., Stewart, S., Berg, R. and Berg, A.: National Hurricane Center Tropical Cyclone
993 Report – Hurricane Dorian. https://www.nhc.noaa.gov/data/tcr/AL052019_Dorian.pdf 74
994 pp, 2020. [accessed 2023-9-30]
995 London Magazine, November, 563-564, 1758.
996 The London Chronicle, July 23-26, 1757.
997 Lamb, H.: Historical storms of the North Sea, British Isles and Northwest Europe. London:
998 Cambridge University Press. 204 pp, 1991.
999 Ludlum, D.: Early American Hurricanes 1492-1870. American Meteorological Society. 198 pp,
1000 1963.
1001 Mann, M.: Little Ice Age *in* The Earth System: Physical and Chemical Dimensions of Global
1002 Environmental Change. MacCracken, M and Perry, S. (eds.) Encyclopedia of Global
1003 Environmental Change. Chichester, UK: Wiley and Sons Ltd., 504-509, 2002.
1004 Matthes, F.: Report of Committee on Glaciers, April 1939. Transactions, American Geophysical
1005 Union. Vol. 20, 518, 1939.
1006 Mazzarella, A. and Scafetta, N.: The Little Ice Age was 1.0-1.5 °C cooler than current warm
1007 period. Climate Dynamics. Vol. 51, 3957-3968, 2018.
1008 McLennan, J.: Louisbourg from its foundation to its fall. Sydney, Nova Scotia: Fortress Press
1009 (1969 Reprint), 207-210, 1918.
1010 Miller, G., Geirsdóttir, Á., Zhong, Y., Larsen, D., Otto-Bliesner, B., Holland, M., Bailey, D.,
1011 Refsnider, K., Lehman, S., Southon, J., Anderson, C., Björnsson, H., and Thordarson, T.:
1012 Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean

1013 feedbacks. *Geophysical Research Letters*. Vol. 39, DOI:10.1029/ 2011GL050168, 5 p,
1014 2012.

1015 Mowat, H., Lieutenant *in* Holland, S.: A plan of the island of Cape Breton reduced from Captn.
1016 Holland's Survey. [soundings and naval observations were taken by Lieut. Henry Mowat,
1017 commander of His Majesty's armed ship Canceaux and the officers of the ship under his
1018 direction], 1776.

1019 Nature Notes, 24 August, 415, 1882. [European heat wave of 1757]

1020 Oliva, H., Peros, M., Viau, A., Reinhardt, E., Nixon, F. and Morin, A. A multi-proxy
1021 reconstruction of tropical cyclone variability during the past 800 years from Robinson
1022 Lake, Nova Scotia, Canada. *Marine Geology*. Vol. 406. 84-97, 2018.

1023 Oliver, J. and Kington, J.: The usefulness of ships' log-books in the synoptic analysis of past
1024 climates. *Weather*. Vol. 25 (12), 520-528, 1970.

1025 Pasche, R., Berg, R., Roberts, D. and Papin, P. National Hurricane Center Tropical Cyclone
1026 Report – Hurricane Laura (AL132020) August 20-29 2020. 75 pp,
1027 https://nhc.noaa.gov/data/tcr/AL132020_Laura.pdf, 2021 [accessed 2023-9-30]

1028 Poey, A.: A chronological table, comprising 400 cyclonic hurricanes which have occurred in the
1029 West Indies in the North Atlantic within 362 years, from 1493 to 1855. *Journal of the*
1030 *Royal Geographical Society* Vol. 25, 291–328, 1855.

1031 Richey, J., Poore, R., Flower, Benjamin P., Quinn, T., and Hollander, D.: Regionally coherent
1032 Little Ice Age cooling in the Atlantic Warm Pool. *Geophysical Research Letters*. Vol. 36,
1033 L21703, doi:10.1029/2009GL040445, 1-5, 2009.

1034 Ruffman, A.: The multidisciplinary rediscovery and tracking of ‘The Great Newfoundland and
1035 Saint-Pierre et Miquelon hurricane of 1775.’ *The Northern Mariner/Le Marin du Nord*.
1036 Vol. 1, 11-23, 1996.

1037 Saegasser, L.: The U.S.S. Macedonian and the hurricane of 1818. *Proceedings of the U.S. Naval*
1038 *Institute*. [https://www.usni.org/magazines/proceedings/1970/january/uss-macedonian-](https://www.usni.org/magazines/proceedings/1970/january/uss-macedonian-and-hurricane-18181970)
1039 [and-hurricane-18181970](https://www.usni.org/magazines/proceedings/1970/january/uss-macedonian-and-hurricane-18181970), 1970. [Accessed 2023-9-30]

1040 Saenger, C., Cohen, A., Oppo, D., Halley, R. and Carilli, J.: Surface-temperature trends and
1041 variability in the low-latitude North Atlantic since 1552. *Nature Geoscience*, 492-495,
1042 2009.

1043 Sicre, M., Jalali, B., Martrat, B., Schmidt, S., Bassetti, M., Kallel, N.: Sea surface temperature
1044 variability in the North Western Mediterranean Sea (Gulf of Lion) during the Common
1045 Era. *Earth and Planetary Science Letters*. Vol. 456, 124-133, 2016.

1046 Smyth, W. *The Sailor’s Word-book: an alphabetical digest of nautical terms, including some*
1047 *more especially military and scientific... as well as archaisms of early voyagers, etc.* by
1048 *the late Admiral W.H. Smyth (2004 Reprint) Toronto: Algrove Publishing Limited. 744*
1049 *pp, 1867.*

1050 Stoetzel, D.: *Encyclopedia of the French and Indian War in North America, 1754-1763*. United
1051 *Kingdom: Heritage Books. 579 pp, 2008.*

1052 Storm, A.: *Seaweed and Gold*. Sydney Nova Scotia: City Printers Limited. 192 pp, 2002.

1053 Studholme, J. Fedorov, A., Gulev, S. Emanuel, K. and Hodges, K. Poleward expansion of
1054 tropical cyclone latitudes in warming climates. *Nature Geoscience*. Vol. 15. p. 14-18.
1055 2022. <https://doi.org/10.1038/s41561-021-00859-1> [Accessed 2023/11/04]

- 1056 Syrett, D.: Shipping and Military Power in the Seven Years' War, 1756-1763: The Sails of
1057 Victory. United Kingdom: Liverpool University Press. 192 pp, 2008.
- 1058 Trouet, V., Scourse, J. and Raible C.: North Atlantic storminess and Atlantic meridional
1059 overturning circulation during the last millennium: reconciling contradictory proxy
1060 records of NAO variability. *Global and Planetary Change*. Vol. 84-85, 48-55, 2012.
- 1061 Trouet, V., Diaz, H., Wahl, E., Viau, H., Graham, R., Graham, N., and Cook, E. A 1500-year
1062 reconstruction of annual mean temperature for temperate North America on decadal to
1063 multi-decadal timescales. *Environmental Research Letters*. Vol. 8, 10 pp. 2013.
1064 Doi:10.1088/1748-9326/8/2/024008
- 1065 U.S. Department of Commerce Environmental Science Services Administration Weather Bureau.
1066 Tropical Cyclone Report - Hurricane Camille August 14-22, 1969. 58 pp.
1067 <https://www.nhc.noaa.gov/pdf/tcr-1969Camille.pdf>, 1969. [accessed 2023-9-30]
- 1068 Van Vliet-Lanoë, B., Goslin, J., Hallégouët, B., Hénaff, A., Delacourt, C., Fernane, A., Franzetti,
1069 M., Le Cornec, E., Le Roy, P., and Penaud, A.: Middle- to late-Holocene storminess in
1070 Brittany (NW France): Part I – morphological impact and stratigraphical record. *The*
1071 *Holocene*. Vol. 24, 413–433, 2014.
- 1072 Vecchi, G., and Knutson, T.: On estimates of historical North Atlantic tropical cyclone activity.
1073 *Journal of Climate*. Vol. 21, 3580-3600, 2008.
- 1074 Virost, E., Ponomarenko, A., Dehandschoewercker, E., Quere, D. and Clanet, C.: Critical wind
1075 speed at which trees break. *Physics Review*. Vol. 93, 7 pp, 2016.
- 1076 Warden, D.: A statistical, political, and historical account of the United States of North America
1077 from the period of their first colonization to the present day. Vol. 1 of 3, 552 pp, 1819.

- 1078 Wheeler, D.: An examination of the accuracy and consistency of ships' logbook weather
1079 observations and records. *Climate Change* Vol. 31, 97-116, 2005.
- 1080 Wheeler, D.: The Great Storm of November 1703 – A new look at the seamen's records.
1081 *Weather*. Vol. 58, 419-427, 2006.
- 1082 Wheeler, D. and Wilkinson, C.: From calm to storm: the origins of the Beaufort Wind Scale. *The*
1083 *Mariner's Mirror*. Vol. 90, 187-201. DOI:10.1080/00253359.2004.10656896, 2004.
- 1084 Wheeler, D., Garcia-Herrera, R. and Wilkinson, C.: Atmospheric circulation and storminess
1085 derived from Royal Navy logbooks: 1685 to 1750. *Climatic Change*. Vol. 101, 257-280,
1086 2010.
- 1087 Winter, A., Ishioroshi, H., Watanabe, T., Oda, T., Christy, J.: Caribbean sea surface
1088 temperatures' two-to-three degrees cooler than present during the Little Ice Age.
1089 *Geophysical Research Letters*. Vol. 27, 3365-3368, 2000.
- 1090 Zinck, J.: *Shipwrecks of Nova Scotia*. 226 pp, 1975.