1 A Major Midlatitude Hurricane in the Little Ice Age

2 John Dickie^{1,2} and Grant Wach¹

¹Basin and Reservoir Lab, Department of Earth and Environmental Sciences

4 Dalhousie University, Halifax, Canada B3H 4R2

5 ²Corresponding Author

6 Contacts: john.dickie@dal.ca; grant.wach@dal.ca


7 Abstract

8 An unusually severe hurricane (Louisbourg Storm) struck Nova Scotia Canada in 1757. Historic records describing storm conditions as well as damage to ships and coastal fortifications indicate 9 an intensity beyond any modern (post-1851) Atlantic cyclones striking the same region, yet this 10 storm struck during a cold climate period known as the Little Ice Age (LIA). Its track and timing 11 coincided with a British naval blockade of a French fleet at Fortress Louisbourg during the Seven 12 Years' War (1756-1763). This provides a unique opportunity to explore growing scientific 13 evidence of heightened storminess in the North Atlantic despite a colder climate expected to 14 suppress hurricane intensification but which research is increasingly showing to have supported 15 North Atlantic storms of exceptional strength. Weather attributes extracted from the logs of 16 naval vessels scattered by the Louisbourg Storm provided multiple hourly observations recorded 17 at different locations. Wave height and wind force estimates at ship locations were compared to 18 extreme storm surge heights calculated for Louisbourg Harbour and a shipwreck site south of 19 Fortress Louisbourg. Comparing these metrics to those of modern analogs that crossed the same 20 bathymetry reflects landfall intensity consistent with a powerful major hurricane. Historical 21 records show this storm originated as a tropical cyclone at the height of hurricane season and 22 23 intensified into the northern midlatitudes along the Gulf Stream. Its intensity at landfall is

consistent with established seasonal climatological models where highly baroclinic westerlies 24 driven by autumn continental cooling encounter intensifying north-tracking tropical cyclones 25 fueled by sea surface temperatures that peak in autumn. Stronger seasonal contrasts from earlier 26 and colder continental westerlies in the Little Ice Age (LIA) may have triggered explosive 27 extratropical transition from a large hurricane resulting in a more severe strike. It suggests that 28 29 tropical cyclones lasting days to weeks and the conditions that generate them are likely masked by cooler historic mean-annual to multi-decadal LIA climate reconstructions. Predictions of 30 warmer midlatitude sea surface temperatures could see powerful hurricanes intensify into higher 31 32 latitudes later into the fall, potentially recreating the strong contrasts that triggered the intensity of the Louisbourg Storm. 33 **1.0 Introduction** 34 On September 25, 1757, a powerful hurricane struck the coast of Cape Breton Island, 35 Nova Scotia, Canada (Fig. 1). There would have had no record of the 'Louisbourg Storm' had it 36

37 not coincided with a British naval blockade of France's Fortress Louisbourg during the Seven

38 Years' War (1756-1763). Three French naval squadrons at Louisbourg and the blockading



39

Figure 1. Study location in Nova Scotia, Canada. Arrow length and orientation represents the 40 distance and direction traveled by the British fleet on September 21-26, 1757. September 25 and 41 26 show the path of the *Invincible* south of the wider dispersal of the British fleet after being 42 43 scattered by the storm (dotted oval). The storm's location off New England is estimated (off map). The estimated storm track (dashed line) shows eye locations for the dates shown Inset 44 45 shows the study area relative to the North Atlantic and the hurricane track based on historic 46 records showing its progressive northward translation seaward of (1) Florida (no date), (2) North Carolina (September 23), (3) New England (September 24) and (4) Cape Breton Canada 47 (September 25-26). Fort Cumberland is 70 km toward 293 Azimuth. 48

British fleet placed 49 sailing battleships and warships in the path of a storm-descriptions of
damage to ships and coastal infrastructure, severe flooding from rainfall and extreme storm surge

51 suggest was more intense than any landfalling storm in Canadian waters since modern records

began in 1851 (Landsea et al., 2004, Finck, 2015). This suggests it had the intensity of a major
hurricane at landfall (Category 3+ on the Saffir-Simpson Hurricane Wind Scale) yet it struck
during the colder climate of the 'Little Ice Age' (LIA; c1300-1850).

Hurricanes are fueled by sea surface temperature (SST) over 28C. They rapidly lose 55 energy as the move north over cooler midlatitude waters, and many tropical cyclones undergo 56 57 extratropical transition which releases tropical energy at increasingly higher latitudes later in hurricane season (Hart and Evans, 2001). Modern tropical cyclone intensity is characterized in 58 59 real time with instruments carried by aircraft, satellites and at ground stations. In contrast, preindustrial metrics must be derived from historical observational records. Subjective interpretation 60 61 and geographic bias can make them less reliable than instrumental data (e.g., Jones and Mann 2004), yet they offer a temporal resolution unavailable in scientific proxies, and they straddle the 62 end of the LIA and the rise of modern anthropogenic emissions. Oliver and Kington (1970) and 63 Lamb (1982) first explored their suitability for weather research. Naval logbooks were 64 65 subsequently found to be a superior source of historical weather data given that hourly ship observations were systematically recorded in real time with a consistent terminology. Logbook 66 67 data have been compiled to assess historical atmospheric circulation patterns (e.g., Garcia et al., 68 2001, Garcia-Herrera et al., 2005a, Wheeler et al., 2010, Barriopedro et al., 2014). CLIWOC, the Climatological Database for the World's Oceans, was compiled from historical British, French, 69 70 Dutch and Spanish naval logbooks. It established a common historical wind force terminology to 71 document ocean surface atmospheric circulation patterns between 1750 and 1850 (Garcia-72 Herrera et al., 2005b).

To date, pooled historical naval records have been used to identify longer-term regional
circulation patterns and extend the multidecadal climate signal into the industrial period (e.g.,

Garcia-Herrera at al., 2005a, 2005b, Wheeler et al., 2010, Barriopedro et al., 2014). In contrast, 75 76 this study takes advantage of an unusual concentration of warships in the path of a single 77 hurricane to characterize its intensity. It seems counterintuitive that the colder LIA climate would generate more powerful midlatitude Atlantic cyclones than in the modern era, yet historical 78 records show the LIA to be generally 'stormier' with unusually powerful midlatitude hurricanes 79 80 despite conditions that dampen hurricane energy. Donnelly et al.'s (2001) historic storm reconstruction from Mattapoisett Pond, Massachusetts, and Oliva et al.'s (2018) historic storm 81 82 reconstruction from Robinson Lake, Nova Scotia, are among a growing number of proxy studies showing that major Atlantic cyclones struck the northeastern seaboard of North America in the 83 LIA. Since winter extratropical cyclones known as Nor'easters cannot be differentiated from 84 Atlantic tropical cyclones and their extratropical derivatives from proxy data alone, historical 85 records can constrain the timing of midlatitude hurricanes and tropical storms. 86 This study utilizes a unique historical data set to characterize the intensity of the 87

Louisbourg Storm using spatial and temporal weather metrics extracted from ship logbooks of both the English and French fleets, British Admiralty records and official documents of both nations, and compares the derived storm metrics to those of modern systems that tracked across the same bathymetry. Characterizing its intensity tests historical descriptions of an unusually severe storm and may help establish a more thorough understanding of LIA hurricane climatology.

94 **2.0 Methodology**

95 *2.1 Historical Records*

96 Eighteenth century navigation and weather data were entered hourly in the daily logs of
97 naval vessels, resulting in reliable records suitable for historical climate research. A noon

98	sighting of the sun fixed latitude and marked the start of the sea day. Britain adopted the
99	Gregorian calendar in 1752 so dates in logs used for this study did not require correction. In 1757
100	a local meridian was used to determine longitude, deduced from logs to have been based on
101	Louisbourg Lighthouse (Fig. 2).
102	Historical British Admiralty Correspondence and Papers (ADM1/481, 1488, 2294)
103	covering storm damage to British vessels on the 'Halifax Station' in 1757 and Fleet Lists
104	(ADM8/31, 32) are preserved at the National Archives at Kew (UK), as are Royal Navy Master's
105	(ADM 51/409, 633,1075) and Captain's (ADM 52/578,819,1064) logbooks. Lieutenant's logs
106	(ADM51) kept at the National Maritime Museum, Greenwich, were often incorporated into
107	Captain's logs with addenda. Master's and Captain's logs of the Royal Navy warships Invincible,
108	Windsor, Sunderland, Eagle, Terrible, Grafton, Newark, and Captain, plus ancillary official
109	correspondence, were used in this study. All logs were consistent in content and format. Letters
110	and logbook entries written in ink were copied from cursive in multiple handwriting styles to a
111	more readable format, interpreted, compiled into a time sequence and cross referenced. Logs
112	from French warships Fleur de Lys, l'Abenaquise, Tonnant, l'Inflexible and Dauphin Royal
113	translated from French describe conditions in Louisbourg Harbour (McLennan, 1918). Wind
114	directions from gimballed ships' compasses reference magnetic north. Bearings and wind
115	directions used the 32 points of the compass (Smyth, 1867, Blake and Lawrence, 1999) and were
116	translated to azimuths. The logs of British ships at sea and French ships moored in Louisbourg
117	Harbour contained: (1) dates and times, (2) position, (3) bearing, (4) wind direction, (5) wind
118	speed terms that evolved into the Beaufort Wind Scale (e.g., Garcia-Herrera et al., 2005a, 2005b,
119	Wheeler, 2005, Wheeler et al., 2010), and (6) descriptions of sea state.
120	2.2 Proxy Climate Context

Major atmospheric circulation patterns that influence Atlantic tropical cyclone behaviour, 121 specifically the El Nino Southern Oscillation (ENSO) and North Atlantic Oscillation (NAO), 122 123 have been reconstructed for the historical period (e.g., Gurgis and Fowler, 2009, Trouet, et al., 2012). These proxy climate patterns provide an overarching context since La Nina years create 124 conditions conducive to driving hurricanes in the Atlantic, and a negative NAO allows Atlantic 125 126 tropical cyclones to enter the Atlantic and potentially reaching the midlatitude eastern seaboard. 127 Atmospheric circulation patterns for 1757 were studied to assess overarching conditions 128 conducive to Atlantic hurricane generation.

129 *2.3 Wind Speed*

Wheeler and Wilkinson's (2004) analysis of the derivation of the Beaufort scale shows 130 terms that vary little from the logbook terms used in this study. A similar approach has been 131 adopted here with adjectives describing primary nomenclature. A 'gale' (Beaufort Force 8) was 132 originally between a breeze (Force 2) and a violent storm (Force 11) and established a 133 134 benchmark (Table 1). A 'near gale,' its diminutive (Smyth, 1867) corresponds to a 'moderate gale.' Wheeler et al. (2010) categorized 'strong gale,' 'hard gale,' 'blew hard' and 'storm' as 135 stronger than 'fresh gale.' Adjectives 'stiff' and 'fresh' indicate winds stronger than a gale 136 137 (Force 9) while a 'severe' or 'hard' gale reflects a 'storm' (Force 10). 'Excessive' and 'extreme' hard gale, necessarily stronger than a 'hard gale' would then correspond to 'violent storm' (Force 138 139 11) which does not appear in the logs used here. 'Hurricane' (Force 12) is mentioned in both 140 French and British records. 'Squall' is a historical term for an increase in wind speed sustained 141 above threshold for at least one minute. The National Oceans and Atmospheric Administration 142 (NOAA) defines it as a sudden increase by at least 16 knots (33 kph) and sustained at over 22 143 knots (41 kph) for one minute. Environment and Climate Change Canada (ECCC) defines

Logbook Term	Beaufort Scale	Rating	Wind (kph)
Hurricane	Hurricane	12	118+
Excessive / Extreme Hard Gale	Violent storm	11	103-117
Severe / Hard Gale	Storm	10	89-102
Strong / Stiff Gale	Strong Gale	9	75-88
Gale	Gale	8	62-74
Moderate Gale	Near Gale	7	50-61
Strong / Stiff Breeze	Strong Breeze	6	39-49

Table 1. Logbook Beaufort Terms and Associated Windspeeds (kph).

squalls as increases of 34 knots (63 kph) or more above prevailing winds sustained for over a 145 146 minute. The World Meteorological Organization (WMO) uses 8 m/s and 11 m/s (29 and 40 kph) 147 above threshold for over one minute while the American Meteorological Association (AMA) notes squalls are of 'several minutes' duration. In considering these definitions 'squall' is taken 148 149 to be a sudden increase in wind speed of 40-60 kph above threshold and sustained for at least one minute. We interpret 'hard' squalls as the upper end of the spectrum in the way adjectives were 150 151 used to create the historic Beaufort scale (Wheeler and Wilkinson, 2004). Heavy rains accompanying squalls noted in the logs appear to be consistent with descriptions of hurricane 152 spiral bands. 153

In this study the Beaufort Wind Force Scale is used to describe wind speeds from gale to hurricane force (63-118 kph). The Saffir-Simpson Hurricane Wind Scale describes hurricane winds greater than 118 kph with peak wind speeds averaged over one minute defining hurricane intensity Categories 1-5. A major hurricane is Category 3 (178-208 kph) or stronger. Wind speeds derived from log entries were plotted from the first southeasterlies noted off Nova Scotia on September 22, 1757, to diminishing westerlies at the storm's end on September 26. Ephemeral squalls of 1 min duration above threshold winds provide an estimate of total wind speed sustained for one minute or longer. Wind speeds at mid-mast height above the deck plus freeboard (distance from the waterline to the upper deck) approximate the 10 m height above ground level for modern hurricane wind speed measurements.

Eighteenth century navies knew hurricanes commonly encountered in the Caribbean 164 sometimes reached North America's eastern seaboard. The modern Saffir-Simpson scale 165 166 provides a 1 to 5 storm intensity rating based on a hurricane's maximum sustained wind speed averaged over one minute. Since no such real time wind force measurement existed in 1757, this 167 168 study has adopted Virot et al.'s (2016) engineering analysis of critical hurricane wind speeds that 169 break trees as a model for estimating threshold wind speeds needed to break ships' masts. Ships' logs indicate they maintained course relative to prevailing storm winds. This placed vessels at a 170 highly oblique angle to wave crests which minimized pitch and yaw, and held masts within a 171 stable plane of reference against which wind applied a sustained force. In addition, large vessels 172 (74-gun third rates) with up to nine feet of flooding in the hold would have a lower center of 173 174 mass that would have affected its righting moment and minimized directional variance in the wind force striking the masts. Rigging designed to stabilize the masts and transfer wind energy 175 176 through the sails would likely have required a higher sustained wind force to achieve failure. 177 2.4 Wind Direction

Wind direction was measured using the ship's magnetic compass and entered in the ships' logs as 'points of the compass.' These entries were translated to azimuths. Compass directions are relative to magnetic north and not corrected for declination given the small study area and short time frame. Eighteenth century navigation was inaccurate but this study benefits from (1) log entries of the fleet relying on smaller vessels sent inshore to establish distance from coastal landmarks, and (2) during the storm ships were driven sufficiently close to land that their

positioning entries were based on triangulation using landmarks which greatly improves
accuracy. Experienced navigators were also able to correct for ship motion in their readings
while the ship's position was typically determined by a Lieutenant plus one or more midshipmen
and the sailing master's mate.

188 2.5 Wave Height

189 Wave height was estimated based on descriptions compared to ship dimensions and is the last accurate metric. Historic references to ship structure in Imperial Units have been converted 190 191 to metric. This includes the distance from the keel to the upper deck and freeboard from the 192 waterline to the upper deck. The depth of water needed to spill over the bow to flood the upper deck and tear away large ship's boats tethered to the deck is estimated. References such as 193 sailors being swept off spars 80' (24 m) above the waterline offers an estimate of peak wave 194 heights. Warships were designed for stability as floating gun platforms and to return to an 'even 195 keel' as quickly as possible after firing. Wave descriptions in Louisbourg Harbour are the least 196 197 reliable since they include storm surge.

198 2.6 Surge

199 Surge is a rise in sea level due to atmospheric pressure and storm winds and is 200 proportional to a tropical cyclone's intensity and translation rate. Coastal surge is a reasonable estimate of storm intensity and can serve as a test of intensity derived from wind data. The surge 201 202 height of modern analogs that struck Nova Scotia after tracking across the Scotian Shelf and 203 whose intensity has been characterized with metrics derived using modern meteorological 204 methods provides a reliable benchmark for comparison to surge calculated for the 1757 storm. In 205 this study, storm surge at known locations and elevations above sea level were described at (1) 206 Battery de la Grave at Fortress Louisbourg, (2) the historic town within the Fortress, and (3) St.

Esprit where the British warship *HMS Tilbury* was stranded in water depths it could not normally
navigate given its displacement. All surge calculations were then corrected for (1) relative sea
level (RSL) rise since 1757, and (2) a mid-tide RSL datum used by Google Earth versus a lowest
low water (tide) datum used by the Canadian Hydrographic Service for a (draft) navigation chart
used for the Tilbury wreck site. In addition, French records noting the tidal change at
Louisbourg allowed for the timing of the tidal cycle to be backed out to determine storm surge
versus storm tide.

214 *Tilbury's* wreck site offered a chance to estimate surge at a second location 45 km southwest of Louisbourg. *Tilbury's* identity was confirmed in 1986 with the discovery of the 215 216 ship's bell, most of its guns, anchors and artifacts (Storm, 2002). Its location remained undisclosed after a letter from the British High Commission in 2006 reminded the Minister of 217 Foreign Affairs Canada of the wreck's sovereign immunity, resulting in Nova Scotia rescinding 218 219 the associated treasure trove license. Relocating the wreck to confirm its water depth required 220 creating a digital bathymetric chart needed to guide a marine magnetometer survey leading to 221 site confirmation by divers.

222 **3.0 Little Ice Age Storminess**

Matthes (1939) named the LIA to explain European glacier expansion during a historically colder climate period. Heightened climate variability saw deeply cold winters and cooler mean annual temperatures primarily in the northern hemisphere (e.g., Kreutz et al., 1997, Mann, 2002, Jones and Mann, 2004). It may have been triggered by late 13th Century volcanic eruptions and a cooling feedback process sustained by Arctic sea-ice expansion (Miller et al., 2012). North Atlantic mean annual SSTs were 1-2°C cooler than today (e.g., Keigwin, 1996, Winter et al., 2000, Richey et al., 2009, Saenger et al., 2009, Cronin et al., 2010, Bertler et al.,

230	2011, Mazzarella and Scaffeta, 2018, Gebbie, 2019). The Maunder Minimum, the coldest part of
231	the LIA, (MM; 1645-1715) saw greater 'storminess' during polar air breakouts from Europe
232	correlating to more frequent easterly gales in the English Channel and Approaches in 1685-1750
233	(Wheeler et al. 2010). Concentrated storm horizons in coastal dunes across western Europe and
234	in Brittany and on France's Mediterranean coast correlate to the coldest part of the LIA
235	(Dezileau et al., 2011, Van Vliet-Lanoe et al., 2014, Sicre et al., 2016, Jackson et al. 2019).
236	Dezileau et al. (2011) attributed LIA storminess to cold-enhanced lower tropospheric
237	baroclinicity modifying prevailing westerlies. In the northwest Atlantic, Donnelly et al. (2015)
238	described major hurricane deposits in New England coastal sediments dating to 1635, 1638 and
239	1815. Ludlum's (1963) compilation of historical northwest Atlantic hurricanes and tropical
240	storms includes the LIA's major 'Independence Hurricane' that struck New England on August
241	29, 1775 and the 'Newfoundland Hurricane' of September 9, 1775, a storm that left 4000 dead to
242	become Canada's deadliest hurricane (Ludlum, 1963, Ruffman, 1996). Lamb's (1991)
243	exhaustive survey of British and European storms includes the Great Storm that devastated the
244	British Isles on November 26, 1703. It was an extratropical cyclone equal to a Category 2
245	hurricane yet Wheeler (2003) notes a far more powerful Atlantic storm on December 1-12, 1792,
246	also late in Atlantic hurricane season. Both were anomalous for a colder climate period.
247	The Scotian Shelf on Canada's Atlantic seaboard (Fig. 1) is dominated by the cold, south-
248	flowing, low-salinity Labrador Current. It originates in the Davis Strait of the Canadian Arctic
249	and hugs the coast to the start of the midlatitudes at Cape Hatteras, North Carolina where it
250	meets, mixes with, and redirects seaward the tropical, north-flowing more saline Gulf Stream.
251	The Labrador Current plays a critical role in hurricane extratropical transition by providing a
252	coastal buffer of cooler sea surface temperatures that effectively cut off the tropical energy of the

Gulf Stream (Hart and Evans, 2001). Summer and fall bring warm water eddies from the Gulf 253 Stream and warmer coastal SST. Sediment cores from the Emerald Basin off Nova Scotia show 254 255 1600 years of cold Labrador Current temperatures and a sudden and sustained warming around 1850 that has continued into the present (Keigwin et al., 2003) and coincides with the end of the 256 LIA. Storm compilations by Landsea et al. (2004) and Chenowith (2006) show a progressive 257 258 increase in the number of historical Atlantic tropical cyclones from 1700 and a sharp increase in 259 the number and percentage reaching New England and eastern Canada beginning around 1850. Vecchi and Knutson (2008) in a study of data from the start of instrumental data collection in 260 261 1880 show a strong correlation between mean annual SST and storm frequency. Historical records offer seasonal weather detail not captured by annual to multidecadal 262 263 proxy trends. Anomalous midlatitude coastal sea surface temperatures (SSTs) over days to weeks, conditions that fuel tropical cyclones, are therefore not likely to appear in annualized data 264 weighted by colder, sustained LIA winters. Jacoby and D'Arrigo's (1989) North American 265 266 northern and Arctic temperature reconstruction shows above normal temperatures in the 1750's. Lieutenant John Knox recorded unusually high temperatures in Halifax Harbour on July 20, 267 268 1757, which fellow officers found hotter than Gibraltar and the Mediterranean (Knox, 1769). 269 This coincided with a heat wave in Britain and southwest Europe from July into early August 270 1757 that set temperature records that stood for over 250 years (The London Chronicle, July 23-271 26, 1757; London Magazine, November 1758 p. 563-4). London on July 16-26 had an average 272 high of 41.2C (Nature Notes, 24 August 1882, p. 415). This does not assume weather conditions 273 in Europe fueled a hurricane tracking into Atlantic Canada, but demonstrates that unusually hot 274 temperatures across the northern hemisphere known to intensify midlatitude hurricanes existed in the summer of 1757. 275

The 1757 hurricane noted by Poey (1855) and Ludlum (1963) was confirmed as a 276 hurricane in Chenowith's (2006) re-assessment compilation. It was first seen off Florida and 277 278 followed the coastline past Cape Hatteras to New England on September 22-24 (Ludlum, 1963). Benjamin Franklin's observations of this specific storm led him to conclude that hurricanes "are 279 produced by currents of cold winds rushing from the north along the Atlantic coast and mingling 280 281 with the warm winds produced by the gulf-stream" (Warden, 1819). It struck the British frigate HMS Winchelsea on September 23 to 24 at 36°45'N 70° 54'W (off North Carolina over the Gulf 282 283 Stream). The log notes gale force east then east-southeast and south winds between 10 p.m. and 284 5 a.m. on September 23 which, 15 minutes later, veered violently to the northeast and then northwest at 'near hurricane' intensity. It split the main sail and broke the main mast. It was also 285 accompanied by a 'great sea' (ADM 52/1105). 286

The storm passed New England on September 24 (Boston Herald, Oct. 17, 1757, Ludlum 287 1963) and struck Nova Scotia as the Louisbourg Storm on September 25, 1757. Its arrival at Fort 288 289 Cumberland on the Nova Scotia border 200 km inland late September 22 included 'violent rain' and 'constant heavy rain' into the 23rd. Knox's journal on the 27th describes September 24-26 290 with ... 'I never saw such storms of wind and rain as we have had for some days past...' 291 292 followed by 'windy, showery and very cold' weather on the 27-28th and 'dry, cold windy weather' on the 29th, followed by frost and snow across Nova Scotia by mid-October (Knox 293 294 1769).

295 **4.0 Historical Context**

The Seven Years' War (1756-1763) arose from unresolved issues following the Treaty of Aix-la-Chappelle that ended the War of the Austian Succession (1740-1748). It began as a European conflict between Great Britain and allies and France and its allies, but soon extended to

the colonial interests of both nations in North America and India. It resulted in significant losses 299 for France including the loss of New France, now Canada, to Great Britain (Syrett, 2008). 300 Britain's overwhelming success in gaining territory at France's expense during the war led 301 France to subsequently support the secession of the American colonies in 1775. 302 Great Britain's 'Grand Plan' for the North American campaign began with John 303 304 Campbell, the 4th Earl of Loudoun, being appointed Commander-in-Chief of the British military in North America. His adversary was Louis-Joseph de Montcalm-Grozon, Marquis de Montcalm 305 306 de Saint-Veran, commander of French forces in North America. To attack Montcalm at Quebec 307 without leaving a powerful French fortress at his rear, Loudoun needed to first seize Fortress 308 Louisbourg in Nova Scotia. On May 22 to 25, 1757, troops boarded 134 transport ships in New York to rendezvous at Halifax with a fleet departing Britain under Vice Admiral Frances 309 Holbourne. Pitt's brief removal as Prime Minister delayed the fleet but his return to power with a 310 coalition government saw it depart Cork, Ireland, on May 8, 1757. The delay allowed France to 311 312 reinforce Louisbourg with three naval squadrons ahead of the British arrival. On May 23 five French battleships and a frigate under Chevalier Joseph de Beauffremont arrived from the West 313 Indies, followed on June 15 by four battleships and two frigates under Joseph Francois de Noble 314 315 du Revest from Toulon. On June 20 nine battleships and two frigates under Vice Admiral Emmanuel-Auguste de Cahideuc (Comte Dubois de la Motte) arrived from Brest. 4000 French 316 317 troops bolstered a garrison of 3200 plus 300 Acadians and Mi'kmaq warriors (McLennan, 1918, 318 Stoetzel, 2008). Holbourne's arrival at Halifax on June 30 bolstered Loudoun's force to create an 319 army of 12 000. HMS Gosport arrived on August 5 with letters intercepted from a French 320 schooner captured off Newfoundland detailing Louisbourg's reinforcement. It rendered the

attack on the fortress untenable. Loudoun returned to New York and on September 11, 1757
Holbourne sailed his fleet north to blockade Louisbourg (Fig 1).

323 5.0 The Louisbourg Storm

The British fleet cruised off the coast of Cape Breton Nova Scotia (Fig. 1) to lure the 324 325 French fleet out of Louisbourg Harbour to do battle. On September 21, the British 80-gun 326 flagship Newark noted fresh westerly gales followed by fair weather and light breezes then calm with fog on the 22nd. That day an officer on the French 28-gun frigate Fleur de Lys saw a low 327 328 mist enter Louisbourg Harbour. The mist was also seen at sea by the British Invincible until it 329 dissipated under a rising southeast breeze. Britain's Newark and France's Fleur de Lys recorded that the breeze veered to the southeast and intensified to moderate gales on September 22. The 330 Invincible recorded strengthening easterlies September 22-26 from otherwise prevailing 331 westerlies through the second half of September (Table 2). 332

	SEPT 16		SEPT 17			SEPT 18			
SW	SW	WSW	SW	W	NNW	NNW	NNW	NNW	
225	225	247.5	225	270	337.5	337.5	337.5	337.5	
	SEPT 19			SEPT 20			SEPT 21		
NNW	NE	WNW	WSW	WSW	W	W	W	NNW	
337.5	45	292.5	247.5	247.5	270	270	270	337.5	
	SEPT 22			SEPT 23			SEPT 24		
SE	SSE	SEBS	SE	SE	SEBS	SEBS	SEBS	EBS	
135	157.5	146.25	135	135	146.25	146.25	146.25	101.25	
	SEPT 25			SEPT 26			SEPT 27		
EBS	SW	W	W	W	NW	SWBW	SEBS	WBS	
101.25	225	270	270	270	315	236.25	146.25	258.75	

333

Table 2. Prevailing Winds (HMS Invincible Logbook)

335	Prevailing wind direction measured for each of three successive 8-hour watches per day and
336	azimuth equivalent on the Invincible. Storm winds, arriving September 22, 1757, off Cape
337	Breton, are shaded and in italics; two watches with easterlies not associated with the storm are
338	shaded only. Mean 250.5 (WSW) prevailing wind direction six days before and five days
339	following storm (continued westerly on 28 and 29). Mean 135 (SE) wind direction during storm.
340	Ships off St. Esprit on September 25 saw prevailing southeasterly winds last until September 26.
341	Ships south of St. Esprit including Invincible, Sunderland and Windsor faced southwesterly
342	winds on September 25.

French naval officers, expecting a storm due to the southeast winds, moored the French fleet in two lines off Royal Battery (Fig. 2) with four 2-ton anchors at the bow of each ship. The southeast winds led the British ships at sea to secure masts and naval guns, weighing as much as 3 tons apiece, anticipating a storm. On the 24th *Invincible* and *Newark* reported increasing cloud, haze and rain beginning under southeast gales.




Figure 2. Louisbourg Harbour showing the French fleet anchorage, Louisbourg Lighthouse,
Royal Battery, Battery de la Grave Guardhouse, and the southeast seawall overlain on chart
image © Canadian Hydrographic Service (2011) Chart Guyon Island to Flint Island 1:37,866
[Issued 2022-11-26]. Shoals (shaded) relative to ship hull displacements of 5.8-7.0 m (19 to 23')
give a general sense of the scale of waves and surge needed to throw battleships on shore and
destroy the southeast facing seawall.

On September 25 fresh southeast gales rose to excessive hard gales with very heavy rain. 355 356 The British *Windsor* noted heavy rain and mist and intensifying strong gales with hard squalls. 357 At 7 p.m. Sunderland faced very hard gales that rose to extreme hard gales by 10 p.m. At 12 358 a.m. Invincible faced strong gales, torrential rains and a 'great sea.' At 2 a.m. on the 25th Invincible noted an 'excessive hard gale' and 'a hurricane of wind' and mountainous waves. 359 Topsails used to control ships in severe weather were 'blown to rags.' Sunderland's main 360 staysail was torn away. Waves 'made a free passage over...' the 70-gun Devonshire and 361 362 smashed in *Lightning*'s stern. The wind tore away the 8-gun *Cruiser* sloop's mizzen mast and three sailors were swept overboard. Cruiser was 'very near foundering having been underwater 363 364 several times' and jettisoned its guns to stay afloat.

Windsor's log records extreme gales with severe squalls, heavy rain and a great sea. Canvas tarpaulins stripped off deck gratings by the wind allowed waves and rain to flood the ships which soon had up to 2.5 m (9') of water in the holds despite the pumps in full operation. *Windsor* and *Sunderland* sailed S across SSW winds. *Grafton's* three-ton 7 m (30') rudder was torn off the ship. *Invincible*'s rudder, also torn free, was only saved by its preventer chains. Sails on all the British ships at sea were torn away by the wind. Captain Bently later reported that *Invincible's* hull planking had opened and broke iron reinforcing brackets and bolts,

allowing the entire gun deck and its tens of tons of heavy naval guns to drop several inches 372 (Captain's Letters, ADM 1/1488). Sunderland's foretopmast, reinforced by ten 5 cm (2") rope 373 374 shrouds plus stays, was torn off the ship and it disappeared into the night with two sailors. Invincible was thrown onto her 'beam ends' (side), forcing it to heave overboard ten 12-pounder 375 upper deck guns and carriages, roughly twenty tons, to right the ship. *Invincible's* main yard was 376 377 ordered taken down but before it could be done the wind broke off the 38" (1 m) diameter mainmast 20' (6 m) above the deck. The falling mast tore down the foretopmast and mizzen mast 378 379 and crushed the starboard gunwale. The wreckage pulled the ship onto its side and swept sailors 380 John Guttredge and Samuel Kirby into the sea. Invincible's sailors cut the tangled mass free before it sank the ship. 381

At Louisbourg, the French military officer at La Grave Battery (Fig. 2) led his troops to 382 383 safety after the sea rose steadily above their knees (Chevalier de Johnstone, 1758). Offshore, the British 14-gun *Ferret* sloop under Francis Upton and a crew of 104 was lost with all hands. 384 385 Around 6 a.m. *Invincible* noted five British ships dangerously close to shore. *Eagle* was blown onto its beam ends and jettisoned ten upper deck guns and cut down its mizzen mast to right the 386 387 ship. Captain's foretopmast was torn away and took its two topmen. Lightning found it was 388 drifting toward offshore breakers less than 200 m away. Captain Faulkner ordered Windsor's guns jettisoned. He noted *Invincible* had lost all but its lower foremast and bowsprit. Sunderland 389 390 was swept by 'a very heavy large sea' that 'passed freely over us.' Barges lashed to the decks of 391 Windsor and Invincible were smashed and swept overboard. Sunderland cut down its main 392 topmast and threw guns overboard to right the ship. The wind snapped its 61 cm (24") diameter 393 mizzen mast as it drifted toward the offshore breakers. Anchors did not slow its drift so the 394 mainmast was cut down. Sunderland stopped close to the breakers and less than a kilometer from

shore. The 74-gun *Terrible* also stopped its drift almost at the breakers. *Eagle's* foretopmast was
cut down to lessen the strain on the ship. It sailed southward narrowly missing the breakers. *Newark* regained control after cutting the anchor cable and heaving guns overboard and barely
cleared the line of breakers. Dawn revealed a signal flag had been raised by the French fishing
village of St. Esprit to give the crews of the British ships hope (Knox Bristol Journal, November
12, 1757).

At Louisbourg the French fleet was pummeled by severe winds and waves. The 70-gun French battleship *Dauphin Royale* fired a gun in distress when its anchor cables snapped under the strain. *Dauphin Royale* collided with the 80-gun *Tonnant*, destroying its bowsprit, figurehead and cutwater, and damaged *Tonnant's* rudder and poop deck. The two ships crossed *l'Abenaquise's* anchor cables and the three entangled ships were heaved on shore at Royal Battery (Fig. 2) along with 25 merchant ships, 50 schooners and 80 small vessels, many high and dry and with many sailors drowned (McLennan, 1918).

At sea, by 10 a.m. the British fleet was dangerously close to the breakers off St. Esprit. Many sailors were certain they were doomed (Knox Bristol Journal, November 12, 1757). *Grafton* struck a rock but floated free and managed to set an anchor. *Windsor* and *Eagle* had been able to sail south of the main British fleet off St. Esprit. *Eagle's* Captain Palliser saw what he judged to be *Nottingham* or *Tilbury* near shore, within the breakers, its bow facing shore with its fore and mizzen masts gone. He also recorded that it was afloat and attempting to wear (turn) but lost sight of it in heavy rain.

Waves tore down sections of the French Fortress Louisbourg's massive southeast facing
stone seawalls. Locals brought news of lakes 10 km inland being reached by the sea. Seawater
rose to flood the streets of the Town of Louisbourg, 'something never before seen' (Chevalier de

Johnstone, 1758). Eventually the beached French battleship *Tonnant* 'floated with the tide' asthe wind veered south and then west at 11 a.m.

At sea the British warship Windsor noted the wind turned to blow from the west at 11:30 420 a.m. but had strengthened. *Eagle* recorded that the squalls had lessened by noon. On the 421 Sunderland massive waves swept sailor George Lancey from the fore yard 24 m (80') above the 422 423 keel. By 3 p.m. waves at Louisbourg fell enough that *l'Inflexible* was able to send sailors to assist other ships. French captains petitioned 74-year-old Admiral Dubois de la Motte to attack the 424 425 stricken British ships off their coast but his orders to defend Louisbourg had been met and he 426 kept his ships in port. James Johnstone, a Scot serving as a French officer, felt that five French warships if they had ventured to sea could have captured the entire British fleet (Chevalier de 427 Johnstone, 1758). This sentiment was subsequently shared by Lady Anson, daughter of a 428 confidante of Lord Newcastle with whom Pitt had formed his coalition government, in an 429 October 31, 1757 letter to the First Lord of the Admiralty, her husband George Anson (Anson, 430 431 1757). On September 27th a boat arrived at Louisbourg from St. Esprit with news that the British warship Tilbury had wrecked there with over 120 lost. Four schooners with 160 French troops 432

TIME	BRITISH AT SEA	WINDS	DESCRIPTION	FRENCHIN PORT	WINDS	DESCRIPTION
7 p.m.	Sunderland	SSE	Very hard gales and hard squalls	Fleet	SE	Moored in Louisbourg Harbour
10 p.m.	Sunderland	SSE	Extreme hard gales		1.0.0	
10 p.m.	Windsor	SSW	Very heavy rain, intensifying strong gales Hard squalls	-		
12 a.m.	Invincible	SW	Strong gales; great sea, torrential rain			
2-4 a.m	Invincible	SW	Excessive hard gale, hurricane of wind seas like mountains.	La Gr <i>a</i> ve Battery	SE	Sea level rises 3.4 m (11')
2-4 a.m.	Sunderland	SSW	topsails and staysails blown to rags	Dauphin Royale		Dauphin Royale collides with Tonnant
2-4 a.m.	Devonshire	SE	Waves swept over the ship	Tonnant		Dauphn Royale and Tonnant driven across
2-4 a.m	Lightning	SE	Waves overrun and destroy stern gallery	l'Abenaquise		l'Abenaquise anchor cable and the three
2-4 a.m.	Cruiser	SE	Waves sweep over the ship			entangled ships are thrown ashore at
		a la fara de la composition	Guns jettisoned to avoid sinking	Royal Battery		Royal Battery
			Mizzen mast torn off ship by wind	Merchant ships		25 merchant ships thrown on shore
2-4 a.m	Windsor	SSW	Severe squalls, heavy rain, great sea	Schooners		50 schooners thrown on shore
2-4 a m	fleet	100000000000000000000000000000000000000	Flooding by rain and waves	Small vessels		80 small vessels thrown on shore
	Grafton	SSE	Rudder torn off ship			
2-4 a.m	Invincible	SW	Rudder torn off ship	SE facing sea wall		Waves tear down fortress stone seawalls
2 4 0.111	manendie	SW	Hull planking sprung; hold flooding	Lakes in region		Lakes 10 km inland flooded by the sea
		SW	Gun deck brackets/bolts snapped	Louisbourg		Seawater floods the Town of Louisbourg
2-4 a.m	Sunderland	SW	Foretopmast torn off ship	Louisbourg		requiring at least 4.4-6.4 m (14.4-21') surge
2 4 u.m.	Invincible	SW	Driven onto its side by wind force			requiring acreases + 0.4 m (14.4 21) surge
		SW	Ten upper deck guns jettisoned			
		SW	Main mast snapped off which tears down			
	-	SW	foretopmast and mizzen mast			
		SW	Ship hauled onto its side by wreckage			
2-4 a.m.	Forret	SE?	Ship swallowed by the sea with all hands	-		
4-6 a.m	Invincible	SW	Near shore, sees five ships close to shore			
4-6 a.m.	Eagle	SE	Driven onto its side by wind force			
+ 0 a.m	Lagio		Jettisons guns and cuts down mizzen			
4-6 a.m	Captain	SE	Foretopmast torn off ship			
4-6 a.m	Lightning	SE	Near offshore breakers 200 m away			
4-6 a.m	Windsor	SSW	Jettisons guns to stay afloat			
4-6 a.m	Sunderland	SSW	Swept by waves			
4 0 4.111	Cundenand	0011	Barge torn off the upper deck by waves			
4-6 a.m.	Windsor	SSW	Barge torn off the upper deck by waves			
4-0 a.m	Sunderland	SSW	Driven onto its side by wind force	-		
		SW	Jettisons guns to stay afloat			
		SW	Mizzen mast torn off ship by wind			
		SW	Anchors at breakers 1 km from shore			
6-8 a.m	Terrible	SE	Anchors at breakers			
0-0 a.m	Newark	SE	Clears breakers			
10 a.m.	Grafton	SE	Strikes rock near St. Esprit			
iv a.m.	Eagle	SE	Notes Tilbury near shore at St. Esprit			
	Tilbury	SE	Aground at St. Esprit			
	fleet	SE	Most ships dangerously close to shore			
11 a.m.	Windsor	W	Winds shifted to westerlies			
12 p.m.	Eagle	W	Squalls lessening in strength			
	Invincible			finflavible	w	Manage and used approach to account other
3 p.m.	Invincible	VV to IVVV	ship under jury rig drifting seaward	l'Inflexible	٧V	Waves reduced enough to assist other ships

Table 3. Timeline of Louisbourg Storm (September 25)

435	Timeline of storm impacts on the British fleet at sea increasingly scattered by the storm and the
436	French fleet moored in Louisbourg Harbour. Relative ship locations, south to north, are blue,
437	orange, green and grey. British ships were relatively static (drifting, sailing under reefed sails or
438	at anchor) but Invincible sailed across storm winds to end up south of Windsor and Sunderland.

439 It is not known when *Ferret* sank but it had been sent ahead of the fleet prior to the storm to440 undertake reconnaissance of the French fleet at Louisbourg.

were unable to counter the heavy seas so they marched to the site across land flooded by the
torrential rain. Mi'kmaq warriors gained the wreck first but informed the shipwrecked British
they would not be harmed since the storm had brought them to their lands (Moreau St. Mery *in*McLennan, 1918).

445 **6.0 Deriving Storm Metrics**

Storm intensity is reflected in key metrics including wind speed and direction, wave
height and surge which is driven by a rise in sea level due to atmospheric pressure and sustained
storm winds and is proportional to a cyclone's intensity, translation rate and the bathymetric
gradient of the continental shelf.


450 *6.1 Estimating Storm Wind Speed*


The wind speed required to cause structural failure in masts was estimated. Virot et al. 451 452 (2016) determined the critical wind force needed to break trees of average integrity is 151 kph irrespective of species with a +9% factor for large diameter trees. 165 kph assumes structural 453 454 defects due to longer tree life offset the structural advantage of size, yet masts were chosen for 455 their lack of defects. Fir and pine trees of superior structural integrity were selectively harvested 456 for Royal Navy masts into the 1770's from North America, Great Britain and the Baltic (Lavery, 457 1984). Masts were also not free-standing (like trees) but reinforced by rigging to effectively transfer wind energy from the sails to the hull. Invincible's masts were secured by sixteen 5 cm 458 459 (2") hemp shrouds per side, each tensioned with paired deadeye blocks, the lower block in an iron band bolted to the ship's frame. Its 1 m (38") diameter lower mainmast stepped against the 460 ship's keelson rose 35.7 m (117') through two decks. Above it stood a 21.3 m (70') 51 cm (20") 461

diameter topmast and above that the 10.7 m (35') 28 cm (11") diameter topgallant mast (Lavery,
1984, 1988). To achieve the critical wind speed of 165 kph, taken as a minimum due to the
factors noted, *Invincible's* motion must be considered.


Invincible sailed SW under SE winds, but gradually encountered SW winds. *Sunderland*and *Windsor* sailed south across SSW winds while most ships of the British fleet to their north
near St. Esprit faced SSE winds. *Invincible* was among the southernmost ships (Fig. 1). It sailed
SW¹/₂W (230°) against EbS (101°) winds on September 24. During the storm its displacement
was 98 km toward 256.7° (22.5 km S; 96 km W). 6 km SE (135°) of Chedabucto Bay it faced W
(270°) winds and SE surface currents estimated at 3.49 kph based on currents of 0.97 m/s based
on currents there during SE winds from Hurricane Juan in 2003 (CBCL Report, 2015).

On September 25 to 26 Invincible sailed 159 km toward 102.75 degrees. The ship spent 472 11 hours under SE winds and another 11 hours under SW winds. The last 2 hours it drifted west 473 474 under jury rig. The strongest winds were SW (225°). Cosine Law (Figure 4) gives a wind speed 475 of 170.62 kph to achieve 165 kph at the mast on the moving vessel. The 5.62 kph difference 476 infers vessel motion played only a minor role in reaching critical force yet is still 18% of the Saffir-Simpson Category 3 wind force range. Ephemeral squalls of 40-60 kph added to sustained 477 478 winds of 170.62 kph suggests peak winds might have reached 211-231 kph. Admittedly an imperfect solution, it assumes a minimum critical force. It does not consider the inherently 479 480 superior structural integrity of masts plus their reinforcement by rigging, suggesting major 481 hurricane threshold winds (178 kph) could have been met even without considering squalls.





distribution of ship logs (*see* Table 3). *Invincible* sailed past *Windsor* and *Sunderland* during the
storm and into the SW winds they had encountered earlier.



Using Cosine Law, we solve for velocity a where α is 122.25 degrees: $a^2 = b^2 + c^2 - 2bc \cos \alpha$ $a^2 = (165)^2 + (10.13)^2 - 2 \times (165 \times 10.13) \times \cos (122.25)$ $a^2 = 27,225 + 102.62 - 2 \times (1671.45) \times (-0.5336)$ $a^2 = 27,327.62 + 1783.77$ a = 170.62 kph from 227.75 degrees (where b = 165 kph and β = 55 degrees)

497

Figure 4. Correction for Vessel Motion. Invincible drifted 159 km toward 102.75° between 498 September 25 and 26 over 24 hours. It experienced SE (11 hours), then SW (11 hours) and 499 500 finally W winds (2 hours). This solution focuses on the 11 hours the ship was under SW winds, the strongest winds closer to the center of the cyclone (Fig. 3). During elapsed hours 59-70 the 501 vessel sailed toward 102.75 under a SW wind (225°) at an average of 6.64 kph based on the total 502 displacement of 159 km toward 102.75°. The incident angle between the wind and the ship 503 displacement vectors is 122.25°. A surface current in Chedabucto Bay during SE winds from 504 505 Hurricane Juan (CBCL Report, 1995) of 0.97 m/s (3.492 kph) is assumed to be a reasonable estimate for this study. The resultant of 6.64 kph toward 102.75° indicates speed relative to 506 surface currents was 10.13 kph. Image not to scale. 507

Anticlockwise wind vectors at ship locations are tangential to concentric cyclonic wind bands.
Normal lines drawn to these vectors converge to identify the location of the eye. Interestingly
they lack the asymmetry diagnostic extratropical cyclone wind fields (Fig. 7). This process,
repeated to plot the eye location on September 26, 1757, indicates the storm crossed Cape Breton
and entered the Gulf of St. Lawrence. Even if the wind field began to collapse, the location of the
storm center suggests the system may have slowed while passing over Cape Breton Island.

514 6.2 Estimating Storm Wave Height

Sunderland's and Devonshire's upper decks were submerged after waves broke over the 515 forecastle. The 12.2 m (40') distance from the keel to the upper deck plus an estimated 3-6 m 516 517 (15-20') to break over the forecastle and tear away ship's boats lashed to the deck requires a wave height of about 18 m (60') (Lavery 1983). Lightning's stern gallery 40-50' above the keel 518 was destroyed by waves striking the ship from astern, also requiring waves of about 12.2 m 519 520 (60'). A sailor swept out of Sunderland's fore yard by a wave necessitates a wave of about 25-30 521 m (80-90'). While carrying considerable uncertainty, these examples provide estimates of significant and maximum wave heights. Waves sufficiently large to tear down stone seawall 522 523 rampart of Fortress Louisbourg are consistent with these estimates, as are waves capable of 524 reaching inland lakes. Descriptions of the sea state in Louisbourg Harbour by French naval 525 officers resulting in extensive damage to ships and boats suggests waves much larger than any 526 recorded in modern times even though wave energy from the southeast would have been partly attenuated by shoals (Fig. 2). 527

528 On September 26-28, 1818, the American frigate *USS Macedonian* met a hurricane off 529 Bermuda (35°N 53°W) and suffered damage nearly identical to *HMS Invincible* in 1757 from 530 waves of 12 m (40') (Saegesser 1970). The dates appear to coincide with Chenowith's (2006)

'Final Storm Number 253' listed as a hurricane in Table IV. Damage to the ship closely parallels 531 that described for the 1757 hurricane except that line of battle ships had a much heavier 532 construction than a frigate. Saegesser (1970) provides a very detailed account based on the ship's 533 log and ancillary damage reports, and notes that in the same storm the Dutch brig De Hoope lost 534 all topmasts and spars, the brig Ann from Nova Scotia was abandoned at sea, the brig Mary from 535 536 Bristol was overturned, the ship Catherine Dawes from Philadelphia sank and a Baltimore schooner and a Nantucket whaler were both dismasted. *Invincible's* substantially more robust 537 538 build than the frigate *Macedonian* implies larger, more powerful waves caused its damage. 6.3 Estimating Surge Height 539

540 *6.3.1 Surge at Louisbourg Harbour*

A Parks Canada coastal erosion study at Fortress Louisbourg National Historic Site 541 542 revealed iron mooring rings set in the remains of a seawall. Modern high tide compared to these rings established historical high tide 0.90 m (3') of sea level rise since 1757 (Duggan, 2010). La 543 Grave Battery (Fig. 2) is 2.0 m (6.6') above sea level (asl; Google Earth mid-tide datum), so sea 544 545 level rise plus flooding to sentries' knees (0.5 m) yields a 3.4 m (11') mid-storm surge. Historic buildings along the waterfront (Fig. 2; 45°53'33.57" N 59°59'07.89" W) are 5 m (16.4') asl 546 while the first street, Rue Royale, is 7 m (22.9') asl. Seawater flooding the town streets at the 547 lowest levels and adjusted for sea level rise indicates 5.9 m (19.4') to 7.9 m (21.4') of surge. 548 Tonnant 'floated with the tide' when the wind veered south at 11 a.m. on September 26 (Fleur de 549 550 Lys log in McLennan, 1918). Louisbourg's 12-hour tidal cycle and assuming low tide around 10 a.m. gives a high tide at 4 a.m. coinciding with storm landfall and creating a storm tide (Fig. 3). 551 Backing out the 1.5 m (5') tidal range gives a 4.4-6.4 m (14.4-21') peak surge, consistent with 552 553 the earlier surge of 3.4 m (11') at La Grave.

554 6.3.2 Surge at St. Esprit (Tilbury Wreck)

555 *HMS Tilbury* was a 58-gun square-rigged warship lost on the coast in the storm. *Eagle's* 556 captain saw either *Tilbury* or *Nottingham* shoreward of the breakers near St. Esprit, 45 km south 557 of Louisbourg. It was deduced to have been *Tilbury* since *Nottingham* survived the storm with a 558 different array of masts than seen on this ship. 'Wreck' appears on a 1776 chart (Fig. 5). Storm 559 (2002) used Zinck's (1975) image of an 18th Century 6-pounder British naval gun at 'Tilbury 560 Rocks' to view Tilbury's wreckage in 4 m (15') from a boat in 1969.





Figure 5. Excerpt from a historic chart of Cape Breton Island showing the general St. Esprit
study area and *HMS Tilbury* wreck site, from Mowat (1776), depicted in Figs. 6a,b.The faint
dotted line right of Barnsley Lake, named for Tilbury's captain, marks a parish boundary.

The historic navigation chart (Fig. 5) showed parish boundaries marked by fieldstone 565 walls of historic St. Esprit (Fig. 6a, b) which helped identify the line of offshore breakers 566 567 described in British naval logs. A draft hydrographic chart (Hanson, 1954) was digitized and gridded with missing data interpolated. Paired depths and locations were entered in a spreadsheet 568 and a grid-plot of local bathymetry supported a marine proton magnetometer survey of Tilbury 569 570 Reef isobaths following best practices for submerged archaeological sites (Cornwall Council Report 2010-R012). Dipole targets investigated by divers led to locating a mid-18th Century 6-571 572 pounder British naval gun in situ in 3 m (10') which was 2.1 m (7') in 1757, near the site of the 573 6-pounder on shore, both interpreted to be from Tilbury's forecastle. In 1757 Tilbury was observed at the time as 'bow in' near shore, landward of the breakers and 'attempting to wear' 574 (turn). It was in water sufficiently deep for its 18' displacement as it was, at the time, afloat and 575 576 under sail. Adding in the hydrographic survey datum offset of 0.6 m (2') between lowest low tide at St. Esprit and the Google Earth WGS84 (World Geodetic Standard 1984) mid-tide datum for 577 578 Louisbourg suggests a minimum 4.0 m (13') surge at St. Esprit. Post-storm relaxation flow stranded the *Tilbury* (Fig. 6b) allowing native warriors to reach it. 579





- to 45°38'31.61" N 60°26'05.28" W) corresponds to Fig. 6b. Dashed line is bedrock reef
- 584 (breakers). Image © Google Earth Pro 7.3.6.9345 (2022) St. Esprit, Nova Scotia Canada.
- 585 $45^{\circ}38'31.54"$ N $60^{\circ}27'37.76"$ W Eye alt 4.50 km TerraMetrics © 2023 MaxarTechnologies ©
- 586 2023.





Figure 6b. Bathymetry of *Tilbury* wreck site at lowest low water adjusted for 1757 relative sea level (solid line) and minimum surge (dashed line) needed to float *Tilbury*. Coastal retreat of 27 m (90') calculated from historic sea level gives the 1757 shoreline. Topographic and bathymetric data were kept in Imperial units for comparison to *Tilbury*'s displacement. X and X' of this block diagram correspond to the same GPS positions on the areal chart in Fig. 6a.



593

Figure 7. Eye location and estimated translation speed. Plots of wind vectors on September
25 at: (1) Invincible, (2) Windsor and Sunderland, (3) Newark and most of the British fleet,
French ships at Louisbourg Harbour. Normal lines taken to wind vectors cluster at the eye.
Estimated translation rates are based on the storm off North Carolina, New England and
Chedabucto Bay on the dates shown, showing increased translation typical of midlatitude
cyclones, yet a similar wind vector reconstruction for September 26 gives an eye location
entering the Gulf of St. Lawrence, suggesting the system slowed over Cape Breton after landfall.

601 **7.0 Modern Analogs**

On September 29, 2003, Hurricane Juan struck Nova Scotia with peak winds of 165 kph (Category 2), a significant wave height of 10 m (32'), a maximum wave height of 19.9 m (65') and a surge at landfall near Halifax of 1.5 m (4.9') (Lixion, 2003). On January 20-22, 2000, an extratropical meteorological 'superbomb' that developed off Cape Hatteras struck Nova Scotia with peak winds of 25-30 m/s (90-108 kph), a significant wave height of 12 m (39'), a peak wave height of 19 m (62') to 23 m (77') at drilling rigs near Sable Island (JD pers. obs.) and a 1.4 m
(4.6') surge at landfall near St. Esprit (Lalbeharry et al., 2009). Both cyclones produced similar
sea states and surge which can be compared to the Louisbourg Storm. On September 24, 2022,
Category 3 Hurricane Fiona began extratropical transition as it crossed the Scotian shelf. A cold
trough over Nova Scotia directed its landfall to the Canso Peninsula. Winds of 140 kph in Nova
Scotia reached 177 kph in Newfoundland and Labrador. Significant and maximum wave heights
were 17 m (56') and 30 m (98') and surge reached 2.4 m (8').

In 1969 Category 5 Hurricane Camille generated a 7.3 m (24') storm tide from 1.8-3.0 m

615 (6-10') surge (U.S. Department of Commerce Environmental Science Services Administration

1969) while Category 5 Katrina in 2005 produced a storm tide of 8.2 m (27') (Knabb et al.,

617 2023). Hurricane Laura (Category 4) in 2020 had a peak 5.2 m (17.2') surge (Pasch et al., 2021)

and a 2.7-4.0 m (9-13') spanning 130 km from Beaumont to Lake Arthur, Texas. In 2018

Hurricane Dorian (Cat 5) slowed to 2 kph over the Bahamas creating an 8.5 m (28') surge (Avila

et al., 2020). Surge from these major hurricanes cannot be readily compared to storm strikes in

621 Nova Scotia due to different coastal bathymetry but they allow a general comparative

622 benchmark.

Hurricane Juan's translation speed before landfall was 1-5 m/s (4-18 kph). Compared to North Atlantic hurricane translation rates of 17.7-19.3 kph (11-12 mph) the Louisbourg Storm slowing from 31 kph over water to 4.6 kph after landfall between September 25-26 may have enhanced surge height, similar to Dorian's impact on the Bahamas as it slowed, resulting in the exceptional surge height at Louisbourg. Prevailing westerlies returned after the storm. The key metrics of wind speed, wave height and surge are summarized in Table 4.

Storm	Year	Date	Peak Wind (kph)	Significant Wave Height (m)	Peak Wave Height (m)	Surge (m)
Louisbourg	1757	25-Sep	171 - 231	12+	25-30	4.4 – 6.4
Unnamed	2000	22-Jan	90 - 108	12	19	1.4
Juan	2003	27-Sep	160 - 165	10	20	1.5
Fiona	2022	24-Sep	155 - 179	17	30	2.4

629

Table 4. Louisbourg Storm Comparison to Modern Nova Scotia Landfalling Storms. The 630 Louisbourg Storm, a winter extratropical storm in 2000, Juan (Category 2 hurricane at landfall), 631 and Fiona, an extratropical cyclone that transitioned from a Category 3 hurricane over the 632 Scotian Shelf crossed the same coastal bathymetry with similar translation rates to strike Nova 633 634 Scotia. Sustained winds for the Louisbourg Storm exceeded 171 kph based on the critical force needed to break main and mizzen masts and break away and carry off topmasts and may have 635 reached 231 kph with squalls. Peak wind is presented as the range between sustained threshold 636 637 and maximum wind speeds.

638 8.0 Discussion

639 Metrics derived from historical data captured during the Louisbourg Storm of 1757 indicate its intensity surpassed any modern (post-1851) Atlantic cyclones striking the same 640 641 region. Historical records show the Louisbourg Storm originated in the tropics to pass Florida, the Carolinas and New England to strike Nova Scotia on September 25, 1757. It developed at 642 643 the height of hurricane season under an optimal NAO index and ENSO conditions for Atlantic hurricanes to form and track up the Atlantic coast of North America into the northern 644 midlatitudes. The already low NAO index also decreases later in the season and may have helped 645 stay over the Gulf Stream which allowed it to intensify into higher latitudes. Its devastating 646 impact on the British and French fleets and coastal infrastructure was due to an unusually violent 647

release of energy over coastal waters. Longer, colder LIA winters skewed mean average 648 temperature profiles but a UK and European heat wave in Europe in 1757, extreme even by 649 modern standards, shows seasonal temperature variability could contribute to warmer SSTs and 650 fuel tropical cyclones in the LIA. A strong correlation between SST and tropical cyclone 651 frequency (Vecchi and Knutson, 2008) suggests that the LIA's cooler SSTs could see fewer 652 653 storms per year. Mean-annual temperature data limited by temporal resolution limitations likely mask peak temperatures that likely existed over smaller areas for shorter periods but historical 654 records clearly show tropical cyclones developed even during the coldest part of the LIA. 655

The large number of British warships scattered along Cape Breton's coast by the Louisbourg Storm provided a spatial resolution of wind vectors not normally available in storm reconstructions. It was partly facilitated by ships sailing across storm winds to avoid being driven ashore. The proximity of many British ships to shore and the severe surge and wave action at Louisbourg led many contemporary naval authorities of both nations to fear the catastrophic loss of the British and French fleets and 20 000 sailors. Only the reversal of wind direction at the last minute as the eye of the storm passed prevented a disaster.

Wind speed is the key metric used in the Saffir Simpson scale to characterize the intensity 663 of modern cyclones. Engineering models are a standard method of determining the force 664 required to trigger structural failure in materials. Trees lacking defects that negate size advantage 665 666 were preferentially selected for masts and so likely required higher wind speeds for structural failure. Rigging not only reinforced masts but redirected wind energy to the hull. Both factors 667 imply that the wind speed estimate of 171 kph determined for *Invincible* to achieve 165 kph at 668 669 the mast is an underestimate. Sustained winds likely exceeded the 178 kph (Cat 3) major hurricane threshold even without considering squalls of 40-60 kph. Extreme winds are reflected 670

in topmasts (along with shrouds and stays) not only being torn off two British ships but being 671 carried off (with sailors) instead of falling to the deck. British ship positions were triangulated 672 against known coastal landmarks which provided greater accuracy in the distribution of wind 673 vectors. Superimposing Invincible's location and the wind vectors that identify the eye location at 674 the height of the storm suggests severe damage was a consequence of proximity to the eye which 675 is the location of a cyclone's strongest winds (Figs. 1,3,7). Peak damage and squalls above 676 hurricane winds lasted 9 hours and hurricane force winds noted by the British ships lasted 15 677 678 hours as the center of the storm passed the coast (Fig. 3). In comparison, Hurricane Juan crossed 679 Nova Scotia in only 3 hours while Fiona crossed the province in under 6 hours, supporting the interpretation derived from eye locations (Fig. 7) that the Louisbourg Storm slowed over land, 680 possibly due to a blocking cold air high. The British warship Tilbury was driven into water 681 depths at St. Esprit it could navigate only under a storm tide. Tidal reversal mid storm stranded 682 the ship near shore (Figs. 3, 6a,b). 683

684 Wind plots also show that the southernmost ships of the British fleet faced southwest winds from the lower right quadrant of the hurricane. British ships to the northeast near St. Esprit 685 faced southeast winds. The French fleet in Louisbourg Harbour also faced southeast winds and 686 687 an anomalously high storm surge which allowed massive waves to drive ships on shore while the surrounding region was flooded by torrential rains, all consistent with the front right quadrant of 688 689 the hurricane where the most severe impacts are felt. There was no suggestion that the air of the storm was cold, but westerlies following the storm were described at Fort Cumberland as very 690 cold and dry. A table of wind directions for the second half of September 1757 (Table 2) shows 691 that, with the exception of the storm, prevailing winds appear to have been continental 692 westerlies. 693

Modern analogs show strong similarities in significant and maximum wave height, but 694 interpreted wind speeds for the Louisbourg storm are greater than those of Category 2 hurricane 695 Juan, a winter extratropical 'superbomb' in 2000, and the extratropical cyclone Fiona in 2022. 696 Surge measured at three locations is consistent with the scale of surge from major hurricanes in 697 the Gulf of Mexico and Caribbean. The 1757 surge greatly exceeds that of modern analogs that 698 699 crossed the same bathymetry with similar translation speeds. This consistent basis of comparison of surge height, closely linked to storm intensity, shows the Louisbourg Storm had an intensity 700 701 far beyond a Category 2 system and was equal to a major hurricane. Surge calculated 702 independently for the lowest streets of the historic town of Louisbourg, Battery de la Grave and the *Tilbury* wreck at St. Esprit were also consistent. Unlike the modern analogs, storm surge at 703 Louisbourg reflects conditions one hundred kms from landfall (Fig. 7). 704

The climatology of tropical cyclones on North America's eastern seaboard renders the 705 simple attribution of 'tropical' vs. 'extratropical' problematic. It is unlikely that a fully tropical 706 707 system with wind speeds equal to a Category 4 hurricane to strike Nova Scotia. Atlantic tropical cyclone extratropical transition is triggered by the interaction of autumn continental westerlies 708 709 pushing strongly baroclinic air eastward toward intensifying tropical cyclones tracking north into 710 the higher midlatitudes of the North American eastern seaboard when SSTs peak in late September into October. This is consistent with climatic drivers interpreted by Dezileau et al. 711 712 (2011) and Jackson et al. (2019) to explain historic European LIA storminess. Storm intensity 713 normally drops following extratropical transition, but not always (Hart and Evans, 2001). The National Hurricane Center (NHC) uses sea surface temperatures plus storm asymmetry in 714 satellite images to indicate the degree of transition. Hart and Evans (2001) also found that 'the 715 NHC declaration (of extratropical transition) typically occurs early in the 1 to 2-day period ... 716

when the storm is just beginning to lose its tropical characteristics.' This is not easy to assess for 717 the Louisbourg Storm whose energy release may have occurred over a short period. The lack of 718 eye asymmetry of the storm at landfall on September 25 based on the convergence of normal 719 lines to vectors at ship locations (Fig. 7) suggests it may have had largely tropical characteristics 720 721 at landfall. It leads to questioning at what point was it 'tropical' (hurricane) vs. 'extratropical' 722 given the NHC's 1 to 2-day range? It was likely both in the coastal zone. The storm's large size is indicated by its winds first being recorded on September 22 by both the British and French 723 fleets at Cape Breton on the same day it struck the British frigate Winchelsea off North Carolina, 724 725 1350 km to the southwest. This may have enabled it to continue to draw tropical energy from the Gulf Stream as it neared the Nova Scotia coastline. Hart and Evans's (2001) extratropical 726 727 climatology based on an analysis of all Atlantic tropical cyclones over a century. It shows that systems can continue to see tropical intensification north of strongly baroclinic conditions that 728 trigger transition, resulting in an explosive release of energy and post-transition intensification. 729 730 Their analysis shows this typically involves hurricanes from south of 20 N that retained an intensely tropical character into the higher midlatitudes. In fact, their analysis of past Atlantic 731 hurricanes shows that the region most conducive to this process in the entire North Atlantic basin 732 733 lies immediately south of Cape Breton, Nova Scotia, where the Louisbourg Storm was in 1757.

734 **9.0 Conclusions**

In 1757 continental westerlies, colder and earlier than today in the LIA, juxtaposed a cold higher pressure air mass against a large, intensifying hurricane approaching Cape Breton. The resulting explosive release of energy gave the Lousibourg Storm its highly destructive power. Its unusual intensity required only an incremental change in the accepted climatology of Atlantic cyclone extratropical transition, that being the early arrival of colder LIA continental westerlies

driving a steeper temperature gradient. The storm slowed over Nova Scotia as it encountered a 740 blocking high, indicated by the short distance between eye locations on September 25 and 26, as 741 well as by the duration of hurricane force winds (15 hours) over the coast, which may have been 742 enhanced by the storm's large diameter. The slowing storm drove an unusually high surge at high 743 tide. Tidal reversal stranded the *Tilbury* close to the historical shoreline. Fall westerlies arriving 744 745 earlier in the LIA would have expanded southward sooner and allowed an intensifying hurricane to enter a zone more baroclinically favourable for transition. In the future, instead of an earlier 746 arrival of colder continental westerlies in fall, a warming North Atlantic could drive tropical 747 748 intensification in to higher latitudes later into the autumn to trigger increasingly destructive storms over coastlines that have seen a meter of sea level rise and extensive coastal growth since 749 the Louisbourg Storm nearly rewrote history two and a half centuries ago. It is a reminder that 750 the past can inform the present, and the future. 751

752 Acknowledgements

The authors would like to thank William Pretel and Antoine LaChance for constructive review comments on the manuscript. Research assistance was provided by Cambria Huff (Dalhousie), John Allison (UK), the National Archives (UK) and the Public Archives of Nova Scotia. Tony Sampson, commercial diver, offshore survival instructor and owner/operator of Salty Dog Sea Tours, and Steve Jennex, Dave Murphy, Steve Dugas and Dana Sheppard of Zodiac Divers coordinated marine operations and underwater exploration.

759 Funding

The authors declare that no funds, grants, or other supports were received during the preparation
of this manuscript and that they have no financial or proprietary interests in any material
discussed in this article.

References

764	Anson, Lady. Letter of October 31, 1757 from Lady Anson to George Anson, First Lord of the
765	Admiralty, British Museum Collections, London UK, Add MSS 35,376 f. 145, 1757.
766	ADM 1/481 Letters from Commanders in Chief North America 1755-1760. (Charles Holmes)
767	The State and Condition of His Majesty's Ships and Sloops under my Command at New
768	York between 3rd of May 1757 and 9th following, The National Archives, UK, 1757.
769	ADM 1/481 Letters from Commanders in Chief North America 1755-1760. (Frances Holbourne)
770	Newark at sea 28 September, The National Archives, UK, 1757. [Letter to the Admiralty
771	outlining his squadron's inability to continue operations and the need to refit]
772	ADM 1/481 Letters from Commanders in Chief North America 1755-1760. (Frances Holbourne)
773	Newark at sea 28 September, The National Archives, UK, 1757 [list of damage to ships
774	sustained in the gale]
775	ADM 1/481 Letters from Commanders in Chief North America 1755-1760 (Frances Holbourne)
776	Newark at Sea 30 September, The National Archives, UK, 1757.
777	ADM 1/481 Letters from Commanders in Chief North America 1755-1760. Newark at Halifax
778	14 October, The National Archives, UK, 1757. [A letter from Frances Holbourne to the
779	Admiralty outlining the state of the squadron and the enemy's ships at Louisbourg]
780	ADM 1/1488 Captain's Letters 1757 (Bently, Jonathon). An account of the damages received on
781	board His Majesty's Ship Invincible in the hurricane on the 25th September, The
782	National Archives, UK, 1757

783	ADM 1/2294 Captain's Letters 1757 (Palliser, Hugh). Sunday 25th September 1757 at 2 am. An
784	account of the Eagle's situation and of the damages she received in the late gale of wind.
785	The National Archives, UK, 1757.
786	ADM 1/2294 Captain's Letters 1757 (Palliser, Hugh). Eagle at sea 30 September, 1757. Account
787	of the Condition of His Majesty's Ship Eagle, The National Archives, UK, 1757.
788	ADM 8/31 Admiralty List Books 1756-1757 Halifax Station, The National Archives, UK, 1757.
789	ADM 8/32 Admiralty List Books 1757-1758 Halifax Station, The National Archives, UK, 1757.
790	ADM 51/409 Captain's Log HMS Grafton (1755 Feb 7–1764 Jun 24), The National Archives,
791	UK,1757.
792	ADM 51/471 Captain's Log HMS Invincible (1756 Aug 7–1758 Mar 6), The National Archives,
793	UK, 1757.
794	ADM 51/633 Captain's Log HMS Newark (1755 Jul 31–1760 Apr 1), The National Archives,
795	UK, 1757.
796	ADM 51/921 Captain's Log HMS Sunderland (1756 Nov 15–1759 Feb 23), The National
797	Archives, UK, 1757.
798	ADM 51/1075 Captain's Log HMS Windsor (1755 Jun 12–1759 May 20), The National
799	Archives, UK, 1757.
800	ADM 52/578 Master's Log HMS Eagle (1757 Apr 28–1759 Mar 3), The National Archives, UK,
801	1757.
802	ADM 52/819 Master's Log HMS Captain 1756 May 21–1760 Feb 21), The National Archives,
803	UK, 1757.

804	ADM 52/1105 Master's Log HMS Winchelsea (1755 Dec 29–1757 Dec 31), The National
805	Archives, UK, 1757.
806	ADM 52/1064 Master's Log HMS Terrible (1756 Feb 22–1758 Sep 30), The National Archives,
807	UK, 1757.
808	Barriopedro, D., Gallego, D., Alvarez-Castro, C., Garcia-Herrera, R., Wheeler, D., Pena-Ortiz,
809	C., Barbosa, S.: Witnessing North Atlantic westerlies variability from ships' logbooks
810	(1685-2008). Climate Dynamics. Vol. 43, 939-955. DOI 10.1007/s00382-013-1957-8,
811	2014.
812	Bertler, N., Mayewski, P., Carter, L.: Cold conditions in Antarctica during the Little Ice Age-
813	Implications for abrupt climate change mechanisms. Earth and Planetary Science
814	Letters, Vol. 308(1-2), 41-51, 2011.
815	Blake, N., and Lawrence, R.: The Illustrated Companion to Nelson's Navy. Great Britain:
816	Chatham Publishing. p. 144, 1999.
817	Boston Herald, Oct. 17, 1757 [account of hurricane]
818	British High Commission, Ottawa, Canada Note 26-06: Letter advising the Minister of Foreign
819	Affairs of the British Government's position respecting the sovereign protection of the
820	HMS Fantome and HMS Tilbury shipwrecks, 2006.
821	Canadian Hydrographic Survey Chart: Guyon Island to Flint Island (2011) 1:37,866 [Issue Date
822	2022-11-26] 2022.
823	CBCL Draft Report Bear Head LNG Terminal Metocean Study. 12 pp with Appendices, 2015.
824	Chenowith, M.: Reassessment of Historical Atlantic Basin Tropical Cyclone Activity, 1700-
825	1855. Climatic Change Vol. 76, 169-240, 2006.

- Corbett, J.: England in the Seven Years' War: A Study in Combined Strategy. 2 Vols. London:
 Longmans, Green and Company. 407 pp, 1907.
- 828 Cornwall Council: Developing Magnetometer Techniques to Identify Submerged Archaeological
 829 Sites. Cornwall Council Report 2010-R01. 221 pp, 2010.
- 830 Cronin, T., Hayo, K., Thunell, R., Dwyer, G., Saenger, C., Willard, D.: The medieval climate
- anomaly and Little Ice Age in Chesapeake Bay and the North Atlantic Ocean.
- Palaeogeography, Palaeoclimatology and Palaeoecology. Vol. 297, 299-310, 2010.
- B33 Dezileau, L., Sabatier, P., Blanchemanche, P., Joly, B., Swingedouw, D., Cassou, C., Castaings,
- J., Martinez, P., Von Grafenstein, U.: Intense storm activity during the Little Ice Age on
 the French Mediterranean coast. Palaeogeography, Palaeoclimatology, Palaeoecology.
 Vol. 299, 289–297, 2011.
- Buggan, R.: Coastal Heritage Planning at the Fortress of Louisbourg Planning it Out *in*Archaeology in Nova Scotia: 2010 News. Halifax: Nova Scotia Museum Collections
 Unit, 1-8, 2010.
- Donnelly, J., Bryant, S., Butler, J. and Dowling, J.: 700 yr. sedimentary record of intense
 hurricane landfalls in Southern New England. Geological Society of America Bulletin
 Vol. 113, 714-727, 2001.
- Finck, P.: A Geological and Coastal Vulnerability Analysis of Point Michaud Provincial Park,
 Richmond County, Nova Scotia. Nova Scotia Natural Resources Open File Report ME
 2015-003. 25 pp. 2015.
- Garcia, R., Diaz, R., Herrera, G., Eischeid, M., Prieto, E., Hernandez, L. Gimeno, F., Duran, R.
 and Bascary, A.: Atmospheric circulation changes in the tropical Pacific inferred from the

848	voyages of the Manila Galleons in the sixteenth-eighteenth centuries, Bulletin of the
849	American Meteorological Society. Vol. 82, 2435–2455, 2001.
850	Garcia-Herrera, R., Wilkenson, C. Koek, F., Prieto, M., Jones, P. and Koek, F.: Description and
851	general background to ships' logbooks as a source of climatic data. Climatic Change.
852	Vol. 73, 13-36, 2005a.
853	García-Herrera R., Können G., Wheeler, D., Prieto, M., Jones, P., Koek, F.: CLIWOC: a
854	climatological database for the world's oceans 1750- 1854. Climatic Change. Vol. 73, 1-
855	12, 2005b.
856	Gebbie, G.: Atlantic warming since the Little Ice Age. Oceanography. Vol. 32, 220–230, 2019.
857	Gurgis, J. and Fowler, A.: A history of ENSO events since A.D. 1525: implications for future
858	climate change. Climatic Change. Vol. 92, 343-387, 2009.
859	Hanson, R: St. Peter's Island to Kelpy Cove, Southeast Coast, Cape Breton Unpublished 1" =
860	3000' field sheet. Canadian Hydrographic Survey, 1954.
861	Hart, R. and Evans, J.: A climatology of extratropical transition of Atlantic tropical cyclones.
862	Journal of Climate. Vol. 14, 546-564, 2001.
863	Jackson, D., Costas, S., Guisado-Pintado. E.: Large-scale transgressive coastal dune behaviour in
864	Europe during the Little Ice Age. Global and Planetary Change. Vol. 175, 82-91, 2019.
865	Johnstone, (James) Chevalier: The campaign of Louisbourg, 1750-'58 [microform]: a short
866	account of what passed at Cape Breton, from the beginning of the last war (1750) until
867	the taking of Louisbourg, by the English, in the year of Our Lord, 1758. Memoirs of the
868	Chevalier de Johnstone Vol. 3 of 3. 33 pp, [Translated in 1871 by Charles Winchester],
869	1758.

- **870** Jones, P. and Mann, M.: Climate over past millennia. American Geophysical Union.
- 871 https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2003RG000143. 42 pp, 2004.
- Keigwin, L.: The Little Ice Age and medieval warming period in the Sargasso Sea. Science, Vol.
 274, 1503-1508, 1996.
- Keigwin, L., Sachs, J., and Rosenthal, Y.: A 1600-year history of the Labrador Current off Nova
 Scotia. *Climate Dynamics*. Vol. 21, 53–62, 2003.
- Knabb, R. Rhone, J. and Brown, D.: Tropical cyclone report Hurricane Katrina 23-30 August
 2005. National Hurricane Center Tropical Cyclone Report. 43 pp,
- https://nhc.noaa.gov/data/tcr/AL122005_katrina.pdf, 2023. [accessed 2023-09-30]
- 879 Knox Bristol Journal, November 12, 1757. [hurricane survivor account]
- 880 Knox, J. (Captain): Historical journal of the campaigns in North America for the years 1757,
- 881 1758, 1759 and 1760. Volume 1 of 3. London: W. Johnston (Ludgate Street), and Dodsly,
 882 J. (Pall Mall), 49, 1769.
- Kreutz, K., Mayewski, P., Meeker, L., Twickler, M., Whitlow, S., and Pittalwa, I.: Bipolar
 changes in atmospheric circulation during the Little Ice Age. Science. Vol. 277 (5330),
 1294-1296, 1997.
- Lalbeharry, R., Bigio, R., Thomas, B. and Wilson.: Numerical simulation of extreme waves
 during the storm of 20-22 January 2000 using winds generated by the CMC weather
 prediction model. Atmosphere-Ocean. Vol. 47, 99-122, 2009.
- Lamb, H.: Climate, history, and the modern world. Methuen: New York. 387 pp, 1982.
- Landsea, C. Anderson, C., Charles, N., Dunion, J., Clark, G., Fernandez-Partag´as, J.,
- 891 Hungerford, P., Neumann, C., and Zimmer, M.: The Atlantic hurricane database re-
- analysis project: Documentation for the 1851–1910 alterations and additions to the

893	HURDAT database, in Murnane, R. J. and Liu, KB. (eds.), Hurricanes and Typhoons:
894	Past, Present, and Future, Columbia University Press, 177–221, 2004.
895	Lavery, B.: The Ship of the Line; Volume 1 – Development of the Battlefleet 1650-1850.
896	London: Conway Maritime Press. 224 pp, 1983.
897	Lavery, B.: The Ship of the Line; Volume 2 – Design, Construction and Fittings. London:
898	Conway Maritime Press. 191 pp, 1984.
899	Lavery, B.: The Royal Navy's First Invincible. Portsmouth, United Kingdom: Invincible
900	Conservations Limited – Burgess and Son (Abingdon) Limited. 119 pp, 1988.
901	Lixion, A.: National Hurricane Center Tropical Cyclone Report – Hurricane Juan. National
902	Hurricane Center Report – Hurricane Juan. https:// www.nhc/noaa.gov/data/tcr/
903	AL152003_Juan.pdf 11 pp, 2003 (Revised 2012) [accessed 2023-9-30]
904	Lixion, A., Stewart, S., Berg, R. and Berg, A.: National Hurricane Center Tropical Cyclone
905	Report – Hurricane Dorian. https://www.nhc.noaa.gov/data/tcr/AL052019_Dorian.pdf 74
906	pp, 2020. [accessed 2023-9-30]
907	London Magazine, November, 563-564, 1758.
908	The London Chronicle, July 23-26, 1757.
909	Lamb, H.: Historical storms of the North Sea, British Isles and Northwest Europe. London:
910	Cambridge University Press. 204 pp, 1991.
911	Ludlum, D.: Early American Hurricanes 1492-1870. American Meteorological Society. 198 pp,
912	1963.
913	Mann, M.: Little Ice Age in The Earth System: Physical and Chemical Dimensions of Global
914	Environmental Change. MacCracken, M and Perry, S. (eds.) Encyclopedia of Global

915 Environmental Change. Chichester, UK: Wiley and Sons Ltd., 504-509, 2002.

- 916 Matthes, F..: Report of Committee on Glaciers, April 1939. Transactions, American Geophysical
 917 Union. Vol. 20, 518, 1939.
- Mazzarella, A. and Scafetta, N.: The Little Ice Age was 1.0-1.5 °C cooler than current warm
 period. Climate Dynamics. Vol. 51, 3957-3968, 2018.
- McLennan, J.: Louisbourg from its foundation to its fall. Sydney, Nova Scotia: Fortress Press
 (1969 Reprint), 207-210, 1918.
- 922 Miller, G., Geirsdóttir, Á., Zhong, Y., Larsen, D., Otto-Bliesner, B., Holland, M., Bailey, D.,
- 923 Refsnider, K., Lehman, S., Southon, J., Anderson, C., Björnsson, H., and Thordarson, T.:
- Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean
- feedbacks. Geophysical Research Letters. Vol. 39, DOI:10.1029/2011GL050168, 5 p,
- 926 2012.
- 927 Mowat, H., Lieutenant *in* Holland, S.: A plan of the island of Cape Breton reduced from Captn.
- Holland's Survey. [soundings and naval observations were taken by Lieut. Henry Mowat,
- 929 commander of His Majesty's armed ship Canceaux and the officers of the ship under his930 direction], 1776.
- 931 Nature Notes, 24 August, 415, 1882. [European heat wave of 1757]
- 932 Oliva, H., Peros, M., Viau, A., Reinhardt, E., Nixon, F. and Morin, A. A multi-proxy
- 933 reconstruction of tropical cyclone variability during the past 800 years from Robinson
- 934 Lake, Nova Scotia, Canada. Marine Geology. Vol. 406. 84-97, 2018.
- Oliver, J. and Kington, J.: The usefulness of ships' log-books in the synoptic analysis of past
 climates. Weather. Vol. 25 (12), 520-528, 1970.

937	Pasche, R., Berg. R., Roberts, D. and Papin, P. National Hurricane Center Tropical Cyclone
938	Report – Hurricane Laura (AL132020) August 20-29 2020. 75 pp,
939	https://nhc.noass.gov/data/tcr/AL132020_Laura.pdf, 2021 [accessed 2023-9-30]
940	Poey, A.: A chronological table, comprising 400 cyclonic hurricanes which have occurred in the
941	West Indies in the North Atlantic within 362 years, from 1493 to 1855. Journal of the
942	Royal Geographical Society Vol. 25, 291–328, 1855.
943	Richey, J., Poore, R., Flower, Benjamin P., Quinn, T., and Hollander, D.: Regionally coherent
944	Little Ice Age cooling in the Atlantic Warm Pool. Geophysical Research Letters. Vol. 36,
945	L21703, doi:10.1029/2009GL040445, 1-5, 2009.
946	Ruffman, A.: The multidisciplinary rediscovery and tracking of 'The Great Newfoundland and
947	Saint-Pierre et Miquelon hurricane of 1775.' The Northern Mariner/Le Marin du Nord.
948	Vol. 1, 11-23, 1996.
949	Saegasser, L.: The U.S.S. Macedonian and the hurricane of 1818. Proceedings of the U.S. Naval
950	Institute. https://www.usni.org/magazines/proceedings/1970/january/uss-macedonian-
951	and-hurricane-18181970, 1970. [Accessed 2023-9-30]
952	Saenger, C., Cohen, A., Oppo, D., Halley, R. and Carilli, J.: Surface-temperature trends and
953	variability in the low-latitude North Atlantic since 1552. Nature Geoscience, 492-495,
954	2009.
955	Sicre, M., Jalali, B., Martrat, B., Schmidt, S., Bassetti, M., Kallel, N.: Sea surface temperature
956	variability in the North Western Mediterranean Sea (Gulf of Lion) during the Common
957	Era. Earth and Planetary Science Letters. Vol. 456, 124-133, 2016.

958	Smyth, W. The Sailor's Word-book: an alphabetical digest of nautical terms, including some
959	more especially military and scientific as well as archaisms of early voyagers, etc. by
960	the late Admiral W.H. Smyth (2004 Reprint) Toronto: Algrove Publishing Limited. 744
961	pp, 1867.
962	Stoetzel, D.: Encyclopedia of the French and Indian War in North America, 1754-1763. United
963	Kingdom: Heritage Books. 579 pp, 2008.
964	Storm, A.: Seaweed and Gold. Sydney Nova Scotia: City Printers Limited. 192 pp, 2002.
965	Syrett, D.: Shipping and Military Power in the Seven Years' War, 1756-1763: The Sails of
966	Victory. United Kingdom: Liverpool University Press. 192 pp, 2008.
967	Trouet, V., Scourse, J. and Raible C.: North Atlantic storminess and Atlantic meridional
968	overturning circulation during the last millennium: reconciling contradictory proxy
969	records of NAO variability. Global and Planetary Change. Vol. 84-85, 48-55, 2012.
970	U.S. Department of Commerce Environmental Science Services Administration Weather Bureau.
971	Tropical Cyclone Report - Hurricane Camille August 14-22, 1969. 58 pp.
972	https://www.nhc.noaa.gov.pdf/tcr-1969Camille.pdf, 1969. [accessed 2023-9-30]
973	Van Vliet-Lanoë, B., Goslin, J., Hallégouët, B., Hénaff, A., Delacourt, C., Fernane, A., Franzetti,
974	M., Le Cornec, E., Le Roy, P., and Penaud, A.: Middle- to late-Holocene storminess in
975	Brittany (NW France): Part I – morphological impact and stratigraphical record. The
976	Holocene. Vol. 24, 413–433, 2014.
977	Vecchi, G., and Knutson, T.: On estimates of historical North Atlantic tropical cyclone activity.
978	Journal of Climate. Vol. 21, 3580-3600, 2008.

979	Virot, E., Ponomarenko, A., Dehandschoewercker, E., Quere, D. and Clanet, C.: Critical wind
980	speed at which trees break. Physics Review. Vol. 93, 7 pp, 2016.
981	Warden, D.: A statistical, political, and historical account of the United States of North America
982	from the period of their first colonization to the present day. Vol. 1 of 3, 552 pp, 1819.
983	Wheeler, D.: An examination of the accuracy and consistency of ships' logbook weather
984	observations and records. Climate Change Vol. 31, 97-116, 2005.
985	Wheeler, D.: The Great Storm of November 1703 – A new look at the seamen's records.
986	Weather. Vol. 58, 419-427, 2006.
987	Wheeler, D. and Wilkinson, C.: From calm to storm: the origins of the Beaufort Wind Scale. The
988	Mariner's Mirror. Vol. 90, 187-201. DOI:10.1080/00253359.2004.10656896, 2004.
989	Wheeler, D., Garcia-Herrera, R. and Wilkinson, C.: Atmospheric circulation and storminess
990	derived from Royal Navy logbooks: 1685 to 1750. Climatic Change. Vol. 101, 257-280,
991	2010.
992	Winter, A., Ishioroshi, H., Watanabe, T., Oda, T., Christy, J.: Caribbean sea surface
993	temperatures' two-to-three degrees cooler than present during the Little Ice Age.
994	Geophysical Research Letters. Vol. 27, 3365-3368, 2000.

995 Zinck, J.: Shipwrecks of Nova Scotia. 226 pp, 1975.