

Intraseasonal and interannual variability of sea temperature in the Arabian Sea Warm Pool

Na Li¹, Xueming Zhu¹, Hui Wang^{2,3}, Shouwen Zhang^{1,*}, Xidong Wang^{4,1}

¹School of marine sciences, Sun Yat-sen University & Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, China

²Key Laboratory of Research on Marine Hazards Forecasting, National Marine Environmental Forecasting Center, Beijing, China

³Institute of Marine Science and Technology, Shandong University, Qingdao, China

⁴Key Laboratory of Marine Hazards Forecasting, Ministry of Natural Resources, Hohai University, Nanjing, China.

10 *Correspondence to:* Shouwen Zhang (zhangshouwen@sml-zhuhai.cn)

Abstract. The Arabian Sea Warm Pool (ASWP) is a part of the Indian Ocean Warm Pool, formed in the Arabian Sea before

the onset of the Indian summer monsoon. The ASWP has a significant impact on climate change in the Indian Peninsula and

globally. In this study, we examined the intraseasonal and interannual variability of sea temperature in the ASWP using the

latest Simple Ocean Data Assimilation (SODA) reanalysis dataset. We quantified the contributions of sea surface heat flux

15 forcing, horizontal advection, and vertical entrainment to the sea temperature using mixed-layer heat budget analysis method.

We also used a lead-lag correlation method to examine the relationship between the interannual variability of the ASWP and

various large-scale modes in the Indo-Pacific Ocean. We found that the ASWP formed in April and decayed in June, its

formation and decay processes were asymmetrical, with the decay rate being twice as fast as the formation rate. During the

ASWP development phase, the sea surface heat flux forcing had the largest impact on the mixed layer temperature with a

20 contribution of up to 85%. Its impact was divided into the net surface heat flux (0.41–0.50°C/5 day) and the shortwave

radiation loss penetrating the mixed layer (from -0.08°C/5 day to -0.17°C/5 day). During the decay phase, the cooling

effect of the vertical entrainment on the temperature variation increased (from -0.05°C/5 day to -0.18°C/5 day) and

dominated the temperature variation jointly with the sea surface heat flux forcing. We also found that the ASWP has strong

interannual variability related to the basin warming of the Indian Ocean. The lead-lag correlation indicated that the ASWP

25 had a good synchronous correlation with the Indian Ocean Dipole. The ASWP had the largest correlation coefficient at a lag

of 5–7 months of the Niño3.4 index, showing the characteristics of modulation by the El Niño-Southern Oscillation (ENSO).

When the El Niño (La Niña) event peaked in the winter of the previous year, the ASWP that occurred before the summer

monsoon was more significant (insignificant) in the following year.

1 Introduction

30 Tropical warm pools greatly affect atmospheric systems and oscillation modes, such as the monsoon, cyclones, El Niño (La Niña), and the Indian Ocean Dipole (IOD) (Lau and Chan, 1988; Webster and Lukas, 1992). Therefore, the study of warm

删除: 1.

删除: I

删除: Arabian Sea Warm Pool

删除: 5-day

删除:

删除: . I

删除: -

删除: -

删除: (from -0.05°C/5 day to -0.18°C/5 day)

删除: consistent

删除: basin

删除:

删除: and

pools and their climate effects is important for the understanding of global climate change. The ASWP is part of the Indian Ocean warm pool (IOWP). It forms in the Arabian Sea (AS) before the onset of the Indian summer monsoon. Sea surface temperature (SST) in the ASWP reaches its maximum before the summer monsoon onset (Kurian and Vinayachandran, 2007; Nagamani et al., 2016). It has a considerable influence on the monsoon onset eddies and other sea-air interactions, representing a remarkable mechanism for the regulation of the regional climate system (Rao and Sivakumar, 1999). A deeper exploration of the regulating factors of the evolution of the ASWP and its relationship with the large-scale modes can not only enhance the scientific knowledge of the changes of the ASWP but also provide a theoretical basis for the prediction of the ASWP.

In addition to being a component of the IOWP, the ASWP also demonstrates evolutionary traits that are distinct from those of the IOWP. Many studies have reported that high SST in the southeastern AS occurs only before the onset of the summer monsoon. In the MONEX-79 monsoon experiment, Seetaramayya and Master (1984) noted a warm pool (SST > 30.8°C, 10°N–13°N, 67°E–72°E) in the southeastern AS one week before the summer monsoon onset. Krishnamurti et al. (1988) noted an extreme value in the northern Indian Ocean before and after the monsoon onset in 1979. Joseph (1990) also suggested that before the onset of the summer monsoon, the warmest area of the warm pool of the tropical oceans is centred over southeast AS.

Rao and Sivakumar (1999) used monthly climatology observation and model simulations to examine plausible mechanisms for the gradual build-up of the ASWP. They found that the heat balance of the mixed layer essentially reflects the distribution of the net heat flux, while advection processes play a secondary modulating role. Debasis Sengupta's (2008) experiments in two different phases during the AS Monsoon Experiment determined that in the warming phase, the temperature rise mainly because of heat absorbed within the mixed layer, while in the cooling phase, the penetrative fluxes of solar radiation and advective cooling are together responsible for the rapid SST cooling. Kumar et al. (2009) used a three-dimensional circulation model to study the growth and decay characteristics of the ASWP in May 2000. The authors found that when the heat flux is not considered, the AS temperature continues to cool and no warm pools appear. When salinity changes are not considered, the ASWP still exists, but the horizontal extent of the warm pools decreases and the warming rate becomes slower. Sanilkumar et al. (2004) demonstrated that the vertical extent and stratification of low salinity waters affect the intensity of heating and thickness of the warm pool, while the extent of low salinity waters regulates the warm pool's horizontal coverage. However, Kurian and Vinayachandran (2007) casted doubt on this theory and proposed an alternative explanation for the warming that occurred in the early phases of the warm pool. They found that the orographic impact of the West Ghats reduces wind speed in the southeastern AS, thereby reducing latent heat loss and leading to positive heat flux entering the ocean.

Using in-situ observations and satellite data, Sabu and Revichandran (2011) examined the relative importance and contribution of various processes to the total heat budget of the mixed layer of the ASWP (29.5°C) during the spring intermonsoon (March–April 2004). They discovered that the advection has a role in transporting the warm water from the coastal region to the far west, while the surface heat flux is the primary factor in the mixed layer warming in the northern part of the

删除: Arabian Sea warm pool (

删除:)

删除: Its s

删除: and gradually dissipates with the onset of the monsoon

删除: –

删除: and forecasting

删除: Seetharamayya

删除: to

删除: to

删除: outbreak

删除: outbreak

删除: .

删除: ed

删除: ed

删除: s

删除: o

删除: . I

删除: were

删除: was

删除: d

删除: e

删除: d

删除: were

删除: ed

删除: d

删除: a

删除: ed

删除: d

warm pool. In the southern part of the warm pool, the eddy-induced horizontal mixing provides a substantial amount of heat spreading, which influences the mixed layer temperature evolution. Shankar et al. (1997) suggested that the formation of the ASWP is related to the Kelvin wave and its radiated Rossby wave in the tropical Indian Ocean and that the fluctuations lead to the formation of the Laccadive High over the AS, thus producing the ASWP.

70 The SST of the ASWP also shows strong interannual variability, which may be linked to the variation of large-scale modes, such as the Indian Ocean Basin mode (IOB), IOD, and the ENSO. According to Lau (2000) and Chowdary et al. (2007), zonal movement of the convergence centers linked to the Walker circulation is the main mechanism by which ENSO effects are transmitted to the Tropical Indian Ocean (TIO). This leads to subsidence in the eastern TIO, which reduces rainfall and causes abnormal easterly winds. Thereafter, the TIO as a whole warmed as a result of an increase in short-wave radiation and a drop in the latent heat flux, thereby affecting the ASWP. This warming is the dominant mode of interannual variability in TIO temperature. When an El Niño event develops, the size of the Indo-Pacific warm pool increases as it extends eastward in the Pacific Ocean sector and westward in the Indian Ocean sector, and vice versa for a La Niña event. The response of the warm pool intensity to ENSO does not reach its peak until about 5 months after ENSO peaks (Lau and Nath, 2003). The displacement and intensity changes in the Indo-Pacific warm pool affect the onset, intensity, and period of ENSO (Picaut et al., 1996).

80 Webster (1999) and Saji et al. (1999) demonstrated that sea-air coupling processes allow the TIO to create its internal modes of SST variability. It is worth noting that in recent years, roughly half of the IOD coincide with El Niño, while the other events are caused by internal TIO development (Meyers et al., 2007). ENSO variability is discovered to have an impact on the periodicity, strength, and formation processes of the IOD during the years of co-occurrence (Annamalai et al., 2003; Behera et al., 2006). IOD was more significant and persistent during the years of co-occurrence, and it was characterized by both eastern cooling and western warming (Chowdary and Gnanaseelan, 2007). For the majority of fluctuations in the area between 10°N–10°S, the observed interannual variability in ocean mixed layer depth—a crucial component that affects the SST variability—is related to the IOD (Thompson et al., 2006; Rao et al., 2015).

90 Even though some researchers have analyzed the climatological features of the ASWP, their studies are mainly based on monthly datasets or sparse data. The ASWP evolutionary features and their related processes were not well-examined due to low data resolution and high variability within that data. However, a variety of marine data with high spatial and temporal resolution and good quality now provides the possibility for more detailed research. Besides, although the effects between the large-scale modes (such as ENSO, IOD) and the Indo-Pacific warm pool have been studied, few studies have explored the relationship between various large-scale modes and smaller-scale seas (such as the AS). Based on the results of previous studies (Rao and Sivakumar, 1999; Kumar et al., 2009; Rao et al., 2015), this paper defines the threshold of the ASWP as 30°C. We investigated the characteristic and causes of the intraseasonal variability in the ASWP using heat budget analysis, and explored the relationship between the interannual variability of the ASWP and large-scale modes using lead-lag correlation. The overall framework of this paper is as follows: Section 2 briefly introduces the data sources used in this paper and the research methods adopted; Section 3 explains the intraseasonal characteristics of the ASWP and analyzes its

删除: ed

删除: d

删除: then

删除: was

删除: s

删除: led

删除: There is some interannual variation in the

删除: El Niño-Southern Oscillation (

删除:)

删除: led

删除: d

删除: d

删除: was

删除: , i.e., the Indian Ocean dipole

删除: d

删除: were

删除: was

删除: area

删除: was

删除: . In the equatorial and southeast Indian Oceans, a ...

删除: work has mostly been limited to monthly averages ...

删除: deeper

删除: A

删除: Using high-resolution satellite remote sensing and ...

100 thermodynamic diagnosis using the mixed layer diagnostic formula; Section 4 analyzes the interannual variability characteristics of the ASWP with IOB, IOD, and ENSO; and Section 5 summarizes the main research results and provides an outlook for future work.

2 Data and Methods

2.1 SODA Ocean Dataset

105 The **SODA3.7.2** ocean dataset is derived from the Simple Ocean Data Assimilation (SODA), which contains various parameters such as temperature, density, salinity, sea surface height, and current velocity, with a resolution of $0.25^\circ \times 0.25^\circ$ in the latitudinal and longitudinal directions and 50 layers in the vertical direction, up to 5400 m. The period of the data is from 1980 to 2020, with a time interval of 5 days (Giese, 2005).

2.2 The Japanese 55-year Reanalysis (JRA-55)

110 JRA-55, the forcing field used in SODA3.7.2 (Carton et al., 2018), is the Japanese second global atmospheric reanalysis project, originating from the Japan Meteorological Agency. Compared with its predecessor, JRA-25, JRA-55 is based on a new data assimilation and prediction system (DA) that improves upon many deficiencies found in the first Japanese reanalysis. These improvements have come about by implementing a higher spatial resolution (TL319L60), a new radiation scheme, four-dimensional variational data assimilation (4D-Var) with variational bias correction (VarBC) for satellite
 115 radiances, and the introduction of greenhouse gases with time-varying concentrations. The data period is from 1958 to the present, with a spatial resolution of $0.5625^\circ \times 0.5625^\circ$. This paper uses daily sea–air heat flux data, including long-wave radiation, short-wave radiation, sensible heat, and latent heat.

2.3 Methods

To investigate the specific processes affecting the temperature variation, this paper uses the SODA and JRA-55 datasets for
 120 the mixed layer thermodynamic diagnostic analysis in the ASWP ($SST > 30^\circ\text{C}$). The changes in mixed layer temperature with time were dominated by sea surface heat flux forcing (SHF), horizontal advection (ADV), and vertical entrainment (ENT). To perform a quantitative and specific analysis of each, we used equations (2.1)–(2.4) (Kurian and Vinayachandran, 2007; Li et al., 2016) for mixed layer diagnosis.

$$\frac{\partial T}{\partial t} = \text{SHF} + \text{ADV} + \text{ENT} + R \quad (2.1)$$

125
$$\text{SHF} = Q_{\text{net}} - Q_{\text{loss}} = \frac{Q_{\text{sw}} - Q_{\text{lw}} \pm Q_{\text{sh}} \pm Q_{\text{lh}}}{\rho_0 c_h} - \frac{Q_{\text{sw}} [r e^{\frac{h}{\beta_R}} + (1-r) e^{-\frac{h}{\beta_B}}]}{\rho_0 c_h} \quad (2.2)$$

$$\text{ADV} = -U \frac{\partial T}{\partial x} - V \frac{\partial T}{\partial y} \quad (2.3)$$

$$ENT = - (W_h + H \frac{\partial h}{\partial t} + U \frac{\partial h}{\partial x} + V \frac{\partial h}{\partial y}) \frac{T - T_h}{h} \quad (2.4)$$

$$ENT1 = - W_h \frac{T - T_h}{h} \quad (2.4.a)$$

$$ENT2 = - H \frac{\partial h}{\partial t} \frac{T - T_h}{h} \quad (2.4.b)$$

$$130 \quad ENT3 = - (U \frac{\partial h}{\partial x} + V \frac{\partial h}{\partial y}) \frac{T - T_h}{h} \quad (2.4.c)$$

The left side of Eq. (2.1) represents the local variation term of mixed layer temperature with time. The SHF, ADV, ENT, and R represent the surface heat flux, horizontal advection, vertical entrainment, and residual term, respectively. T represents the

average mixed layer temperature; $\frac{\partial T}{\partial x}$ and $\frac{\partial T}{\partial y}$ represent the zonal and meridional variation of mixed layer temperature,

respectively. Q_{net} represents the surface net heat flux, and its value is the sum of short-wave radiation flux (Q_{sw}), long-wave

135 radiation flux (Q_{lw}), sensible heat (Q_{sh}), and latent heat (Q_{lh}) at the sea surface. Q_{loss} represents the penetrating shortwave

radiation at the bottom of the mixed layer. $r=0.62$ is the red light fraction, $\beta_R=0.6$ m is the penetration depth scale of red

light, and $\beta_B=10$ m is the penetration depth scale of blue light. h is the depth of the mixed layer. In this paper, we select the

criterion to be a density difference of 0.03 kg/m^3 between the reference depth (10 m) and the bottom of the mixed layer (De

Boyer Montégut et al., 2004; Jofia et al., 2023). U, V is the average horizontal current velocity of the mixed layer. W_h is the

140 vertical velocity at the base of the mixed layer. T_h is the temperature at a depth of 10 m below the bottom of the mixed layer;

H is a dimensionless number that equals 0 when $\frac{\partial h}{\partial t} \leq 0$ and 1 when $\frac{\partial h}{\partial t} > 0$. ENT1, ENT2, and ENT3 in equations (2.4.a)–(2.4.c)

denote the entrainment due to the vertical velocity, the local changes of the mixed layer depth, and the “advection of the

mixed layer depth,” respectively (Nogueira Neto et al., 2018). The term R is the residual term, representing unresolved

processes such as the horizontal and vertical diffusion and errors in the estimation of terms. The units in the heat budget

145 equation are all $^{\circ}\text{C} / \text{s}$. Since the time interval of the SODA data set used in this paper is 5 days, in order to ensure the

accuracy of calculation, we converted the unit into $^{\circ}\text{C}/5$ days when calculating and **making plots**.

This study also used empirical orthogonal function (EOF) analysis to separate the spatial and temporal characteristics of the

ASWP and study the relationship between the ASWP and IOD/ENSO using lead-lag correlation. Previous studies have used

the above methods for related studies, such as Li (2017), and Kim et al (2012). However we focus on analyzing the results

150 every 5 days from April to June in order to gain new insights.

删除:

删除:

删除:

删除:

删除: drawing

删除:

删除: But


删除: some

删除: some

删除:

3 Intraseasonal variability of the ASWP

3.1 Intraseasonal variation characteristics of the ASWP



155 **Figure 1: Every 5-day evolution of SST (unit:°C) in April–June, where the black contour is the 30°C contour. The dashed rectangle represents the selected range of the ASWP (55.25°E-77.25°E, 5.25°N-20.25°N). The data is based on a gridded dataset for the period of 1980-2016 from SODA3.7.2. (<https://www2.atmos.umd.edu/~ocean/>)**

删除: s
删除: at
删除: v




Figure 2: Time series of the area and maximum temperature of the climatological ASWP, where the blue bars represent the warm pool area (km²) and the solid line represents the maximum warm pool temperature (°C). The area of the ASWP was the sea area within the rectangular range with a SST greater than 30°C. The maximum temperature was the highest temperature of the sea area with SST greater than 30°C within the rectangular sea area.

This paper defined a rectangle range of 55.25°E–77.25°E and 5.25°N–20.25°N based on the distribution of SST in the AS (As shown in Figure 1, marked with a dashed rectangle). ASWP was defined as the sea area within the rectangular range with a SST greater than 30°C. The entire formation and extinction of the ASWP are generally completed in April, May, and June. Therefore, using the climatological SST from April to June, we calculated the area and maximum temperature of the ASWP in Figures 1 and 2. It can be seen that, on average, the SST at the southern edge of the Indian Peninsula began to exceed 30°C on April 1, after which the high-temperature area continued to extend westward and northward. The warm pools then expanded to their maximum extent (1.63 × 10⁶ km²) on May 11, when the warm pools were in a matured stage. Thereafter, the warm pools decayed rapidly and disappeared completely in early June (Rao et al., 2015). From Figure 2, it can be seen that the development and decay processes of ASWP are not symmetrical. From the formation to the mature stage of the ASWP, it takes approximately 1.5 months (6 weeks). While from the mature stage to disappear completely, it only takes 3 weeks. The rate of decline is twice that of the development. The mechanism influencing this asymmetry in development and decay of the ASWP is discussed below.

3.2 ASWP mixed layer heat budget analysis




Figure 3: Contribution of different processes to the temperature variation of the warm pool mixed layer in the AS (°C/5 day), where $\frac{\partial T}{\partial t}$ (solid red line) is the temperature variation of the warm pool mixed layer with time, SHF (solid blue line) represents the sea surface heat flux forcing, ADV (solid green line) represents the horizontal advection, ENT (solid black line) represents the

- 删除:
- 删除: -
- 删除: -
- 删除: sea
- 删除: surface temperature
- 删除: is
- 删除: SODA
- 删除: sea
- 删除: surface temperature
- 删除: 1980–2016
- 删除: its
- 删除: were calculated to obtain
- 删除: sea
- 删除: surface temperature
- 删除: tip
- 删除: After that, t
- 删除: *
- 删除: almost
- 删除: than

180 vertical entrainment, and R (black dashed line) represents the residual. The red dashed line represents the sum of SHF, ADV, and ENT. The red dotted line represents the sum of SHF and ENT. 删除: d

To investigate the influence and contribution of different factors during the warm pool evolution, a mixed layer heat budget analysis was performed using SODA3.7.2 reanalysis data and JRA-55 heat flux dataset. Throughout the evolution of the ASWP, the sums of SHF, ADV, and ENT were consistent with the trend of mixed layer temperature. The residual term R was also maintained within 0.1°C/5 day, indicating that the heat balance equation was almost closed and the variation of the ASWP temperature was mainly related to these three terms. Therefore, these three terms can be used to explore the evolutionary mechanism of the ASWP qualitatively and quantitatively (Figure 3). 删除: The results are shown in Figure 3. 删除: basically

The variation in the ASWP mixed layer temperature was asymmetric. From April 1 to May 16, $\frac{\partial T}{\partial t}$ was greater than zero, and the mixed layer temperature gradually increased as the ASWP developed. After May 21, $\frac{\partial T}{\partial t}$ was less than zero, and the mixed layer temperature rapidly decreased, corresponding to the rapid decay of the warm pool. SHF consistently dominated the warm pool throughout the warming phase, with a contribution of up to 0.34°C/5 day. The overall trend of the temperature variation was consistent with SHF. ADV and ENT had less influence on the temperature variation, and the ENT played a weak cooling role. In the cooling phase, the SHF gradually decreased and the ENT strengthened, up to -0.27°C/5 day. The effect of ADV was much smaller, not more than 0.1°C/5 day. The following is a detailed analysis of the mechanism and effect of each process. 删除: n 删除: influence 删除: always 删除: specific and




Figure 4: Contribution of the sea surface heat flux forcing to the mixed layer temperature ($^{\circ}\text{C}/5$ day), where blue bars represent the total sea surface heat flux forcing, red bars represent the contribution of the net surface heat flux, and yellow bars represent the contribution of penetrating shortwave radiation at the bottom of the mixed layer.

200 The contribution of SHF to $\frac{\partial T}{\partial t}$ ranged from $0.11^{\circ}\text{C}/5$ day to $0.34^{\circ}\text{C}/5$ day (Figure 4), always playing a heating role. From April to early May, the contribution of SHF to temperature was around $0.3^{\circ}\text{C}/5$ day, indicating that SHF provided substantial thermal support for the increased temperature. By early June, the contribution of SHF to temperature dropped below $0.2^{\circ}\text{C}/5$ day. The effect of SHF on the warm pool temperature can be divided into two parts, the net surface heat flux ($\text{SHF}_{Q_{net}}$) and the penetrating shortwave radiation at the bottom of the mixed layer ($\text{SHF}_{Q_{loss}}$). In the upper layer of the mixed layer, the temperature change is related to the heat exchange between the surface seawater and the atmosphere, so it is related to the net heat flux at the sea surface ($\text{SHF}_{Q_{net}}$). In the lower layer of the mixed layer, the heat flux that can penetrate the bottom of the mixed layer is mainly shortwave radiation. Therefore, the heat loss at the bottom of the mixed layer is mainly related to penetrative shortwave radiation ($\text{SHF}_{Q_{loss}}$). The contribution of $\text{SHF}_{Q_{net}}$ gradually increased from $0.41^{\circ}\text{C}/5$ day to $0.50^{\circ}\text{C}/5$ day in April. It increased slowly but remained at a relatively high level. Then it gradually decreased to less than $0.2^{\circ}\text{C}/5$ day.

210 Similar to $\text{SHF}_{Q_{net}}$, $\text{SHF}_{Q_{loss}}$ also gradually increased to $-0.17^{\circ}\text{C}/5$ day in April and then decreased.

删除: s

删除: s

删除: s

删除: the

删除: that are caused by

删除: a

删除: -




Figure 5: The upper panel: Monthly climatology of shortwave radiation flux (shading; W/m^2) and mixed layer temperature (black contours; $^{\circ}C$). The lower panel: the distribution of total cloud coverage.

删除: color

SHF_{Q_{net}} is related to sensible heat, latent heat, longwave radiation flux, and shortwave radiation flux (SWR). In climatology, the SWR was the largest contributor ($190\text{--}283\text{ W/m}^2$) and was always positive, indicating that the main source of heat in the warm pool was shortwave radiation from the sun. Therefore, we focused on the variation in SWR in the AS in combination with cloud coverage (Figure 5). In April, the clear sky and lack of clouds over the AS mean the sea surface can receive more SWR than at other times. The SWR was above 240 W/m^2 throughout the AS, with a clear high value in the center of the AS. In May, when the shortwave radiation flux had decreased (230 W/m^2), the temperature of the warm pool reached its highest. When the summer monsoon broke out, the atmospheric convective activity was strong and the cloud coverage over the sea was high, leading to a decrease in the shortwave radiation entering the mixed layer. The shortwave radiation dropped to below 220 W/m^2 , with a clear low-value center ($<180\text{ W/m}^2$) in the southeast of the AS. In addition, although the variation of the mixed layer temperature was roughly in line with the trend of SWR, the variation of temperature had a lag time of about one month compared with SWR. This trend occurred because seawater has a large specific heat capacity and there is a lag of one month in the variation of temperature.

删除:





Figure 6: Monthly climatology of shortwave radiation loss penetrating the mixed layer (color shading; W/m^2) and the depth of the mixed layer (black contours; m).

SHF_{Q_{loss}} was ~~negative correlated with~~ the mixed layer depth (Figure 6). In April, the wind speed over the AS was low, making the mixed layer depth shallow (<15 m). Shortwave radiation can penetrate the mixed layer at this time ($-55 W/m^2$) and heat the subsurface seawater. In May, when the summer monsoon was about to ~~onset~~, increased wind strengthened the ~~ocean~~ mixing, leading to a deeper mixed layer (25 m) and a decrease in the shortwave radiation penetrating the mixed layer ($-37 W/m^2$). In June, when the summer monsoon was fully formed, ~~the mixed layer was further deepened (>30m) and~~ the penetrating shortwave radiation ~~through the mixed layer~~ was further decreased ($-9 W/m^2$). In addition, the study by Liu et al (2007) found that MLD is forced by penetrative shortwave radiation, and the impact of penetrative shortwave radiation on the depth of the mixed layer is controlled by wind speed and net shortwave radiation at the sea surface.

删除: related to
删除: and was negatively correlated with it
删除: break
删除: out
删除: across the AS
删除: and the mixed layer was further deepened (>30m)
删除: Zhiliang

3.2.2 Vertical entrainment



240 **Figure 7: Variation of the vertical entrainment and mixed layer depth, where the blue bars represent the vertical entrainment (°C/5 day) and the red line represents the mixed layer depth (m).** 删除: s




Figure 8: Contribution of each term in the vertical entrainment to the temperature change of the mixed layer.

The contribution of ENT to temperature ranged from $-0.04^{\circ}\text{C}/5$ day to $-0.28^{\circ}\text{C}/5$ day (Figure 7), acting as a cooling effect and intensifying over time. Together with the gradual decrease in SHF, it led to a rapid cooling of the ASWP. During the development phase of the warm pool, the cooling effect of ENT was weak, accounting for only 16%–20%. After May 1, the cooling effect intensified up to 58%. The effect of ENT on temperature can be divided into three parts: the entrainment due to the vertical velocity (ENT1), the entrainment due to the local tendency of the mixed layer depth (ENT2), and the entrainment due to “advection of the mixed layer depth” (ENT3). The cooling effect due to the local variation of the mixed layer was the strongest and almost dominated, ranging from $-0.05^{\circ}\text{C}/5$ day to $-0.18^{\circ}\text{C}/5$ day (Figure 8).

The vertical entrainment is closely related to the vertical thermohaline structure of seawater and the depth of the mixed layer. When the mixing of the upper seawater in terms of temperature and salinity is inconsistent, the isothermal layer and mixed layer will differ, and a barrier layer will be formed between the bottom of the mixed layer and the top of the thermocline (Sprintall and Tomczak, 1992). The barrier layer has the characteristics of a strong salinity gradient and high gravitational stability, which makes it difficult to transport heat from top to bottom by mixing. On the one hand, a strong and stable salinity stratification can effectively inhibit the transfer of non-shortwave radiation flux to the interior of the ocean, which leads to the warming of the upper mixed layer. On the other hand, the existence of the barrier layer can inhibit the entry of the cold water of the thermocline into the mixed layer, which is not conducive to the exchange of heat, momentum, mass, and nutrients between the mixed layer and the thermocline (Pang Shanshan, 2021). Therefore, In the development stage of the ASWP, the low wind speed made the mixed layer shallower, which made the seawater within the mixed layer more easily heated and thus more sensitive to heat flux. Then the weak wind speed and the existence of anticyclonic circulation (Shankar et al., 2002) associated with the Lakshadweep High trapped the fresh water, contributing to the formation of the barrier layer, which maintain~~ing~~ the ASWP by effectively inhibiting the vertical mixing of the upper ocean, and causing ~~the~~ warming of the upper mixed layer. In the late stage of ASWP evolution, with the imminent ~~onset~~ of the summer monsoon, the stirring effect of wind started to strengthen, the mixing layer became deeper rapidly (Figure 6), and the cooling effect of entrainment was enhanced, which accelerated the decay of the ASWP (Liu Yanliang, 2013).

3.2.3 Horizontal advection

The contribution of ADV to $\frac{\partial T}{\partial t}$ ranges from $-0.05^{\circ}\text{C}/5$ day to $0.01^{\circ}\text{C}/5$ day, which is much smaller than the contribution of other processes. From April 1 to May 1, the contribution of ADV was positive, indicating that the advection process played a weak heating role during the development phase. After May 1, ADV started to become negative and played an increasingly important cooling role in the warm pool. Due to heavy rainfall and river runoff over evaporation, the salinity of the seawater in the Bay of Bengal was very low. During November and December, there is an equatorward-flowing East India Coastal Current (EICC) off the east coast of India. This current flows around Sri Lanka and continues as a poleward-flowing West India Coastal Current off the west coast of India (Shetye et al., 1996; Rao and Sivakumar, 1999; Mukhopadhyay et al., 2020). The EICC carries a large amount of low-salt seawater from the Bay of Bengal into the AS. This coastal current system is

删除: -

删除: -

删除: always

删除: -

删除: -

删除: e

删除: and

删除:

删除: a

删除: outbreak

275 further strengthened between December and February by the westward North Equatorial Current (NEC). These low-saline
waters create a near-surface halocline, which enhances stratification. Then the stratification prevents the vertical
redistribution of the turbulent and radiative heat fluxes, amplifying the change in SST (Rao and Sivakumar, 1999). Due to
the low wind speed in the AS from April to May, the sea surface flow field is very weak. Coupled with the anticyclonic
circulation (Shankar et al., 2002) associated with the Lakshadweep High (Bruce et al., 1994), the low-salinity water is
280 trapped in the AS and can be maintained until May, promoting the formation of the barrier layer, which is conducive to the
development and maintenance of the ASWP. In June, the summer monsoon is about to break out, the sea surface wind field
changes, the sea surface flow field also turns, and the salinity stratification weakens, promoting the disappearance of the
ASWP.

Therefore, during the development and peak stages of the ASWP, strong shortwave radiation flux causes sea temperature to
285 continuously rise, gradually forming a warm pool. The presence of weak wind and the barrier layer allows for the
maintenance of high temperature. During this time, the ASWP is mainly influenced by the sea surface heat flux forcing, with
little impact from vertical entrainment and horizontal advection. In the decline stage of the ASWP, the rapid decrease in
shortwave radiation flux weakens the heating effect of the sea surface heat flux forcing, while the strengthening of mixing
leads to an increase in vertical cooling. The combined effect of these two factors results in a sharp decrease in temperature,
290 leading to the rapid decline of the warm pool.

4 Interannual variability of the ASWP

4.1 Characteristics of the interannual variability of ASWP




Figure 9: Time series of ASWP area from April to June 1980–2016 (the area calculated is daily mean every 5 days; red dots indicate the average area of ASWP per year). The horizontal black continuous line represents the seasonal mean, and the dashed lines represent ± 0.5 standard deviations.

The ASWP does not occur every year, its intensity and extent vary from year to year. As shown in Figure 9, there has been a significant interannual variation in the area of the ASWP. The ASWP was a large area in 1980, 1991, 1998, 2003, 2005, 2010, 2014, 2015, and 2016, exceeding 0.5 standard deviations of the area and reaching a maximum of 3.1×10^6 km². In other years, the area of the warm pool has been almost negligible (1984, 1989, 1992, 1999, 2000, 2008), indicating that the ASWP is weak in these years. A basin-scale warming/cooling is the leading mode of tropical Indian Ocean SST variability on interannual time scales. It peaks in late winter and persists into the following spring and summer (Klein et al., 1999; Li et al., 2008). The years with a large (small) ASWP area correspond to positive (negative) IOB periods, indicating that the ASWP area is mainly controlled by IOB (Guo et al., 2018).

删除: . I

删除: *

删除: s

删除: at this time

305




Figure 10: Variation of the area and maximum temperature of different types of ASWP (calculated every 5 days), where the bars represent the area of the warm pool (km²), the solid lines represent the maximum temperature of the warm pool (°C), red line represents those during a weak ASWP, blue represents those during a normal ASWP, and yellow represents those during a strong ASWP.

删除: -
删除:

310




Figure 11: ASWP mixed layer heat budget diagnosis, with weak ASWP years on the left, normal ASWP years in the middle, and strong ASWP years on the right.

Table 1. Contribution of each term to temperature variation.

	SHF(%)			ENT(%)			ADV(%)		
	Weak	Normal	Strong	Weak	Normal	Strong	Weak	Normal	Strong
Heating phase	80.20	78.51	81.52	19.74	19.7	17.04	0.06	1.71	1.44
Cooling phase	31.87	35.27	38.35	53.59	54.20	52.15	14.55	10.52	9.50

315 Based on a 0.5 sigma criterion, we estimated the average area of the ASWP for each year and identified the weak, normal, and strong ASWP years. As shown in Figure 10 and Figure 11, the warm pool area and maximum temperature were always the highest in strong ASWP years, with an average area of 1.3×10^6 km² and a maximum temperature of 30.7°C. In normal years, the average area was 0.53×10^6 km², and the maximum temperature was 30.3°C. In weak ASWP years, the average area was only 7.8×10^3 km², and the maximum temperature was only 29.9°C. The diagnostic results of its mixed layer

320 synthesized separately indicated similarity to the diagnostic results of the climatology (Figure 3). During the warming phase, the temperature changes were faster in strong ASWP years ($>0.2^\circ\text{C}/5_{\text{day}}$), while the temperature changes were smaller in weak ASWP years ($<0.2^\circ\text{C}/5_{\text{day}}$). This is consistent with the faster expansion and larger area of ASWP in strong years. However, regardless of the year, the contribution of SHF to the temperature change was largest (78.5% to 81.5%), and the contributions of ENT and ADV were relatively small. Among them, the weaker warm pool in weak ASWP years may be related to the greater cooling effect of ENT at this time (19.7%). In the cooling phase, the contribution of SHF decreased (31.9% to 38.3%), the cooling effect of ENT increased (52.2% to 54.2%), and the contribution of ADV increased slightly (9.5% to 14.6%). SHF and ENT together dominated the temperature change of the ASWP. The larger (smaller) warm pool in

325 strong (weak) ASWP years may be related to the larger (smaller) heating effect of SHF and the smaller (larger) cooling effect of ENT and ADV.





Figure 12: EOF1 (the first column), EOF2 (the second column), and PC1 and PC2 (the third column) of SST anomalies in the AS for April-June.



335 **Figure 13: Lead-lag correlation between PC1 and the IOD index DMI (left) and ENSO index niño3.4 (right) in April, May, and June.**

The EOF analysis was conducted separately for temperatures in the AS from April to June by using SST anomaly from the SODA3.7.2 dataset, and the first two modes were used for the analysis (Figure 12). In April, May, and June, the variance contributions of the first mode were all greater than 50% (60.86%, 54.80%, and 51.83%). The spatial distributions were similar, showing a consistent change of the temperature in the AS. This was mainly related to the Indian Ocean basin model (Guo et al., 2018). The Principal Component (PC) time series in April and May showed large positive peaks in 1987, 1991, 1998, 2010, and 2016, indicating that the temperature in April and May in the AS was anomalously warm in these five years. This corresponds to several years with large ASWP areas, indicating that the large ASWP ranges in these five years were mostly due to the consistent warming of the IOB. Similarly, the years with negative peaks in the PC time series (1989, 1992, 1999, 2000, and 2008) coincide with the years with smaller ASWP areas, suggesting that these five years with smaller ASWP ranges were controlled by the IOB. The variance contribution of the second mode (10.71%, 12.51%, and 11.52%) was much smaller than the first mode. The spatial distribution pattern varied by months, but there are at least two antiphase extreme centers.

340
345
350 The most dominant interannual climate modes in the TIO are the IOB mode and the IOD mode (Schott et al., 2009). Therefore, we further investigated the effects of IOB and IOD on the interannual variability of the ASWP. When the IOB is in a positive phase, there is consistent warming across the AS with a large range and intensity in the ASWP. This is consistent with ASWP's larger area in 1987, 1991, 1998, 2005, 2010, 2014, 2015, and 2016. The IOD is an inherent interannual climate mode in the TIO. During a positive phase, warm waters are pushed to the western part of the Indian Ocean, while cold deep waters are brought up to the surface in the eastern Indian Ocean. This pattern is reversed during the

- 删除:
- 删除: temperature
- 删除: ,
- 删除:
- 删除: was
- 删除: And t
- 删除: and time
- 删除: changes
- 删除: which shows
- 删除: the
- 删除:
- 删除: The amplitude of PC2 change every month is smaller than the amplitude of PC1.
- 删除: The ASWP is part of the Indian Ocean.
- 删除:
- 删除: W
- 删除: E

355 negative phase of the IOD (Saji et al., 1999; Kumar et al., 2020). The AS is located in the western side of the Indian Ocean, which is part of the western IOD. Its local sea-air interaction and interannual variation are closely related to the IOD. The left panel of Figure 13 shows that the highest correlation coefficient between the PC1 and IOD reached 0.56 ($p < 0.01$), indicating that the ASWP had a significant correlation with the IOD. There are three peaks in the lead-lag correlation between PC1 and IOD. The first peak appears at -5, -7, and -8 months ($r = 0.48$ - 0.56), indicating that the correlation is highest when the sea temperature lags the IOD 5-8 months, this is consistent with that the IOD peaks in winter (Li et al., 2008). The second peak appears at -2, -1, and 0 months ($r = 0.36$ - 0.49), indicating that the IOD can also regulate the temperature in near real-time. The third peak appears at 7 months ($r = 0.14$ - 0.34), indicating that the temperature is not only regulated by IOD, but can also affect the IOD mode, although the effect is relatively small. This is consistent with the research results of Li et al (2008). The Indian Ocean is directly and indirectly influenced by global or other regional climate patterns, such as the ENSO. Therefore, we also considered the impact of the ENSO on the interannual variability of the ASWP. Previous research has indicated that the ENSO has an impact on SST over the Indian Ocean (Klein et al., 1999; Thandlam et al., 2020). Both observations and model studies indicate that the Indian Ocean warms (cools) throughout the ocean basin in March-May following an El Niño (La Niña) year (Song et al., 2007; Dommengeset and Jansen, 2009; Venugopal et al., 2018; Thandlam et al., 2023). Kim et al. (2012) investigated the warm pool features in the Indian and Pacific Ocean and discovered that the seasonal variations in the Indian Ocean were greater than in the Pacific, which was due mostly to ENSO. The intensity of the warm pool was associated to a 5-6 months delay after the ENSO. The right panel of Figure 13 shows the lead-lag correlation between the PC1 of ASWP and Niño3.4. The correlation coefficient between PC1 and the Niño3.4 index is highest at a lag of 5-7 months ($r = 0.61$ - 0.72), and then sharply decreases after -2 months. This indicates that the temperature is modulated by the ENSO. In other words, after the El Niño (La Niña) peaks in winter, the ASWP before the summer monsoon in the upcoming year was more (less) pronounced, which is consistent with the strength of the IOWP peaking about five months after the ENSO peaked. This delay is because that the ENSO-induced teleconnection between oceans takes some time to develop (Lau and Nath, 2003). ENSO can not only directly affect the temperature changes in the AS through atmospheric bridges (Lau and Nath, 2000; Chowdary and Gnanaseelan, 2007), but also indirectly regulate the temperature changes by influencing the IOD (Behera et al., 2006). So, the correlation coefficient between the ENSO index and PC1 is higher than that of IOD, and the impact is stronger.

- 删除: .
- 删除: its
- 删除: -
- 删除: -
- 删除: -
- 删除: -
- 删除: ~
- 删除: seawater
- 删除: behind
- 删除: for
- 删除: -
- 删除: . It also means that the temperature from April to Ju ...
- 删除: -
- 删除: -
- 删除: ~
- 删除: ~
- 删除: was
- 删除: ~
- 删除: -
- 删除: -
- 删除: -
- 删除: ~
- 删除: ~
- 删除: -
- 删除: ,
- 删除: ing
- 删除: it
- 删除: was
- 删除: due to the fact
- 删除: On the one hand




Figure 14: ASWP mixed layer heat budget with El Niño year on the left, normal year in the middle, and La Niña year on the right.

Can changes in the ENSO affect the role of different processes in the evolution of the ASWP? To explore this question, we divided the ASWP into El Niño, La Niña, and normal years, and diagnosed them separately. The results are shown in Figure 14. It can be seen from the overall trend that the change in the ENSO did not change the dominance of SHF and ENT in the evolution of ASWP. However, the surface heat flux forcing still dominated in the warming phase, and the surface heat flux and the vertical entrainment together dominated the temperature change of ASWP in the cooling phase. It is noteworthy that although the temperature warmed up during El Niño, the effect of surface heat flux forcing did not increase significantly. However, the effect of vertical entrainment was enhanced, which may be related to the enhanced convective activity and mixing effect during El Niño (Sullivan et al., 2019).

5 Summary and Discussion

In this study, we analyzed the causes of the intraseasonal variability in the ASWP using heat budget analysis and explored the relationship between the interannual variability of the ASWP and large-scale modes using lead-lag correlation. By using the latest 5-day SODA reanalysis dataset, we found that the formation and extinction of the ASWP were completed in April, May, and June. Its formation and decay rates showed asymmetry, with a decay rate twice as fast as the development rate.

The results of the heat budget analysis show that the sea surface heat flux forcing had the largest effect on the ASWP's mixed layer temperature. The next most important factor was the vertical entrainment, and then the horizontal advection had the smallest effect, which is consistent with previous results. The effect of SHF on the warm pool temperature was divided

删除: I

删除:

into two parts, $SHF_{Q_{net}}$ and $SHF_{Q_{loss}}$. $SHF_{Q_{net}}$ was mainly related to shortwave radiation, which is at a maximum in April
400 due to the lack of clouds and gradually decreases with an increase in cloudiness in May and June. $SHF_{Q_{loss}}$ was related to
the depth of the mixed layer and increases with the onset of the summer monsoon in June. The ENT always played a cooling
role. The increasing cooling effect made the ASWP decay rapidly, and the effect due to the local changes in the mixed layer
was the strongest.

ASWP, as part of the Indian Ocean, has strong interannual variability. The diagnostic analysis revealed that during the
405 warming phase, temperature changes were faster in strong ASWP years. While the temperature changes were smaller in
weak ASWP years. This is consistent with the faster expansion and larger area of the ASWP in strong years. Then we found
that the ASWP had a positive correlation ($r = 0.56$) between its PC1 and IOD most of the time. Concerning the ENSO, the
ASWP had the largest correlation at a lag of 5–7 months for the Niño3.4 index, showing that it was modulated by the ENSO.
In other words, after the El Niño (La Niña) peaks in winter, the ASWP before the summer monsoon in the upcoming year
410 was more (less) pronounced.

However, there are many interacting factors affecting the variability of temperature in AS. In this paper, we investigated the
effects of only three processes: sea surface heat flux forcing, horizontal advection, and vertical entrainment. We **did not**
consider other factors, such as horizontal and vertical diffusion, which require further optimization of the diagnostic formula
and more comprehensive diagnostic methods. In addition, the variation of ASWP is influenced by various global scale
415 modes. **Its development, extinction, and interannual variation should be regarded as a coupled ocean–land–air process. In**
this way, we can capture its variation characteristics, dynamic mechanisms, and influence on various sea–air interaction
processes more comprehensively and accurately.

删除: With respect to

删除: ignored

删除: e

删除: to

Data availability. The SODA and JRA-55 dataset described in this paper is available to download at the following address:
420 <https://dsrs.atmos.umd.edu/DATA/soda3.7.2/ORIGINAL/ocean/>; <https://rda.ucar.edu/datasets/ds628.0/dataaccess/>.

Author contributions. SWZ, XMZ, and XDW designed the study. NL conducted and made the analysis. All authors contributed to the
discussion of the analysis and the final manuscript.

425 *Competing interests.* The authors declare that they have no conflict of interest.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and
institutional affiliations.

430 *Acknowledgments.* We thank Hui Wang and Xidong Wang for their helpful discussions.

Financial support. This research was supported by the NSFC (42206029), the National Key R&D Program of China (2020YFA0608804),
and Southern Marine and Engineering Guangdong Laboratory (Zhuhai) (SML2020SP008).

References

- 435 Annamalai, H., Murtugudde, R., Potemra, J., Xie, S. P., Liu, P., and Wang, B.: Coupled dynamics over the Indian Ocean: spring initiation of the Zonal Mode, *Deep-Sea Res., Part II.*, 50, 2305-2330, [https://doi.org/10.1016/S0967-0645\(03\)00058-4](https://doi.org/10.1016/S0967-0645(03)00058-4), 2003.
- Behera, S. K., Luo, J. J., Masson, S., Rao, S. A., Sakuma, H., and Yamagata, T.: A CGCM Study on the Interaction between IOD and ENSO, *J. Clim.*, 19, 1688-1705, <https://doi.org/10.1175/JCLI3797.1>, 2006.
- 440 Bruce, J. G., Johnson, D. R., and Kindle, J. C.: Evidence for eddy formation in the eastern Arabian Sea during the northeast monsoon, *J. Geophys. Res.: Oceans.*, 99, 7651-7664, <https://doi.org/10.1029/94JC00035>, 1994.
- Carton, J. A., Chepurin, G. A., and Chen, L.: SODA3: A New Ocean Climate Reanalysis, *J. Clim.*, 31, 6967-6983, <https://doi.org/10.1175/JCLI-D-18-0149.1>, 2018.
- Chowdary, J. S. and Gnanaseelan, C.: Basin-wide warming of the Indian Ocean during El Niño and Indian Ocean dipole years, *Int J Climatol.*, 27, 1421-1438, <https://doi.org/10.1002/joc.1482>, 2007.
- 445 de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *J. Geophys. Res.: Oceans.*, 109, <https://doi.org/10.1029/2004JC002378>, 2004.
- Dommenget, D. and Jansen, M.: Predictions of Indian Ocean SST Indices with a Simple Statistical Model: A Null Hypothesis, *J. Clim.*, 22, 4930-4938, <https://doi.org/10.1175/2009JCLI2846.1>, 2009.
- Giese, J. A. C. a. B. S.: SODA: A Reanalysis of Ocean Climate, *Journal of Geophysical Research-Oceans*, 2005.
- Guo, F., Liu, Q., Yang, J., and Fan, L.: Three types of Indian Ocean Basin modes, *Clim. Dyn.*, 51, 4357-4370, <https://doi.org/10.1007/s00382-017-3676-z>, 2018.
- Jofia, J., Girishkumar, M. S., Ashin, K., Sureshkumar, N., Shivaprasad, S., and Pattabhi Ram Rao, E.: Mixed Layer Temperature Budget in the Arabian Sea During Winter 2019 and Spring 2019: The Role of Diapycnal Heat Flux, *J. Geophys. Res.: Oceans.*, 128, e2022JC019088, <https://doi.org/10.1029/2022JC019088>, 2023.
- Joseph, P. V.: Warm pool in the Indian Ocean and monsoon onset, *Tropical Ocean-Atmos. News Let.*, 53,1-55, 1990.
- Kim, S. T., Yu, J. Y., and Lu, M.-M.: The distinct behaviors of Pacific and Indian Ocean warm pool properties on seasonal and interannual time scales, *J. Geophys. Res.*, 117, <https://doi.org/10.1029/2011JD016557>, 2012.
- 460 Klein, S. A., Soden, B. J., and Lau, N.-C.: Remote Sea Surface Temperature Variations during ENSO: Evidence for a Tropical Atmospheric Bridge, *J. Clim.*, 12, 917-932, [https://doi.org/10.1175/1520-0442\(1999\)012<0917:RSSTVD>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<0917:RSSTVD>2.0.CO;2), 1999.
- Krishnamurti, T. N., Oosterhof, D. K., and Mehta, A. V.: Air–Sea Interaction on the Time Scale of 30 to 50 Days, *J. Atmos. Sci.*, 45, 1304-1322, [https://doi.org/10.1175/1520-0469\(1988\)045<1304:AIOTTS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<1304:AIOTTS>2.0.CO;2), 1988.
- 465 Kumar, P., Hamlington, B., Cheon, S.-H., Han, W., and Thompson, P.: 20th Century Multivariate Indian Ocean Regional Sea Level Reconstruction, *J. Geophys. Res.: Oceans.*, 125, e2020JC016270, <https://doi.org/10.1029/2020JC016270>, 2020.
- Kumar, P. V. H., Joshi, M., Sanilkumar, K. V., Rao, A. D., Anand, P., Kumar, K. M. A., and Rao, C. V. K. P.: Growth and decay of the Arabian Sea mini warm pool during May 2000: Observations and simulations, 528-540, <https://doi.org/10.1016/j.dsr.2008.12.004>,
- 470 Kurian, J. and Vinayachandran, P. N.: Mechanisms of formation of the Arabian Sea mini warm pool in a high-resolution Ocean General Circulation Model, *J. Geophys. Res.: Oceans.*, 112, <https://doi.org/10.1029/2006JC003631>, 2007.
- Lau, K.-M. and Chan, P. H.: Intraseasonal and Interannual Variations of Tropical Convection: A Possible Link between the 40–50 Day Oscillation and ENSO?, *J. Atmos. Sci.*, 45, 506-521, [https://doi.org/10.1175/1520-0469\(1988\)045<0506:IAIVOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<0506:IAIVOT>2.0.CO;2), 1988.
- 475 Lau, N.-C. and Nath, M. J.: Impact of ENSO on the Variability of the Asian–Australian Monsoons as Simulated in GCM Experiments, *J. Clim.*, 13, 4287-4309, [https://doi.org/10.1175/1520-0442\(2000\)013<4287:IOEOTV>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<4287:IOEOTV>2.0.CO;2), 2000.
- Lau, N.-C. and Nath, M. J.: Atmosphere–Ocean Variations in the Indo-Pacific Sector during ENSO Episodes, *J. Clim.*, 16, 3-20, [https://doi.org/10.1175/1520-0442\(2003\)016<0003:AOVITI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0003:AOVITI>2.0.CO;2), 2003.
- Li, S., Lu, J., Huang, G., and Hu, K.: Tropical Indian Ocean Basin Warming and East Asian Summer Monsoon: A Multiple AGCM Study, *J. Clim.*, 21, 6080-6088, <https://doi.org/10.1175/2008JCLI2433.1>, 2008.
- 删除: p

- Li Xiaohui, X. F., Chen Hongying, et al.: Correlation analysis of the cycle process between the western Pacific warm pool and ENSO during 1980-2016, *Journal of Marine Meteorology*, 37(3), 85-94, 1019513/i.cnki.issn2096-3599.2017.03.010, 2017.
- 485 Li, Y., Han, W., Wang, W., and Ravichandran, M.: Intraseasonal Variability of SST and Precipitation in the Arabian Sea during the Indian Summer Monsoon: Impact of Ocean Mixed Layer Depth, *J. Clim.*, 29, 7889-7910, <https://doi.org/10.1175/JCLI-D-16-0238.1>, 2016.
- Liu Yanliang, Y. W., Li Kuiping: Mixed layer heat budget in Bay of Bengal: Mechanism of the generation and decay of spring warm pool, *Acta Oceanol. Sin.*, 35, 1-8, 2013.
- 490 Liu Zhiliang, C. Z.: The effects of penetration radiation and salinity on mixed layer depth, *Transactions of Oceanology and Limnology*, 2, 19-25, 2007.
- Meyers, G., McIntosh, P., Pigot, L., and Pook, M.: The Years of El Niño, La Niña, and Interactions with the Tropical Indian Ocean, *J. Clim.*, 20, 2872-2880, <https://doi.org/10.1175/JCLI4152.1>, 2007.
- Mukhopadhyay, S., Shankar, D., Aparna, S. G., Fernando, V., and Kankonkar, A.: Observed variability of the East India Coastal Current on the continental shelf during 2010–2018, *Journal of Earth System Science*, 129, 106, 10.1007/s12040-020-1353-9, 2020.
- 495 Nagamani, P. V., Ali, M. M., Goni, G. J., Udaya Bhaskar, T. V. S., McCreary, J. P., Weller, R. A., Rajeevan, M., Gopala Krishna, V. V., and Pezzullo, J. C.: Heat content of the Arabian Sea Mini Warm Pool is increasing, *Atmospheric Sci. Lett.*, 17, 39-42, <https://doi.org/10.1002/asl.596>, 2016.
- Nogueira Neto, A. V., Giordani, H., Caniaux, G., and Araujo, M.: Seasonal and Interannual Mixed-Layer Heat Budget Variability in the Western Tropical Atlantic From Argo Floats (2007–2012), *J. Geophys. Res.: Oceans.*, 123, 5298-5322, <https://doi.org/10.1029/2017JC013436>, 2018.
- PANG Shanshan, W. X., LIU Hailong, SHAO Caixia: Multi-Scale Variations of Barrier Layer in the Tropical Ocean and Its Impacts on Air-Sea Interaction: A Review, *Adv. Earth Sci.*, 36, 139-153, 10.11867/j.issn.1001-8166.2021.022, 2021.
- 500 Picaut, J., Ioualalen, M., Menkes, C. E., Delcroix, T., and McPhaden, M. J.: Mechanism of the Zonal Displacements of the Pacific Warm Pool: Implications for ENSO, *Science*, 274, 1486 - 1489, DOI: 10.1126/science.274.5292.1486, 1996.
- 505 Rao, R. and Sivakumar, R.: On the possible mechanisms of the evolution of a mini-warm pool during the pre-summer monsoon season and the genesis of onset vortex in the South-Eastern Arabian Sea, *Q. J. R. Meteorol. Soc.*, 125, <https://doi.org/10.1002/qj.49712555503>, 1999.
- Rao, R. R., Jitendra, V., GirishKumar, M. S., Ravichandran, M., and Ramakrishna, S. S. V. S.: Interannual variability of the Arabian Sea Warm Pool: observations and governing mechanisms, *Clim. Dyn.*, 44, 2119-2136, 10.1007/s00382-014-2243-0, 2015.
- 510 Sabu, P. and Ravichandran, C.: Mixed layer processes of the Arabian Sea Warm Pool during spring intermonsoon: a study based on observational and satellite data, *Int. J. Remote. Sens.*, 32, 5425 - 5441, <https://doi.org/10.1080/01431161.2010.501350>, 2011.
- 515 Saji, N. H., Goswami, B. N., Vinayachandran, P. N., and Yamagata, T.: A dipole mode in the tropical Indian Ocean, *Nature*, 401, 360-363, 10.1038/43854, 1999.
- Sanilkumar, K. V. P., Hareesh Kumar & Joseph, Jossia & Panigrahi, Jitendra: Arabian Sea mini warm pool during May 2000, *Curr. Sci.*, 86, <http://www.jstor.org/stable/24109531>, 2004.
- Schott, F. A., Xie, S.-P., and McCreary Jr., J. P.: Indian Ocean circulation and climate variability, *Rev. Geophys.*, 47, <https://doi.org/10.1029/2007RG000245>, 2009.
- 520 Seetaramayya, P. and Master, A.: Observed air-sea interface conditions and a monsoon depression during MONEX-79, *Arch. Meteorol., Geophys. Bioklimatol., Ser. A.*, 33, 61-67, 10.1007/BF02265431, 1984.
- Sengupta, D., Parampil, S. R., Bhat, G. S., Murty, V. S. N., Ramesh Babu, V., Sudhakar, T., Premkumar, K., and Pradhan, Y.: Warm pool thermodynamics from the Arabian Sea Monsoon Experiment (ARMEX), *J. Geophys. Res.: Oceans.*, 113, <https://doi.org/10.1029/2007JC004623>, 2008.
- 525 Shankar, D. and Shetye, S. R.: On the dynamics of the Lakshadweep high and low in the southeastern Arabian Sea, *J. Geophys. Res.: Oceans.*, 102, 12551-12562, <https://doi.org/10.1029/97JC00465>, 1997.
- Shankar, D., Vinayachandran, P. N., and Unnikrishnan, A. S.: The monsoon currents in the north Indian Ocean, *Prog. Oceanogr.*, 52, 63-120, [https://doi.org/10.1016/S0079-6611\(02\)00024-1](https://doi.org/10.1016/S0079-6611(02)00024-1), 2002.

删除: :

删除: t

- 530 Shetye, S. R., Gouveia, A. D., Shankar, D., Shenoi, S. S. C., Vinayachandran, P. N., Sundar, D., Michael, G. S., and Nampoothiri, G.: Hydrography and circulation in the western Bay of Bengal during the northeast monsoon, *J. Geophys. Res.: Oceans.*, 101, 14011-14025, <https://doi.org/10.1029/95JC03307>, 1996.
- Song, Q., Vecchi, G. A., and Rosati, A. J.: Indian Ocean Variability in the GFDL Coupled Climate Model, *J. Clim.*, 20, 2895-2916, <https://doi.org/10.1175/JCLI4159.1>, 2007.
- 535 Sprintall, J. and Tomczak, M.: Evidence of the barrier layer in the surface layer of the tropics, *J. Geophys. Res.: Oceans.*, 97, 7305-7316, <https://doi.org/10.1029/92JC00407>, 1992.
- Sullivan, S. C., Schiro, K. A., Stubenrauch, C., and Gentine, P.: The Response of Tropical Organized Convection to El Niño Warming, *Journal of Geophysical Research: Atmospheres*, 124, 8481-8500, <https://doi.org/10.1029/2019JD031026>, 2019.
- Thandlam, V., Rahaman, H., Rutgersson, A., Sahlee, E., Ravichandran, M., and Ramakrishna, S. S. V. S.: Quantifying the role of antecedent Southwestern Indian Ocean capacitance on the summer monsoon rainfall variability over homogeneous regions of India, *Scientific Reports*, 13, 5553, [10.1038/s41598-023-32840-w](https://doi.org/10.1038/s41598-023-32840-w), 2023.
- 540 Thandlam, V., T.V.S, U. B., Hasibur, R., Luca, P. D., Sahlée, E., Rutgersson, A., M, R., and S.S.V.S, R.: A sea-level monopole in the equatorial Indian Ocean, *npj Climate and Atmospheric Science*, 3, 25, [10.1038/s41612-020-0127-z](https://doi.org/10.1038/s41612-020-0127-z), 2020.
- Thompson, B., Gnanaseelan, C., and Salvekar, P. S.: Variability in the Indian Ocean circulation and salinity and its impact on SST anomalies during dipole events, *J.Mar.Res*, 64, 853-880, 2006.
- 545 Venugopal, T., Ali, M. M., Bourassa, M. A., Zheng, Y., Goni, G. J., Foltz, G. R., and Rajeevan, M.: Statistical Evidence for the Role of Southwestern Indian Ocean Heat Content in the Indian Summer Monsoon Rainfall, *Scientific Reports*, 8, 12092, [10.1038/s41598-018-30552-0](https://doi.org/10.1038/s41598-018-30552-0), 2018.
- Webster, P. J. and Lukas, R.: TOGA COARE: The Coupled Ocean–Atmosphere Response Experiment, *Bull. Am. Meteorol. Soc.*, 73, 1377-1416, [https://doi.org/10.1175/1520-0477\(1992\)073<1377:TCTCOR>2.0.CO;2](https://doi.org/10.1175/1520-0477(1992)073<1377:TCTCOR>2.0.CO;2), 1992.
- 550 Webster, P. J., Moore, A. M., Loschnigg, J., and Leben, R. R.: Coupled ocean–atmosphere dynamics in the Indian Ocean during 1997–98, *Nature*, 401, 356-360, [10.1038/43848](https://doi.org/10.1038/43848), 1999.

555