

Climatologically invariant scale-invariance of cloud horizontal sizes

Thomas D. DeWitt¹, Timothy J. Garrett¹, Karlie N. Rees¹, Corey Bois¹, Steven K. Krueger¹, and Nicolas Ferlay²

¹Department of Atmospheric Sciences, University of Utah, 135 S 1460 E, Salt Lake City, UT 84112, USA;

²Université de Lille, UMR 8518 - LOA - Laboratoire d'Optique Atmosphérique, F-59000, Lille, France;

Correspondence: tim.garrett@utah.edu

Abstract.

Cloud area distributions are a defining feature of Earth's radiative exchanges with outer space. Cloud perimeter distributions $n(p)$ are also interesting because the shared interface between clouds and clear-sky determines exchanges of buoyant energy and air. Here, we test using detailed model output and a wide range of satellite datasets, a first-principles derivation that perimeter distributions follow a scale invariant power-law $n(p) \propto p^{-(1+\beta)}$ with $\beta = 1$, for perimeters evaluated within moist isentropic atmospheric layers. In model analyses, the value of β is closely reproduced. In satellite data β is remarkably robust to latitude, season, and land/ocean contrasts, which suggests that, at least statistically speaking, cloud perimeter distributions are determined more by atmospheric stability than Coriolis forces, surface temperature, or contrasts in aerosol loading between continental and marine environments. However, the satellite-measured value of β is found to be 1.29 ± 0.05 rather than $\beta = 1$. The reason for the discrepancy is unclear but comparison with a model reproduction of the satellite perspective suggests it may owe to cloud overlap. Satellite observations also show scale invariance governs cloud areas for a range at least as large as $\sim 3\text{km}^2$ to $\sim 3 \times 10^5\text{km}^2$. Many prior studies observed a much smaller range and we argue this difference is due to inappropriate treatments of the statistics of clouds that are truncated by the edge of the measurement domain.

1 Introduction

Since the first numerical global climate models (GCMs) were developed in the 1960s, there have been exponential advances in computational capabilities that have led to spectacular simulations of cloud structures. The next generation of climate models is expected to resolve individual clouds at kilometer scales (Schär et al., 2020). The strategy behind this “bottom-up” approach to representing the role of clouds in climate is that pursuing ever finer spatial resolution and improved model physics will lead to more accurate predictions, accepting the necessary evil of increased computational expense (Slingo et al., 2022). Yet, perhaps alarmingly, it has not been clear that this approach has been successful in its goal given that the spread in GCM predictions of the climate sensitivity to greenhouse gases has, if anything, only increased (Palmer, 2016; Arias et al., 2021; Lovejoy, 2022).

In some sense, time-dependent deterministic simulations are not obviously well suited for obtaining a statistical time-independent climatology. An alternative approach might be to derive the statistics directly, using principles of statistical thermodynamics, from bulk physical constraints (Arakawa, 2004; Procyk et al., 2022). A familiar example is the simplicity of the derivation of the Maxwell-Boltzmann statistics characterizing the distribution of speeds of molecules in an ideal gas, obtained

knowing only the average energy per molecule and without deterministically simulating individual particles and the extraordinary complexities of their quantum mechanical interactions (Schroeder, 2021). There is some evidence that this “top-down” philosophy may work for convective cloud fields. An exponential distribution of mass fluxes can be derived for non-interacting clouds by considering only the large-scale vertical mass flux (Cohen and Craig, 2006; Craig and Cohen, 2006). Another study
30 by Garrett et al. (2018) took a similar top-down approach but allowed for cloud interactions. It obtained a distribution of cloud horizontal sizes at steady-state that follows an exponential in saturated static energy and a power-law with respect to cloud perimeter.

In this study, we use a range of satellite observations to test the validity of the cloud perimeter distribution derived by Garrett et al. (2018). We show that both cloud perimeters and cloud areas do indeed follow a power-law, but that the power-law
35 exponent appears to be a function of perspective, agreeing well with theory in thin horizontal layers in cloud-resolving models but not to satellite observations of cloud fields looking down from space. We also find that the choice of domain size and treatment of clouds that are truncated by the domain edge can introduce spurious scale breaks in power-law size distributions. We suggest previous results that do not account for these subtle effects should be interpreted with caution.

This paper is organized as follows. In Sect. 2, we first overview the theoretical arguments presented by Garrett et al. (2018)
40 that led to the predicted cloud perimeter distribution. With this necessary background, prior empirical measurements of the related cloud area distribution are then discussed, along with the subtleties involved in measuring distributions of cloud sizes. The methods are presented in Sect. 3 and results from satellite observations in Sect. 4. In Sect. 5, we examine the role of perspective in measuring cloud size distributions and finally conclude in Sect. 6.

2 A steady-state thermodynamic model for cloud size distributions

45 To begin, we justify why it is physically meaningful to look at cloud perimeters by summarizing the derivation of the cloud perimeter number distribution $n(p)$ presented by Garrett et al. (2018). The foundation follows a parcel through an idealized thermodynamic cycle around cloud edges – what was termed a “mixing engine” – defined by four “legs”:

1. Moist adiabatic ascent inside cloud
2. Diabatic mixing with clear air across cloud edge that dries the parcel and reduces cloud perimeter
- 50 3. Dry adiabatic clear-sky descent
4. Diabatic mixing with cloudy air across cloud edge that moistens the parcel and lengthens cloud perimeter

The cycle is analogous to the familiar Carnot cycle, used to describe hurricanes (Emanuel, 1991), but with entropy generation associated with mixing at cloud edge rather than with energetic exchanges with the oceans or outer space. In observations of tropical convection, Heus and Jonker (2008) found that shallow cumulus tend to have a neutrally buoyant cloud edge and
55 a “subsiding shell” of descending clear air adjacent to cloud edge. A similar pattern was later observed in local circulations around simulated deep convection (Glenn and Krueger, 2014). These observations appear to support the mixing engine framework, at least for actively convecting clouds.

Representing mass fluxes across cloud edges in 4-D space-time coordinates, as is typically done in detailed cloud numerical simulations, is difficult because turbulent mixing changes both the location and length of the cloud edge itself, and over a very wide range of time and space scales. However, while cloud edge may deform during mixing, it maintains its position as a point of approximate neutral buoyancy, and in this sense can serve as a fixed reference point in a related coordinate system. For this purpose, we use the moist static energy, which is given by

$$h = gz + c_p T + L_v q, \quad (1)$$

where g , c_p , and L_v are the gravitational acceleration, the specific heat of air at constant pressure, and the latent heat of vaporization of water, respectively, and z , T , and q are height, temperature, and the water vapor mixing ratio, respectively. At cloud edge, air is just saturated, so the moist static energy is equal to the *saturated* static energy, defined as $h^* = h(q = q^*)$ where q^* is the saturated mixing ratio. At a given height, perturbations in saturated static energy can be related to temperature (and hence buoyancy) perturbations T' through $h^{*'} = c_p(1 + \gamma)T'$ where $\gamma = L/c_p \partial q^*/\partial T$ (Randall, 1980).

In a tropical atmosphere, variability in h^* between horizontal levels dominates variability within a given level, so a constant h^* surface can be approximated as lying along a surface of constant z (Xu and Emanuel, 1989). Supposing a thin atmospheric layer of thickness δz , clouds within this layer can be partitioned into discrete bins j of mean perimeter p_j . For a number n_j of such clouds, each bin has a total cloud perimeter $n_j p_j$ and a total surface area $\sigma = n_j p_j \delta z$. σ measures the component of the overall cloud surface area which is vertically oriented. Fick's law suggests that for bin j , the total rate of dissipation of potential energy across cloud edge Q_j due to diabatic turbulent mixing is proportional to the product of the energy gradient between cloudy and clear air ∇h and the total surface area σ (Garrett, 2012; Garrett et al., 2018). Provided the perturbation from the domain mean δh is much smaller than the mean value $\langle h^* \rangle$, a constraint that is satisfied even over the entire depth of the troposphere, and that turbulence around cloud edges is approximately isotropic (Heus and Jonker, 2008; Heus et al., 2009; Wang et al., 2009), the vertical and horizontal legs of the mixing engine are approximately the same size, so $\delta x \approx \delta z$ and $\nabla h \approx \delta h/\delta z = S$ where S is the stability. Thus the rate of dissipation of energy due to horizontal mixing across cloud edge in any given size bin j is

$$Q_j \propto n_j p_j \delta h. \quad (2)$$

In any cloud field, clouds continually grow and shrink due to turbulent mixing processes, and so cloud number is passed from one perimeter bin j to the next $j + 1$ or $j - 1$. At steady-state, however, which can be defined as a time-invariant perimeter distribution, there must be no net convergence of cloud number or energy into any bin j . This implies that $dQ/dp = 0$, or in discretized form from Eq. 2, that $n_j p_j = \text{const}$. The steady-state perimeter distribution $n(p)$ can therefore be expected to follow the power-law (Garrett et al., 2018)

$$n(p) \equiv \frac{dn}{dp} \propto p^{-(1+\beta)}, \quad \beta = 1, \quad p_{\min} < p < p_{\max}. \quad (3)$$

Power-laws such as Eq. 3 are generally considered to be “scaling”, since a rescaling of p by some constant factor c results in a constant rescaling of $n(p)$ by a constant factor $c^{-(1+\beta)}$. Of course, it is impossible for any physical system to exhibit scale

90 invariance over an infinite range of scales, and so such scaling behavior can only be valid over a finite range $p_{\min} < p < p_{\max}$. Beyond these “scale breaks”, the value of β changes or the functional form of the distribution changes. As an example, a common feature of power-law distributions describing many other natural and social systems is an exponential cutoff at large scales (Newman, 2005; Clauset et al., 2009).

For clouds, as a guess, the smallest possible size defining p_{\min} might be the Kolmogorov microscale for turbulent circulations
 95 such as in the mixing engine, whose order of magnitude is ~ 1 mm (Tennekes and Lumley, 1972). The largest possible clouds are of course limited by the Earth’s circumference of $\sim 10^5$ km, but might be more reasonably constrained by the Rossby radius of deformation $\sim 10^3$ km where Coriolis forces limit horizontal spreading.

Since cloud edges are fractal (Lovejoy, 1982), calculated perimeter lengths depend on the chosen measurement resolution, so p_{\max} can be orders of magnitude larger than the distance from one end of a cloud to the other. A measure of maximum cloud
 100 size that is less resolution dependent is maximum cloud area, which is roughly $\mathcal{O}(\text{cloud length})^2$.

The continuous function $n(p)$ can be discretized into linearly spaced bins with constant Δp , in which case the slope on a plot with two logarithmic axes would be $-(1 + \beta)$. If logarithmically-binned, with constant $\Delta \ln p$, the slope of the power-law would be $-\beta$ because $dn/d \ln p = pn(p)$. We favor logarithmically-spaced bins as being better suited to describe the vast range of cloud sizes because linearly-spaced bins can be associated with poor sampling in large bins where a power-law is anticipated
 105 (White et al., 2008).

2.1 The challenge of measuring cloud size distributions

For the power law exponent β , Garrett et al. (2018) found $\beta = 1.06 \pm 0.02$ in a comparison with a highly detailed numerical simulation of a tropical cloud field, in close agreement with the theoretically expected value of $\beta = 1$. Cloud perimeter distributions have yet to be assessed observationally although cloud area distributions have been widely studied, generally revealing
 110 power-law distributions in both satellite observations (Cahalan and Joseph, 1989; Kuo et al., 1993; Benner and Curry, 1998; Koren et al., 2008; Wood and Field, 2011) and models (Neggers et al., 2003; Yamaguchi and Feingold, 2013; Neggers et al., 2019; Christensen and Driver, 2021), although not in every study (López, 1977). Assuming both cloud areas and perimeters are power-law distributed, the two quantities can be related by the scaling relationship

$$p = \text{const.} \times a^{D/2}. \quad (4)$$

115 D is often interpreted to be the fractal dimension D_f as it is formally defined by the relation $l \propto \xi^{1-D_f}$ relating how the measured length l of a fractal line such as cloud perimeter depends on the “ruler length” (or resolution) ξ used to measure it (Mandelbrot, 1982). Assuming the relationship $D = D_f$ is valid, the fractal dimension can be determined by fitting a linear regression between observations of $\ln \sqrt{a}$ and $\ln p$ (e.g. Lovejoy, 1982; Cahalan and Joseph, 1989; Siebesma and Jonker, 2000; Christensen and Driver, 2021).

120 Subsequent work has shown that for clouds D is not in fact strictly equivalent to the fractal dimension D_f . Batista-Tomás et al. (2016) and Peters et al. (2009) pointed out that adopting the equivalence $D = D_f$ requires holes in clouds to be excluded

from contributing to the cloud’s perimeter, as the fractal dimension is a property of a single curve (a cloud’s exterior perimeter) rather than an ensemble of curves (a cloud’s exterior perimeter, and the perimeter of each hole).

125 Furthermore, Imre (1992) showed that the constant pre-factor in Eq. 4 often itself scales with a . In this case, fitting a regression line to a scatterplot of $\ln \sqrt{a}$ vs. $\ln p$ would yield a value for D that implicitly includes a scaling contribution from the supposed “constant”.

Setting aside these details, a scaling of the form Eq. 4 nonetheless can be used to empirically relate cloud areas and perimeters, making the expression useful regardless of any particular interpretation of D . Here, it permits the perimeter size distribution Eq. 3 to be converted to a distribution in cloud area. Since $d \ln a \propto d \ln p$, $dn/d \ln a \propto dn/d \ln p$, and so

$$130 \quad n(a) \equiv \frac{dn}{da} \propto a^{-(1+\alpha)}, \quad \alpha = \frac{D\beta}{2}, \quad a_{\min} < a < a_{\max}. \quad (5)$$

Adopting $D \approx 4/3$ (Lovejoy, 1982; Siebesma and Jonker, 2000), noting that both higher (Cahalan and Joseph, 1989; Christensen and Driver, 2021) and lower (Cahalan and Joseph, 1989; Batista-Tomás et al., 2016) values have been measured, and $\beta = 1$ as proposed by (Garrett et al., 2018), then Eq. 5 yields $\alpha \approx 2/3$. By contrast, widely conflicting values are observed for α , as well as for the location of the scale break a_{\max} . Cahalan and Joseph (1989) and Benner and Curry (1998) found in satellite
135 observations values for a_{\max} ranging from 4 km^2 to between 0.28 km^2 and 0.62 km^2 , respectively. In large eddy simulations, Neggers et al. (2003) found scale breaks between $0.16 \text{ km}^2 \leq a_{\max} \leq 1.6 \text{ km}^2$.

For larger domains considered in other studies a_{\max} tends to be larger. Wood and Field (2011) found using MODIS satellite data that $\alpha = 0.87 \pm 0.03$ and $a_{\max} \gtrsim 10^6 \text{ km}^2$. Peters et al. (2009) found a scale break in mesoscale convective clusters at $a_{\max} \sim 10^5 \text{ km}^2$, although it depended on the value of a threshold based on column water vapor. Conversely, Christensen and
140 Driver (2021) found $a_{\max} \sim 10^6 \text{ km}^2$ for tropical deep convection. There is also variation in calculated values for α , with Koren et al. (2008) finding $\alpha = 0.3 \pm 0.1$ and Yamaguchi and Feingold (2013) finding $\alpha = 0.59$. Neither found evidence for a scale break a_{\max} although they considered smaller domains.

These conflicting results could reflect meteorological differences, as there is some evidence D , or α through Eq. 5, is itself dependent on cloud type and size (Cahalan and Joseph, 1989; Batista-Tomás et al., 2016). However, a largely overlooked
145 explanation for the surprising variance in values for α and a_{\max} is one of sampling bias. Larger clouds are more likely to be truncated by the edge of the measurement domain than small clouds, and if they are removed from the analysis, as is sometimes done, there can be a spurious scale break introduced to the size distribution. Such a scale break would depend only on the size of domain considered, rather than some intrinsic physical property of the cloud field itself. This spurious effect of domain size has also been found to influence the measured power-law exponent for idealized 1-D cloud sizes (Wood and Field, 2011).

150 Past studies generally do not mention how clouds truncated by the domain edge are treated or, in some cases, they simply remove them from analysis (e.g. Peters et al., 2009; Christensen and Driver, 2021). Plausibly, some of the inconsistencies seen in measured values of a_{\max} and α could owe to this measurement problem. For example, one seeming solution is to retain clouds that are truncated by the domain edge but measure only the portion of the cloud area that lies within the domain. With this approach, a portion of the given cloud’s area is necessarily omitted, likely placing it in a smaller size bin where counts are
155 consequently oversampled.

3 Methods

Our goal here is to test in satellite observations and models the hypotheses proposed by Garrett et al. (2018), namely that $\beta = 1$ as specified by Eq. 3 and that the value of β is the same for any cloud field at steady-state. Second, we attempt to address inconsistencies in previous observations of α and a_{\max} by appropriately accounting for bias introduced by the treatment of clouds truncated by the edge of a satellite measurement domain.

3.1 Satellite datasets

The satellite platforms used to image clouds in this study fall into two broad categories: full-disk and polar-orbiting. Full-disk images are effectively a snapshot of Earth taken from geostationary orbit or, in the case of EPIC, the L1 Lagrange point. Polar-orbiting sensors continuously scan a rectangular swath as they move poleward. Details about the datasets are summarized in Table 1.

Table 1. Satellite datasets used in this study.

Dataset name	Sensor name	View Type	Approx. nadir resolution	Longitude at nadir	Dates examined	Description of cloud mask algorithm
GOES -137°	ABI	Full-Disk	2 km	137° W	01 January 2021 to 01 January 2022	Derrien and Gléau (2005, 2010)
GOES -75°	ABI	Full-Disk	2 km	75° W	01 January 2021 to 01 January 2022	Derrien and Gléau (2005, 2010)
MSG 0°	SEVIRI	Full-Disk	3 km	0°	01 January 2021 to 01 January 2022	Derrien and Gléau (2005, 2010)
MSG 42°	SEVIRI	Full-Disk	3 km	42° E	01 January 2021 to 01 January 2022	Derrien and Gléau (2005, 2010)
Himawari 141°	AHI	Full-Disk	2 km	141° E	01 January 2021 to 01 January 2022	Derrien and Gléau (2005, 2010)
EPIC	EPIC	Full-Disk	8 km	-	01 January 2017 to 01 January 2018	Yang et al. (2019)
VIIRS	VIIRS	Polar-Orbiting	0.75 km	-	01 January 2021 to 01 January 2022	Kopp et al. (2014)
MODIS 1 km	MODIS	Polar-Orbiting	1 km	-	01 January 2012 to 01 January 2013	Ackerman et al. (1998, 2008)
MODIS 0.25 km	MODIS	Polar-Orbiting	0.25 km	-	01 January 2021 to 10 January 2021	Section 3.1
POLDER	POLDER	Polar-Orbiting	1/18°	-	01 January 2012 to 01 January 2013	Buriez et al. (1997)

For most of the satellite datasets described in Table 1, individual clouds are identified from pre-processed binary cloud masks designed to distinguish cloudy and clear sky. The definition of a cloud is somewhat subjective, and so inevitable differences in cloud identification algorithms and sensor capabilities lead to variations in global cloud coverage estimates between datasets. Even for a given satellite dataset, estimates of global cloud fraction depend on the choice of viewing angle, increasing with
170 more oblique perspectives (Maddux et al., 2010). To mitigate this concern, images are truncated to exclude cloud imagery where the sensor zenith angle is greater than 60° , a choice intended as a compromise between limiting sensitivity to viewing angle while retaining a large domain area.

To test the sensitivity of measured distributions of cloud sizes to cloud definition, we also use a simple cloud mask based on MODIS band 1 optical reflectance R , which is sensitive to wavelengths between 620nm and 670nm and has a resolu-
175 tion at nadir of 0.25km. We examine 13 tropical maritime granules, each centered between approximately 10°S and 20°N and 115°W to 140°W and covering an area approximately 1950km wide by 2030km long. Images from each granule were visually inspected for artifacts from sun glint, and several additional granules were omitted from the analysis due to sun glint contamination. Figure 1 compares several example cloud masks generated using various thresholds in R alongside the pre-processed cloud mask and an RGB image.

180 3.2 SAM numerical simulations

For numerical simulations of cloud fields, we use output from the System for Atmospheric Modeling (SAM) (Khairoutdinov and Randall, 2003). SAM was initialized and forced by large-scale thermodynamic tendencies derived from mean conditions during the GATE Phase III field experiment (Khairoutdinov et al., 2009) and run with prescribed radiative heating, diagnostic subgrid-scale turbulence, and two prognostic hydrometeor variables (precipitating and non-precipitating) from which cloud
185 water, cloud ice, rain, snow, and graupel are diagnosed. The simulation’s domain size is $204.8\text{km} \times 204.8\text{km}$ with 100m horizontal grid spacing and a 2s time step. The vertical grid spacing is 50m below $z = 1.2\text{km}$ and increases to 100m at $z = 5\text{km}$. There are a total of 210 vertical levels.

Shallow cumulus form in the first hour of the simulation, gradually deepening into deep convection by hour 6. Beyond approximately hour 12, a steady-state period is reached where the convection is in quasi-equilibrium with the prescribed large-scale
190 forcing (Arakawa and Schubert, 1974; Lord and Arakawa, 1980; Lord, 1982). During this steady-state period, the precipitation rate and cloud cover fluctuate without significant trends and the simulation does not self-aggregate (Khairoutdinov et al., 2009). We analyze hourly 3-D model output from hours 12 to 24. Output from this simulation was also used in Garrett et al. (2018) and is described in full detail in Khairoutdinov et al. (2009).

A cloud mask for each horizontal layer in the simulation was applied by setting all grid cells with non-precipitating cloud
195 condensate mixing ratios q_n (including both liquid and ice) in excess of 1% of the saturated mixing ratio q^* to cloudy and the remainder to clear. Once every grid cell is defined as either cloudy or clear, 2-D images were created by isolating each individual height level in the domain, creating 210 images for every time step. These images were then analyzed separately using the same method as the satellite imagery (described below). Isolating individual horizontal layers in this manner provides an approximate method of isolating constant h^* surfaces (Garrett et al., 2018). After perimeters were calculated and binned for




Figure 1. Example RGB image, pre-processed cloud mask, and cloud masks created from various thresholds in optical reflectance R for a single MODIS granule. In the reflectance-based cloud masks, pixels with reflectance higher than the threshold are set to cloudy (white), while the others are set to clear (dark blue). The image is centered at approximately 1° S, 130° W and was taken at approximately 01 January 2021, 19:05 UTC. Note that pixels are depicted here as being uniform in size but that cloud size calculations account for pixel size increasing away from nadir.

200 each layer, counts were summed over all layers. We also create a “satellite-like” image from the simulation, which is described in Section 5.

3.3 Cloud identification and filtering

Both the satellite and model datasets yield 2-D binary cloud masks where grid cells or pixels are either cloudy or clear. Individual clouds are defined as connected cloudy regions, identified by applying a convention that adjacent cloudy pixels are
205 connected whereas diagonal cloudy pixels are not (termed “4-connectivity”; Wood and Field (2011); Christensen and Driver (2021)), although the analysis is not significantly affected by this choice (Kuo et al., 1993). In the satellite datasets, the pixel lengths in the x and y directions are determined independently as a function of satellite distance and sensor zenith angle.

Cloud perimeter is then computed by summing all pixel side lengths along the edge of the cloud, and cloud area by summing the areas of each individual cloudy pixel. Cloud holes add to the clouds’ perimeter but reduce its area, which as described
210 above implies $D \neq D_f$ in Eq. 4. Clouds consisting of a small number of pixels are more Euclidean than fractal (Christensen and Driver, 2021), which leads to an inaccurate estimate of the small portion of the size distribution. We therefore truncate number distributions to exclude cloud perimeters $\leq 10 \times (\text{resolution at nadir})$ or areas $\leq 10 \times (\text{resolution at nadir})^2$.

Unexpectedly the smaller portion of the size distributions obtained using EPIC display non-power-law behavior, in contrast to all other satellite datasets over similar scales. To ensure values for α and β were calculated over only the power-law
215 regime, thresholds used for EPIC clouds were increased to exclude perimeters $\leq 30 \times (\text{nadir resolution})$ or areas $\leq 1000 \times (\text{nadir resolution})^2$. Conceivably, the discrepancy is caused by a compression algorithm that averages 2×2 pixel regions before data transmission. The regions are subsequently interpolated back to the original resolution, which may smooth cloud perimeters (see Appendix A for further discussion).

To account for possible scale breaks in size distributions introduced by clouds truncated by the edge of the measurement
220 domain, area or perimeter bins in which the number of clouds truncated by the edge is greater than 50% of the total in that bin are removed from consideration. For the observed cloud fields, such bins tend to be those in the larger end of the size spectrum as large clouds are most likely to touch the domain edge. The threshold choice of 50% represents a compromise, removing bins most sensitive to truncation effects while allowing for a large range of cloud sizes to be studied. Calculated values for α and β are relatively insensitive to more stringent thresholds less than 50%.

225 We calculate the power-law exponents α and β by performing a linear regression in logarithmic space since from Eqs. 3 and 5 $\ln n(p) = -\beta \ln p + \text{const.}$ and $\ln n(a) = -\alpha \ln a + \text{const.}$ It has been argued this method can lead to underestimates of the exponent (Clauset et al., 2009), but there is no straightforward alternative when presented, as is the case here, with a power-law that has both an upper and lower bound (Hanel et al., 2017). We evaluate uncertainties as 95% confidence intervals corresponding to two standard errors of the regression. We report numerical values of plotted data in the Supplemental
230 Information.




Figure 2. Left: Logarithmically-binned histograms of cloud areas for satellite datasets. Right: Measured values of the power-law exponent α (Eq. 5) with associated mean (gray line) and 95% confidence interval (gray box). Counts have been vertically offset for clarity. Uncertainties represent 95% confidence intervals, derived from a linear regression standard error analysis.

4 Measured cloud size distributions

In satellite observations examined here, both cloud areas and perimeters are well described by a power-law distribution. For cloud areas, Fig. 2 shows measured values of α ranging between $\alpha = 0.90 \pm 0.02$ (POLDER and MODIS 0.25 km) and $\alpha = 0.99 \pm 0.02$ (GOES -75° and Himawari 141°) and a mean value, across all satellite datasets, of $\langle \alpha \rangle = 0.95 \pm 0.08$. These values
 235 are largely in agreement with several previous studies (e.g. Cahalan and Joseph (1989), Benner and Curry (1998), and Wood and Field (2011)).

For cloud perimeters, Fig. 3 shows values of β ranging from $\beta = 1.22 \pm 0.02$ (MODIS 1 km and 0.25 km) to $\beta = 1.316 \pm 0.008$ (GOES -75°), with a mean across all satellite datasets of $\langle \beta \rangle = 1.26 \pm 0.06$. This value differs from that found in SAM horizontal levels, where $\beta = 0.98 \pm 0.03$, and from the theoretically-derived value of $\beta = 1$ (Eq. 3).

240 These mean values of α and β imply, from Eq. 4, that $D = 1.5 \pm 0.1$, which is in good agreement with prior studies that have generally found values of D slightly greater than $4/3$, e.g. $D = 1.35$ (Lovejoy, 1982), $1.25 \leq D \leq 1.59$ (Cahalan and Joseph, 1989), or $D = 1.4$ (Christensen and Driver, 2021).

After omitting bins containing 50% or more clouds truncated by the domain edge, a scale break a_{\max} is no longer evident in the area distributions. In several cases, the distributions exhibit scale invariance extending to areas larger than 10^5 km^2 , with the largest to at least $\sim 3 \times 10^5 \text{ km}^2$ (EPIC). We find that a_{\max} must therefore have a value larger than roughly $3 \times 10^5 \text{ km}^2$,
 245 corresponding to an effective diameter of $\sim 600 \text{ km}$, substantially larger than some have previously suggested (e.g. Cahalan and Joseph, 1989; Benner and Curry, 1998; Neggers et al., 2003), with Wood and Field (2011) extending a_{\max} to 10^6 km^2 .




Figure 3. As in Fig. 2, but considering cloud perimeters (Eqn 3) and including results from SAM horizontal levels. The grey line and box on the right plot represent the mean and 95% confidence interval, respectively, across only the satellite datasets (excluding SAM).

4.1 Variability with seasonality, latitude, and surface type

The perimeter size distribution, given by Eq. 3, was derived without explicit consideration of local climatological characteristics such as season, surface type, or latitudinal location (Garrett et al., 2018). In satellite observations, the sensitivity of β to such considerations is shown in Figs. 4, 5, and 6. Comparisons between latitude bands shown in Figs. 4 and 5 are restricted to observations using the polar-orbiting satellites MODIS and VIIRS because imagery from these sensors, regardless of latitudinal location, is both similar in domain area and always centered directly below the satellite. These conditions reduce the likelihood of bias due to differing viewing geometry, and neither condition holds for the full disk images. POLDER has been omitted from Figs. 4, 5, and 6 because, when limited to smaller domains, its smaller sample size introduces significant statistical variability.

Independent of sensor, measured values of β appear robust across latitudinal regions, land/ocean contrasts, and seasons. Fig. 5 does show modest variability in the value of β by month in the midlatitude regions 60°S - 30°S and 30°N - 60°N between a minimum value of $\beta = 1.21 \pm 0.03$ (MODIS, March, May, June, northern midlatitudes) and a maximum value of 1.32 ± 0.02 (MODIS, June, July, southern midlatitudes). Annual mean values of β for the midlatitude and equatorial regions in Fig. 4 show similar values ranging from $\beta = 1.22 \pm 0.03$ for MODIS at northern midlatitudes to $\beta = 1.28 \pm 0.03$ for VIIRS in all regions and MODIS at southern midlatitudes. Separating clouds by marine and continental regions in Fig. 6, mean values for β are 1.25 ± 0.05 for land and 1.28 ± 0.04 for ocean. All values are consistent with the global mean value across datasets of $\langle \beta \rangle = 1.26 \pm 0.06$.




Figure 4. Measured values of β (Eq. 3), separated into the northern midlatitude region (left), the equatorial region (middle), and the southern midlatitude region (right). The grey line and box represent the global mean and 95% confidence interval, respectively, from Fig. 3.




Figure 5. Measured values of β (Eq. 3), for the northern midlatitude region (left) and the southern midlatitude region (right), separated by month. The equatorial region (not shown) shows similar variability around a mean value shown in Fig. 4. The grey line and box represent the global mean and 95% confidence interval, respectively, from Fig. 3.




Figure 6. Measured values of β (Eq. 3), separated into clouds over land (left) and clouds over ocean (right). The grey line and box represent the global mean and 95% confidence interval, respectively, from Fig. 3.

5 Discussion

265 After accounting for spurious scale breaks introduced by the problem of attempting to measure the extent of scale invariance with a finite domain, we find that a power-law describes the distributions of both cloud perimeters and areas for a size range spanning four and five orders of magnitude, respectively, and likely extends even further. This result is perhaps all the more remarkable for the fact that the value of the exponent β appears to be robust to such local climatological characteristics as season, latitude, land/ocean contrasts, or latitude as might be related to surface temperature, the Coriolis force, dominant cloud
270 type, or aerosol loading. In this sense, the observations appear to lend support to the general theoretical “mixing-engine” approach employed by Garrett et al. (2018) to obtain Eq. 3, where β was derived only by considering mixing processes at cloud edge.

However, a puzzle remains: the global mean value of $\langle\beta\rangle = 1.26 \pm 0.06$ in satellite observations is higher than the value of $\beta \simeq 1$ obtained both theoretically (Eq. 3) and from SAM model simulations (Fig. 3). The difference is significant given
275 the range of scales in cloud sizes involved. For example, for a roughly three order of magnitude measured range for cloud perimeters, the discrepancy would imply an order of magnitude difference in cloud counts.

The most obvious inference is that the theory is missing something fundamental about what determines cloud perimeters, even if it produced perimeter distribution values of β very close to those seen in a highly detailed numerical cloud model. Alternatively, one important distinction that may be made between the two approaches is simply one of perspective. Perimeter
280 distributions from the numerical model SAM shown in Fig. 3 and previously in Garrett et al. (2018) were calculated by treating every individual horizontal layer in the SAM volume as an independent 2D image. Only after each cloud perimeter was calculated and binned were the counts summed over all layers to create a single histogram, with no account made for cloud overlap. We term this method “layers”.




Figure 7. Example comparison of two methods of measuring perimeters of a 3-D cloud in SAM. The “compress” method creates a 2-D image by vertically summing cloud properties, resulting in a single image for each volume representing how clouds might be seen from above. In contrast, the “layers” method used in Garrett et al. (2018) considers each horizontal slice as a separate image such that for n horizontal layers, n individual images would be produced and analyzed as independent images. In the example, the “compressed” method would produce one cloud with $p = 12$ pixels and the “layers” method would produce three clouds, one with $p = 10$ and two with $p = 4$.

Satellite imagery differs, as it offers a two-dimensional representation of a cloud field as seen from above rather than within. Any vertical cloud overlap is effectively “compressed” into a single horizontal plane, *before* individual cloud perimeters are calculated. No distinction is made between overlapping clouds and vertically continuous clouds.

For example, the idealized cloud field in Fig. 7 yields a single cloud with $p = 12$ in the compressed satellite view (b), whereas a layers analysis would see three clouds, two in layer (c), each with $p = 4$, and one in layer (d), with $p = 10$. A priori, we might therefore expect a compressed image to yield relatively fewer small clouds than the layers case, as is the case in the example. This would result in a smaller value of β for the compressed case relative to the layers case. Counterintuitively, however, the opposite appears true: the value of β is larger in the compressed satellite perimeter distribution than in the layered SAM distribution.

The difference between the two perspectives can be manufactured in SAM by creating vertically compressed images as they might be seen by a satellite from above. Here, this is accomplished by creating a 2D vertically-summed optical depth (τ) field, to which a range of optical depth thresholds is applied to create a selection of cloud masks. Once binary cloud masks are created, clouds are identified and analyzed as described in Sect. 3.3.




Figure 8. Left: Measurements of β for individual horizontal layers (“layers”) in SAM for varying thresholds in total cloud condensate q_n , normalized by the saturated mixing ratio q^* . Middle: Measured values of β for “compressed” images in SAM for varying model τ thresholds, created by vertically summing modeled τ . The middle inset displays β vs. τ for the compressed SAM data, using a logarithmically scaled abscissa, over a larger range in τ . Right: Measured values of β for cloud masks of varying reflectance (R) thresholds for MODIS 0.25 km data. See Figure 7 for a visualization of the difference between “compressed” images and “layers”. The grey line and box indicate the global mean $\langle\beta\rangle = 1.26 \pm 0.06$ (Figure 3). Histograms from which values for β are calculated are shown in S2.

Fig. 8 shows that values of β , calculated using the “layers” method in SAM, are consistent with the theoretical prediction $\beta = 1$ regardless of the threshold used to define cloud, but that the value of β does indeed increase when the perspective is switched to one in which the clouds are vertically compressed as they might be seen from space.

300 However, as the compressed threshold in τ grows, β decreases, reaching a value of roughly 1 at $\tau \gtrsim 10$. Such sensitivity of β to optical depth threshold is at odds with observations, given that cloud masks specified by thresholds in reflectance between $R = 0.1$ and $R = 0.7$ for MODIS 0.25 km data show very little trend in calculated β . Note, for comparison, that the range of reflectance thresholds considered is roughly equivalent to a range of optical depths between $\tau = 1$ and $\tau = 10$, and that the reflectance thresholds in MODIS 0.25 km data generally produce values of β that are consistent with the global mean value
 305 derived from the pre-processed MODIS cloud mask.

6 Conclusions

By considering cloud edges as a surface across which cloudy and clear skies compete for available convective potential energy through small-scale mixing processes in a “mixing-engine”, Garrett et al. (2018) derived a cloud perimeter distribution that follows a power-law $n(p) \propto p^{-(1+\beta)}$, where $\beta = 1$ (Eq. 3), for perimeters evaluated within thin isentropic layers. The predic-

310 tion is independent of such considerations as the details of cloud microphysics or climatological state. We find in a detailed numerical simulation of a tropical cloud field that $\beta = 0.98 \pm 0.03$, which is consistent with the prediction.

In a wide range of satellite observations, however, the picture is more nuanced. Within measurement uncertainty, values of β are insensitive to zonal band, land/ocean contrasts, or season, conditionally supporting the small-scale mixing engine hypothesis. However, the globally averaged value across all satellite datasets is significantly higher than predicted by either
315 theory or model with a value of $\beta = 1.26 \pm 0.06$.

The discrepancy likely owes to a difference in perspective between cloud size distributions measured within individual quasi-horizontal moist isentropic layers, as was done with the numerical simulation, and those seen looking from above, as was calculated using satellite observations. The precise explanation remains a puzzle. We do see that values of β are higher in numerical simulations when the perspective is changed to one looking from above where clouds are defined by a threshold
320 in vertically summed optical depth. This may seem to help resolve the matter. But even here the picture is unclear since β approaches unity as the optical depth threshold increases and there is no similar sensitivity to reflectance threshold seen in MODIS observations.

Our results also suggest a warning for how future satellite missions are designed. The data compression algorithm used prior to transmission of EPIC data averages 2×2 pixel regions and then interpolates them back to the original resolution in post-
325 processing. We argue this approach may produce erroneous cloud size distributions that do not follow a power-law. Further work could determine whether the interpolation adversely impacts other calculated cloud properties.

Regardless, scale invariance appears to be a defining feature of clouds over at least four orders of magnitude in perimeter and five in area. We find, in satellite observations, that the upper limit of scale invariance in cloud area distributions a_{\max} has a value larger than $3 \times 10^5 \text{ km}^2$, a scale much larger than some other studies have suggested (e.g. Cahalan and Joseph, 1989;
330 Benner and Curry, 1998; Neggers et al., 2003) and close to that found in Wood and Field (2011).

The distribution of cloud areas at large scales remains difficult to measure due to domain size limitations. An intriguing possibility might be to synthesize geostationary data to produce a quasi-global cloud mask product. The product would be similar to existing aerosol optical depth maps (Ceamanos et al., 2021).

With a better understanding of cloud size distributions, at least statistically speaking, it may only be necessary to simulate
335 the counts of the largest clouds to predict the numbers of the smallest.

Code and data availability. Python code to analyze all data and generate all Figs. is available from the first author upon request. The VIIRS and EPIC datasets were downloaded from NASA Earthdata (NASA) and all others from the ICARE Data Center in Lille, France (ICARE).

Appendix A: EPIC data

Due to the inaccuracy of measuring cloud perimeters and areas consisting of a small number of pixels (Christensen and Driver,
340 2021), we remove all clouds with perimeters $\leq 10 \times (\text{nadir resolution})$ or areas $\leq 10 \times (\text{nadir resolution})^2$ (Sec. 3.3). If these




Figure A1. Cloud perimeter (left) and area (right) histograms for EPIC, omitting data where perimeters $\leq 10 \times (\text{nadir resolution})$ or areas $\leq 10 \times (\text{nadir resolution})^2$. These thresholds are the same as those used for other datasets; however, EPIC displays non-power-law behavior over the range left of the grey lines. In Figs. 2 and 3, we instead use the grey lines as minimum thresholds (that is, omit perimeters $\leq 30 \times (\text{nadir resolution})$ or areas $\leq 1000 \times (\text{nadir resolution})^2$).

same minimum thresholds are used for EPIC’s cloud size distributions, results show non-power-law size distributions for both area and perimeter at the small end of the size distribution (Fig. A1). This is in contrast to all other satellite datasets over similar size ranges (Figs. 2 and 3).

As a possible explanation for this discrepancy, EPIC imagery is compressed prior to transmission to Earth by averaging 2×2 pixel regions. These regions are then interpolated back to their original resolution in post-processing, artificially smoothing out the details of cloud perimeters. Since cloud perimeter lengths are resolution dependent, this results in an inaccurate perimeter measurement given EPIC’s resolution. Likewise, if the cloud signal in a cloudy pixel is “spread out” into neighboring clear pixels by the smoothing process, pixels that were originally cloudy may become clear or vice versa, so individual cloud areas are likely not conserved through the compression process. Conceivably, these effects may explain the anomalous EPIC area and perimeter distributions, though other differences in e.g. the cloud masking algorithm used may contribute.

It appears interpolation predominately affects measurements of area and perimeter in small clouds. To account for this inconsistency, we instead truncate EPIC’s size distributions where perimeters $\leq 30 \times (\text{nadir resolution})$ or areas $\leq 1000 \times (\text{nadir resolution})^2$. With these revised thresholds, results from EPIC roughly agree with those from other datasets (Figs. 2 and 3).

Author contributions. T.D.D.: Formal analysis, Methodology, Writing - original draft preparation. T.J.G.: Conceptualization, Funding Ac-
355 quisition, Supervision, Methodology, Writing - review and editing. K.N.R.: Methodology, Writing - review & editing. C.B.: Methodology,
Writing - review & editing. S.K.K: Funding acquisition, Writing - review & editing. N.F.: Formal analysis.

Competing interests. T.J.G. is a senior editor for Atmospheric Chemistry and Physics.

Acknowledgements. The Laboratoire d'Optique Atmosphérique kindly hosted visits by T.D.D and T.J.G. to l'Université de Lille in support
of scientific collaboration. The AERIS/ICARE Data and Services Center at l'Université de Lille and the Center for High Performance
360 Computing at the University of Utah provided data storage and computing services.

References

- Ackerman, S. A., Strabala, K., Menzel, W., Frey, R., Moeller, C., and Gumley, L.: Discriminating clear sky from clouds with MODIS, *J. Geophys. Res.*, 103, 32 141–32 157, 1998.
- Ackerman, S. A., Holz, R. E., Frey, R., Eloranta, E. W., Maddux, B. C., and McGill, M.: Cloud detection with MODIS. Part II: Validation, *J. Atmos. Oceanic Technol.*, 25, 1073–1086, 2008.
- 365 Arakawa, A.: The cumulus parameterization problem: Past, present, and future, *Journal of climate*, 17, 2493–2525, 2004.
- Arakawa, A. and Schubert, W. H.: Interaction of a Cumulus Cloud Ensemble with the Large-Scale Environment, Part I, *Journal of Atmospheric Sciences*, 31, 674 – 701, [https://doi.org/https://doi.org/10.1175/1520-0469\(1974\)031<0674:IOACCE>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1974)031<0674:IOACCE>2.0.CO;2), 1974.
- Arias, P., Bellouin, N., Coppola, E., Jones, R., Krinner, G., Marotzke, J., Naik, V., Palmer, M., Plattner, G.-K., Rogelj, J., Rojas, M., Sillmann, J., Storelvmo, T., Thorne, P., Trewin, B., Achuta Rao, K., Adhikary, B., Allan, R., Armour, K., Bala, G., Barimalala, R., Berger, S., Canadell, J., Cassou, C., Cherchi, A., Collins, W., Collins, W., Connors, S., Corti, S., Cruz, F., Dentener, F., Dereczynski, C., Di Luca, A., Diongue Niang, A., Doblas-Reyes, F., Dosio, A., Douville, H., Engelbrecht, F., Eyring, V., Fischer, E., Forster, P., Fox-Kemper, B., Fuglested, J., Fyfe, J., Gillett, N., Goldfarb, L., Gorodetskaya, I., Gutierrez, J., Hamdi, R., Hawkins, E., Hewitt, H., Hope, P., Islam, A., Jones, C., Kaufman, D., Kopp, R., Kosaka, Y., Kossin, J., Krakovska, S., Lee, J.-Y., Li, J., Mauritsen, T., Maycock, T., Meinshausen, M., Min, S.-K., Monteiro, P., Ngo-Duc, T., Otto, F., Pinto, I., Pirani, A., Raghavan, K., Ranasinghe, R., Ruane, A., Ruiz, L., Sallée, J.-B., Samset, B., Sathyendranath, S., Seneviratne, S., Sörensson, A., Szopa, S., Takayabu, I., Tréguier, A.-M., van den Hurk, B., Vautard, R., von Schuckmann, K., Zaehle, S., Zhang, X., and Zickfeld, K.: Technical Summary, pp. 33–144, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.002>, 2021.
- 370
- Batista-Tomás, A., Díaz, O., Batista-Leyva, A., and Altshuler, E.: Classification and dynamics of tropical clouds by their fractal dimension, *Quarterly Journal of the Royal Meteorological Society*, 142, 983–988, 2016.
- 380
- Benner, T. C. and Curry, J. A.: Characteristics of small tropical cumulus clouds and their impact on the environment, *Journal of Geophysical Research: Atmospheres*, 103, 28 753–28 767, 1998.
- Buriez, J. C., Vanbauce, C., Parol, F., Goloub, P., Herman, M., Bonnel, B., Fouquart, Y., Couvert, P., and Seze, G.: Cloud detection and derivation of cloud properties from POLDER, *International Journal of Remote Sensing*, 18, 2785–2813, <https://doi.org/10.1080/014311697217332>, 1997.
- 385
- Cahalan, R. F. and Joseph, J. H.: Fractal statistics of cloud fields, *Monthly weather review*, 117, 261–272, 1989.
- Ceamanos, X., Six, B., and Riedi, J.: Quasi-Global Maps of Daily Aerosol Optical Depth From a Ring of Five Geostationary Meteorological Satellites Using AERUS-GEO, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034 906, <https://doi.org/https://doi.org/10.1029/2021JD034906>, e2021JD034906 2021JD034906, 2021.
- 390
- Christensen, H. M. and Driver, O. G.: The Fractal Nature of Clouds in Global Storm-Resolving Models, *Geophysical Research Letters*, p. e2021GL095746, 2021.
- Clauset, A., Shalizi, C. R., and Newman, M. E.: Power-law distributions in empirical data, *SIAM review*, 51, 661–703, 2009.
- Cohen, B. G. and Craig, G. C.: Fluctuations in an equilibrium convective ensemble. Part II: Numerical experiments, *Journal of the atmospheric sciences*, 63, 2005–2015, 2006.
- 395
- Craig, G. C. and Cohen, B. G.: Fluctuations in an equilibrium convective ensemble. Part I: Theoretical formulation, *Journal of the atmospheric sciences*, 63, 1996–2004, 2006.

- Derrien, M. and Gléau, H. L.: MSG/SEVIRI cloud mask and type from SAFNWC, *International Journal of Remote Sensing*, 26, 4707–4732, <https://doi.org/10.1080/01431160500166128>, 2005.
- Derrien, M. and Gléau, H. L.: Improvement of cloud detection near sunrise and sunset by temporal-differencing and region-growing techniques with real-time SEVIRI, *International Journal of Remote Sensing*, 31, 1765–1780, <https://doi.org/10.1080/01431160902926632>, 2010.
- Emanuel, K. A.: The theory of hurricanes, *Annual Review of Fluid Mechanics*, 23, 179–196, 1991.
- Garrett, T. J.: Modes of growth in dynamic systems, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 468, 2532–2549, <https://doi.org/10.1098/rspa.2012.0039>, 2012.
- 405 Garrett, T. J., Glenn, I. B., and Krueger, S. K.: Thermodynamic constraints on the size distributions of tropical clouds, *Journal of Geophysical Research: Atmospheres*, 123, 8832–8849, 2018.
- Glenn, I. B. and Krueger, S. K.: Downdrafts in the near cloud environment of deep convective updrafts, *Journal of Advances in Modeling Earth Systems*, 6, 1–8, 2014.
- Hanel, R., Corominas-Murtra, B., Liu, B., and Thurner, S.: Fitting power-laws in empirical data with estimators that work for all exponents, *PloS one*, 12, e0170920, 2017.
- 410 Heus, T. and Jonker, H. J.: Subsiding shells around shallow cumulus clouds, *Journal of the Atmospheric Sciences*, 65, 1003–1018, 2008.
- Heus, T., J. Pols, C. F., J. Jonker, H. J., A. Van den Akker, H. E., and H. Lenschow, D.: Observational validation of the compensating mass flux through the shell around cumulus clouds, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 135, 101–112, 2009.
- 415 ICARE: ICARE Data and Services Center, <https://www.icare.univ-lille.fr/>, accessed: 2023-03-01.
- Imre, A.: Problems of measuring the fractal dimension by the slit-island method, *Scripta metallurgica et materialia*, 27, 1713–1716, 1992.
- Khairoutdinov, M. F. and Randall, D. A.: Cloud resolving modeling of the ARM summer 1997 IOP: Model formulation, results, uncertainties, and sensitivities, *Journal of the Atmospheric Sciences*, 60, 607–625, 2003.
- Khairoutdinov, M. F., Krueger, S. K., Moeng, C.-H., Bogenschutz, P. A., and Randall, D. A.: Large-Eddy Simulation of Maritime Deep Tropical Convection, *Journal of Advances in Modeling Earth Systems*, 1, <https://doi.org/https://doi.org/10.3894/JAMES.2009.1.15>, 2009.
- 420 Kopp, T. J., Thomas, W., Heidinger, A. K., Botambekov, D., Frey, R. A., Hutchison, K. D., Iisager, B. D., Brueske, K., and Reed, B.: The VIIRS Cloud Mask: Progress in the first year of S-NPP toward a common cloud detection scheme, *Journal of Geophysical Research: Atmospheres*, 119, 2441–2456, <https://doi.org/https://doi.org/10.1002/2013JD020458>, 2014.
- Koren, I., Oreopoulos, L., Feingold, G., Remer, L., and Altartatz, O.: How small is a small cloud?, *Atmospheric Chemistry and Physics*, 8, 3855–3864, 2008.
- 425 Kuo, K.-S., Welch, R. M., Weger, R. C., Engelstad, M. A., and Sengupta, S.: The three-dimensional structure of cumulus clouds over the ocean: 1. Structural analysis, *Journal of Geophysical Research: Atmospheres*, 98, 20685–20711, 1993.
- López, R. E.: The lognormal distribution and cumulus cloud populations, *Monthly Weather Review*, 105, 865–872, 1977.
- Lord, S. J.: Interaction of a Cumulus Cloud Ensemble with the Large-Scale Environment. Part III: Semi-Prognostic Test of the Arakawa-Schubert Cumulus Parameterization, *Journal of Atmospheric Sciences*, 39, 88 – 103, [https://doi.org/https://doi.org/10.1175/1520-0469\(1982\)039<0088:IOACCE>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1982)039<0088:IOACCE>2.0.CO;2), 1982.
- 430 Lord, S. J. and Arakawa, A.: Interaction of a Cumulus Cloud Ensemble with the Large-Scale Environment. Part II, *Journal of Atmospheric Sciences*, 37, 2677 – 2692, [https://doi.org/https://doi.org/10.1175/1520-0469\(1980\)037<2677:IOACCE>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1980)037<2677:IOACCE>2.0.CO;2), 1980.
- Lovejoy, S.: Area-perimeter relation for rain and cloud areas, *Science*, 216, 185–187, 1982.

- 435 Lovejoy, S.: The Future of Climate Modelling: Weather Details, Macroweather Stochastics – Or Both?, *Meteorology*, 1, 414–449, <https://doi.org/10.3390/meteorology1040027>, 2022.
- Maddux, B., Ackerman, S., and Platnick, S.: Viewing geometry dependencies in MODIS cloud products, *Journal of Atmospheric and Oceanic Technology*, 27, 1519–1528, 2010.
- Mandelbrot, B. B.: *The fractal geometry of nature*, vol. 1, WH freeman New York, 1982.
- 440 NASA: NASA Earthdata, <https://www.earthdata.nasa.gov/>, accessed: 2023-03-01.
- Negggers, R., Griewank, P., and Heus, T.: Power-law scaling in the internal variability of cumulus cloud size distributions due to subsampling and spatial organization, *Journal of the Atmospheric Sciences*, 76, 1489–1503, 2019.
- Negggers, R. A., Jonker, H. J., and Siebesma, A.: Size statistics of cumulus cloud populations in large-eddy simulations, *Journal of the atmospheric sciences*, 60, 1060–1074, 2003.
- 445 Newman, M. E.: Power laws, Pareto distributions and Zipf’s law, *Contemporary physics*, 46, 323–351, 2005.
- Palmer, T. N.: A personal perspective on modelling the climate system, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 472, 20150772, 2016.
- Peters, O., Neelin, J. D., and Nesbitt, S. W.: Mesoscale convective systems and critical clusters, *Journal of the atmospheric sciences*, 66, 2913–2924, 2009.
- 450 Procyk, R., Lovejoy, S., and Hébert, R.: The fractional energy balance equation for climate projections through 2100, *Earth System Dynamics*, 13, 81–107, <https://doi.org/10.5194/esd-13-81-2022>, 2022.
- Randall, D. A.: Conditional Instability of the First Kind Upside-Down, *Journal of Atmospheric Sciences*, 37, 125 – 130, [https://doi.org/https://doi.org/10.1175/1520-0469\(1980\)037<0125:CIOTFK>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1980)037<0125:CIOTFK>2.0.CO;2), 1980.
- Schär, C., Fuhrer, O., Arteaga, A., Ban, N., Charpiilloz, C., Di Girolamo, S., Hentgen, L., Hoefler, T., Lapillonne, X., Leutwyler, D., et al.: Kilometer-scale climate models: Prospects and challenges, *Bulletin of the American Meteorological Society*, 101, E567–E587, 2020.
- 455 Schroeder, D.: *An Introduction to Thermal Physics*, Oxford University Press, <https://physics.weber.edu/thermal/>, 2021.
- Siebesma, A. and Jonker, H.: Anomalous scaling of cumulus cloud boundaries, *Physical review letters*, 85, 214, 2000.
- Slingo, J., Bates, P., Bauer, P., Belcher, S., Palmer, T., Stephens, G., Stevens, B., Stocker, T., and Teutsch, G.: Ambitious partnership needed for reliable climate prediction, *Nature Climate Change*, 12, 499–503, 2022.
- 460 Tennekes, H. and Lumley, J. L.: *A first course in turbulence*, MIT press, 1972.
- Wang, Y., Geerts, B., and French, J.: Dynamics of the cumulus cloud margin: An observational study, *Journal of the atmospheric sciences*, 66, 3660–3677, 2009.
- White, E. P., Enquist, B. J., and Green, J. L.: On Estimating the Exponent Of Power-Law Frequency Distributions, *Ecology*, 89, 905–912, <https://doi.org/https://doi.org/10.1890/07-1288.1>, 2008.
- 465 Wood, R. and Field, P. R.: The distribution of cloud horizontal sizes, *Journal of Climate*, 24, 4800–4816, 2011.
- Xu, K.-M. and Emanuel, K. A.: Is the Tropical Atmosphere Conditionally Unstable?, *Monthly Weather Review*, 117, 1471 – 1479, [https://doi.org/https://doi.org/10.1175/1520-0493\(1989\)117<1471:ITTACU>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0493(1989)117<1471:ITTACU>2.0.CO;2), 1989.
- Yamaguchi, T. and Feingold, G.: On the size distribution of cloud holes in stratocumulus and their relationship to cloud-top entrainment, *Geophysical Research Letters*, 40, 2450–2454, 2013.
- 470 Yang, Y., Meyer, K., Wind, G., Zhou, Y., Marshak, A., Platnick, S., Min, Q., Davis, A. B., Joiner, J., Vasilkov, A., Duda, D., and Su, W.: Cloud products from the Earth Polychromatic Imaging Camera (EPIC): algorithms and initial evaluation, *Atmospheric Measurement Techniques*, 12, 2019–2031, <https://doi.org/10.5194/amt-12-2019-2019>, 2019.