

Surface Rupture Kinematics of the 2020 M_w 6.6 Masbate (Philippines) Earthquake determined from Optical and Radar Data

Khelly Shan Sta. Rita¹, Sotiris Valkaniotis², and Alfredo Mahar Francisco Lagmay^{1,3}

¹National Institute of Geological Sciences, University of the Philippines, Diliman, Quezon City, Philippines 1101

²Department of Civil Engineering, Democritus University of Thrace, Xanthi, Greece 67100

³UP Resilience Institute, University of the Philippines, Diliman, Quezon City, Philippines 1101

Correspondence: Khelly Shan Sta. Rita (kcstarita@up.edu.ph)

Abstract. Optical correlation, interferometry, and field investigation of laterally offset features were undertaken to analyze the kinematics of the 2020 M_w 6.6 Masbate earthquake. Coseismic displacement fields ~~from optical correlation show a maximum displacement of 0.61~~ show a peak left lateral offset of 0.6 m corresponding to M_w ~~6.64~~ 6.6 geodetic moment magnitude ~~and a lone asperity in Cataingan, with an average 0.4 m left lateral slip. The slip distributions also indicate a single asperity located~~ ~200 m SE of the centroid. Post-seismic deformation estimates from interferometry highlights ~~a maximum~~ an at least 0.14 m ~~sinistral displacement equivalent to left lateral offset consistent with a~~ M_w ~~6.15~~ 6.2 post-seismic moment magnitude, ~~with coincident afterslip and coseismic slip distributions. The measured slip decreased towards the north, suggesting the presence of a slip barrier where stress can accumulate. The coseismic and post-seismic slip distributions coincide with each other, with both peaks adjacent to the centroid.~~ Slip measurements and rupture length ~~estimates from the field and remote sensing datasets~~ characterize the Masbate segment as capable of producing unusually long ruptures with significant offsets despite the presence of creep. Post-seismic interferograms resolved the rupture far better than optical correlation, which was degraded due to high amplitude noise from sensor and environmental sources. Nevertheless, this review of the 2020 M_w 6.6 Masbate earthquake provides a comprehensive slip measurement of the surface rupture and demonstrates the ~~resultant surface rupture morphology, as observed in optical correlation outputs and interferograms, demonstrated the~~ presence of two transtensional basins in the ~~north and south of the province, interlinked by a stepover of the respective Riedel shear zones. This review of the 2020 M_w 6.6 Masbate earthquake reveals~~ Masbate province, revealing new insights into the seismic hazard and seismotectonic setting of ~~Masbate province in the~~ Central Philippines.

~~Graphical Abstract~~

1 Introduction

Philippine earthquake geodesy traditionally relies on interferometry, global navigation satellite system (GNSS), waveform modeling, and field investigation (Velasco et al., 1996; Yen et al., 2018). The techniques have been used to investigate the rupture kinematics of significant earthquakes in the Philippines. Earthquake kinematics describe the three-dimensional motion of the ground surface during an earthquake event, and is characterized by the spatial distribution of ground displacements

25 along the fault rupture. It is crucial in seismic hazard assessments since it can be used in estimating the possible ground shaking intensity and fault rupture potential of a particular location (Morell et al., 2020).

A M_w 6.6 earthquake jolted the island of Masbate on the 18th of August 2020, and was followed by a ~~strong~~ $M5.7$ aftershock in about 10 minutes (Aurelio et al., 2021). In the span of 32 hours after the mainshock, 244 aftershocks were recorded ranging from $M1.6$ to $M5.1$ (PHIVOLCS, 2020). The focal mechanism solution of the mainshock illustrates a strike-slip movement with ~~epicenter~~ hypocenter at 7 km S29°E of Cataingan, and a depth of 21 km (PHIVOLCS, 2020). The earthquake resulted in 30 devastating losses in terms of human lives and properties (NDRRMC, 2020). The heavy impacts of the earthquake warrants a thorough evaluation of the event to support the assessment of seismic hazards to mitigate losses and damages for the next earthquakes in the island of Masbate.

Preliminary assessments using Sentinel-1 ~~descending track interferograms show butterfly~~ interferograms showed butterfly-shaped fringes and phase shifts between 3 to 30 cm along the descending track line-of-sight (LOS) (PHIVOLCS, 2020) and >15 cm 35 LOS ground deformation along the ascending track (Tiongson and Ramirez, 2022). The ascending interferograms used a 12-day (~~15-27~~ 15 to 27 Aug. 2020) temporal baseline, ~~indicating the combination of coseismic and post-seismic slip components alongside the reduced coherence due to the interval~~ resulting in low phase coherence of the InSAR pairs. Spectral analysis of seismic waveforms (Simborio et al., 2022) outlined a 3.0 km $/s$ rupture velocity beginning on a shallow region of the southern fault plane with peak ground displacements ≥ 1.0 m. The InSAR and seismological models underestimate and over- 40 estimate the slip, respectively, compared to the seismic moment predicted slip from global scaling relationships based on moment magnitude (Wells and Coppersmith, 1994).

~~This study provides first-order measurements of the coseismic surface offsets using the novel~~ We introduce the application of optical image correlation (OIC) ~~and the post-seismic deformation using time-series InSAR analysis. The application of OIC in~~ the Philippines ~~is introduced~~ to highlight its importance when used alongside interferometry. The simultaneous application of 45 OIC and InSAR provides complementary information on the decorrelated portions of the coseismic interferogram to improve the understanding of the 2020 M_w 6.6 Masbate earthquake since the event is still relatively unstudied relative to the 2003 ~~event (Besana and Ando, 2005; Lai et al., 2019; Tsutsumi and Perez, 2011, 2013). The acquired slip distributions offer new information on the seismotectonics of Masbate, which is critical to assess consequent seismic hazards. Finally, remote sensing data are coupled with~~ M_w 6.2 event that occurred on the same fault segment (Besana and Ando, 2005; Lai et al., 2019; Tsutsumi and Perez, 2011, 2013).

50 This study presents first-order coseismic offsets and post-seismic slip from the Masbate segment of the Philippine Fault. The resulting slip distributions and surface rupture parameters refine our understanding of the Masbate segment, shed light on the relationship of rupture initiation with mesoscale structures, topography, and bedrock lithology, and provide insights into the underlying faulting mechanisms. The accompanying morphological interpretations of ~~observed surface ruptures to accentuate~~ the stress regime that operates surface ruptures highlight the known stress regime in Masbate, expanding upon ~~the known our~~ understanding of the structural characteristics of the central Philippines. As such, these techniques and findings can help bolster community resilience against earthquake hazards upon practical application and widespread adoption.

2 Geologic Setting and Tectonic Framework

The Philippines is a region of active tectonics and volcanism arising from the interaction of the Sunda Block of the Eurasian Plate, and the Philippine Sea Plate (Aurelio, 2000a; Bird, 2003) (Fig. 1a). Oppositely dipping subduction margins flank the archipelago: the east-dipping Manila, Sulu, Negros, and Cotabato Trenches in the west; and the west-dipping Philippine Trench and East Luzon Trough in the east (Aurelio, 2000a; Barrier et al., 1991). The eastern margin accommodates the northwesterly advance of the Philippine Sea Plate at a rate of 3-9 cm yr⁻¹ relative to the Eurasian Plate (Seno, 1977), while the western end consumes the subducting Sunda Block moving at a rate of ~~10 mm~~ 1 cm yr⁻¹ eastward relative to the Eurasian Plate (Chamot-Rooke and Le Pichon, 1999). The interaction and oblique convergence of the tectonic plates that bind the country resulted in the formation of the approximately 1,200 km long ~~sinistral~~ left lateral Philippine Fault (PF) in the Middle Miocene (Allen, 1962; Fitch, 1972; Tsutsumi and Perez, 2013; Pinet and Stephan, 1990) as the Philippine Sea Plate's motion shifted counterclockwise from northward to a northwestward direction (Aurelio, 2000a). An ~~ASEAN-wide~~ Association of Southeast Asian Nations (ASEAN)-wide GPS network assessment (Aurelio et al., 1997; Aurelio, 1998, 2000b; Rangin et al., 1999) ~~corroborated an extant measured a~~ 2-3 cm yr⁻¹ current slip rate throughout the fault zone. An approximately 580 km ~~northeast~~ convex-curved section of the PF traverses the central Philippines from the southeastern portion of Quezon province to Leyte island (Fig. 1b). Coined the Philippine Fault Bend, the deformation zone in the midsection of the PF is characterized by a N50°W fault strike and forms a releasing bend structure relative to the approximately N20°W general strike north and south of the central section (Lagmay et al., 2005).

2.1 Geology of Masbate

The PF and Sibuyan Sea Fault (SSF) are the prominent structural features in the Masbate island (PHIVOLCS, 2020; Tsutsumi and Perez, 2013) (Fig. 1b). The Masbate segment of the PF extends from Burias and Ticao islands in the north, and strikes N40°W as it crosses the southeastern tip of the mainland (Aurelio et al., 1991). In terms of morphology, it spans approximately 30 km along a linear trough parallel to the coastline (PHIVOLCS, 2020; Tsutsumi and Perez, 2013). Complexities in the fault geometry include ~~an approximately 400 m narrow stepover, delineated a fault complexity~~ between 12°08' N and 12°06' N latitude, and a bifurcation around 12°03' N. The northeastern leg of the bifurcated structure ~~has a northeast convex trace is~~ curved at 12° N and connects to the Leyte segment, whereas the southwestern leg strikes at N20°W and appears to terminate at Cataingan Bay (PHIVOLCS, 2018). Offshore seismic profiling in the bay (Llamas and Marfito, 2022) show left-stepping faults and negative flower structures that denote transtensional deformation in the region.

The SSF is another ~~sinistral~~ left lateral strike-slip fault with a normal component (Aurelio et al., 1991; Bischke et al., 1988). It runs parallel to the PF in mainland Masbate then deflects westward into the Sibuyan Sea (Fig. 1b). The ~~extension~~ extensional component associated with the SSF is represented by tilted blocks formed from normal faulting and bathymetric depressions in the Sibuyan Sea (Bischke et al., 1990). The SSF ~~alters the stress regime in Masbate and is manifested as~~ influences the local stress regime expressed by alternating extensive and compressive structures ~~in seismic profiles between the junction of the SSF and~~, as it interacts with the PF (Aurelio et al., 1997). The SSF was also described as separate from the PF and is a




Figure 1. A) Regional geologic setting of the Philippine archipelago showing the adjacent Philippine Sea Plate (PSP) and Eurasian Plate (EU). The large-scale tectonic features are also depicted, including the east-dipping subduction zones - the Manila Trench (MT), Negros Trench (NT), Cotabato Trench (CT), and Sulu Trench (ST) on the western side; west-dipping trenches - East Luzon Trough (ELT) and Philippine Trench (PT) on the eastern side; and the resultant Philippine Fault running across the archipelago. White arrow indicates the movement direction of the Philippine Sea Plate. B) Major structural features directly affecting Masbate including the central Philippine Fault Zone (Guinayangan, Masbate, and Leyte segments), their approximate offshore projections, the Legaspi Lineament, and the Sibuyan Sea Fault. C) Southeastern Masbate area showing the involved lithologic units and traces of the Philippine Fault and Uson Fault. Black-white bounding box highlights the primary region of interest. Lithology adapted from Manalo et al. (2015). Trace of active structural features adapted from PHIVOLCS. D) Historical seismicity of the Masbate island from 1917 to 2022 showing the moderate earthquakes ($M4\sim6.6$) with focal mechanism solutions of the February 2003 and August 2020 events. Background topography and bathymetry from GEBCO. Earthquake data acquired from USGS-NEIC. Focal mechanism solutions sourced from GlobalCMT.

manifestation the transtensional deformation with the σ_3 oriented perpendicular to the PF (Aurelio, 1992; Aurelio et al., 1991). GPS measurements further substantiate this regime on the local scale (Bacolcol, 2003; Duquesnoy et al., 1994).

Ophiolites comprise the basement complex of Masbate island and are unconformably overlain by a series of sedimentary and igneous units (Fig. 1c). The Cretaceous Panguiranan Chert represents the pelagic cover of the ophiolite sequence (Aurelio and Peña, 2010). The Late Oligocene Nabangig Formation is the oldest lithologic unit underlying the area of interest and is composed of a clastic sequence with recrystallized limestone. The Middle Miocene Lanang Formation lies unconformably over the Nabangig Formation, and consists of highly deformed sedimentary rocks ranging from conglomerates to mudstones (Manalo et al., 2015). Majority of the study area is underlain by the Miocene-Pliocene Buyag Formation. It overlies the Lanang Formation and is composed of conglomerates with fine-grained igneous clasts and interbedded shale and siliceous sediments. The largely coralline Pleistocene Port Barrera Limestone is the youngest stratigraphic unit in the province (Porth et al., 1989).

2.2 Seismotectonics of the Masbate Segment

Bacolcol (2003) and Lai et al. (2019) described the Masbate segment as the transition zone between the northern and southern segments of the central PF. Historical earthquake information revealed recurrence intervals of as short as five years (Besana and Ando, 2005) for moderate magnitude ($M_{5.5-6.2}$) earthquakes to as much as 238 years using slip deficit rates between the 2003 event and interseismic velocities (Lai et al., 2019). A marked absence of strong events beyond $M > 7.0$ in the past four centuries is also apparent (Bautista and Oike, 2000; SEASEE, 1985). A map (Fig. 1d) of moderate seismicity between 1917 and 2022 culled from the United States Geological [Service Survey](#) - National Earthquake Information Center (USGS-NEIC) ([USGS, 2017](#)) highlights the earthquake ~~centers~~-[epicenters](#) related to the Masbate segment and the SSF during this period. Observations and discussions on the 2003 event led Besana and Ando (2005) to characterize the seismic activity of the segment as capable of generating moderate to large events that produce ruptures larger than expected and can be succeeded by post-seismic deformation or creep.

The 2003 $M_s 6.2$ event is the previous notable earthquake in the province that produced an approximately 20 km long rupture with epicenter in the northern portion of the Masbate segment. It was accompanied by either post-seismic deformation or a slow creep component (Besana and Ando, 2005; Lai et al., 2019; Tsutsumi and Perez, 2011, 2013). Kinematic offsets were measured at a maximum of 47 to 50 cm, which is beyond the expected peak offsets for a $M 6.2$ earthquake, based on empirical magnitude scaling equations (Wells and Coppersmith, 1994). Lai et al. (2019) attributed the post-seismic deformation to the significant excess moment released by the earthquake. Aftershock analysis of earthquakes within two months of the 2003 temblor showed a cluster of $M 5.0-6.0$ epicenters at the southern portion of the fault. The aftershocks were accompanied by the development of hairline fractures near Cataingan that expanded to as much as 8 mm (Besana and Ando, 2005). In addition, Bacolcol et al. (2005) recorded 10 cm GPS deviations within six months after the mainshock, supporting observations of post-seismic deformation. Coulomb stress transfer (CST) analysis showed that the 2003 event raised the stress in the southern part of Masbate by approximately 0.3 to 1 bar (Legaspi et al., 2018), which is presumed to ~~cause the~~ [have reduced the time elapsed before the 2020 moderate earthquake along this segment earthquake](#).

The apparent absence of strong earthquakes ($M > 7.0$) in the Masbate segment may be explained by the presence of creep (Scholz, 1998). Aseismic slip during interseismic periods was measured at $2.3 \pm 1.1 \text{ cm yr}^{-1}$ through GPS (Bacolcol et al., 2005) and $0.7\text{-}1.7 \text{ cm yr}^{-1}$ through alignment arrays (Tsutsumi et al., 2016). A moderate background seismicity can be observed along the Masbate segment (Fig. 1d) and is further described in PHIVOLCS reports (1999).

3 Data and Methods

The ~~ground-motion-surface displacement~~ components of the Masbate earthquake were measured using optical and radar satellite data. Pixel offset tracking on PlanetLabs image pairs with 3,125 m resolution was accomplished using the `mm2dpossism` binary of the MicMac program (Rosu et al., 2015) for the horizontal coseismic displacements. Interferometric Synthetic Aperture Radar (InSAR) was done using the European Space Agency's (ESA) Sentinel-1A and 1B radar data with a 6-day temporal baseline for the coseismic slip and a 31-day time series after the mainshock for the post-seismic deformation following the rapid aftershock decay duration (this study). ~~Corresponding slip information was used for the detailed observation and analysis of the ground deformation caused by the 2020 Masbate earthquake event.~~

3.1 Spatiotemporal Seismicity Analysis

Earthquake data from 2019 to 2021 with magnitudes $M \geq 1.0$ were gathered from the PHIVOLCS online earthquake archive (Supplement S3). Foreshock and aftershock patterns were analyzed by plotting the distribution of both magnitude and frequency in time and space. All seismicity information was primarily used to define the earthquake periods (Farrell et al., 2009; Mogi, 1963) related to the 2020 Masbate event. Focal mechanism solutions acquired from PHIVOLCS, USGS, and GlobalCMT were used for the analysis of the nodal planes present in the double-couple as a seismological basis of the sense of movement of the fault plane responsible for the mainshock.

3.2 Optical Image Correlation

PlanetScope satellite imagery (Planet Team, 2017–) was the primary dataset due to the daily data acquisition and high spatial resolution of available products. Images with the least cloud cover covering the region of interest (Fig. 1c) with a maximum 15 % obstruction threshold were selected. Ortho Tile products with a 3.125 m ground resolution, processed at level 3A and corrected for surface reflectance (SR) were utilized. When Ortho Tile products were unavailable, we used Ortho Scene products downsampled to 3.125 m ground resolution and corrected for top-of-atmosphere (TOA) reflectance.

To observe the influence of time on the quality of the output offset rasters, two sets of temporal ranges were tested for the Masbate event (Fig. 2). Five short-term pairs were selected from visually feasible images sampled months apart in 2020, whereas ten *annual* pairs were roughly one year apart (Table 1). This was done to minimize the effect of environmental variables following the approach of Elliott et al. (2020a).

The image pairs were coregistered to a Sentinel-2 basemap using the global coregistration function of the Automated and Robust Open-Source Image Co-Registration Software (AROSICS) (Scheffler et al., 2017) to increase the registration accuracy

Table 1. Satellite geometry and acquisition parameters of the PlanetScope data used in the study. Values of the satellite azimuth (satAzi) are unavailable for the older images in the table. Available tiles for 23 June 2020 are merged to create a single mosaic to maximize the extent. The two datasets for 2 April 2020 are analyzed separately.

Acquisition	satID	satAzi	view \angle	sunAzi	sunElev
2019-11-28T01:53:43	103e	-	1.2	143.9	48.7
2020-04-09T00:33:57	100d	-	0	90.3	42.7
2020-04-17T02:35:36	106e	-	2.1	92.5	73.1
2020-06-06T00:26:15	0f32	-	3	69.9	41.8
2020-06-23T01:29:47	2263	272.7	2	66.4	55.4
2020-06-23T01:54:27	100a	102.6	1	63.2	61.1
2020-07-07T00:19:38	1048	204.5	0.1	70.4	39.2
2020-11-18T02:03:10	105c	98.7	5	146.2	52.6
2020-11-29T01:51:59	1010	98.0	5	143.8	48.8
2020-12-24T02:21:12	1069	261.0	3	149.9	49.4
2021-04-01T02:16:14	241c	99.1	2	106.1	66.2
2021-04-02T02:13:12	1105	168.7	0.1	104.3	65.6
2021-04-02T01:54:40	1013	276.8	5	101.2	60.8
2021-04-11T01:33:11	2276	98.6	4.1	93.2	57.7
2021-04-12T02:15:35	227c	98.4	4.9	96.1	68.1
2021-06-05T01:51:14	1038	98.2	4.9	63.9	61.7
2021-06-10T23:23:03	1049	84.5	3	70	27.5
2021-07-08T02:15:18	2274	12.8	0.2	61.7	65.1
2021-07-11T01:26:43	2434	277.1	5	68.8	54.1

155 while preserving the deformation signal. The pairs were clipped to their overlapping extents with a Universal Transverse Mercator (UTM) projection and either 8- or 16-bit integer pixel data type as required by MicMac. Manual cloud masking was accomplished using the `saisiemasqQT` binary of MicMac for precision masking.

Correlations in the spatial domain were performed with MicMac on the prepared image pairs using the following parameters:

- 7x7 window size ($SzW=3$) for a 21.875 m search grid on each side to compensate for feature-related noise and expected displacement,
 - 0.5 regularization factor ($Reg=0.5$) to provide distance-based correlation weights on the neighboring pixels within the window for noise reduction, and,
 - pixel subsampling parameter of 2 ($SsResolOpt=2$) as recommended by Canizares et al. (2020) for earthquake events to resolve displacement fields close to the ground resolution of the source imagery.
- 160




Figure 2. Two classes of Planet satellite imagery pairs evaluated through optical correlation. A) Short-term pairs taken within the same year as the Masbate earthquake ranging from five to nine months apart and as close to the event as possible. B) Approximately annual span pairs acquired at about 365 days apart where environmental interference is expected to be at a minimum (Elliott et al., 2020a).

165 The resulting output files were georeferenced using `gdal_translate` with the `-a_ullr` flag (GDAL/OGR contributors, 2021) to ensure a 1:1 alignment with the input image pairs without warping the rasters. To retain most deformation signals, the correlation threshold was set to 50 %. A non-local median (NLM) filter with a low H-noise value was applied to prevent flattening real ground deformation signals while improving the signal-to-noise ratio (SNR). The outputs were then translated into the spatial domain by multiplying the rasters with the ground resolution of the input files. ~~Primary offset rasters subjected~~
170 ~~to profiling was~~ The rasters chosen to extract profiles and calculate fault-parallel offsets were selected based on noise levels, atmospheric obstruction, pre-processing accuracy, ROI coverage, and quality of slip visualization ([Supplement S1.1](#)).

The post-processed data generated coherent pixel offset maps along the N-S ($D_{N_{oic}}$) and E-W ($D_{E_{oic}}$) horizontal axes with a 3.125 m pixel resolution. Slip measurements with respect to the general fault strike azimuth ($\theta = 325^\circ$) of the Masbate segment were acquired by transposing the N-S and E-W slip into fault-oriented components using the following equations (Elliott et al.,
175 2020a):

$$D_{\parallel} = D_{N_{oic}} \cdot \cos(\theta) + D_{E_{oic}} \cdot \sin(\theta) \quad (1)$$

$$D_{\perp} = D_{N_{OIC}} \cdot \sin(\theta) - D_{E_{OIC}} \cdot \cos(\theta) \quad (2)$$

180 [Measurements were acquired by taking profiles of the pixel offset data across the fault to determine the net fault-parallel offset \(Supplement S1.2\).](#) The fault-parallel (D_{\parallel}) direction refers to movements parallel to the fault trace represented by laterally offset features on the field. This component was the sole subject of profiling due to the stronger horizontal offset signal in the field. The tension-compression (D_{\perp}) aspect detectable from the fault-normal component was not analyzed due to the minimal deformation along the vertical axis, further complicated by high noise amplitudes.

185 Wide swath profiles (321 x 161 px, roughly ~1.0 x ~0.5 km) along the fault trace were measured with StackProf (Delorme, 2021) developed for the MicMac project. The slip values from profiles aided the measurement of the near-fault slip distribution for comparison with offsets measured in the field. The [offsets-displacements](#) were calculated from the intercepts of the regression of median [displacements/offsets](#), excluding the apparent noise signals and zero correlation pixels to highlight the genuine earthquake slip in the arctangent model. Validated slip data were plotted along the fault strike to show the spatial trends. Absolute standard deviations on the eastern and western fault blocks represent the measurement uncertainties in individual box profiles. Slip uncertainties represented by the standard error of intercepts of the regression lines for the east and 190 western fault blocks were calculated automatically using StackProf. The coseismic displacement curve was calculated using a [robust](#) weighted moving mean approach with minimal smoothing ($q = 0.15$) to prevent the removal of small deviations along the strike.

3.3 Small Baseline Subset (SBAS) Interferometry

Coseismic and post-seismic interferograms were processed with NASA's Alaska Satellite Facility Distributed Active Archive 195 Center (ASF DAAC) Hybrid Pluggable Processing Pipeline (HyP3) (Kennedy et al., 2021). Sentinel-1A/B single look complex (SLC) C-band radar data acquired through the interferometric wide (IW) swath mode was used. Data with 6 to 12-day intervals from paths 61 (descending track) and 69 (ascending track) (Fig. 1d) imaged between 14 to 21 August 2020 for the coseismic slip and 20 August to 20 September 2020 for the afterslip were multilooked by 20 in the azimuth direction and 4 in the range resulting to a pixel size of 80 x 80 m to minimize the effect of land cover (Table 2).

200 Small baseline time series analysis (SBAS) was employed using MintPy (Yunjun et al., 2019) to quantify the post-seismic deformation in the aftershock observation period. Atmospheric corrections were not applied, since the study area is highly localized relative to the global dataset implemented in available databases. Furthermore, the unwrapping reference point was set to a common high coherence pixel for both tracks. The ascending (V_A) and descending (V_D) LOS velocities were projected into the azimuthal fault strike ($\theta = 325^\circ$) (Dianala et al., 2020; Lindsey et al., 2014; Tymofyeyeva and Fialko, 2018) to estimate 205 the fault-parallel (V_{\parallel}) and vertical (V_z) post-seismic velocities. The approach assumes a zero third component acceptable for areas of relatively simple faulting. Equation 3 was used to derive the fault-oriented afterslip LOS velocities within the `asc_desc2horz_vert` script in MintPy. The pixel unit vector directed to the satellite was decomposed into e , n , and u representing the east, north, and upward directions of both tracks, respectively.

Table 2. Summary of reference and secondary [granules scenes](#) processed within Alaska Satellite Facility’s Hyp3 processor to create interferograms.

Track	Reference	Path/Frame ₁	Secondary	Path/Frame ₂	Perp. baseline (m)	Interval (days)	Stage
desc	20200814	61/551	20200820	61/549	-76.4	6	co
asc	20200815	69/32	20200821	69/34	107.1	6	co
desc	20200820	61/549	20200901	61/549	50.4	12	post
asc	20200821	69/34	20200827	69/32	-173.3	6	post
desc	20200826	61/551	20200901	61/549	-76.4	6	post
asc	20200827	69/32	20200902	69/34	126.0	6	post
asc	20200827	69/32	20200908	69/32	42.6	12	post
desc	20200901	61/549	20200907	61/551	49.6	6	post
desc	20200901	61/549	20200913	61/549	68.5	12	post
asc	20200902	69/34	20200908	69/32	-83.1	6	post
asc	20200908	69/32	20200920	69/32	49.9	12	post
desc	20200913	61/549	20200919	61/551	-39.4	6	post

$$\begin{bmatrix} V_A \\ V_D \end{bmatrix} = \begin{bmatrix} e_A \sin(\theta) + n_A \cos(\theta) & u_A \\ e_D \sin(\theta) + n_D \cos(\theta) & u_D \end{bmatrix} \begin{bmatrix} V_{\parallel} \\ V_Z \end{bmatrix} \quad (3)$$

210 The total afterslip was obtained by multiplying the fault-parallel component with the temporal range of the time series and smoothed the data using an NLM filter to arrest pixel outliers. Swath profiles (Krambach, 2015) were taken at 0.5 km intervals with a 3.5 km swath width across the fault strike ~~Slip error (Supplement S4.1)~~. [Slip uncertainty](#) estimates come from the sum of the standard error of regression intercept on both sides of the fault, following the StackProf approach (Delorme, 2021). The spatial distribution of afterslip along the strike was evaluated by applying a low smoothing factor ($q = 0.15$) on the

215 moving mean displacement curve. The vertical post-seismic displacement was not investigated due to the primarily horizontal ground movement. The coseismic interferogram was qualitatively compared with the optical correlation data due to the lack of resolvable slip in the near-fault region due to decorrelation [due to higher magnitudes of coseismic slip](#).

3.4 Ground Truthing

Field investigation was conducted from 18 to 21 May 2022 to identify and measure the horizontal offsets ~~of across~~ the ground

220 rupture. Locations of known ground rupture (dela Cruz, per. comm., 2022) were complemented by points of interest identified from the image correlation outputs and InSAR. ~~Significant contrast on the horizontal offset rasterson~~ [Due to the two-year interval between the earthquake and fieldwork, geomorphic or human modifications of offset features may have occurred.](#) [Additionally, many offsets and evidence of surface rupture may have been erased given the small magnitude of surface slip.](#)

225 The significance of these changes can be assessed by correlating field data with other measurement techniques to determine the slip.

We selected points of interest for field checking based on the pixel offset rasters, prioritizing locations with significant contrast along the approximate and certain fault traces~~was the primary criterion to select the points of interest for field checking.~~ Field offsets were measured with a steel tape and structural orientations were determined with a base plate compass. Measurements of on-fault offsets were limited to individual rupture strands. Due to the scarcity of remnant ruptures, most sites
230 had a single measurement. Uncertainty was estimated by comparing values at these locations to earlier measurements acquired closer to the earthquake (PHIVOLCS, 2020; Miraballes et al., 2020). In areas with complex, closely spaced ruptures, multiple measurements were acquired and statistically assessed to estimate the uncertainty.

Observation points with measurable feature offset and ruptures were plotted along the strike to show the attitude and slip distribution along the fault (Supplement S2). The measured horizontal offsets were used to validate the results of the remote
235 sensing analysis. Finally, the surface rupture length, and maximum and average displacements were derived from the conjoined remote sensing outputs and field data to estimate the seismic moment.

4 Results

4.1 Seismicity

4.1.1 Earthquake Frequency and Temporal Relationships

240 The frequency-time distribution ~~plot of hypocenters of earthquakes~~ between 25 August 2019 to 18 August 2021 shows the immediate seismicity related to the 18 August 2020 M_w 6.6 event (Fig. 3). The foreshocks, mainshock, and aftershocks are identifiable in the probability density subplot (Fig. 3a) as a distinct bell-shaped curve relative to the generally quiescent background activity. The ~~typical daily earthquake activity of Masbate island is~~ island of Masbate typically experiences less than ten
~~events earthquakes~~ per day, corresponding to a moderate background seismicity~~with multiple days without recorded earthquake~~
245 ~~activity. The increasing.~~ An increase in earthquake frequency defines the foreshock sequence ~~in the probability density subplot (Farrell et al., 2009) that began (Farrell et al., 2009) at around~~ 32 days before the mainshock, as shown by the distinct trend increase in the probability density subplot. The peaks of the frequency plots (Figs. 3a and b) correspond to 141 earthquakes which were mostly aftershocks on 18 August 2020. A negative slope in the probability density subplot depicts the aftershock sequence. The decaying earthquake frequency is initially symmetrical to the foreshock trend with an inflection point around 30
250 days after the mainshock, which ~~then~~ transitioned to a more gradual decrease. The aftershocks persisted for 166 days or up to 31 January 2021. Succeeding return to background levels ~~is characterized by low seismic activity and days without earthquakes are closer to each other relative to the~~ saw a similar seismic activity to the previous background activity from 2019 to early 2020.




Figure 3. A) Earthquake probability density subplot showing the normalized population of earthquakes per day, highlighting the contrast between the periods of the earthquake sequence. B) Plot showing the number of earthquakes with magnitudes $M \geq 1$ related to the Masbate segment and their temporal distribution relative to the 18 August 2020 mainshock.

4.1.2 Spatiotemporal Earthquake Distribution

255 In general, the background seismicity ($M \leq 4$) is well distributed throughout the overall length of the Masbate segment of the PF traversing the islands of Ticao, Burias, and the provincial mainland (Fig. 4a). However, an absence of seismic activity is noticeable in the northern portion of the fault in mainland Masbate during this period. On the contrary, an earthquake cluster occurred near the location of the mainshock. Foreshock activity is concentrated in southeastern Masbate adjacent to the mainshock and is composed of mostly weak ($M 2 \sim 3$) and shallow hypocenters peaking at $M 4.9$ (Fig. 4b).

260 Three ~~(3)~~ moment tensor solutions from GlobalCMT, PHIVOLCS, and USGS highlight the mainshock's dominant left-lateral strike-slip ~~motion-movement~~ (Fig. 4c; ~~Table ??~~). The solutions of PHIVOLCS and USGS show a noticeable, albeit minimal, dip-slip component towards the NNE. The epicentral location of the mainshock from GlobalCMT and PHIVOLCS coincided with the fault trace, while USGS' focal mechanism solution deviated to the northeast of the fault.

~~Fault-mechanism parameters from PHIVOLCS, USGS, and GlobalCMT for the $M_w 6.6$ Masbate earthquake mainshock on 18 August 2020.~~

Source	Time	Long	Lat	Depth (UTC)	(E)	(N)	(km)	Str	Dip	Rake	Str	Dip	Rake
PHIVOLCS	00:03:00	124.00	12.00	5	23564	17414285	26	USGS	00:03:48	124.13	12.03	10	23186167322774
GlobalCMT	00:03:53	123.98	12.04	12	23389177323871								

Aftershocks that transpired on the same day as the mainshock (Fig. 4d) are shallow (1-30 km) and densely grouped in the southeastern offshore extension of the segment and deepen (10-50 km) as they taper toward the onshore portion. The aftershocks have weak magnitudes ranging from $M 2.0$ to $M 4.4$. Succeeding first-month aftershocks are weak to moderate,

270




Figure 4. Spatiotemporal earthquake distribution for the Masbate segment. A) Background seismicity between 25 Aug 2019 to 10 Jul 2020. B) Foreshocks from 19 Jul to 17 Aug 2020. C) Moment tensors of the mainshock from GlobalCMT, PHIVOLCS, and USGS. D) Same-day aftershocks. E) First-month aftershocks from 19 Aug to 17 Sept 2020. F) Second-month aftershocks from 18 Sept to 17 Oct 2020. G) Third-month aftershocks from 19 Oct to 16 Nov 2020. H) Fourth-month aftershocks from 18 Nov to 14 Dec 2020. I) Last aftershocks from 22 Dec 2020 to 9 Feb 2021. J) Return to background seismicity covering 11 Feb to 18 Aug 2021.

with a wide range of magnitudes between $M1.7$ to $M5.1$, mostly occurring at depths above 30 km. A monthly progressive dispersal of hypocenters in space is also observed, with the first month showing a dense cluster in the southeastern section that tapers to the northwest (Fig. 4e). The pattern and geographic extent of the first-month aftershocks appear to be an expansion of the same-day aftershocks. The succession of the aftershock patterns in the following months continued to scatter along the Masbate segment and the SSF while retaining a perceptible concentration in the offshore southeastern extension of the causative fault segment (Figs. 4f, 4g, 4h, and 4i). The last two months of aftershock activity (Figs. 4h and 4i) feature the least number of hypocenters throughout the aftershock sequence.

Background levels returned on 2 February 2021 defined by the well-distributed hypocenters on the fault segments with moderate magnitudes (Fig. 4j) similar to the pre-event seismicity in 2019 (Fig. 4a). The frequency and distribution of earthquakes in time and space enabled the distinction of various stages of the earthquake sequence relative to the 2020 Masbate event (Table ??).

~~Duration and date range of earthquake sequences of the 2020 Masbate event derived from the probability density plot of the frequency-time distribution. The aftershock sequence is subdivided into two following the inflection point separating the sharp and gradual decay of activity.~~
Sequence Start date End date Duration
foreshocks 17 Jul 2020 17 Aug 2020 32 days
mainshock 18 Aug 2020 18 Aug 2020
aftershocks (rapid decrease) 18 Aug 2020 17 Sept 2020 30 days
aftershocks (gradual decay) 17 Sept 2020 31 Jan 2021 136 days

4.2 Optical Image Correlation

4.2.1 Long and Short Temporal Baseline

The acquired ~~optical correlation displacement maps~~ pixel offset maps (Fig. 5) illustrate the fault rupture and the known ~~sinistral left lateral~~ movement of the Masbate segment. ~~However, high noise values tend to obscure the rupture trace, despite the presence of noise that obscures portions of the rupture.~~ Only one of the eight image pairs from the short timespan (Figs. 2a) produced ~~limited regions of outputs with~~ perceptible rupture due to the low SNR. ~~The resolved slip is comparatively clearer in the annual timespan pairs despite the persistent low SNR.~~ Annual timespan pairs resolved the rupture more clearly compared the short timespan image pairs. The north-south slip components show better representations of the rupture, whereas the east-west components tend to be noisier. Nevertheless, the valid image pairs ~~with resolvable displacements (Supplement S1.1) with~~ resolvable slip and subjected to stack profiling exhibit reliable estimates of the slip in the near-field region.

4.2.2 Coseismic ~~Offsets~~ Offset Estimates

~~Supplemented by the known trace of the PF from PHIVOLCS, a~~ A generalized fault trace (yellow line; Fig. 5) was delineated from the valid ~~optical correlation displacement rasters and served as a reference~~ pixel offset rasters and the known fault trace, which served as the baseline for slip profiling. The fault trace is mostly straight, except for an apparent northeasterly strike rotation towards the north, where it follows the piedmont of a topographic high in Dimasalang. No measurements were acquired in the area due to anomalously high values, which are probably terrain residuals. Topographic artifacts and decreasing




Figure 5. Optical correlation results showing the representative data from the 9 April 2020 and 2 April 2021 image pair. Blue lines in panels E and G are the absolute median fault-parallel values across the profile box. Gray shaded region in panels E and G feature the 1σ absolute standard deviation of the fault-parallel values across the fault in each box. A) N-S component. Blue refers to southward movement, while yellow refers to northward motion. B) E-W component. Blue refers to westward movement, while yellow refers to eastward motion. C) Fault-parallel component. Pink refers to movements toward the NW while blue denotes motion toward the SE. D) Closer view of box 32 showing the fault-parallel offset. E) Swath profile of box 32 highlighting the 43 cm slip. Lower regression intercept of the western block relative to the eastern block indicates left lateral movement. F) Closer view of box 47 displaying the fault-parallel offset in Cataingan. G) Swath profile of box 47 exhibiting the peak 59.2 cm left lateral offset in this particular image pair.




Figure 6. Along strike distribution of the fault-parallel offsets from 126 profile boxes across six image pairs. Error bars refer to the 1σ sum of standard errors of the intercepts of regressions of the east and west blocks providing slip error estimates.

displacements slip away from the epicenter posed difficulties in delineating the northern rupture section near Naro Bay. The In the southwestern section, the offset rasters show a new splay striking NW-SE that bisects the bifurcating fault trace in Cataingan, identified by PHIVOLCS (2020) and Tsutsumi and Perez (2013) in their active faults map previously recognized bifurcation in Cataingan. The rupture traverses the deltaic area and possibly continues southward across Cataingan Bay, then resurfaces on the opposite side of the bay. Smaller fractures and minute rupture components are not visible due to ubiquitous noise.

Sinistral Left lateral offsets were quantified along the ~25 km onshore length of the fault from Dimasalang to Cataingan, given the inland spatial extents and continuity where optical correlation was possible. Slip measurements were not derived in the area feasible. Edge artifacts and the obscured western block in the south of the bay due to the presence of a water body, which obscured the western block and promoted edge artifacts hindered the acquisition of slip profiles.

Fifty-one (51) wide swath profile boxes were generated throughout the fault rupture, with the first box in Dimasalang and the last in Cataingan to determine the along-strike slip distribution (Fig. 6). The optical correlation results generated a total of 126 left-lateral measurements over six image pairs (Supplement S1.2). Using a 325° general strike azimuth of the Masbate segment, the east-west and north-south displacement components (Figs. 5a and b) were transposed into the fault-parallel direction (Fig. 5c).

The 9 April 2020 - 2 April 2021 pair was selected as the representative dataset due to its wide spatial extent and reliable representation of the fault rupture. The north-south component (Fig. 5a) shows a continuous linear rupture trace from Dimasalang to the deltaic region area in Cataingan, although the Dimasalang area and its border with Palanas exhibited unusually high values due to noise topographic artifacts. The east-west component (Fig. 5b) appears is less coherent, likely due to smaller ground movements in this direction or overprinting information caused by low SNR overprinted data from the noise. The outputs show noise from both artificial and environmental sources, represented by topographic artifacts in the southwestern block, an ap-

parent seamline, and reduced coherence due to differences in land cover between the images. Nevertheless, the fault-parallel component (Fig. 5c) show good estimates of the ~~sinistral-left lateral~~ slip, further supported by the near-field offset models.

325 Box 32 (Fig. 5d), located ~6 km north of the GlobalCMT surface projection, shows a 43 cm ~~displacement-slip~~ in the near-field (Fig. 5e). Pixel offset values increase in the far-field for the western block (~~negative x-axis~~), which may ~~be due to noise~~ either represent noise or distributed deformation. Box 47, situated ~1.5 km south of the GlobalCMT epicenter, exhibits a 59.2 cm ~~sinistral-left lateral~~ offset. The box represents the peak offset in this image pair and displays a clear contrast between the east and western blocks ~~despite the noise~~.

330 The plot of the slip distribution along strike (Fig. 6) displays the 126 ~~sinistral-left lateral~~ offsets compiled from the valid offset rasters to characterize the displacement field. The slip minima ~~are measured to range ranges~~ from 6 to 9 cm, located 4.5 km from the northern shoreline in Dimasalang. The displacement increased to ~50 cm around the vicinity of the GlobalCMT projection of the centroid. Slip values remained constant within a ~5 km span in Cataingan, ~~followed by an increase and~~ increases to a peak ~60 cm ~~sinistral slip. The anomaly is followed by a recession southward to the~~ at around ~1 km southeast
335 of the centroid. South of the peak value, the displacement declines toward the shoreline. The skewed semi-elliptical weighted moving mean curve (solid gray line; Fig. 6) ~~yields a mean sinistral displacement~~ shows the average left lateral offset of 37.7 ± 10 cm ~~with a maximum of~~, while the individual measurements indicate a peak 60.6 ± 9.8 cm ~~around the GlobalCMT offset~~ adjacent to the surface projection of the centroid in Cataingan.

4.3 Interferometry

340 4.3.1 Coseismic Interferograms

The Sentinel-1 coseismic wrapped interferograms in the ascending and descending tracks (Fig. 7a and c) show the ~~butterfly fringe pattern and oppositely directed LOS displacement vergence~~ butterfly-shaped fringe patterns and oppositely verging LOS displacements, which is typical of strike-slip events. The fringe lobe patterns illustrate a range increase from the ascending track and a range decrease from the descending track for the western side of the fault. Since the orbit of the ascending track is
345 subparallel to the regional fault strike, the ~~sinistral-left lateral~~ movement is indicated with a possible subsidence component. ~~This is~~ This is consistent with the moment tensor solutions (~~Table ?? Fig. 4c~~), and agrees with the sense of movement ~~observed from~~ shown by the optical correlation outputs and field observations.

Low phase coherence persists in the near-fault region. Widespread decorrelation leads to mostly unusable information upon phase unwrapping due to the likelihood of creating unwrapping errors. Noise is similarly dominant and overprints most of the
350 deformation signals in an approximately 18 km-wide region across the fault trace, hence, the lack of near-fault information in preliminary published interferograms (PHIVOLCS, 2020; Tiongson and Ramirez, 2022). In addition, the rupture is unresolved upon phase unwrapping (Fig. 7b and d) due to decorrelation. This loss of resolvable information in the near-field ultimately hinders slip measurements in the coseismic interferograms.




Figure 7. Sentinel-1 coseismic interferograms in ascending and descending tracks. A) Wrapped ascending interferogram. B) Unwrapped LOS displacement in the ascending track. C) Wrapped descending track interferogram. D) Unwrapped LOS displacement in the descending track.




Figure 8. Post-seismic deformation between 20 August to 20 September 2020. A) Ascending track time series showing range increase (yellow) and decrease (blue) for the eastern and western blocks, respectively. B) Descending track time series showing range decrease (blue) and increase (yellow) for the eastern and western blocks, respectively. C) Projected fault-parallel post-seismic deformation field. D) Fault-parallel afterslip for box 4 in Dimasalang showing the 5.4 cm left lateral afterslip. Lower regression intercept of the western block relative to the eastern block indicates left lateral movement. Median afterslip value is shown by the blue line. Gray background exhibits the 1σ uncertainty across the box. E) Swath profile of box 13 indicating an 8.4 cm left lateral afterslip around the Dimasalang-Palanas boundary. F) Swath profile of box 25 showing a 6.9 cm afterslip in Palanas. G) Swath profile of box 39 in Cataingan showing a 12.0 cm left lateral afterslip.

4.3.2 Post-seismic Deformation Estimates

355 Post-seismic interferograms capture an improved overall manifestation of the fault rupture compared to the coseismic ones (Fig. 8). This is a consequence of the expected smaller afterslip values relative to the coseismic slip, which decreases the number of fringes and increases the coherence. The temporal coverage of the interferogram stack ~~subsequently provides a means to quantify the~~ allows us to quantify afterslip to complement the ~~coseismic slip derived estimated coseismic slip~~ from optical correlation. However, the first post-earthquake Sentinel-1 revisit occurred two days after the mainshock, resulting in all
360 measurements to be lower bound estimates of the afterslip.

Unwrapping errors are less frequent on the ascending and descending track interferograms and are further minimized upon time-series stacking. The remaining phase unwrapping errors upon stacking and filtering were manually excluded in profiling. These unwrapping errors occur as significant high amplitude phase shifts and polarity reversals, noticeable in the Dimasalang and Palanas areas.

365 The fault-parallel afterslip deformation field (Fig. 8c) shows a ~~strong~~ strongly linear and simple fault trace that accommodated the post-seismic deformation. Deviations from the regional fault strike are observed around 5-13 km from Naro Bay in the north, represented by multiple short NNE-SSW oriented ($\sim 290\text{-}300^\circ$ azimuth) bends which are adjacent to and coincide with the narrow stepover in the Palanas-Dimasalang area. However, the noticeable unwrapping errors and low ground resolution (80 m) precludes the validity and recognition of these rupture complexities.

370 The afterslip is ~~measured~~ resolved from Naro Bay in the north towards Cataingan Bay in the south along the ~ 25 km simplified fault trace. ~~Forty-nine (Supplement S4.1). Forty-nine~~ (49) swath profile boxes were generated along this trace. Individual ~~afterslip~~ measurements reveal centimetric fault-parallel afterslip with minimal across-fault variability and minor 1σ uncertainties, except where the boxes overlap with the phase unwrapping error patches. The deformation was detected up to the far-field extents of the profiles from most swath boxes.

375 Boxes 4, 13, 25, and 39 (Figs. 8d, e, f, and g) illustrate select wide swath profiles roughly equidistant from each other. Box 4, situated in Dimasalang, shows a 5.4 cm left lateral fault-parallel afterslip and a reasonably low variability across the fault. Unwrapping errors in the far-field at 1.5 km from the fault on both sides caused irregularities in the profile. Box 13, located around the boundary of Palanas and Dimasalang, depicts an 8.4 cm fault-parallel slip. Both sides of the fault exhibit exceedingly even slip across the fault, except for a noticeable dip in the eastern edge. The dip is likely due to data loss
380 since the box overlapped with the water body to the east. The profile also appears to be offset by about ~ 100 m to the west, which corresponds to one of the abovementioned bends that may be an unwrapping artifact. Still, the measured slip from the arctangent model remains valid. Box 25 shows a 6.9 cm afterslip in Palanas, located at the southern edge of the ~~stepover~~ fault complexity. Multiple across-fault deviations are visible around 200 and 500 m east of the fault and in the western far-field. The eastern anomalies could either be unwrapping or topography related, while the western patch comes from unwrapping issues.
385 Box 39 in Cataingan displays a 12.0 cm ~~sinistral~~ left lateral afterslip with good symmetry and even across-fault distribution.

The along-strike afterslip distribution graph (Fig. 9) depicts the computed fault-parallel offsets from north to south on the mainland. The smallest value is measured in Dimasalang, adjacent to Naro Bay, amounting to 4.3 cm of ~~sinistral~~ left lateral




Figure 9. Along strike distribution of the fault-parallel afterslip formed by the 49 profile boxes. Error bars refer to the 1σ standard errors of the intercepts of regressions of the east and west blocks. Solid blue line shows the weighted moving mean afterslip curve.

afterslip. The [ground](#) offsets exhibit a generally increasing trend to around ~5 km from the north. Fluctuating values are observed between 6.1 to 13.8 km from the north in Dimasalang and Palanas. The measured displacements oscillate between 5.0 to 9.6 cm in this span. The increasing trend continued to a 14.3 ± 0.9 cm peak afterslip in Cataingan, located 24 km from Naro Bay and ~3 km southwest of the GlobalCMT surface projection. The displacements decrease from the [maximum largest](#) value to 11.5 cm on the shoreline of Cataingan Bay. The weighted moving mean curve (solid blue line) indicates a mean [sinistral-left lateral](#) afterslip of 8.8 ± 0.7 cm.

4.4 Ground Truthing

4.4.1 Observed Surface Ruptures

Remnant earthquake effects were observed on laterally offset cultural and geomorphic features from Dimasalang to Cataingan following the known fault trace ([Supplement S2](#)). Assisted by the remote sensing data, the observed ground manifestation of the earthquake from fifty-seven (57) stations (Fig. 10a and b) include fractures, reported earthquake-induced landslides (EIL), indications of liquefaction features and lateral spreading, secondary gravitational features, and ruptures with quantifiable displacements. Earthquake effects recorded during the field activity include reported ground shaking from interviews with locals.

The observed rupture zone of the 2020 Masbate earthquake persisted from Dimasalang to Cataingan and crossed Cataingan Bay. Measured surface ruptures strike [both-subparallel-and](#) at a low angle with respect to the regional fault trace (Fig. 10c) with a mean strike azimuth of 313° . The [northern-terminus-of-the-surface-rupture-northernmost-resolvable-ground-offset](#) was observed on a road in Dimasalang, around ~3 km from Naro Bay, [with-an-An](#) apparent 0.5 cm offset fracture oriented parallel to the fault trace [was recorded in this point, possibly representing the minimum northward constraint of the field rupture](#). A 2-5




Figure 10. Compiled ground-truthing data. A) Earthquake hazards aside from ruptures. B) Ruptures symbolized by size showing the amount of slip. C) Rupture orientations. D) Ruptured tree showing cumulative 87 cm slip from 2003 and 2020 events with estimated 27 ± 10 cm offset from the 2020 earthquake. E) Soil mound cut by the fault highlighting a 48 ± 3 cm left lateral offset. F) Rice field embankment traversed by fault showing 60 ± 15 cm displacement. G) Ruptured fishpond dike recording a 34 ± 5 cm offset. H) Abandoned septic tank showing a 30 cm displacement south of Cataingan Bay.

cm displacement oriented 325° NW occurring on a residential structure on top of the fault was observed on the border between Dimasalang and Palanas. The rupture displacement increased further ~~down~~ south ~7 km from the northern shoreline, where a 7 cm left lateral offset was seen coupled with spreading of the road pavement. The fault bisected a coconut tree at the southern
410 limit of the ~~stepever~~ fault complexity in Palanas (Fig. 10d) (Aurelio, 2022), resulting in a cumulative 87 cm opening from the 2003 and 2020 events. The newer slip is estimated at 27 ± 10 cm on a 275° W oriented rupture. Right-stepping en echelon fractures occurred adjacent to the coconut tree, most of which were eroded at the time of visit.

The rupture ~~continued southwards to Cataingan, is continuous southward, up to~~ 4 km from the GlobalCMT centroid projection. ~~Fault~~ The fault rupture traversed an abandoned house and deflected ~~a wall for its wall by~~ roughly ~50 cm. A soil mound
415 at the base of the wall (Fig. 10e), ~~located at the base of the wall,~~ shows a 297° W-oriented rupture with a 48 ± 3 cm left lateral offset. Residents reported a ~200 m long rupture in the area, which was already mostly eroded. Nevertheless, the remnant right-stepping en echelon fractures formed a linear rupture zone connected to the displaced wall. ~~The surface rupture began to bisect the bifurcating fault trace, expressed as a~~ A 30 cm ~~rupture across~~ offset oriented 279° E on a road in Cataingan : ~~The 279E-oriented rupture continued south~~ likely represents the northern indication of the rupture bisecting the bifurcation in
420 Cataingan. The rupture continued to an agricultural area (Fig. 10f), where the embankments are displaced by ~~~60~~ 60 ± 15 cm, representing the peak field offset measurements. Long, continuous ruptures were observed ~~in~~ on the rice fields immediately after the event but were erased by the time of the field investigation. Another southerly continuation was identified, where ~~the~~ a 341° NW-oriented rupture displaced fishpond levees (Fig. 10g) by about 34 ± 5 cm. Locals stated that the visible rupture appeared lengthy and continuous in the deltaic areas immediately after the earthquake. The rupture was evident again south of
425 Cataingan Bay, manifested on a collinear set of points through several residential structures. A septic tank diagonally cut by a 279° NW-oriented rupture, shows a 30 cm left lateral displacement (Fig. 10h). Offset measurements adjacent to the septic tank vary from 5 to 30 cm.

4.4.2 Left-lateral Offset Distribution

The plot of offset features along the fault strike (Fig. 11) shows the minimum 0.5 cm left lateral displacement ~3 km from
430 Naro Bay in Dimasalang. This point corresponds to the ~~northern terminus of the rupture~~ northernmost resolvable ground offset. The measurements continually increased to 50 cm beginning at 17 km from Naro Bay, around the boundary of Palanas and Cataingan. The offset measurements remained constant for a ~4 km span followed by an abrupt dip to 30 cm, which increases again to the peak 60 cm offset. The local 30 cm minimum was measured on a ruptured road. This single discordant measurement could be due to off-fault displacement, physical reduction due to post-earthquake repairs, or an inaccurate measurement from
435 a cultural feature. The displacement tapered to 30 cm near the shoreline of Cataingan Bay. Spatially coincident offsets ranging from 5 to 30 cm resulted in a ~~peculiar trend~~ wide range of uncertainties at the southeastern limitation of the plot. The weighted moving mean curve of the slip distribution (~~solid green~~ broken red line) for the 2020 Masbate event ~~suggests a 24.5 cm mean~~ sinistral displacement with a indicates a 25.3 cm mean left lateral displacement, while the individual observations denote the maximum of 60 cm displacement near the GlobalCMT centroid projection.




Figure 11. Along strike distribution of measured field offsets.

440 5 Discussion

5.1 Comparison of Optical Correlation and InSAR

5.1.1 Displacement Threshold

Independently applying optical correlation and interferometry to analyze the coseismic and post-seismic periods of the 2020 Masbate event provided a basis to assess the technical capabilities and shortcomings of the methods in a tropical setting. Peak coseismic displacements of the Masbate earthquake amount to ~60 cm, denoting a high interferometric phase gradient for Sentinel-1. SAR interferograms lose coherence when offsets exceed half the interferometric fringe per pixel (Michel et al., 1999), highlighting the effectiveness of the method for small deformations in mm and cm scales (Tronin, 2006). The underestimated slip estimates of PHIVOLCS (2020) and Tiongson and Ramirez (2022) are likely due to a loss of information because of the high phase gradients and near-field decorrelation on the coseismic interferograms (Fig. 7).

450 The ability of InSAR to detect centimetric displacements was evident when comparing the post-seismic and coseismic interferograms. The rupture trace is conspicuous and the displacement offset is observable in both the near-field and far-field across the fault. Sinistral afterslip was measured from of the post-seismic interferograms. Left lateral afterslip measurements are at least 4 to 14 cm with low an average 1σ uncertainties uncertainty of 1.5 cm. Millimeter-scale offsets were not detected, owing to the resultant 80 m pixel resolution, which is too large to discern fine rupture information. The underestimated coseismic slip of PHIVOLCS (2020) and Tiongson and Ramirez (2022) can be attributed to this limitation, where information is lost due to high phase gradients and near-field decorrelation (Fig. 7).

Optical correlation is useful to assess the near-field region (Avouac et al., 2006) and larger displacements equivalent to at least one-tenth of the pixel resolution (Elliott et al., 2020b; Leprince et al., 2007). Results show that near-field displacements are resolved (Figs. 5 and 6) despite noticeable high noise levels. OIC, using PlanetScope data with 3.125 m pixel resolution,

460 was able to detect ~~sinistral coseismic displacements~~ left lateral offsets between 7 to 60 cm in the near-field, but with ~~high-a~~
larger 19.9 cm 1σ ~~uncertainties~~ average uncertainty than InSAR due to noise. ~~However~~ Thus, careful interpretation of swath
profiles is crucial to identify valid measurements because noise easily overprints tectonic signals. Noise signals are amplified
by misaligned acquisition geometry, unequal solar illumination angle, and insufficient geometric corrections (Elliott et al.,
2020b; Stumpf et al., 2017). The representative 9 April 2020 - 2 April 2021 (satellite ID 1105) images had decent illumination
465 and acquisition geometry agreement (Table 1) and support the validity of the measurements.

5.1.2 Temporal Baseline

The pair classes of PlanetScope orthoimages used in the study showed that images roughly one year apart resolved the rupture
far better than those imaged months apart primarily due to the contrasting vegetative cover. ~~Cyclical changes in land cover~~
~~(Elliott et al., 2020a) necessitate the use of satellite images taken during the same season of the year because similar features~~
470 ~~assists~~ Satellite images from the same seasons are critical for optical correlation because feature similarity helps the algorithm
match features ~~for better correlation~~ more accurately (Elliott et al., 2020a). Feature matching in agricultural and developing
areas can be difficult in the subpixel domain given the difference in crop patterns and rapid anthropogenic development,
leading to differing pixel intensity values in the pre-event and post-event images (Barnhart et al., 2011).

Images that are one day apart generate deformation fields with minimal seasonal and anthropogenic artifacts. However, this
475 may have exceptions such that correlations for the 2016 Fukushima earthquake using one-day interval drone images resulted in
noisy deformation fields with low correlation (Valkaniotis, per. comm., 2021). Single-day intervals were not tested in Masbate
due to the lack of usable orthoimagery in terms of study area coverage, relatively similar solar illumination, and cloud cover.

DInSAR relies on coherence to resolve the deformation along the LOS direction, which is controlled by the similarity of point
scatterers between the acquisitions. Good coherence in urban areas and dry deserts that remain constant throughout several
480 years produce quality interferograms (Wang and Fialko, 2014; Wei and Sandwell, 2010), which indicate the capability of
interferometry to function for longer periods. However, in other types of surface cover such as vegetation and snow, coherence
~~exhibits an indirect~~ has an inverse relationship with temporal baseline, which reduces the usable time interval (Kervyn, 2001).
The six-day revisit of Sentinel-1 conveniently dictates the pre-event and post-event radar data pairing in areas affected by sub-
optimal land cover. The baseline can be increased for L-band satellites, such as ALOS-2, for they can detect the bare ground
485 irrespective of surface cover. Interferograms with severe unwrapping errors are manually excluded in the time-series analysis
(Yunjun et al., 2019), effectively limiting the temporal baseline range to six to 12 days.

5.2 Accuracy of Optical Correlation

The ~~obtained coseismic slip~~ temporal range of the pixel offset rasters implies a potential inclusion of interseismic and post-seismic
slip in the optical correlation measurements. We preferentially co-interpreted the stacked optical correlation results with the
490 field data to describe the rupture kinematics and displacements, which may be inferred to represent the surface coseismic slip.

The slip distribution from optical correlation agrees with the offset feature measurements from the field survey (Fig. 12).
The weighted moving mean of the coseismic offsets from optical correlation unifies the measurements from the six valid pairs




Figure 12. Combined afterslip and coseismic slip distributions. Solid gray line shows the weighted moving mean of the six valid pairs. Dashed red line represents the weighted moving mean of the field measurements. Blue line and markers represent the afterslip placed in a secondary axis.

to compensate for data gaps that particular pairs are unable to resolve. The variance between the two coseismic slip measures ~~for Masbate~~ is typically at ~ 2 cm with a maximum ~ 17 cm. The larger difference is interpreted to be a local variation or measurement error since the specific point was taken on a repaired road section, which the remote sensing approach may have missed as it generally detects the overall deformation and is less sensitive to local slip.

Previous studies show varying degrees of misfit between field offsets and remote sensing data (Cheloni et al., 2014; Gold et al., 2021). These were attributed to diffuse displacement, secondary faulting, and shallow fault complexities. Our measured coseismic displacements, however, show consistency and ~~lack of minimal~~ short wavelength variability. This ~~is may be~~ a consequence of the two-year interval between the earthquake event and the field survey resulting in the eradication of most surface traces and smoothing of the field offset distribution. Furthermore, ruptures on unconsolidated overburden ~~is are~~ easily erased due to active weathering and erosion, leaving only the most prominent traces to remain. Similarly, abundant vegetation, challenging surface conditions, and relatively small tectonic displacements compromise the clarity of the resolved rupture and efficacy of optical correlation. Nevertheless, OIC outputs ~~co-validate the field offset measurements and~~ indicate that the remote sensing results ~~are not merely arbitrary results~~ accurately capture ground displacements and provide greater along-strike coverage.

The extent of detectable rupture is another fundamental difference between optical correlation outputs and field measurements. The northern rupture terminations are situated 4.6 km and 3.2 km from Naro Bay for optical correlation and field survey, respectively. To the south, OIC measurements cease at the shore of Cataingan Bay since no slip measurements are attainable on the opposite side of the water body, where the field survey revealed more ruptures as the fault resurfaced therein. ~~However, both methods are unable to acquire information in the~~ Neither method can resolve slip on submerged fault segments, requiring the use of complementary marine survey techniques and datasets that cover the ocean floor.

5.3 Surface Rupture Kinematics

5.3.1 Coseismic Offsets

515 The amount of coseismic slip is a key factor in the assessment of seismic moment, rupture mechanism, and degree of seismic hazard. The results of this study show a unimodal curve with a characteristic long wavelength and isolated short wavelength variability for the regional slip distribution (Fig. 12). The pattern suggests a low generalized resolution due to temporally-influenced smoothing and noise. The long wavelength offset trend imply overall maturity (Allam et al., 2019), which is a credible interpretation for the Masbate segment of the PF given its Middle Miocene initiation [and rate of activity](#) (Fitch, 1972; 520 Tsutsumi and Perez, 2013; Pinet and Stephan, 1990). Furthermore, the known fault trace in Masbate is morphologically linear, supporting the general maturity estimate (Manighetti et al., 2021).

The peak of the unimodal distribution lies ~2 km southeast of the centroid, which implies a relatively simple fault zone (Gold et al., 2021) where the slip is concentrated in the shallow crust. This interpretation, however, requires validation through geodetic inversion. Such peaks usually define the location of the ruptured asperity (Kaneda et al., 2008) where the highest 525 stress drop occurs due to peak moment release from the accumulated potential energy (Freymueller et al., 1994).

~~By inferring a variable slip signature in the submerged segment coupled with the observed offset variations as the fault resurfaces to the southeast, a mostly vertical rupture propagation with a southwesterly component is suggested (Xu et al., 2010) ; in accordance with source spectral fitting (Simborio et al., 2022) and indicates a relatively complex shallow submarine fault structure (Llamas and Marfito, 2022)~~
530 [A variable slip distribution can be inferred on the submerged sections of the Masbate segment of the PF due to the lack of submarine offset measurements. This slip trend may portray a complex shallow fault structure in this section as the slip is distributed on multiple fault segments \(Avouac et al., 2006; Allam et al., 2019; Wang et al., 2021\) . Moreover, recent marine geophysical surveys in the vicinity of Cataingan Bay \(Llamas and Marfito, 2022\) revealed the geometric complexity of the fault, which supports our interpretation.](#)

Offset values gradually decay to zero towards the northwest. The smooth slip trend, devoid of short wavelength varia- 535 tion, ~~may suggest~~ [suggests](#) a systematic rupture (Treiman, 2002) on a relatively simple sub-vertical fault plane (Allam et al., 2019; Chen et al., 2021). Tapering and deepening aftershock pattern along with the focal mechanism solutions could support this observation. ~~The decreasing slip trend marks the presence of a slip barrier in the north (Xu et al., 2010) due to velocity-strengthening behavior or complex fault geometry (Biasi and Wesnousky, 2017; Bischke et al., 1990) where strain accumulates. This aligns with the known stress accumulation at rupture terminations of most faults (Stein, 2003).~~

540 ~~Topography and bedrock geology may also control the rupture (Kaneda et al., 2008), such that the southwestern rupture termination in the study is on an alluvial plain, succeeded by rugged topography northwards. The narrow stepover in Palanas and the bifurcation point in Cataingan coincide with the stratigraphic contacts of the Late Oligocene Nabangig and Miocene-Pliocene Buyag Formations. The fault complexities may reflect the rheological contrast between the elastic limestones of the Nabangig Formation and conglomeratic Buyag Formation.~~

545 5.3.2 Post-Seismic Offsets

Post-seismic deformation is usually accommodated as aseismic afterslip, viscoelastic relaxation, or poroelastic rebound (He et al., 2021; Tomita et al., 2020). The coseismic displacement distribution suggests that post-seismic deformation is stress-driven (He et al., 2021) and is mechanically operated as aseismic afterslip (Johanson, 2006) to accommodate the excess coseismic stress. Our interferometry results revealed a 14 cm peak-largest afterslip, which accounts for at least 23 % of the peak coseismic displacement over the rapid aftershock decay interval. This lower limit estimate is a consequence of the six-day revisit cycle of Sentinel-1, however, lead to incomplete measurements resulting in incomplete data since the first radar data image in the time-series stack was from 20 August 2020 and was not able to capture does not include the earliest post-seismic deformation immediately after the 18 August 2020 mainshock.

The distribution of afterslip (Fig. 12) along the fault demonstrates a general slip increase southwards, which can be divided into two distinct sections. The northern ~15 km section is characterized by short-wavelength variations and average sinistral-left lateral offsets of 5-6 cm, whereas the southern section includes the peak afterslip without distinctive variability. The northern trend may indicate the occurrence of diffuse afterslip on a wide zone, which may or may not have breached the surface. Chen et al. (2021) suggested an increased relative friction such that the released post-seismic moment is typically less compared to the mainshock, resulting in heterogeneous loading. However, we cannot exclude the possibility that variability may stem from surface complexities or data processing artifacts, as indicated by the observed unwrapping errors and the lack of data from the earliest post-seismic period.

Post-seismic deformation usually occurs outside the regions of peak coseismic slip in normal and thrust faults as stress transfer as the shear stress is transferred by the mainshock cause and causes stress loading at the locations of incomplete rupture and unruptured sections (Cheloni et al., 2014; Johanson, 2006; Ozawa et al., 2011; Yagi et al., 2001). However, the (Johanson, 2006; Bürgli et al., 2019) Our results reflect the coincidence of the post-seismic and coseismic slip distributions on the surface whose maxima are aligned and equidistant from the centroid accompanied by a gradual slip recession northwards (Fig. 12). This is similar to the 1999 Izmit (Reilinger et al., 2000), 1999-1996 Biak (Das and Henry, 2003), and 2011 Maduo (Wang et al., 2019) earthquakes, which also exhibit spatially overlapping coseismic and post-seismic slip distributions.

The findings primarily suggest We interpret this overlap of the slip distributions to indicate a vertical stress migration along the fault surface to either deeper or shallower sections of the fault. However, we

Aseismic afterslip, viscoelastic relaxation, and poroelastic rebound are the common mechanisms of post-seismic deformation (He et al., 2021; Tomita et al., 2020). The existence of resolvable afterslip is a clear indication of post-seismic activity as the stress is released by the mainshock. Our observed coincidence of the coseismic and post-seismic slip distributions probably indicates a localized stress transfer to the adjacent velocity strengthening vicinity of the primary asperity in accordance with the rate-and-state friction model (Marone et al., 1991). This would suggest that the post-seismic deformation is likely stress-driven (He et al., 2021) and operated as aseismic afterslip (Johanson, 2006) to accommodate the excess coseismic stress. This also agrees with the hypothesized vertical migration of stress along the fault plane.

We recognize that the data is limited to surface measurements and more information is required to accurately describe the occurrence of the afterslip along the fault plane. Determining the cause-mechanics of the superposed slip character of the 2020 Masbate earthquake event is of interest, and warrants investigation into potential factors such as non-steady state friction

(Helmstetter and Shaw, 2009), residual stress heterogeneity (Wang et al., 2019), stress reorganization, or shallow material redistribution (He et al., 2021). Additionally, the contribution of known creep associated with the Masbate segment to the slip distributions merits further investigation.

5.3.3 Surface Rupture Length

585 Surface rupture length estimates from remote sensing data were assumed to be symmetric about the peak displacement zone (Table 3). The coseismic displacement distributions from optical correlation and field survey (Fig. 12) implies that the northern rupture terminus is located approximately 3.2 km from the northern shoreline with the peak slip values occurring between 17.3 to 23.5 km, resulting to rupture length estimates ranging from 28.2 to 41 km. The absence of offset measurements in the submerged portions of the fault necessitated the use of interpolation. Given the asymmetrical nature of strike-slip
590 profiles distributions, which can take on either elliptical or triangular forms (Perrin et al., 2016), caution is critical when interpreting the interpolated values. The rupture lengths are also measured from the total length of the field survey and its extension towards the stepover in Cataingan Bay (Llamas and Marfito, 2022) resulting in 25 and ~~35.59~~35.6 km rupture lengths.

5.3.4 Seismic Moment Estimate

Moment magnitude (M_w) and seismic moment (M_o) estimates for the 2020 Masbate event were calculated from the surface
595 rupture parameters (Table 3). M_w calculations were conducted using the corresponding empirical equations from regression of rupture parameters and moment magnitudes (Wells and Coppersmith, 1994). The M_o was derived as a function of M_w from scaling relationships (Hanks and Kanamori, 1979). This approach was preferred over directly calculating the M_o from the rupture area and material rigidity due to the lack of reliable estimates of rupture width and subsurface fault rupture parameters.

The M_w estimates from the maximum and average displacements range between M_w ~~6.50~~6.5 to M_w ~~6.66~~6.66, ~~and are similar to~~
600 6.7, and closely fit the instrumental 6.6 moment magnitude. All rupture length estimates returned high M_w values between M_w ~~6.73~~6.7 to M_w ~~6.97~~7.0 which agrees with the 2003 earthquake observations, wherein the fault rupture is longer than expected relative to the magnitude. ~~Comparing the~~However, a downward counterfactual approach (Woo and Mignan, 2018; Woo, 2019)
given the fault uncertainties at depth, suggests the probability of the larger M_w estimates. Nevertheless, comparing the individually calculated M_w values show that the moment magnitude estimates from maximum displacement provide the closest fit
605 with instrumentally determined moment. The ~~accepted M_w 6.64 from the maximum displacement is equivalent to 6.6 estimate~~
from the inferred maximum displacement parameter is consistent with a seismic moment of ~~1.15×10^{19}~~ 1.2×10^{19} N·m.

The ~~moment released during lower limit of the~~ post-seismic ~~deformation moment~~ was assessed using the ~~maximum largest~~
and average afterslip values (Table 3). The ~~maximum largest~~ afterslip, which measured 0.14 m, ~~corresponded to a moment~~
~~magnitude of 6.15~~corresponds to M_w 6.2, while the average afterslip of 0.09 m ~~yielded a moment magnitude of~~ is consistent
610 with M_w 6.1. Given that the source parameter of the ~~accepted preferred~~ coseismic moment magnitude is the maximum displacement, the M_w estimate from the ~~maximum afterslip is preferred largest afterslip is similarly selected~~
among the two afterslip parameters. The ~~maximum afterslip is equivalent to largest afterslip is consistent with~~ a M_o of ~~2.12×10^{18}~~ 2.1×10^{18} N·m, indicating that the afterslip resulted in the release of energy equal to at least 18 % of the mainshock.

Table 3. Measured rupture parameters with corresponding moment magnitude (Wells and Coppersmith, 1994) and seismic moment (Hanks and Kanamori, 1979).

Parameter	Value	M_w	Mo (N·m)
Surface rupture length	34.40 <u>34.4</u> km ¹	6.88 <u>6.9</u>	2.64×10^{19} <u>2.6×10^{19}</u>
	28.20 <u>28.2</u> km ²	6.78 <u>6.8</u>	1.89×10^{19} <u>1.9×10^{19}</u>
	41.00 <u>41.0</u> km ³	6.97 <u>7.0</u>	3.54×10^{19} <u>3.5×10^{19}</u>
	39.50 <u>39.5</u> km ⁴	6.95 <u>6.9</u>	3.33×10^{19} <u>3.3×10^{19}</u>
	25.00 <u>25.0</u> km ⁵	6.73 <u>6.7</u>	1.54×10^{19} <u>1.5×10^{19}</u>
	35.59 <u>35.6</u> km ⁶	6.90 <u>6.9</u>	2.79×10^{19} <u>2.8×10^{19}</u>
Maximum displacement	0.61 <u>0.6</u> m ⁷	6.64 <u>6.6</u>	1.15×10^{19} <u>1.2×10^{19}</u>
	0.60 <u>0.6</u> m ⁸	6.64 <u>6.6</u>	1.14×10^{19} <u>1.1×10^{19}</u>
Average displacement	0.38 <u>0.4</u> m ⁹	6.66 <u>6.7</u>	1.24×10^{19} <u>1.2×10^{19}</u>
	0.25 <u>0.3</u> m ¹⁰	6.50 <u>6.5</u>	6.99×10^{18} <u>7.3×10^{18}</u>
Maximum <u>Largest</u> afterslip	0.14 m	6.15 <u>6.2</u>	2.12×10^{18} <u>2.1×10^{18}</u>
Average afterslip	0.09 m	6.10 <u>6.1</u>	1.78×10^{18} <u>1.8×10^{18}</u>

¹ symmetrical about the midpoint of the zone of peak displacement

² symmetrical about the lower limit of the zone of peak displacement

³ symmetrical about the upper limit of the zone of peak displacement

⁴ symmetrical about the peak displacement

⁵ cumulative rupture length from the field survey

⁶ cumulative rupture length from the field survey extended to the onshore stepover identified by Llamas and Marfito (2022)

⁷ from optical correlation

⁸ from field survey

⁹ from the weighted moving mean curve of optical correlation outputs

¹⁰ from the weighted moving mean curve of field measurements

5.4 Seismotectonic Implications

615 5.4.1 Surface Rupture Morphology

The fault geomorphology (Fig. 10c) and geodetic observations (Figs. 5 and 8) highlight the linearity of the Masbate segment. However, the ~~narrow stepover~~ fault complexity (Tsutsumi and Perez, 2013) around 12°07'N was not clearly defined due to unwrapping errors in interferometry and high-amplitude noise in optical correlations.

620 The ruptures in Palanas form a right-stepping Riedel shear zone, with the ruptures occurring at a low angle with respect to the regional fault strike. Furthermore, a possible new splay was observed to the south, which potentially represents the primary fault plane, cutting across the bifurcating trace and traversing Cataingan Bay. Antithetic shears were observed at the bifurcation point and along the cross-cutting splay.




Figure 13. Morphology of the transtensional basins linked by right-stepping shear zone. Solid line indicates the regional fault. Lines with hachures indicate the basin sidewall boundary.

625 Topography and bedrock geology may also control the rupture (Kaneda et al., 2008), such that the southwestern rupture termination is observed on an alluvial plain, succeeded by rugged topography northwards. The fault complexity in Palanas and the bifurcation point in Cataingan coincide with the stratigraphic contacts of the Late Oligocene Nabangig and Miocene-Pliocene Buyag Formations. The previously identified fault complexities in this area may reflect the rheological contrast between the clastic limestones of the Nabangig Formation and conglomeratic Buyag Formation.

5.4.2 Developing Transtensional Basins

630 The morphology of mainland Masbate following the 2020 event (Fig. 13) was the basis of assigning a late relative development stage in a modelled transtensional basin development sequence (Wu et al., 2009). The basin boundary is characterized by floodplains surrounded by topographic highs. Apparent asymmetric depocenters were ~~also~~ recognized, with the shallower depocenter occurring on the floodplain and the nearshore section of Cataingan Bay, while the second one extends further southwest.

635 A separate transtensional basin is discernible in Dimasalang given the topography, rupture occurrence, and extensional structures in north Masbate (Bischke et al., 1990). The northern and southern basins appear to be linked ~~along the narrow stepover by the Riedel shear zone~~ in Palanas. Fault complexities in the principal deformation zone of a pull-apart basin (Wu et al., 2009) impede rupture propagation (Biasi and Wesnousky, 2017) in line with the implications of the observed decreasing coseismic and post-seismic slip towards the north. However, the scarcity of observed ruptures inhibits further analysis of the northern transtensional basin.

640 Fitting the development stage of the southern basin is constrained by the absence of information regarding local strain partitioning. Nonetheless, the expressed fault maturity and identified negative flower structures (Bischke et al., 1990; Llamas and Marfito, 2022) in the northern and southern bays highlight the presence of oblique extensional stress bounding the Masbate mainland. The orientation of the regional fault implies a WNW-ESE oriented σ_1 . However, real-world conditions are more complex compared to the simplified parameters in ~~the analog model (Wu et al., 2009)~~ analog models.

645 6 Conclusions

The necessity of coupling optical correlation and interferometry is demonstrated by investigating the surface rupture of the 2020 M_w 6.6 Masbate earthquake along the Masbate segment of the Philippine Fault. Seismicity data showed that foreshocks began 32 days prior to the mainshock on 18 August 2020. The first 30 days of subsequent aftershocks are characterized by a rapid decrease in earthquake frequency, followed by a transition to a gradual decay until 31 January 2021.

650 Optical correlation and InSAR are directly compared through the corresponding deformation rasters in the coseismic period. Measuring the ~~displacements~~ ground offset from the near-field region of the interferogram is hindered by decorrelation, whereas the far-field suffers because of the submerged northeastern block. The optical correlation method revealed an average ~~displacement of~~ 37.7 ± 10 cm ~~and a maximum displacement of~~ displacement and 60.6 ± 9.8 cm ~~sinistral offsets~~ peak left lateral offset in the near-field of the onshore fault segments. ~~Both methods are not capable of measuring offshore displacements. The peak sinistral~~ The preferred peak left lateral coseismic offset is ~~equivalent to~~ consistent with a geodetic moment magnitude of ~~M_w 6.64 (1.15×10^{19})~~ 6.6 (1.2×10^{19} N·m).

The capacity of InSAR to detect smaller displacement amplitudes is highlighted by ~~assessing~~ the post-seismic ~~deformation~~ interferograms. The SBAS time-series stack outlined ~~a~~ at least 14.3 ± 0.9 cm ~~peak largest~~ afterslip and 8.8 ± 0.7 cm average ~~sinistral~~ left lateral afterslip. The measurements correspond to a ~~M_w 6.15 (2.12×10^{18})~~ 6.2 (2.1×10^{18} N·m) energy release equivalent to at least 660 18 % of the coseismic moment estimate. The resolved rupture from interferometry is clearer relative to optical correlation. Multimodal rupture length estimates translate to overestimated moment magnitude values. Hence, the Masbate segment is characterized as capable to produce significant slip and ruptures that are longer than expected, despite the short interval and continuous aseismic stress release during interseismic periods.

~~In terms of accuracy, the~~ Our optical correlation data agrees with the laterally offset features from field investigation. ~~We interpreted,~~ highlighting the capacity of the method to accurately capture ground displacements despite the difficult tropical terrain. We view this to be a consequence of the time interval between the 2020 Masbate earthquake and the field investigation. Weathering and erosion of fine ruptures ~~was expected~~ had transpired, leading to a smoother field offset distribution.

The coseismic slip distribution shows the presence of a single asperity adjacent to the GlobalCMT centroid. ~~A possible slip barrier exists in the north that may become a strain accumulation zone due to velocity-strengthening properties and shallow fault complexities. Furthermore~~ In conjunction, the surface distribution of the afterslip coincides with the coseismic slip distribution. 670 Based on the stress transfer theorem and the rate-and-state friction model, we infer that the ~~afterslip involved post-seismic~~

deformation is fundamentally stress-driven and migrated vertically in either the downdip or updip direction along the sub-vertical fault plane.

675 The surface rupture of the 2020 Masbate event cut across the extant bifurcating trace of the Philippine Fault in the island, possibly reflecting the primary fault plane. Comparing the rupture morphology with analog models reveal the presence of two transtensional basins in Cataingan and Dimasalang, located in the south and north, respectively. The low-angle orientation of ruptures in the midsection correspond to a Riedel shear zone, ~~whereas the stepover~~ which is interpreted to link the ~~two~~ identified transtensional basins.

680 The findings and the methods we used for this study in Masbate leads us and future scientists to further analyze the geometry, kinematics, mechanics, and dynamics of fault movement in the area for improved seismic hazard mitigation. Specifically, the comprehensive slip data and observed unusually long rupture lengths along the fault measured through the optical correlation method agree with previous characterizations by Besana and Ando (2005), indicating significant movement despite the presence of creep.

685 Usage of optical correlation alongside interferometry provides high-resolution surface displacement measurements that can aid in the assessment of earthquake hazards to assist the development of mitigation strategies. We recommend the widespread ~~use~~ adoption of these methods in studying other active faults in the Philippines. In addition, future studies could incorporate other datasets such as UAV data and explore the method for other surface processes. By leveraging the capabilities of these technologies, overall community resilience against earthquake hazards can be improved and ultimately help mitigate the loss of lives and infrastructure damage caused by earthquakes.

690 *Code availability.* MicMac is available from github.com/micmacIGN/micmac. StackProf is available from github.com/micmacIGN/stackprof. MintPy is available from github.com/insarlab/MintPy.

Data availability. Sentinel-2 datasets are accessed from ESA/EC Copernicus Sentinels Scientific Data Hub and various repositories. Dataset in this study are retrieved from peps.cnes.fr. Sentinel-1A/B radar data and cloud interferogram processing is available at search.asf.alaska.edu. Planet Labs data are not openly accessible, but academic/scientific access can be requested. Outputs are rendered with Scientific Colour Maps (Crameri, 2021) available from fabiocrameri.ch/colourmaps.
695

Author contributions. **Khelly Shan C. Sta. Rita:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Validation, Visualization, Writing - original draft, Writing - review & editing. **Sotiris N. Valkaniotis:** Formal analysis, Resources, Software, Supervision, Methodology, Writing - review & editing. **Alfredo Mahar Francisco A. Lagmay:** Conceptualization, Funding acquisition, Resources, Supervision, Writing - review & editing.

700 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. ~~TEXT~~ We extend our wholehearted appreciation to Dr. Arturo Daag for his suggestions and comments. We would like to thank the National Mapping and Resource Information Authority for providing the elevation data. We would also like to thank the Philippine Institute of Volcanology and Seismology as well as the Disaster Risk Management office of Dimasalang, Masbate for their contributions to the field investigation. We are also grateful for the comments of Dr. Austin Elliot, Dr. Gordon Woo, and the anonymous reviewer for their input and contributions towards improving this work.

705

References

- Allam, A. A., Kroll, K. A., Milliner, C. W. D., and Richards-Dinger, K. B.: Effects of fault roughness on coseismic slip and earthquake locations, *Journal of Geophysical Research: Solid Earth*, 124, 11 336–11 349, <https://doi.org/10.1029/2018JB016216>, 2019.
- Allen, C. R.: Circum-Pacific faulting in the Philippines-Taiwan Region, *Journal of Geophysical Research (1896-1977)*, 67, 4795–4812, <https://doi.org/https://doi.org/10.1029/JZ067i012p04795>, 1962.
- Aurelio, M.: Tectonique du segment central de la faille philippine : etude structurale, cinématique et evolution géodynamique, Ph.D. thesis, <http://www.theses.fr/1992PA066699>, thèse de doctorat dirigée par Rangin, Claude Terre, océan, espace Paris 6 1992, 1992.
- Aurelio, M.: Displacement rates and block rotation in and around the Philippines-results from GEODYSSSEA data Part II, Program and Abstracts : GEOSEA 98-Ninth Regional Congress on Geology Mineral and Energy Resources of Southeast Asia, 238, <https://cir.nii.ac.jp/crid/1570572700614632960>, 1998.
- Aurelio, M.: Tectonics of the Philippines revisited, *Journal of the Geological Society of the Philippines*, 55, 119–183, 2000a.
- Aurelio, M.: Shear partitioning in the Philippines: Constraints from Philippine Fault and global positioning system data, *Island Arc*, 9, 584 – 597, <https://doi.org/10.1111/j.1440-1738.2000.00304.x>, 2000b.
- Aurelio, M.: Masbate's three-legged coconut tree spurs scientific interest, <https://coverstory.ph/masbates-three-legged-coconut-tree-spurs-scientific-interest/>, 2022.
- Aurelio, M. and Peña, R.: *Geology of the Philippines*, 2nd Edition (2010), Mines and Geosciences Bureau, 2010.
- Aurelio, M., Barrier, E., Rangin, C., and Müller, C.: The Philippine Fault in the Late Cenozoic tectonic evolution of the Bondoc-Masbate-N. Leyte area, Central Philippines, *Journal of Southeast Asian Earth Sciences*, 6, 221–238, [https://doi.org/10.1016/0743-9547\(91\)90069-a](https://doi.org/10.1016/0743-9547(91)90069-a), 1991.
- Aurelio, M., Barrier, E., Gaulon, R., and Rangin, C.: Deformation and stress states along the central segment of the Philippine Fault: implications to wrench fault tectonics, *Journal of Asian Earth Sciences*, 15, 107–119, [https://doi.org/10.1016/s0743-9547\(97\)00001-9](https://doi.org/10.1016/s0743-9547(97)00001-9), 1997.
- Aurelio, M., Lagmay, M., Escudero, J. A., and Catugas, S.: Latest Philippine earthquake reveals tectonic complexity, *Temblor*, <https://doi.org/10.32858/temblor.191>, 2021.
- Avouac, J.-P., Ayoub, F., Leprince, S., Konca, O., and Helmberger, D. V.: The 2005, Mw 7.6 Kashmir earthquake: Sub-pixel correlation of ASTER images and seismic waveforms analysis, *Earth and Planetary Science Letters*, 249, 514–528, <https://doi.org/10.1016/j.epsl.2006.06.025>, 2006.
- Bacolcol, T. C.: Etude géodésique de la faille Philippine dans les Visayas, Ph.D. thesis, Université Pierre et Marie Curie, <http://www.theses.fr/2003PA066404>, thèse de doctorat dirigée par Barrier, Éric Géologie Paris 6 2003, 2003.
- Bacolcol, T. C., Aguilar, A., Jorgio, R., and de la Cruz, R.: Interseismic Deformation and Horizontal Displacements Associated with the February 15, 2003 Masbate Earthquake, Tech. rep., Philippine Institute of Volcanology and Seismology, Philippines, 2005.
- Barnhart, W. D., Willis, M. J., Lohman, R. B., and Melkonian, A. K.: Insar and optical constraints on fault slip during the 2010-2011 new zealand earthquake sequence, *Seismological Research Letters*, 82, 815–823, <https://doi.org/10.1785/gssrl.82.6.815>, 2011.
- Barrier, E., Huchon, P., and Aurelio, M.: Philippine fault: A key for Philippine kinematics, *Geology*, 19, 32, [https://doi.org/10.1130/0091-7613\(1991\)019<#60;0032:pfakfp>#62;2.3.co;2](https://doi.org/10.1130/0091-7613(1991)019<#60;0032:pfakfp>#62;2.3.co;2), 1991.
- Bautista, M. L. P. and Oike, K.: Estimation of the magnitudes and epicenters of Philippine historical earthquakes, *Tectonophysics*, 317, 137–169, [https://doi.org/10.1016/s0040-1951\(99\)00272-3](https://doi.org/10.1016/s0040-1951(99)00272-3), 2000.

- Besana, G. M. and Ando, M.: The central Philippine Fault Zone: Location of great earthquakes, slow events, and creep activity, *Earth Planets Space*, 57, 987–994, <https://doi.org/10.1186/BF03351877>, 2005.
- 745 Biasi, G. and Wesnousky, S.: Bends and Ends of Surface Ruptures, *Bulletin of the Seismological Society of America*, 107, 2543–2560, <https://doi.org/10.1785/0120160292>, 2017.
- Bird, P.: An updated digital model of plate boundaries, *Geochemistry, Geophysics, Geosystems*, 4, <https://doi.org/https://doi.org/10.1029/2001GC000252>, 2003.
- Bischke, R., Suppe, J., and Del Pilar, R.: Implications of a newly discovered branch of the Philippine fault system, in: *International Symposium on the Geodynamic Evolution of Eastern Eurasian Margin*, 1988.
- 750 Bischke, R., Suppe, J., and del Pilar, R.: A new branch of the Philippine fault system as observed from aeromagnetic and seismic data, *Tectonophysics*, 183, 243–264, [https://doi.org/https://doi.org/10.1016/0040-1951\(90\)90419-9](https://doi.org/https://doi.org/10.1016/0040-1951(90)90419-9), geodynamic evolution of the Eastern Eurasian Margin, 1990.
- Bürgmann, R., Fielding, E., Liu, Z., and Wang, K.: Postseismic deformation due to the 2019 Ridgecrest earthquakes and its impact on the CGM, Tech. rep., https://files.scec.org/s3fs-public/reports/2020/20016_report.pdf?CUhbHuJtXfV9IPyOGtLr.ZvyIuNq91Fg, 2021.
- 755 Canizares, C., Motagh, M., and Haghshenas Haghghi, M.: Resolving 3D coseismic deformation of the 2019 Mw 7.1 Ridgecrest earthquake using radar and optical data, in: *EGU General Assembly Conference Abstracts*, p. 13659, 2020.
- Chamot-Rooke, N. and Le Pichon, X.: GPS determined eastward Sundaland motion with respect to Eurasia confirmed by earthquakes slip vectors at Sunda and Philippine trenches, *Earth and Planetary Science Letters*, 173, 439–455, [https://doi.org/https://doi.org/10.1016/S0012-821X\(99\)00239-3](https://doi.org/https://doi.org/10.1016/S0012-821X(99)00239-3), 1999.
- 760 Cheloni, D., Giuliani, R., D’Anastasio, E., Atzori, S., Walters, R., Bonci, L., D’Agostino, N., Mattone, M., Calcaterra, S., Gambino, P., Deninno, F., Maseroli, R., and Stefanelli, G.: Coseismic and post-seismic slip of the 2009 L’Aquila (Central Italy) MW 6.3 earthquake and implications for seismic potential along the Campotosto fault from joint inversion of high-precision levelling, InSAR and GPS data, *Tectonophysics*, 622, 168–185, <https://doi.org/10.1016/j.tecto.2014.03.009>, 2014.
- 765 Chen, H., Qu, C., Zhao, D., Ma, C., and Shan, X.: Rupture kinematics and coseismic slip model of the 2021 mw 7.3 Maduo (China) earthquake: implications for the seismic hazard of the Kunlun fault, *Remote Sensing*, 13, 3327, <https://doi.org/10.3390/rs13163327>, 2021.
- Cramer, F.: Scientific colour maps, <https://doi.org/10.5281/zenodo.5501399>, 2021.
- Das, S. and Henry, C.: Spatial relation between main earthquake slip and its aftershock distribution: AFTERSHOCK DISTRIBUTIONS, *Reviews of Geophysics*, 41, <https://doi.org/10.1029/2002RG000119>, 2003.
- 770 Delorme, A.: StackProf, <https://github.com/micmacIGN/stackprof>, 2021.
- Dianala, J. D. B., Jolivet, R., Thomas, M. Y., Fukushima, Y., Parsons, B., and Walker, R.: The Relationship Between Seismic and Aseismic Slip on the Philippine Fault on Leyte Island: Bayesian Modeling of Fault Slip and Geothermal Subsidence, *Journal of Geophysical Research: Solid Earth*, 125, e2020JB020052, <https://doi.org/https://doi.org/10.1029/2020JB020052>, e2020JB020052 10.1029/2020JB020052, 2020.
- 775 Duquesnoy, T., Barrier, E., Kasser, M., Aurelio, M., Gaulon, R., Punongbayan, R. S., and Rangin, C.: Detection of creep along the Philippine Fault: First results of geodetic measurements on Leyte Island, central Philippine, *Geophysical Research Letters*, 21, 975–978, <https://doi.org/https://doi.org/10.1029/94GL00640>, 1994.
- Elliott, A., Elliott, J., Hollingsworth, J., Kulikova, G., Parsons, B., and Walker, R.: Satellite imaging of the 2015 M7.2 earthquake in the Central Pamir, Tajikistan, elucidates a sequence of shallow strike-slip ruptures of the Sarez-Karakul fault, *Geophysical Journal International*, 221, 1696–1718, <https://doi.org/10.1093/gji/ggaa090>, 2020a.
- 780

- Elliott, J., De Michele, M., and Gupta, H. K.: Earth observation for crustal tectonics and earthquake hazards, *Surveys in Geophysics*, 41, 1355–1389, <https://doi.org/10.1007/s10712-020-09608-2>, 2020b.
- Farrell, J., Husen, S., and Smith, R. B.: Earthquake swarm and b-value characterization of the Yellowstone volcano-tectonic system, *Journal of Volcanology and Geothermal Research*, 188, 260–276, <https://doi.org/https://doi.org/10.1016/j.jvolgeores.2009.08.008>, the Track of the
785 Yellowstone Hotspot, 2009.
- Fitch, T. J.: Plate convergence, transcurrent faults, and internal deformation adjacent to Southeast Asia and the western Pacific, *Journal of Geophysical Research*, 77, 4432–4460, <https://doi.org/10.1029/jb077i023p04432>, 1972.
- Frey Mueller, J., King, N. E., and Segall, P.: The co-seismic slip distribution of the Landers earthquake, *Bulletin of the Seismological Society of America*, 84, 646–659, <https://doi.org/10.1785/BSSA0840030646>, 1994.
- 790 GDAL/OGR contributors: GDAL/OGR Geospatial Data Abstraction software Library, <https://gdal.org>, 2021.
- Gold, R. D., DuRoss, C. B., and Barnhart, W. D.: Coseismic surface displacement in the 2019 ridgecrest earthquakes: comparison of field measurements and optical image correlation results, *Geochemistry, Geophysics, Geosystems*, 22, <https://doi.org/10.1029/2020GC009326>, 2021.
- Hanks, T. C. and Kanamori, H.: A moment magnitude scale, *Journal of Geophysical Research: Solid Earth*, 84, 2348–2350,
795 <https://doi.org/https://doi.org/10.1029/JB084iB05p02348>, 1979.
- He, L., Feng, G., Wu, X., Lu, H., Xu, W., Wang, Y., Liu, J., Hu, J., and Li, Z.: Coseismic and early postseismic slip models of the 2021 mw 7.4 maduo earthquake (Western china) estimated by space-based geodetic data, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2021GL095860>, 2021.
- Helmstetter, A. and Shaw, B. E.: Afterslip and aftershocks in the rate-and-state friction law, *Journal of Geophysical Research: Solid Earth*,
800 114, 2007JB005077, <https://doi.org/10.1029/2007JB005077>, 2009.
- Johanson, I. A.: Coseismic and postseismic slip of the 2004 parkfield earthquake from space-geodetic data, *Bulletin of the Seismological Society of America*, 96, S269–S282, <https://doi.org/10.1785/0120050818>, 2006.
- Kaneda, H., Nakata, T., Tsutsumi, H., Kondo, H., Sugito, N., Awata, Y., Akhtar, S. S., Majid, A., Khattak, W., Awan, A. A., Yeats, R. S., Hussain, A., Ashraf, M., Wesnousky, S. G., and Kausar, A. B.: Surface rupture of the 2005 kashmir, pakistan, earthquake and its active
805 tectonic implications, *Bulletin of the Seismological Society of America*, 98, 521–557, <https://doi.org/10.1785/0120070073>, 2008.
- Kennedy, J. H., Hogenson, K., Johnston, A., Kristenson, H., Lewandowski, A., Logan, T. A., Meyer, F. J., and Rine, J.: Get Hyp3! SAR processing for everyone, Tech. Rep. EGU21-8973, Copernicus Meetings, <https://doi.org/10.5194/egusphere-egu21-8973>, 2021.
- Kervyn, F.: Modelling topography with SAR interferometry: illustrations of a favourable and less favourable environment, *Computers & Geosciences*, 27, 1039–1050, [https://doi.org/10.1016/S0098-3004\(00\)00158-8](https://doi.org/10.1016/S0098-3004(00)00158-8), 2001.
- 810 Krambach, M.: swath profile, <https://plugins.qgis.org/plugins/swathProfile/>, 2015.
- Lagmay, A. M., Tengonciang, A., and Uy, H.: Structural setting of the Bicol Basin and kinematic analysis of fractures on Mayon Volcano, Philippines, *Journal of Volcanology and Geothermal Research*, 144, 23–36, <https://doi.org/https://doi.org/10.1016/j.jvolgeores.2004.11.015>, the Tectonics and Physics of Volcanoes, 2005.
- Lai, L. C., Rau, R. J., Bacolcol, T., Ker, C. M., Sapla, G., Ching, K. E., and Solidum, Jr., R.: Earthquake Potential of the Creeping-to-locked
815 Masbate Segment of the Philippine Fault Based on GPS Observations, in: AGU Fall Meeting Abstracts, vol. 2019, pp. G43B–0748, 2019.
- Legaspi, C. J., Ortega, J. P., Reyes, P. R., and Riza, E. G.: Investigation of Stress Changes along the Philippine Fault Zone Using Coulomb Stress Transfer Modeling, in: GEOCON 2018, 2018.

- Leprince, S., Ayoub, F., Klingler, Y., and Avouac, J.-P.: Co-Registration of Optically Sensed Images and Correlation (COSI-Corr): an operational methodology for ground deformation measurements, in: 2007 IEEE International Geoscience and Remote Sensing Symposium, p. 1943–1946, <https://doi.org/10.1109/IGARSS.2007.4423207>, 2007.
- Lindsey, E. O., Fialko, Y., Bock, Y., Sandwell, D. T., and Bilham, R.: Localized and distributed creep along the southern San Andreas Fault, *Journal of Geophysical Research: Solid Earth*, 119, 7909–7922, <https://doi.org/https://doi.org/10.1002/2014JB011275>, 2014.
- Llamas, D. and Marfito, B.: Geometry and structure of the offshore segments of the Philippine Fault in Central Philippines, in: *GeoCon 2022: Disruptions, Transformations—Geosciences at the Forefront, Philippines, 2022*.
- 825 Manalo, P. C., Dimalanta, C. B., Faustino-Eslava, D. V., Payot, B. D., Ramos, N. T., Queaño, K. L., Perez, A. D. C., and Yumul Jr., G. P.: Geochemical and Geophysical Characteristics of the Balud Ophiolitic Complex (BOC), Masbate Island, Philippines: Implications for its Generation, Evolution and Emplacement, *Terrestrial, Atmospheric and Oceanic Sciences*, 26, 687, [https://doi.org/10.3319/tao.2015.05.19.01\(tc\)](https://doi.org/10.3319/tao.2015.05.19.01(tc)), 2015.
- Manighetti, I., Mercier, A., and De Barros, L.: Fault Trace Corrugation and Segmentation as a Measure of Fault Structural Maturity, *Geophysical Research Letters*, 48, e2021GL095372, <https://doi.org/https://doi.org/10.1029/2021GL095372>, e2021GL095372 2021GL095372, 2021.
- 830 Marone, C., Scholz, C., and Bilham, R.: On the Mechanics of Earthquake Afterslip, *Journal of Geophysical Research*, 96, 8441–8452, <https://doi.org/10.1029/91JB00275>, 1991.
- Michel, R., Avouac, J.-P., and Taboury, J.: Measuring ground displacements from SAR amplitude images: Application to the Landers Earthquake, *Geophysical Research Letters*, 26, 875–878, <https://doi.org/10.1029/1999g1900138>, 1999.
- 835 Miraballes, M., Mirana, R., Ng, J., Rejuso, A., and Villanueva, F.: Report on the Rapid Assessment on the effects of the Magnitude 6.6 Earthquake in the Municipalities of Cataingan and Pio V. Corpuz, Province of Masbate, Tech. rep., Mines and Geosciences Bureau - Region V, 2020.
- Mogi, K.: Some Discussions on Aftershocks, Foresocks and Earthquake Swarms : the Fracture of a Semi-infinite Body Caused by an Inner Stress Origin and Its Relation to the Earthquake Phenomena, *Bulletin of the Earthquake Research Institute, University of Tokyo*, 41, 615–658, <https://doi.org/https://doi.org/10.15083/0000033716>, 1963.
- 840 Morell, K. D., Styron, R., Stirling, M., Griffin, J., Archuleta, R., and Onur, T.: Seismic Hazard Analyses From Geologic and Geomorphic Data: Current and Future Challenges, *Tectonics*, 39, e2018TC005365, <https://doi.org/https://doi.org/10.1029/2018TC005365>, e2018TC005365 10.1029/2018TC005365, 2020.
- 845 NDRRMC: Sitrep re Magnitude 6.6 Earthquake in Cataingan, Masbate, <https://ndrrmc.gov.ph/index.php/20-incidents-monitored/4096-sitrep-re-magnitude-6-6-earthquake-in-cataingan,-masbate.html>, 2020.
- Ozawa, S., Nishimura, T., Suito, H., Kobayashi, T., Tobita, M., and Imakiire, T.: Coseismic and postseismic slip of the 2011 magnitude-9 Tohoku-Oki earthquake, *Nature*, 475, 373–376, <https://doi.org/10.1038/nature10227>, 2011.
- 850 Perrin, C., Manighetti, I., Ampuero, J.-P., Cappa, F., and Gaudemer, Y.: Location of largest earthquake slip and fast rupture controlled by along-strike change in fault structural maturity due to fault growth, *Journal of Geophysical Research: Solid Earth*, 121, 3666–3685, <https://doi.org/10.1002/2015JB012671>, 2016.
- PHIVOLCS: Historical Earthquakes in the Philippines, 1:5,000,000 map, 1999.
- PHIVOLCS: Active Faults Map - Province of Masbate, <https://gisweb.phivolcs.dost.gov.ph/gisweb/earthquake-volcano-related-hazard-gis-information>, 2018.

- 855 PHIVOLCS: 2020 Masbate Earthquake, <https://www.phivolcs.dost.gov.ph/index.php/2-uncategorised/10490-2020-masbate-earthquake>, 2020.
- Pinet, N. and Stephan, J. F.: The Philippine wrench fault system in the Ilocos Foothills, northwestern Luzon, Philippines, *Tectonophysics*, 183, 207–224, [https://doi.org/https://doi.org/10.1016/0040-1951\(90\)90417-7](https://doi.org/https://doi.org/10.1016/0040-1951(90)90417-7), geodynamic evolution of the Eastern Eurasian Margin, 1990.
- Planet Team: Planet Application Program Interface: In *Space for Life on Earth*, <https://api.planet.com>, 2017–.
- 860 Porth, H., Müller, C., and von Daniels, C. H.: The sedimentary formations of the Visayan Basin, Philippines, *Geol. Jahrbuch Reihe*, 1989.
- Rangin, C., Le Pichon, X., Mazzotti, S., Pubellier, M., Chamot-Rooke, N., Aurelio, M., Walpersdorf, A., and Quebral, R.: Plate convergence measured by GPS across the Sundaland/Philippine Sea Plate deformed boundary: the Philippines and eastern Indonesia, *Geophysical Journal International*, 139, 296–316, <https://doi.org/10.1046/j.1365-246X.1999.00969.x>, 1999.
- Reilinger, R. E., Ergintav, S., Burgmann, R., McClusky, S., Lenk, O., Barka, A., Gurkan, O., Hearn, L., Feigl, K. L., Cakmak, R., Aktug, B., Ozener, H., and Toksoz, M. N.: Coseismic and postseismic fault slip for the 17 august 1999, $m = 7.5$, izmit, turkey earthquake, *Science*, 289, 1519–1524, <https://doi.org/10.1126/science.289.5484.1519>, 2000.
- 865 Rosu, A.-M., Pierrot-Deseilligny, M., Delorme, A., Binet, R., and Klinger, Y.: Measurement of ground displacement from optical satellite image correlation using the free open-source software MicMac, *ISPRS Journal of Photogrammetry and Remote Sensing*, 100, 48–59, <https://doi.org/10.1016/j.isprsjprs.2014.03.002>, 2015.
- 870 Scheffler, D., Hollstein, A., Diedrich, H., Segl, K., and Hostert, P.: AROSICS: An Automated and Robust Open-Source Image Co-Registration Software for Multi-Sensor Satellite Data, *Remote Sensing*, 9, 676, <https://doi.org/10.3390/rs9070676>, 2017.
- Scholz, C.: Earthquakes and friction laws, *Nature*, 391, 37–42, <https://doi.org/10.1038/34097>, 1998.
- SEASSEE: Series on Seismology Volume IV Philippines, Government Printing Office, 1985.
- Seno, T.: The instantaneous rotation vector of the Philippine sea plate relative to the Eurasian plate, *Tectonophysics*, 42, 209–226, [https://doi.org/https://doi.org/10.1016/0040-1951\(77\)90168-8](https://doi.org/https://doi.org/10.1016/0040-1951(77)90168-8), 1977.
- 875 Simborio, T., Yokoi, T., and Hara, T.: Strong ground motion simulation of the 2020 Masbate, Philippines earthquake (Mw6.6) using Empirical Green's Function method, <https://iisee.kenken.go.jp/synodb/data/20230117f70f6a30.pdf>, synopsis of IISEE-GRIPS, 2022.
- Stein, R.: Earthquake Conversations, *Scientific American Sp*, 15, 82–89, <https://doi.org/10.1038/scientificamerican0705-82sp>, 2003.
- Stumpf, A., Malet, J.-P., and Delacourt, C.: Correlation of satellite image time-series for the detection and monitoring of slow-moving landslides, *Remote Sensing of Environment*, 189, 40–55, <https://doi.org/10.1016/j.rse.2016.11.007>, 2017.
- 880 Tiongson, S. and Ramirez, R.: Mapping of ground surface deformations and its associated damage using SAR interferometry: a case study of the 2020 Masbate earthquake, *E3S Web Conf.*, 347, 03 014, <https://doi.org/10.1051/e3sconf/202234703014>, 2022.
- Tomita, F., Iinuma, T., Ohta, Y., Hino, R., Kido, M., and Uchida, N.: Improvement on spatial resolution of a coseismic slip distribution using postseismic geodetic data through a viscoelastic inversion, *Earth, Planets and Space*, 72, 84, <https://doi.org/10.1186/s40623-020-01207-0>, 2020.
- 885 Treiman, J. A.: Primary surface rupture associated with the mw 7. 1 16 october 1999 hector mine earthquake, san bernardino county, california, *Bulletin of the Seismological Society of America*, 92, 1171–1191, <https://doi.org/10.1785/0120000923>, 2002.
- Tronin, A.: Remote sensing and earthquakes: A review, *Physics and Chemistry of the Earth, Parts A/B/C*, 31, 138–142, <https://doi.org/10.1016/j.pce.2006.02.024>, 2006.
- 890 Tsutsumi, H. and Perez, J.: Along strike variation of seismic behavior of the Philippine fault based on historical- and paleo-seismicity, in: *AGU Fall Meeting Abstracts*, vol. 2011, pp. T43D–2366, 2011.

- Tsutsumi, H. and Perez, J.: Large-scale active fault map of the Philippine fault based on aerial photograph interpretation, *Active Fault Research*, 39, 29–37, https://doi.org/10.11462/afr.2013.39_29, 2013.
- 895 Tsutsumi, H., Perez, J., and Lienkaemper, J.: Variation of Surface Creep Rate Along the Philippine Fault Based on Surveys of Alinement Arrays and Offset Cultural Features, in: American Geophysical Union, Fall Meeting 2016, 2016.
- Tymofeyeva, E. and Fialko, Y.: Geodetic Evidence for a Blind Fault Segment at the Southern End of the San Jacinto Fault Zone, *Journal of Geophysical Research: Solid Earth*, 123, 878–891, <https://doi.org/https://doi.org/10.1002/2017JB014477>, 2018.
- USGS: Advanced National Seismic System (ANSS) Comprehensive Catalog, <https://doi.org/10.5066/F7MS3QZH>, 2017.
- 900 Velasco, A. A., Ammon, C. J., Lay, T., and Hagerty, M.: Rupture process of the 1990 Luzon, Philippines (Mw = 7.7), earthquake, *Journal of Geophysical Research: Solid Earth*, 101, 22 419–22 434, <https://doi.org/https://doi.org/10.1029/96JB02290>, 1996.
- Wang, K. and Fialko, Y.: Space geodetic observations and models of postseismic deformation due to the 2005 M7.6 Kashmir (Pakistan) earthquake, *Journal of Geophysical Research: Solid Earth*, 119, 7306–7318, 2014.
- Wang, S., Xu, W., Xu, C., Yin, Z., Burgmann, R., Liu, L., and Jiang, G.: Changes in groundwater level possibly encourage shallow earthquakes in central australia: the 2016 petermann ranges earthquake, *Geophysical Research Letters*, 46, 3189–3198, 905 <https://doi.org/10.1029/2018GL080510>, 2019.
- Wang, S., Song, C., Li, S., and Li, X.: Resolving co- and early post-seismic slip variations of the 2021 MW 7.4 Maduo earthquake in east Bayan Har block with a block-wide distributed deformation mode from satellite synthetic aperture radar data, *Earth and Planetary Physics*, 6, <https://doi.org/10.26464/epp2022007>, 2021.
- Wei, M. and Sandwell, D. T.: Decorrelation of L-Band and C-Band Interferometry Over Vegetated Areas in California, *IEEE Transactions on Geoscience and Remote Sensing*, 48, 2942–2952, <https://doi.org/10.1109/TGRS.2010.2043442>, 2010.
- 910 Wells, D. and Coppersmith, K.: New Empirical Relationships among Magnitude, Rupture Length, Rupture Width, Rupture Area, and Surface Displacement, *Bulletin of the Seismological Society of America*, 84, 974–1002, 1994.
- Woo, G.: Downward Counterfactual Search for Extreme Events, *Frontiers in Earth Science*, 7, <https://doi.org/10.3389/feart.2019.00340>, 2019.
- 915 Woo, G. and Mignan, A.: Counterfactual Analysis of Runaway Earthquakes, *Seismological Research Letters*, 89, 2266–2273, <https://doi.org/10.1785/0220180138>, 2018.
- Wu, J. E., McClay, K., Whitehouse, P., and Dooley, T.: 4D analogue modelling of transtensional pull-apart basins, *Marine and Petroleum Geology*, 26, 1608–1623, <https://doi.org/https://doi.org/10.1016/j.marpetgeo.2008.06.007>, 2009.
- 920 Xu, C., Liu, Y., Wen, Y., and Wang, R.: Coseismic slip distribution of the 2008 mw 7. 9 wenchuan earthquake from joint inversion of gps and insar data, *Bulletin of the Seismological Society of America*, 100, 2736–2749, <https://doi.org/10.1785/0120090253>, 2010.
- Yagi, Y., Kikuchi, M., and Sagiya, T.: Co-seismic slip, post-seismic slip, and aftershocks associated with two large earthquakes in 1996 in Hyuga-nada, Japan, *Earth, Planets and Space*, 53, 793–803, <https://doi.org/10.1186/BF03351677>, 2001.
- Yen, J., Lu, C., Dorsey, R. J., Kuo-Chen, H., Chang, C., Wang, C., Chuang, R. Y., Kuo, Y., Chiu, C., Chang, Y., Bovenga, F., and Chang, W.: Insights into Seismogenic Deformation during the 2018 Hualien, Taiwan, Earthquake Sequence from InSAR, GPS, and Modeling, 925 *Seismological Research Letters*, 90, 78–87, <https://doi.org/10.1785/0220180228>, 2018.
- Yunjun, Z., Fattahi, H., and Amelung, F.: Small baseline InSAR time series analysis: Unwrapping error correction and noise reduction, *Computers & Geosciences*, 133, 104 331, <https://doi.org/https://doi.org/10.1016/j.cageo.2019.104331>, 2019.