

# Surface Rupture Kinematics of the 2020 $M_w$ 6.6 Masbate (Philippines) Earthquake determined from Optical and Radar Data

Khelly Shan Sta. Rita<sup>1</sup>, Sotiris Valkaniotis<sup>2</sup>, and Alfredo Mahar Francisco Lagmay<sup>1,3</sup>

<sup>1</sup>National Institute of Geological Sciences, University of the Philippines, Diliman, Quezon City, Philippines 1101

<sup>2</sup>Department of Civil Engineering, Democritus University of Thrace, Xanthi, Greece 67100

<sup>3</sup>UP Resilience Institute, University of the Philippines, Diliman, Quezon City, Philippines 1101

**Correspondence:** Khelly Shan Sta. Rita (kcstarita@up.edu.ph)

**Abstract.** Optical correlation, interferometry, and field investigation of laterally offset features were undertaken to analyze the kinematics of the 2020  $M_w$ 6.6 Masbate earthquake. Ground displacement fields show a peak left lateral offset of 0.6 m corresponding to  $M_w$ 6.6 geodetic moment magnitude, with an average 0.4 m left lateral slip. The slip distributions also indicate a single asperity located ~200 m SE of the centroid. Post-seismic deformation estimates from interferometry highlight an at least 0.14 m left lateral offset consistent with a  $M_w$ 6.2 post-seismic moment magnitude. The total and post-seismic slip distributions coincide with each other, with both peaks adjacent to the centroid. Slip measurements and the ~28.2 to 41 km rupture length estimates from field and remote sensing datasets characterize the Masbate segment as capable of producing long ruptures with significant offsets despite the presence of interseismic creep. Post-seismic interferograms resolved the rupture far better than optical correlation, which was degraded due to high-amplitude noise from sensor and environmental sources. Nevertheless, this review of the 2020  $M_w$ 6.6 Masbate earthquake provides a comprehensive slip measurement of the surface rupture and demonstrates the presence of two transtensional basins in the Masbate province, revealing new insights into the seismic hazard and seismotectonic setting of the Central Philippines.

## 1 Introduction

Philippine earthquake geodesy traditionally relies on interferometry, global navigation satellite system (GNSS), waveform modeling, and field investigation (Velasco et al., 1996; Yen et al., 2018). The techniques have been used to investigate the rupture kinematics of significant earthquakes in the Philippines. Earthquake kinematics describe the three-dimensional motion of the ground surface during an earthquake event, and is characterized by the spatial distribution of ground displacements along the fault rupture. It is crucial in seismic hazard assessments since it can be used in estimating the possible ground shaking intensity and fault rupture potential of a particular location (Morell et al., 2020).

A  $M_w$ 6.6 earthquake jolted the island of Masbate on the 18<sup>th</sup> of August 2020, and was followed by a  $M$ 5.7 aftershock in about 10 minutes (Aurelio et al., 2021). In the span of 32 hours after the mainshock, 244 aftershocks were recorded ranging from  $M$ 1.6 to  $M$ 5.1 (PHIVOLCS, 2020). The focal mechanism solution of the mainshock illustrates a strike-slip movement with hypocenter at 7 km S29°E of Cataingan, and a depth of 21 km (PHIVOLCS, 2020). The earthquake resulted in devastating losses in terms of human lives and properties (NDRRMC, 2020). The heavy impacts of the earthquake warrants a thorough

25 evaluation of the event to support the assessment of seismic hazards to mitigate losses and damages for the next earthquakes in the island of Masbate.


Preliminary assessments using Sentinel-1 interferograms showed butterfly-shaped fringes and phase shifts between 3 to 30 cm along the descending track line-of-sight (LOS) (PHIVOLCS, 2020) and >15 cm LOS ground deformation along the ascending track (Tiongson and Ramirez, 2022). The ascending interferograms used a 12-day (15 to 27 Aug 2020) temporal  
30 baseline, resulting in low phase coherence of the InSAR pairs. Spectral analysis of seismic waveforms (Simborio et al., 2022) outlined a  $3.0 \text{ km s}^{-1}$  rupture velocity beginning on a shallow region of the southern fault plane with peak ground displacements  $\geq 1.0 \text{ m}$ .

We introduce the application of optical image correlation (OIC) in the Philippines to highlight its importance when used alongside interferometry. The simultaneous application of OIC and InSAR provides complementary information on the decor-  
35 related portions of the coseismic interferogram to improve the understanding of the 2020  $M_w 6.6$  Masbate earthquake since the event is still relatively unstudied compared to the 2003  $M_s 6.2$  event that occurred on the same fault segment (Besana and Ando, 2005; Lai et al., 2019; Tsutsumi and Perez, 2011, 2013).

This study presents first-order ground displacements and post-seismic slip from the Masbate segment of the Philippine Fault. The resulting slip distributions and surface rupture parameters refine our understanding of the Masbate segment, shed light on  
40 the relationship of rupture initiation with mesoscale structures, topography, and bedrock lithology, and provide insights into the underlying faulting mechanisms. The accompanying morphological interpretations of surface ruptures highlight the known stress regime in Masbate, expanding upon our understanding of the structural characteristics of the central Philippines. These findings contribute to the enhancement of earthquake models for use in seismic hazard assessments in the region.

## 2 Geologic Setting and Tectonic Framework

45 The Philippines is a region of active tectonics and volcanism arising from the interaction of the Sunda Block of the Eurasian Plate, and the Philippine Sea Plate (Aurelio, 2000a; Bird, 2003) (Fig. 1a). Oppositely dipping subduction margins flank the archipelago: the east-dipping Manila, Sulu, Negros, and Cotabato Trenches in the west; and the west-dipping Philippine Trench and East Luzon Trough in the east (Aurelio, 2000a; Barrier et al., 1991). The eastern margin accommodates the northwesterly advance of the Philippine Sea Plate at a rate of  $3\text{-}9 \text{ cm yr}^{-1}$  relative to the Eurasian Plate (Seno, 1977), while the western end  
50 consumes the subducting Sunda Block moving at a rate of  $1 \text{ cm yr}^{-1}$  eastward relative to the Eurasian Plate (Chamot-Rooke and Le Pichon, 1999). The interaction and oblique convergence of the tectonic plates that bind the country resulted in the formation of the approximately 1,200 km long left lateral Philippine Fault (PF) in the Middle Miocene (Allen, 1962; Fitch, 1972; Tsutsumi and Perez, 2013; Pinet and Stephan, 1990) as the Philippine Sea Plate's motion shifted counterclockwise from northward to a northwestward direction (Aurelio, 2000a). An Association of Southeast Asian Nations (ASEAN)-wide  
55 GPS network assessment (Aurelio et al., 1997; Aurelio, 1998, 2000b; Rangin et al., 1999) measured a  $2\text{-}3 \text{ cm yr}^{-1}$  current slip rate throughout the fault zone. An approximately 580 km curved section of the PF traverses the central Philippines from the southeastern portion of Quezon province to Leyte island (Fig. 1b). Coined the Philippine Fault Bend, the deformation



**Figure 1.** A) Regional geologic setting of the Philippine archipelago showing the adjacent Philippine Sea Plate (PSP) and Eurasian Plate (EU). The large-scale tectonic features are also depicted, including the east-dipping subduction zones - the Manila Trench (MT), Negros Trench (NT), Cotabato Trench (CT), and Sulu Trench (ST) on the western side; west-dipping trenches - East Luzon Trough (ELT) and Philippine Trench (PT) on the eastern side; and the resultant Philippine Fault running across the archipelago. White arrow indicates the movement direction of the Philippine Sea Plate. B) Major structural features directly affecting Masbate including the central Philippine Fault Zone (Guinayangan, Masbate, and Leyte segments), their approximate offshore projections, the Legaspi Lineament, and the Sibuyan Sea Fault. C) Southeastern Masbate area showing the involved lithologic units and traces of the Philippine Fault and Uson Fault. Black-white bounding box highlights the primary region of interest. Lithology adapted from Manalo et al. (2015). Trace of active structural features adapted from PHIVOLCS. D) Historical seismicity of the Masbate island from 1917 to 2022 showing the moderate earthquakes ( $M4\sim 6.6$ ) with focal mechanism solutions of the February 2003 and August 2020 events. Background topography and bathymetry from GEBCO. Earthquake data acquired from USGS-NEIC. Focal mechanism solutions sourced from GlobalCMT.

zone in the midsection of the PF is characterized by a N50°W fault strike and forms a releasing bend structure relative to the approximately N20°W general strike north and south of the central section (Lagmay et al., 2005).

## 60 2.1 Geology of Masbate

The PF and Sibuyan Sea Fault (SSF) are the prominent structural features in the Masbate island (PHIVOLCS, 2020; Tsutsumi and Perez, 2013) (Fig. 1b). The Masbate segment of the PF extends from Burias and Ticao islands in the north, and strikes N40°W as it crosses the southeastern tip of the mainland (Aurelio et al., 1991). In terms of morphology, it spans approximately 30 km along a linear trough parallel to the coastline (PHIVOLCS, 2020; Tsutsumi and Perez, 2013). Complexities in the fault geometry include a fault complexity between 12°08' N and 12°06' N latitude, and a bifurcation around 12°03' N. The northeastern leg of the bifurcated structure is curved at 12° N and connects to the Leyte segment, whereas the southwestern leg strikes at N20°W and appears to terminate at Cataingan Bay (PHIVOLCS, 2018). Offshore seismic profiling in the bay (Llamas and Marfito, 2022) show left-stepping faults and negative flower structures that denote transtensional deformation in the region.

70 The SSF is another left lateral strike-slip fault with a normal component (Aurelio et al., 1991; Bischke et al., 1988). It runs parallel to the PF in mainland Masbate then deflects westward into the Sibuyan Sea (Fig. 1b). The extensional component associated with the SSF is represented by tilted blocks formed from normal faulting and bathymetric depressions in the Sibuyan Sea (Bischke et al., 1990). The SSF influences the local stress regime expressed by alternating extensive and compressive structures, as it interacts with the PF (Aurelio et al., 1997). The SSF was also described as separate from the PF and is a manifestation the transtensional deformation with the  $\sigma_3$  oriented perpendicular to the PF (Aurelio, 1992; Aurelio et al., 1991). GPS measurements further substantiate this regime on the local scale (Bacolcol, 2003; Duquesnoy et al., 1994).

Ophiolites comprise the basement complex of Masbate island and are unconformably overlain by a series of sedimentary and igneous units (Fig. 1c). The Cretaceous Panguiranan Chert represents the pelagic cover of the ophiolite sequence (Aurelio and Peña, 2010). The Late Oligocene Nabangig Formation is the oldest lithologic unit underlying the area of interest and is composed of a clastic sequence with recrystallized limestone. The Middle Miocene Lanang Formation lies unconformably over the Nabangig Formation, and consists of highly deformed sedimentary rocks ranging from conglomerates to mudstones (Manalo et al., 2015). Majority of the study area is underlain by the Miocene-Pliocene Buyag Formation. It overlies the Lanang Formation and is composed of conglomerates with fine-grained igneous clasts and interbedded shale and siliceous sediments. The largely coralline Pleistocene Port Barrera Limestone is the youngest stratigraphic unit in the province (Porth et al., 1989).

## 85 2.2 Seismotectonics of the Masbate Segment

Bacolcol (2003) and Lai et al. (2019) described the Masbate segment as the transition zone between the northern and southern segments of the central PF. Historical earthquake information revealed recurrence intervals of as short as five years (Besana and Ando, 2005) for moderate magnitude ( $M_{5.5-6.2}$ ) earthquakes to as much as 238 years using slip deficit rates between the 2003 event and interseismic velocities (Lai et al., 2019). A marked absence of strong events beyond  $M>7.0$  in the past four centuries is also apparent (Bautista and Oike, 2000; SEASEE, 1985). A map (Fig. 1d) of moderate seismicity between 1917 and 2022 culled

from the United States Geological Survey - National Earthquake Information Center (USGS-NEIC) (USGS, 2017) highlights the earthquake epicenters related to the Masbate segment and the SSF during this period. Observations and discussions on the 2003 event led Besana and Ando (2005) to characterize the seismic activity of the segment as capable of generating moderate to large events that produce ruptures larger than expected and can be succeeded by post-seismic deformation or creep.

95 The 2003  $M_s 6.2$  event is the previous notable earthquake in the province that produced an approximately 20 km long rupture with epicenter in the northern portion of the Masbate segment. It was accompanied by either post-seismic deformation or a slow creep component (Besana and Ando, 2005; Lai et al., 2019; Tsutsumi and Perez, 2011, 2013). Kinematic offsets were measured at a maximum of 47 to 50 cm, which is beyond the expected peak offsets for a  $M 6.2$  earthquake, based on empirical magnitude scaling equations (Wells and Coppersmith, 1994). Lai et al. (2019) attributed the post-seismic deformation to the  
100 significant excess moment released by the earthquake. Aftershock analysis of earthquakes within two months of the 2003 temblor showed a cluster of  $M 5.0-6.0$  epicenters at the southern portion of the fault. The aftershocks were accompanied by the development of hairline fractures near Cataingan that expanded to as much as 8 mm (Besana and Ando, 2005). In addition, Bacolcol et al. (2005) recorded 10 cm GPS deviations within six months after the mainshock, supporting observations of post-seismic deformation. Coulomb stress transfer (CST) analysis showed that the 2003 event raised the stress in the southern part  
105 of Masbate by approximately 0.3 to 1 bar (Legaspi et al., 2018), which is presumed to have reduced the time elapsed before the 2020 earthquake.

The apparent absence of strong earthquakes ( $M > 7.0$ ) in the Masbate segment may be explained by the presence of creep (Scholz, 1998). Aseismic slip during interseismic periods was measured at  $2.3 \pm 1.1$  cm yr<sup>-1</sup> through GPS (Bacolcol et al., 2005) and 0.7-1.7 cm yr<sup>-1</sup> through alignment arrays (Tsutsumi et al., 2016). A moderate background seismicity can be observed  
110 along the Masbate segment (Fig. 1d) and is further described in PHIVOLCS reports (1999).

### 3 Data and Methods

The surface displacement components of the Masbate earthquake were measured using optical and radar satellite data. Pixel offset tracking on PlanetLabs image pairs with 3.125 m resolution was accomplished using the `mm2dpossism` binary of the MicMac program (Rosu et al., 2015) for the horizontal ground displacements. Interferometric Synthetic Aperture Radar  
115 (InSAR) was done using the European Space Agency's (ESA) Sentinel-1A and 1B radar data with a 6-day temporal baseline (14 - 21 Aug 2020) for the coseismic slip and a 31-day (20 Aug - 20 Sept 2020) time series after the mainshock for the post-seismic deformation following the rapid aftershock decay duration.

#### 3.1 Spatiotemporal Seismicity Analysis

Earthquake data from 2019 to 2021 with magnitudes  $M \geq 1.0$  were gathered from the PHIVOLCS online earthquake archive  
120 (Supplement S3). Foreshock and aftershock patterns were analyzed by plotting the distribution of both magnitude and frequency in time and space. All seismicity information was primarily used to define the earthquake periods (Farrell et al., 2009; Mogi, 1963) related to the 2020 Masbate event. Focal mechanism solutions acquired from PHIVOLCS, USGS, and Glob-

alCMT were used for the analysis of the nodal planes present in the double-couple as a seismological basis of the sense of movement of the fault plane responsible for the mainshock.

## 125 3.2 Optical Image Correlation

PlanetScope satellite imagery (Planet Team, 2017–) was the primary dataset due to the daily data acquisition and high spatial resolution of available products. Images with the least cloud cover covering the region of interest (Fig. 1c) with a maximum 15 % obstruction threshold were selected. Ortho Tile products with a 3.125 m ground resolution, processed at level 3A and corrected for surface reflectance (SR) were utilized. When Ortho Tile products were unavailable, we used Ortho Scene products  
130 downsampled to 3.125 m ground resolution and corrected for top-of-atmosphere (TOA) reflectance.

We realize the necessity for the shortest temporal baseline to properly delimit the slip components. However, no workable Planet datasets in terms of cloud cover, ROI coverage, or compatibility of acquisition parameters were available that more closely bracketed the earthquake. To observe the influence of time on the quality of the output offset rasters, two sets of temporal ranges were tested for the Masbate event (Fig. 2). Five short-term pairs were selected from visually feasible images  
135 sampled months apart in 2020, whereas ten *annual* pairs were roughly one year apart (Table 1). This was done to minimize the effect of environmental variables following the approach of Elliott et al. (2020a).

The image pairs were coregistered to a Sentinel-2 basemap using the global coregistration function of the Automated and Robust Open-Source Image Co-Registration Software (AROSICS) (Scheffler et al., 2017) to increase the registration accuracy while preserving the deformation signal. The pairs were clipped to their overlapping extents with a Universal Transverse  
140 Mercator (UTM) projection and either 8- or 16-bit integer pixel data type as required by MicMac. Manual cloud masking was accomplished using the `saisiemasqQT` binary of MicMac for precision masking.

Correlations in the spatial domain were performed with MicMac on the prepared image pairs using the following parameters:

- 7x7 window size ( $SzW=3$ ) for a 21.875 m search grid on each side to compensate for feature-related noise and expected displacement,
- 145 – 0.5 regularization factor ( $Reg=0.5$ ) to provide distance-based correlation weights on the neighboring pixels within the window for noise reduction, and,
- pixel subsampling parameter of 2 ( $SsResolOpt=2$ ) as recommended by Canizares et al. (2020) for earthquake events to resolve displacement fields close to the ground resolution of the source imagery.

The resulting output files were georeferenced using `gdal_translate` with the `-a_ullr` flag (GDAL/OGR contributors, 2021) to ensure a 1:1 alignment with the input image pairs without warping the rasters. To retain most deformation signals, the correlation threshold was set to 50 %. A non-local median (NLM) filter with a low H-noise value was applied to prevent flattening real ground deformation signals while improving the signal-to-noise ratio (SNR). The outputs were then translated into the spatial domain by multiplying the rasters with the ground resolution of the input files. The rasters chosen to extract profiles and calculate fault-parallel offsets were selected based on noise levels, atmospheric obstruction, pre-processing accuracy, ROI  
155 coverage, and quality of slip visualization (Supplement S1.1).

**Table 1.** Satellite geometry and acquisition parameters of the PlanetScope data used in the study. Values of the satellite azimuth (satAzi) are unavailable for the older images in the table. Available tiles for 23 June 2020 are merged to create a single mosaic to maximize the extent. The two datasets for 2 April 2020 are analyzed separately.


Acquisition	satID	satAzi	view $\angle$	sunAzi	sunElev
2019-11-28T01:53:43	103e	-	1.2	143.9	48.7
2020-04-09T00:33:57	100d	-	0	90.3	42.7
2020-04-17T02:35:36	106e	-	2.1	92.5	73.1
2020-06-06T00:26:15	0f32	-	3	69.9	41.8
2020-06-23T01:29:47	2263	272.7	2	66.4	55.4
2020-06-23T01:54:27	100a	102.6	1	63.2	61.1
2020-07-07T00:19:38	1048	204.5	0.1	70.4	39.2
2020-11-18T02:03:10	105c	98.7	5	146.2	52.6
2020-11-29T01:51:59	1010	98.0	5	143.8	48.8
2020-12-24T02:21:12	1069	261.0	3	149.9	49.4
2021-04-01T02:16:14	241c	99.1	2	106.1	66.2
2021-04-02T02:13:12	1105	168.7	0.1	104.3	65.6
2021-04-02T01:54:40	1013	276.8	5	101.2	60.8
2021-04-11T01:33:11	2276	98.6	4.1	93.2	57.7
2021-04-12T02:15:35	227c	98.4	4.9	96.1	68.1
2021-06-05T01:51:14	1038	98.2	4.9	63.9	61.7
2021-06-10T23:23:03	1049	84.5	3	70	27.5
2021-07-08T02:15:18	2274	12.8	0.2	61.7	65.1
2021-07-11T01:26:43	2434	277.1	5	68.8	54.1

The post-processed data generated coherent pixel offset maps along the N-S ( $D_{N_{OIC}}$ ) and E-W ( $D_{E_{OIC}}$ ) horizontal axes with a 3.125 m pixel resolution. Slip measurements with respect to the general fault strike azimuth ( $\theta = 325^\circ$ ) of the Masbate segment were acquired by transposing the N-S and E-W slip into fault-oriented components using the following equations (Elliott et al., 2020a):

$$160 \quad D_{\parallel} = D_{N_{OIC}} \cdot \cos(\theta) + D_{E_{OIC}} \cdot \sin(\theta) \quad (1)$$

$$D_{\perp} = D_{N_{OIC}} \cdot \sin(\theta) - D_{E_{OIC}} \cdot \cos(\theta) \quad (2)$$

Measurements were acquired by taking profiles of the pixel offset data across the fault to determine the net fault-parallel offset (Supplement S1.2). The fault-parallel ( $D_{\parallel}$ ) direction refers to movements parallel to the fault trace represented by laterally



**Figure 2.** Two classes of Planet satellite imagery pairs evaluated through optical correlation. A) Short-term pairs taken within the same year as the Masbate earthquake ranging from five to nine months apart and as close to the event as possible. B) Approximately annual span pairs acquired at about 365 days apart where environmental interference is expected to be at a minimum (Elliott et al., 2020a).

offset features on the field. This component was the sole subject of profiling due to the stronger horizontal offset signal in the  
 165 field. The tension-compression ( $D_{\perp}$ ) aspect detectable from the fault-normal component was not analyzed due to the minimal  
 deformation along the vertical axis, further complicated by high noise amplitudes.

Wide swath profiles (321 x 161 px, roughly ~1.0 x ~0.5 km) along the fault trace were acquired with StackProf (Delorme,  
 2021) developed for the MicMac project. The slip values from profiles aided in the measurement of the near-fault slip distribu-  
 170 tion for comparison with offsets measured in the field. The displacements were calculated from the intercepts of the regression  
 of weighted median offsets across the fault trace. We used the coherence values to remove zero correlation pixels to highlight  
 the genuine earthquake slip in the arctangent model. We selected the regression limits by hand to exclude areas of decorrelation,  
 apparent noise signals, and obvious spurious pixel offset data that remained after filtering. As a result, most regression lines  
 were limited to 300 to 600 m across the fault trace. Validated slip data were plotted along the fault strike to show the spatial  
 trends.

175 Absolute standard deviations ( $1\sigma$ ) from the weighted median values across the transect indicate the measurement uncertain-  
 ties in individual box profiles. The individual slip uncertainties are the sum of the standard errors of intercepts of the regression



**Table 2.** Summary of reference and secondary scenes processed within Alaska Satellite Facility’s Hyp3 processor to create interferograms.

Track	Reference	Path/Frame <sub>1</sub>	Secondary	Path/Frame <sub>2</sub>	Perp. baseline (m)	Interval (days)	Stage
desc	20200814	61/551	20200820	61/549	-76.4	6	co
asc	20200815	69/32	20200821	69/34	107.1	6	co
desc	20200820	61/549	20200901	61/549	50.4	12	post
asc	20200821	69/34	20200827	69/32	-173.3	6	post
desc	20200826	61/551	20200901	61/549	-76.4	6	post
asc	20200827	69/32	20200902	69/34	126.0	6	post
asc	20200827	69/32	20200908	69/32	42.6	12	post
desc	20200901	61/549	20200907	61/551	49.6	6	post
desc	20200901	61/549	20200913	61/549	68.5	12	post
asc	20200902	69/34	20200908	69/32	-83.1	6	post
asc	20200908	69/32	20200920	69/32	49.9	12	post
desc	20200913	61/549	20200919	61/551	-39.4	6	post

of the eastern and western blocks (Delorme, 2021). The displacement curve was calculated using a robust weighted moving mean approach with minimal smoothing ( $q = 0.15$ ) to prevent the removal of small deviations along the strike.

### 3.3 Small Baseline Subset (SBAS) Interferometry

180 Coseismic and post-seismic interferograms were processed with NASA’s Alaska Satellite Facility Distributed Active Archive  
Center (ASF DAAC) Hybrid Pluggable Processing Pipeline (HyP3) (Kennedy et al., 2021). Sentinel-1A/B single look complex  
(SLC) C-band radar data acquired through the interferometric wide (IW) swath mode was used. Data with 6 to 12-day intervals  
from paths 61 (descending track) and 69 (ascending track) (Fig. 1d) imaged between 14 to 21 August 2020 for the coseismic  
slip and 20 August to 20 September 2020 for the afterslip were multilooked by 20 in the azimuth direction and 4 in the range  
185 resulting to a pixel size of 80 x 80 m to minimize the effect of land cover (Table 2).

Small baseline time series analysis (SBAS) was employed using MintPy (Yunjun et al., 2019) to quantify the post-seismic  
deformation in the aftershock observation period. Atmospheric corrections were not applied, since the study area is highly  
localized relative to the global dataset implemented in available databases. Furthermore, the unwrapping reference point was  
set to a common high coherence pixel for both tracks. The ascending ( $V_A$ ) and descending ( $V_D$ ) LOS velocities were projected  
190 into the azimuthal fault strike ( $\theta = 325^\circ$ ) (Dianala et al., 2020; Lindsey et al., 2014; Tymofyeyeva and Fialko, 2018) to estimate  
the fault-parallel ( $V_{||}$ ) and vertical ( $V_z$ ) post-seismic velocities. The approach assumes a zero third component acceptable  
for areas of relatively simple faulting. Equation 3 was used to derive the fault-oriented afterslip LOS velocities within the  
asc\_desc2horz\_vert script in MintPy. The pixel unit vector directed to the satellite was decomposed into  $e$ ,  $n$ , and  $u$   
representing the east, north, and upward directions of both tracks, respectively.

$$195 \begin{bmatrix} V_A \\ V_D \end{bmatrix} = \begin{bmatrix} e_A \sin(\theta) + n_A \cos(\theta) & u_A \\ e_D \sin(\theta) + n_D \cos(\theta) & u_D \end{bmatrix} \begin{bmatrix} V_{\parallel} \\ V_Z \end{bmatrix} \quad (3)$$

The total afterslip was obtained by multiplying the fault-parallel component with the temporal range of the time series and smoothed the data using an NLM filter to arrest pixel outliers. Swath profiles (Krambach, 2015) were taken at 0.5 km intervals with a 3.5 km swath width across the fault strike (Supplement S4.1). We determined the across-fault  $1\sigma$  uncertainties from the absolute deviation from the weighted median in each swath. Slip uncertainty estimates come from the sum of the standard error of regression intercepts of the eastern and western blocks, following the StackProf approach (Delorme, 2021). Similar to optical correlation, the regression lines for the eastern and western blocks were manually delimited in order to exclude the remaining phase unwrapping artifacts and other noise signals.


The spatial distribution of the afterslip along the strike was evaluated using a smoothed moving mean displacement curve ( $q = 0.15$ ). The vertical post-seismic displacement was not investigated due to the primarily horizontal ground movement. The coseismic interferogram was limited to qualitative comparisons with optical correlation. No quantifiable comparisons can be made since slip is unresolved in the near-fault region as a result of decorrelation due to high magnitudes of coseismic slip.

### 3.4 Ground Truthing

Field investigation was conducted from 18 to 21 May 2022 to identify and measure the horizontal offsets across the ground rupture. Locations of known ground rupture (de la Cruz, per. comm., 2022) were complemented by points of interest identified from the image correlation outputs and InSAR. Due to the two-year interval between the earthquake and fieldwork, geomorphic or human modifications of offset features may have occurred. Additionally, many offsets and evidence of surface rupture may have been erased given the small magnitude of surface slip. The significance of these changes can be assessed by correlating field data with other measurement techniques to determine the slip.

We selected points of interest for field checking based on the pixel offset rasters, prioritizing locations with significant contrast along the approximate and certain fault traces. Field offsets were measured with a steel tape and structural orientations were determined with a base plate compass. Measurements of on-fault offsets were limited to individual rupture strands. Due to the scarcity of remnant ruptures, most sites had a single measurement. Uncertainty was estimated by comparing values at these locations to earlier measurements acquired a week after the earthquake (PHIVOLCS, 2020; Miraballes et al., 2020). In areas with complex, closely spaced ruptures, multiple measurements were acquired and statistically assessed to estimate the uncertainty.

Observation points with measurable feature offset and ruptures were plotted along the strike to show the attitude and slip distribution along the fault (Supplement S2). The measured horizontal offsets were used to validate the results of the remote sensing analysis. Finally, the surface rupture length, and maximum and average displacements were derived from the conjoined remote sensing outputs and field data to estimate the seismic moment.



**Figure 3.** A) Earthquake probability density subplot showing the normalized population of earthquakes per day, highlighting the contrast between the periods of the earthquake sequence. B) Plot showing the number of earthquakes with magnitudes  $M \geq 1$  related to the Masbate segment and their temporal distribution relative to the 18 August 2020 mainshock.

## 225 4 Results

### 4.1 Seismicity

#### 4.1.1 Earthquake Frequency and Temporal Relationships

The frequency-time distribution of earthquakes between 25 August 2019 to 18 August 2021 shows the immediate seismicity related to the 18 August 2020  $M_w$  6.6 event (Fig. 3). The foreshocks, mainshock, and aftershocks are identifiable in the probability density subplot (Fig. 3a) as a distinct bell-shaped curve relative to the generally quiescent background activity. The island of Masbate typically experiences less than ten earthquakes per day, corresponding to a moderate background seismicity. An increase in earthquake frequency defines the foreshock sequence (Farrell et al., 2009) at around 32 days before the mainshock, as shown by the distinct trend increase in the probability density subplot. The peaks of the frequency plots (Figs. 3a and b) correspond to 141 earthquakes which were mostly aftershocks on 18 August 2020. A negative slope in the probability density subplot depicts the aftershock sequence. The decaying earthquake frequency is initially symmetrical to the foreshock trend with an inflection point around 30 days after the mainshock, which transitioned to a more gradual decrease. The aftershocks persisted for 166 days or up to 31 January 2021. Succeeding return to background levels saw a similar seismic activity to the previous background activity from 2019 to early 2020.

## 4.1.2 Spatiotemporal Earthquake Distribution

240 In general, the background seismicity ( $M \leq 4$ ) is well distributed throughout the overall length of the Masbate segment of the PF traversing the islands of Ticao, Burias, and the provincial mainland (Fig. 4a). However, an absence of seismic activity is noticeable in the northern portion of the fault in mainland Masbate during this period. On the contrary, an earthquake cluster occurred near the location of the mainshock. Foreshock activity is concentrated in southeastern Masbate adjacent to the mainshock and is composed of mostly weak ( $M2 \sim 3$ ) and shallow hypocenters peaking at  $M4.9$  (Fig. 4b).

245 Three moment tensor solutions from GlobalCMT, PHIVOLCS, and USGS highlight the mainshock's dominant left-lateral strike-slip movement (Fig. 4c). The solutions of PHIVOLCS and USGS show a noticeable, albeit minimal, dip-slip component towards the NNE. The epicentral location of the mainshock from GlobalCMT and PHIVOLCS coincided with the fault trace, while USGS' focal mechanism solution deviated to the northeast of the fault.


Aftershocks that transpired on the same day as the mainshock (Fig. 4d) are shallow (1-30 km) and densely grouped in  
250 the southeastern offshore extension of the segment and deepen (10-50 km) as they taper toward the onshore portion. The aftershocks have weak magnitudes ranging from  $M2.0$  to  $M4.4$ . Succeeding first-month aftershocks are weak to moderate, with a wide range of magnitudes between  $M1.7$  to  $M5.1$ , mostly occurring at depths above 30 km. A monthly progressive dispersal of hypocenters in space is also observed, with the first month showing a dense cluster in the southeastern section that tapers to the northwest (Fig. 4e). The pattern and geographic extent of the first-month aftershocks appear to be an expansion  
255 of the same-day aftershocks. The succession of the aftershock patterns in the following months continued to scatter along the Masbate segment and the SSF while retaining a perceptible concentration in the offshore southeastern extension of the causative fault segment (Figs. 4f, 4g, 4h, and 4i). The last two months of aftershock activity (Figs. 4h and 4i) feature the least number of hypocenters throughout the aftershock sequence.

Background levels returned on 2 February 2021 defined by the well-distributed hypocenters on the fault segments with moderate magnitudes (Fig. 4j) similar to the pre-event seismicity in 2019 (Fig. 4a). The frequency and distribution of earthquakes  
260 in time and space enabled the distinction of various stages of the earthquake sequence relative to the 2020 Masbate event.

## 4.2 Optical Image Correlation

### 4.2.1 Long and Short Temporal Baseline

The acquired pixel offset maps (Fig. 5) illustrate the fault rupture and the known left lateral movement of the Masbate segment, despite the presence of noise that obscures portions of the rupture. Only one of the eight image pairs from the short timespan  
265 spaced months apart (Figs. 2a) produced outputs with perceptible rupture due to the low SNR. Annual timespan pairs resolved the rupture more clearly compared the short timespan image pairs. The north-south slip components show better representations of the rupture, whereas the east-west components tend to be noisier. Nevertheless, the valid image pairs (Supplement S1.1) with resolvable slip and subjected to stack profiling exhibit reliable estimates of the slip in the near-field region.



**Figure 4.** Spatiotemporal earthquake distribution for the Masbate segment. A) Background seismicity between 25 Aug 2019 to 10 Jul 2020. B) Foreshocks from 19 Jul to 17 Aug 2020. C) Moment tensors of the mainshock from GlobalCMT, PHIVOLCS, and USGS. D) Same-day aftershocks. E) First-month aftershocks from 19 Aug to 17 Sept 2020. F) Second-month aftershocks from 18 Sept to 17 Oct 2020. G) Third-month aftershocks from 19 Oct to 16 Nov 2020. H) Fourth-month aftershocks from 18 Nov to 14 Dec 2020. I) Last aftershocks from 22 Dec 2020 to 31 Jan 2021. J) Return to background seismicity covering 09 Feb to 18 Aug 2021.

A generalized fault trace (yellow line; Fig. 5) was delineated from the valid pixel offset rasters and the known fault trace, which served as the baseline for slip profiling. The fault trace is mostly straight, except for an apparent northeasterly strike rotation towards the north, where it follows the piedmont of a topographic high in Dimasalang. No measurements were acquired in the area due to anomalously high values, which are probably terrain residuals. Topographic artifacts and decreasing slip away from the epicenter posed difficulties in delineating the northern rupture section near Naro Bay. In the southwestern section, the offset rasters show a new splay striking NW-SE that bisects the previously recognized bifurcation in Cataingan. The rupture traverses the deltaic area and possibly continues southward across Cataingan Bay, then resurfaces on the opposite side of the bay. Smaller fractures and minute rupture components are not visible due to ubiquitous noise.


Left lateral offsets were quantified along the ~25 km onshore length of the fault from Dimasalang to Cataingan, given the inland spatial extents and continuity where optical correlation was feasible. Edge artifacts and the obscured western block in the south of the bay hindered the acquisition of slip profiles.

Fifty-one (51) wide swath profile boxes were generated throughout the fault rupture, with the first box in Dimasalang and the last in Cataingan to determine the along-strike slip distribution (Fig. 6). The optical correlation results generated a total of 126 left-lateral measurements over six image pairs (Supplement S1.2). Using a 325° general strike azimuth of the Masbate segment, the east-west and north-south displacement components (Figs. 5a and b) were transposed into the fault-parallel direction (Fig. 5c).


The 9 April 2020 - 2 April 2021 pair was selected as the representative dataset due to its wide spatial extent and reliable representation of the fault rupture. The north-south component (Fig. 5a) shows a continuous linear rupture trace from Dimasalang to the deltaic area in Cataingan, although the Dimasalang area and its border with Palanas exhibited unusually high values due to topographic artifacts. The east-west component (Fig. 5b) is less coherent, likely due to smaller ground movements in this direction or overprinted data from the noise. The outputs show noise from both artificial and environmental sources, represented by topographic artifacts in the southwestern block, an apparent seamline, and reduced coherence due to differences in land cover between the images. Nevertheless, the fault-parallel component (Fig. 5c) shows good estimates of the left lateral slip, further supported by the near-field offset models.

Box 32 (Fig. 5d), located ~6 km north of the GlobalCMT surface projection, shows a 43 cm slip in the near-field (Fig. 5e). Pixel offset values increase in the far-field for the western block, which may either represent noise or distributed deformation. Box 47, situated ~1.5 km south of the GlobalCMT epicenter, exhibits a 59.2 cm left lateral offset. The box represents the peak offset in this image pair and displays a clear contrast between the east and western blocks.

The plot of the slip distribution along strike (Fig. 6) displays the 126 left lateral offsets compiled from the valid offset rasters to characterize the displacement field. The slip minima ranges from 6 to 9 cm, located 4.5 km from the northern shoreline in Dimasalang. The displacement increased to ~50 cm around the vicinity of the GlobalCMT projection of the centroid. Slip values remained constant within a ~5 km span in Cataingan, and increases to a peak ~60 cm at around ~1 km southeast of the centroid. South of the peak value, the displacement declines toward the shoreline. The skewed semi-elliptical weighted moving



**Figure 5.** Optical correlation results showing the representative data from the 9 April 2020 and 2 April 2021 image pair. Blue lines in panels E and G are the absolute median fault-parallel values across the profile box. Gray shaded region in panels E and G feature the  $1\sigma$  absolute standard deviation of the fault-parallel values across the fault in each box. A) N-S component. Blue refers to southward movement, while yellow refers to northward motion. B) E-W component. Blue refers to westward movement, while yellow refers to eastward motion. C) Fault-parallel component. Pink refers to movements toward the NW while blue denotes motion toward the SE. D) Closer view of box 32 showing the fault-parallel offset. E) Swath profile of box 32 highlighting the 43 cm slip. Lower regression intercept of the western block relative to the eastern block indicates left lateral movement. F) Closer view of box 47 displaying the fault-parallel offset in Cataingan. G) Swath profile of box 47 exhibiting the peak 59.2 cm left lateral offset in this particular image pair.



**Figure 6.** Along strike distribution of the fault-parallel offsets from 126 profile boxes across six image pairs. Error bars refer to the  $1\sigma$  sum of standard errors of the intercepts of regressions of the east and west blocks providing slip uncertainty estimates.

mean curve (solid gray line; Fig. 6) shows the average left lateral offset of  $37.7 \pm 10$  cm, while the individual measurements  
 305 indicate a peak  $60.6 \pm 9.8$  cm offset adjacent to the surface projection of the centroid in Cataingan.


### 4.3 Interferometry

#### 4.3.1 Coseismic Interferograms


The Sentinel-1 coseismic wrapped interferograms in the ascending and descending tracks (Fig. 7a and c) show the butterfly-shaped fringe patterns and oppositely verging LOS displacements, which is typical of strike-slip events. The fringe lobe patterns  
 310 illustrate a range increase from the ascending track and a range decrease from the descending track for the western side of the fault. Since the orbit of the ascending track is subparallel to the regional fault strike, the left lateral movement is indicated with a possible subsidence component. This is consistent with the moment tensor solutions (Fig. 4c), and agrees with the sense of movement shown by the optical correlation outputs and field observations.

Low phase coherence persists in the near-fault region. Widespread decorrelation leads to mostly unusable information upon  
 315 phase unwrapping due to the likelihood of creating unwrapping errors. Noise is similarly dominant and overprints most of the deformation signals in an approximately 18 km-wide region across the fault trace, hence, the lack of near-fault information in preliminary published interferograms (PHIVOLCS, 2020; Tiongson and Ramirez, 2022). In addition, the rupture is unresolved upon phase unwrapping (Fig. 7b and d) due to decorrelation. This loss of resolvable information in the near-field ultimately hinders slip measurements in the coseismic interferograms.





**Figure 7.** Sentinel-1 coseismic interferograms in ascending and descending tracks. A) Wrapped ascending interferogram. B) Unwrapped LOS displacement in the ascending track. C) Wrapped descending track interferogram. D) Unwrapped LOS displacement in the descending track.



**Figure 8.** Post-seismic deformation between 20 August to 20 September 2020. A) Ascending track time series showing range increase (yellow) and decrease (blue) for the eastern and western blocks, respectively. B) Descending track time series showing range decrease (blue) and increase (yellow) for the eastern and western blocks, respectively. C) Projected fault-parallel post-seismic deformation field. D) Fault-parallel afterslip for box 4 in Dimasalang showing the 5.4 cm left lateral afterslip. Lower regression intercept of the western block relative to the eastern block indicates left lateral movement. Median afterslip value is shown by the blue line. Gray background exhibits the  $1\sigma$  uncertainty across the box. E) Swath profile of box 13 indicating an 8.4 cm left lateral afterslip around the Dimasalang-Palanas boundary. F) Swath profile of box 25 showing a 6.9 cm afterslip in Palanas. G) Swath profile of box 39 in Cataingan showing a 12.0 cm left lateral afterslip.

### 320 4.3.2 Post-seismic Deformation Estimates

Post-seismic interferograms capture an improved overall manifestation of the fault rupture compared to the coseismic ones (Fig. 8). This is a consequence of the expected smaller afterslip values relative to the coseismic slip, which decreases the number of fringes and increases the coherence. The temporal coverage of the interferogram stack allows us to quantify afterslip to complement the estimated coseismic slip from optical correlation. However, the first post-earthquake Sentinel-1 revisit occurred  
325 two days after the mainshock, resulting in all measurements to be lower bound estimates of the afterslip.


Unwrapping errors are less frequent on the ascending and descending track interferograms and are further minimized upon time-series stacking. The remaining phase unwrapping errors upon stacking and filtering were manually excluded in profiling. These unwrapping errors occur as significant high amplitude phase shifts and polarity reversals, noticeable in the Dimasalang and Palanas areas.

330 The fault-parallel afterslip deformation field (Fig. 8c) shows a strongly linear and simple fault trace that accommodated the post-seismic deformation. Deviations from the regional fault strike are observed around 5-13 km from Naro Bay in the north, represented by multiple short NNE-SSW oriented ( $\sim 290\text{-}300^\circ$  azimuth) bends which are adjacent to and coincide with the narrow stepover in the Palanas-Dimasalang area. However, the noticeable unwrapping errors and low ground resolution (80 m) precludes the validity and recognition of these rupture complexities.

335 The afterslip is resolved from Naro Bay in the north towards Cataingan Bay in the south along the  $\sim 25$  km simplified fault trace (Supplement S4.1). Forty-nine (49) swath profile boxes were generated along this trace. Individual measurements reveal centimetric fault-parallel afterslip with minimal across-fault variability and minor  $1\sigma$  uncertainties, except where the boxes overlap with the phase unwrapping error patches. The deformation was detected up to the far-field extents of the profiles from most swath boxes.

340 Boxes 4, 13, 25, and 39 (Figs. 8d, e, f, and g) illustrate select wide swath profiles roughly equidistant from each other. Box 4, situated in Dimasalang, shows a 5.4 cm left lateral fault-parallel afterslip and a reasonably low variability across the fault. Unwrapping errors in the far-field at 1.5 km from the fault on both sides caused irregularities in the profile. Box 13, located around the boundary of Palanas and Dimasalang, depicts an 8.4 cm fault-parallel slip. Both sides of the fault exhibit exceedingly even slip across the fault, except for a noticeable dip in the eastern edge. The dip is likely due to data loss since  
345 the box overlapped with the water body to the east. The profile also appears to be offset by about  $\sim 100$  m to the west, which corresponds to one of the abovementioned bends that may be an unwrapping artifact. Still, the measured slip from the arctangent model remains valid. Box 25 shows a 6.9 cm afterslip in Palanas, located at the southern edge of the fault complexity. Multiple across-fault deviations are visible around 200 and 500 m east of the fault and in the western far-field. The eastern anomalies could either be unwrapping or topography related, while the western patch comes from unwrapping issues. Box 39 in Cataingan  
350 displays a 12.0 cm left lateral afterslip with good symmetry and even across-fault distribution.

The along-strike afterslip distribution graph (Fig. 9) depicts the computed fault-parallel offsets from north to south on the mainland. The smallest value is measured in Dimasalang, adjacent to Naro Bay, amounting to 4.3 cm of left lateral afterslip. The ground offsets exhibit a generally increasing trend to around  $\sim 5$  km from the north. Fluctuating values are observed between



**Figure 9.** Along strike distribution of the fault-parallel afterslip formed by the 49 profile boxes. Error bars refer to the  $1\sigma$  standard errors of the intercepts of regressions of the east and west blocks. Solid blue line shows the weighted moving mean afterslip curve.

6.1 to 13.8 km from the north in Dimasalang and Palanas. The measured displacements oscillate between 5.0 to 9.6 cm in this span. The increasing trend continued to a  $14.3 \pm 0.9$  cm peak value in Cataingan, located 24 km from Naro Bay and ~3 km southwest of the GlobalCMT surface projection. The displacements decrease from the largest value to 11.5 cm on the shoreline of Cataingan Bay. The weighted moving mean curve (solid blue line; Fig. 9) indicates a mean left lateral afterslip of at least  $8.8 \pm 0.7$  cm.

## 4.4 Ground Truthing


### 4.4.1 Observed Surface Ruptures

Remnant effects after about two years post-earthquake were observed on laterally offset cultural and geomorphic features from Dimasalang to Cataingan following the known fault trace (Supplement S2). Assisted by the remote sensing data, the observed ground manifestation of the earthquake from fifty-seven (57) stations (Fig. 10a and b) include fractures, reported earthquake-induced landslides (EIL), indications of liquefaction features and lateral spreading, secondary gravitational features, and ruptures with quantifiable displacements. Earthquake effects recorded during the field activity include reported ground shaking from interviews with locals.

The observed rupture zone of the 2020 Masbate earthquake persisted from Dimasalang to Cataingan and crossed Cataingan Bay. Measured surface ruptures strike at a low angle with respect to the regional fault trace (Fig. 10c) with a mean strike azimuth of  $313^\circ$ . Our northernmost resolvable ground offset was observed on a road in Dimasalang, around ~3 km from Naro Bay. An apparent 0.5 cm offset fracture oriented parallel to the fault trace was recorded in this point, possibly representing the minimum northward constraint of the field rupture. A 2-5 cm displacement oriented  $325^\circ$  NW occurring on a residential structure on top of the fault was observed on the border between Dimasalang and Palanas. The rupture displacement increased



**Figure 10.** Compiled ground-truthing data. A) Earthquake hazards aside from ruptures. B) Ruptures symbolized by size showing the amount of slip. C) Rupture orientations. D) Ruptured tree showing cumulative 87 cm slip from 2003 and 2020 events with estimated  $27 \pm 10$  cm offset from the 2020 earthquake. E) Soil mound cut by the fault highlighting a  $48 \pm 3$  cm left lateral offset. F) Rice field embankment traversed by fault showing  $60 \pm 15$  cm displacement. G) Ruptured fishpond dike recording a  $34 \pm 5$  cm offset. H) Abandoned septic tank showing a 30 cm displacement south of Cataingan Bay.



**Figure 11.** Along strike distribution of measured field offsets acquired on May 2022.

further south ~7 km from the northern shoreline, where a 7 cm left lateral offset was seen coupled with spreading of the road pavement. The fault bisected a coconut tree at the southern limit of the fault complexity in Palanas (Fig. 10d) (Aurelio, 2022),  
 375 resulting in a cumulative 87 cm opening from the 2003 and 2020 events. The newer slip is estimated at  $27 \pm 10$  cm on a  $275^\circ$  W oriented rupture. Right-stepping en echelon fractures occurred adjacent to the coconut tree, most of which were eroded at the time of visit.

The rupture is continuous southward, up to 4 km from the GlobalCMT centroid projection. The fault rupture traversed an abandoned house and deflected its wall by roughly ~50 cm. A soil mound at the base of the wall (Fig. 10e), shows a  $297^\circ$   
 380 W-oriented rupture with a  $48 \pm 3$  cm left lateral offset. Residents reported a ~200 m long rupture in the area, which was already mostly eroded. Nevertheless, the remnant right-stepping en echelon fractures formed a linear rupture zone connected to the displaced wall. A 30 cm offset oriented  $279^\circ$  E on a road in Cataingan likely represents the northern indication of the rupture bisecting the bifurcation in Cataingan. The rupture continued to an agricultural area (Fig. 10f), where the embankments are displaced by  $60 \pm 15$  cm, representing the peak field offset measurements. Long, continuous ruptures were observed on  
 385 the rice fields immediately after the event but were erased by the time of the field investigation. Another southerly continuation was identified, where a  $341^\circ$  NW-oriented rupture displaced fishpond levees (Fig. 10g) by about  $34 \pm 5$  cm. Locals stated that the visible rupture appeared lengthy and continuous in the deltaic areas immediately after the earthquake. The rupture was evident again south of Cataingan Bay, manifested on a collinear set of points through several residential structures. A septic tank diagonally cut by a  $279^\circ$  NW-oriented rupture, shows a 30 cm left lateral displacement (Fig. 10h). Offset measurements  
 390 adjacent to the septic tank vary from 5 to 30 cm.

#### 4.4.2 Left-lateral Offset Distribution

The plot of offset features along the fault strike (Fig. 11) shows the minimum 0.5 cm left lateral displacement ~3 km from Naro Bay in Dimasalang. This point corresponds to the northernmost resolvable ground offset. The measurements continually increased to 50 cm beginning at 17 km from Naro Bay, around the boundary of Palanas and Cataingan. The offset measurements remained constant for a ~4 km span followed by an abrupt dip to 30 cm, which increases again to the peak 60 cm offset. The local 30 cm minimum was measured on a ruptured road. This single discordant measurement could be due to off-fault displacement, physical reduction due to post-earthquake repairs, or an inaccurate measurement from a cultural feature. The displacement tapered to 30 cm near the shoreline of Cataingan Bay. Spatially coincident offsets ranging from 5 to 30 cm resulted in a wide range of uncertainties at the southeastern limitation of the plot. The weighted moving mean curve of the slip distribution (broken red line; Fig. 11) for the 2020 Masbate event indicates a 25.3 cm mean left lateral displacement, while the individual observations denote the maximum of 60 cm displacement near the GlobalCMT centroid projection.

## 5 Discussion

### 5.1 Comparison of Optical Correlation and InSAR

#### 5.1.1 Displacement Threshold

Independently applying optical correlation and interferometry to analyze the 2020 Masbate event provided a basis to assess the technical capabilities and shortcomings of the methods in a tropical setting. Peak displacements of the Masbate earthquake amount to ~60 cm, denoting a high interferometric phase gradient for Sentinel-1. SAR interferograms lose coherence when offsets exceed half the interferometric fringe per pixel (Michel et al., 1999), highlighting the effectiveness of the method for small deformations in mm and cm scales (Tronin, 2006). The LOS measurements of PHIVOLCS (2020) and Tiongson and Ramirez (2022) from coseismic interferograms are likely underestimated due to data loss because of the high phase gradients and near-field decorrelation on the coseismic interferograms (Fig. 7).

The ability of InSAR to detect centimetric displacements was evident when comparing the post-seismic and coseismic interferograms. The rupture trace is conspicuous and the offset is observable in both the near-field and far-field of the post-seismic interferograms. Left lateral afterslip measurements are at least 4 to 14 cm with an average  $1\sigma$  uncertainty of 1.5 cm. Millimeter-scale offsets were not detected given the coarse 80 m pixel resolution and noise of the resultant interferograms.

Optical correlation is useful to assess the near-field region (Avouac et al., 2006) and larger displacements equivalent to at least one-tenth of the pixel resolution (Elliott et al., 2020b; Leprince et al., 2007). Results show that near-field displacements are resolved (Figs. 5 and 6) despite noticeable high noise levels. OIC, using PlanetScope data with 3.125 m pixel resolution, was able to detect left lateral offsets between 7 to 60 cm in the near-field, but with a larger 19.9 cm  $1\sigma$  average uncertainty than InSAR due to noise. Thus, careful interpretation of swath profiles is crucial to identify valid measurements because noise easily overprints tectonic signals. Noise signals are amplified by misaligned acquisition geometry, unequal solar illumination

angle, and insufficient geometric corrections (Elliott et al., 2020b; Stumpf et al., 2017). The representative 9 April 2020 - 2 April 2021 (satellite ID 1105) images had decent illumination and acquisition geometry agreement (Table 1) and support the validity of the measurements.

### 425 **5.1.2 Temporal Baseline**

The pair classes of PlanetScope orthoimages used in the study showed that images roughly one year apart resolved the rupture far better than those imaged months apart primarily due to the contrasting vegetative cover. Satellite images from the same seasons are critical for optical correlation because feature similarity helps the algorithm match features more accurately (Elliott et al., 2020a). Feature matching in agricultural and developing areas can be difficult in the subpixel domain given the difference  
430 in crop patterns and rapid anthropogenic development, leading to differing pixel intensity values in the pre-event and post-event images (Barnhart et al., 2011).

Images that are one day apart generate deformation fields with minimal seasonal and anthropogenic artifacts. However, this may have exceptions such that correlations for the 2016 Fukushima earthquake using one-day interval drone images resulted in noisy deformation fields with low correlation (Valkaniotis, per. comm., 2021). Single-day intervals were not tested in Masbate  
435 due to the lack of usable orthoimagery in terms of study area coverage, compatible acquisition parameters, and cloud cover.


DInSAR relies on coherence to resolve the deformation along the LOS direction, which is controlled by the similarity of point scatterers between the acquisitions. Good coherence in urban areas and dry deserts that remain constant throughout several years produce quality interferograms (Wang and Fialko, 2014; Wei and Sandwell, 2010), which indicate the capability of interferometry to function for longer periods. However, in other types of surface cover such as vegetation and snow, coherence  
440 has an inverse relationship with temporal baseline, which reduces the usable time interval (Kervyn, 2001). The six-day revisit of Sentinel-1 conveniently dictates the pre-event and post-event radar data pairing in areas affected by sub-optimal land cover. The baseline can be increased for L-band satellites, such as ALOS-2, for they can detect the bare ground irrespective of surface cover. Interferograms with severe unwrapping errors are manually excluded in the time-series analysis (Yunjun et al., 2019), effectively limiting the temporal baseline range to six to 12 days.

### 445 **5.2 Accuracy of Optical Correlation**

The slip distribution from optical correlation agrees with the offset feature measurements from the field survey (Fig. 12a). The weighted moving mean of the pixel offset measurements unifies the six valid pairs to compensate for data gaps that particular pairs are unable to resolve. The variance between the pixel offsets and field measures is typically at  $\sim 2$  cm with a maximum  $\sim 17$  cm. The larger difference is interpreted to be a local variation or measurement error since the specific point was taken on a  
450 repaired road section, which the remote sensing approach may have missed as it generally detects the overall deformation and is less sensitive to local slip.

Previous studies show varying degrees of misfit between field offsets and remote sensing data (Cheloni et al., 2014; Gold et al., 2021). These were attributed to diffuse displacement, secondary faulting, and shallow fault complexities. Our measurements, however, show consistency and minimal short wavelength variability. This may be a consequence of the two-year inter-





**Figure 12.** Combined slip distribution plots. A) Solid gray line shows the weighted moving mean of the six valid optical correlation pairs. Dashed red line represents the weighted moving mean of the field measurements. Blue line and markers represent the afterslip from InSAR placed in a secondary axis. B) Residual curves of the weighted moving mean pixel offsets and field measurements with InSAR to estimate the coseismic slip.

455 val between the earthquake event and the field survey as most surface traces have been modified by active surface processes. Furthermore, ruptures on unconsolidated overburden are easily erased due to active weathering and erosion, leaving only the most prominent traces to remain. Similarly, abundant vegetation, challenging surface conditions, and relatively small tectonic displacements compromise the clarity of the resolved rupture and efficacy of optical correlation. Nevertheless, OIC outputs indicate that the remote sensing results accurately capture ground displacements and provide greater along-strike coverage.

460 The extent of detectable rupture is another fundamental difference between optical correlation outputs and field measurements. The northern rupture terminations are situated 4.6 km and 3.2 km from Naro Bay for optical correlation and field survey, respectively. To the south, OIC measurements cease at the shore of Cataingan Bay since no slip measurements are attainable on the opposite side of the water body, where the field survey revealed more ruptures as the fault resurfaced therein.

Neither method can resolve slip on submerged fault segments, requiring the use of complementary marine survey techniques  
465 and datasets that cover the ocean floor.

### 5.3 Surface Rupture Kinematics

#### 5.3.1 Constraining the Coseismic Slip

The temporal range of the pixel offset rasters and field investigation date implies the presence of both coseismic and post-  
seismic slip components in the optical correlation measurements and field measurements. To estimate the peak coseismic  
470 offsets, we subtracted the largest afterslip value from the peak pixel offsets and field measures. The average values were  
derived from the mean of residuals of the moving mean curves of the two total slip measures with InSAR (Fig. 12b). From  
this, the estimated peak coseismic slip amount to  $46.3 \pm 10.7$  cm from optical correlation and  $45.7 \pm 15.9$  cm from the field  
measure residuals. The average values are 27.7 cm and 24.3 cm from pixel offsets and field measures, respectively.

#### 5.3.2 Coseismic Offsets

475 The amount of coseismic slip is a key factor in the assessment of seismic moment, rupture mechanism, and degree of seismic  
hazard. The results of this study show a unimodal curve with a characteristic long wavelength and isolated short wavelength  
variability for the regional slip distribution (Fig. 12). The pattern suggests a low generalized resolution due to temporally-  
influenced smoothing and noise. In addition, the long wavelength trend can also indicate fault maturity (Allam et al., 2019),  
which is a credible interpretation for the Masbate segment of the PF given its Middle Miocene initiation and rate of activity  
480 (Fitch, 1972; Tsutsumi and Perez, 2013; Pinet and Stephan, 1990). Moreover, the known fault trace in Masbate is morphologi-  
cally linear, supporting the general maturity estimate (Manighetti et al., 2021).

Slip distributions can give clues into the fault complexity and mechanical properties of the fault plane (Allam et al., 2019;  
Bruhat et al., 2020; Perrin et al., 2016). This suggests that the observed unimodal slip distribution with a singular peak implies  
a relatively simple fault zone at depth, subject to validation through geodetic inversion. Such peaks usually define the location  
485 of the ruptured asperity (Kaneda et al., 2008) where the highest stress drop occurs due to peak moment release from the  
accumulated potential energy (Freymueller et al., 1994).

Our seismotectonic interpretations (Section 5.4) imply a variable slip distribution on the submerged sections of the Masbate  
segment of the PF since it represents the primary depression of the identified transtensional basin. This inferred slip trend may  
portray a complex shallow fault structure in this section as the slip is distributed on multiple fault segments (Avouac et al.,  
490 2006; Allam et al., 2019; Wang et al., 2021). Moreover, recent marine geophysical surveys in the vicinity of Cataingan Bay  
(Llamas and Marfito, 2022) revealed the geometric complexity of the fault, which supports our interpretation.

Offset values gradually decay to zero towards the northwest. The smooth slip trend, devoid of short wavelength variation,  
suggests a systematic rupture (Treiman, 2002) on a relatively simple sub-vertical fault plane (Allam et al., 2019; Chen et al.,  
2021). Tapering and deepening aftershock pattern along with the focal mechanism solutions could support this observation.

Our interferometry results denote a 14 cm largest afterslip, which accounts for at least 30 % of the estimated peak coseismic displacement over the rapid aftershock decay interval. This lower limit estimate is a consequence of the six-day revisit cycle of Sentinel-1, resulting in incomplete data since the first radar image in the time-series stack was from 20 August 2020 and does not include the earliest post-seismic deformation immediately after the 18 August 2020 mainshock.

500 The distribution of afterslip (Fig. 12) along the fault demonstrates a general slip increase southwards, which can be divided into two distinct sections. The northern ~15 km section is characterized by short-wavelength variations and average left lateral offsets of 5-6 cm, whereas the southern section includes the peak afterslip without distinctive variability. The northern trend may indicate the occurrence of diffuse afterslip on a wide zone, which may or may not have breached the surface. Chen et al. (2021) suggested an increased relative friction such that the released post-seismic moment is typically less compared to the  
505 mainshock, resulting in heterogeneous loading. However, we cannot exclude the possibility that variability may stem from surface complexities or data processing artifacts, as indicated by the observed unwrapping errors and the lack of data from the earliest post-seismic period.

Post-seismic deformation usually occurs outside the regions of peak coseismic slip as the shear stress is transferred by the mainshock and causes stress loading at the locations of incomplete rupture and unruptured sections (Johanson, 2006;  
510 Bürgmann et al., 2021). Our results reflect the coincidence of the post-seismic and coseismic slip distributions on the surface, whose maxima are aligned and equidistant from the centroid, accompanied by a gradual slip recession northwards (Fig. 12). This is similar to the 1999 Izmit (Reilinger et al., 2000), 1996 Biak (Das and Henry, 2003), 2011 Maduo (Wang et al., 2019), and 2014 South Napa (DeLong et al., 2016) earthquakes, which also exhibit spatially overlapping coseismic and post-seismic slip distributions. We interpret this overlap of the slip distributions to indicate a vertical stress migration along the fault surface  
515 to either deeper or shallower sections of the fault.

Aseismic afterslip, viscoelastic relaxation, and poroelastic rebound are the common mechanisms of post-seismic deformation (He et al., 2021; Tomita et al., 2020). The existence of resolvable afterslip is a clear indication of post-seismic activity as the stress is released by the mainshock. Our observed coincidence of the coseismic and post-seismic slip distributions probably indicates a localized stress transfer to the adjacent velocity strengthening vicinity of the primary asperity in accordance with the  
520 rate-and-state friction model (Marone et al., 1991). This would suggest that the post-seismic deformation is likely stress-driven (He et al., 2021) and operated as aseismic afterslip (Johanson, 2006) to accommodate the excess coseismic stress. This also agrees with the hypothesized vertical migration of stress along the fault plane.

We recognize that the data is limited to surface measurements and more information is required to accurately describe the occurrence of the afterslip along the fault plane. Determining the mechanics of the superposed slip character of the 2020  
525 Masbate earthquake event is of interest, and warrants investigation into potential factors such as non-steady state friction (Helmstetter and Shaw, 2009), residual stress heterogeneity (Wang et al., 2019), stress reorganization, or shallow material redistribution (He et al., 2021). Additionally, the contribution of known interseismic creep associated with the Masbate segment to the slip distributions merits further investigation.

### 5.3.4 Surface Rupture Length

530 Surface rupture length estimates from remote sensing data were assumed to be symmetric about the peak displacement zone (Table 3). The ground displacement distributions from optical correlation and field survey (Fig. 12) imply that the northern rupture terminus is located approximately 3.2 km from the northern shoreline, with peak slip values occurring between 17.3 to 23.5 km, resulting in rupture length estimates ranging from 28.2 to 41 km. The absence of offset measurements in the submerged portions of the fault necessitated the use of interpolation. Given the asymmetrical nature of strike-slip distributions, 535 which can take on either elliptical or triangular forms (Perrin et al., 2016), caution is critical when interpreting the interpolated values. The rupture lengths are also measured from the total length of the field survey and its extension towards the stepover in Cataingan Bay (Llamas and Marfito, 2022) resulting in 25 and 35.6 km rupture lengths, respectively.

### 5.3.5 Seismic Moment Estimates

Moment magnitude ( $M_w$ ) and seismic moment ( $M_o$ ) estimates for the 2020 Masbate event were calculated from the surface rupture parameters (Table 3).  $M_w$  calculations used the corresponding empirical equations from regression of rupture parameters and moment magnitudes (Wells and Coppersmith, 1994). The  $M_o$  was derived as a function of  $M_w$  from scaling relationships (Hanks and Kanamori, 1979). This approach was preferred over directly calculating the  $M_o$  from the rupture area and material rigidity due to the lack of reliable estimates of rupture width and subsurface fault rupture parameters. 540

The  $M_w$  estimates from the maximum and average displacements range between  $M_w$ 6.5 to  $M_w$ 6.7. The estimated maximum and average coseismic offsets all equate to  $M_w$ 6.5. The values closely fit the instrumental 6.6 moment magnitude. All rupture length estimates returned high  $M_w$  values between  $M_w$ 6.7 to  $M_w$ 7.0 which agrees with the 2003 earthquake observations, wherein the fault rupture is longer than expected relative to the magnitude. However, a downward counterfactual approach (Woo and Mignan, 2018; Woo, 2019) given the fault uncertainties at depth, suggests the probability of the larger  $M_w$  estimates. Nevertheless, the estimated coseismic displacements reasonably fit with the instrumental seismic moment of the mainshock. 545 The calculated  $M_w$ 6.5 value from the estimated coseismic slip parameter is consistent with a seismic moment of  $8.4 \times 10^{18}$  N·m.

The lower limit of the post-seismic moment was assessed using the largest and average afterslip values (Table 3). The largest afterslip, which measured 0.14 m, corresponds to  $M_w$ 6.2, while the average afterslip of 0.09 m is consistent with  $M_w$ 6.1. By comparing the largest afterslip value ( $M_o = 2.1 \times 10^{18}$  N·m) with the peak coseismic moment, we estimate that the afterslip 555 resulted in the release of energy equal to at least 25 % of the mainshock.

**Table 3.** Measured rupture parameters with corresponding moment magnitude (Wells and Coppersmith, 1994) and seismic moment (Hanks and Kanamori, 1979).

Parameter	Value	$M_w$	Mo (N·m)
Surface rupture length	34.4 km <sup>1</sup>	6.9	$2.6 \times 10^{19}$
	28.2 km <sup>2</sup>	6.8	$1.9 \times 10^{19}$
	41.0 km <sup>3</sup>	7.0	$3.5 \times 10^{19}$
	39.5 km <sup>4</sup>	6.9	$3.3 \times 10^{19}$
	25.0 km <sup>5</sup>	6.7	$1.5 \times 10^{19}$
	35.6 km <sup>6</sup>	6.9	$2.8 \times 10^{19}$
Maximum displacement	0.6 m <sup>7</sup>	6.6	$1.2 \times 10^{19}$
	0.6 m <sup>8</sup>	6.6	$1.1 \times 10^{19}$
Average displacement	0.4 m <sup>9</sup>	6.7	$1.2 \times 10^{19}$
	0.3 m <sup>10</sup>	6.5	$7.3 \times 10^{18}$
Largest afterslip	0.14 m <sup>11</sup>	6.2	$2.1 \times 10^{18}$
Average afterslip	0.09 m <sup>12</sup>	6.1	$1.8 \times 10^{18}$
Est. peak coseismic offset	0.5 m <sup>13</sup>	6.5	$8.4 \times 10^{18}$
	0.4 m <sup>14</sup>	6.5	$8.3 \times 10^{18}$
Est. average coseismic offset	0.3 m <sup>15</sup>	6.5	$8.2 \times 10^{18}$
	0.2 m <sup>16</sup>	6.5	$6.9 \times 10^{18}$

<sup>1</sup> symmetrical about the midpoint of the zone of peak displacement

<sup>2</sup> symmetrical about the lower limit of the zone of peak displacement

<sup>3</sup> symmetrical about the upper limit of the zone of peak displacement

<sup>4</sup> symmetrical about the peak displacement

<sup>5</sup> cumulative rupture length from the field survey

<sup>6</sup> cumulative rupture length from the field survey extended to the onshore stepover identified by Llamas and Marfito (2022)

<sup>7</sup> from optical correlation

<sup>8</sup> from field survey

<sup>9</sup> from the weighted moving mean curve of optical correlation outputs

<sup>10</sup> from the weighted moving mean curve of field measurements

<sup>11</sup> highest individual measurement from InSAR


<sup>12</sup> from the weighted moving mean curve of InSAR

<sup>13</sup> peak optical correlation value minus peak InSAR value

<sup>14</sup> peak field measurement minus peak InSAR value

<sup>15</sup> average of the residuals between the weighted moving mean curves of optical correlation and InSAR

<sup>16</sup> average of the residuals between the weighted moving mean curves of field measurements and InSAR



**Figure 13.** Morphology of the transtensional basins linked by right-stepping shear zone. Solid line indicates the regional fault. Lines with hachures indicate the basin sidewall boundary.

## 5.4 Seismotectonic Implications

### 5.4.1 Surface Rupture Morphology

The fault geomorphology (Fig. 10c) and geodetic observations (Figs. 5 and 8) highlight the linearity of the Masbate segment. However, the fault complexity (Tsunami and Perez, 2013) around 12°07'N was not clearly defined due to unwrapping errors in interferometry and high-amplitude noise in optical correlations.

The ruptures in Palanas form a right-stepping Riedel shear zone, with the ruptures occurring at a low angle with respect to the regional fault strike. Furthermore, a possible new splay was observed to the south, which potentially represents the primary fault plane, cutting across the bifurcating trace and traversing Cataingan Bay. Antithetic shears were observed at the bifurcation point and along the cross-cutting splay.

Topography and bedrock geology may also control the rupture (Kaneda et al., 2008), such that the southwestern rupture termination is observed on an alluvial plain, succeeded by rugged topography northwards. The fault complexity in Palanas and the bifurcation point in Cataingan coincide with the stratigraphic contacts of the Late Oligocene Nabangig and Miocene-Pliocene Buyag Formations. The previously identified fault complexities in this area may reflect the rheological contrast between the clastic limestones of the Nabangig Formation and conglomeratic Buyag Formation.

### 5.4.2 Developing Transtensional Basins

The morphology of mainland Masbate following the 2020 event (Fig. 13) was the basis for assigning a late relative development stage in a modelled transtensional basin development sequence (Wu et al., 2009). The basin boundary is characterized by

floodplains surrounded by topographic highs. Apparent asymmetric depocenters were recognized, with the shallower depocenter occurring on the floodplain and the nearshore section of Cataingan Bay, while the second one extends further southwest.

575 A separate transtensional basin is discernible in Dimasalang given the topography, rupture occurrence, and extensional structures in north Masbate (Bischke et al., 1990). The northern and southern basins appear to be linked by the Riedel shear zone in Palanas. Fault complexities in the principal deformation zone of a pull-apart basin (Wu et al., 2009) impede rupture propagation (Biasi and Wesnousky, 2017) in line with the implications of the observed decreasing coseismic and post-seismic slip towards the north. However, the scarcity of observed ruptures inhibits further analysis of the northern transtensional basin.

580 Fitting the development stage of the southern basin is constrained by the absence of information regarding local strain partitioning. Nonetheless, the expressed fault maturity and identified negative flower structures (Bischke et al., 1990; Llamas and Marfito, 2022) in the northern and southern bays highlight the presence of oblique extensional stress bounding the Masbate mainland. The orientation of the regional fault implies a WNW-ESE oriented  $\sigma_1$ . However, real-world conditions are more complex compared to the simplified parameters in analog models.

## 585 6 Conclusions

The necessity of coupling optical correlation and interferometry is demonstrated by investigating the surface rupture of the 2020  $M_w$ 6.6 Masbate earthquake along the Masbate segment of the Philippine Fault. Seismicity data showed that foreshocks began 32 days prior to the mainshock on 18 August 2020. The first 30 days of subsequent aftershocks are characterized by a rapid decrease in earthquake frequency, followed by a transition to a gradual decay until 31 January 2021.

590 Optical correlation and InSAR are directly compared through the corresponding generated offset maps. Measuring the ground offset from the near-field region of the interferogram is hindered by decorrelation, whereas the far-field suffers because of the submerged northeastern block. The optical correlation method revealed an average  $37.7 \pm 10$  cm displacement and  $60.6 \pm 9.8$  cm peak left lateral offset in the near-field of the onshore fault segments. Estimates of the peak left lateral coseismic slip from the total slip and afterslip are consistent with a geodetic moment magnitude of  $M_w$ 6.5 ( $8.4 \times 10^{18}$  N·m).

595 The capacity of InSAR to detect smaller displacement amplitudes is highlighted by the post-seismic interferograms. The SBAS time-series stack outlined at least  $14.3 \pm 0.9$  cm largest afterslip and  $8.8 \pm 0.7$  cm average left lateral afterslip. The measurements correspond to a  $M_w$ 6.2 ( $2.1 \times 10^{18}$  N·m) energy release equivalent to at least 25 % of the coseismic moment estimate. The resolved rupture from interferometry is clearer relative to optical correlation. Multimodal rupture length estimates translate to overestimated moment magnitude values. Hence, the Masbate segment is characterized as capable of producing  
600 significant slip and ruptures that are longer than expected, despite the short interval and continuous aseismic stress release during interseismic periods.

Our optical correlation data agrees with the laterally offset features from field investigation, highlighting the capacity of the method to accurately capture ground displacements despite the difficult tropical terrain. We view this as a consequence of the time interval between the 2020 Masbate earthquake and the 2022 field investigation. Weathering and erosion of fine ruptures  
605 had transpired, possibly influencing the field offset distribution.

The coseismic slip distribution shows the presence of a single asperity adjacent to the GlobalCMT centroid. In conjunction, the surface distribution of the afterslip coincides with the coseismic slip distribution. Based on the stress transfer theorem and the rate-and-state friction model, we infer that the involved post-seismic deformation is fundamentally stress-driven and migrated vertically in either the downdip or updip direction along the sub-vertical fault plane.

610 The surface rupture of the 2020 Masbate event cut across the extant bifurcating trace of the Philippine Fault on the island, possibly reflecting the primary fault plane. Comparing the rupture morphology with analog models reveals the presence of two transtensional basins in Cataingan and Dimasalang, located in the south and north, respectively. The low-angle orientation of ruptures in the midsection corresponds to a Riedel shear zone, which is interpreted to link the identified transtensional basins. Recognition of this structure adds valuable insights into zones vulnerable to future deformation, informing targeted land-use  
615 planning and mitigation strategies.

The findings and the methods we used for this study in Masbate lead us and future scientists to further analyze the geometry, kinematics, mechanics, and dynamics of fault movement in the area for improved seismic hazard mitigation. Specifically, the comprehensive slip data and observed unusually long rupture lengths along the fault measured through the optical correlation method agree with previous characterizations by Besana and Ando (2005), indicating significant movement despite the pres-  
620 ence of creep. This challenges current seismic hazard models of Masbate and underscores the need to re-evaluate potential earthquake magnitudes and ground shaking intensities in the region.

Our use of optical correlation alongside interferometry provides high-resolution surface displacement measurements that can aid in the assessment of earthquake hazards to assist the development of mitigation strategies. We recommend the widespread adoption of these methods in studying other active faults in the Philippines. Future studies could incorporate other datasets  
625 such as UAVs and explore the method for other surface processes. By reevaluating existing hazard models, refinement of the geological characterization, and scrutinizing slip estimates, we can gain a deeper understanding of the region's seismotectonic configuration and enhance earthquake preparedness in the region.

*Code availability.* MicMac is available from [github.com/micmacIGN/micmac](https://github.com/micmacIGN/micmac). StackProf is available from [github.com/micmacIGN/stackprof](https://github.com/micmacIGN/stackprof). MintPy is available from [github.com/insarlab/MintPy](https://github.com/insarlab/MintPy).

630 *Data availability.* Sentinel-2 datasets are accessed from ESA/EC Copernicus Sentinels Scientific Data Hub and various repositories. Dataset in this study are retrieved from [peps.cnes.fr](https://peps.cnes.fr). Sentinel-1A/B radar data and cloud interferogram processing is available at [search.asf.alaska.edu](https://search.asf.alaska.edu). Planet Labs data are not openly accessible, but academic/scientific access can be requested. Outputs are rendered with Scientific Colour Maps (Cramer, 2021) available from [fabiocrameri.ch/colourmaps](https://fabiocrameri.ch/colourmaps).

*Author contributions.* **Khelly Shan C. Sta. Rita:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project  
635 administration, Resources, Software, Validation, Visualization, Writing - original draft, Writing - review & editing. **Sotiris N. Valkaniotis:**



Formal analysis, Resources, Software, Supervision, Methodology, Writing - review & editing. **Alfredo Mahar Francisco A. Lagmay:**  
Conceptualization, Funding acquisition, Resources, Supervision, Writing - review & editing.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We extend our wholehearted appreciation to Dr. Arturo Daag for his suggestions and comments. We would like to thank  
640 the National Mapping and Resource Information Authority for providing the elevation data. We would also like to thank the Philippine  
Institute of Volcanology and Seismology as well as the Disaster Risk Management office of Dimasalang, Masbate for their contributions to  
the field investigation. We are also grateful for the comments of Dr. Austin Elliot, Dr. Gordon Woo, and the anonymous reviewer for their  
input and contributions towards improving this work.

## References

- 645 Allam, A. A., Kroll, K. A., Milliner, C. W. D., and Richards-Dinger, K. B.: Effects of fault roughness on coseismic slip and earthquake locations, *Journal of Geophysical Research: Solid Earth*, 124, 11 336–11 349, <https://doi.org/10.1029/2018JB016216>, 2019.
- Allen, C. R.: Circum-Pacific faulting in the Philippines-Taiwan Region, *Journal of Geophysical Research (1896-1977)*, 67, 4795–4812, <https://doi.org/https://doi.org/10.1029/JZ067i012p04795>, 1962.
- Aurelio, M.: Tectonique du segment central de la faille philippine : etude structurale, cinématique et evolution geodynamique, Ph.D. thesis, 650 <http://www.theses.fr/1992PA066699>, thèse de doctorat dirigée par Rangin, Claude Terre, océan, espace Paris 6 1992, 1992.
- Aurelio, M.: Displacement rates and block rotation in and around the Philippines-results from GEODYSSSEA data Part II, Program and Abstracts : GEOSEA 98-Ninth Regional Congress on Geology Mineral and Energy Resources of Southeast Asia, 238, <https://cir.nii.ac.jp/crid/1570572700614632960>, 1998.
- Aurelio, M.: Tectonics of the Philippines revisited, *Journal of the Geological Society of the Philippines*, 55, 119–183, 2000a.
- 655 Aurelio, M.: Shear partitioning in the Philippines: Constraints from Philippine Fault and global positioning system data, *Island Arc*, 9, 584 – 597, <https://doi.org/10.1111/j.1440-1738.2000.00304.x>, 2000b.
- Aurelio, M.: Masbate’s three-legged coconut tree spurs scientific interest, <https://coverstory.ph/masbates-three-legged-coconut-tree-spurs-scientific-interest/>, 2022.
- Aurelio, M. and Peña, R.: *Geology of the Philippines*, 2nd Edition (2010), Mines and Geosciences Bureau, 2010.
- 660 Aurelio, M., Barrier, E., Rangin, C., and Müller, C.: The Philippine Fault in the Late Cenozoic tectonic evolution of the Bondoc-Masbate-N. Leyte area, Central Philippines, *Journal of Southeast Asian Earth Sciences*, 6, 221–238, [https://doi.org/10.1016/0743-9547\(91\)90069-a](https://doi.org/10.1016/0743-9547(91)90069-a), 1991.
- Aurelio, M., Barrier, E., Gaulon, R., and Rangin, C.: Deformation and stress states along the central segment of the Philippine Fault: implications to wrench fault tectonics, *Journal of Asian Earth Sciences*, 15, 107–119, [https://doi.org/10.1016/s0743-9547\(97\)00001-9](https://doi.org/10.1016/s0743-9547(97)00001-9), 665 1997.
- Aurelio, M., Lagmay, M., Escudero, J. A., and Catugas, S.: Latest Philippine earthquake reveals tectonic complexity, *Temblor*, <https://doi.org/10.32858/temblor.191>, 2021.
- Avouac, J.-P., Ayoub, F., Leprince, S., Konca, O., and Helmberger, D. V.: The 2005, Mw 7.6 Kashmir earthquake: Sub-pixel correlation of ASTER images and seismic waveforms analysis, *Earth and Planetary Science Letters*, 249, 514–528, 670 <https://doi.org/10.1016/j.epsl.2006.06.025>, 2006.
- Bacolcol, T. C.: Etude géodésique de la faille Philippine dans les Visayas, Ph.D. thesis, Université Pierre et Marie Curie, <http://www.theses.fr/2003PA066404>, thèse de doctorat dirigée par Barrier, Éric Géologie Paris 6 2003, 2003.
- Bacolcol, T. C., Aguilar, A., Jorgio, R., and de la Cruz, R.: Interseismic Deformation and Horizontal Displacements Associated with the February 15, 2003 Masbate Earthquake, Tech. rep., Philippine Institute of Volcanology and Seismology, Philippines, 2005.
- 675 Barnhart, W. D., Willis, M. J., Lohman, R. B., and Melkonian, A. K.: Insar and optical constraints on fault slip during the 2010-2011 new zealand earthquake sequence, *Seismological Research Letters*, 82, 815–823, <https://doi.org/10.1785/gssrl.82.6.815>, 2011.
- Barrier, E., Huchon, P., and Aurelio, M.: Philippine fault: A key for Philippine kinematics, *Geology*, 19, 32, [https://doi.org/10.1130/0091-7613\(1991\)019&#60;0032:pfakfp&#62;2.3.co;2](https://doi.org/10.1130/0091-7613(1991)019&#60;0032:pfakfp&#62;2.3.co;2), 1991.
- Bautista, M. L. P. and Oike, K.: Estimation of the magnitudes and epicenters of Philippine historical earthquakes, *Tectonophysics*, 317, 680 137–169, [https://doi.org/10.1016/s0040-1951\(99\)00272-3](https://doi.org/10.1016/s0040-1951(99)00272-3), 2000.

- Besana, G. M. and Ando, M.: The central Philippine Fault Zone: Location of great earthquakes, slow events, and creep activity, *Earth Planets Space*, 57, 987–994, <https://doi.org/10.1186/BF03351877>, 2005.
- Biasi, G. and Wesnousky, S.: Bends and Ends of Surface Ruptures, *Bulletin of the Seismological Society of America*, 107, 2543–2560, <https://doi.org/10.1785/0120160292>, 2017.
- 685 Bird, P.: An updated digital model of plate boundaries, *Geochemistry, Geophysics, Geosystems*, 4, <https://doi.org/https://doi.org/10.1029/2001GC000252>, 2003.
- Bischke, R., Suppe, J., and Del Pilar, R.: Implications of a newly discovered branch of the Philippine fault system, in: *International Symposium on the Geodynamic Evolution of Eastern Eurasian Margin*, 1988.
- Bischke, R., Suppe, J., and del Pilar, R.: A new branch of the Philippine fault system as observed from aeromagnetic and seismic  
690 data, *Tectonophysics*, 183, 243–264, [https://doi.org/https://doi.org/10.1016/0040-1951\(90\)90419-9](https://doi.org/https://doi.org/10.1016/0040-1951(90)90419-9), geodynamic evolution of the Eastern Eurasian Margin, 1990.
- Bruhat, L., Klinger, Y., Vallage, A., and Dunham, E. M.: Influence of fault roughness on surface displacement: from numerical simulations to coseismic slip distributions, *Geophysical Journal International*, 220, 1857–1877, <https://doi.org/10.1093/gji/ggz545>, 2020.
- Bürgmann, R., Fielding, E., Liu, Z., and Wang, K.: Postseismic deformation due to the 2019 Ridgecrest earthquakes and its impact on the  
695 CGM, Tech. rep., [https://files.sceec.org/s3fs-public/reports/2020/20016\\_report.pdf?CUhbHuJtXfv9IPyOGtLr.ZvyIuNq91Fg](https://files.sceec.org/s3fs-public/reports/2020/20016_report.pdf?CUhbHuJtXfv9IPyOGtLr.ZvyIuNq91Fg), 2021.
- Canizares, C., Motagh, M., and Haghshenas Haghghi, M.: Resolving 3D coseismic deformation of the 2019 Mw 7.1 Ridgecrest earthquake using radar and optical data, in: *EGU General Assembly Conference Abstracts*, p. 13659, 2020.
- Chamot-Rooke, N. and Le Pichon, X.: GPS determined eastward Sundaland motion with respect to Eurasia confirmed by earthquakes slip vectors at Sunda and Philippine trenches, *Earth and Planetary Science Letters*, 173, 439–455,  
700 [https://doi.org/https://doi.org/10.1016/S0012-821X\(99\)00239-3](https://doi.org/https://doi.org/10.1016/S0012-821X(99)00239-3), 1999.
- Cheloni, D., Giuliani, R., D’Anastasio, E., Atzori, S., Walters, R., Bonci, L., D’Agostino, N., Mattone, M., Calcaterra, S., Gambino, P., Deninno, F., Maseroli, R., and Stefanelli, G.: Coseismic and post-seismic slip of the 2009 L’Aquila (Central Italy) MW 6.3 earthquake and implications for seismic potential along the Campotosto fault from joint inversion of high-precision levelling, InSAR and GPS data, *Tectonophysics*, 622, 168–185, <https://doi.org/10.1016/j.tecto.2014.03.009>, 2014.
- 705 Chen, H., Qu, C., Zhao, D., Ma, C., and Shan, X.: Rupture kinematics and coseismic slip model of the 2021 mw 7.3 Maduo (China) earthquake: implications for the seismic hazard of the Kunlun fault, *Remote Sensing*, 13, 3327, <https://doi.org/10.3390/rs13163327>, 2021.
- Crameri, F.: Scientific colour maps, <https://doi.org/10.5281/zenodo.5501399>, 2021.
- Das, S. and Henry, C.: Spatial relation between main earthquake slip and its aftershock distribution: AFTERSHOCK DISTRIBUTIONS, *Reviews of Geophysics*, 41, <https://doi.org/10.1029/2002RG000119>, 2003.
- 710 DeLong, S. B., Donnellan, A., Ponti, D. J., Rubin, R. S., Lienkaemper, J. J., Prentice, C. S., Dawson, T. E., Seitz, G., Schwartz, D. P., Hudnut, K. W., Rosa, C., Pickering, A., and Parker, J. W.: Tearing the terroir: Details and implications of surface rupture and deformation from the 24 August 2014 M6.0 South Napa earthquake, California, *Earth and Space Science*, 3, 416–430, <https://doi.org/https://doi.org/10.1002/2016EA000176>, 2016.
- Delorme, A.: StackProf, <https://github.com/micmacIGN/stackprof>, 2021.
- 715 Dianala, J. D. B., Jolivet, R., Thomas, M. Y., Fukushima, Y., Parsons, B., and Walker, R.: The Relationship Between Seismic and Aseismic Slip on the Philippine Fault on Leyte Island: Bayesian Modeling of Fault Slip and Geothermal Subsidence, *Journal of Geophysical Research: Solid Earth*, 125, e2020JB020052, <https://doi.org/https://doi.org/10.1029/2020JB020052>, e2020JB020052  
[10.1029/2020JB020052](https://doi.org/10.1029/2020JB020052), 2020.

- Duquesnoy, T., Barrier, E., Kasser, M., Aurelio, M., Gaulon, R., Punongbayan, R. S., and Rangin, C.: Detection of creep along the Philippine Fault: First results of geodetic measurements on Leyte Island, central Philippine, *Geophysical Research Letters*, 21, 975–978, <https://doi.org/https://doi.org/10.1029/94GL00640>, 1994.
- 720 Elliott, A., Elliott, J., Hollingsworth, J., Kulikova, G., Parsons, B., and Walker, R.: Satellite imaging of the 2015 M7.2 earthquake in the Central Pamir, Tajikistan, elucidates a sequence of shallow strike-slip ruptures of the Sarez-Karakul fault, *Geophysical Journal International*, 221, 1696–1718, <https://doi.org/10.1093/gji/ggaa090>, 2020a.
- 725 Elliott, J., De Michele, M., and Gupta, H. K.: Earth observation for crustal tectonics and earthquake hazards, *Surveys in Geophysics*, 41, 1355–1389, <https://doi.org/10.1007/s10712-020-09608-2>, 2020b.
- Farrell, J., Husen, S., and Smith, R. B.: Earthquake swarm and b-value characterization of the Yellowstone volcano-tectonic system, *Journal of Volcanology and Geothermal Research*, 188, 260–276, <https://doi.org/https://doi.org/10.1016/j.jvolgeores.2009.08.008>, the Track of the Yellowstone Hotspot, 2009.
- 730 Fitch, T. J.: Plate convergence, transcurrent faults, and internal deformation adjacent to Southeast Asia and the western Pacific, *Journal of Geophysical Research*, 77, 4432–4460, <https://doi.org/10.1029/jb077i023p04432>, 1972.
- Frey Mueller, J., King, N. E., and Segall, P.: The co-seismic slip distribution of the Landers earthquake, *Bulletin of the Seismological Society of America*, 84, 646–659, <https://doi.org/10.1785/BSSA0840030646>, 1994.
- GDAL/OGR contributors: GDAL/OGR Geospatial Data Abstraction software Library, <https://gdal.org>, 2021.
- 735 Gold, R. D., DuRoss, C. B., and Barnhart, W. D.: Coseismic surface displacement in the 2019 ridgecrest earthquakes: comparison of field measurements and optical image correlation results, *Geochemistry, Geophysics, Geosystems*, 22, <https://doi.org/10.1029/2020GC009326>, 2021.
- Hanks, T. C. and Kanamori, H.: A moment magnitude scale, *Journal of Geophysical Research: Solid Earth*, 84, 2348–2350, <https://doi.org/https://doi.org/10.1029/JB084iB05p02348>, 1979.
- 740 He, L., Feng, G., Wu, X., Lu, H., Xu, W., Wang, Y., Liu, J., Hu, J., and Li, Z.: Coseismic and early postseismic slip models of the 2021 mw 7.4 maduo earthquake (Western china) estimated by space-based geodetic data, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2021GL095860>, 2021.
- Helmstetter, A. and Shaw, B. E.: Afterslip and aftershocks in the rate-and-state friction law, *Journal of Geophysical Research: Solid Earth*, 114, 2007JB005 077, <https://doi.org/10.1029/2007JB005077>, 2009.
- 745 Johanson, I. A.: Coseismic and postseismic slip of the 2004 parkfield earthquake from space-geodetic data, *Bulletin of the Seismological Society of America*, 96, S269–S282, <https://doi.org/10.1785/0120050818>, 2006.
- Kaneda, H., Nakata, T., Tsutsumi, H., Kondo, H., Sugito, N., Awata, Y., Akhtar, S. S., Majid, A., Khattak, W., Awan, A. A., Yeats, R. S., Hussain, A., Ashraf, M., Wesnousky, S. G., and Kausar, A. B.: Surface rupture of the 2005 kashmir, pakistan, earthquake and its active tectonic implications, *Bulletin of the Seismological Society of America*, 98, 521–557, <https://doi.org/10.1785/0120070073>, 2008.
- 750 Kennedy, J. H., Hogenson, K., Johnston, A., Kristenson, H., Lewandowski, A., Logan, T. A., Meyer, F. J., and Rine, J.: Get HyP3! SAR processing for everyone, Tech. Rep. EGU21-8973, Copernicus Meetings, <https://doi.org/10.5194/egusphere-egu21-8973>, 2021.
- Kervyn, F.: Modelling topography with SAR interferometry: illustrations of a favourable and less favourable environment, *Computers & Geosciences*, 27, 1039–1050, [https://doi.org/10.1016/S0098-3004\(00\)00158-8](https://doi.org/10.1016/S0098-3004(00)00158-8), 2001.
- Krambach, M.: swath profile, <https://plugins.qgis.org/plugins/swathProfile/>, 2015.

- 755 Lagmay, A. M., Tengonciang, A., and Uy, H.: Structural setting of the Bicol Basin and kinematic analysis of fractures on Mayon Volcano, Philippines, *Journal of Volcanology and Geothermal Research*, 144, 23–36, <https://doi.org/https://doi.org/10.1016/j.jvolgeores.2004.11.015>, the Tectonics and Physics of Volcanoes, 2005.
- Lai, L. C., Rau, R. J., Bacolcol, T., Ker, C. M., Sapla, G., Ching, K. E., and Solidum, Jr., R.: Earthquake Potential of the Creeping-to-locked Masbate Segment of the Philippine Fault Based on GPS Observations, in: *AGU Fall Meeting Abstracts*, vol. 2019, pp. G43B–0748, 2019.
- 760 Legaspi, C. J., Ortega, J. P., Reyes, P. R., and Riza, E. G.: Investigation of Stress Changes along the Philippine Fault Zone Using Coulomb Stress Transfer Modeling, in: *GEOCON 2018*, 2018.
- Leprince, S., Ayoub, F., Klingler, Y., and Avouac, J.-P.: Co-Registration of Optically Sensed Images and Correlation (COSI-Corr): an operational methodology for ground deformation measurements, in: *2007 IEEE International Geoscience and Remote Sensing Symposium*, p. 1943–1946, <https://doi.org/10.1109/IGARSS.2007.4423207>, 2007.
- 765 Lindsey, E. O., Fialko, Y., Bock, Y., Sandwell, D. T., and Bilham, R.: Localized and distributed creep along the southern San Andreas Fault, *Journal of Geophysical Research: Solid Earth*, 119, 7909–7922, <https://doi.org/https://doi.org/10.1002/2014JB011275>, 2014.
- Llamas, D. and Marfito, B.: Geometry and structure of the offshore segments of the Philippine Fault in Central Philippines, in: *GeoCon 2022: Disruptions, Transformations—Geosciences at the Forefront, Philippines*, 2022.
- Manalo, P. C., Dimalanta, C. B., Faustino-Eslava, D. V., Payot, B. D., Ramos, N. T., Queaño, K. L., Perez, A. D. C., and Yumul Jr., G. P.: Geochemical and Geophysical Characteristics of the Balud Ophiolitic Complex (BOC), Masbate Island, Philippines: Implications for its Generation, Evolution and Emplacement, *Terrestrial, Atmospheric and Oceanic Sciences*, 26, 687, [https://doi.org/10.3319/tao.2015.05.19.01\(tc\)](https://doi.org/10.3319/tao.2015.05.19.01(tc)), 2015.
- 770 Manighetti, I., Mercier, A., and De Barros, L.: Fault Trace Corrugation and Segmentation as a Measure of Fault Structural Maturity, *Geophysical Research Letters*, 48, e2021GL095372, <https://doi.org/https://doi.org/10.1029/2021GL095372>, e2021GL095372 2021GL095372, 2021.
- 775 Marone, C., Scholz, C., and Bilham, R.: On the Mechanics of Earthquake Afterslip, *Journal of Geophysical Research*, 96, 8441–8452, <https://doi.org/10.1029/91JB00275>, 1991.
- Michel, R., Avouac, J.-P., and Taboury, J.: Measuring ground displacements from SAR amplitude images: Application to the Landers Earthquake, *Geophysical Research Letters*, 26, 875–878, <https://doi.org/10.1029/1999g1900138>, 1999.
- 780 Miraballes, M., Mirana, R., Ng, J., Rejuso, A., and Villanueva, F.: Report on the Rapid Assessment on the effects of the Magnitude 6.6 Earthquake in the Municipalities of Cataingan and Pio V. Corpuz, Province of Masbate, Tech. rep., Mines and Geosciences Bureau - Region V, 2020.
- Mogi, K.: Some Discussions on Aftershocks, Foresocks and Earthquake Swarms : the Fracture of a Semi-infinite Body Caused by an Inner Stress Origin and Its Relation to the Earthquake Phenomena, *Bulletin of the Earthquake Research Institute, University of Tokyo*, 41, 615–658, <https://doi.org/https://doi.org/10.15083/0000033716>, 1963.
- 785 Morell, K. D., Styron, R., Stirling, M., Griffin, J., Archuleta, R., and Onur, T.: Seismic Hazard Analyses From Geologic and Geomorphic Data: Current and Future Challenges, *Tectonics*, 39, e2018TC005365, <https://doi.org/https://doi.org/10.1029/2018TC005365>, e2018TC005365 10.1029/2018TC005365, 2020.
- NDRRMC: Sitrep re Magnitude 6.6 Earthquake in Cataingan, Masbate, <https://ndrrmc.gov.ph/index.php/20-incidents-monitored/4096-sitrep-re-magnitude-6-6-earthquake-in-cataingan,-masbate.html>, 2020.
- 790

- Perrin, C., Manighetti, I., Ampuero, J.-P., Cappa, F., and Gaudemer, Y.: Location of largest earthquake slip and fast rupture controlled by along-strike change in fault structural maturity due to fault growth, *Journal of Geophysical Research: Solid Earth*, 121, 3666–3685, <https://doi.org/10.1002/2015JB012671>, 2016.
- PHIVOLCS: Historical Earthquakes in the Philippines, 1:5,000,000 map, 1999.
- 795 PHIVOLCS: Active Faults Map - Province of Masbate, <https://gisweb.phivolcs.dost.gov.ph/gisweb/earthquake-volcano-related-hazard-gis-information>, 2018.
- PHIVOLCS: 2020 Masbate Earthquake, <https://www.phivolcs.dost.gov.ph/index.php/2-uncategorised/10490-2020-masbate-earthquake>, 2020.
- Pinet, N. and Stephan, J. F.: The Philippine wrench fault system in the Ilocos Foothills, northwestern Luzon, Philippines, *Tectonophysics*, 183, 207–224, [https://doi.org/https://doi.org/10.1016/0040-1951\(90\)90417-7](https://doi.org/https://doi.org/10.1016/0040-1951(90)90417-7), geodynamic evolution of the Eastern Eurasian Margin, 1990.
- 800 Planet Team: Planet Application Program Interface: In *Space for Life on Earth*, <https://api.planet.com>, 2017–.
- Porth, H., Müller, C., and von Daniels, C. H.: The sedimentary formations of the Visayan Basin, Philippines, *Geol. Jahrbuch Reihe*, 1989.
- Rangin, C., Le Pichon, X., Mazzotti, S., Pubellier, M., Chamot-Rooke, N., Aurelio, M., Walpersdorf, A., and Quebral, R.: Plate convergence measured by GPS across the Sundaland/Philippine Sea Plate deformed boundary: the Philippines and eastern Indonesia, *Geophysical Journal International*, 139, 296–316, <https://doi.org/10.1046/j.1365-246X.1999.00969.x>, 1999.
- 805 Reilinger, R. E., Ergintav, S., Burgmann, R., McClusky, S., Lenk, O., Barka, A., Gurkan, O., Hearn, L., Feigl, K. L., Cakmak, R., Aktug, B., Ozener, H., and Toksoz, M. N.: Coseismic and postseismic fault slip for the 17 august 1999,  $m = 7.5$ , izmit, turkey earthquake, *Science*, 289, 1519–1524, <https://doi.org/10.1126/science.289.5484.1519>, 2000.
- Rosu, A.-M., Pierrot-Deseilligny, M., Delorme, A., Binet, R., and Klinger, Y.: Measurement of ground displacement from optical satellite image correlation using the free open-source software MicMac, *ISPRS Journal of Photogrammetry and Remote Sensing*, 100, 48–59, <https://doi.org/10.1016/j.isprsjprs.2014.03.002>, 2015.
- 810 Scheffler, D., Hollstein, A., Diedrich, H., Segl, K., and Hostert, P.: AROSICS: An Automated and Robust Open-Source Image Co-Registration Software for Multi-Sensor Satellite Data, *Remote Sensing*, 9, 676, <https://doi.org/10.3390/rs9070676>, 2017.
- Scholz, C.: Earthquakes and friction laws, *Nature*, 391, 37–42, <https://doi.org/10.1038/34097>, 1998.
- 815 SEASEE: Series on Seismology Volume IV Philippines, Government Printing Office, 1985.
- Seno, T.: The instantaneous rotation vector of the Philippine sea plate relative to the Eurasian plate, *Tectonophysics*, 42, 209–226, [https://doi.org/https://doi.org/10.1016/0040-1951\(77\)90168-8](https://doi.org/https://doi.org/10.1016/0040-1951(77)90168-8), 1977.
- Simborio, T., Yokoi, T., and Hara, T.: Strong ground motion simulation of the 2020 Masbate, Philippines earthquake (Mw6.6) using Empirical Green’s Function method, <https://iisee.kenken.go.jp/syndb/data/20230117f70f6a30.pdf>, synopsis of IISEE-GRIPS, 2022.
- 820 Stumpf, A., Malet, J.-P., and Delacourt, C.: Correlation of satellite image time-series for the detection and monitoring of slow-moving landslides, *Remote Sensing of Environment*, 189, 40–55, <https://doi.org/10.1016/j.rse.2016.11.007>, 2017.
- Tiongson, S. and Ramirez, R.: Mapping of ground surface deformations and its associated damage using SAR interferometry: a case study of the 2020 Masbate earthquake, *E3S Web Conf.*, 347, 03 014, <https://doi.org/10.1051/e3sconf/202234703014>, 2022.
- Tomita, F., Iinuma, T., Ohta, Y., Hino, R., Kido, M., and Uchida, N.: Improvement on spatial resolution of a coseismic slip distribution using postseismic geodetic data through a viscoelastic inversion, *Earth, Planets and Space*, 72, 84, <https://doi.org/10.1186/s40623-020-01207-0>, 2020.
- 825 Treiman, J. A.: Primary surface rupture associated with the mw 7. 1 16 october 1999 hector mine earthquake, san bernardino county, california, *Bulletin of the Seismological Society of America*, 92, 1171–1191, <https://doi.org/10.1785/0120000923>, 2002.

- Tronin, A.: Remote sensing and earthquakes: A review, *Physics and Chemistry of the Earth, Parts A/B/C*, 31, 138–142, <https://doi.org/10.1016/j.pce.2006.02.024>, 2006.
- 830 Tsutsumi, H. and Perez, J.: Along strike variation of seismic behavior of the Philippine fault based on historical- and paleo-seismicity, in: *AGU Fall Meeting Abstracts*, vol. 2011, pp. T43D–2366, 2011.
- Tsutsumi, H. and Perez, J.: Large-scale active fault map of the Philippine fault based on aerial photograph interpretation, *Active Fault Research*, 39, 29–37, [https://doi.org/10.11462/afr.2013.39\\_29](https://doi.org/10.11462/afr.2013.39_29), 2013.
- 835 Tsutsumi, H., Perez, J., and Lienkaemper, J.: Variation of Surface Creep Rate Along the Philippine Fault Based on Surveys of Alinement Arrays and Offset Cultural Features, in: *American Geophysical Union, Fall Meeting 2016*, 2016.
- Tymofeyeva, E. and Fialko, Y.: Geodetic Evidence for a Blind Fault Segment at the Southern End of the San Jacinto Fault Zone, *Journal of Geophysical Research: Solid Earth*, 123, 878–891, <https://doi.org/https://doi.org/10.1002/2017JB014477>, 2018.
- USGS: Advanced National Seismic System (ANSS) Comprehensive Catalog, <https://doi.org/10.5066/F7MS3QZH>, 2017.
- 840 Velasco, A. A., Ammon, C. J., Lay, T., and Hagerty, M.: Rupture process of the 1990 Luzon, Philippines (Mw = 7.7), earthquake, *Journal of Geophysical Research: Solid Earth*, 101, 22 419–22 434, <https://doi.org/https://doi.org/10.1029/96JB02290>, 1996.
- Wang, K. and Fialko, Y.: Space geodetic observations and models of postseismic deformation due to the 2005 M7.6 Kashmir (Pakistan) earthquake, *Journal of Geophysical Research: Solid Earth*, 119, 7306–7318, 2014.
- Wang, S., Xu, W., Xu, C., Yin, Z., Burgmann, R., Liu, L., and Jiang, G.: Changes in groundwater level possibly encourage shallow earthquakes in central australia: the 2016 petermann ranges earthquake, *Geophysical Research Letters*, 46, 3189–3198, <https://doi.org/10.1029/2018GL080510>, 2019.
- 845 Wang, S., Song, C., Li, S., and Li, X.: Resolving co- and early post-seismic slip variations of the 2021 MW 7.4 Maduo earthquake in east Bayan Har block with a block-wide distributed deformation mode from satellite synthetic aperture radar data, *Earth and Planetary Physics*, 6, <https://doi.org/10.26464/epp2022007>, 2021.
- 850 Wei, M. and Sandwell, D. T.: Decorrelation of L-Band and C-Band Interferometry Over Vegetated Areas in California, *IEEE Transactions on Geoscience and Remote Sensing*, 48, 2942–2952, <https://doi.org/10.1109/TGRS.2010.2043442>, 2010.
- Wells, D. and Coppersmith, K.: New Empirical Relationships among Magnitude, Rupture Length, Rupture Width, Rupture Area, and Surface Displacement, *Bulletin of the Seismological Society of America*, 84, 974–1002, 1994.
- Woo, G.: Downward Counterfactual Search for Extreme Events, *Frontiers in Earth Science*, 7, <https://doi.org/10.3389/feart.2019.00340>, 2019.
- 855 Woo, G. and Mignan, A.: Counterfactual Analysis of Runaway Earthquakes, *Seismological Research Letters*, 89, 2266–2273, <https://doi.org/10.1785/0220180138>, 2018.
- Wu, J. E., McClay, K., Whitehouse, P., and Dooley, T.: 4D analogue modelling of transtensional pull-apart basins, *Marine and Petroleum Geology*, 26, 1608–1623, <https://doi.org/https://doi.org/10.1016/j.marpetgeo.2008.06.007>, 2009.
- 860 Yen, J., Lu, C., Dorsey, R. J., Kuo-Chen, H., Chang, C., Wang, C., Chuang, R. Y., Kuo, Y., Chiu, C., Chang, Y., Bovenga, F., and Chang, W.: Insights into Seismogenic Deformation during the 2018 Hualien, Taiwan, Earthquake Sequence from InSAR, GPS, and Modeling, *Seismological Research Letters*, 90, 78–87, <https://doi.org/10.1785/0220180228>, 2018.
- Yunjun, Z., Fattahi, H., and Amelung, F.: Small baseline InSAR time series analysis: Unwrapping error correction and noise reduction, *Computers & Geosciences*, 133, 104 331, <https://doi.org/https://doi.org/10.1016/j.cageo.2019.104331>, 2019.