

Modeling the interannual variability of Maipo and Rapel river plumes off central Chile

Julio Salcedo-Castro^{1,2,3}, Antonio Olita⁴, Freddy Saavedra^{5,6,7}, Gonzalo S. Saldías^{8,9,10}, Raúl C. Cruz-Gómez¹¹, and Cristian D. De la Torre Martínez¹¹

¹School of Earth and Atmospheric Sciences, Faculty of Science. Queensland University of Technology. Brisbane, Australia

²Institute for Marine and Antarctic Studies, College of Sciences and Engineering, University of Tasmania, Hobart, Australia

³Sino-Australian Research Consortium for Coastal Management, School of Science, UNSW, Canberra, ACT, Australia

⁴National Research Council - Institute of Atmospheric Sciences and Climate, Cagliari, Italy

⁵Geografía, Facultad de Ciencias Naturales y Exactas, Universidad de Playa Ancha, Valparaíso, Chile

⁶Laboratorio de Teledetección Ambiental (TeleAmb), Universidad de Playa Ancha, Valparaíso, Chile

⁷HUB Ambiental, Universidad de Playa Ancha, Valparaíso, Chile

⁸Departamento de Física, Facultad de Ciencias, Universidad del Bío-Bío, Concepción, Chile

⁹Instituto Milenio en Socio-Ecología Costera (SECOS), Santiago, Chile

¹⁰Centro de Investigación Oceanográfica COPAS COASTAL, Universidad de Concepción, Chile

¹¹Departamento de Física, Universidad de Guadalajara. Blvd. Marcelino García Barragán y Calzada Olímpica C.P. 44840. Guadalajara, Jalisco, México

Correspondence: Julio Salcedo-Castro (julio.salcedocastro@qut.edu.au)

Abstract. River plumes have a direct influence on coastal environments, impacting coastal planktonic and benthic communities, including fishery resources. In general, the main drivers of river plume dynamics are the river discharge and the alongshore wind stress, whereas the tides and topography play a secondary role. In central Chile, rivers flowing into the eastern Pacific have a relatively short path on land, with a high slope and a mixed snow-rain regime. This study aims to understand the interannual variability of the plumes of the Maipo and Rapel rivers in the coastal/shelf area off central Chile and their influence on local ocean dynamics. We used the Coastal and Regional Ocean Community model (CROCO), with 1 km of horizontal resolution and 20 sigma levels, to simulate the ocean dynamics for the period 2003-2011. The results show that the plume's area coverage and coastal ocean salinity are strongly correlated with the river discharges. The predominant northeastward winds control the plumes orientation toward the northwest. However, episodes of southeastward winds in winter can reverse the plumes direction, promoting their attachment to the coast and southward transport. Results also show a salification trend linked to the severe droughts hitting central Chile during the studied period. This salification determines a change in local dynamics which could be more frequent in future scenarios of climate change with a significant lack of rain and river discharges along central Chile.

Copyright statement. TEXT

1 Introduction

15 Among coastal ecosystems, river plumes are relevant marine areas because of their impact on physical and biogeochemical processes driving the seasonal and spatial dynamics of planktonic communities (D'sa and Miller, 2003; Mestres et al., 2003; Masotti et al., 2018). The most evident characteristics of river plumes are their low salinity, strong stratification, the generation of buoyancy-driven currents around the frontal area, and the higher turbidity associated with suspended solids. Although there are many forcings influencing the dispersion and dynamics of river plumes (e.g. tides, topography, inertia, local circulation, Earth rotation and buoyancy) the river discharge and wind stress dominate the river plume dynamics (Fong and Rockwell Geyer, 2001; Lentz and Largier, 2006; Fernández-Nóvoa et al., 2015; Horner-Devine et al., 2015). For instance, Hetland (2010) demonstrated that the magnitude and length scale of cross-shore plume density changes are directly proportional to the river discharge. On the other hand, a recent study demonstrates that infra-gravity wave forcing has considerable influence on the near-field plume dynamics (Flores et al., 2022), which could influence mixing and cross-shore dispersion.

25 A proper description about the variability and trend of variables like salinity, nutrients and suspended solids in river plumes allows to estimate the condition of the catchment-coast system, where activities like deforestation, agriculture, and urban inputs can change the temporal pattern and influence of river discharges (Acker et al., 2009; Bainbridge et al., 2012; Martínez et al., 2018, 2022), in addition to the effects associated with climate change. For instance, by comparing different river systems, Acker et al. (2009) concluded that a decreasing trend in chlorophyll concentration within the river-influenced area is associated with a reduction of the river discharges.

River plumes usually have high nutrient content and support primary production and algal biomass (Mallin et al., 2005; Peterson and Peterson, 2008; Kudela and Peterson, 2009). The influence of river plumes on larval transport and survival is related with their tolerance to osmotic shocks and their capacity to move vertically through the water column and density gradients (Bloodsworth et al., 2015). Different taxonomic groups will have different conditions to move and survive within river-influenced environments (Bloodsworth et al., 2015). Similarly, the influence of the river plume on sediments and benthos can be observed several (Forrest et al., 2007) to hundred (Grimes and Kingsford, 1996) kilometers from the river mouth. Other studies have demonstrated that the turbidity associated with the plume can influence the predation mortality of larval fish (Carreon-Martinez et al., 2014) and phytoplanktonic communities (Chakraborty and Lohrenz, 2015). The reduction of river discharges can have a negative impact on the fishery resources, biodiversity and ecological functions (Fan et al., 2022). For instance, in central Chile, Vargas et al. (2006) described the influence of the Maipo river plume on the distribution of chlorophyll and barnacle larvae on the inner shelf.

Several studies have demonstrated the importance of wind forcing on river plume dynamics by combining numerical modeling results with remote sensing and *in situ* observations. Choi and Wilkin (2007) demonstrated the strong influence of buoyancy and wind forcing on the Hudson River plume. Similar findings were described for the Yukon River plume by Clark and Mannino (2022). The change of river plume direction caused by winds associated with the passage of low atmospheric pressure systems was investigated using the Navy Coastal Ocean Model (Cobb et al., 2008). The plume extension and orientation driven by the river discharge-wind interplay was described by Dong et al. (2004) for the Pearl River plume. An

interesting modeling study showed the influence of Columbia River plume on the continental shelf during upwelling conditions, affecting the alongshore and cross-shore momentum transport as well as the vertical turbulence structure (Fulton, 2007).
50 Normally, the river plume flows northward along the Washington coast, but in spring-summer the upwelling-favorable winds and coastal circulation forces the plume farther south and offshore off Oregon (Hickey et al., 2005; Saldías et al., 2016). In general, the role of the wind forcing is enhanced on small river plumes (Osadchiev et al., 2021; Basdurak and Largier, 2022), which is the case for most river outflows along central-southern Chile (Saldías et al., 2016).

The relevance of combining hydrodynamic models with remote sensing and *in situ* data has been highlighted by Devlin et al. (2015), who studied the river plumes on the Great Barrier Reef using MODIS imagery. In another study, Bai et al. (2010) used MERIS sensor data to characterize the Changjiang Estuary. In a recent study, Bainbridge et al. (2012) used MODIS images to describe the influence of river discharge, wind and Coriolis forcing on the Burdekin River plume and the transport of fine sediments and nutrients. In a study focused on the Amur River, Abrosimova et al. (2009) compared direct observations with MODIS images and concluded that the plume is highly dynamic and is mostly controlled by the Earth's rotation in comparison
60 with the plume inertial effects.

Central Chile is a region characterized by the presence of several rivers that discharge freshwater and sediment mostly in winter; this region exhibits almost permanent upwelling-favorable southwesterly winds and the presence of Subantarctic Water (SAAW) and some times with remnants of Subtropical Water (STW), with salinities higher than 34.4. Recent studies of river plumes off central Chile have been conducted using satellite imagery (Saldías et al., 2012, 2016) and numerical modeling
65 results (Salcedo-Castro et al., 2020; Rojas et al., 2023; Vergara et al., 2023). These studies described river plumes with sea surface salinity (SSS) values typically lower than 33.9 (Piñones et al., 2005; Saldías et al., 2012; Salcedo-Castro et al., 2020; Vergara et al., 2023), a high seasonality in plume spreading and turbidity signals associated with the river discharges. In austral winter, a larger areal extent and the merging of the plumes can be observed after storms (coalescence events), whereas smaller plumes restricted to the nearfield region are observed in austral spring-summer. A recent study combining remote sensing
70 and numerical modeling results confirms that the river plumes and flow field are primarily modulated by the wind forcing in winter (Rojas et al., 2023). This work also highlights that the geostrophic component of the flow is associated with the wind modulation of the plume's shape on a synoptic scale.

This study aims to describe the interannual variability of the circulation and hydrographic conditions (and stratification) in the coastal area influenced by two rivers off central Chile: Maipo and Rapel rivers, complementing the previous study by
75 Salcedo-Castro et al. (2020) which was focused on the plume spreading climatology and vertical structure. To the best of our knowledge, this is the first interannual modeling study of these river plumes.

2 Methods

2.1 Study area

The study area, delimited by 32°30'S-34°S latitude, is representative of the Mediterranean climate of central Chile, where
80 the Maipo and Rapel rivers are discharged (Fig. 1). These are mixed rivers with snow- and rain-fed regimes, having higher

discharges in winter and late spring. As most of Chilean rivers, they cover a relatively short distance across transverse and longitudinal valleys between the mountains and the coast, and have a small watershed (Saldías et al., 2012). However, their discharges are intervened by activities associated with mining, agriculture, industries and urban development, as there are several cities with a large population. The lower part of Rapel River is downstream of Rapel Dam, a reservoir finished in 1968.

85 In the lower coastal-estuarine region, the river discharges are under the influence of tides and local topography, characterized by a partially opened sandbar, with a strong seasonal wind influence (Flores et al., 2022).

2.2 Numerical model

We used the Coastal and Regional Ocean Community model (CROCO) and CROCO_TOOLS package (<http://www.croco-ocean.org>). This version solves the primitive hydrostatic equations of ocean dynamics, uses the terrain following coordinate and is an

90 adaptation of ROMS_AGRIF (Penven et al., 2006; Debreu et al., 2012), which is based on a new nonhydrostatic and a non-Boussinesq solver developed within the former ROMS kernel (Shchepetkin and McWilliams, 2005), for an optimal accuracy and cost efficiency (Hilt et al., 2020). The model was configured with a 1 km horizontal resolution (Arakawa C-grid) and 20 vertical levels, with higher resolution toward the surface and bottom levels. This configuration allows us to resolve sub-

95 mesoscale features of the river plumes and their interaction with mesoscale processes. The model momentum and buoyancy fluxes were forced with the Scatterometer Climatology of Ocean Winds (SCOW) and the Comprehensive Ocean-Atmosphere Data Set (COADS), which have a 25 km resolution. Boundary conditions were obtained from the 10 km resolution Ocean General Circulation Model for the Earth Simulator (OFES), which was forced with fluxes from the National Centers for Environmental Prediction (NCEP). Daily river discharges from Maipo and Rapel rivers were obtained from the General Direction of Waters (Dirección General de Aguas, Chile) and complemented with the CAMELS-CL dataset (Alvarez-Garreton et al.,

100 2018). The Rapel River gauge is located 25 km downstream of the dam and 16.5 km from the river mouth. The Maipo River gauge (Cabimbao Station) is 21 km upstream from the river mouth. The monthly mean wind forcing and river discharges in the study area are shown in Fig. 2. The modeled period was 2003-2011, which is probably not just enough to reflect the effect of long-term climatic events like the Pacific Interdecadal Oscillation (PDO); however, our simulations encompassed contrasting years with El Niño/La Niña affecting the river discharges and the plumes in the coastal ocean.

105 The horizontal distribution of salinity was described to study the spatial-temporal variability of the plumes. A salinity value of 33.8 was used to delimit the plumes from ambient waters after computing monthly averages; this value is consistent with the reference value described by Rojas et al. (2023). To estimate the plume's area, a finer grid was generated by linear interpolation and then exported as Geotiff images. These images were processed on ArcGIS Pro to measure the plume area and mean plume surface salinity (SSS).

110 Along with the description of salinity distribution, we also estimated the horizontal gradient of salinity (Yu, 2015; Freeman and Lovendus 2016; Saldías and Lara, 2020; Bao et al., 2021) as an indication of the strength of the plume's frontal characteristics. We computed the meridional ($S_{grad(x)}$) and zonal ($S_{grad(y)}$) gradients according to eqn. 1 and eqn. 2, respectively, which were combined to obtain the gradient magnitude at the center of each 1 km² grid cell (S_{grad}) (eqn. 3).

$$\text{Sgrad}_{(x)} = \frac{1}{2} \left(\frac{\text{S}_{(i+1,j)} - \text{S}_{(i,j)}}{\text{Lon}_{(i+1,j)} - \text{Lon}_{(i,j)}} + \frac{\text{S}_{(i+1,j+1)} - \text{S}_{(i,j+1)}}{\text{Lon}_{(i+1,j+1)} - \text{Lon}_{(i,j+1)}} \right) \quad (1)$$

$$115 \quad \text{Sgrad}_{(y)} = \frac{1}{2} \left(\frac{\text{S}_{(i,j+1)} - \text{S}_{(i,j)}}{\text{Lat}_{(i,j+1)} - \text{Lat}_{(i,j)}} + \frac{\text{S}_{(i+1,j+1)} - \text{S}_{(i+1,j)}}{\text{Lat}_{(i+1,j+1)} - \text{Lat}_{(i+1,j)}} \right) \quad (2)$$

$$\text{Sgrad} = \sqrt{\text{Sgrad}_{(x)}^2 + \text{Sgrad}_{(y)}^2} \quad (3)$$

We used eqn. 4 to compute the area-averaged salinity gradient (AASG) in the whole domain (Salcedo-Castro et al., 2015), where "Area" is the surface of each 1 km² grid cell and "Total area" is the model's total domain surface.

$$\text{AASG} = \frac{1}{\text{Total area}} \sum_{i=1, j=1}^{i=m, j=n} \text{Sgrad}_{i,j} \times \text{Area}_{i,j} \quad (4)$$

120 The stratification in terms of the contribution by river discharges was assessed through the potential energy anomaly (PEA, J m⁻³) (Simpson et al., 1978; O'Donnell, 2010; Rojas et al., 2023) (eqn. 5):

$$\phi = \int_{-H}^{\eta} g(\rho - \bar{\rho}) z dz, \quad (5)$$

where ϕ is the potential energy anomaly, g is the gravity acceleration, ρ is the local density at depth z , and the term $\bar{\rho}$ corresponds to the depth-averaged density (eqn. 6):

$$125 \quad \bar{\rho} = \frac{1}{H} \int_{-H}^{\eta} \rho(z) dz, \quad (6)$$

where H is the total depth and η is the free surface.

We evaluated the PEA between 1 and 20 m along the transect depicted in Fig. 1 to represent the strength of stratification, as this represents the equivalent work to homogenize the water column (Simpson, 1981).

2.3 Empirical orthogonal functions (EOFs) and wavelet analysis

130 We performed an empirical orthogonal functions (EOF) analysis (Emery and Thomson, 2004) to the daily model outputs to evaluate the interannual variability of the plumes, reducing the dimensions of large datasets to a few significant orthogonal (uncorrelated) modes of variability and their associated time series (represented by the principal components (PC)). Considering that geophysical time series can be hard to be interpret because of the presence, even in the first modes, of complex non-periodic signals (e.g. Olita et al., 2011a), we analyzed the spectrum of the PCs by means of the continuous wavelet trans-
135 form (CWT). The CWT allows the localization of a signal in the time domain neglecting some localization in frequency

(Torrence and Compo, 1998). The rectification of the wavelet power spectra was calculated following Liu et al. (1998). We used the Morlet wavelet after removing the trends. Thus, spurious low-frequency signals are not considered in the analysis. The trend identification was performed by least squares linear fit. The main variable considered for this analysis was the surface salinity, considering our focusing on river plumes that by the way have a strong signal in salinity. We also analysed surface
140 currents, in order to investigate where these currents could be influenced by surface plumes, or vice versa. This was done by considering separately meridional and zonal components of the flow.

3 Results

3.1 Horizontal plume pattern

Previous to analyze the horizontal plume pattern, a qualitative validation of the model was carried out. Considering the un-
145 availability of direct observations (mooring), we used altimetry data from Aviso (Archiving, Validation and Interpretation of Satellite Oceanographic, www.aviso.altimetry.fr) to estimate geostrophic currents and compare with those computed from the model, following Aguirre et al. (2012). As shown in Fig. 3, the mean seasonal geostrophic currents compare well between the model and Aviso, considering the differences in the spatial resolution and the problems normally associated with the satellite sensor close to the coast and the smoothing caused by gridding fields from multiple altimeters (Aguirre et al., 2012). The
150 strong contrast observed in summer is probably associated with the coastal upwelling and varying interannual forcing during the study period (Aguirre et al., 2012). Moreover, this difference is not relevant as the plume dynamics mostly occurs during Winter-Spring. Additionally, we evaluated the model's performance by means of the reflectance associated with suspended sediment, given that the sea surface salinity (SSS) is tightly correlated with this variable in river plume regions. According to previous studies (Saldías et al., 2012, 2016; Masotti et al., 2018), total suspended solids are better sensed in the 645 nm band;
155 therefore, we worked with the same band. We computed the 99th percentile (P99) of the 645nm reflectance in the model domain to compare it with the total river plume area and SSS. Firstly, we tested the validity and consistency of MODIS imagery by comparing the 99th percentile of the 645nm reflectance in the model domain against the river's discharge, obtaining a $r^2=0.76$. Similarly, when comparing the monthly plume area and mean SSS within the plumes versus MODIS (P99), the correlation was $r^2=0.56$ and $r^2=0.46$, respectively.

160 The sea surface salinity (SSS), averaged over the 9-year period, and its standard deviation (STD) are shown in Fig. 4. The plumes are restricted to a relatively short area next to the river mouths and mostly oriented in the NW-NNW direction. This also reflects the mean direction of surface currents driven by northeastward winds. The STD field reflects the largest variability associated with the lowest mean salinity near the river mouths – the plume's signal responds to the pulses of river discharges.

The strong dependence of the plume horizontal pattern on the wind direction is especially evidenced during some winter
165 months (Fig. 5), when wind is able to reverse the plume's northwestward direction. This is the situation observed in June 2008 and July 2010, corresponding to the two largest peaks observed in wind stress direction shown in Fig. 2. The plumes presented contrasting spreading in the coastal ocean, in response to the wind forcing; in winter 2008, the mean wind stress was 0.05 N m^{-2} , whereas the wind stress was predominantly 0.07 N m^{-2} in July 2010.

We defined the limit of the plume's area with a salinity value of 33.8 in order to separate the plumes from ambient waters. A practical reason for choosing 33.8 as a limit was that the plumes kept a maximum regular shape within the domain when using this value. However, the main reason to define the 33.8 threshold is that this is typical value to identify the Subantarctic Water Mass (salinity: 33.8-33.9), the characteristic surface water mass in this region. Consequently, the rationale is that all waters fresher than that salinity must be from the plumes and not from a regional water mass. In a similar approach, this limit coincided with the values defined by other authors that have studied river plumes in Chile (Saldías et al., 2012; Flores et al., 2022). Other authors have applied other criteria, depending on the objectives of the study and the dynamic characteristics of the area under study (Falcieri et al., 2014). The variation of the plume area (< 33.8), total river discharge (Maipo + Rapel) and mean surface salinity is shown in Fig. 6 where it is possible to see a direct and inverse relationship, respectively. These relationships are clearly observed in Fig. 7.

The minimum and maximum salinity in the domain shows a clear increasing salinity trend over the entire period Fig. 8, which is related to a gradual decrease in river discharge. The interannual variability revealed major plume events with reduced salinity during the winters of 2005 and 2008.

The variation of the strength of the area-averaged salinity gradient (AASG) (Fig. 9) is consistent with the extension of the plumes and correlates with the total river discharges (Fig. 6), which involves stronger gradients and fronts during fall-winter.

Besides the variation in the total area, the alongshore extension was also evaluated in order to identify interannual variability of plume influence along the coast (Fig. 1). Episodes of high river discharges are normally accompanied by an extension of up to 30 km southward along the coast and coalescence of the plumes (Fig. 10), which is consistent with changes in wind direction (Fig. 2 and Fig. 5). The years 2005, 2006 and 2008 presented a marked plume extension during austral winter, whereas the summers of years 2005, 2007 and 2010 were characterized by a minimum or absence of the plumes along the entire coast.

3.2 Vertical plume pattern

The effect of river discharges on the vertical distribution of salinity along the longitudinal transect (Fig. 1) is shown in Fig. 11. We can observe that both plumes can distribute individually and there are episodes of coalescence, especially during high river discharges. Also, during some summer-fall months the plumes are so reduced that they are not detected along this transect. If we use a salinity of 34.2 as a vertical boundary value, the plume thickness increased up to 15 m during high river discharge events. These events also involve a stronger stratification. The time variation along the longitudinal transect of stratification strength (Fig. 12), represented by the potential energy anomaly (PEA, $J m^{-3}$), agrees with variations of sea surface salinity (Fig. 10) and river discharges (Fig. 6).

3.3 EOFs and wavelet

The first 3 modes of variability (Fig. 13 to Fig. 15) of the surface salinity are presented as it follows: a map representing the EOF mode, thus, below, the EOF expansion coefficient (PC time series) with the related wavelet power spectrum and, on the right, the cumulative wavelet spectrum. The first 4 modes of variability explain the 66.7%, 9.1%, 5.5%, and 4.9% of variability, respectively. The first mode explains the seasonal variability, as it can clearly be deduced by observing the related PC and its

wavelet decomposition. The whole domain is in phase in this mode of variability, i.e. the whole system varies in the same direction (Fig. 13).

In the second mode we can observe an interesting feature, probably related to meteorological events. Here, the time series shows a high frequency signature, with two distinct and relevant events, having a period of few months, centered in 2005 and 2010, that are probably related to particular drought events. This feature is of interest also looking at the spatial counter-phase character, as the southernmost area of the domain shows a different sign with respect to the estuaries area. The shape of the southern plume forms a front with a southern area having a positive sign. This is linked to a low-precipitations mode that reduces the area of influence of the estuaries with their plume, with special reference to the southern one (Fig. 14). Also the direction of the plumes for this mode of variability seems to be more northward than the usual climatological NW direction, which is related to changes in wind direction and intensity for this two particular events.

The third mode is related again to a high frequency signal superimposed on a longer signal at the end of the series. This corresponds to a reduction of this signature of the plumes, where a high frequency signature characterizes a longer-period signal most likely related to the beginning of the 2010-2015 mega-drought (Fig. 15). The most noticeable events are those shown in the second mode of variability. This feature in the spatial mode corresponds to two distinct minima of surface salinity related, especially to the Rapel River plume.

For what concerns zonal and meridional velocities, we just show the first three modes of variability of the meridional component (v), i.e. the north-south part of the flow. The zonal component analysis does not show significant results, and/or easy to be interpreted probably because the geography and morphology of the central Chile coast that is dominated by the meridional component of the flow.

The three modes, shown in Fig. 16 to Fig. 18 explain the 42, 15 and 12 percent of variability, respectively, linked to the annual cycle, the semiannual cycle and to some mesoscale variability (high frequency) linked on its turn to part semiannual and annual frequencies (i.e. the analysis put in relation such scales, that actually are interlinked by means of energy cascades). What is evident is, for all the 3 modes, a clear influence of the river runoff area on the spatial variability of the meridional transport.

On the other side, at first sight, there is not a clear similarity between PCs of the salinity with the meridional flow PCs. Considering the wavelet analysis, above all in terms of power spectrum and cumulative spectrum, the first mode of the meridional transport Fig.16 shows some similarity with the spectra shown by the third EOF surface salinity variability (Fig.15). So, a good part of the meridional transport variability, especially in the "blue" area of this first mode, could be influenced by processes linked to the third mode of the salinity which seems related to interannual events, as it is observable by frequencies of the cumulative power spectrum (panel c). It also seems that the other part of such interannual variability is also enclosed in the second mode of salinity variability, in particular for the mid-2005 event, that in salinity is evident in the second mode (Fig. 14), while in terms of meridional velocity is separated in the third variability mode. Also in this case, the event, with a semi-annual signal in terms of cumulative spectrum, seems to have a correspondence in terms of meridional transport variations.

During the 9-year period of study, the river plumes were normally restricted to a short distance from the river mouths, where they also exhibited larger horizontal variability but a more homogeneous vertical structure close to the river mouth. In an analytical study, Hetland (2005) stated that mixing is more intense near the river mouth while wind efficiency is higher farther away, which is consistent with the plume shape and orientation in our results. Similarly, other authors have pointed out that in the near-field the plume behaves as a buoyant jet with stronger effect of the bottom, while in the far-field the horizontal density gradient is weaker and strongly affected by the wind and Coriolis force (Chao, 1988; Chen et al., 2009; Horner-Devine et al., 2015). In our study, we observed a strongly stratified plume even in the far field during winter months. This contrast is explained by the combination of larger river discharge and weaker wind efficiency.

Considering their relatively steep slopes and small watersheds, compared to larger rivers, the Maipo and Rapel rivers are similar to systems of small mountainous rivers that exhibit a rapid response to episodes of increase in freshwater, generating river plumes with a strong stratification near the river mouth (Warrick et al., 2004). An interesting feature observed in river plumes is the occurrence of "rooted" plumes in shallow areas (Zhi et al., 2022). It is possible that this process can also occur off central Chile in winter when there is a predominance of northwesterly winds.

The plume extension and mean surface salinity is strongly correlated with the river discharge, with wind playing a secondary role mostly influencing the plume shape and orientation, as observed during two winter episodes with northwesterly winds also described elsewhere by Rojas et al. (2023). Comparing different forcing, Hickey et al. (2005) state that the extension and shape of the river plumes mostly depend on river runoff, wind and surface currents over the continental shelf. Although there are other factors that influence the spread of a river discharge, like tides, characteristics of the discharge, bathymetry, and Coriolis acceleration (Archetti and Mancini, 2012). For instance, Chen et al. (2017a) state that, outside the micro-tidal estuary, wind is the main forcing contributing energy for mixing the Pearl River plume over the shelf. This explains the vertical pattern observed in the Maipo-Rapel river area, in agreement with Rojas et al. (2023), where the higher PEA and stratification are observed in winter, when wind is weaker and river discharge is the most important driver of the plume dynamics, except during the northwesterly wind events.

Fernández-Nóvoa et al. (2015) showed that the river discharge had a higher variability during the period of higher discharge and when landward, downwelling-favorable winds pushed the plume to the coast, making it flow along the coast. Something similar is observed during episodes of strong NNW winds off Maipo-Rapel rivers occurring in winter. This seasonal change in the plume extension and direction had already been described by other authors (Fiedler and Laurs, 1990; García Berdeal et al., 2002).

Subtidal currents associated with wind-influenced currents and mesoscale eddies are also an important forcing linked to the transport of river plume sediments over long distances and long time scales (Blaas et al., 2007). However, the intra-annual and seasonal variability remain driven by wind and river discharges, respectively, as it has also been observed in other systems where river discharge controls the plume dynamics in the long term while wind is more relevant in the short term (Falcieri et al., 2014). In this sense, consistent with our results, Piñones et al. (2005), assert that the Maipo River plume is mostly driven by

the river discharge in winter but the influence of wind is more important during spring-summer. Normally, the predominant
270 wind in this region is from southwest, decreasing its intensity in fall-winter (Strub et al., 1998) and with episodic strong storms
during winter (Hernández-Miranda et al., 2003) and periods of upwelling and relaxation and intrusion of oceanic waters during
summer (Letelier et al., 2009; Aguirre et al., 2012).

It is worth to mention that 9 years is a short period to capture any significant impact by the PDO in the study area (although
it shows a clear interannual variability which correlates with the ENSO). However, the impact of El Niño and La Niña showed
275 marked contrasting effects on turbid river plumes off central-southern Chile during the period of study (e.g. Saldías et al.,
2016). In fact, the years of 2005 and 2006 were very wet periods with anomalously high rainfall and peaks in river discharges.
The river plumes were anomalously big during those years in response to the freshwater availability. In fact, these conditions
promoted flooding events in central Chile. In contrast, periods such as 2007 and 2011 were characterized by anomalously low
signatures of freshwater plumes from the satellite observations which coincided with the influence of La Niña and a lack of
280 rainfall in central-southern Chile (Saldías et al., 2016). In this sense, Hernandez et al. (2022) pointed out that, even though
some catchments are strongly correlated to ENSO with related hydro-climatic anomalies, mixed regimes do not exhibit a clear
connection. On the other hand, Alvarez-Garreton et al. (2021) stated that snow-dominated catchments are especially vulnerable
to long term droughts, showing the accumulated effect from previous years; this could be reflected especially in a reduction
of the plumes extension during spring-summer, as observed after 2008 (Fig. 6). This long-term drought has been described by
285 other authors (Winckler et al., 2020), which is attributed to the anthropogenic climate change, with an evident decrease in river
discharges in central Chile since 2010.

The pattern here described is opposite to the interannual and seasonal variability of the Columbia River. Here, Burla et al.
(2010) and Chen et al. (2017a) described a northward plume attached to the coast in winter and a detached plume in summer,
detaching that is normally observed during events of wind relaxation or wind reversal. Moreover, Chen et al. (2017b) as-
290 serted that upwelling jets are able to transport river plumes long distances along the coast. However, whereas Columbia River
has a snow-dominated regime where a maximum river discharge coincides with the period of intense upwelling-favorable
winds, river plumes in central Chile are characterized by a phase difference between higher freshwater discharge and stronger
upwelling-favorable winds. The detachment of the plumes (northwestward direction) observed during summer in this study is
in agreement with the Ekman theory, as described by Rojas et al. (2023) and Saldías et al. (2012), where upwelling-favorable
295 wind forces the detachment and direction of the buoyancy-driven plume.

An interesting point to consider is the conclusion by Berghuijs et al. (2014), who stated that a shift from snow- to rain-
dominated regimes in some catchments would lead to a decrease in the mean streamflow. In pluvio-nival regimes like Rapel
and Maipo rivers, this would mean that plumes extending on coastal areas would mostly depend on the river discharges
occurring in winter. In this sense, Döll and Schmied (2012) modeled climate change projections and asserted that low flows
300 could decrease up to 50% and some systems could even change their regime from perennial to intermittent.

As described by Garreaud and Falvey (2009), it is expected that future conditions in this region will be characterized by
stronger southerly winds, which would involve that Rapel and Maipo rivers could extend further north and closer to the coast.
On the other hand, projections also predict a southward extension of the semi-arid climate (Winckler et al., 2020), which

means that the river discharges in this region would tend to decrease over time. Thus, we expect that smaller river plumes
305 would extend closer to the coast and probably having an impact on the distribution of benthic communities and larval stages
(Grimes and Kingsford, 1996), besides a possible impact on the sand supply to beaches that strongly depend on these rivers
and that already show a progressive erosion and shoreline retreat (Martínez et al., 2018, 2022). Although not explored in this
study, it is evident that ENSO hydro-climatic anomalies have a strong influence on the hydrological regime (Hernandez et al.,
2022) and the river plume structure along the continental shelf off central-southern Chile (Saldías et al., 2016). Consequently,
310 it is expected to observe changes in the pattern described here during the coming decades.

Although the model does not include the wave effect, further studies should consider this variable as most of the plume
remains attached to the coasts, especially, during high-energy wave events in winter. In this sense, Delpy et al. (2014) describes
the strong effect that waves can have on the river plume dynamics, including alongshore currents and flushing time. A recent
study on Maipo River plume also demonstrates the strong influence that waves can have on plume dynamics on shallow areas
315 (Flores et al., 2022).

5 Conclusions

The hydrography of the area influenced by Maipo and Rapel River plumes was modeled for the period 2003-2011, where a
strong dependence of the plumes features on freshwater discharge and wind forcing could be evidenced. Unlike other systems
like Columbia River, the larger extent, lower mean salinity and stronger stratification is observed in winter time, when wind
320 is weaker and some times downwelling-favorable; in spring-summer, when upwelling-favorable wind is stronger, these river
plumes are smaller and their stratification is weaker, which is consistent with previous studies. Strong, downwelling-favorable,
events are able to reverse the river plume direction and push it southward to form a narrow band attached to the coast. The
EOFs analysis confirmed the strong seasonality of the Maipo-Rapel river plume system. A second mode showed an intra-annual
signal, likely associated with meteorological events, which also exhibited a contrasting north-south spatial sign.

325 An increasing trend of mean salinity was observed in the study domain, associated with a decreasing trend in river discharges.
This trend corresponds to the beginning of the 2010-2015 mega-drought exhibited by this region. No correlation was found
between the plumes characteristics and El Niño-Southern Oscillation (ENSO) and Pacific Interdecadal Oscillation (PDO)
indices, which would be explained by the complexity of mixed regimes and the fact that snow melting regimes show a lag and
cumulative response from previous years. It is possible to speculate that a shift from snow- to rain-dominated regimes, along
330 with changes in wind and precipitation patterns will be reflected on smaller river plumes and more attached to the coast, along
with an increase in local mean surface salinity, which would affect planktonic and benthic communities.

Author contributions. TEXT

JSC elaborated the idea and general organization of the manuscript; AO undertook the EOF and wavelets analysis along with the respective discussion and the analysis of the long term trends. FS and GS contributed to the remote sensing review, analysis, and discussion. RCG and CDLTM collaborated with the altimeter data analysis, and in the edition and discussions.

Competing interests. TEXT

No competing interests are present.

Disclaimer. TEXT

Acknowledgements. This study was funded by CONICYT FONDECYT Chile, grant 11160309 “Numerical modeling of river plumes in central Chile (32S-34S) and assessment of climate change scenarios” and ANID Millennium Science Initiative Program-Code ICN2019_015 (Millennium Institute SECOS). JSC thanks the valuable collaboration of Mr. David Donoso for configuring and running the numerical experiments. GSS is partially funded by COPAS COASTAL ANID FB210021 and Fondecyt 1220167. We acknowledge the National Laboratory for High Performance Computing – Chile for providing the computational capabilities to run the numerical simulations for this project.

References

- 345 Abrosimova, A., Zhabin, I., and Dubina, V.: Influence of the Amur River runoff on the hydrological conditions of the Amur Liman and Sakhalin Bay (Sea of Okhotsk) during the spring-summer flood, Tech. rep., V.I. Il'ichev Pacific Oceanological Institute, FEB RAS, Vladivostok, Russia, <https://doi.org/10.3103/S1068373910040084>, 2009.
- Acker, J. G., McMahon, E., Shen, S., Hearty, T., and Casey, N.: Time-series analysis of remotely-sensed SeaWiFS chlorophyll in river-influenced coastal regions, in: *EARSel eProceedings*, 8(2), pp. 114–139, 2009.
- 350 Aguirre, C., Pizarro, Ó., Strub, P. T., Garreaud, R., and Barth, J. A.: Seasonal dynamics of the near-surface alongshore flow off central Chile, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/10.1029/2011JC007379>, 2012.
- Alvarez-Garreton, C., Mendoza, P. A., Pablo Boisier, J., Addor, N., Galleguillos, M., Zambrano-Bigiarini, M., Lara, A., Puelma, C., Cortes, G., Garreaud, R., McPhee, J., and Ayala, A.: The CAMELS-CL dataset: Catchment attributes and meteorology for large sample studies-Chile dataset, *Hydrology and Earth System Sciences*, 22, 5817–5846, <https://doi.org/10.5194/hess-22-5817-2018>, 2018.
- 355 Alvarez-Garreton, C., Pablo Boisier, J., Garreaud, R., Seibert, J., and Vis, M.: Progressive water deficits during multiyear droughts in basins with long hydrological memory in Chile, *Hydrology and Earth System Sciences*, 25, 429–446, <https://doi.org/10.5194/hess-25-429-2021>, 2021.
- Archetti, R. and Mancini, M.: Freshwater dispersion plume in the sea: Dynamic description and case study, in: *Hydrodynamics - Natural Water Bodies*, edited by Schulz, H. E., Simoes, A. L. A., and Lobosco, R. J., IntechOpen, London, <https://doi.org/10.5772/28390>, 2012.
- 360 Bai, Y., He, X., Pan, D., Zhu, Q., Lei, H., Tao, B., and Hao, Z.: The extremely high concentration of suspended particulate matter in Changjiang Estuary detected by MERIS data, in: *Remote Sensing of the Coastal Ocean, Land, and Atmosphere Environment*, vol. 7858, p. 78581D, SPIE, <https://doi.org/10.1117/12.869632>, 2010.
- Bainbridge, Z. T., Wolanski, E., Álvarez-Romero, J. G., Lewis, S. E., and Brodie, J. E.: Fine sediment and nutrient dynamics related to particle size and floc formation in a Burdekin River flood plume, Australia, *Marine Pollution Bulletin*, 65, 236–248, <https://doi.org/10.1016/j.marpolbul.2012.01.043>, 2012.
- 365 Bao, S., Wang, H., Zhang, R., Yan, H., Chen, J., and Bai, C.: Application of Phenomena-Resolving Assessment Methods to Satellite Sea Surface Salinity Products, *Earth and Space Science*, 8, 1–14, <https://doi.org/10.1029/2020EA001410>, 2021.
- Basdurak, N. and Largier, J.: Wind Effects on Small-Scale River and Creek Plumes, *Journal of Geophysical Research: Oceans*, p. e2021JC018381, 2022.
- 370 Berghuijs, W. R., Woods, R. A., and Hrachowitz, M.: A precipitation shift from snow towards rain leads to a decrease in streamflow, *Nature Climate Change*, 4, 583–586, <https://doi.org/10.1038/nclimate2246>, 2014.
- Blaas, M., Dong, C., Marchesiello, P., McWilliams, J. C., and Stolzenbach, K. D.: Sediment-transport modeling on Southern Californian shelves: A ROMS case study, *Continental Shelf Research*, 27, 832–853, <https://doi.org/10.1016/j.csr.2006.12.003>, 2007.
- Bloodsworth, K. H., Tilburg, C. E., and Yund, P. O.: Influence of a River Plume on the Distribution of Brachyuran Crab and Mytilid Bivalve Larvae in Saco Bay, Maine, *Estuaries and Coasts*, 38, 1951–1964, <https://doi.org/10.1007/s12237-015-9951-5>, 2015.
- 375 Burla, M., Baptista, A. M., Zhang, Y., and Frolov, S.: Seasonal and interannual variability of the Columbia River plume: A perspective enabled by multiyear simulation databases, *Journal of Geophysical Research*, 115, <https://doi.org/10.1029/2008JC004964>, 2010.
- Carreon-Martinez, L. B., Wellband, K. W., Johnson, T. B., Ludsins, S. A., and Heath, D. D.: Novel molecular approach demonstrates that turbid river plumes reduce predation mortality on larval fish, *Molecular Ecology*, 23, 5366–5377, <https://doi.org/10.1111/mec.12927>, 2014.
- 380

- Chakraborty, S. and Lohrenz, S. E.: Phytoplankton community structure in the river-influenced continental margin of the northern Gulf of Mexico, *Marine Ecology Progress Series*, 521, 31–47, <https://doi.org/10.3354/meps11107>, 2015.
- Chao, S.-Y.: River-forced estuarine plumes, *Journal of Physical Oceanography*, 18, 72–88, 1988.
- Chen, F., MacDonald, D. G., and Hetland, R. D.: Lateral spreading of a near-field river plume: Observations and numerical simulations, *Journal of Geophysical Research: Oceans*, 114, <https://doi.org/10.1029/2008JC004893>, 2009.
- 385 Chen, Z., Gong, W., Cai, H., Chen, Y., and Zhang, H.: Dispersal of the Pearl River plume over continental shelf in summer, *Estuarine, Coastal and Shelf Science*, 194, 252–262, <https://doi.org/10.1016/j.ecss.2017.06.025>, 2017a.
- Chen, Z., Pan, J., Jiang, Y., and Lin, H.: Far-reaching transport of Pearl River plume water by upwelling jet in the northeastern South China Sea, *Journal of Marine Systems*, 173, 60–69, <https://doi.org/10.1016/j.jmarsys.2017.04.008>, 2017b.
- 390 Choi, B. J. and Wilkin, J. L.: The effect of wind on the dispersal of the Hudson River plume, *Journal of Physical Oceanography*, 37, 1878–1897, <https://doi.org/10.1175/JPO3081.1>, 2007.
- Clark, J. B. and Mannino, A.: The Impacts of Freshwater Input and Surface Wind Velocity on the Strength and Extent of a Large High Latitude River Plume, *Frontiers in Marine Science*, 8, <https://doi.org/10.3389/fmars.2021.793217>, 2022.
- Cobb, M., Keen, T. R., and Walker, N. D.: Modeling the circulation of the Atchafalaya Bay system. Part 2: River plume dynamics during cold fronts, *Journal of Coastal Research*, 24, 1048–1062, <https://doi.org/10.2112/07-0879.1>, 2008.
- 395 Debreu, L., Marchesiello, P., Penven, P., and Cambon, G.: Two-way nesting in split-explicit ocean models: Algorithms, implementation and validation, *Ocean Modelling*, 49–50, 1–21, <https://doi.org/10.1016/j.ocemod.2012.03.003>, 2012.
- Delpey, M. T., Arduin, F., Otheguy, P., and Jouon, A.: Effects of waves on coastal water dispersion in a small estuarine bay, *Journal of Geophysical Research: Oceans*, 119, 70–86, <https://doi.org/10.1002/2013JC009466>, 2014.
- 400 Devlin, M. J., Petus, C., da Silva, E., Tracey, D., Wolff, N. H., Waterhouse, J., and Brodie, J.: Water quality and river plume monitoring in the Great Barrier Reef: An overview of methods based on ocean colour satellite data, *Remote Sensing*, 7, 12 909–12 941, <https://doi.org/10.3390/rs71012909>, 2015.
- Döll, P. and Schmied, H. M.: How is the impact of climate change on river flow regimes related to the impact on mean annual runoff? A global-scale analysis, *Environmental Research Letters*, 7, <https://doi.org/10.1088/1748-9326/7/1/014037>, 2012.
- 405 Dong, L., Su, J., Ah Wong, L., Cao, Z., and Chen, J. C.: Seasonal variation and dynamics of the Pearl River plume, *Continental Shelf Research*, 24, 1761–1777, <https://doi.org/10.1016/j.csr.2004.06.006>, 2004.
- D’sa, E. J. and Miller, R. L.: Bio-optical properties in waters influenced by the Mississippi River during low flow conditions, *Remote Sensing of Environment*, 84, 538–549, www.elsevier.com/locate/rse, 2003.
- Emery, W. J. and Thomson, R. E.: *Data Analysis Methods in Physical Oceanography*, Elsevier, Amsterdam, 2nd edn., 2004.
- 410 Falcieri, F. M., Benetazzo, A., Sclavo, M., Russo, A., and Carniel, S.: Po River plume pattern variability investigated from model data, *Continental Shelf Research*, 87, 84–95, <https://doi.org/10.1016/j.csr.2013.11.001>, 2014.
- Fan, Y., Zhang, S., Du, X., Wang, G., Yu, S., Dou, S., Chen, S., Ji, H., Li, P., and Liu, F.: The effects of flow pulses on river plumes in the Yellow River Estuary, in spring, *Journal of Hydroinformatics*, 00, 1–15, <https://doi.org/10.2166/hydro.2022.049>, 2022.
- Fernández-Nóvoa, D., Mendes, R., DeCastro, M., Dias, J. M., Sánchez-Arcilla, A., and Gómez-Gesteira, M.: Analysis of the influence of river discharge and wind on the Ebro turbid plume using MODIS-Aqua and MODIS-Terra data, *Journal of Marine Systems*, 142, 40–46, <https://doi.org/10.1016/j.jmarsys.2014.09.009>, 2015.
- 415 Fiedler, P. C. and Laurs, R. M.: Variability of the Columbia River plume observed in visible and infrared satellite imagery, *International Journal of Remote Sensing*, 11, 999–1010, 1990.

- Flores, R. P., Williams, M. E., and Horner-Devine, A. R.: River Plume Modulation by Infragravity Wave Forcing, *Geophysical Research Letters*, 49, <https://doi.org/10.1029/2021GL097467>, 2022.
- 420 Fong, D. A. and Rockwell Geyer, W.: Response of a river plume during an upwelling favorable wind event, *Journal of Geophysical Research: Oceans*, 106, 1067–1084, <https://doi.org/10.1029/2000JC900134>, 2001.
- Forrest, B. M., Gillespie, P. A., Cornelisen, C. D., and Rogers, K. M.: Multiple indicators reveal river plume influence on sediments and benthos in a New Zealand coastal embayment, *New Zealand Journal of Marine and Freshwater Research*, 41, 13–24, 2007.
- 425 Freeman, N. M. and Lovenduski, N. S.: Mapping the Antarctic Polar Front: Weekly realizations from 2002 to 2014, *Earth System Science Data*, 8, 191–198, <https://doi.org/10.5194/essd-8-191-2016>, 2016.
- Fulton, D. P.: Modeling the Columbia River Plume on the Oregon Shelf during Summer Upwelling, Tech. rep., Cooperative Institute for Oceanographic Satellite Studies, <http://www.orcoos.org>, 2007.
- García Berdeal, I., Hickey, B. M., and Kawase, M.: Influence of wind stress and ambient flow on a high discharge river plume, *Journal of Geophysical Research: Oceans*, 107, <https://doi.org/10.1029/2001jc000932>, 2002.
- 430 Garreaud, R. D. and Falvey, M.: The coastal winds off western subtropical South America in future climate scenarios, *International Journal of Climatology*, 29, 543–554, <https://doi.org/10.1002/joc.1716>, 2009.
- Grimes, C. B. and Kingsford, M. J.: How do riverine plumes of different sizes influence fish larvae: do they enhance recruitment?, *Marine and Freshwater Research*, 47, 191–208, 1996.
- 435 Hernandez, D., Mendoza, P. A., Boisier, J. P., and Ricchetti, F.: Hydrologic Sensitivities and ENSO Variability Across Hydrological Regimes in Central Chile (28°–41°S), *Water Resources Research*, 58, <https://doi.org/10.1029/2021wr031860>, 2022.
- Hernández-Miranda, E., Palma, A. T., and Ojeda, F. P.: Larval fish assemblages in nearshore coastal waters off central Chile: Temporal and spatial patterns, *Estuarine, Coastal and Shelf Science*, 56, 1075–1092, [https://doi.org/10.1016/S0272-7714\(02\)00308-6](https://doi.org/10.1016/S0272-7714(02)00308-6), 2003.
- Hetland, R. D.: Relating river plume structure to vertical mixing, *Journal of Physical Oceanography*, 35, 1667–1688, 2005.
- 440 Hetland, R. D.: The effects of mixing and spreading on density in near-field river plumes, *Dynamics of Atmospheres and Oceans*, 49, 37–53, <https://doi.org/10.1016/j.dynatmoce.2008.11.003>, 2010.
- Hickey, B., Geier, S., Kachel, N., and MacFadyen, A.: A bi-directional river plume: The Columbia in summer, *Continental Shelf Research*, 25, 1631–1656, <https://doi.org/10.1016/j.csr.2005.04.010>, 2005.
- Hilt, M., Auclair, F., Benshila, R., Bordoais, L., Capet, X., Debreu, L., Dumas, F., Jullien, S., Lemarié, F., Marchesiello, P., Nguyen, C., and 445 Roblou, L.: Numerical modelling of hydraulic control, solitary waves and primary instabilities in the Strait of Gibraltar, *Ocean Modelling*, 151, <https://doi.org/10.1016/j.ocemod.2020.101642>, 2020.
- Horner-Devine, A. R., Hetland, R. D., and MacDonald, D. G.: Mixing and transport in coastal river plumes, <https://doi.org/10.1146/annurev-fluid-010313-141408>, 2015.
- Kudela, R. M. and Peterson, T. D.: Influence of a buoyant river plume on phytoplankton nutrient dynamics: What controls standing stocks and productivity?, *Journal of Geophysical Research: Oceans*, 114, <https://doi.org/10.1029/2008JC004913>, 2009.
- 450 Lentz, S. J. and Largier, J.: The influence of wind forcing on the Chesapeake Bay buoyant coastal current, *Journal of Physical Oceanography*, 36, 1305–1316, 2006.
- Letelier, J., Pizarro, O., and Nuñez, S.: Seasonal variability of coastal upwelling and the upwelling front off central Chile, *Journal of Geophysical Research: Oceans*, 114, <https://doi.org/10.1029/2008JC005171>, 2009.
- 455 Liu, W. T., Tang, W., and Polito, P. S.: NASA scatterometer provides global ocean-surface wind fields with more structures than numerical weather prediction, *Geophysical Research Letters*, 25, 761, <https://doi.org/10.1029/98GL00544>, 1998.

- Mallin, M. A., Cahoon, L. B., and Durako, M. J.: Contrasting food-web support bases for adjoining river-influenced and non-river influenced continental shelf ecosystems, *Estuarine, Coastal and Shelf Science*, 62, 55–62, <https://doi.org/10.1016/j.ecss.2004.08.006>, 2005.
- 460 Martínez, C., Contreras-López, M., Winckler, P., Hidalgo, H., Godoy, E., and Agredano, R.: Coastal erosion in central Chile: A new hazard?, *Ocean and Coastal Management*, 156, 141–155, <https://doi.org/10.1016/j.ocecoaman.2017.07.011>, 2018.
- Martínez, C., Grez, P. W., Martín, R. A., Acuña, C. E., Torres, I., and Contreras-López, M.: Coastal erosion in sandy beaches along a tectonically active coast: The Chile study case, *Progress in Physical Geography*, 46, 250–271, <https://doi.org/10.1177/03091333211057194>, 2022.
- Masotti, I., Aparicio-Rizzo, P., Yevenes, M. A., Garreaud, R., Belmar, L., and Farías, L.: The influence of river discharge on nutrient export and phytoplankton biomass off the Central Chile Coast (33°-37°S): Seasonal cycle and interannual variability, *Frontiers in Marine Science*, 5, <https://doi.org/10.3389/fmars.2018.00423>, 2018.
- 465 Mestres, M., Sierra, J. P., Sánchez-Arcilla, A., González del Río, J., Wolf, T., Rodríguez, A., and Ouillo, S.: Modelling of the Ebro River plume. Validation with field observations, *Scientia Marina*, 67, 379–391, 2003.
- O'Donnell, J.: The dynamics of estuary plumes and fronts, in: *Contemporary Issues in Estuarine Physics*, edited by Valle-Levinson, A., p. 186–246, Cambridge University Press, <https://doi.org/10.1017/CBO9780511676567.009>, 2010.
- 470 Olita, A., Ribotti, A., Sorgente, R., Fazioli, L., and Perilli, A.: SLA - chlorophyll-a variability and covariability in the Algero-Provençal Basin (1997-2007) through combined use of EOF and wavelet analysis of satellite data, *Ocean Dynamics*, 61, 89–102, <https://doi.org/10.1007/s10236-010-0344-9>, 2011a.
- Osadchiv, A., Sedakov, R., and Barymova, A.: Response of a small river plume on wind forcing, *Frontiers in Marine Science*, 2021.
- 475 Penven, P., Debreu, L., Marchesiello, P., and McWilliams, J. C.: Evaluation and application of the ROMS 1-way embedding procedure to the central california upwelling system, *Ocean Modelling*, 12, 157–187, <https://doi.org/10.1016/j.ocemod.2005.05.002>, 2006.
- Peterson, J. O. and Peterson, W. T.: Influence of the Columbia River plume (USA) on the vertical and horizontal distribution of mesozooplankton over the Washington and Oregon shelf, *ICES Journal of Marine Science*, 65, 477–483, <http://icesjms.oxfordjournals.org/>, 2008.
- 480 Piñones, A., Valle-Levinson, A., Narváez, D. A., Vargas, C. A., Navarrete, S. A., Yuras, G., and Castilla, J. C.: Wind-induced diurnal variability in river plume motion, *Estuarine, Coastal and Shelf Science*, 65, 513–525, <https://doi.org/10.1016/j.ecss.2005.06.016>, 2005.
- Rojas, C. M., Saldías, G. S., Flores, R. P., Vásquez, S. I., Salas, C., and Vargas, C. A.: A modeling study of hydrographic and flow variability along the river-influenced coastal ocean off central Chile, *Ocean Modelling*, 181, 102–155, <https://doi.org/10.1016/j.ocemod.2022.102155>, 2023.
- 485 Salcedo-Castro, J., de Camargo, R., Marone, E., and Sepúlveda, H.: Using the mean pressure gradient and NCEP/NCAR reanalysis to estimate the strength of the South Atlantic Anticyclone, *Dynamics of Atmospheres and Oceans*, 71, <https://doi.org/10.1016/j.dynatmoce.2015.06.003>, 2015.
- Salcedo-Castro, J., Saldías, G., Saavedra, F., and Donoso, D.: Climatology of Maipo and Rapel river plumes off Central Chile from numerical simulations, *Regional Studies in Marine Science*, 38, <https://doi.org/10.1016/j.rsma.2020.101389>, 2020.
- 490 Saldías, G. S. and Lara, C.: Satellite-derived sea surface temperature fronts in a river-influenced coastal upwelling area off central – southern Chile, *Regional Studies in Marine Science*, 37, 101–322, <https://doi.org/10.1016/j.rsma.2020.101322>, 2020.
- Saldías, G. S., Sobarzo, M., Largier, J., Moffat, C., and Letelier, R.: Seasonal variability of turbid river plumes off central Chile based on high-resolution MODIS imagery, *Remote Sensing of Environment*, 123, 220–233, <https://doi.org/10.1016/j.rse.2012.03.010>, 2012.

- 495 Saldías, G. S., Largier, J. L., Mendes, R., Pérez-Santos, I., Vargas, C. A., and Sobarzo, M.: Satellite-measured interannual variability of turbid river plumes off central-southern Chile: Spatial patterns and the influence of climate variability, *Progress in Oceanography*, 146, 212–222, <https://doi.org/10.1016/j.pocean.2016.07.007>, 2016.
- Saldías, G. S., Shearman, R. K., Barth, J. A., and Tuffillaro, N.: Optics of the offshore Columbia River plume from glider observations and satellite imagery, *Journal of Geophysical Research: Oceans*, 121, 2367–2384, 2016.
- Shchepetkin, A. F. and McWilliams, J. C.: The regional oceanic modeling system (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Modelling*, 9, 347–404, <https://doi.org/10.1016/j.ocemod.2004.08.002>, 2005.
- 500 Simpson, J.: The shelf-sea fronts: implications of their existence and behaviour, *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 302, 531–546, 1981.
- Simpson, J. H., Allen, C. M., and Morris, N. C. G.: Fronts on the continental shelf, *Journal of Geophysical Research: Oceans*, 83, 4607–4614, <https://doi.org/https://doi.org/10.1029/JC083iC09p04607>, 1978.
- 505 Strub, P. T., Mesías, J. M., Montecino, V., Rutlant, J., and Salinas, S.: Coastal ocean circulation off western South America, in: *The Sea*, Volume 11, edited by Robinson, A. R. and Brink, K. H., pp. 273–313, John Wiley and Sons Inc, 1998.
- Torrence, C. and Compo, G. P.: A practical guide to wavelet analysis, *Bull. Am. Meteor. Soc.*, 79, 61–78, 1998.
- Vargas, C. A., Narva, D. A., Piñones, A., Navarrete, S. A., and Lagos, N. A.: River plume dynamic influences transport of barnacle larvae in the inner shelf off central Chile, *Journal of the Marine Biological Association of the United Kingdom*, 86, 1057–1065, 2006.
- 510 Vergara, O. A., Echevin, V., Sobarzo, M., Sepúlveda, H. H., Castro, L., and Soto-Mendoza, S.: Impacts of the freshwater discharge on hydrodynamical patterns in the Gulf of Arauco (central-southern Chile) using a high-resolution circulation model, *Journal of Marine Systems*, 240, 103 862, <https://doi.org/10.1016/j.jmarsys.2023.103862>, 2023.
- Warrick, J. A., A.k. Mertes, L., Washburn, L., and A. Siegel, D.: A conceptual model for river water and sediment dispersal in the Santa Barbara Channel, California, *Continental Shelf Research*, 24, 2029–2043, <https://doi.org/10.1016/j.csr.2004.07.010>, 2004.
- 515 Winckler, P., Aguirre, C., Farías, L., Contreras-López, M., and Masotti, Í.: Evidence of climate-driven changes on atmospheric, hydrological, and oceanographic variables along the Chilean coastal zone, *Climatic Change*, 163, 633–652, <https://doi.org/10.1007/s10584-020-02805-3>, 2020.
- Yu, L.: Sea-surface salinity fronts and associated salinity-minimum zones in the tropical ocean, *Journal of Geophysical Research: Oceans*, 120, 4205–4225, <https://doi.org/10.1002/2015JC010790>.Received, 2015.
- 520 Zhi, H., Wu, H., Wu, J., Zhang, W., and Wang, Y.: River Plume Rooted on the Sea-Floor: Seasonal and Spring-Neap Variability of the Pearl River Plume Front, *Frontiers in Marine Science*, 9, 1–17, <https://doi.org/10.3389/fmars.2022.791948>, 2022.




Figure 1. Study area showing Maipo and Rapel Rivers. The red dashed line represents the transect to describe the temporal variation of sea surface salinity (SSS) and vertical structure across the study area (see Fig. 10, Fig. 11 and Fig. 12).




Figure 2. Monthly mean wind stress (OFES-NCEP) and river discharges (monthly averages) used to force the model in the study area.




Figure 3. Comparison of mean seasonal geostrophic currents from AVISO and CROCO model.




Figure 4. Maps of (left panel) averaged sea surface salinity and (right panel) sea surface salinity standard deviation over the 9 years period.




Figure 5. Salinity distribution in June 2008 and July 2010, corresponding to the changes in wind stress direction to SSE – see Fig. 2.




Figure 6. Variation of (upper panel) river plume area and total river discharge, and (lower panel) mean plume salinity for the period 2003-2011.




Figure 7. Scatter plot of (left panel) river plume area and (right panel) mean plume salinity as function of total river discharge.




Figure 8. Time series and trends of the (top panel) maximum and (bottom panel) minimum surface salinity over the domain




Figure 9. Variation of the area-averaged salinity gradient (AASG) in the Maipo-Rapel plume area.




Figure 10. Hovmöller diagram for sea surface salinity (SSS) along a longitudinal transect (see Fig. 1) extending across Maipo and Rapel river plumes.




Figure 11. Vertical distribution of salinity along the longitudinal transect (Fig. 1) in the Maipo-Rapel River plumes area.




Figure 12. Hovmöller diagram for potential energy anomaly (PEA, J m^{-3}) along the longitudinal transect (see Fig. 1) extending across Maipo and Rapel river plumes.




Figure 13. (Left panel) EOF Mode 1 map. (a) EOF PC 1 time series, (b) Wavelet spectrum of the PC series and (c) cumulative global wavelet spectrum.




Figure 14. (Left panel) EOF Mode 2 map. (a) EOF PC 2 time series, (b) Wavelet spectrum of the PC series and (c) cumulative global wavelet spectrum.




Figure 15. (Left panel) EOF Mode 3 map. (a) EOF PC 3 time series, (b) Wavelet spectrum of the PC series and (c) cumulative global wavelet spectrum.




Figure 16. (Left panel) Meridional velocity EOF Mode 1 map. (a) EOF PC 1 time series, (b) Wavelet spectrum of the PC series and (c) cumulative global wavelet spectrum.




Figure 17. (Left panel) Meridional Velocity EOF Mode 2 map. (a) EOF PC 2 time series, (b) Wavelet spectrum of the PC series and (c) cumulative global wavelet spectrum.




Figure 18. (Left panel) Meridional velocity EOF Mode 3 map. (a) EOF PC 3 time series, (b) Wavelet spectrum of the PC series and (c) cumulative global wavelet spectrum.