

Preliminary evaluation of the effect of electro-coalescence with conducting sphere approximation on the formation of warm cumulus clouds using SCALE-SDM version 0.2.5-2.3.0

5 Ruyi Zhang¹, Limin Zhou^{1,2}, Shin-ichiro Shima^{3,4}, Huawei Yang¹

¹Key Laboratory of Geographic Information Science, Ministry of Education, East China Normal University, Shanghai, 200241, China

²Key Laboratory of Numerical Modeling for Atmospheric Science and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, CAS, Beijing, 100029, China

10 ³Graduate School of Simulation Studies, University of Hyogo, Kobe, 6512103, Japan

⁴RIKEN Center for Computational Science, Kobe, 6500047, Japan

Correspondence to: Limin Zhou(lmzhou@geo.ecnu.edu.cn)

Abstract. The phenomenon of electric fields applied to droplets inducing droplet coalescence is called the electro-coalescence effect. An The analytic expression for electro-coalescence with the accurate electrostatic force for a pair of droplets with opposite sign charges is established by treating the droplets as conducting spheres (CSs). To investigate this effect, we applied a weak electric field to a cumulus cloud using a cloud model that employs the super-droplet method (SDM), a probabilistic particle-based microphysics method. Then, the weak electric effect on a cumulus cloud is investigated by a size-resolved cloud model with a particle-based microphysics method: super-droplet method. This study employs a two-dimensional large eddy simulation (LES) in a flow-coupled model to examine aerosol microphysics (such as collision-coalescence enhancement, velocity fluctuations, and supersaturation fluctuations) in warm cumulus clouds without relying on subgrid dynamics. In the simulation, we assume that droplets carry opposite sign charges and are well mixed within the cloud. The charge is not treated as an individual particle attribute. This study employs 2D simulation in flow-coupled model that captures aerosol microphysics in warm cumulus case without relying on subgrid dynamics process, we assume droplets are always with opposite charge and well mixed in the cloud and charge is not a particle attribute in the simulation. To assess fluctuation effects, we conducted 50 simulations with varying pseudo-random number sequences for each electro-coalescence treatment. The results show that with CS treatment, the electrostatic force contributes a larger effect on cloud evolution than previous research. With lower charge limit of ~~of~~ the maximum charge amount ~~on~~ of the droplet, the domain total precipitation with CS treatment for droplets with opposite signs is higher than that with the no charge (NC) setting. Compared to previous work, the multi-image-dipole treatment of CS results in higher precipitation. ~~Compared with previous work, with the multi image dipole treatment of CS, the amount of precipitation is~~

15
20
25
30
35

~~higher.~~ It is found that the electro-coalescence effect could affect rain formation even when the droplet charge is at the lower charge limit. High pollution levels result in greater sensitivity to electro-coalescence. The results show that when the charges ratio between two droplets is over 100, the short-range attractive electric force due to the multi-image dipole would also significantly enhance precipitation for the cumulus. It is indicated that although the accurate treatment of the electrostatic force with CS method would require 30% longer computation time than before, it is worthwhile to include it in cloud, weather, and climate models.

45 1 Introduction

Clouds are considered to play a key role in climate systems, and the collision-coalescence of cloud droplets plays a key role in ~~cloud-rain~~ formation. Droplet coalescence is one of the main processes leading to precipitation ~~and even cloud chemistry~~, affecting cloud microphysics and thereby changing the global radiation budget (Pruppacher and Klett, 2010, Chapter 15; Grabowski and Wang, 2013; IPCC AR6 WG1 Ch7, 2021). Several studies have reported that the electrostatics force on charged droplets could significantly influence the droplet coalescence and droplet-aerosol coagulation in weakly electrified clouds (Rayleigh, 1879; ~~Pruppacher and Klett, 2010~~; Tinsley et al., 2001, 2006; Zhou et al., 2009; Tripathi et al., 2008; Pruppacher and Klett, 2010; Zhang et al., 2018; Guo and Xue, 2021). This electrostatic force induced effect is called electro-coalescence or electro-anti-coalescence (Tinsley, 2008) and could even explain the link between solar wind fluctuations and changes in atmospheric parameters, such as cloud cover, polar surface pressure and the effective radiation in polar regions (Kniveton et al., 2008; Lam et al., 2014; Frederick & Tinsley 2018; Frederick et al., 2019).

In weakly electrified clouds, the accumulation of space charges on droplets is controlled by the diffusion of atmospheric ions produced by the cosmic ray flux, and the concentration is dependent on the ratio of attachment and recombination and the downward ionosphere-earth current density (J_z). When the J_z penetrates the cloud, the gradients of the electric field at the cloud boundary could generate net positively charged droplets at the upper cloud boundary and net negatively charged droplets at the lower boundary (Zhou and Tinsley, 2007; Nicoll and Harrison, 2016). The observations of Beard et al. (2004) revealed that with a J_z of 1-6 pAm⁻² in stratocumulus and altostratus clouds, a cloud droplet with the radius of 10 ~~um~~ ~~um~~ can accept approximately 100 elemental charges, which is consistent with the theoretical calculations by Zhou and Tinsley (2007). In cumulus clouds, vertical convection causes positive charge droplets from the upper boundary and negative charge droplets from the lower boundary to mix, leading to electro-coalescence. ~~In the cumulus, with vertical convection, the charged droplets at the boundaries can be mixed, droplets with opposite sign charged affect by electro-coalescence.~~ The maximum charge on the droplets is determined by the air breakdown voltage for corona discharge (Meek and Carggs, 1953) and is a quadratic function of the droplet radius (Khain et al., 2004; Andronache, 2004).

Numerous studies have focused on parameterizing the microphysics of the electro-coalescence of particles, the challenge is to approximate calculation one the electrostatic force between charged droplets.

In the 1970s, the collision efficiency of oppositely charged droplets evaluated with a centered Coulomb force indicated that only in strongly electrified clouds can the charge on droplets significantly affect cloud droplet coagulation (Wang et al., 1978). ~~The trajectory simulation studies by Tinsley et al. (2001, 2006), Tripathi et al. (2006), and Zhou et al. (2009)~~The series of trajectory simulation work by Tinsley et al. (2001, 2006), Zhou et al., (2009) and Tripathi et al., (2006) revealed that in a weakly electrified cloud, when taking into account the image charge force, the collision rate coefficient between the charged droplets could be different. Even with droplet charges of the same sign, the collision rate coefficient could be enhanced as a function of the charge on the particles with radii ranging from 0.1 ~~μm microns~~ to 10 ~~μm microns~~ (Zhou et al., 2009). The so-called Greenfield gap, identified by Greenfield (1957), describes the reduced concentrations of particles in the 0.1 to 1 ~~μm micrometer~~ size range. Greenfield gap could be ~~eliminated-reduced~~ with sufficient charging of the droplets. Simulation results showed that for particles with radii smaller than 0.1 ~~μm microns~~, when the particles obtain a large charge due to the evaporation of highly charged droplets, the collision rate coefficient is significantly decreased due to the repulsive electric force of droplets with charges of the same sign and is increased for charges of the opposite sign (Tinsley and Leddon, 2013). The updated simulation by Zhou et al. (2009) with an exact electric force treatment with the conducting sphere (CS) method indicated that the collision efficiency is a factor of two higher in the Greenfield Gap than that from the results of single image charge (IM) treatment. A few laboratory experiment results were consistent with these theoretical simulations (Ardon-Dryer et al., 2015). These findings highlight the need to represent coagulation due to droplet and aerosol charges in the cloud model. Khain et al. (2004) (hereafter Khain04) conducted a 0-dimension simulation to study the effect of seeding charged droplets on a cumulus cloud using the spectral bin cloud model with a 4-dimensional (mass and charging rate of two droplets) collision efficiency lookup table based on the static electric force between charged droplets. The results showed a significant response in the evolution of clouds due to charged droplets. ~~Khain04 set a charging rate equal to 5% of the maximum charge of natural droplets, which is 2.5 times larger than the values reported by Zhou et al. (2007), to study electro-coalescence impact on rain enhancement and fog elimination. -5% of maximum charge amounts of natural droplets, which is 2.5 times larger than the results from Zhou et al. (2007), was used in Khain04 to investigate their influence on rain enhancement and fog elimination.~~ Andronache (2004) and Wang et al. (2015) claimed that charged droplets significantly contribute to below cloud scavenging according to the analytical formula suggested by Davenport and Peters (1978), where the minimum amount of charge on droplets is 7% of the maximum limit. However, only Coulomb force (CB) treatment was used in Andronache (2004)'s and Wang et al. (2015)'s simulations.

Lagrangian particle-based approaches accurate solutions for the collision-coalescence process compared to bin microphysics schemes, as they overcome the limitations imposed by the assumptions of bin schemes (Grabowski et al., 2019; Liu et al., 2023). In this study, we estimate the effect of electro-coalescence from J_z on warm cumulus clouds by an exact treatment of electric forces using the conducting sphere (CS) method, using the Super-Droplet Method (SDM), a Lagrangian particle-based cloud microphysics scheme.

~~In this work, the effect of the electro-coalescence from J_z on a warm cumulus with an exact electric force treatment with the CS method is estimated based on particle-based cloud modeling with the real-time~~

~~collision efficiency calculation using the super droplet method.~~ The lower charging rate threshold for electro-coalescence is discussed. The extreme assumption of the droplet charging scenario of opposite sign charge is investigated. The electro-anti-coalescence (Tinsley and Zhou 2015) between charged droplets and particles could also be important for deep convection and stratus cloud evolution, ~~and this will be addressed in future work.~~

2. Description of the cloud model

~~In this study, we assume that the charged droplets are well-mixed in the warm cumulus cloud and focus on the electro-coalescence effect. The particle based microphysics method, which calculates the electro collision-coalescence kernel in real time, offers more detailed insights into droplet behavior influenced by electrostatic forces, surpassing the bin method that relies on lookup tables (Khain et al., 2004), while also demanding less computational resources.~~ A Lagrangian particle-based cloud model is used with the particle size resolved treatment following the ~~Super Droplet Method~~SDM-(SDM) by Shima et al. (2009, 2020). ~~Compared to bin microphysics schemes, the Super-Droplet Method (SDM) eliminates numerical diffusion and provides more accurate solutions for well-mixed volumes (Grabowski et al., 2019). Despite its sensitivity to super-droplet initialization and a higher variance than observed in reality (Liu et al., 2023), SDM is well-suited for this study.~~ This section provides a description of the ~~super droplets method~~SDM, how we generalize the exact electric force treatment with the CS method approach for the cloud model, and the numerical simulation setup.

2.1 Definition of super-droplets

Super-droplets have been defined in detail by Shima et al. (2009, 2020). A super-droplet represents multiple droplets with the same attributes and position, and this multiplicity is denoted by the positive integer $\xi_i(t)$, which can be different in each super-droplet and is time-dependent due to the definition of coalescence. Then, each super-droplet has its own position $x_i(t)$ and its own attributes $a_i(t)$ that characterize the $\xi_i(t)$ identical droplets represented by super-droplet i . In this study, we assume that the attributes consist of the equivalent radius ~~of water~~ and the ~~ammonium bisulfate~~ mass in the droplet $a_i(t) = [R_i(t), M_i(t)]$. Since each real droplet takes different positions and attributes, a super-droplet is a kind of coarse-grained view of droplets both in real space and attribute space. Assume that $N_s(t)$ is the number of super-droplets in the domain at time t . Then, the super-droplets represent $N_r(t) = \sum_{i=1}^{N_s(t)} \xi_i(t)$ real droplets in total.

2.2 Motion of a super-droplet

The advection and sedimentation ~~processes was/were~~ described in detail by Shima et al. (2009, 2020) as follows:

$$\frac{d(m_i v_i)}{dt} = F_i^{drg} - m_i g \hat{z} \quad (1)$$

150 where $m_i = (4\pi/3) R_i^3 \rho_{liq}$ is the mass of droplet i , and $\rho_{liq} = 1.0 \text{ g} \cdot \text{cm}^{-3}$ is the density of liquid water. $F_i^{drg} = m_i g \hat{z} + d(m_i v_i)/dt$ is the drag force from moist air, g is the gravity of Earth, and \hat{z} is the unit vector in the direction of the z-axis. $-F_i^{drg}$ gives the reaction force acting on the moist air (Montero-Martínez et al. 2009). Considering that the relaxation to the terminal velocity is instantaneous, and the equation of motion becomes:

$$155 \quad v_i = U_i - \hat{z} v_i^\infty, \frac{dx_i}{dt} = v_i \quad (2)$$

where $U_i = U(x)$ is the ambient wind velocity of the i -th particle, and v_i^∞ is the terminal velocity, which in general is a function of the attributes a_i and the state of the ambient air.

The motion of a super-droplet is the same as that of a droplet, which is described in equation (2), and $v_i(t)$ is equal to the terminal velocity.

160 2.3 Condensation and evaporation

The condensation/evaporation process is based on Köhler's theory, which takes into account the solution and curvature effects on the droplet's equilibrium vapour pressure (Köhler 1936; Rogers & Yau 1989; Pruppacher and Klett, 2010, Chapter 13; Yau & Rogers 1989). The growth equation of radius R_i is derived as follows:

$$165 \quad R_i \frac{dR_i}{dt} = \frac{(S-1) - \frac{a}{R_i} + \frac{b}{R_i^3}}{F_k + F_d} \quad (3)$$

$$F_k = \left(\frac{L}{R_v T} - 1 \right) \frac{L \rho_{liq}}{KT}, F_d = \frac{\rho_{liq} R_v T}{De_s(T)} \quad (4)$$

where S is the ambient saturation ratio; F_k represents the thermodynamic term associated with the latent heat release; F_d represents the term associated with vapour diffusion; the term a/R_i represents the curvature effect, which expresses the increase in the saturation ratio over a droplet compared with that of a plane surface; the term b/R_i^3 represents the reduction in the vapour pressure due to the presence of a dissolved substance, where b depends on the mass of solute M_i dissolved in the droplet; $a \approx 3.3 \times 10^{-5} \text{ cm K/T}$ and $b \approx 4.3 \text{ cm}^3 i M_i / m_s$, where T is the temperature, $i \approx 2$ is the degree of ionic dissociation and m_s is the molecular weight of the solute. R_v is the individual gas constant for water vapour, K is the coefficient of thermal conductivity of air, D is the molecular diffusion coefficient, L is the latent heat of vaporization, $e_s(T)$ is the saturation vapour pressure. Note the charge-induced reduction in surface tension decreases the equilibrium vapor pressure (Weon & Je 2010).

175

2.4 Collision-coalescence and the electric effect

In warm clouds, the collision-coalescence of two droplets to form a larger droplet is responsible for precipitation and ~~the~~ cloud lifetime. The droplet growth due to the coalescence is controlled by the net action of various forces impacting the relative motion of the two droplets. The effective collision-coalescence of droplets can be evaluated by the collision-coalescence kernel K , which can be described as follows:

$$K = \pi E \left((R+r)^2 |v_R^\infty - v_r^\infty| \right) + K_B \quad (5)$$

where $E = E_0(R+r) + E_{es}(R, Q_R, r, q_r)$ is the collision-coalescence efficiency, and K_B is the Brownian coagulation kernel. Respectively, R represents the radius of the larger droplet, r is the radius of smaller droplets of given pair (R, r) . Similarly, Q_R represents the charge of larger droplets, q_r is the charge of smaller droplets. And v_R represents the terminal velocity of larger droplets, v_r is the terminal velocity of smaller droplets. In this study, we assume

$E_0(R, r)$ takes into account the effect of a small droplet/particle being swept by the stream flow around a larger droplet or bouncing on the surface by front, side or rear collection, or droplets of similar size collide on the downstream side and are caught (Davis, 1972; Hall, 1980; Jonas, 1972; Pruppacher and Klett, 2010, chapter 14). Following Seeßelberg et al. (1996) and Bott (1998), the collision efficiency of Davis (1972) and Jonas (1972) for small droplets and the collision efficiency of Hall (1980) for large droplets are adopted. We assume the coalescence efficiency is unity in this study.

The Brownian coagulation kernel K_B is given by Seinfeld and Pandis (2006, chapter 13) using Fuchs (1964) corrected factor to correct the boundary condition of absorbing particles surface. The Fuchs Form of Brownian coagulation coefficient is derived as follows:

$$K_B = 2\pi(D_1 + D_2)(D_{p1} + D_{p2}) \left(\frac{D_{p1} + D_{p2}}{D_{p1} + D_{p2} + 2(g_1^2 + g_2^2)^{1/2}} + \frac{8(D_1 + D_2)}{(\bar{c}_1^2 + \bar{c}_2^2)^{1/2}(D_{p1} + D_{p2})} \right)^{-1} \quad (6)$$

Where:

$$\bar{c}_i = \left(\frac{8kT}{\pi m_i} \right)^{1/2} \quad (7)$$

$$\ell_i = \frac{8D_i}{\pi \bar{c}_i} \quad (8)$$

$$g_i = \frac{\sqrt{2}}{3D_{pi}\ell_i} \left[(D_{pi} + \ell_i)^3 - (D_{pi} + \ell_i^2)^{3/2} \right] - D_{pi} \quad (9)$$

$$D_i = \frac{kTC_c}{3\pi\mu D_{pi}} \quad (10)$$

ℓ_i represents particle mean free path, D_i represents Brownian diffusivity; D_{pi} represents diameters of particles, m_i is particle mass, $k = 1.381 \times 10^{-23}$ J/K is the Boltzmann constant, μ represents the dynamic viscosity ~~viscosity rate~~ of air and C_c is a slip correction factor.

~~In this study~~ Referring to Andronache (2004), we propose a parameterization of the collision efficiency due to the electric force $E_{es}(R, Q_R, r, q_r)$ based on the work by Zhou et al. (2009) and Tinsley and Zhou (2015). The induced charge on the droplet is involved in our E_{es} . Based on the trajectory model simulation, the electric force with the IM treatment (Tinsley et al., 2006) and the CS treatment (Zhou et al., 2009) can significantly contribute to the collision efficiency. For droplets with opposite sign charges, in the front and side collision ranges, the short-range attractive electric force due to the induced image charge provides additional force to balance the repulsive force. In the rear collision range, this short-range attractive force contributes to balancing the inertia. The rear collision range is relevant for droplets smaller than 0.1 micrometers, the droplets typically accept fewer than 1 elemental charge, meaning the electric force does not significantly impact the collision process. the droplet accepts less than 1 elemental charge and the electric force does not have a significant effect on the collision process. Therefore, the main electric force remains in the side and front collision range and droplets accept more than 1 elemental charge in this study.

The analytical parameterization for the collision efficiency with the electric force suggested by Davenport and Peters (1978) is used with modification to include the image charge effect of opposite charged droplets in our study. Tinsley and Zhou (2015) developed same charged droplets charge effect.

$$E_{es} = \frac{4c_f}{6\pi\mu Rv} \cdot F_{es} \quad (11)$$

where c_f is the Cunningham correction factor, ~~μ_a is the air viscosity, and v is the terminal velocity~~ of the droplet. F_{es} is the electric force between the colliding droplets.

In this study, F_{es} is calculated in four different ways, namely, CB, IM, Khain04, and CS, which are given by eqs. (13-16), respectively. ~~The distance parameter r_{nt} is needed to calculate F_{es} . Based on the trajectory simulation results by Zhou et al. (2009), r_{nt} is fitted as follows:~~

$$r_{nt} = \frac{r}{R} \left[1 + r_{ref} \cdot \left(\frac{R}{r/2} \right) + \frac{R}{r} \right] \quad (12)$$

~~where $r_{ref} = 0.01$, R and r represents the radius of larger and smaller droplet of a pair droplets. Q_R and q_r are charge of large and small particles~~

CB treatment considers only the Coulomb force between the centre points of the droplets. Then, F_{es} is given by

$$F_{es} = \frac{1}{4\pi\epsilon_0} \frac{Q_R q_r}{R_b^2} \quad (13)$$

Where $\epsilon_0 = 8.854 \times 10^{-12} \text{ Fm}^{-1}$ is the dielectric permittivity of free space. Q_R and q_r are charge of large and small particles. R_b is the distance between the centre of two droplets.

Khain04 used the superposition method to calculate a four-dimensional (with respect to droplet size and charge) lookup table for collision efficiency, and present an approximated solution for electrostatic forces of droplet by following formula:

$$F_{es} \approx \frac{Q_R q_r}{4\pi\epsilon_0 R_b^2} + \frac{1}{4\pi\epsilon_0} \left\{ Q_R^2 \left[\frac{1}{R_b^3} - \frac{R_b}{(R_b^2 - r^2)^2} \right] + q_r^2 R \left[\frac{1}{R_b^3} - \frac{R_b}{(R_b^2 - R^2)^2} \right] \right. \\ \left. + Q_R q_r R r \left[\frac{1}{R_b^4} + \frac{1}{(R_b^2 - R^2 - r^2)^2} - \frac{1}{(R_b^2 - R^2)^2} - \frac{1}{(R_b^2 - r^2)^2} \right] \right\} \quad (143)$$

R and r represents the radius of larger and smaller droplet of a pair droplet. –Note that in this study, we calculate collision-coalescence kernel of Khain04 method by eq (134) for electrostatic forces and eq (11) for charge effect, whereas Khain04 used a 4-dimensional lookup table for collision efficiency.

The distance parameter r_{nt} is needed to calculate F_{es} of IM and CS treatment. Based on the trajectory simulation results by Zhou et al. (2009), r_{nt} is fitted as follows:

$$r_{nt} = \frac{r}{R} \left[1 + r_{ref} \cdot \left(\frac{R}{r/2} \right) + \frac{R}{r} \right] \quad (124)$$

where $r_{ref} = 0.01$.

When the large droplet radius is 100 times larger than the small droplet radius, the IM treatment, is accurate enough. F_{es} for IM treatment is given by:

$$F_{es} = \frac{4Cq_r^2}{(d_p/2)^2} \left[\frac{r_{nt}}{(r_{nt}^2 - 1)^2} + \frac{1}{r_{nt}^2} \cdot \left(\frac{Q_R}{q_r} - \frac{1}{r_{nt}} \right) \right] \quad (15)$$

where $C = 9 \times 10^9$ (in Nm^2C^{-2}).

If the ratio between the droplet and particle is less than 100, the electric force is treated by the CS method according to Zhou et al. (2009), which originates from Davis (1964a, 1964b):

$$F_{es} = 4C \cdot \left(q_r Q_R \frac{F_6}{r^2} + q_r^2 \frac{F_7}{r^2} + Q_R^2 \frac{F_5}{r^2} \right) \quad (16)$$

where F_5 , F_6 , and F_7 are dimensionless complex polynomial expressions given by Davis (1964a, 1964b) that depend only on the radii of the two droplets and their distance parameter r_{nt} .

In this study, we assume that J_z charges the droplets. Zhou and Tinsley (2012) observed that droplets with a 10 μm radius achieve 70% of their charge in 680 seconds. However, following Andronache (2004) simplification of the complex charging process, we assume that the charge on droplets resulting from collision-coalescence reaches equilibrium instantaneously. We also consider an extreme scenario where the charge polarity of two colliding droplets is always opposite. The assumption of instantaneous charging might lead to an overestimation of the electro-coalescence effect. Although the charging process of droplets of 10 μm radius completes 70% in 680s (Zhou & Tinsley 2012), in this study, we consider that the charge amount of the resultant droplets after collision-coalescence relaxes to their equilibrium value instantaneously. Regarding the charge polarity, convective mixing within cumulus clouds introduces oppositely charged droplets from the cloud boundary into the cloud interior. These droplets retain their opposite charges due to the relatively long discharge timescale, significantly impacting the early stages of raindrop formation after convective mixing of inner cumulus clouds, the opposite charge droplets from the cloud boundary get into the cloud and keep opposite charge amount because of the relatively long timescale of discharge, with a significant impact on the early stage of rain droplets

~~formation. The coalescence of large rain droplets is dominated by gravity settling. We will consider the extreme case that the charge polarity of the two droplets is always opposite. Notably, Khain (2004) addressed charge differences by subtracting the charge of opposite polarity particles and adding the charge of same polarity particles after collision-coalescence. Note that Khain04 subtracted the charge amount of opposite polarity particles after collision-coalescence, added the charge amount of same polarity particles respectively.~~

The voltage near a charged spherical particle is described by $U = q / 4\pi\epsilon_0 r^2$ (Bleaney and Bleaney 1993); here $\epsilon_0 = 8.854 \times 10^{-12}$ is the dielectric permittivity of free space. The air breakdown voltage: The maximum charge that cloud droplets can carry is determined by the air breakdown voltage $U_b \sim 3 \times 10^6 \text{ Vm}^{-1}$, determines the maximum charge that cloud droplets can carry (Meek and Craggs, 1953). (Meek and Craggs 1953). Consequently, the maximum charge that droplets can carry is as follows:

$$q_{\max} = 4\pi U_b \epsilon_0 r^2 \quad (17)$$

~~To simulate droplets in a weak electric field, we followed Andronache (2004) and described the mean charges on the larger and smaller droplets in a pair as a function of their radii as follows: To simulate droplets in weakly electric field, we following Andronache (2004), described the mean charges on the larger droplet and smaller droplet in a pair of droplets as a function of the droplet radii as follows:~~

$$Q_R = 4\alpha AR^2, q_r = 4\alpha Ar^2 \quad (18)$$

Here $A = \pi \times U_b \times \epsilon_0 \times 10^{-2} = 0.83 \times 10^{-6}$ is two orders of magnitude smaller than the maximum particle charge, representing weakly charge condition, and the charging rate α is an empirical parameter (α is referred to herein as the droplet charging rate) that varies between 0, which represents neutral particles, and 7, which represents highly electrified clouds associated with thunderstorms (Andronache, 2004). In our work, the α value ranges from 0.1 to 0.6, which represents a weakly electrified cloud.

Compared with the maximum charge of the droplet method used by Khain04, when α ranges from 0.1 to 0.6 cm^{-2} , the charge on the droplet reaches 0.3% up to 2% of the maximum charge, which is 10 times to 2 times smaller than the lowest value used by Khain04 and Wang et al. (2015). The minimum limit for the droplet charge is equivalent to 1 elemental charge. This could be a reliable estimation for the accumulated charge on droplets with the downward current density (J_z) since a droplet with a radius of 10 μm can accept 200 elemental charges when $\alpha=0.1$, which is consistent with the stratus cloud charge distribution simulation by Zhou and Tinsley (2007, 2012).

Figure 1 displays a comparison of the collision-coalescence kernel for droplet radii of 40 μm (black lines), 20 μm (green lines), and 10 μm (red lines) across different calculation methods. The plots vary by line style to represent different analytical treatments and the inclusion or absence of static electric forces, with specific settings for the droplet charging rate shown in Figures 1(a) and 1(b). The results indicate that the primary range for electro-coalescence is approximately 0.1 μm to 10 μm , which encompasses the Greenfield Gap. The results show that the main electro-coalescence range is approximately 0.1 μm to 10 μm . When the small droplet radius is less than 0.1 μm , the collision process is controlled by Brownian motion due to an excessively small number of charges on the small droplet. Conversely, when the radius of the droplets exceeds 10 μm , the collision process is primarily governed by

~~droplet is larger than 10 μm , the collision process is controlled by~~ gravity collision. The electric force has a larger effect on the smaller droplet. The electric force treated with the CS method has a larger effect on the collision-coalescence kernel than that of the IM method and the CB method. In the range of the Greenfield gap, the collision-coalescence kernels from the analytical method fit well with those with the trajectory method result. Note that CB, Khain04 and IM method does not take into account the collision of same size droplets, for CS method, the Q^2 term provide attractive or repulsive force between same size droplets ensures collision. For the range of droplets smaller than 10 μm , when the particle radius is close to 10 μm , CB, Khain04 and IM method deviates from the trajectory result, but the result of the CS method becomes over 2 times less than that of the trajectory method, where the collision process is controlled by the interception effect. The interception effect in particle collision coalescence refers to the process where smaller particles are captured by a larger droplet's boundary layer and swept into it, even without direct contact, due to the aerodynamic airflow around the falling droplet. Although the analytical method cannot reproduce the interception effect, it can give the lower limit of estimation to the effect of the electric force effect with the conducting sphere method.

~~2.5 Fluid dynamics of moist air~~

~~In our model, the warm cloud dynamics is described by the fully compressible non-hydrostatic equation as follows:~~

2.5 Numerical setup and schemes

~~Shima et al. (2009, 2020) constructed a particle-based cloud model SCALE-SDM by implementing the SDM into SCALE, which is a library of weather and climate models of the Earth and other planets (Nishizawa et al., 2015; Sato et al., 2015). Because of its efficient Monte Carlo algorithm for coalescence, the particle-base scheme SDM requires less computational cost to accurately simulate clouds and precipitation compare to bin scheme (Shima et al., 2009). This study concentrates on warm-rain microphysics. We developed a numerical simulation using the latest version of SCALE-SDM, specifically employing the SDM warm rain algorithm from Shima et al. (2009), rather than the SDM mixed-phase extension presented by Shima et al. (2020). We implemented the electro-coalescence process into SDM's coalescence scheme as defined by eqs. (5)-(18). The moist air fluid dynamics in this study are computed using eqs. (71)-(81) of Shima et al. (2020). The calculations utilize SCALE's dynamical core, employing a forward temporal integration scheme. This approach is implemented on an Arakawa-C staggered grid (Arakawa and Lamb, 1977) using a finite volume method. For the initialization of the super-particle, the "uniform sampling method" is applied as in previous works (Arabas and Shima, 2013; Shima et al., 2014, 2020; Sato et al., 2017, 2018). Unterstrasser et al. (2017) found that the uniform sampling method is more efficient than the "constant multiplicity method". Then, the multiplicity of the super-droplets becomes proportional to the initial distribution function of real particles:~~

$$\xi(a, x) = n(a, x, t = 0) / (N_s(0)p) \quad (19)$$

$$p(a, x) = p = \text{constant} \quad (20)$$

In SCALE-SDM, moist air dynamics and cloud microphysics processes are integrated separately by using the 1st-order operator splitting scheme. Δt is set as the common time step. We set Δt_{adv} , $\Delta t_{end/evp}$, and Δt_{coal} as the time steps for the advection and sedimentation of particles, condensation/evaporation, and collision-coalescence. We set Δt_{dyn} as the time step for the fluid dynamics of moist air, which has to fulfil the Courant-Friedrichs-Lewy (CFL) condition of acoustic waves. All these time steps are divisors of the common time step Δt . The order of calculation in the model is as follows: 1) calculate the fluid dynamics without the coupling terms from the particles to moist air, and update the moist air; 2) update the super-droplets $\{\{\zeta_i, x_i, a_i\}\}$ from t to $t + \Delta t$. 3) We integrate one cloud microphysics process one time step forward and then moves on to the next process. Processes with shorter timescales are prioritized in the computation to ensure they accurately capture their subsequent impacts. Simultaneously, the feedback from particles to moist air comes through the coupling terms of Eqs. (75)-(79) of Shima et al. (2020), and we update the moist air from $G_{lim}(t)$ to $G_{lim}(t + \Delta t)$.

In our simulation, the domain of the simulation is two dimensional (x-z), 10 km in the horizontal and vertical directions with 50 m grid spacing, and the calculation time steps are $\Delta t = 0.4$ s, $\Delta t_{dyn} = 0.05$ s, $\Delta t_{adv} = 0.4$ s, $\Delta t_{end/eva} = 0.1$ s, and $\Delta t_{coal} = 0.2$ s. Initial super droplet number concentration per grid cell is 128. We employ a subgrid-scale (SGS) turbulence model for dynamic processes but exclude it for cloud microphysics processes, such as collision-coalescence enhancement, velocity fluctuations, and supersaturation fluctuations. This approach might lead to an underestimation of the collision rate of charged droplets, as noted by Lu and Shaw (2015). Our simulations use two-dimensional Large Eddy Simulation (LES) methodology. To assess the impact of fluctuations, we conduct a 50-member ensemble of simulations, varying the pseudo-random number sequence for each run.

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho U) = \frac{\partial \rho}{\partial t} \Big|_{cm} \quad (19)$$

$$\frac{\partial \rho q_v}{\partial t} + \nabla \cdot (\rho q_v U) = \frac{\partial \rho q_v}{\partial t} \Big|_{cm} + D_v \nabla^2 (\rho q_v) \quad (20)$$

$$\frac{\partial \rho U}{\partial t} + \nabla \cdot (\rho U \otimes U) = -\nabla P - \rho g \hat{z} + \frac{\partial \rho U}{\partial t} \Big|_{cm} + \mu \nabla^2 (\rho U) \quad (21)$$

$$\frac{\partial \rho \theta}{\partial t} + \nabla \cdot (\rho \theta U) = \frac{\partial \rho \theta}{\partial t} \Big|_{cm} + \frac{\kappa}{c_p} \nabla^2 \theta \quad (22)$$

$$P = \rho R T = P_0 \left(\frac{\rho \theta R}{P_0} \right)^{c_p / (c_p - R)} \quad (23)$$

Here, $U = (u, v, w)$ represent wind velocity, ρ_d is density of dry air, ρ_v is density of water vapor, the density of moist air $\rho := \rho_d + \rho_v$, $q_v := \rho_v / \rho$ is specific humidity, $q_d := \rho_d / \rho$ is mass of dry

air per unit mass of moist air, T is temperature and P is pressure, $\theta := T/\Pi := T/(P/P_0)^{R/c_p}$ is
 380 potential temperature of moist air, where $P_0 = 1000$ hPa is the reference pressure;
 $R := q_d R_d + q_v R_v$, R_d , R_v are the gas constants of moist air, dry air and water vapor; ditto,
 $c_p := q_d c_{pd} + q_v c_{pv}$, c_{pd} , c_{pv} are the isobaric specific heats of moist air, dry air and water vapor. The
 model employed variable $G := \{U, \rho, q_v, \theta, P, T\}$ to represent the state of moist air.

The four terms with the form $\partial / \partial t|_{cm}$ indicated cloud microphysics coupling terms.

385 $\partial \rho / \partial t|_{cm} = \partial \rho q_v / \partial t|_{cm}$ represents source of vapor:

$$\frac{\partial \rho}{\partial t}|_{cm} = \frac{\partial \rho q_v}{\partial t}|_{cm} = s_v \quad (24)$$

where s_v indicates the source of vapor through condensation/evaporation. s_v can be evaluated by
 the microphysics variables as follows:

$$s_v(x, t) = \sum_{i \in I_r(t)} \delta^3(x - x_i(t)) \frac{dm_i}{dt}|_{cnd/evp} \quad (25)$$

390 Here $\delta^3(x)$ represent the three dimensional Dirac delta function, and the time derivatives for
 condensation/evaporation.

$$\frac{\partial \rho \theta}{\partial t}|_{cm} = -\frac{L_v s_v}{c_p \Pi} \quad (26)$$

$\partial \rho U / \partial t|_{cm}$ represents the drag force from the particles, as mentioned in 2.3:

$F_i^{drg} = m_i g \hat{z} + d(m_i v_i) / dt$. The value of second term of F_i^{drg} is much smaller than first term:

$$\begin{aligned} \frac{\partial \rho U}{\partial t}|_{cm} &= -\sum_{i \in I_{c(t)}} \delta^3(x - x_i(t)) F_i^{drg} \\ &\approx -\left[\sum_{i \in I_{r(t)}} \delta^3(x - x_i(t)) m_i(t) \right] g \hat{z} \end{aligned} \quad (27)$$

2.6.6 Design of our numerical experiment

To evaluate the effect of electro-coalescence on warm clouds, 2D simulation of an isolated cumulus is
 performed following the setup of Lasher-Trapp et al. (2005). Note the original study of Lasher-Trapp et
 400 al. (2005) was conducted in 3D, but 2D simulation is used in this study to save computational resources.
 The initial profile of the atmosphere is horizontally uniform. The vertical profile of the moist air is given
 by 1545 UTC 22 July sounding data from the Small Cumulus Microphysics Study (SCMS) in Florida.
 The cloud base is steady at 1050 m, and the maximum cloud top height is 5350 m. As suggested by
 Lasher-Trapp et al. (2005), wind shear is assumed to be absent, and random velocity perturbation is
 405 applied (maximum of 0.5 ms^{-1}) in the lowest kilometre of the model.

In general, there are different types of soluble/insoluble aerosols in a droplet. In the model, only one
 soluble substance ($(\text{NH}_4)\text{HSO}_4$ aerosol) is applied for simplicity. Initially, the aerosols are uniformly
 distributed in the simulation domain. The aerosol number concentration and size distribution were based
on the data provided by Van Zanten et al. (2011) for the RICO intercomparison case. The aerosol number

410 ~~concentration and size distribution were adjusted to 3, 6 or 9 times from that given in Van Zanten et al. (2011) for RICO intercomparison case.~~ The aerosol number concentration and size distribution is given by a bimodal log-normal distribution: The particle number concentrations of the two modes are $N_1 = 90 \text{ cm}^{-3}$ and $N_2 = 15 \text{ cm}^{-3}$, respectively. Note that aerosol concentrations are multiplied by factors of 3, 6, or 9, depending on the aerosol background conditions. The geometric mean radii are
 415 $r_1 = 0.03 \text{ }\mu\text{m}$ and $r_2 = 0.14 \text{ }\mu\text{m}$, with geometric standard deviations of $\sigma_1 = 1.28$ and $\sigma_2 = 1.75$, respectively.

2.7 Numerical setup and schemes

420 ~~Shima et al. (2009, 2020) constructed a particle-based cloud model SCALE-SDM by implementing the SDM into SCALE, which is a library of weather and climate models of the Earth and other planets (Nishizawa et al., 2015; Sato et al., 2015; <https://scale.riken.jp>). The SDM code is not accessible through this site. Because of its efficient Monte Carlo algorithm for coalescence, the particle base scheme SDM requires less computational cost to accurately simulate clouds and precipitation compare to bin scheme (Shima et al., 2009). This study concentrates on warm rain microphysics. We developed a numerical simulation using the latest version of SCALE-SDM, specifically employing the SDM warm rain algorithm from Shima et al. (2009), rather than the SDM mixed phase extension presented by Shima et al. (2020). We implemented the electro-coalescence process into SDM's coalescence scheme as defined by eqs. (5)–(18). The moist air fluid dynamics are calculated by eqs. (19)–(27) using SCALE's dynamical core using a forward temporal integration scheme with an Arakawa-C staggered grid (Arakawa and Lamb, 1977) using a finite volume method.~~

425 ~~For the initialization of the super-particle, the “uniform sampling method” is applied as in previous works (Arabas and Shima, 2013; Shima et al., 2014, 2020; Sato et al., 2017, 2018). Unterstrasser et al. (2017) found that the uniform sampling method is more efficient than the “constant multiplicity method”. Then, the multiplicity of the super-droplets becomes proportional to the initial distribution function of real particles:~~

$$\xi(a, x) = n(a, x, t=0) / (N_s(0)p) \quad (28)$$

$$p(a, x) = p = \text{constant} \quad (29)$$

430 ~~In SCALE-SDM, moist air dynamics and cloud microphysics processes for aerosol/cloud/precipitation particles are integrated separately by using the 1st order operator splitting scheme. Δt is set as the common time step. We set Δt_{adv} , $\Delta t_{cnd/evp}$, and Δt_{coal} as the time steps for the advection and sedimentation of particles, condensation/evaporation, and collision-coalescence. We set Δt_{dyn} as the time step for the fluid dynamics of moist air, which has to fulfil the Courant-Friedrichs-Lewy (CFL) condition of acoustic waves. All these time steps are divisors of the common time step Δt . The order of calculation in the model is as follows: 1) calculate the fluid dynamics without the coupling terms from the particles to moist air, and update the moist air;~~

440
445

2) update the super droplets $\{\{C_i, x_i, a_i\}\}$ from t to $t + \Delta t$. 3) We integrate one cloud microphysics process one time step forward and then moves on to the next process. Lagging processes are prioritized in computational priorities to ensure their synchronization and accurately describe their subsequent impact, consistent with overall system dynamics. Simultaneously, the feedback from particles to moist air comes through the coupling terms of Eqs. (23)–(26), and we update the moist air from $G_{lim}(t)$ to $G_{lim}(t + \Delta t)$.

In our simulation, the domain of the simulation is two dimensional ($x-z$), 10 km in the horizontal and vertical directions with 50 m grid spacing, and the calculation time steps are $\Delta t = 0.4$ s, $\Delta t_{dyn} = 0.05$ s, $\Delta t_{adv} = 0.4$ s, $\Delta t_{end/eva} = 0.1$ s, and $\Delta t_{coal} = 0.2$ s. Initial super droplet number concentration per grid cell is 128. We are employing a subgrid scale (SGS) turbulence model for dynamic processes, but not for cloud microphysics processes such as collision-coalescence enhancement, velocity fluctuations, and supersaturation fluctuations. This approach may lead to an underestimation of the collision rate of charged droplets (Lu & Shaw 2015). The simulations conducted are based on the Large Eddy Simulation (LES) methodology. To evaluate the fluctuation effect, a 50 member ensemble of simulations is conducted by changing the pseudo random number sequence.

3. Results

3.1 The effect of charged droplets on cloud evolution

Figure 2 presents snapshots of cloud structures at 1500 seconds, 2100 seconds, and 2700 seconds from a single simulation, illustrating the temporal changes in the mixing ratios of cloud water and rainwater. The results show that with electro-coalescence by the CS setting (Figure 2(d)-(i)), the cumulus takes a shorter time to form rain droplets than with that by the CB and NC settings. Comparing Figure 2(d)-(f) and Figure 2(g)-(i) shows that the electric force with the CS setting has a much stronger impact on the cloud evolution than that with the CB setting. For the CS setting, there is heavy precipitation at 2700 s, while there is only haze for the CB and NC settings.

Figure 3(a) shows the time evolution of the domain and 50 ensembles averaged accumulated precipitation amount. Figure 3(b)-(d) shows the domain-averaged pathway, including the total liquid water pathway (b), rainwater pathway (c), and cloud water pathway (d). The grey shades indicate the standard error, which is also calculated from the 50 members of the ensemble. An unbiased estimator is used to calculate the standard deviation error. The results show that the accumulated precipitation amount in the CS setting is 52.5% higher than that in NC setting, 34.9% higher than CB setting, and 8.4% higher than IM settings. There is significant difference between the accumulated precipitation amounts in NC setting, CS setting, IM setting and CB setting. The initial precipitation time for four settings all start at 2100 seconds. However, the total liquid water path and cloud path of the CB and NC settings are significantly higher than those of the CS setting because higher precipitation eliminates cloud evolution.

Figure 4 presents the droplet mass density distributions during three stages of cloud development for the NC, CB, IM, and CS settings. Figure 4 shows the evolution of the mass density distribution of droplets for NC setting (solid line), CB setting (dashed line), IM setting (dotted line) and CS setting (dashed-dotted line) at 1500s, 2100s and 2700s. At these three stages, the droplet size distribution in the CS setting is much wider and rain droplets are much coarser than in the NC, IM and CB settings. At 2100 seconds, there are two mass density peaks of 10 μm and 1000 μm droplets for the NC, IM, CS, and CB settings, while the CS setting shows the highest mass density at 1000 μm . There are two mass density peaks of 10 μm droplets and 1000 μm droplets for the NC, IM, CS and CB settings, while for the CS setting have highest mass density of 1000 μm , which is consistent with the results of Figure 3.

3.2 The effect of charge on droplets

Figure 5 shows the results of droplet time evolution (a)-(c) and the water fraction path (d)-(f) for charging rate (α) is 0.1 (solid line), 0.2 (dashed line), and 0.6 (dashed-dotted line); the grey line represents the NC setting. The results show that there are no significant differences between the results of the NC and CS settings with the charging rate $\alpha=0.1$, which gives 0.3% of the maximum charge on the droplets. With the enhancement of the charge on the droplets, clouds can form more rapidly. When the charging rate $\alpha=0.6$, at 1500 second, there are larger droplets with radius over 1000 μm and even droplets sized 5000 μm . However, cloud elimination is faster in higher charging rate condition, at 2700 second, a lower charging rate condition results in a higher droplet mass density at a peak of approximately 1000 μm , which indicates that a higher charging rate results in a shorter lifetime of the cumulus cloud. The results of the domain water path, averaged over 50 ensembles. The results of the domain and 50 ensembles averaged water path in Figure 5(d)-(f) are consistent with those in Figures 5(a)-(c).

Figure 6 shows the domain and 50 ensembles averaged precipitation amount as a function of the droplets charging rate in the CS setting. Similar to the results in Figure 5, clouds with higher charging rate condition produce precipitation earlier than those under low charging rate conditions. With the enhancement of the charging rate, the precipitation amount at 3500 second does not simultaneously increase under all conditions. When the charging rate is 0.6, the final precipitation amount decreases due to more liquid and cloud water loss in the early stage of cloud formation. In Figure 6, the result of the CS setting charging rate α equal to 0.05, which is 0.16% of the maximum charge on the droplets, is given by the solid orange line, which the precipitation amount is 9.5% higher than that of the NC setting.

3.3 The effect of the aerosol concentration

Figure 7 illustrates the average precipitation amounts for 50 ensemble simulations under the CS setting, plotted as a function of aerosol concentrations. The results are shown for low (solid line), medium (dotted line), and high (dash-dotted line) aerosol concentrations, with a charging rate of 0.2. The grey lines represent the results of LA, MA and HA conditions with the NC setting. Under NC settings, the Twomey effect demonstrates that higher aerosol concentrations lead to smaller particle radii in clouds, reducing precipitation efficiency. Conversely, when electrostatic forces are introduced, these higher aerosol concentrations substantially enhance precipitation across different scenarios. Specifically, in high aerosol (HA) conditions, the precipitation enhancement reaches 782% over the no charge (NC) setting; for

medium aerosol (MA) conditions, it's 467% higher; and for low aerosol (LA) conditions, the increase is 110%. This illustrates the significant role electrostatic forces play in modulating cloud dynamics and precipitation responses to aerosol variations.

3.4 Comparison of different electrostatic force calculations

525 Figure 8 presents the domain-averaged precipitation amounts under different electrostatic force settings
with a charging rate of 0.3, illustrated by various line styles for each setting. The yellow dashed line
represents the result of droplets with opposite sign charges and the setting based on Khain04 method.
The upper limit of the charge on the small droplets is 50 elemental charges, the lower limit of the limit
of the charge on the droplets is 1 elemental charges and the orange dashed-dotted line represents the
530 special setting of only a large droplet charged with electrostatic force by the CS method (CS-q0). For the
CS, Khain04 and IM setting, precipitation increases at 2100 s, which is 300 s before the NC setting. The
domain and ensembles averaged precipitation amount with the CS setting is 52.5% higher than that with
the NC setting; with the Khain04 setting, it is 5.42% larger; with the CS-q0 setting, it is 9.6% larger; and
with the IM setting, it is 8.45% larger.

535 4. Discussion

When the two droplets move together and coalesce, there are three sites where collisions can occur, the
front, side and rear. The radius ratio between a large droplet and a small droplet (RARA) controls the
collision site, and when the radius of the small droplet is less than $0.1 \mu\text{m}$ and the RARA is larger than
100, the collision is a rear collision. For front and side collisions, in clouds where the droplet size is less
540 than $40 \mu\text{m}$ and the relative humidity is 100%, the droplet collision is controlled by the balance of the
Stokes drag of the air flow and electric force. The analytic expression in our work suggested by
Davenport and Peters (1978) can give a good estimation, especially for the Greenfield gap part, although
in the front and side collision regions when the RARA is close to 1, the additional contribution due to
the interception associated with the electric effect cannot be fully reproduced by this method. For the rear
545 collision, the flow drag, electric force and Brownian motion of the small droplet can impact the collision
process. In the present work, because the charge on the droplet varies as a function of the droplet radius,
there is less than 1 elemental charge on a droplet with a radius less than $0.1 \mu\text{m}$. Therefore, the electric
force does not have a significant effect on the collision process even for droplets of opposite signs, and
the Brownian collision efficiency could be good enough for estimation under these conditions. When the
550 amount of charge on a small droplet is over several elemental charges due to the evaporation of a large
droplet with a large amount of charge, the rear collision could be significantly affected by the electric
force (Tinsley and Leddon, 2013). The net attractive force of droplets with opposite signs increases the
collision efficiency, and the net repulsive force of droplets with the same sign decreases the collision
efficiency; this is called electro-anti-coalescence. As Tinsley (2001) mentioned, below the cloud bottom
555 boundary, there could be a highly charged nucleus or small droplet with tens to hundreds of elemental
charges due to the evaporation of a highly charged droplet. These highly charged small droplets or nuclei

could be moved into the cloud by upward air flow, which is not considered in this paper ~~and will be evaluated in our next paper.~~

In clouds, there are several ways to charge droplets, and in the cloud boundary, due to charging by the vertical electric current density (J_z) from the ionosphere to the ground surface, a droplet in the cloud top boundary accumulates a positive charge, and that in the cloud bottom boundary accumulates a net negative charge; this has been shown by simulations (Zhou and Tinsley, 2007, 2012) and field observations (Nicoll et al., 2016). With the charging rate of 0.05, there are on the order of 100 ~~403~~ elemental charges on a droplet with a radius of 10 μm , which is consistent with observation (Beard et al., 2004) and simulation (Zhou and Tinsley, 2007) results. Therefore, in the stratus cloud, most droplet collisions occur between droplets of the same sign or between one charged droplet and one uncharged droplet. Using the CS method, the additional electrostatic force due to the multiple image dipoles between the colliding droplets can be addressed, even if the droplets have the same sign charges or a small droplet is uncharged. Khain et al. (2004) evaluated the electro-coalescence effect on warm clouds rain enhancement and fog formation based on the image charge method from one induced dipole on each droplet by bin scheme. Zhou et al. (2009) claimed that when the RARA is close to 1, the collision efficiency calculated by the CS method, which treats the multiple induced dipoles on each droplet, is twice as large as is two factors larger than those calculated with the IM method. Therefore, for the Greenfield gap region and interception region, the evaluation of the charge effect with the CS method is more accurate. And the particle-based approach SDM provide superior performance than bin schemes (Li et al., 2017). In Figure 8, due to the additional induced image charge on droplets, the maximum averaged precipitation amount of the CS setting is 8.45% larger than that of the IM setting. Despite a 30% increase in computation burden, the CS method for electrostatic force should be incorporated into the cloud microphysics scheme. CS method provides superior numerical stability and accuracy in simulating charge droplet interactions, particularly for charge droplets of similar size. It could be suggested that the CS method for electrostatic force should still be involved in the cloud model, although its computation time is 30% longer. Khain et al. (2004) evaluated electro-coalescence at a low charging rate of 5% of the maximum charge on droplets. In our simulation, we tested charging rates (α) ranging from 0.05 to 0.6, equivalent to 0.15% to 1.8% of the maximum charge. At a charging rate of 0.3, the electric force evaluated by the CS method increased domain and ensemble-averaged precipitation by approximately 5.42% compared to the Khain04 setting. The results indicate that even with weak charging, the electro-coalescence effect significantly increases precipitation. ~~The previous work by Khain et al. (2004) evaluated electro-coalescence at the lowest charging rate of 5% of the maximum charge on droplets. In our simulation, a charging rate (α) of 0.05 up to 0.6 is tested, which represents 0.15% to 1.8% of the maximum charge on the droplet. With a charging rate (α) equal to 0.05, the electric force evaluated by the CS setting can increase the domain and ensembles averaged precipitation by approximately 9.5% compared to that of the NC setting, which provides the lower limit of the effect of electrostatic force on cloud formation.~~

Tinsley et al. (2001, 2006) and Zhou et al. (2009) claimed that the induced charge on droplets of the same sign could produce a short-range attractive electrostatic force that increases the collision efficiencies. The charge on the large droplets could exert an additional short-range attraction on the small droplet,

even if there is no charge or the same charge on the small droplets. However, for droplets of the same sign, the short-range electrostatic force has a significant effect only if the charge ratio between the large droplet and small droplet $Q:q$ is greater than 100 or $q:Q$ is greater than 1. For $Q:q$ ratios larger than 100, the additional image charge effect on the small droplet due to the large charged droplet controls the collision process. For $q:Q$ greater than 1, the additional image charge effect is due to the small charged droplet.

~~The particle-based approach SDM provides explicitly cloud-aerosol interaction simulation, such as the role of CCN in rain formation (Grabowski et al., 2019). According to our simulation results, the electro-coalescence effect on precipitation is sensitive to the aerosol concentration. An increase in the aerosol concentration decreases the effective radius by increasing the concentration of small droplets, which could have a significant impact on cloud formation (Twomey, 1974). According to our simulation, the effect of electro-coalescence is sensitive to the aerosol concentration.~~ With a high aerosol concentration, the average precipitation with an electric effect could be a factor of 4 higher than that of the NC condition. A much higher aerosol concentration corresponds to a more sensitive cloud response to the electrostatic force. Then, under high aerosol concentration conditions, a small variation in J_z could have a significant effect on cloud formation. Alternatively, in highly polluted clouds, placing a small number of charged aerosols or droplets accelerates rain enhancement due to electro-coalescence.

5 Conclusion

The electro-coalescence effect on a weakly electrified warm cumulus cloud was revisited. Assuming droplets with opposite signs are charged instantaneously by J_z , the amount of charge is determined by the size of the droplets. A new simulation with the exact treatment of the electrostatic force for opposite sign charge case based on the particle-based approach SDM provides a good estimation of the effect of electro-coalescence in the Greenfield gap region. In the simulation, droplets smaller than $0.1 \mu\text{m}$ are controlled by Brownian motion. The results show that for droplets of opposite signs with the same treatment of the electrostatic force, the cloud evolution can be significantly changed as a function of the arbitrarily prescribed charging rate $\propto \alpha$. The same sign charge droplets case (Tinsley and Zhou 2015) and charge amount prediction are necessary for accurate simulation, ~~we leave them for the future work.~~

Electro-coalescence has a larger impact on highly polluted warm cumulus clouds.

~~Cloud radiation feedback is one of the sources of uncertainty in the climate model (Zelinka et al. 2017). The electrostatic force effect parameterization for different cloud types should be indicated to improve climate model accuracy. This study reveals the electrostatic force effect on warm cumulus clouds, contributing to the parameterization of electrostatic microphysical processes. This indicates that the effect of the electrostatic force with exact treatment should be included in cloud, weather and even climate models to improve the simulation accuracy.~~

635 **Code and data availability.** The source code of SCALE is available from <https://scale.riken.jp>. Note
the SDM code is not accessible through this site. The source code of SCALE-SDM 0.2.5-2.3.0 and single
simulation data of $\alpha=0.3$ in four settings of NC, CB, IM, and CS are available from
<https://zenodo.org/records/11058066>. All the data used for this study can be reproduced by following the
instructions included in the above repository. The data are also deposited in local storage at the University
of Hyogo, Japan.

640

Author contributions. All the authors designed the model and numerical experiments. LM and SS
developed the model code and RZ performed the simulations. RZ prepared the manuscript with
contributions from all co-authors.

645

Competing interests. The authors declare that they have no conflicts of interest.

650

Financial support. This work was funded in part by the Strategic Priority Research Program of CAS
(Grant No. XDB 41000000), the National Science Foundation of China (41971020), MEXT KAKENHI
(grant no. 18H04448), JSPS KAKENHI (grant nos. 26286089, 20H00225, 23H00149), JST [Moonshot
R & D][Grant Number JPMJMS2286], China Scholarship Council (202106140113).

References

- Andronache, C.: Diffusion and electric charge contributions to below-cloud wet removal of atmospheric ultra-fine aerosol particles, *Journal of Aerosol Science*, 35, 1467-1482, 10.1016/j.jaerosci.2004.07.005, 2004.
- 655 Arabas, S. and Shima, S.-i.: Large-eddy simulations of trade wind cumuli using particle-based microphysics with monte carlo coalescence, *Journal of the Atmospheric Sciences*, 70, 2768-2777, 10.1175/JAS-D-12-0295.1, 2013.
- Arakawa, A. and Lamb, V. R.: Computational design of the basic dynamical processes of the UCLA general circulation model, *Methods in Computational Physics: Advances in Research and Applications*, 173-265, 10.1016/B978-0-12-460817-7.50009-4, 1977.
- 660 Ardon-Dryer, K., Huang, Y. W., and Cziczo, D. J.: Laboratory studies of collection efficiency of sub-micrometer aerosol particles by cloud droplets on a single-droplet basis, *Atmospheric Chemistry and Physics*, 15, 9159-9171, 10.5194/acp-15-9159-2015, 2015.
- 665 Beard, K. V., Ochs III, H. T., and Twohy, C. H.: Aircraft measurements of high average charges on cloud drops in layer clouds, *Geophysical Research Letters*, 31, 10.1029/2004GL020465, 2004.
- Bleaney, B. I., and Bleaney, B.: *Electricity and Magnetism, Volume 1*. Oxford University Press, 676 pp, 1993.
- Bott, A.: A flux method for the numerical solution of the stochastic collection equation, *Journal of the Atmospheric Sciences*, 55, 2284-2293, 10.1175/1520-0469(1998)055<2284:afmftn>2.0.co;2, 1998.
- 670 Craggs, J. M. M. a. J. D.: Electrical breakdown of gases, *Quarterly Journal of the Royal Meteorological Society*, 80, 282-283, 10.1002/qj.49708034425, 1954.
- Davenport, H. M. and Peters, L. K.: Field studies of atmospheric particulate concentration changes during precipitation, *Atmospheric Environment* (1967), 12, 997-1008, 10.1016/0004-6981(78)90344-X, 1978.
- 675 ~~Davis, M.: Two charged spherical conductors in a uniform electric field: Forces and field strength, RM 3860 PR, 1964a.~~
- Davis, M. H.: Two charged spherical conductors in a uniform electric field: Forces and field strength, *The Quarterly Journal of Mechanics and Applied Mathematics*, 17, 499-511, [10.1093/qjmam/17.4.499](https://doi.org/10.1093/qjmam/17.4.499), 1964b.
- 680 Davis, M. H.: Collisions of small cloud droplets: gas kinetic effects, *Journal of Atmospheric Sciences*, 29, 911-915, 10.1175/1520-0469(1972)029<0911:COCDG>2.0.CO;2, 1972.
- Frederick, J. E. and Tinsley, B. A.: The response of longwave radiation at the South Pole to electrical and magnetic variations: Links to meteorological generators and the solar wind, *Journal of Atmospheric and Solar-Terrestrial Physics*, 179, 214-224, 10.1016/j.jastp.2018.08.003, 2018.
- 685 Frederick, J. E., Tinsley, B. A., and Zhou, L.: Relationships between the solar wind magnetic field and ground-level longwave irradiance at high northern latitudes, *Journal of Atmospheric and Solar-Terrestrial Physics*, 193, 10.1016/j.jastp.2019.105063, 2019.
- 690 [Grabowski, W. W., Morrison, H., Shima, S.-I., Abade, G. C., Dziekan, P., and Pawlowska, H.: Modeling of Cloud Microphysics: Can We Do Better?, *Bulletin of the American Meteorological Society*, 100, 655-672, <https://doi.org/10.1175/BAMS-D-18-0005.1>, 2019.](https://doi.org/10.1175/BAMS-D-18-0005.1)

- Greenfield, S. M.: Rain scavenging of radioactive particulate matter from the atmosphere, *Journal of Atmospheric Sciences*, 14, 115-125, 10.1175/1520-0469(1957)014<0115:RSORPM>2.0.CO;2, 1957.
- Hall, W. D.: A detailed microphysical model within a two-dimensional dynamic framework: model description and preliminary results, *Journal of Atmospheric Sciences*, 37, 2486-2507, 10.1175/1520-0469(1980)037<2486:ADMMWA>2.0.CO;2, 1980.
- 695 Jonas, P. R.: The collision efficiency of small drops, *Quarterly Journal of the Royal Meteorological Society*, 98, 681-683, 10.1002/qj.49709841717, 1972.
- Khain, A., Arkipov, V., Pinsky, M., Feldman, Y., and Ryabov, Y.: Rain enhancement and fog elimination by seeding with charged droplets. part I: theory and numerical simulations, *Journal of Applied Meteorology*, 43, 1513-1529, 10.1175/JAM2131.1, 2004.
- 700 Kniveton, D. R., Tinsley, B. A., Burns, G. B., Bering, E. A., and Troshichev, O. A.: Variations in global cloud cover and the fair-weather vertical electric field, *Journal of Atmospheric and Solar-Terrestrial Physics*, 70, 1633-1642, 10.1016/j.jastp.2008.07.001, 2008.
- Köhler, H.: The nucleus in and the growth of hygroscopic droplets, *Transactions of the Faraday Society*, 32, 1152-1161, 10.1039/TF9363201152, 1936.
- 705 Fuchs, N. A.: *The Mechanics of Aerosols*, Revised and Enlarged, Pergamon Press Oxford, Oxford, 1964.
- Lam, M. M., Chisham, G., and Freeman, M. P.: Solar wind-driven geopotential height anomalies originate in the Antarctic lower troposphere, *Geophysical Research Letters*, 41, 6509-6514, 10.1002/2014GL061421, 2014.
- 710 Lasher-Trapp, S. G., Cooper, W. A., and Blyth, A. M.: Broadening of droplet size distributions from entrainment and mixing in a cumulus cloud, *Quarterly Journal of the Royal Meteorological Society*, 131, 195-220, 10.1256/qj.03.199, 2005.
- [Li, X.-Y., Brandenburg, A., Haugen, N. E. L., and Svensson, G.: Eulerian and Lagrangian approaches to multidimensional condensation and collection, *Journal of Advances in Modeling Earth Systems*, 9, 1116-1137, <https://doi.org/10.1002/2017MS000930>, 2017.](#)
- 715 [Liu, Y., Yau, M.-K., Shima, S.-i., Lu, C., and Chen, S.: Parameterization and Explicit Modeling of Cloud Microphysics: Approaches, Challenges, and Future Directions, *Advances in Atmospheric Sciences*, 40, 747-790, 10.1007/s00376-022-2077-3, 2023.](#)
- Lu, J. and Shaw, R. A.: Charged particle dynamics in turbulence: Theory and direct numerical simulations, *Physics of Fluids*, 27, 10.1063/1.4922645, 2015.
- 720 Meek, J. M. and Craggs, J. D.: Electrical breakdown of gases., *Quarterly Journal of the Royal Meteorological Society*, 80, 282-283, 10.1002/qj.49708034425, 1954.
- Montero-Martínez, G., Kostinski, A. B., Shaw, R. A., and García-García, F.: Do all raindrops fall at terminal speed?, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2008GL037111>, 2009.
- 725 Nicoll, K. A. and Harrison, R. G.: Stratiform cloud electrification: comparison of theory with multiple in-cloud measurements, *Quarterly Journal of the Royal Meteorological Society*, 142, 2679-2691, 10.1002/qj.2858, 2016.
- Nishizawa, S., Yashiro, H., Sato, Y., Miyamoto, Y., and Tomita, H.: Influence of grid aspect ratio on planetary boundary layer turbulence in large-eddy simulations, *Geoscientific Model Development*, 8, 3393-3419, 10.5194/gmd-8-3393-2015, 2015.
- 730

- Pruppacher, H. R. and Klett, J. D.: Microphysics of clouds and precipitation, Springer Dordrecht, 10.1007/978-0-306-48100-0, 2010.
- Sato, Y., Shima, S.-i., and Tomita, H.: A grid refinement study of trade wind cumuli simulated by a Lagrangian cloud microphysical model: the super-droplet method, Atmospheric Science Letters, 18, 359-365, 10.1002/asl.764, 2017.
- 735 Rayleigh, L.: The influence of electricity on colliding water drops, Proceedings of the Royal Society of London, 28, 405-409, <https://www.jstor.org/stable/113853>, 1878.
- ~~Rogers, R. R., Yau, M. K. and Yau, M. K.: A Short Course in Cloud Physics, Elsevier Science, 1989.~~
- 740 Sato, Y., Nishizawa, S., Yashiro, H., Miyamoto, Y., Kajikawa, Y., and Tomita, H.: Impacts of cloud microphysics on trade wind cumulus: which cloud microphysics processes contribute to the diversity in a large eddy simulation?, Progress in Earth and Planetary Science, 2, 23, 10.1186/s40645-015-0053-6, 2015.
- 745 Sato, Y., Shima, S.-i., and Tomita, H.: Numerical convergence of shallow convection cloud field simulations: comparison between double-moment ~~Eulerian eulerian~~ and particle-based ~~Lagrangian lagrangian~~ microphysics coupled to the same dynamical core, Journal of Advances in ~~ModelingModelling~~ Earth Systems, 10, 1495-1512, 10.1029/2018MS001285, 2018.
- Seeßelberg, M., Trautmann, T., and Thorn, M.: Stochastic simulations as a benchmark for mathematical methods solving the coalescence equation, Atmospheric Research, 40, 33-48, 10.1016/0169-8095(95)00024-0, 1996.
- 750 Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics: From Air Pollution to Climate Change, Wiley, 2006.
- Shima, S.-i., Hasegawa, K., and Kusano, K.: Preliminary numerical study on the cumulus-stratus transition induced by the increase of formation rate of aerosols, Low Temperature Science, 72, 249-264, <http://hdl.handle.net/2115/55063>, 2014. (In Japanese)
- Shima, S.-i., Sato, Y., Hashimoto, A., and Misumi, R.: Predicting the morphology of ice particles in deep convection using the super-droplet method: development and evaluation of SCALE-SDM 0.2.5-2.2.0, -2.2.1, and -2.2.2, Geoscientific Model Development, 13, 4107-4157, 10.5194/gmd-13-4107-2020, 2020.
- 760 Shima, S., Kusano, K., Kawano, A., Sugiyama, T., and Kawahara, S.: The super-droplet method for the numerical simulation of clouds and precipitation: a particle-based and probabilistic microphysics model coupled with a non-hydrostatic model, Quarterly Journal of the Royal Meteorological Society, 135, 1307-1320, 10.1002/qj.441, 2009.
- Tinsley, B. A.: The global atmospheric electric circuit and its effects on cloud microphysics, Reports on Progress in Physics, 71, 066801, 10.1088/0034-4885/71/6/066801, 2008.
- 765 Tinsley, B. A. and Leddon, D. B.: Charge modulation of scavenging in clouds: Extension of Monte Carlo simulations and initial parameterization, Journal of Geophysical Research: Atmospheres, 118, 8612-8624, 10.1002/jgrd.50618, 2013.
- Tinsley, B. A. and Zhou, L.: Initial results of a global circuit model with variable stratospheric and tropospheric aerosols, Journal of Geophysical Research, 111, D16205, 10.1029/2005jd006988, 2006.
- 770

Tinsley, B. A. and Zhou, L.: Parameterization of aerosol scavenging due to atmospheric ionization, *Journal of Geophysical Research: Atmospheres*, 120, 8389-8410, 10.1002/2014jd023016, 2015.

Tinsley, B. A., Rohrbaugh, R. P., and Hei, M.: Electroscavenging in clouds with broad droplet size distributions and weak electrification, *Atmospheric Research*, 59-60, 115-135, 10.1016/s0169-8095(01)00112-0, 2001.

775 Tripathi, S. N., Vishnoi, S., Kumar, S., and Harrison, R. G.: Computationally efficient expressions for the collision efficiency between electrically charged aerosol particles and cloud droplets, *Quarterly Journal of the Royal Meteorological Society*, 132, 1717-1731, <https://doi.org/10.1256/qj.05.125>, 2006.

Tripathi, S. N., Michael, M., and Harrison, R. G.: Profiles of Ion and Aerosol Interactions in Planetary
780 *Atmospheres*, *Space Science Reviews*, 137, 193-211, 10.1007/s11214-008-9367-7, 2008.

Unterstrasser, S., Hoffmann, F., and Lerch, M.: Collection/aggregation algorithms in Lagrangian cloud microphysical models: rigorous evaluation in box model simulations, *Geoscientific Model Development*, 10, 1521-1548, 10.5194/gmd-10-1521-2017, 2017.

VanZanten, M. C., Stevens, B., Nuijens, L., Siebesma, A. P., Ackerman, A. S., Burnet, F., Cheng, A.,
785 Couvreux, F., Jiang, H., Khairoutdinov, M., Kogan, Y., Lewellen, D. C., Mechem, D., Nakamura, K., Noda, A., Shipway, B. J., Slawinska, J., Wang, S., and Wyszogrodzki, A.: Controls on precipitation and cloudiness in simulations of trade-wind cumulus as observed during RICO, *Journal of Advances in Modeling Earth Systems*, 3, M06001, 10.1029/2011MS000056, 2011.

Wang, F., Zhang, Y., and Zheng, D.: Impact of updraft on neutralized charge rate by lightning in
790 thunderstorms: A simulation case study, *Journal of Meteorological Research*, 29, 997-1010, 10.1007/s13351-015-5023-9, 2015.

Wang, P. K., Grover, S. N., and Pruppacher, H. R.: On the effect of electric charges on the scavenging of aerosol particles by clouds and small raindrops, *Journal of Atmospheric Sciences*, 35, 1735-1743, 10.1175/1520-0469(1978)035<1735:OTEOEC>2.0.CO;2, 1978.

795 [Weon, B. M. and Je, J. H.: Charge-induced wetting of aerosols, *Applied Physics Letters*, 96, 194101, 10.1063/1.3430007, 2010.](#)

[Zelinka, M. D., Randall, D. A., Webb, M. J., and Klein, S. A.: Clearing clouds of uncertainty, *Nature Climate Change*, 7, 674-678, 10.1038/nclimate3402, 2017.](#)

~~[Yau, M. K. and Rogers, R. R.: A Short Course in Cloud Physics, Elsevier Science, 1989.](#)~~

800 Zhang, L., Tinsley, B. A., and Zhou, L.: Parameterization of in-cloud aerosol scavenging due to atmospheric ionization: Part 3. effects of γ -varying droplet radius, *Journal of Geophysical Research: Atmospheres*, 123, 10546–10567, 10.1029/2018jd028840, 2018.

Zhou, L. and Tinsley, B. A.: Production of space charge at the boundaries of layer clouds, *Journal of Geophysical Research: Atmospheres*, 112, D11203, 10.1029/2006jd007998, 2007.

805 Zhou, L. and Tinsley, B. A.: Time dependent charging of layer clouds in the global electric circuit, *Advances in Space Research*, 50, 828-842, 10.1016/j.asr.2011.12.018, 2012.

Zhou, L., Tinsley, B. A., and Plemmons, A.: Scavenging in weakly electrified saturated and subsaturated clouds, treating aerosol particles and droplets as conducting spheres, *Journal of Geophysical Research*, 114, D18201, 10.1029/2008jd011527, 2009.

810

Figures and Captions:

815





Figure 1: Comparison of the effect of electric charge on the collision kernel for droplets sized 40 μm , 20 μm and 10 μm with small droplet radii between 10⁻² μm and 10 μm . The charging rate α is 0.2 for panel (a) and 0.3 for panel (b). The solid line represents the results where the collision kernel is calculated by the analytical expression and treats the charged droplets as CS setting. The long-dashed line represents the results calculated by the analytical expression and treats the charge droplets as Khain04 setting. The dashed-dotted-dotted line represents the results calculated by the analytical expression and treats the electrostatic electric force by the IM setting. The dashed-dotted line represents the results calculated by the analytical expression and treats the electrostatic electric force by the CB setting. The dotted line shows the results with NC setting. The dashed line represents the results of the trajectory simulation according to Zhou et al. (2009).


820

825



830


Figure 2: A comparison of the spatial structure of the mixing ratio of hydrometeors of the cumulus with NC setting, the electric force evaluated by CB setting and the electrostatic electric force evaluated by CS setting at times of 1500s, 2100s and 2700s. The charging rate α is 0.3.



835 **Figure 3: The time evolution of the domain-averaged precipitation amount (a) and the domain-averaged water path of the liquid water path (b), rainwater path (c) and cloud water path (d), which is consistent with Figure 2. The solid line represents NC setting, the dashed line represents CB setting, and the dashed-dotted line represents CS setting. The grey shade indicates the standard deviation calculated from 50 members of the random ensemble.**


840

845




850

Figure 4: The mass density distribution evolution of the droplets at 1500s (a), 2100s (b) and 2700s (c), which is consistent with Figure 2 and Figure 3. The solid line represents NC setting, the dashed line represents CB setting, the dotted line represents IM settings and the dashed-dotted line represents CS setting.




860 **Figure 5: Comparison of the cloud evolution for variable charging rates. The mass density distribution of droplets at 1500s (a), 2100s (b) and 2700s (c) and time evolution of the domain-averaged water path of liquid water path (LWP) (d), cloud water path (CWP) (e), and rain water path (RWP) (f) are presented for charging rate α is 0.1 (solid line), 0.2 (dashed-dotted line) and 0.6 (dotted line).**



870 **Figure 6: A comparison of the evolution of the domain-averaged precipitation amount with variable droplet charging rate α is 0.05 (orange solid line), 0.1 (black solid line), 0.2 (black dashed line), and 0.6 (black dashed-dotted line), and the electric force is evaluated with the CS setting. The grey solid line represents the NC setting.**

875



880

Figure 7: The time evolution of the domain-averaged precipitation of the aerosol concentration represented by solid lines ($3.15 \times 10^8/m^3$), dotted lines ($9.45 \times 10^8/m^3$) and dashed-dotted lines ($15.75 \times 10^8/m^3$). The black lines represent the simulation with the electric force, which is evaluated with the CS setting, and grey lines represent the simulation with NC setting. The charging rate α is 0.2.

885

890




Figure 8: Comparison of the time evolution of the domain-averaged precipitation amount for variable evaluation of the electric force. The grey solid line represents the NC setting. The dashed-dotted lines represent the results of the electric force evaluated by the CS method. The black dashed-dotted line represents the result of droplets of with opposite sign charges. The orange dashed-dotted line represents the condition where the charge is only on the large droplet, the yellow dashed line represents the result of droplets with opposite sign charges and the CS method based on Khain et al. (2004). The black dotted line is the result of droplets with opposite sign charges, and the electric force is evaluated by the image charge method.

895

900