



Measurement Report: Effects of transition metal ions on the optical properties of humic-like substances revealing structural preference

Juanjuan Qin^{1,2}, Leiming Zhang³, Yuanyuan Qin¹, Shaoxuan Shi¹, Jingnan Li¹, Zhao Shu¹, Yuwei Gao¹, Ting Qi⁴, Jihua Tan¹, Xinming Wang²

5 ¹College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China


²Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

³Air Quality Research Division, Science & Technology Branch, Environment and Climate Change Canada, Toronto, Canada

⁴College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China

Correspondence to: Jihua Tan (tanjh@ucas.ac.cn); Xinming Wang (wangxm@gig.ac.cn)

10 **Abstract.** Humic-like substances (HULIS) are complex macromolecules in water-soluble organic compounds containing multiple functional groups, and transition metal ions (TMs) are ubiquitous in atmospheric particles. In this study, potential physical and chemical interactions between HULIS and four TM species including Cu²⁺, Mn²⁺, Ni²⁺, and Zn²⁺ were analyzed by optical method under acidic, weakly acidic and neutral conditions. The results showed that Cu²⁺, Mn²⁺, and Zn²⁺ only slightly enhanced mass absorption efficiency (MAE₃₆₅) of HULIS in winter, and had indiscernible effects on absorption
15 Ångström exponent (AAE) of HULIS in both seasons under all acidity conditions. All four TMs had fluorescence quenching effects on winter HULIS, and only Cu²⁺ had similar effects on summer HULIS, with the highest quenching coefficients found under weakly acidic condition in both seasons. The ¹H-nuclear magnetic resonance (¹H-NMR) and Fourier-transform infrared (FT-IR) spectra revealed that Cu²⁺ mainly bound with aromatic species and tightened the molecule structures of HULIS. The parallel factor analysis (PARAFAC) results extracted four components of HULIS, including low-oxidized
20 humic-like substances (C1), N-containing compounds (C2), highly-oxidized humic-like substances (C3), and the mixing residentials (C4), from the fluorescence spectra in both winter and summer. The divergent variations of HULIS spectral components with Cu²⁺ additions under three acidity conditions indicated that electron-donating groups of HULIS mainly corresponded to C1 and C3, with Cu²⁺ binding with HULIS by replacing proton, while electron-withdrawing groups of HULIS could correspond to C2, with its connection with Cu²⁺ through electrostatic adsorption or colliding induced energy
25 transfer.



1 Introduction

Atmospheric particles, consisting of both organic and inorganic substances, have significant impacts on the natural environment, climate, and human health. Humic-like substances (HULIS) and transition metal ions (TMs) are among the major organic and inorganic fractions, and can substantially affect the physical and chemical properties of atmospheric particles (Hawkins et al., 2016; Zanca et al., 2017; Frka et al., 2018; Ma et al., 2019). Besides, HULIS and many TMs, like Fe, Cu, Mn, V, Cr, and Co, have negative impacts on human health since they can promote the generation of reactive oxygen species (ROS), which cause inflammatory response of human respiratory system (Verma et al., 2012; Gali et al., 2015; Lin and Yu 2019). HULIS are known as a mixture of macromolecular organic compounds, containing aromatics and aliphatic species with multiple oxygenated functional groups like carbonyls, hydroxyl, nitrate, and nitroxy-organosulfate (Win et al., 2018). TMs can transfer electrons and participate in chemical reactions or serve as catalyst, especially in atmospheric photochemistry (Mao et al., 2013; Guasco et al., 2014).

Considering that HULIS and TMs are both active chemical components of atmospheric particles, the significance of their co-effects is apparently conceivable. Early studies have demonstrated that HULIS can enhance the solubility of TMs by organic complexation (Paris and Desboeufs 2013; Scheinhardt et al., 2013), and TMs might mediate organic transformation processes and induce the formation of light absorbing secondary water-soluble organics and water insoluble abiotic polymers (Slikboer et al., 2015). More recent studies focused on the oxidation potentials, and ROS formation or other metal induced photochemical reactions of TMs (Fe^{2+} , Cu^{2+} , and Mn^{2+}) in combination with HULIS (Gonzalez et al., 2017; Lin and Yu 2019; 2020; Ye et al., 2021). The mixture of TMs and HULIS could exhibit synergistic or antagonistic effects on ROS formation



45 associated with the chemical composition of HULIS. Moreover, TMs can also bind with HULIS and affect the light absorption properties of HULIS (Fan et al., 2021; Wang et al., 2021).

To reveal the linkage between HULIS and TMs in atmospheric chemistry, this study characterized the varying optical and chemical properties of HULIS with addition of different TMs species under three different levels of acidity. The effects of TMs on light absorption of HULIS and the TMs-sensitive fraction of HULIS were determined by optical spectra, and parallel factor analysis (PARAFAC) were performed to unveil the susceptible fluorescent components. The interaction between Cu^{2+} and HULIS were further analyzed for its prominent effects, and the potential reaction mechanisms were deduced based on the structural and optical variation scenarios. Results from the present study advanced our understanding on the physical and chemical characteristics and transformations of HULIS, especially the physical and chemical connections between HULIS and TMs.

55 2 Methods

2.1 Sample extraction and titration

In the present research, two concentrated HULIS solutions were extracted from $\text{PM}_{2.5}$ samples collected in a winter and summer month in Beijing, and the detailed sample information can be found elsewhere (Qin et al., 2022). The HULIS isolation was mainly based on the method established by Lin et al., (2010) with some adjusted procedures for practical reasons. Firstly, WSOC was ultrasonically extracted from 30 or 60 daily $\text{PM}_{2.5}$ filter samples ($D = 2$ cm, 30 pieces for winter and 60 pieces for summer), using 200 mL Milli-Q water three times, and filtered using $0.22 \mu\text{m}$ membranes. The liquids were then separated by SPE cartridges (Oasis HLB, 6cc, 500 mg/cartridge, Waters, USA), rinsed with 10 mL water, and pumped to dry. The cartridges were eluted with 10 mL methanol afterwards. Finally, the eluants were evaporated to dryness by N_2 at room temperature and redissolved into 200 mL ultrapure water to obtain a HULIS stock solution.

65 Before metal ion titration, the HULIS stock solutions were diluted by 10 folds, and the total organic carbon (TOC) concentration was determined by a TOC analyzer (Multi N/C 3100, Germany), with the carbon concentrations of 1.9 mg/L for winter HULIS and 1.6 mg/L for summer HULIS. The original acidity of HULIS was determined by a pH meter (Mettler Toledo, Swiss) before and after dilution. In both winter and summer, triplicate 600 mL of diluted HULIS solutions were separately added to 1 L sealed vials to obtain weakly acidic HULIS ($\text{pH} = 5.65$ for winter and $\text{pH} = 5.92$ for summer), acidic HULIS ($\text{pH} \approx 3.0$, by $0.05\text{M H}_2\text{SO}_4$), and neutral ($\text{pH} \approx 7.5$, by 0.1M NaOH), respectively. Considering the instability of natural pH gradient, excessive base solution was added to ensure the solution pH to be around neutral before measurement. Four transition metals ions including Cu^{2+} , Zn^{2+} , Mn^{2+} , and Ni^{2+} were chosen as representative metal ions that might be reactive to HULIS, considering their richness in atmospheric environment or significant health risks (Fan et al., 2021; Wang et al., 2021). Titration experiments were conducted by adding 0 – 526.32 μl corresponding metal sulfate solutions (0.01 M) and 9 ml diluted HULIS solution to a 10 ml amber volumetric flask, and ultrapure water was then added to obtain samples containing TMs concentrations of 0, 20, 50, 100, 200, 300 and 500 μM , respectively. The titrated solutions were shaken for



12 h at room temperature to ensure complexation equilibrium, and pH of all titrated samples were detected again before spectral analysis.

2.2 Spectral analysis

80 The optical properties of HULIS with addition of TMs (0–500 μM) were analyzed by UV-Vis and fluorescence spectrophotometers under acidic, neutral, and neutral pH. The chemical properties of HULIS before and after adding TMs were analyzed by ^1H -nuclear magnetic resonance (^1H -NMR) and Fourier-transform infrared (FT-IR).

The UV-vis absorption spectra of WSOC and HULIS were measured by UV-VIS photometer in Agilent Technologies covering wavelength range of 200 nm–500 nm with an interval of 1nm, using a scanning speed of 200 nm/min. The sample 85 pool is a quartz colorimetric dish with an optical path of 1 cm.

The three-dimensional fluorescence spectra were measured by an Agilent Technologies photometer under the following conditions: excitation wavelength (EX) 200 nm–400 nm, step length 5 nm, slit width 10 nm; emission wavelength (EM) 250 nm–500 nm, step length 5 nm, slit width 5 nm; scanning speed: 2400 nm/min; and response time: automatic. The sample pool uses a quartz colorimeter with a light path of 1cm.

90 The infrared spectra of HULIS for wavenumber between 4000-400 cm^{-1} were measured by an FTIR spectrometer (Thermal Fisher, Nicolet IS10) at room temperature. The essential extraction of HULIS for FT-IR measurements was the same as those described in section 2.1 above, except that around 1 mg of the HULIS (dried by gentle stream of N_2 at room temperature 25 $^\circ\text{C}$) was directly mixed with 60 mg KBr and dried in a drying oven, rather than redissolved by water. The mixtures were then ground and pressed into firm tablets for analysis. A 600 MHz NMR spectrometer (Bruker, AVANCE III 95 600 M) was applied for ^1H -NMR spectra measurements. About 5 mg HULIS samples were redissolved into 1 mg D_2O for detection.

2.3 Data Analysis

2.3.1 Mass absorption efficiency and absorption Ångström exponent

100 The mass absorption efficiency (MAE, $10^3 \cdot \text{AU} \cdot \text{cm}^2/\text{mg}$) of HULIS and the absorption Ångström exponent (AAE) were obtained as follows (Kirillova et al., 2014; Saleh et al., 2014):

$$\text{MAE} = (A_\lambda - A_{700}) \times \frac{\ln(10)}{C_{\text{species}} \times L} \quad (1)$$

$$A_\lambda = K\lambda^{-\text{AAE}} \quad (2)$$

Where A_λ is light absorbance at wavelength λ , C_{species} is the chemical concentration of organic compounds (WSOC and HULIS in the present research), L is the light path length (1 cm), K is a scaling constant, and the fitting wavelength of AAE 105 is 330-400 nm.



2.3.2 Fluorescence indices

Fluorescence indices based on intensity ratios may provide clues about the condensation state of WSOC. The average fluorescence intensity is the mean intensity of an excitation-emission matrix (EEM) spectra; the specific fluorescence intensity (SFI) is the fluorescence intensity per unit TOC (mg/L) (Kalbitz et al., 2000). The biological index (BIX) reflects the freshness of microbially produced DOM (Parlanti et al., 2000), with a high BIX value (> 1) corresponding to a predominantly microbial derived organic matter, and a value of <0.6 indicating fewer microbial originated DOM (Qin et al., 2018). BIX is calculated as follows:

$$BIX = \frac{EEM_{Ex200-310,Em250-380}}{EEM_{(max(Ex200-310,Em420-435))}} \quad (3)$$

The fluorescence lifetime for HULIS was estimated by a formula derived from the Strickler–Berg equation (Strickler and Berg 1962). The τ_0/ρ ratio comprehensively reflects the fluorescence intrinsic lifetime (related to the fluorophore type) and fluorophore density (associated with the molecular structure), a large τ_0/ρ can be resulted from a long fluorescence lifetime or a small fluorophore (Xiao et al., 2018), it was a representation of fluorescence lifetime in this research.

$$\tau_0/\rho = c2.88 \times 10^{-12} n^2 \frac{\int_{Em} FI(v)dv}{\int_{Em} v^{-3} FI(v)dv} \int_{Ex} UVA(v)dv^{-1} \quad (4)$$

Where τ_0 is the intrinsic lifetime (ns), ρ is the density of fluorophore in the organics (mmol-fluorophore per gC), c is the total organic concentration (mgC/L), n is the refractive index of the solution ($n^2 \approx 1.8$), v is the excitation or emission wavenumber (cm^{-1}), UV is the UV-Vis absorbance (A.U.), and FI is the fluorescence intensity (R.U.).

Stokes shift (SS) can reflect the fluorescence energy of organics (Xiao et al., 2020; Yu et al., 2020), and is defined as $Ex^{-1} - Em^{-1} \text{ nm}^{-1}$, demonstrating the energy loss of the excited fluorophore due to internal conversion during relaxation (Lakowicz 2006). AvgSS were defined as the intensity-weighted average Stokes shifts of HULIS (Xiao et al., 2019).

2.3.3 Parallel factor analysis (PARAFAC)

PARAFAC model can decompose complex EEM spectra into several main components by statistical method. The excitation spectrum, emission spectrum, and scores of each component are as follows:

$$x_{ijk} = \sum_{f=1}^F a_{if} b_{jf} c_{kf} + \varepsilon_{ijk} \quad i=1,\dots,I, j=1,\dots,J, k=1,\dots,K \quad (5)$$

Where x represents the fluorescence intensity, f is the number of components resolved by PARAFAC, a is proportional to the concentration of the f -th component, and b and c are the scaled estimating of the emission and excitation spectra. The subscript i is the sample number, and j and k represent emission and excitation wavelength, respectively. Before performing PARAFAC, all EEM data were normalized to unit norm to reduce concentration-related collinearity and avoid extremely different leverages (Wang et al., 2020). Tucker congruence coefficient (TCC) was determined for each excitation spectrum and emission spectrum, and a threshold of 0.95 was applied to confirm the spectral congruence. The model was determined by half-split validation.



3 Results and discussion

The light absorption and fluorescence efficiencies of winter and summer HULIS under acidic, weakly acidic, and neutral environments are depicted in Figure 1, and the spectra derived indices are listed in

140 Table 1. Generally, the overall light absorption efficiencies (represented by MAE) of winter HULIS were higher than those in summer, and the AAE showed a reverse trend, indicating higher light absorption abilities of winter than summer samples (Mo et al., 2021). The MAE and AAE of HULIS under three acidity levels in both seasons were consistent with those reported in our previous research, with the average MAE₃₆₅ at 0.011±0.00 in winter and 0.005±0.001 in summer, and the corresponding average AAE_{300–400nm} at 6.46±0.86 and 6.97± 0.83, respectively (Qin et al., 2022). The fluorescence efficiencies of winter HULIS (represented by SFI and AFI/TOC) were higher than those of summer HULIS as well (Figure 145 1), and the fluorescence indices revealed some structural differences. The SFI spectra of HULIS mainly exhibited two fluorescence peaks at wavelength of Ex = 225–230 nm, Em = 380–400 nm, and Ex = 325–330 nm, Em = 395–400 nm in both seasons, with the peaks being able to be roughly characterized as fulvic-acid like group and protein-like groups, respectively. The Stokes shift index of AvgSS could be associated with the hydrophobicity/hydrophilicity of organics, with a low AvgSS implying a relatively high hydrophilicity and a high one implying a high hydrophilicity (aromaticity or 150 humification degree) (Xiao et al., 2019). Thus, the higher AvgSS of winter HULIS indicated a higher hydrophilicity compared to that in summer.

The different optical properties of HULIS between winter and summer were mainly resulted from divergent chemical structures. Our previous research had revealed that winter HULIS contained more unsaturated structures and were relatively hydrophobic, leading to the higher light absorption and fluorescence efficiencies, while summer HULIS were rich with 155 oxidized species and functional groups and were more hydrophilic (Qin et al., 2022). Baduel et al., (2011) evaluated the varying optical properties of HULIS during O₃ oxidation, and found the C=O and COOH groups of HULIS increased with oxidation. However, Yang et al., (2021) found that O₃ oxidation of humic acid could consume some functional groups like aromatic C=C, OH, C=O, COOH and COO⁻, causing an decrease in MAE and an increase in AAE, and these divergences were ascribed to the different structures of HULIS. The different structures of HULIS between winter and summer could not 160 only lead to the varied optical properties but also the distinctive behaviors of HULIS with other coexist species in atmospheric environment. Therefore, it is necessary to analyze the relations between TMs and HULIS for winter and summer HULIS separately.




Figure 1 The MAE (a-b) and SFI (c-h) spectra of winter and summer HULIS under three acidity conditions.

165 **Table 1** The light absorption and fluorescence indices of HULIS under three acidity conditions.

	Winter				Summer			Mean ± S.D.
	Acidic	Weakly acidic	Neutral	Mean ± S.D.	Acidic	Weakly acidic	Neutral	
MAE ₃₆₅ ^a	0.011	0.011	0.011	0.011 ± 0.00	0.006	0.004	0.006	0.005 ± 0.001
AAE	7.05	6.86	5.47	6.46 ± 0.86	6.97	7.80	6.15	6.97 ± 0.83
AFI/TOC ^b	0.79	0.62	0.58	0.66 ± 0.11	0.10	0.10	0.09	0.10 ± 0.01
BIX	1.07	0.96	0.99	1.00 ± 0.05	0.98	0.90	0.90	0.93 ± 0.05




AvgSS	1.54	1.54	1.54	1.54 ± 0.00	1.01	0.90	0.91	0.94 ± 0.06
τ_0/ρ	0.15	0.24	0.13	0.17 ± 0.06	0.56	0.92	0.60	0.69 ± 0.19

a: $10^3 \cdot \text{AU} \cdot \text{cm}^2/\text{mgC}$ b. $\text{RU} \cdot \text{L}/\text{mgC}$

3.2 Effects of TMs on the light absorption properties of HULIS

The gradients of $\text{MAE}_{365_i}/\text{MAE}_{365_0}$ and $\text{AAE}_{300-400\text{nm}}$ for HULIS with TM (Cu^{2+} , Mn^{2+} , Ni^{2+} , and Zn^{2+}) concentrations ranging from 0 to 500 μM under three acidity conditions are depicted in Figure 2 and the whole MAE spectra are exhibited in SI Figures S1 and S2. For winter HULIS, under acidic environment, the addition of Mn^{2+} , Ni^{2+} , and Zn^{2+} at low concentrations ($< 50 \mu\text{M}$) could induce a decrease of 14%–16% in MAE_{365} , and further increasing TM concentrations had little effects on MAE_{365} , whereas Cu^{2+} showed minimal effects on MAE_{365} , which only decreased MAE_{365} by 5% with increasing TM concentrations. Under weakly acidic environment, the addition of a small amount of any of the four TMs (0–50 μM) could induce evident increase of MAE_{365} at a range of 16% to 20%, and then MAE_{365} became steady or slightly decreased with further increasing TM concentrations. As for the neutral environment, the addition of Cu^{2+} could lead to up to 9% increase in MAE_{365} , but Cu^{2+} showed inhibiting effects on MAE_{365} in cases with additions of Mn^{2+} , Ni^{2+} , or Zn^{2+} , with the maximum decrease at 11%, 7%, and 14%, respectively. Nevertheless, TMs could only change the $\text{AAE}_{300-400\text{nm}}$ of HULIS slightly, and the solution acidity sieved the initial $\text{AAE}_{300-400\text{nm}}$ of HULIS; besides, the TM concentrations had inconspicuous effects on the regularity. An exception was observed for Ni^{2+} because of its self-absorption, with $\text{AAE}_{300-400\text{nm}}$ continuously reducing with increasing additions of $[\text{Ni}^{2+}]$.

For summer HULIS, Cu^{2+} , Mn^{2+} , and Zn^{2+} could cause fluctuations in MAE_{365} with increasing metal ion concentrations under acidic and neutral environments, and they inhibited the light absorption efficiencies of HULIS under weakly acidic environment. Likewise, Ni^{2+} could enhance the MAE_{365} of HULIS under all acidity conditions because of the increasing absorption generated by increasing Ni^{2+} concentrations. The $\text{AAE}_{300-400\text{nm}}$ of HULIS had subtle variation when adding Cu^{2+} , Mn^{2+} , and Zn^{2+} under acidic and neutral environments, indicating that the TMs-HULIS mixtures might not compose complexations or their complexes don't have structural effects on HULIS. Yet when the solution was weakly acidic, the $\text{AAE}_{300-400\text{nm}}$ exhibited sharp fluctuations with increasing Mn^{2+} and Zn^{2+} , and even increased with increasing Cu^{2+} concentrations.



190 **Figure 2** The light absorption indices of HULIS with additions of TMs under different acidity conditions.

3.3 Effects of TMs on the fluorescence properties of HULIS

The effects of TMs on the fluorescence indices of HULIS under different acidity environments are shown in Figure 3 and Figure 4, and the fluorescence spectral variations are provided in SI Figures S3 and S4. Obviously, all four TMs showed quenching effects on the AFI of HULIS with fiddling seasonal or acidity dependent differences, with the effectiveness ranking in the order of $\text{Cu}^{2+} > \text{Mn}^{2+} \sim \text{Ni}^{2+} > \text{Zn}^{2+}$. At the maximum TMs concentration of 500 μM , AFI of winter HULIS reduced by 5%–21%, 5%–11%, and 7%–28%, and of summer HULIS reduced by 8%–22%, 12%–39%, and 5%–26% under acidic, weakly acidic, and neutral environments, respectively.

The effects of TMs on fluorescent HULIS could be evaluated by the Stern-Volmer equation. Supposing metal ions only singly bonded with HULIS molecules, the quenching constants (k) of TMs species to the total fluorescence of HULIS were obtained and are listed in Table 2. Briefly, for winter HULIS, all TMs species had significant liner correlations with AFI/AFI_0 ($R^2 > 0.6$). Cu^{2+} and Mn^{2+} had largest k constants under acidic and neutral environments and relatively low k under weakly acidic environment, indicating that Cu^{2+} and Mn^{2+} might mainly bond with organic groups that dissociated under acidic or neutral environments. Ni^{2+} had the same k under weakly acidic and acidic environments and larger k under neutral environment, indicating that Ni^{2+} might prefer to bond with neutral HULIS. Similarly, Zn^{2+} had larger k under acidic




205 environment, indicating a preference of acidic environment for the complexation of Zn^{2+} and HULIS. For summer HULIS, Cu^{2+} still had significant and large k under acidic and neutral environments and relatively low k under neutral environment. Mn^{2+} only had a relatively large k under acidic environment and a low k under neutral environment, but no statistically significant k under weakly acidic environment. Besides, AFI/AFI_0 of HULIS with Ni^{2+} and Zn^{2+} also had non-significant k , indicating negligible interactions between fluorescent HULIS and these two ions.

210 Although TMs showed quenching effects on HULIS, the locations of fluorescence peaks rarely moved with increasing TM concentrations. Thus, the fluorescence spectra-based indices only show minimal variations under different acidity conditions with increasing metal ions concentrations. In Figure 4, the Stokes shift spectra of HULIS separated into groups under three acidity conditions for both seasons. Only addition of Cu^{2+} could hierarchically distribute the spectra slightly. The other three TMs showed inconspicuous effects, indicating that they could not cause substantial structural variations on fluorescent

215 HULIS, as can be more visually presented by the ratio of $AvgSS_0/AvgSS_i$ (Figure 5). Additionally, the addition of Mn^{2+} , Ni^{2+} , and Zn^{2+} didn't change the τ_0/ρ of HULIS either, which might confirm that they did not bound with HULIS molecules since fluorescent lifetime is a sensitive indicator vulnerable to the micro-environment surrounding detected objects (Coble, 2014).

Table 2 The fluorescence quenching coefficients of TMs and HULIS obtained from Stern-Volmer equation

Stern-Volmer	Acidity	Cu^{2+}		Mn^{2+}		Ni^{2+}		Zn^{2+}	
		$k \cdot 10^3$	R^2	$k \cdot 10^3$	R^2	$k \cdot 10^3$	R^2	$k \cdot 10^3$	R^2
Winter	Acidic	0.60	0.99	0.10	0.98	0.10	0.95	0.20	0.98
	Weakly Acidic	0.10	0.86	0.08	0.91	0.10	0.95	0.07	0.83
	Neutral	0.60	1.00	0.10	0.68	0.20	0.90	0.07	0.58
Summer	Acidic	0.60	0.99	0.20	0.38	-0.02	0.79	0.01	0.02
	Weakly Acidic	1.00	0.82	-0.10	0.02	-0.04	0.05	-0.20	0.13
	Neutral	0.40	0.88	0.05	0.63	0.05	0.08	-0.20	0.16



220 **Figure 3 The average fluorescence intensities of HULIS with adding of Cu^{2+} , Zn^{2+} , Mn^{2+} , and Ni^{2+} under different acidity conditions in winter and summer.**





Figure 4 The Stokes shift spectra of HULIS with adding of Cu^{2+} , Zn^{2+} , Mn^{2+} , and Ni^{2+} under different acidity conditions.



225 **Figure 5** The fluorescence indices of HULIS with increasing Cu^{2+} , Zn^{2+} , Mn^{2+} , and Ni^{2+} concentrations under different acidity conditions.



3.4 Distinguishing the varying fractions of HULIS with additions of TMs

To further identify the response of specific HULIS compounds to TMs, PARAFAC methods were applied to distinguish the varying groups of HULIS with addition of different TM species. In general, four compounds (C1-C4) were sorted out from EEM spectra of winter and summer HULIS, and they were characterized as low-oxidized humic-like substances (C1), N-containing compounds (C2), highly-oxidized humic-like substances (C3), and the mixing residentials (C4), respectively (Chen et al., 2016). The constitution of C1-C4 in winter and summer HULIS under different acidity levels are listed in Table 3.

Form the correlations between F/F_0 ratio and TM concentrations of fluorophores C1-C4 shown in Figure 6 and Figure 7, it was obvious that the intensities of four fluorophores had increasing or decreasing tendencies in response to the acidity and TM species because of the divergent acid/base groups of HULIS. In the case of winter HULIS under acidic environment, Cu^{2+} could induce a substantial fluorescence quenching effects on C1 and C4 and a concurrent modest fluorescence increasing effects on C3, Mn^{2+} and Ni^{2+} exhibited weak fluorescence quenching effects on C1 and C3, and Zn^{2+} exhibited weak fluorescence quenching effects on C1 and C4. Under weakly acidic environment, all of the four TMs exhibited discernible quenching effects on C2 and C3, with Cu^{2+} manifest as the most potent quencher. Under neutral environment, Cu^{2+} significantly quenched the fluorescence intensities of C1, C2, and C4, Ni^{2+} displayed modest quenching effects only on C3, and other metal ions showed inconspicuous fluorescence quenching/enhancement effects.

As for summer HULIS, only Cu^{2+} exhibited notable quenching effects on fluorophores C1-C4, while Mn^{2+} , Ni^{2+} , and Zn^{2+} showed rather minimal or erratic effects under all acidity conditions. Under acidic and weakly acidic environments, Cu^{2+} could reduce the fluorescence intensity of C1-C3, and contributions of C4 to total fluorescence intensities gradually increased. For neutral environment, Cu^{2+} exhibited pronounced fluorescence quenching effects on C2, and mild reduction tendencies on C1 and C3. Because of the minimal quenching effects of Mn^{2+} , Ni^{2+} , and Zn^{2+} on C1-C4, most of the correlation coefficients (R^2) between F/F_0 and metal ions concentrations were small, indicating the insignificant effects of Mn^{2+} , Ni^{2+} , and Zn^{2+} on HULIS.

Table 3 The PARAFAC decomposed fractions of HULIS for winter and summer under three acidity conditions.

		C1	C2	C3	C4	Total fluorescence intensity
Winter	Acidic	42%	19%	24%	15%	805.52
	Weakly acidic	29%	24%	27%	19%	869.69
	Neutral	22%	27%	29%	22%	587.49
Summer	Acidic	50%	28%	20%	2%	192.09
	Weakly acidic	23%	26%	18%	32%	179.17
	Neutral	16%	32%	16%	36%	161.00




Figure 6 The changing patterns of four fluorophores with TMs under different acidity conditions for winter HULIS.


In light of the above analysis, it was obvious that Cu²⁺ demonstrated much more significant effects on optical properties of HULIS than the other three metal ions, which might be because of their disparate complexing abilities (Fan et al., 2021). Li et al., (2022) found that with increasing Fe³⁺, Al³⁺, and Cu²⁺ concentrations, the fluorescence intensities of highly oxidized HULIS decreased, and those of low oxidized HULIS gradually increased, and they ascribed these phenomena to the fluorophore blue-shifts of highly oxidized HULIS induced by complexation. Likewise, some other studies observed strong bondage between Cu²⁺ and atmospheric organic matter (Wang et al., 2021; Ma et al., 2022). However, the detailed mechanisms causing the differed complexation results of TMs remained unclear.

It is noticed that studies on the effects of TMs on optical properties of fulvic acid and humic acid had been conducted in earlier years in surface water than atmospheric environment, which provided an inspiration to the present study. For example, it was reported in 1980s that Cu²⁺ posed greater fluorescence quenching effects on fulvic-acid than Co²⁺ and Mn²⁺, and it was thus proposed that Cu²⁺ bound internally to fulvic-acid, while Co²⁺ and Mn²⁺ might only externally attached to organics via electrostatic attraction (Ryan et al., 1983). Other studies found that Cu²⁺ had stronger complexation capabilities



265 compared to the other divalent transition metal ions, such as Pb^{2+} , Cd^{2+} , Ni^{2+} , and Zn^{2+} (Yamashita and Jaffe 2008; Chen et al., 2015; Huang et al., 2018). With application of two-dimensional infrared correlation spectroscopy and synchronous/asynchronous fluorescence spectroscopy analyze, carboxyl and polysaccharide groups were found to bind fast with Cu^{2+} ; and bondage of Cu^{2+} with phenolic and aromatic carboxyl groups altered the molecule vibration of organics, inducing fluorescence quenching. Besides, Cu^{2+} could bind with several amide and aliphatic groups (protein-like), leading to

270 fluorescence quenching. More recently, Zhu et al. (2021) suggested that the small atomic radius and paramagnetic properties might result in the unique response of Cu^{2+} to carboxyl groups, thus a strong complexation ability with DOM. Considering the smaller molecules and aromatic structures of HULIS compared to humic substances in water, Cu^{2+} continued its specialty of significant fluorescence quenching effects on HULIS, whereas bondages between Zn^{2+} (or Mn^{2+} and Ni^{2+}) and responsive HULIS might be too weak to be observed.



275

Figure 7 The changing patterns of four fluorophores with TMs under different acidity conditions for summer HULIS.



3.5 The hidden mechanism of HULIS with addition of TMs - taking Cu²⁺ as an example

To further understand the complexation mechanisms of Cu²⁺ and HULIS, the structural variations of HULIS before and after adding 100 μM Cu²⁺ were evaluated by ¹H-NMR and FT-IR methods, respectively (Figure 8 and Figure 9). The ¹H-NMR results showed that HULIS were rich with protons of H-C, H-C=C-, Ar-H, and H-C-O and functional groups of -OH or -NH, Ar-C=O, Ar-C-O-C, and C-O in both seasons. The addition of Cu²⁺ mainly affected the protons bound with aromatics and oxygen in HULIS, with the signals located between δ7-δ8 and δ3.4-δ4 ppm trailing off, for which the Ar-H signal decreased by 8.9% and 2.1%, and H-C-O signal decreased by 1.4% and 0.8% in winter and summer, respectively. According to the basic principles of NMR spectroscopy, coordination or chemical exchange reactions between HULIS molecules and Cu²⁺ might cause the chemical shift signals vanished or observed as new signals (Günther 2013). Meanwhile, the functional groups of HULIS, revealed by FT-IR spectra, mainly included -OH/-NH, -CH, Ph-C=O, C=O, Ph-C-O-C, and -C-O in both seasons, and the summer HULIS contained more oxygen-containing functional groups than winter HULIS. With the addition of Cu²⁺, the whole FTIR spectra of winter and summer HULIS blue shifted to higher energy, especially for O-containing aromatic groups, but the peak shapes remained unchanged, indicating that Cu²⁺ might not directly lead to the transformation of functional groups, but could result in an enhancement of chemical bonds in HULIS molecules.




Figure 8 The ¹H-NMR spectra of HULIS and HULIS with addition of 100 μM Cu²⁺ in winter and summer.




Figure 9 The FT-IR spectra of HULIS and HULIS with addition of 100 μM Cu^{2+} (light colors) in winter and summer.

295 Our previous research had characterized the pK_a and corresponding acidic/basic functional groups of winter and summer HULIS, and proposed a hypothetical HULIS structure of unsaturated aromatic main body with pH responsive functional groups (Qin et al., 2022). Briefly, the pK_a of 3.7 – 3.8 and 8.0 – 8.1 were assigned as electron-donating groups (represented by H_1A_1 and H_3A_3 , respectively), and the pK_a of 5.0 – 6.3 were assigned as electron-withdrawing groups (represented by H_2A_2) for winter HULIS. Likewise, the pK_a of 6.4 was assigned as electron-donating groups (represented by H_4A_4) and pK_a of 8.4 – 8.5 was assigned as electron-withdrawing groups (represented by H_5A_5) for summer HULIS. By synthetically considering the versatile optical properties of HULIS with addition of Cu^{2+} under three acidity conditions, and its inherent acid/base groups, the Cu^{2+} and HULIS complexation or interaction mechanisms were explored and the schematic diagram is depicted in Figure 10.

(1) Firstly, for winter HULIS under acidic environment, the acidity-sensitive HULIS molecules were hard to deprotonate when H^+ was crowding around. Most of HULIS existed as forms of H_1A_1 , H_2A_2 , and H_3A_3 . Cu^{2+} could easily be absorbed by the electron-donating groups of HULIS to form $\text{Cu-A}_1\text{H}_1$ complex, or replace the H atom to form Cu-A_1 , causing static fluorescence quenching of C1 and C4 which slightly increased the lifetime of HULIS. Simultaneously, Cu could alter the charge densities of the fluorescent aromatic molecules, leading to an enlargement of conjugated structures and enhancing the fluorescence of C3.

(2) Secondly, for winter HULIS under weakly acidic environment, some of the acidity-sensitive HULIS deprotonated and the solution contained A_1^- , H_2A_2 , and H_3A_3 . When added with Cu^{2+} , A_1^- complexed with Cu^{2+} to form Cu-A_1 , whereas the sensitive structures of H_2A_2 were electron-donating groups which could hardly provide electrons to Cu^{2+} . However, Cu^{2+} could still be attracted by protons in the unsaturated aromatic structures of HULIS, causing dynamic fluorescence quenching of C2 and C3. With regard to the fast-decreasing lifetime and steady AvgSS of HULIS, chemical reactions might not be




315 involved in these processes, and thus flexible structures of Cu-A₂H₂ by electrostatic adsorption or colliding induced energy transfer could be the possible explanation.

(3) Thirdly, for winter HULIS under neutral environment, the acidity-sensitive HULIS existed as A₁⁻, A₂⁻, and H₃A₃. The deprotonated HULIS had become non-fluorescent, and collision of A₁⁻ and A₂⁻ with added Cu²⁺ might not be observed. The fluorescent H₃A₃ could provide electrons to Cu²⁺, leading to partially fluorescence quenching effects on C1 and C2, and the potential complexation mechanism might be similar to H₁A₁.

(4) Lastly, for summer HULIS, the quenching mechanism of Cu²⁺ and acidity-sensitive HULIS were similar to those of winter HULIS. Under acidic and weakly acidic environment, H₄A₄ and H₅A₅ of summer HULIS were not deprotonated, the quenching effects could be resulted from the replacement of H atom in H₄A₄ (electron-donating) with Cu²⁺, causing decrease of C1-C3. Under neutral environment, Cu²⁺ complexed with deprotonated A₄⁻ and aromatic structure of H₅A₅ (electron-withdrawing), mainly causing decrease of C2.

Thus, it was speculated that HULIS with electron-donating groups mainly exhibit fluorescence signals in C1 and C3, which were mainly low and highly oxidized humic-like substances, respectively. Whereas for HULIS with electron-withdraw groups, the fluorescence signals were mainly exhibited in C2, which was N-containing species or protein-like species.



330 **Figure 10** The schematic diagram of HULIS complexation with Cu²⁺ under different acidity conditions

Conclusions and Implications

In this study, the interrelations between four TMs and HULIS were analyzed by optical methods under acidic, weakly acidic, and near neutral environments. The results elucidated that the light absorption and fluorescence properties of HULIS were



substantially influenced by the solution acidity. TMs either enhance or attenuate the signal, contingent on the metal species, but with negligible impact on the spectral indices. Cu^{2+} demonstrated the most conspicuous influence on the optical properties of HULIS, while Mn^{2+} , Ni^{2+} , and Zn^{2+} proved to be less responsive. The $^1\text{H-NMR}$ and FT-IR spectra of HULIS and HULIS-Cu indicated that Cu^{2+} mainly bound with aromatic HULIS, resulting in more compacted molecule structures of HULIS. The distinct tendencies of PARAFAC-separated fluorophores with increasing Cu^{2+} under three acidity conditions indicated that electron-donating groups might correspond to low and highly oxidized humic-like substances, and Cu^{2+} coupled with HULIS by substituting H atom, while electron-withdraw groups were N-containing species or protein-like species, with their fluorescence quenching likely caused by electrostatic adsorption or colliding induced energy transfer. Results from this study not only revealed that the complexations of HULIS and TMs could be fragile, but also connected the fluorescence spectra of HULIS with structural characterizations through the micro-effects of TMs on HULIS. HULIS and TMs are both well-known ROS generative species, and the combined effects of HULIS and TMs on ROS generation could be synergistic or antagonistic depending on the TMs species (Yu et al., 2018; Lin and Yu 2020). The physical and chemical interaction between TMs and HULIS might prohibit ROS generation because of their competitive reactions on ROS generation, but the ROS generation abilities of newly formed TM-HULIS groups could be either accelerated or reduced depending on the HULIS structures, which are difficult to be distinguished by ROS detection methods. Our results could provide some perception from chemical analysis perspective on this important issue.

350 **Code availability**

Data availability

The data used in this study are available on the Zenodo data repository platform: <https://doi.org/10.5281/zenodo.10460562>

Author contribution

TJ and WX designed the experiments, QJ and QY carried them out, SZ and GY helped with the experimental issues, LJ and SS finished the data analysis. QJ prepared the manuscript with contributions from all co-authors, ZL, TJ and WX supervised the writing, and QT provided professional advices on the discussion section.

Competing interests

At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.



Acknowledgements

360 This work was supported by the National Key Research and Development Program of China (2022YFC370300, 2023YFE0102400, 2022YFC3703402, and 2022YFC3701103), and the Provincial Natural Science Foundation of Hunan (2023JJ30004).

References

- 365 Baduel C, Monge M E, Voisin D, et al., Oxidation of atmospheric humic like substances by ozone: a kinetic and structural analysis approach. *Environmental Science & Technology*, 2011, 45: 5238-5244.
- Chen Q C, Miyazaki Y, Kawamura K, et al., Characterization of chromophoric water-soluble organic matter in urban, forest, and marine aerosols by HR-TOF-MS analysis and excitation emission matrix spectroscopy. *Environmental Science & Technology*, 2016, 50: 10351-10360.
- Chen W, Habibul N, Liu X Y, et al., FTIR and synchronous fluorescence heterospectral two-dimensional correlation analyses on the binding characteristics of copper onto dissolved organic matter. *Environmental Science & Technology*, 2015, 49: 2052-2058.
- 370 Fan X, Liu C, Yu X, et al., Insight into binding characteristics of copper(II) with water-soluble organic matter emitted from biomass burning at various pH values using EEM-PARAFAC and two-dimensional correlation spectroscopy analysis. *Chemosphere*, 2021, 278: 130439.
- 375 Frka S, Grgić I, Turšič J, et al., Seasonal variability of carbon in humic-like matter of ambient size-segregated water soluble organic aerosols from urban background environment. *Atmospheric Environment*, 2018, 173: 239-247.
- Gali N K, Yang F, Jiang S Y, et al., Spatial and seasonal heterogeneity of atmospheric particles induced reactive oxygen species in urban areas and the role of water-soluble metals. *Environmental Pollution*, 2015, 198: 86-96.
- Gonzalez D H, Cala C K, Peng Q, et al., HULIS enhancement of hydroxyl radical formation from Fe(II): kinetics of fulvic acid-Fe(II) complexes in the presence of lung antioxidants. *Environmental Science & Technology*, 2017, 51: 7676-7685.
- 380 Guasco T L, Cuadra-Rodriguez L A, Pedler B E, et al., Transition metal associations with primary biological particles in sea spray aerosol generated in a wave channel. *Environmental Science & Technology*, 2014, 48: 1324-1333.
- Günther H, *NMR Spectroscopy: Basic Principles, Concepts and Applications in Chemistry* (3rd edition). Germany, 2013.
- Hawkins L N, Lemire A N, Galloway M M, et al., Maillard chemistry in clouds and aqueous aerosol as a source of atmospheric humic-like substances. *Environmental Science & Technology*, 2016, 50: 7443-7452.
- 385 Huang M, Li Z, Huang B, et al., Investigating binding characteristics of cadmium and copper to DOM derived from compost and rice straw using EEM-PARAFAC combined with two-dimensional FTIR correlation analyses. *Journal of Hazardous Materials*, 2018, 344: 539-548.
- Kalbitz K, Geyer S and Geyer W, A comparative characterization of dissolved organic matter by means of original aqueous samples and isolated humic substances. *Chemosphere*, 2000, 40: 1305-1312.
- 390



- Kirillova E N, Andersson A, Han J, et al., Sources and light absorption of water-soluble organic carbon aerosols in the outflow from northern China. *Atmospheric Chemistry and Physics*, 2014, 14: 1413-1422.
- Lakowicz J R, *Principles of Fluorescence Spectroscopy* Springer. US, 2006.
- Li J, Chen Q, Sha T, et al., Significant promotion of light absorption ability and formation of triplet organics and reactive oxygen species in atmospheric hulis by Fe(III) ions. *Environmental Science & Technology*, 2022,
- 395 Lin M and Yu J Z, Effect of metal-organic interactions on the oxidative potential of mixtures of atmospheric humic-like substances and copper/manganese as investigated by the dithiothreitol assay. *Science of the Total Environment*, 2019, 697: 134012.
- Lin M and Yu J Z, Assessment of interactions between transition metals and atmospheric organics: ascorbic acid depletion and hydroxyl radical formation in organic-metal mixtures. *Environmental Science & Technology*, 2020, 54: 1431-1442.
- 400 Lin P, Engling G and Yu J Z, Humic-like substances in fresh emissions of rice straw burning and in ambient aerosols in the Pearl River Delta Region, China. *Atmospheric Chemistry and Physics*, 2010, 10: 6487-6500.
- Ma L, Li Z, Yabo S D, et al., Insight into the interaction between heavy metals and water-soluble organic compounds in PM_{2.5} affected by heavy haze using ultraviolet-visible and fluorescence spectra combined with two-dimensional correlation spectroscopy. *Journal of Cleaner Production*, 2022, 362: 132476.
- 405 Ma Y, Cheng Y, Qiu X, et al., Optical properties, source apportionment and redox activity of humic-like substances (HULIS) in airborne fine particulates in Hong Kong. *Environmental Pollution*, 2019, 255: 113087.
- Mao J, Fan S, Jacob D J, et al., Radical loss in the atmosphere from Cu-Fe redox coupling in aerosols. *Atmospheric Chemistry and Physics*, 2013, 13: 509-519.
- 410 Mo Y, Li J, Cheng Z, et al., Dual carbon isotope-based source apportionment and light absorption properties of water-soluble organic carbon in PM_{2.5} over China. *Journal of Geophysical Research: Atmospheres*, 2021, 126: e2020JD033920.
- Paris R and Desboeufs K V, Effect of atmospheric organic complexation on iron-bearing dust solubility. *Atmospheric Chemistry and Physics*, 2013, 13: 4895-4905.
- Parlanti E, Worz K, Geoffroy L, et al., Dissolved organic matter fluorescence spectroscopy as a tool to estimate biological activity in a coastal zone submitted to anthropogenic inputs. *Organic Geochemistry*, 2000, 31: 1765-1781.
- 415 Qin J, Zhang L, Qin Y, et al., pH-Dependent chemical transformations of humic-like substances and further cognitions revealed by optical methods. *Environmental Science & Technology*, 2022, 56: 7578-7587.
- Qin J, Zhang L, Zhou X, et al., Fluorescence fingerprinting properties for exploring water-soluble organic compounds in PM_{2.5} in an industrial city of northwest China. *Atmospheric Environment*, 2018, 184: 203-211.
- 420 Ryan, David, K., et al., Comparison of Mn²⁺, Co²⁺, and Cu²⁺ binding to fulvic acid as measured by fluorescence quenching. *Canadian Journal of Chemistry*, 1983, 61: 1505-1509.
- Saleh R, Robinson E S, Tkacik D S, et al., Brownness of organics in aerosols from biomass burning linked to their black carbon content. *Nature Geoscience*, 2014, 7: 647-650.



- Scheinhardt S, Müller K, Spindler G, et al., Complexation of trace metals in size-segregated aerosol particles at nine sites in
425 Germany. *Atmospheric Environment*, 2013, 74: 102-109.
- Slikboer S, Grandy L, Blair S L, et al., Formation of light absorbing soluble secondary organics and insoluble polymeric
particles from the dark reaction of catechol and guaiacol with Fe(III). *Environmental Science & Technology*, 2015, 49:
7793-7801.
- Strickler S J and Berg R A, Relationship between absorption intensity and fluorescence lifetime of molecules. *The Journal of*
430 *Chemical Physics*, 1962, 37: 814-822.
- Verma V, Rico-Martinez R, Kotra N, et al., Contribution of water-soluble and insoluble components and their
hydrophobic/hydrophilic subfractions to the reactive oxygen species-generating potential of fine ambient aerosols.
Environmental Science & Technology, 2012, 46: 11384-11392.
- Wang H, Zhang L, Huo T, et al., Application of parallel factor analysis model to decompose excitation-emission matrix
435 fluorescence spectra for characterizing sources of water-soluble brown carbon in PM_{2.5}. *Atmospheric Environment*, 2020,
223: 117192.
- Wang X, Qin Y, Qin J, et al., Spectroscopic insight into the pH-dependent interactions between atmospheric heavy metals
(Cu and Zn) and water-soluble organic compounds in PM_{2.5}. *Science of the Total Environment*, 2021, 767: 145261.
- Win M S, Tian Z, Zhao H, et al., Atmospheric HULIS and its ability to mediate the reactive oxygen species (ROS): A review.
440 *Journal of Environmental Science (China)*, 2018, 71: 13-31.
- Xiao K, Han B, Sun J, et al., Stokes shift and specific fluorescence as potential indicators of organic matter hydrophobicity
and molecular weight in membrane bioreactors. *Environmental Science & Technology*, 2019, 53: 8985-8993.
- Xiao K, Shen Y, Sun J, et al., Correlating fluorescence spectral properties with DOM molecular weight and size distribution
in wastewater treatment systems. *Environmental Science: Water Research & Technology*, 2018, 4: 1933-1943.
- 445 Xiao K, Yu J, Wang S, et al., Relationship between fluorescence excitation-emission matrix properties and the relative
degree of DOM hydrophobicity in wastewater treatment effluents. *Chemosphere*, 2020, 254: 126830.
- Yamashita Y and Jaffe R, Characterizing the interactions between trace metals and dissolved organic matter using
excitation-emission matrix and parallel factor analysis. *Environmental Science & Technology*, 2008, 42: 7374-7379.
- Yang W, Zhang T, Han C, et al., Photoenhanced heterogeneous reaction of O₃ with humic acid: Focus on O₃ uptake and
450 changes in the composition and optical property. *Environmental Pollution*, 2021, 268: 115696.
- Ye C, Chen H, Hoffmann E H, et al., Particle-phase photoreactions of hulis and tmis establish a strong source of H₂O₂ and
particulate sulfate in the winter north china plain. *Environmental Science & Technology*, 2021, 55: 7818-7830.
- Yu H, Wei J, Cheng Y, et al., Synergistic and antagonistic interactions among the particulate matter components in
generating reactive oxygen species based on the dithiothreitol assay. *Environmental Science & Technology*, 2018, 52: 2261-
455 2270.



Yu J, Xiao K, Xue W, et al., Excitation-emission matrix (EEM) fluorescence spectroscopy for characterization of organic matter in membrane bioreactors: Principles, methods and applications. *Frontiers of Environmental Science & Engineering*, 2020, 14: 31.

460 Zanca N, Lambe A T, Massoli P, et al., Characterizing source fingerprints and ageing processes in laboratory-generated secondary organic aerosols using proton-nuclear magnetic resonance ($^1\text{H-NMR}$) analysis and HPLC HULIS determination. *Atmospheric Chemistry and Physics*, 2017, 17: 10405-10421.

Zhu Y, Jin Y, Liu X, et al., Insight into interactions of heavy metals with livestock manure compost-derived dissolved organic matter using EEM-PARAFAC and 2D-FTIR-COS analyses. *Journal of Hazardous Materials*, 2021, 420: 126532.