

Numerical Simulation of Aerosol Concentration Effects on Cloud Droplet Size Spectra Evolutions of Warm Stratiform Clouds in Jiangxi, China

Yi Li^{1,2}, Xiaoli Liu^{1,2*}, Hengjia Cai^{1,2}

¹ China Meteorological Administration Aerosol-Cloud and Precipitation Key Laboratory, Nanjing University of Information Science and Technology, Nanjing 210044, China.

² College of Atmospheric Physics, Nanjing University of Information Science and Technology, Nanjing 210044, China.

Correspondence to: Xiaoli Liu (liuxiaoli2004y@nuist.edu.cn)

Abstract.

Changes in aerosol amount and size distribution significantly impact cloud droplet size distribution, as aerosols act as cloud condensation nuclei (CCN) and influence the relative dispersion (ϵ) of cloud droplet spectra. Relative dispersion plays a key role in parameterizing cloud processes in general circulation models (GCMs) and microphysical schemes, affecting precipitation estimates and climate predictions. However, the effect of varying aerosol modes on cloud microphysics remains debated, depending on thermodynamic conditions and cloud type. This study simulates a warm stratiform cloud in Jiangxi,

China, using the WRF-SBM model from 18:00 on December 24, 2014, to 06:00 (UTC) on December 25, 2014. Satellite and aircraft observations were used to validate the simulation, showing good agreement in cloud structure. Sensitivity experiments were conducted by increasing nucleation, accumulation, and coarse-mode aerosols by fivefold and reducing the total aerosol concentration to one-fifth of the control. Results show that higher aerosol concentrations enhance cloud formation and broaden droplet spectra, while lower concentrations suppress cloud development. Accumulation mode aerosols increase small droplet concentrations, while nucleation and coarse-mode aerosols favor larger droplets. The correlation between ϵ and volume-weighted radius (R_v) shifts from positive to negative as R_v increases. This transition is driven by cloud droplet collision-coalescence, condensation, and activation. Increased accumulation mode aerosol concentrations shift the ϵ - R_v correlation from

批注【懿李1】: Referee2: 1) The title is not clear. What is the meaning of spectral evolutions?

批注【懿李2】: 2) The Abstract: The abstract needs to be rewritten in a more precise and accurate manner, as it contains several unclear and incomplete statements. For example, consider the first sentence: "Aerosols, as cloud condensation nuclei (CCN), may impact cloud droplet spectrum relative dispersion (ϵ), affecting precipitation and climate change." This should be rewritten to provide more detail and accuracy, such as: "Changes in aerosol amount and size distribution may impact the size distribution of cloud droplets (since they serve as CCN), as well as their relative dispersion. Relative dispersion is a key factor used to parameterize cloud processes in general circulation models (GCMs) and bulk microphysical schemes, and thus influences precipitation estimates and various aspects of climate predictions." Another example is the second sentence: "However, the influence of various aerosol modes on cloud microphysics remains controversial, and this effect varies with area and cloud type." Again, this statement is unclear and inaccurate. It should be rewritten as: "However, the influence of changes in the concentration of various aerosol modes on cloud microphysical processes is still a subject of debate, and it depends significantly on thermodynamic conditions and cloud type."

negative to positive in the R_v range of 4.5-8 μm , while reduced aerosol concentrations strengthen the negative correlation. Regardless of different coalescence intensity, ε converges with the increase in number concentration of cloud droplet (N_c).

25 1 Introduction

According to Lau and Wu (2003), warm clouds account for 32% of total precipitation in tropical regions and cover 72% of the total precipitation area. Warm clouds play a critical role in evaluating cloud-precipitation-climate feedback, making the understanding of their formation, development, and cloud microphysical processes a crucial topic in cloud physics (Zhao and Ishizaka, 2004; Seifert and Onishi, 2010).

30 Grosvenor et al. (2018) identified a significant relationship between cloud droplet number concentration (N_c), cloud optical thickness, and cloud top temperature, proposing an improved remote sensing retrieval algorithm for cloud droplet effective radius to reduce the errors in satellite measurements of N_c . Zheng et al. (2021) utilized merged Cloud Sat-CALIPSO-MODIS products to compare the macro- and microphysical properties of precipitating and non-precipitating clouds during the warm season in central-eastern China, focusing on parameters such as cloud optical thickness and the effective radius of cloud
35 droplets. Their findings indicated that the probability of precipitation increased with the increasing of cloud optical thickness, liquid water path, and ice water path, but showed a decreasing trend when the cloud droplet effective radius exceeded 22 micrometres. However, these studies have overlooked the impact of changes in particle size distribution in clouds, which may be critical for parameterization of cloud droplet effective radius and is an essential factor that cannot be ignored during cloud-rain auto-conversion processes, affecting macroscopic and microscopic physical processes in clouds (Wang and Lu, 2022; Xie
40 et al., 2015).

Cloud droplet spectral relative dispersion (ε) is an important parameter that describes the width and distribution of cloud droplet sizes. It is represented as the ratio between the standard deviation (σ) and the mean radius (R_{ave}) of the droplets (Wang

and Lu, 2022). On one hand, ϵ (the relative dispersion of cloud droplet spectra) influences the effective radius (R_e) of cloud droplets by altering their size distribution. A higher ϵ typically leads to a broader droplet size distribution, which increases the effective radius, thereby enhancing the auto-conversion process that drives the formation of precipitation from cloud droplets (Liu et al., 2005, 2006; Zhu et al., 2020; Lu and Xu, 2021; Wang et al., 2022; Wang et al., 2023; Yang et al., 2023). On the other hand, ϵ modulates cloud-aerosol interactions by affecting cloud microphysical properties, such as droplet concentration and liquid water content, which in turn influences cloud radiative properties and, consequently, climate (Xie et al., 2017).

Many researchers have conducted causal analyses on the uncertainty of the effect of cloud microphysical properties on ϵ . The results indicate that the variability of ϵ is influenced by various factors, such as atmospheric temperature, humidity, and entrainment (Lu et al., 2013). Zhu et al. (2020) analysed data from a flight observation conducted in Monterey, California, in July 2008 as part of the US POST (Physics of Stratocumulus Top) project, found that in adiabatic clouds, vertical velocity plays a dominant role, and an increase in vertical velocity promotes the activation of cloud condensation nuclei (CCN), leading to an increase in N_c and facilitating droplet coalescence and growth. On the other hand, Kumar et al. (2017) conducted idealized simulation experiments using direct numerical simulation (DNS) to study the mixing dynamics at cloud edges and their impact on the droplet size distribution (DSD). They showed that ϵ is also related to turbulent mixing and variations in vertical velocity within the cloud.

However, as Lu et al. (2020) pointed out, existing studies on ϵ primarily rely on empirical data from observations, leading to significant uncertainty in characterizing the ϵ within clouds. In addition, the relationship between the ϵ and the volume-mean radius (R_v) has shown varied conclusions in different studies: some indicate a negative correlation (Liu et al., 2008; Pandithurai et al., 2012), while others suggest a positive correlation (Tas et al., 2012). It is also found that as R_v increases, the ϵ exhibits a converging trend (Chen et al., 2016). Meanwhile, the correlation between the ϵ and N_c also shows uncertainty. Jin et al. (2021) conducted aircraft observational studies on stratiform warm clouds in Jiangxi, China, indicating that ϵ in both

批注【审稿3】: Referee2: 3) The introduction: line 47 states: “On the one hand, ϵ influences the effective radius of cloud droplets and the auto-conversion process, thereby affecting cloud precipitation processes On the other hand, impacting climate” This description is partial and unclear. Phrases like “ ϵ influences the effective radius of cloud droplets” and “ ϵ affects cloud-aerosol interactions” lack specificity. These statements need to be rewritten for clarity and accuracy.

批注【审稿4】: Referee2: 3) The introduction: line 47 states: “On the one hand, ϵ influences the effective radius of cloud droplets and the auto-conversion process, thereby affecting cloud precipitation processes On the other hand, impacting climate” This description is partial and unclear. Phrases like “ ϵ influences the effective radius of cloud droplets” and “ ϵ affects cloud-aerosol interactions” lack specificity. These statements need to be rewritten for clarity and accuracy.

precipitating and non-precipitating warm clouds is negatively correlated with N_c . But, some studies report a positive correlation (Pandithurai et al., 2012; Chen et al., 2016), while others indicate a negative correlation (Cecchini et al., 2017; Wang et al., 2011). Some studies even suggest that no significant correlation is observed between ε and N_c (Tas et al., 2015). Meanwhile, the correlation between ε and N_c also shows uncertainty. Jin et al. (2021) conducted aircraft observational studies on stratiform warm clouds in Jiangxi, China, indicate that ε in both precipitating and non-precipitating warm clouds is negatively correlated with N_c . Similarly, Cecchini et al. (2017) and Wang et al. (2011) reported negative correlations between ε and N_c . However, some studies report a positive correlation (Pandithurai et al., 2012; Chen et al., 2016). While some studies even suggest that no significant correlation is observed between ε and N_c (Tas et al., 2015).

Studies by Ma et al. (2010) and Wang et al. (2011, 2019) have shown that changes in ε are highly sensitive to aerosol concentration and its activation process. Additionally, alterations in aerosol concentration or size distribution significantly impact the cloud-rain auto-conversion process through ε changes. Consequently, ε becomes a critical link connecting the aerosol-cloud interaction effects (Liu and Daum, 2002).

Liu et al. (2003) compared aircraft observations and satellite retrievals for stratiform warm clouds in both the northern and southern hemispheres and found that an increase in aerosol concentration leads to a decrease in cloud droplet effective radius and narrowing of the droplet spectrum, thus suppressing warm precipitation processes. Fan et al. (2012) conducted a numerical simulation on variations of aerosol concentration in Eastern China, demonstrating that an increase in CCN leads to an increase in N_c and cloud droplet mass concentration, reduces the number concentration of raindrops, and delays the onset of precipitation. Yang et al. (2017) analysed aerosol concentration and cloud droplet spectrum distribution in the high-altitude region of eastern China during summer, and the results showed that increased aerosol concentration inhibits the cloud-rain auto-conversion process, resulting in more cloud water remaining in the atmosphere and reducing warm precipitation. By analysing the aerosol observations in India from 2000 to 2017, Kant et al. (2019) found that strong updrafts with abundant

批注 [龔季5]: Referee2: 4) The introduction: Throughout the entire introduction, it is crucial to specify the exact type of cloud being studied in each referenced work. While "warm shallow clouds" is a broad term encompassing various cloud types, it is important to clearly identify the specific cloud types under investigation. This distinction is necessary because different cloud types are governed by unique controlling processes, which significantly influence their formation, behavior, and interaction with aerosols. Moreover, throughout the paper, it should be clear what specific type of cloud is being simulated and studied. Simply referring to "warm shallow clouds" or warm processes is insufficient and could lead to ambiguity in interpreting the results. Precise terminology should be used consistently to describe the cloud type under investigation in this study. This will ensure that the findings are accurately interpreted and compare with previous studies on similar cloud types. Adjustments should be made to the manuscript to address this issue.

85 mineral dust aerosols can activate more cloud droplets, leading to competition for water vapor and narrowing the droplet spectrum, limiting the growth of high-level liquid droplets. It is suggesting that an increase in aerosol concentration leads to a reduction in ϵ , thereby inhibiting the cloud-rain auto-conversion process (Chandrakar et al., 2016, 2018; Desai et al., 2019).

However, there are also studies indicating that an increase in aerosol concentration results in an increase in ϵ and enhances droplet collision-coalescence processes (Rotstajn and Liu, 2003; Yum and Hudson, 2005; Rotstajn and Liu, 2009; Prabha et al., 2012; Liu et al., 2020). For instance, Liu et al. (2020) found that increasing aerosol concentration in clean tropical or marine regions can prolong cloud lifetimes and enhance precipitation by modifying the cloud droplet spectrum distribution. Moreover, it is found that the influence of aerosol concentrations on cloud droplet size distribution exhibits strong regional dependence, varies according to cloud types and geographical regions (Chandraka et al., 2016; 2018).

In addition, the impact of aerosol concentrations on cloud droplet spectrum varies for different size ranges of aerosols. Liu et al. (2022), using satellite data to investigate the influence of aerosols on warm rain processes, found that fine particles with diameters ranging from 0.1 to 2.5 micrometres, acting as cloud condensation nuclei, can suppress precipitation and prolong the lifetime of warm stratiform clouds and shallow maritime cumulus clouds, like the conclusions of Kovačević (2018) and Lerach and Cotton (2018). On the other hand, an increase in coarse-mode marine condensation nuclei with larger particle sizes leads to a noticeable increase in cloud droplet effective radius and warm rain intensity. It is found that large particles with diameters exceeding 2 micrometres, acting as giant cloud condensation nuclei, can increase ϵ and facilitate cloud droplet growth during the collision-coalescence process (Yin et al., 2000; Jensen and Nugent, 2017). However, Wehbe et al. (2020) analysed aircraft observations over the United Arab Emirates in 2019, and found that although giant cloud condensation nuclei were present, no significant collision-coalescence process was observed in marine warm stratiform clouds.

批注 [懿李6]: Referee2: 4) The introduction: Throughout the entire introduction, it is crucial to specify the exact type of cloud being studied in each referenced work. While "warm shallow clouds" is a broad term encompassing various cloud types, it is important to clearly identify the specific cloud types under investigation. This distinction is necessary because different cloud types are governed by unique controlling processes, which significantly influence their formation, behavior, and interaction with aerosols. Moreover, throughout the paper, it should be clear what specific type of cloud is being simulated and studied. Simply referring to "warm shallow clouds" or warm processes is insufficient and could lead to ambiguity in interpreting the results. Precise terminology should be used consistently to describe the cloud type under investigation in this study. This will ensure that the findings are accurately interpreted and compare with previous studies on similar cloud types. Adjustments...

批注 [懿李7]: Referee2: 4) The introduction: Throughout the entire introduction, it is crucial to specify the exact type of cloud being studied in each referenced work. While "warm shallow clouds" is a broad term encompassing various cloud types, it is important to clearly identify the specific cloud types under investigation. This distinction is necessary because different cloud types are governed by unique controlling processes, which significantly influence their formation, behavior, and interaction with aerosols. Moreover, throughout the paper, it should be clear what specific type of cloud is being simulated and studied. Simply referring to "warm shallow clouds" or warm processes is insufficient and could lead to ambiguity in interpreting the results. Precise terminology should be used consistently to describe the cloud type under investigation in this study. This will ensure that the findings are accurately interpreted and compare with previous studies on similar cloud types. Adjustments...

Furthermore, Rosenfeld et al. (2001) attributed the reduction in cloud droplet effective radius over the Sahara Desert to numerous submicron-sized cloud condensation nuclei (CCN), which decreased ε exacerbated the decrease in precipitation over the Sahara region. Numerical experiments by Flossmann and Wobrock (2010) yielded similar conclusions.

In summary, under the context of climate change, changes in the physicochemical properties of aerosols significantly affect the microphysical characteristics of warm clouds. Existing studies often rely on exploring the relationships between aerosol concentration and microphysical cloud quantities such as N_c and R_v , and further research on ε , a key factor affecting the cloud-aerosol effect, is still needed. However, the response of warm clouds to aerosol physicochemical properties depends on the region and cloud type, and due to limitations in observational methods, the response of ε to changes in aerosol concentration varies significantly across studies, making this issue a crucial and controversial scientific question in climate prediction.

This study utilizes the SBM-FAST bin microphysics scheme within the Weather Research and Forecasting (WRF) model to simulate a stratiform warm cloud event in Jiangxi, China. The numerical experiments aim to explore the impacts of changes in nucleation, accumulation, coarse, and total mode aerosol concentrations on the macroscopic and microscopic characteristics of stratiform warm clouds in this region. The paper is organized as follows: Section 2 outlines the numerical simulation setup, aircraft, and satellite observations to validate simulation results, and the computational formulas used in the analysis, the third section 3 conducts validations of the control experiment's simulation results through comparisons with concurrent aircraft and satellite cloud top temperature observations, uncovering the effects of different aerosol modes on the macroscopic and microscopic physical properties of clouds, with a particular focus on the correlation between ε and cloud microphysical properties. The last two section include the discussion and conclusions.

2 Model Introduction and Experiment Design

2.1Simulation Setup and Weather Conditions

125 This paper selects a stratiform warm cloud process that occurred in the Jiangxi, China on December 25, 2014, and conducts simulations using the WRF (Weather Research and Forecasting) 4.2 version. The experiment comprises one control and five aerosol spectrum modification experiments. Except for aerosol concentrations, all groups keep the initial field data and simulation settings consistent. The simulations use the fifth generation of ECMWF atmospheric reanalyses of the global climate (ERA5) hourly data on pressure levels as the initial field, with a resolution of $0.25^{\circ} \times 0.25^{\circ}$.


130 The simulations employ a two-layer nesting approach with 3 km and 1 km grid resolutions. The model is divided vertically into 57 layers, reaching a top pressure level of 50 hPa. In the first two kilometers above ground level, the vertical grid resolution varies as follows: the lowest layer is at approximately 401 meters, with subsequent layers at 457 meters, 528 meters, 618 meters, 733 meters, 876 meters, 1053 meters, 1270 meters, 1533 meters, 1845 meters, and 2209 meters. And the innermost layer grid measure contains 376×376 grid points. The microphysics scheme used is the new version of SBM-fast bin scheme (FSBM-2) under the WRF 4.2 version. The boundary layer scheme selected is the Mellor-Yamada-Janjic (Eta) Turbulence Kinetic Energy (TKE) scheme, and the near-surface layer scheme uses the Monin-Obukhov (Janjic Eta) scheme. The land surface process adopts the unified Noah land-surface model. The (old) Goddard shortwave radiation scheme is used, and the Rapid Radiative Transfer Model (RRTM) scheme is chosen for longwave radiation.

The simulation region is illustrated in Figure 1, and the simulation duration is from 18:00 on December 24, 2014, to 06:00 on December 25, 2014 (UTC), with no precipitation was observed at the ground during the simulation period. The simulated area, Ganzhou City, is in the southern part of eastern China's Jiangxi province. It is located upstream of the Gan River and in the transitional zone between the southeastern coastal and central inland regions. The city is surrounded by mountains, with

批注 [章节8]: Referee1: L131 It is still unclear what is the vertical grid resolution in meters, this is relevant to understand how many grid points there are in the cloud region. If not constant you could give a range for the first kilometers. The discussion of this potential modeling issue could be valuable, and motivate future work to elucidate its importance.

145 faulted basins traversed. The predominant topographical features are mountains, hills, and basins. The area is located at the southern edge of the subtropical zone and falls under the subtropical monsoon climate region.

150 As shown in Figure 2, at the 500 hPa level, the isobars over southern Jiangxi are relatively flat, indicating the absence of significant trough or ridge systems. This suggests that the upper atmosphere in this region is under stable airflow control. A strong westerly jet is observed at this level, indicating the presence of a notable westerly jet stream aloft. At the 850 hPa level, a cyclonic circulation is evident, pointing to convergence in the lower atmosphere. As the warm cloud system develops, the convergence intensifies, and the associated low-pressure system shifts eastward, providing favorable upward motion in the region. At the surface, a pronounced low-pressure system is present. As the process evolves, wind speeds increase, and the low-pressure system progresses southward. These features indicate that southern Jiangxi is dominated by a low-pressure system at the lower levels during this period, with dynamic conditions favorable for cloud development.



155 **Figure 1: Simulated Region and Nesting Configuration.** The shading represents the elevation (m) of the terrain, and the area within the red box is the analysis range.

批注 [懿李9]: Referee2: 6) To enhance the clarity of the study, I recommend including more comprehensive meteorological data for the event under investigation. The current version of them anuscript presents a single snapshot of the 500 mb height and 700 mb winds, which is insufficient for a thorough analysis of the conditions affecting cloud formation and development. Adding detailed meteorological context will greatly assist in interpreting the results and understanding the processes involved in cloud dynamics. Adding 850 mb and surface pressure maps will provide insights into low-level dynamics and synoptic influences affecting cloud formation. Instead of a single snapshot, present meteorological data at multiple time points to show how conditions evolve during the event, correlating with cloud development. Include radiosonde data to provide thermal and humidity profiles, helping to understand atmospheric stability and conditions relevant to cloud formation.




Figure 2: The 500 hPa, 850 hPa, and surface geopotential height fields (blue contour lines, unit: dagpm) and the wind fields (wind barbs) at 00:00 (UTC) on December 25, 2014. The area within the red box indicates the starting point of the flight.

2.2 Introduction to Microphysics Scheme

The SBM-fast scheme was initially developed by Khain and Lynn (2009) as a simplified version of the SBM-full bin scheme based on the original microphysics scheme included in the Hebrew University Cloud Model (HUCM) (FSBM-1) (Khain and Sednev, 1996; Khain et al., 2000). The FSBM-2 used in this study is an improvement over FSBM-1 by Shpund et al. (2019)

and has been verified to exhibit better simulation performance (Han et al., 2019).

In FSBM-2, cloud and rain droplets are described using a unified liquid droplet bin scheme, which is divided into 33 bins. The aerosol scheme is divided into marine and continental components, and the aerosol spectrum distribution is described using 43 or 33 mass bins. Regardless of whether 33 or 43 aerosol bins are used, the maximum dry aerosol radius is set to 2 μ m. The scheme activates aerosols into liquid droplets under supersaturation conditions (cloud nucleation: Pinsky and Khain, 2018).
170 In the model, the minimum CCN size is assumed to be 0.003 μ m, and the initial aerosol distribution is represented by the sum of three lognormal distributions, corresponding to the nucleation mode (centered at 0.008 μ m), accumulation mode (centered at 0.034 μ m), and coarse mode (centered at 0.46 μ m). The calculation of cloud droplet nucleation considers the effect of supersaturation, and the algorithm's accuracy is verified through comparison with large-eddy simulation results (Iltoviz et al., 2015).

175 **2.3 Sensitivity Experiment Configuration**

This paper includes five aerosol concentration modification experiments and one control experiment (ORG). The initial aerosol concentrations set in the control experiment, are shown in Table 1. The initial aerosol concentrations are modified for the other five experiment, as shown in Table 2. According to the aircraft observational study on the impact of aerosol concentration changes on precipitation in Eastern China (Qian et al., 2009) and the numerical simulation study on the effect of aerosol concentration changes on clouds and precipitation in Eastern China (Fan et al., 2012), increasing the initial aerosol concentration to five times realistically reflects the background concentration of continental aerosols under polluted conditions in Eastern China. This adjustment is beneficial for demonstrating the realistic impacts of aerosol concentration changes on stratiform warm clouds in eastern China. Experiments 1, 2, and 3, modify the aerosol concentrations of the nucleation mode (NM), accumulation mode (AM), and coarse mode (CM) to five times their original values, respectively. Experiment 4 (ITM) simultaneously modifies the aerosol concentrations of the nucleation, accumulation, and coarse modes to five times their
185

original values, and experiment 5 (DTM) reduces the aerosol concentrations by five times compared to the original group. The initial background aerosol spectrums in the simulations are shown in Figure 3.

Table 1: Initial Aerosol Concentration in the Control Experiment.

Aerosol Types	Number Concentration (cm ⁻³)	Mean Particle Size (μm)
Nucleation Mode	1000	0.008
Accumulation Mode	800	0.034
Coarse Mode	0.720	0.460

190

Table 2: Initial Aerosol Concentration Settings in Sensitivity Experiments.

	Nucleation Mode (cm ⁻³)	Accumulation Mode (cm ⁻³)	Coarse Mode (cm ⁻³)
Experiment 1 (NM)	5000	800	0.720
Experiment 2 (AM)	1000	4000	0.720
Experiment 3 (CM)	1000	800	3.600
Experiment 4 (ITM)	5000	4000	3.600
Experiment5 (DTM)	200	160	0.144




Figure 3: Initial aerosol number concentration (unit: cm^{-3}) as a function of particle diameters (unit: nm).

195 2.4 Calculation of Cloud Droplet Spectrum Parameters

In this study, the changes in cloud droplet spectrum and cloud droplet spectral parameters were analysed. The mean cloud droplet Radius (R_m), R_v , liquid water path (LWP), cloud-rain auto-conversion threshold (T), ε , and cloud droplet activation intensity (Fbs) were calculated as shown in Supplement.

2.5 Introduction of Data

200 2.5.1 Introduction of Aircraft Observation Data

The aircraft observation data used in this study were sourced from a flight observation mission conducted in Jiangxi, China on December 25, 2014. Observations were carried out using the Yun-12 aircraft equipped with a comprehensive set of aerosol-cloud-precipitation detectors. The cloud microphysical data were obtained from the Cloud-Aerosol Spectrometer (CAS), while

flight altitude and path information were obtained from the Aircraft Integrated Meteorological Measurement System (AIMMS-
205 20). To ensure the accuracy and reliability of the observation data, all probes and the observation platform were precisely
calibrated prior to the observations, and outliers were removed from the post-observation data.

The observation flight area was located above Ganzhou City in Jiangxi Province, spanning coordinates from 114.0°E to
117.0°E and from 25°N to 27°N. The aircraft took off from Ganzhou Airport and followed a flight pattern that included
ascending, cruising, and spiralling down. The flight lasted from 01:29 to 04:45 (UTC), reaching a maximum altitude of 4126
210 meters. To exclude data from non-cloud areas during the observation period, a cloud region criterion of cloud liquid water
content (Clw) > 0.001 g/m³ and number concentration of cloud droplets (Nc) > 10 particles cm⁻³ was applied (Jin et al. 2021,
Wang et al., 2024).

2.5.2 Introduction to Satellite-Observed Cloud Top Temperature Data

This study utilizes the standard format cloud-top temperature scan data from the Modis (Aqua) satellite, with the
215 satellite passing over the simulation area at 05:30 (UTC) on December 25, at a resolution of 5° × 5°. The Aqua satellite
is part of NASA's Earth Observing System (EOS) and focuses on monitoring the Earth's water cycle. Equipped with
the Modis sensor, which has multispectral imaging capabilities ranging from visible to infrared wavelengths, Aqua
provides high-resolution information on the Earth's surface and atmosphere. It is widely used in meteorological,
oceanographic, and environmental research (Platnick et al., 2015).3 Results and analysis

220 3.1 Simulation Results Validation

To verify the simulation performance of the control experiment, we compared the simulated results with cloud-top temperature
observations from the Modis (Aqua) satellite and aircraft observations on December 25, 2014.

Aircraft observation data on December 25, 2014, in Jiangxi region was chosen to further validate the simulated vertical
distribution of cloud microphysical characteristics. The data was obtained from the CAS probe onboard the aircraft, which
225 measures aerosols and cloud particles with diameters ranging from 0.51 to 50 μm, covering 30 bins with varying size bins.

批注 [懿李10]: Referee1: Fig 4 is the same as Fig 6 a, right? Maybe
remove Fig 4.

The observation period was from 01:35 to 04:45 (UTC) on December 25, 2014. During the observation period, the stratiform warm cloud within the flight area had a maximum horizontal extent of over 50 kilometres, and it was characterized as a stratiform warm cloud process. For the control experiment, the cloud water content, cloud droplet number concentration, and average cloud droplet diameter were compared in the same observation duration and flight regions.

Figure 4 shows the flight trajectory and the cloud liquid water content along the observation path. To validate the control experiment's simulation results, a comprehensive cloud penetration segment from 04:10 to 04:20 UTC was selected. During this period, the Clw and Nc were calculated and compared with the simulation results in the same region and time frame, as illustrated in Figure 5. To minimize the impact of differing vertical resolutions between aircraft observation data and model simulations, the cloud base height within the validation interval was set to 0 and the cloud top height to 1, thus achieving height normalization. In the control experiment, the vertical distribution of Clw aligns well with the aircraft observations, both exhibiting a trend of increasing with height at first and then decreasing. The Clw values are relatively close between the simulation and observations below the mid-cloud region, but near the cloud top, the simulated Clw is slightly higher than the aircraft observations. Nevertheless, the maximum Clw in both cases occurs in the mid-cloud region. Additionally, the Nc in the control experiment is similar to the aircraft observations, with both showing no significant variation with height. The largest discrepancy appears in the Clw. This discrepancy may be due to the inherent errors in numerical simulations, as well as differences in the resolution and the cloud sampling between the simulation and the aircraft observations. The spatial resolution of aircraft measurements can be as fine as a few meters to tens of meters, while the numerical model's minimum grid resolution is 1 km. Therefore, the simulation results represent an average over a larger area within a similar temporal and spatial range. Moreover, the aircraft's flight path through the cloud may not fully penetrate the entire cloud structure, with data possibly being collected from a narrower spatial range or from the cloud's edges. As a result, the observations may reflect the cloud's internal microphysical properties.

Figure 6 compares the cloud-top temperatures between the control experiment and the simulation results. During the simulation period, the Modis (Aqua) satellite passed over the study region at 5:30 (UTC). To ensure consistency with the area observed by the aircraft, the comparison of cloud-top temperatures focuses on the region with concentrated warm clouds in southwestern Jiangxi. Both the control experiment and satellite observations show cloud-top temperatures in the range of 0-10°C, with higher temperature areas (5-10°C) located west of 114°E. The results indicate that the observed cloud-top temperature distribution is consistent with the simulation results.

Figure 7 also compares the liquid water path (LWP) values observed by the ground-based microwave radiometer during the period of vigorous cloud development between 04:00 and 05:00 UTC. The RPG-HATPRO microwave radiometer features two bands: 22-31 GHz (7-channel filter-bank humidity profiler and LWP radiometer) and 51-58 GHz (7-channel filter-bank temperature profiler) (Liu et al. 2014). The microwave radiometer was located at 114.95°E, 25.85°N, and the simulation results were taken from the corresponding area centered on the radiometer. The results show a slight difference in the average LWP between the control experiment and the radiometer observations. However, the observed LWP values generally fall within the range of the simulated LWP, and their temporal distribution is largely consistent with the simulation. Overall, regarding the distribution of stratiform warm clouds and the vertical distribution of cloud microphysical properties, the simulation results are generally consistent with the observed data. Therefore, the simulation results are reliable.

批注 [懿李11]: Referee2: 7) Results: Fig 5: The figure presents the validation of the simulated cloud top temperature against satellite measurements. There are significant differences between the two that should be addressed and explained. The validation is a bit questionable.




Figure 4: Aircraft flight trajectory and cloud liquid water content (Clw) within the cloud region along the observation path. The red box indicates a comprehensive cloud penetration process from 04:10 to 04:20 UTC.




Figure 5: Aircraft observations (scatter points) of cloud liquid water content (Clw, in g/m^3), cloud droplet number concentration (N_c , in cm^{-3}) on December 25, 2014, compared with model simulations (black dashed lines) of Clw and N_c .




Figure 6: Cloud-top temperatures observed by the Modis (Aqua) satellite (05:30 UTC) and simulated by the control experiment on December 25, 2014.




Figure 7: Comparison of hourly simulated regionally averaged liquid water path (LWP, unit: kg/m²) and the liquid water path observed by the microwave radiometer.

3.2 Analysis of the Impact of Background Aerosols on Warm Cloud Properties

3.2.1 Vertical Distribution of Cloud Microphysical Properties

275 Figure 8-10 reflects that the cloud thickness significantly increases as the cloud system develops. Both Clw and Rm increase with height, show high consistency. In contrast, Nc exhibits different trends with height at different times. From 00:00 to 02:00 UTC, when the cloud system is in initial stage of development, Nc decreases with height, and many small cloud droplets appear at the cloud base, which is the main area for droplet activation. As the cloud system further develops, from 03:00 to 04:00 UTC, Nc shows relative uniform distribution with height. From 04:00 to 05:00 UTC, this trend changes again, with the maximum Nc appearing at the cloud base. Large numbers of small cloud droplets present at the cloud base, the primary area for droplet activation. The peak of Clw appears at higher cloud layers. In contrast, the maximum cloud droplet radius occurs in the middle to upper cloud layers, indicating that the main region of cloud droplet size increasing is near the top and middle-upper parts of cloud regions.

Compared to the control experiment, the increase in aerosol concentration promotes cloud development. This phenomenon is consistent with the findings of Khain et al. (2005) and Morrison et al. (2018). When the accumulation mode aerosol concentration increases, this "promoting" effect becomes most evident. On the other hand, when aerosol concentration decreases, cloud development is suppressed, resulting in a noticeable decrease in cloud-top height.

In terms of cloud microphysical properties, with an increase in aerosol concentration, Nc noticeably increases. As a result, more cloud droplets of small sizes compete for water vapor, reducing cloud droplet size. The maximum Nc and minimum cloud droplet size are observed in the ITM and AM experiments. However, with aerosol concentration decreased, despite cloud development being restrained, the DTM experiment exhibits the largest cloud droplet size.




Figure 8: The variations of averaged cloud liquid water content (Clw, in g/m^3) with time (UTC) and altitude (km) within the study area of different experiments.




Figure 9: The variations of averaged cloud droplet radius (R_m , in μm) with time (UTC) and altitude (km) within the study area of different experiments.




Figure 10: The variations of averaged cloud droplet number concentration (N_c , in cm^{-3}) with time (UTC) and altitude (km) within the study area of different experiments.

3.2.2 Temporal Distribution of Regionally Averaged Cloud Fraction

Figure 11 shows the hourly mean cloud fraction as a function of height. The cloud fraction in the model is a dimensionless ratio, representing the proportion of cloud cover within a grid cell. Throughout the simulation period from 00:00 to 05:00 UTC, the cloud fraction exhibits noticeable variations with both height and time, responding differently to changes in aerosol concentrations across the different modes. In the control experiment (ORG), the cloud fraction peaks between 2 km and 3.5 km, with a maximum value exceeding 0.6, occurring during the later stages of cloud development. This indicates strong cloud development in the mid-cloud layers.

In the NM and AM experiments, increased aerosol concentrations significantly enhance cloud formation, particularly between 03:00 and 05:00 UTC, where the peak cloud fraction above 2 km increases. In the AM experiment, the cloud fraction remains elevated above 3 km, suggesting extended cloud formation and persistence in the upper cloud layers. This result is

批注 [龔李12]: Referee2: 5) In the results section, it is important in my opinion to provide some information about changes in cloud fraction across the different simulations. Cloud fraction is a significant factor in cloud-aerosol interactions and can substantially influence the interpretation of results related to cloud properties and processes. Understanding how cloud fraction varies with different aerosol concentrations is crucial for assessing the broader implications of the study.

consistent with the tendency of accumulation mode aerosols to increase cloud droplet number concentrations and extend cloud lifetime, as reported by Liu et al. (2022). In the ITM experiment, the increase in cloud fraction is most prominent between 03:00 and 05:00 UTC, especially concentrated in the middle cloud layers at 2 to 3 km.

In contrast, the DTM experiment shows a significant reduction in cloud fraction, particularly in the upper cloud layers. The maximum cloud fraction only reaches about 0.4, with most cloud formation concentrated below 3 km. This suppression of cloud formation indicates that lower aerosol concentrations limit cloud development, reducing droplet activation and cloud water content.




Figure 11: The variations of averaged cloud fraction with time (UTC) and altitude (km) within the study area of different experiments.

3.2.3 Cloud Droplet Size Distribution

Figure 12 represents the hourly probability distribution of N_c concerning D . As the cloud system develops, the cloud droplet spectrum widens and exhibits a unimodal distribution. When aerosol concentration increases, the cloud droplet spectrum broadens earlier, and the maximum N_c appears in the AM and ITM experiments. Additionally, the distribution characteristic of the droplet spectrum differs among the experiments. The AM and ITM experiments have their peaks in the 9-15 μm size range, while the NM and CM experiments have their peaks concentrated in the 15-24 μm size range. Meanwhile, with aerosol concentration decreased in the DTM experiment, a tendency of spectrum broadening is observed. However, the spectrum width is smaller than that in the control experiment, and the N_c is lower.

This analysis shows that increased aerosol concentration promotes cloud development and leads to an earlier widening of the cloud droplet spectrum. The increase in accumulation mode aerosols tends to increase the number concentration of small-sized cloud droplets. In contrast, an increase in nucleation and coarse mode aerosols favors the production of large-size cloud droplets. In the NM experiment, although the particle size of nucleation mode aerosols is small, the increase in aerosol concentration still leads to an increase in cloud droplet number concentration because aerosol particle sizes follow a normal distribution in the WRF-SBM scheme. Therefore, aerosol particles with larger sizes within the nucleation mode range can still participate in cloud droplet activation.




Figure 12: The probability distribution of the averaged cloud droplet number concentration (N_c , in cm^{-3}) with respect to the mean diameter (D , in μm), the subfigures represent 01:00, 02:00, 03:00, 04:00, and 05:00 (UTC) on the 25th, respectively. The shading represents the probability magnitude.

3.3 Analysis of Cloud Droplet Spectrum Characteristics

3.3.1 Vertical profiles of cloud droplet spectrum characteristics

To analyse the impact of aerosols on cloud droplet spectrum and cloud microphysical processes, Figure 13-15 given out the variations of hourly averaged ε , cloud-rain auto-conversion intensity (T), and R_v with altitude. The T value represents the probability of auto-conversion occurrence, which can be used to assess the intensity of collision-coalescence processes during cloud and precipitation (Liu et al., 2005, 2006). In the early development stage, the collision-coalescence intensity within the cloud is low. As the cloud system develops, at the vigorous development stage, the T value increases significantly, and the intensity increases with altitude. The intense collision-coalescence processes (with T values > 0.5) are primarily located in the

middle to upper parts of the cloud, consistent with the distribution trend of R_v with altitude. It can be found from Figure 14 that the relative dispersion ε does not change monotonically with R_v or T . The correlation between them will be discussed in the next section.

Compared to the control experiment, the ITM and AM experiments have significantly smaller R_v values, resulting in smaller cloud droplet sizes and lower collision-coalescence intensities than the other experiments. When the aerosol concentration decreases, the R_v in the DTM experiment increases, leading to higher collision - coalescence intensity with respect to other experiments. Additionally, fewer small cloud droplets are activated due to the lower aerosol concentration in the DTM experiment, resulting in lower relative dispersion of cloud droplet spectrum than the other experiments.





Figure13: Distribution of cloud droplet collision-coalescence intensity (T) over time (UTC) and altitude (km). The color shading indicates the collision-coalescence intensity values.



360 **Figure 14: Distribution of cloud droplet volume-weighted mean diameter (R_v , in μm) over time (UTC) and altitude (km). The color shading indicates the magnitude of R_v values.**




Figure 15: Distribution of relative dispersion (ϵ) of cloud droplet spectrum over time (UTC) and altitude (km). The color shading indicates the magnitude of relative dispersion values.

3.3.2 Relationship between ϵ - R_v

Figure 16 reflects the correlation between ϵ and R_v in experiments involving changes in concentration of aerosol modes within the cloud area from 01:00 to 05:00, illustrating the variation of ϵ during the growth of cloud droplet sizes. Fbs indicates the activation intensity corresponding to the fitted correlation in specific droplet size ranges. The ϵ - R_v correlation coefficient table is in the Supplement (Table S1). It is shown that ϵ does not vary monotonically with R_v . There is a significant transition in cloud droplet collision-coalescence intensity around 8 μm radius of cloud droplet. When the R_v is smaller than 8 μm , cloud droplet growth mainly depends on the condensation process. At this stage, there exists a critical radius (R_c) of 4.2 μm . When $R_v < 4.2 \mu\text{m}$, ϵ shows a positive correlation with R_v . While $R_v > 4.2 \mu\text{m}$, ϵ shows a negative correlation with R_v . This trend is close to Lu et al. (2020), but with the value of R_c differs. Among the experiments, increased aerosol concentration enhances

批注 [懿李13]: 8) Fig. 15,16,17: What time in the simulation do these presented results represent? How general they are?

the positive correlation between ε and R_v with $R_v < 4.2 \mu\text{m}$. In the ITM and AM experiments, when R_v is between $4.2\mu\text{m}$ and
375 $8\mu\text{m}$, the negative correlation trend changes to a positive one. In contrast, decreasing aerosol concentration strengthens the negative correlation trend between ε and R_v within the same size range ($4.2 \mu\text{m} < R_v < 8 \mu\text{m}$).

Cloud droplets primarily grow through condensation within the radius range (R_v) of $2\text{-}8 \mu\text{m}$. Figure 17 illustrates the variation of cloud droplet number concentration (N_c) with R_v during the same stage as the ε - R_v correlation, reflecting the concurrent changes in N_c during the growth of cloud droplet sizes. As shown in Figure 17, when R_v is less than $4.2 \mu\text{m}$,
380 accompanied by higher intensity of cloud droplet activation, N_c increases with R_v , and ε shows a positive correlation with R_v . When R_v ranges between 4.2 and $8 \mu\text{m}$, strong collision-coalescence processes have not yet been initiated, and activation intensity is lower. At this stage, N_c does not exhibit significant changes with increasing R_v . Due to the negative correlation between condensation growth efficiency and droplet size, smaller droplets grow rapidly through condensation, whereas larger droplets experience slower growth rate. As R_v increases, ε exhibits a negative correlation with R_v , leading to a more uniform
385 droplet size distribution and a narrower cloud droplet spectrum. This finding aligns with the results of Liu et al. (2006) and Peng et al. (2007). When R_v exceeds $8 \mu\text{m}$, as R_v increases, higher collision-coalescence intensity rapidly depletes smaller droplets (Figure 17), with ε shows a converging trend, ultimately approaching the range of $0.3\text{-}0.4$, consistent with the findings of Lu et al. (2020).

For the sensitivity experiments, an increase in aerosol concentration enhances the activation of cloud droplets, enhancing
390 the positive correlation between ε and R_v when $4.2 < R_v < 8 \mu\text{m}$. Among different aerosol modes, an increase in accumulation mode aerosol contributes to the prolonged maintenance of cloud droplet activation and significantly increases N_c (Figure 17). When $4.2 < R_v < 8 \mu\text{m}$, ε shows a positive correlation with R_v . However, when cloud droplet size increases above $8 \mu\text{m}$, cloud droplet collision-coalescence intensity increases with particle size, while cloud droplet number concentration decreases as R_v increases. Therefore, in this situation, dominant cloud droplet coalescence promotes the rapid growth of cloud droplet size,

395 increasing large-sized cloud droplets while simultaneously consuming small-sized cloud droplets. As a result, ε tends to converge with droplet size.

converge with droplet size.

As it is shown in Figure 17 that the correlation between N_c and cloud microphysical processes is more complex. Regions with the same N_c may be dominated by condensation growth or coalescence processes. Furthermore, the ε - N_c correlation, which is significantly influenced by cloud droplet activation, condensation, and collision-coalescence processes, may exhibit

400 even more complex variations.




Figure 16: The variation of relative dispersion (ε) of cloud droplet spectrum against the cloud droplet volume-weighted radius (R_v , in μm) for different experiments. FBS indicates the cloud droplet activation intensity, and the shading represents the coalescence intensity.





Figure 17: The variation of cloud droplet number concentration (N_c , in cm^{-3}) against the cloud droplet volume-weighted radius (R_v , in μm) for different experiments. The shading represents the coalescence intensity.

3.3.3 Relationship between ε - N_c

Figure 18 shows the relationship between ε and N_c in experiments involving changes in aerosol concentration modes within the cloud area from 01:00 to 05:00. As shown in Figure 18 as N_c increases, ε tends to converge, consistent with the findings of Zhao et al. (2006) and Jin et al. (2021). Additionally, the coalescence intensity does not significantly impact the ε - N_c correlation. With increased coalescence intensity, the dispersion of ε in the low N_c region decreases, but the ε - N_c relationship still shows a converging trend.

批注 [懿李14]: 8) Fig. 15,16,17: What time in the simulation do these presented results represent? How general they are?

415 Compared to the control experiment, changes in aerosol concentration did not affect the ε - N_c correlation. When the aerosol concentration increased, N_c significantly increased, and in the AM and ITM experiments, the dispersion of ε slightly increased in the low coalescence intensity region. On the other hand, a decrease in aerosol concentration led to a significant reduction in N_c and increased cloud droplet size. In the region with $T > 0.8$, the dispersion of ε was higher.



420 **Figure 18: The variation of cloud droplet spectral relative dispersion (ε) against cloud droplet number concentration (N_c , in cm^{-3}).**
The shading represents the coalescence intensity.

4 Discussion

In this study, the increase in coarse-mode aerosol concentration resulted in an increase in R_v and enhanced collision-coalescence intensity, consistent with the findings of Liu et al. (2022). However, unlike Liu et al. (2022), the increase in

425 nucleation-mode aerosol concentration in this study also promoted the early development of cloud tops above 3 km. Compared to the control experiment, both R_v and collision-coalescence intensity at the cloud top region were enhanced. This difference may stem from the classification of aerosol particle sizes; in the WRF-SBM scheme, the distribution of different aerosol modes is assumed to follow a normal distribution. Therefore, for the nucleation mode, some aerosol particles also reach the size scale of the accumulation mode, promoting an increase in N_c and a rise in T values.

430 Moreover, due to the different physical mechanisms involved in the growth of cloud droplet sizes, aerosols of different modes exhibit varying effects. Here, we provide an additional summary based on the schematic mechanism in Figure 19. When cloud droplet sizes are smaller than $8\text{ }\mu\text{m}$, the collision-coalescence process is less active due to the smaller droplet sizes. The increase in accumulation-mode aerosol concentrations has the most pronounced effect on the activation of small cloud droplets, which enhances both the N_c and the condensational growth of droplets within the $4.2\text{--}8\text{ }\mu\text{m}$ range. This results in a shift in the correlation between the ε and the R_v from negative to positive. Aerosols in other modes have a less significant impact on the increase of N_c , which is why they do not alter the ε - R_v relationship. In contrast, a reduction in aerosol concentration weakens cloud droplet activation. As the droplets grow through condensation, their sizes become more uniform, thereby strengthening the negative correlation between ε and R_v when R_v is in the range of $4.2\text{--}8\text{ }\mu\text{m}$.

The relationship between ε and cloud microphysical properties differs from previous studies. In this study, ε shows a convergence trend as N_c increases, and changes in aerosol concentration do not alter this trend but rather affect the degree of dispersion, like the findings of Deng et al. (2009) and Yu et al. (2018). In contrast, study on non-precipitating stratiform clouds in northern China using aircraft observational data (Ma et al., 2010) shown that with an increase in aerosol concentration, ε tended to decrease with increasing N_c , whereas Anil et al. (2016) observed the opposite trend, with ε showing a positive correlation with N_c .

批注 [第15]: 9) Fig. 18: Very informative schematic figure that should be explained more clearly in the text.

445 The complex variations in the ε -Nc relationship are mainly due to the sensitivity of Nc and ε to many microphysical processes, such as updraft strength, aerosol properties, or condensation-coalescence processes (Lu et al., 2012; Peng et al., 2007). During the fitting process of the ε -Nc relationship, it is challenging to determine the corresponding relationship between Nc and cloud microphysical processes. In regions with low Nc, it may correspond to strong collision/coalescence initiation zones, while in regions with high Nc, it may be in the condensation/coalescence dominant zone. As Liu et al. (2008) and Tas
450 et al. (2012) have stated, compared to Nc, Rv considers the synergistic relationship between Nc and water content, providing a more explicit mapping to cloud microphysical processes. Therefore, this study explored the ε -Rv relationship to provide a more systematic understanding of the stratiform warm clouds in Eastern China. The ε -Rv correlation is summarized in Figure 16.




Figure 19: Differences in the effects of aerosol modes on microphysical processes. Cloud Droplet Collision-Coalescence reduces droplet numbers represents the process where cloud droplet size increases while droplet number concentration decreases due to collision-coalescence growth. Cloud Droplet Condensational Growth represents the process of cloud droplets growing by condensation. Cloud Droplet Activation refers to the activation of aerosol particles to form small cloud droplets. Upward red arrows indicate an enhancement of physical quantities or processes, while downward blue arrows indicate suppression..

5 Conclusions

This study used the SBM-FAST bin scheme in the WRF model to simulate a stratiform warm cloud process in Jiangxi, China. Numerical experiments were further conducted to investigate the impact of changes in nucleation mode, accumulation mode, coarse mode, and total aerosol concentrations on the macroscopic and microscopic characteristics of stratiform warm clouds.

The variations in cloud microphysical parameters with aerosol concentrations were analysed, the ε -Rv and ε -Nc relationships
465 were fitted to explore the influence of microphysical processes on ε . Specific conclusions are as follows:

(1) The numerical simulation with bin microphysics scheme reproduces stratiform warm clouds' macro- and microscopic characteristics in Jiangxi, China. Aerosols in the accumulation mode enhance cloud fraction by promoting cloud droplet nucleation and increasing cloud water content, while reduced aerosol concentrations suppress cloud fraction, particularly in the upper cloud layers. As the cloud system develops, Rv and T values gradually increase. Vertically, Rv increases with height,
470 and T also strengthens synchronously with the enlargement of cloud droplet size. The relationship between ε and Rv is not strictly monotonic; as Rv increases, ε initially increases and then decreases. Furthermore, it is found that variations in aerosol concentrations exert a significant influence on cloud development. With an increase in the aerosol concentration of any mode, the cloud droplet spectrum widens earlier. Specifically, higher aerosol concentrations promote cloud growth, increasing cloud-top height. In comparison, lower aerosol concentrations impede cloud droplet activation, decreasing the concentration of cloud
475 droplets and leading to a notable reduction in ε and increased Rv and higher T values.

(2) In contrast, different modes of aerosol concentration variations impact cloud microphysical properties differently. An increase in accumulation mode aerosol tends to increase the concentration of small-size cloud droplets, leading to decreased Rv and a lower collision and coalescence intensity concerning the control experiment. An increase in nucleation mode and coarse mode aerosols favors the production of large cloud droplets. As a result, the increase in accumulation mode aerosol has
480 the most significant impact on Nc enhancement. On the other hand, increases in nucleation mode and coarse mode aerosol concentrations result in an increase in Rv and an enhancement of collision and coalescence intensity.

(3) The variation of ε in the cloud is closely related to cloud microphysical processes. Fitting the ε with Rv and Nc reveals that as Rv increases, the correlation between ε and Rv changes from positive to negative, eventually converging. This transformation is mainly related to cloud droplet activation, condensation, and collision-coalescence processes within the cloud.

485 When T values are less than 0.5, as cloud droplet condensation growth becomes more active and nucleation weakens, the cloud
droplet spectrum relative dispersion transitions from an increasing trend to a decreasing trend with the increase in Rv. With
the enhanced coalescence between cloud droplets, ε primarily decreases with the increase in Rv. Increasing accumulation mode
aerosol concentration contributes to the prolonged cloud droplet activation, causing the correlation between ε and Rv to shift
from negative to positive. On the other hand, a decrease in aerosol concentration leads to a reduction in cloud droplet activation
490 intensity, making the negative correlation trend between ε and Rv more pronounced. In addition, regardless of different T
values, ε converges with the increase in Nc. As Nc increases, ε converges to a range of 0.2-0.4. Changes in aerosol
concentration for different modes do not alter the converging trend of ε with Nc but only affect the dispersion degree of ε at
low Nc values.

Lastly, in this study, due to computational power limitations, the vertical resolution of our simulation setup is relatively
495 coarse. Future research could consider enhancing the resolution to reveal the variations of cloud-aerosol effects more
effectively within the vertical profile of clouds. Moreover, while this study has explored the effects of variations in aerosol
concentrations across different modes on the macroscopic and microscopic characteristics of stratiform warm clouds, mainly
focusing on the influence of these variations on the relationship between ε and cloud microphysical properties, the interaction
between clouds and aerosols is a complex process influenced by multiple factors, including cloud dynamics and supersaturation
500 levels. Therefore, future research should investigate other vital factors affecting cloud-aerosol interactions further. Additionally,
incorporating case studies from diverse regions could effectively reduce the regional dependency of cloud-aerosol effect
research, thereby enhancing our comprehensive understanding of these complex interactions on a global scale.

6 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be
505 construed as a potential conflict of interest.

7 Acknowledgments

In addition, we acknowledge the High Performance Computing Center of Nanjing University of Information Science and
Technology for their support of this work.

8 Funding

510 This work was supported by the National Natural Science Foundation of China (Grant Nos. 42061134009 and 41975176) and
2023 Jiangsu Provincial College Students' Innovative Entrepreneurial Training Program (Grant Nos.202310300108Y).

9 Data Availability Statement

The data used in this study can be accessed at the following link: <https://doi.org/10.57760/sciencedb.11210>. The data link
includes the satellite-observed cloud top temperature data, WRF model simulation results, and simulated initial aerosol
515 spectrum information used in this study.

The cloud top temperature data used in this study is obtained from NASA's Earth Observing System (EOS), specifically
from the Aqua satellite within the MODIS instrument. The data has a horizontal resolution of 5 km × 5 km and is provided
in .nc file format.

The WRF model simulation configurations are described in the previous section. The data format is .netcdf, and details
520 about the data and its dimensions can be found in the data description.

The initial aerosol spectrum data includes the distribution information of aerosol spectra within the first hour of the simulation for one control group and five experimental groups mentioned in the article. The temporal resolution is 10 minutes. Data details can be found in the data description.

In addition, the initial fields used in the numerical simulations are based on the Fifth generation of ECMWF atmospheric reanalysis of the global climate (ERA5) hourly data on pressure levels. These data can be accessed at the following link: <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels?tab=overview>. The study utilized all height variables for every 6 hours from December 24th, 2014, 18:00 to December 25th, 2014, 06:00.

If the manuscript is accepted, the data will be publicly available through the aforementioned link (<https://doi.org/10.57760/sciencedb.11210>). To access the data, you only need to use the database link and provide your name, affiliation, and purpose of the data request to the authors for download.

10 References

Anil Kumar, V. et al.: Investigation of aerosol indirect effects on monsoon clouds using ground-based measurements over a high-altitude site in Western Ghats, *Atmospheric Chemistry and Physics*, 16(13), pp. 8423–8430. doi:10.5194/acp-16-8423-2016, 2016.

Cecchini, M.A. et al.: Sensitivities of amazonian clouds to aerosols and updraft speed, *Atmospheric Chemistry and Physics*, 17(16), pp. 10037–10050. doi:10.5194/acp-17-10037-2017, 2017.

Chandrakar, K.K. et al.: Aerosol indirect effect from turbulence-induced broadening of cloud-droplet size distributions, *Proceedings of the National Academy of Sciences*, 113(50), pp. 14243–14248. doi:10.1073/pnas.1612686113, 2016.

Chandrakar, K.K., Cantrell, W. and Shaw, R.A.: Influence of turbulent fluctuations on cloud droplet size dispersion and aerosol indirect effects, *Journal of the Atmospheric Sciences*, 75(9), pp. 3191–3209. doi:10.1175/jas-d-18-0006.1, 2018.

Chen, J. et al.: New understanding and quantification of the regime dependence of aerosol-cloud interaction for studying aerosol indirect effects, *Geophysical Research Letters*, 43(4), pp. 1780–1787. doi:10.1002/2016gl067683, 2016.

Desai, N. et al.: Search for microphysical signatures of stochastic condensation in marine boundary layer clouds using airborne digital holography, *Journal of Geophysical Research: Atmospheres*, 124(5), pp. 2739–2752. doi:10.1029/2018jd029033, 2019.

Fan J, Leung L R and Li Z, et al. Aerosol impacts on clouds and precipitation in eastern China: Results from bin and bulk microphysics [J]. *Journal of Geophysical Research: Atmospheres*, 117(D16), doi: 10.1029/2011JD016537, 2012.

Flossmann, A.I. and Wobrock, W.: A review of our understanding of the aerosol–cloud interaction from the perspective of a bin resolved cloud scale modelling, *Atmospheric Research*, 97(4), pp. 478–497. doi:10.1016/j.atmosres.2010.05.008, 2010.

Grosvenor, D. P., Sourdeval, O., Zuidema, P., Ackerman, A., Alexandrov, M. D., Bennartz, R., et al.: Remote sensing of droplet number concentration in warm clouds: A review of the current state of knowledge and perspectives. *Reviews of Geophysics*, 56, 409–453. <https://doi.org/10.1029/2017RG000593>, 2018.

Han, B. et al.: Cloud-Resolving Model Intercomparison of an MC3E squall line case: Part II. Stratiform precipitation properties, *Journal of Geophysical Research: Atmospheres*, 124(2), pp. 1090–1117. Available at: <https://doi.org/10.1029/2018jd029596>, 2019.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J.-N.: ERA5 hourly data on pressure levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), DOI: 10.24381/cds.bd0915c6, 2023.

Iltoviz, E. et al.: Effect of aerosols on freezing drops, hail, and precipitation in a midlatitude storm, *Journal of the Atmospheric Sciences*, 73(1), pp. 109–144. doi:10.1175/jas-d-14-0155.1, 2015.

Jensen, J.B. and Nugent, A.D.: Condensational growth of drops formed on giant sea-salt aerosol particles, *Journal of the Atmospheric Sciences*, 74(3), pp. 679–697. doi:10.1175/jas-d-15-0370.1, 2017.

Jin Yuchen, Niu Shengjie and Lü Jingjing, et al.: Study of the Microphysical Structural Characteristics and Cloud–Rain Autoconversion Threshold Function of Stratiform Warm Clouds in Jiangxi [J]. *Chinese Journal of Atmospheric Sciences*, 45(5): 981–993. doi: 10.3878/j.issn.1006-9895.2102.20166, 2021.

Kant, S., Panda, J. and Gautam, R.: A seasonal analysis of aerosol-cloud-radiation interaction over Indian region during 2000–2017, *Atmospheric Environment*, 201, pp. 212–222. doi:10.1016/j.atmosenv.2018.12.044, 2019.

Khain, A. and Sednev, I.: Simulation of precipitation formation in the Eastern Mediterranean Coastal Zone using a spectral microphysics cloud ensemble model, *Atmospheric Research*, 43(1), pp. 77–110. Available at: [https://doi.org/10.1016/s0169-8095\(96\)00005-1](https://doi.org/10.1016/s0169-8095(96)00005-1), 1996.

Khain, A. et al.: Notes on the state-of-the-art numerical modeling of Cloud Microphysics, *Atmospheric Research*, 55(3–4), pp. 159–224. Available at: [https://doi.org/10.1016/s0169-8095\(00\)00064-8](https://doi.org/10.1016/s0169-8095(00)00064-8), 2000.

Khain A P, Rosenfeld D and Pokrovsky A, et al.: Effects of Atmospheric Aerosols on Precipitation From Deep Convective Clouds As Seen From Simulations Using A Spectral Microphysics Cloud Model[J]. Available at: 2002EGSGA.27.6326K, 2002.

Khain, A., Rosenfeld, D. and Pokrovsky, A.: Aerosol impact on the dynamics and microphysics of deep convective clouds. *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 131(611), pp.2639–2663, 2005.

Khain, A. and Lynn, B.: Simulation of a supercell storm in clean and dirty atmosphere using weather research and forecast model with spectral bin Microphysics, *Journal of Geophysical Research*, 114(D19). Available at: <https://doi.org/10.1029/2009jd011827>, 2009.

Kovačević, N.: Hail suppression effectiveness for varying solubility of natural aerosols in water, *Meteorology and Atmospheric Physics*, 131(3), pp. 585–599. doi:10.1007/s00703-018-0587-4, 2018.

585 Kumar, B. et al.: Cloud-edge mixing: Direct numerical simulation and observations in Indian Monsoon clouds, *Journal of Advances in Modeling Earth Systems*, 9(1), pp. 332–353. doi:10.1002/2016ms000731, 2017.

Lau, K.M. and Wu, H.T.: Warm rain processes over tropical oceans and climate implications, *Geophysical Research Letters*, 30(24). Available at: <https://doi.org/10.1029/2003gl018567>, 2003.

Lerach, D.G. and Cotton, W.R.: Simulating southwestern U.S. desert dust influences on supercell thunderstorms, *Atmospheric Research*, 204, pp. 78–93. doi:10.1016/j.atmosres.2017.12.005, 2018.

590 Liu, F. et al.: Opposing comparable large effects of fine aerosols and coarse sea spray on marine warm clouds, *Communications Earth & Environment*, 3(1). doi:10.1038/s43247-022-00562-y, 2022.

Liu, G.: Retrieval of cloud droplet size from visible and microwave radiometric measurements during INDOEX: Implication to aerosols’ indirect radiative effect, *Journal of Geophysical Research*, 108(D1). doi:10.1029/2001jd001395, 2003.

595 Liu, M., Li, L., Xu, L. P., Yu, J. Y., and Liu, P.: Usage and Maintenance of RPG-HATPRO Ground-Based Multi-Channel Microwave Radiometer (in Chinese), *Anal. Instrum.*, 35(5), 89–92, doi:10.3969/j.issn.1001-232x.2014.05.018, 2014.

Liu, Y. and Daum, P.H.: Indirect warming effect from dispersion forcing, *Nature*, 419(6907), pp. 580–581. doi:10.1038/419580a, 2002.

Liu, Y.: Size truncation effect, threshold behavior, and a new type of autoconversion parameterization, *Geophysical Research Letters*, 32(11). doi:10.1029/2005gl022636, 2005.

600 Liu, Y. et al.: Generalized threshold function accounting for effect of relative dispersion on threshold behavior of autoconversion process, *Geophysical Research Letters*, 33(11). doi:10.1029/2005gl025500, 2006.

Liu, Y. et al.: Dispersion bias, dispersion effect, and the aerosol–cloud conundrum, *Environmental Research Letters*, 3(4), p. 045021. doi:10.1088/1748-9326/3/4/045021, 2008.

605 Liu, Y. et al.: Tibetan Plateau driven impact of Taklimakan dust on northern rainfall, *Atmospheric Environment*, 234, p. 117583. doi:10.1016/j.atmosenv.2020.117583, 2020.

Lu, C. et al.: Observed impacts of vertical velocity on cloud microphysics and implications for aerosol indirect effects, *Geophysical Research Letters*, 39(21). doi:10.1029/2012gl053599, 2012.

Lu, C. et al.: Empirical relationship between entrainment rate and microphysics in cumulus clouds’, *Geophysical Research Letters*, 40(10), pp. 2333–2338. doi:10.1002/grl.50445, 2013.

610 Lu, C. et al.: Reconciling contrasting relationships between relative dispersion and volume-mean radius of cloud droplet size distributions, *Journal of Geophysical Research: Atmospheres*, 125(9). doi:10.1029/2019jd031868, 2020.

Lu, C. S., and Xu, X. Q.: Research Progress on Cloud Entrainment-Mixing Processes. *Torrential Rain and Disasters (in China)*, 40(3), 271-279. DOI: 10.3969/j.issn.1004-9045.2021.03.005, 2021.

615 Ma, J. et al.: Strong air pollution causes widespread haze-clouds over China, *Journal of Geophysical Research*, 115(D18). doi:10.1029/2009jd013065, 2010.

Morrison, H. et al.: Broadening of modeled cloud droplet spectra using bin microphysics in an Eulerian spatial domain, *Journal of the Atmospheric Sciences*, 75(11), pp. 4005–4030. doi:10.1175/jas-d-18-0055.1, 2018.

Pandithurai, G. et al.: Aerosol effect on droplet spectral dispersion in warm continental cumuli, *Journal of Geophysical Research: Atmospheres*, 117(D16). doi:10.1029/2011jd016532, 2012.

620 Peng, Y. et al.: An investigation into the aerosol dispersion effect through the activation process in marine stratus clouds, *Journal of Geophysical Research*, 112(D11). doi:10.1029/2006jd007401, 2007.

Pinsky, M. and Khain, A.: Theoretical Analysis of the Entrainment–mixing process at cloud boundaries. part I: Droplet size distributions and humidity within the interface zone, *Journal of the Atmospheric Sciences*, 75(6), pp. 2049–2064. doi:10.1175/jas-d-17-0308.1, 2018.

Platnick, S., Ackerman, S. and King, M., et al.: MODIS Atmosphere L2 Cloud Product (06_L2), NASA MODIS Adaptive Processing System, Goddard Space Flight Center, USA, 2015. doi:10.5067/MODIS/MOD06_L2.006.

Prabha, T.V. et al.: Spectral width of premonsoon and monsoon clouds over Indo-Gangetic Valley, *Journal of Geophysical Research: Atmospheres*, 117(D20). doi:10.1029/2011jd016837, 2012.

Qian Y, Gong D and Fan J, et al. Heavy pollution suppresses light rain in China: Observations and modeling [J]. *Journal of Geophysical Research: Atmospheres*, 114(D7), doi: 10.1029/2008jd011575, 2009.

Rosenfeld, D., Rudich, Y. and Lahav, R.: Desert dust suppressing precipitation: A possible desertification feedback loop, *Proceedings of the National Academy of Sciences*, 98(11), pp. 5975–5980. doi:10.1073/pnas.101122798, 2001.

Rotstajn, L.D. and Liu, Y.: Sensitivity of the first indirect aerosol effect to an increase of cloud droplet spectral dispersion with droplet number concentration, *Journal of Climate*, 16(21), pp. 3476–3481. doi:10.1175/1520-0442(2003)016<3476: sotfia>2.0.co;2, 2003.

Rotstajn, L.D. and Liu, Y.: Cloud droplet spectral dispersion and the indirect aerosol effect: Comparison of two treatments in a GCM, *Geophysical Research Letters*, 36(10). doi:10.1029/2009gl038216, 2009.

Seifert, A., Nuijens, L. and Stevens, B.: Turbulence effects on warm-rain autoconversion in precipitating shallow convection. *Quarterly Journal of the Royal Meteorological Society*, 136(652), pp.1753-1762, 2010.

Shpund, J. et al.: Simulating a mesoscale convective system using WRF with a new spectral bin Microphysics: 1: Hail vs graupel, *Journal of Geophysical Research: Atmospheres*, 124(24), pp. 14072–14101. doi:10.1029/2019jd030576, 2019.

Tas, E., Koren, I. and Altaratz, O.: On the sensitivity of droplet size relative dispersion to warm cumulus cloud evolution, *Geophysical Research Letters*, 39(13). doi:10.1029/2012gl052157, 2012.

645 Tas, E. et al.: The relative dispersion of cloud droplets: Its robustness with respect to key cloud properties, *Atmospheric Chemistry and Physics*, 15(4), pp. 2009–2017. doi:10.5194/acp-15-2009-2015, 2015.

Wang Fei and Lu Chunsong.: Advances of Theoretical, Observational, and Numerical Studies on Relative Dispersion of Cloud Droplet Spectral (in China). *Plateau Meteorology*. DOI: 10.7522/j.issn.1000-0534.00067, 2022.

Wang, F. et al.: An airborne study of the aerosol effect on the dispersion of cloud droplets in a drizzling marine stratocumulus
650 cloud over eastern China, *Atmospheric Research*, 265, p. 105885. doi:10.1016/j.atmosres.2021.105885, 2022.

Wang, X. et al.: A study of shallow cumulus cloud droplet dispersion by large eddy simulations, *Acta Meteorologica Sinica*, 25(2), pp. 166–175. doi:10.1007/s13351-011-0024-9, 2011.

Wang, Y. et al.: An observational study on cloud spectral width in North China, *Atmosphere*, 10(3), p. 109. doi:10.3390/atmos10030109, 2019.

655 Wang, Y. et al.: Diverse dispersion effects and parameterization of relative dispersion in urban fog in eastern China, *Journal of Geophysical Research: Atmospheres*, 128(6). doi:10.1029/2022jd037514, 2023.

Wang Yuan, Jia Hailing and Zhang Ping, et al.: Sensitivity of Cloud Microphysics to Aerosol is Highly Associated with Cloud Water Content: Implications for Indirect Radiative Forcing [J]. *Atmospheric Research*, 309: 107552, 2024. DOI: 10.1016/j.atmosres.2024.107552.

660 Wehbe, Y., Temimi, M. and Adler, R.F.: Enhancing precipitation estimates through the fusion of weather radar, satellite retrievals, and surface parameters, *Remote Sensing*, 12(8), p. 1342. doi:10.3390/rs12081342, 2020.

Xie, X. N., Liu, X. D., and Wang, Z. S.: Research Progress on the Impact of Cloud Droplet Spectrum Dispersion on Aerosol Indirect Effects (in China). *Journal of Earth Environment*, 6(2), 8. DOI: 10.7515/JEE201502008, 2015.

Xie, X. et al.: Sensitivity study of cloud parameterizations with relative dispersion in CAM5.1: Impacts on aerosol indirect
665 effects, *Atmospheric Chemistry and Physics*, 17(9), pp. 5877–5892. doi:10.5194/acp-17-5877-2017, 2017.

Yang, F. et al.: Evaluation of multiple forcing data sets for precipitation and shortwave radiation over major land areas of
China, *Hydrology and Earth System Sciences*, 21(11), pp. 5805–5821. doi:10.5194/hess-21-5805-2017, 2017.

Yang, S. et al.: Effects of aerosol number concentration and updraft velocity on relative dispersion during the collision–
coalescence growth stage of warm clouds, *Atmosphere*, 14(5), p. 828. doi:10.3390/atmos14050828, 2023.

670 Yin, Y. et al.: The effects of giant cloud condensation nuclei on the development of precipitation in convective clouds — a
numerical study, *Atmospheric Research*, 53(1–3), pp. 91–116. doi:10.1016/s0169-8095(99)00046-0, 2000.

Yu Guohang, Yang Suying and Hu Cheng-rong, et al.: Simulation on Impacts of Aerosol Number Concentration on Physical
Characteristics of Warm Clouds during Different Growth Stages [J]. *Journal of Meteorology and Environment*, 38(3):
52-64, 2022.

675 Yum, S.S. and Hudson, J.G.: Adiabatic predictions and observations of cloud droplet spectral broadness, *Atmospheric
Research*, 73(3–4), pp. 203–223. doi:10.1016/j.atmosres.2004.10.006, 2005.

Zhao, C. and Ishizaka, Y.: Modeling marine stratocumulus with a detailed microphysical scheme, *Advances in Atmospheric
Sciences*, 21(1), pp. 61–74. Available at: <https://doi.org/10.1007/bf03342546>, 2004.

Zhao, C. et al.: Aircraft measurements of cloud droplet spectral dispersion and implications for indirect aerosol radiative
680 forcing, *Geophysical Research Letters*, 33(16). doi:10.1029/2006gl026653, 2006.

Zheng, X. et al.: Comparison of macro- and microphysical properties in precipitating and non-precipitating clouds over
Central-eastern China during warm season, *Remote Sensing*, 14(1), p. 152. doi:10.3390/rs14010152, 2021.

Zhu Lei, Lu Chunsong, Gao Sinan and Yum Seong Soo.: Spectral Width of Cloud Droplet Spectra and Its Impact Factors in Marine Stratocumulus[J]. Chinese Journal of Atmospheric Sciences, 44(3): 575-590. doi: 10.3878/j.issn.1006-6859.1905.19115, 2020.