

1 Spring-water temperature suggests widespread occurrence of 2 Alpine permafrost in pseudo-relict rock glaciers

3 Luca Carturan¹, Giulia Zuecco^{1,2}, Angela Andreotti¹, Jacopo Boaga³, Costanza Morino¹, Mirko
4 Pavoni³, Roberto Seppi⁴, Monica Tolotti⁵, Thomas Zanon⁴, Matteo Zumiani⁶

5 ¹Department of Land, Environment, Agriculture and Forestry, University of Padova, Legnaro, Italy

6 ²Department of Chemical Sciences, University of Padova, Padova, Italy

7 ³Department of Geosciences, University of Padova, Padova, Italy

8 ⁴Department of Earth and Environmental Sciences, Pavia, Italy

9 ⁵Fondazione Edmund Mach - Istituto Agrario San Michele All'Adige, S. Michele all'Adige, Italy

10 ⁶Geological Service, Autonomous Province of Trento, Trento, Italy

11 *Correspondence to:* Luca Carturan (luca.carturan@unipd.it)

12 Abstract

13 Runoff originating from ground ice contained in ~~landforms like~~ rock glaciers ~~and talus slopes~~ represents ~~an important~~
14 ~~significant~~ water supply for the lowlands. Pseudo-relict rock glaciers host patchy permafrost, but appear to be visually
15 relict, and therefore can be misinterpreted by using standard classification approaches. Permafrost content, spatial
16 distribution and frequency of this type of rock glaciers are poorly known. Therefore, identifying pseudo-relict rock
17 glaciers that might still host permafrost, and potentially ice, is crucial for understanding their hydrological role in a climate
18 change context.

19 This work analyses rock-glacier spring-water temperature in a 795 km² catchment in the Eastern Italian Alps to understand
20 how many rock glaciers classified as relict could have spring-water temperature comparable to ~~in~~~~taet~~ ~~active or transitional~~
21 rock glaciers, as possible evidence of their pseudo-relict nature. Spring-water temperature, often used as auxiliary to other
22 approaches for specific sites, was used for a preliminary estimate of the permafrost presence in 50 rock glaciers classified
23 as relict. In addition, we present electrical resistivity tomography (ERT) results on two relict rock glaciers with opposing
24 spring-water temperature and surface characteristics to constrain spring-water temperature results at local scale.

25 The results show that about 50% of rock glaciers classified as relict might be pseudo-relict, thus potentially containing
26 permafrost. Both supposedly relict rock glaciers investigated by geophysics contain frozen sediments. The majority of
27 cold springs are mainly associated with rock glaciers with blocky and sparsely vegetated surface, but geophysics suggest
28 that permafrost may also exist in rock glaciers below 2000 m a.s.l., entirely covered by vegetation and with spring-water
29 temperature up to 3.7°C. We estimate that pseudo-relict rock glaciers might contain a significant portion (20%) of all the
30 ice stored in the rock glaciers in the study area. These results highlight the relevance of pseudo-relict rock glaciers in
31 periglacial environments. Even if not a conclusive method, spring-water-temperature analyses can be used to preliminarily
32 distinguish between relict and pseudo-relict rock glaciers in wide regions.

34

35 1. Introduction

36 Timings and magnitude of cryosphere runoff have high climatic sensitivity and are impacted by the current changes of
37 Earth's climate (Engelhardt et al., 2014; Zemp et al., 2015; Carturan et al., 2019). Moreover, a deterioration of the water
38 quality has been reported for springs fed by melting permafrost (Thies et al., 2013; Ilyashuk et al., 2014). Due to glacier
39 decline, in the last decades growing attention has been given to other water reservoirs, such as subsurface ice, including
40 debris-covered glacier ice and, in particular, ground ice stored in periglacial landforms such as rock glaciers and glacial-
41 permafrost composite landforms (e.g., Brighenti et al., 2019; Jones et al., 2019; Schaffer et al., 2019; Seppi et al., 2019;
42 Wagner et al., 2021). Projection of ice loss rates indicates that in the second half of the 21st century more subsurface ice
43 may be preserved than glacier surface ice because of their different response times to atmospheric warming (Haeberli et
44 al., 2017). Subsurface ice is therefore expected to significantly contribute to stream runoff under future climate warming
45 (Janke et al., 2015, 2017).

46 Jones et al. (2018) assessed the importance of ice contained in rock glaciers at global scale, estimating that 62.02 ± 12.40
47 Gt of ice is contained in intact rock glaciers. With the adjective 'intact' the Authors refer to the traditional categorization
48 of rock glaciers, which distinguish between intact rock glaciers (containing ice) and relict rock glaciers (not containing
49 ice). According to the most updated classification (RGIK, 2023), rock glaciers should be categorized into active,
50 transitional and relict, referring exclusively to the efficiency of sediment conveyance (expressed by the surface
51 movement) at the time of observation. This classification should not be used to infer any ground ice content.

52 Even though relict rock glaciers should not contain ice (Haeberli, 1985; Barsch, 1996), more recent studies showed that
53 some relict rock glaciers can preserve permafrost and ice far below the regional lower limit of discontinuous permafrost
54 (e.g., Delaloye, 2004; Strozzi et al., 2004; Lewkowicz et al., 2011; Bollati et al., 2018; Colucci et al., 2019). This evidence
55 raises the question whether a significant fraction of rock glaciers classified as relict is actually to be considered 'pseudo-
56 relict', i.e. *"rock glaciers which appear to be visually relict but still contain patches of permafrost"* (Kellerer-Pirklbauer
57 et al., 2012; Kellerer-Pirklbauer, 2008, 2019). This question is relevant because landforms classified as relict in some
58 regions can be up to an order of magnitude larger and more numerous than ~~intact~~-active and transitional rock glaciers
59 (e.g., Seppi et al., 2012; Scotti et al., 2013; Kofler et al., 2020), with potentially significant ecological and hydrological
60 impacts (e.g., Brenning, 2005a; Millar and Westfall, 2019; Brighenti et al., 2021; Sannino et al., 2021). According to
61 Jones et al. (2019), identifying and establishing the activity state of rock glaciers is an important initial step in determining
62 their potential hydrological significance.

63 Previous investigations on the possible permafrost content of relict rock glaciers looked at -single case studies or small
64 groups of landforms (e.g., Delaloye, 2004; Kellerer-Pirklbauer et al., 2014; Popescu, 2018; Colucci et al., 2019; Pavoni
65 et al., 2023). ~~whereas~~ Studies considering a larger number of relict rock glaciers, at the regional scale, were mainly
66 focussed on the past distribution of mountain permafrost and on the reconstruction of related paleoclimatic conditions
67 (e.g., Frauenfelder et al., 2001; Seppi et al., 2010; Charton et al., 2021; Dlabáčková et al., 2023).

68 As a result, the actual distribution, frequency, and ice content of pseudo-relict rock glaciers might be underestimated,
69 with the latter being essential for implementing worldwide estimates of water resources stored in periglacial landforms
70 (e.g., Jones et al., 2018). Detailed geophysical investigation of selected landforms is certainly suitable as a first step
71 towards a better knowledge of pseudo-relict rock glaciers and their ice content. However, due to logistic constraints, this

72 approach cannot be applied to a large number of rock glaciers at the catchment or regional scale. A recent and
73 commendable advance on this topic has been achieved by the proposition of operational guidelines on the InSAR-based
74 kinematic characterization of rock glaciers (Bertone et al., 2022), which can be used for thorough studies of wide areas.
75 However, this approach is not suitable for distinguishing between relict and pseudo-relict rock glaciers, because their
76 surface ~~is motionless~~has no movement or the movement is very slow and in the same range as the uncertainty of the
77 method.

78 A possible way to investigate the presence of permafrost in these landforms over large areas is by analysing spring-water
79 temperature measured downslope of rock glaciers. Haeberli (1975) proposed the monitoring of spring-water temperature
80 in late summer as useful evidence of permafrost, and various authors employed such method as auxiliary permafrost
81 evidence (e.g., Frauenfelder et al., 1998; Scapozza, 2009; Imhof et al., 2000; Strozzi et al., 2004; Cossart et al., 2008).
82 Carturan et al. (2016) demonstrated that this method can be used successfully for mapping permafrost distribution at the
83 catchment scale. All these works are based on the evidence that, in late summer, spring water affected by permafrost has
84 lower temperature compared to those unaffected, with upper thresholds ranging between 0.9 and 1.1°C for probable
85 permafrost, and between 1.8 and 2.2°C for possible permafrost.

86 In this work, we analyse the spatial variability of spring-water temperature in a 795 km² catchment located in the Eastern
87 Italian Alps, where 338 rock glaciers were inventoried (Seppi et al., 2012), to better understand permafrost distribution.
88 We hypothesise that a significant portion of rock glaciers classified as relict have spring-water temperature comparable
89 to those of ~~inactive~~active and transitional rock glaciers, as possible evidence of their permafrost content and of their pseudo-
90 relict nature. The specific objectives of this study are to:


- 91 i) analyse the influence of topographic and geomorphological factors on spring-water temperature,
- 92 ii) investigate the main controls on water temperature for springs downslope of rock glaciers, and particularly of relict
93 rock glaciers,
- 94 iii) investigate via geophysical analyses the presence of permafrost in two rock glaciers selected for their different spring-
95 water temperature and surface characteristics, to constrain spring-water temperature results at local scale,
- 96 iv) preliminarily estimate and compare the ice content of rock glaciers and glaciers in the study area.-

97

98 2. Study area

99 Val di Sole is located in the upper part of the Noce River catchment, a tributary of the Adige River, which is the main
100 river system in northeastern Italy (Fig. 1). The catchment is 795 km² wide, with elevation ranging between 520 m a.s.l.
101 at the outlet (Mostizzolo) and 3769 m a.s.l. at the summit of Mt. Cevedale, averaging 1705 m a.s.l. (Fig. 1). Metamorphic
102 rocks (mica schists, paragneiss and orthogneiss) prevail in the northern side of the valley, whereas tonalite is found in the
103 southwestern part and dolomites and limestones prevail in the southeastern part (Dal Piaz et al., 2007; Martin et al., 2009;
104 Chiesa et al., 2010, Montrasio et al., 2012).

105



106
107 Figure 1: Geographic location of the study area and of sampled springs. The background is the hillshaded Lidar 2014
108 DEM surveyed by the Provincia Autonoma di Trento (<https://siat.provincia.tn.it/stem/>).

109
110 The catchment includes a glacierised area of 16 km² (in 2006, Salvatore et al., 2015). Bare bedrock and debris are found
111 outside the glaciers down to an elevation of 2700 m, which is the lower regional limit of discontinuous permafrost
112 (Boeckli et al., 2012). A discontinuous cover of alpine meadows and shrubs is present between 2200 m and 2700 m, while
113 below 2000-2200 m forests are dominant. The valley bottom is covered by cultivations and settlements.

114 Val di Sole lies in a transition zone between the “inner dry alpine zone” in the north (Frei and Schär, 1998) and the wetter
115 area under the influence of the Mediterranean Sea in the south. At the valley floor, the mean annual precipitation in the
116 period between 1971 and 2008 averages is ~900 mm. Precipitation increases with elevation and in the southern part, with
117 a maximum of 1500 mm in the Adamello-Presanella Group (Carturan et al., 2012; Isotta et al., 2014). The mean annual
118 0°C isotherm is located at 2500 m. The mean annual air temperature variability is dominated by elevation, whereas
119 latitudinal and longitudinal variations are negligible.

120 Seppi et al. (2012) mapped 338 rock glaciers in Val di Sole. Based on evidence visible in the orthophotos and digital
121 elevation models (DEMs). The largest part of rock glaciers was classified as relict (229, 68% of the total), whereas of
122 the 109 intact rock glaciers whereas only 42 of the remaining 109 can be classified as active based on multitemporal high-

123 resolution ~~digital elevation models (DEMs), and the other 67 can be considered transitional~~. Most rock glaciers (302, 89%
124 of the total) are composed of deposits of metamorphic rocks in the orographic left side of the valley.

125

126 3. Materials and methods

127 3.1 Experimental design

128 We focussed our investigations on the northern part of Val di Sole because it has a rather homogeneous lithology
129 (metamorphic rocks with predominant micaschists) and mean annual precipitation ~~of~~ 1233 mm at 2600 m- (Carturan et
130 al., 2016). This was done to minimise the effects of different lithologies and annual precipitation on the spatial variability
131 of spring-water temperature, and to highlight the role of other variables related to their catchment, upslope area or upslope
132 rock glaciers.

133 To obtain statistically meaningful and generalisable results, we designed a sampling scheme for rock-glacier spring-water
134 temperature considering the variability of permafrost-related characteristics in the study area, namely vegetation cover
135 (related to ground temperature and fine debris infill), size (length, area), elevation, slope, aspect, and lithology (Barsch,
136 1996; Haeberli, 1985; Lambiel and Reynard, 2001; Boekli et al., 2012).

137 We inspected these variables, reported for each rock glacier of Val di Sole in the database of Seppi et al. (2012), using a
138 correlation matrix and the Principal Component Analysis. The aim was to evaluate their possible covariance and to
139 optimise the number of variables ~~and their combinations~~, to be included in the sampling scheme. The analysis revealed
140 high positive covariance between length and area (both related to size). Negative covariance was found between elevation
141 and vegetation cover, and between slope and length/area.

142 Based on these outcomes and considering accessibility of springs, we built a sampling scheme around four variables: i)
143 rock glacier activity, ii) length, iii) mean elevation, and iv) vegetation cover. The last two variables are correlated because
144 ~~intact-active and transitional~~ rock glaciers are at high elevation and almost free from vegetation, and the opposite is true
145 for relict rock glaciers. Vegetation cover is probably one of the few variables that may aid at identifying rock-glacier
146 activity (Ikeda and Matsuoka, 2002; Strozzi et al., 2004, Kofler et al., 2020), and it can vary greatly among rock glaciers
147 at similar elevation. For this reason, we kept both elevation and vegetation, applying a modification to the vegetation-
148 cover classification proposed by Seppi et al. (2012). We distinguish between two classes, namely 'vegetated' and 'non
149 vegetated' for both ~~intact-active/transitional~~ and relict rock glaciers (see Table 1 for threshold values). The vegetation
150 cover was visually estimated in the field and in orthophotos for each rock glacier. Our sampling scheme ensured that at
151 least one rock glacier was sampled for each combination of variables (Table 2). The frequency distribution of rock glacier
152 length and mean elevation was used to identify three terciles, employed for grouping them into short-mid-long rock
153 glaciers and into low-mid-high elevation rock glaciers. Frequency distributions and terciles of ~~intact-active/transitional~~
154 and relict rock glaciers were calculated separately (Table 2).

155

156

157

158

159 Table 1 - Classification of ~~intact~~ active/transitional and relict rock glaciers in two different classes of vegetation cover.

Rock glacier category	Vegetation cover class	Meaning
Intact Active/transitional	Vegetated	Vegetation cover >10%
	Non vegetated	Vegetation cover <10%
Relict	Vegetated	Vegetation cover >50%
	Non vegetated	Vegetation cover <50%

160

161

162

163 Table 2: Sampling scheme used for water temperature measurements at rock glaciers springs.

Activity state	Length	Elevation	Vegetation cover	Number of sampled rock glaciers
Intact Active/transitional	Short (<142 m)	Low (<2634 m)	Non vegetated	2
			Vegetated	none
		Mid (>2634 and <2811 m)	Non vegetated	2
			Vegetated	none
		High (>2811 m)	Non vegetated	1
			Vegetated	none
	Mid (>142 and <251 m)	Low (<2596 m)	Non vegetated	1
			Vegetated	none
		Mid (>2596 and <2817 m)	Non vegetated	1
			Vegetated	3
		High (>2817 m)	Non vegetated	2
			Vegetated	none
	Long (>251 m)	Low (<2655 m)	Non vegetated	none
			Vegetated	1
Mid (>2655 and <2779 m)		Non vegetated	1	
		Vegetated	none	
High (>2779 m)		Non vegetated	3	
		Vegetated	none	
Relict	Short (<180 m)	Low (<2267 m)	Non vegetated	3
			Vegetated	4
		Mid (>2267 and <2453 m)	Non vegetated	1
			Vegetated	2
	High (>2453 m)	Non vegetated	2	
		Vegetated	2	
		Low (<2255 m)	Non vegetated	3

	Mid (>180 and <340 m)		Vegetated	4
		Mid (>2255 and <2425 m)	Non vegetated	1
			Vegetated	2
		High (>2425 m)	Non vegetated	2
			Vegetated	3
		Long (>340 m)	Low (<2222 m)	Non vegetated
	Vegetated			4
	Mid (>2222 and <2388 m)		Non vegetated	3
			Vegetated	5
	High (>2388 m)		Non vegetated	5
			Vegetated	3
				Total:

164

165

166 3.2 Data collection

167 Water temperature was measured at 220 springs, 133 of which are located downslope of rock glaciers ([multiple springs](#)
168 [were often measured downslope of the same rock glacier](#)), 81 are located downslope of other deposits, and 8 are located
169 in bedrock. Springs were sampled from mid-August to mid-October, after the end of the snowmelt. Most springs have
170 been measured once per year from 2018 to 2020, and a small group of them was also measured in 2021. In these four
171 years, the total number of single measurements is 540.

172 Based on the sampling scheme (Table 2), we measured spring-water temperature at 17 [intact-active/transitional](#) rock
173 glaciers and 50 relict rock glaciers, which corresponds to 22% of all rock glaciers existing in the study area. All variables'
174 combinations defined for relict rock glaciers have been sampled, whereas several combinations for [intact](#)
175 [active/transitional](#) rock glaciers lack samplings. This was due to the inexistence of single combinations (e.g., there are no
176 short and vegetated [intact-active/transitional](#) rock glaciers at low elevation) or to the lack of springs and inaccessibility of
177 some rock glaciers.


178 Measurements of spring-water temperature were carried out using a WTW Cond3310 (WTW GmbH, Weilheim,
179 Germany) and a Testo 110 (Testo AG, Lenzkirch, Germany). These instruments have both 0.1°C resolution, but the WTW
180 has higher accuracy ($\pm 0.1^\circ\text{C}$) compared to the Testo ($\pm 0.2^\circ\text{C}$), which was used for back-up/validation. Water temperature
181 measurements were carried out shading the spring from direct sunlight and avoiding probe contact with sediments, rocks,
182 and vegetation. The calibration of the two instruments was checked at the beginning and at the end of the annual
183 campaigns using an ice bath. [In addition, we assessed runoff by a quick visual estimation \(always the same operator\)](#)
184 [similar to Strobl et al. \(2020\), who considered average width, mean depth and velocity of the flow downslope of the](#)
185 [spring. This approach was used to rule out springs with very low runoff \(<0.1 l/s\).](#)~~[In addition, runoff was visually](#)~~
186 ~~[estimated at each spring.](#)~~

187

188 3.3 Data analysis

189 Before proceeding with statistical analyses, we preliminary filtered field data to exclude problematic or redundant
190 measurements. First, we discarded measurements that were clearly affected by very low runoff (<0.1 l/s), responsible of
191 large temperature fluctuations during the day (Seppi, 2006). We then selected one measurement site for each rock glacier
192 and for groups of springs separated less than 10 m from each other. Spring selection was carried out favouring springs
193 with higher-highest runoff, repeated readings in the four years, eloser-closest location to rock glacier fronts, and with
194 lower-lowest interannual temperature variability.

195 After this selection, 131 springs were retained. We characterise the springs using different variables (Table 3), namely
196 the topographic characteristics of the catchments draining to the springs, the activity-state, topographic,
197 geomorphological, and vegetation characteristics of rock glaciers, and the topographic, geomorphological, geological,
198 vegetation and permafrost characteristics of the area immediately upslope of the springs. The latter is defined by the
199 intersection of the catchment perimeter with a circular buffer zone with a radius of 100 m (Fig. 2; Carturan et al., 2016).
200 Details on these variables, the methodology and the data sources (e.g., DEMs, orthophotos, geological maps and literature)
201 employed to derive them are listed and described in Table 3.



203 Figure 2: Delimitation of the spring upslope area, defined by the intersection of a circular buffer zone with a radius of
204 100 m over the catchment perimeter. The methodology was introduced and tested in Carturan et al. (2016).

205
206
207
208
209

Formattato: Tipo di carattere: (Predefinito) Times New Roman, 10 pt

210
211
212
213
214
215
216
217
218

Table 3: Quantitative and qualitative variables used for characterizing spring areas and for statistical analyses.

Spatial scale	Variable type	Variable	Classes/acronym	Meaning
Catchment	Quantitative	Minimum elevation (m a.s.l.) ^a	\	Spring elevation
		Maximum elevation (m a.s.l.) ^a	\	
		Mean elevation (m a.s.l.) ^a	\	Half sum of minimum and maximum elevations
		Planimetric length (m) ^a	\	
	Qualitative	Mean aspect ^a	+NW-NE 2NE-SE SW-NW 3SE-SW	NW-NE (from 315° to 45°) NE-SE (from 45° to 135°) and SW-NW (from 225° to 315°) SE-SW (from 135° to 225°)
Spring upslope area	Qualitative	Geomorphology ^{b,s}	VERver	Slope deposit (scree slope or debris cone)
			GLACglac	Glacial deposit
			RGrgr	Rock glacier
			PRpr	Protalus rampart
			RPrp	Bedrock
			DFdf	Debris flow deposit
			LSls	Solifluction lobe
		Lithology ^b	TTP	Sillimanite paragneiss (Tonale Unit)
			TUG	Granate and cyanite paragneiss (Ultimo Unit)
			TUO	Orthogneiss (Ultimo Unit)
OME	Chlorite e sericite micascists (Peio Unit)			

			OMI	Granate and staurolite micascists (Peio Unit)
			OOG	Orthogneiss (Peio Unit)
			TPN	Metapegmatites (Tonale Unit)
			TTM	Marbles (Tonale Unit)
		Vegetation cover^c	1	0-10% covered by vegetation
			2	10-50% covered by vegetation
			3	50-90% covered by vegetation
			4	90-100% covered by vegetation
		Permafrost evidence^{a,c,h}	weqt	winter equilibrium temperature measured by temperature data loggers
			geophys	geophysical investigations (this work)
			snow	perennial snowfields
			movement	surface displacement visible in multi-temporal DEMs
			none	no evidence available
		APIM^d	Blue	permafrost in nearly all conditions
			Purple	permafrost mostly in cold conditions
			Yellow	permafrost only in very favorable conditions
			White	no permafrost
		Open work deposit^{e,g}	y Open work	present
			a No open work	absent (includes boulder deposits with fine infill and/or widespread vegetation cover)
Rock glacier	Quantitative	Front slope (degrees) ^a	\	
	Qualitative	Activity ^{f,g}	Intact Active/ transitional	Intact Active/transitional rock glacier (active or inactive)
			Relict	Relict rock glacier
		Length ^a	Short	Short rock glacier length class (as defined in Sect. 3)
			Mid	Mean rock glacier length class (as defined in Sect. 3)
			Long	Long rock glacier length class (as defined in Sect. 3)
		Elevation ^a	Low	Low rock glacier elevation class (as defined in Sect. 3)
			Mid	Mean rock glacier elevation class (as defined in Sect. 3)
			High	High rock glacier elevation class (as defined in Sect. 3)

	Vegetation cover^c	Vegetated <u>Vegetation</u>	Vegetated rock glacier (as defined in Sect. 3, Table 1)
		Non vegetated <u>No vegetation</u>	Non vegetated rock glacier (as defined in Sect. 3, Table 1)
	Front characteristics^g	1 <u>I</u>	No vegetation, evidence of recent instability, outcrop of fine material, little or no surface weathering, weathering degree lower than the surface of the rock glacier
		2 <u>II</u>	Very little or no vegetation (<20%), very little or no fine material, weathering and lichen cover comparable to the surface of the rock glacier
		3 <u>III</u>	Scarce or discontinuous and cold-adapted vegetation (≤50%), abundant debris, weathering similar to the surface of the rock glacier, cold air draining from voids among blocks
		4 <u>IV</u>	Completely vegetated, little outcropping debris, without voids and cold air drainage
	Subdued topography^{a-g}	y	The lateral and frontal ridges are clearly evident and the central part of the rock glacier is depressed with respect to them (concave contour lines)
		n	Lateral ridges are absent or evident only in the upper part of the rock glacier, from halfway down the morphology is convex or almost flat

219

220 ^a Derived from the 2006 and 2014 LiDAR DEM of the Trento Province (siat.provincia.tn.it)

221 ^b Derived from the 1:10000 geological map of the Trento Province (protezionecivile.tn.it)

222 ^c Derived from the 2014 orthophoto of the Trento Province (siat.provincia.tn.it)

223 ^d Derived from the Boeckli et al. (2012) Alpine Permafrost Index Map

224 ^e Derived from the hillshaded 2014 LiDAR DEM of the Trento Province (siat.provincia.tn.it)

225 ^f Derived from the Seppi et al. (2012) rock glacier inventory

226 ^g Derived from field observations

227 ^h Ground surface temperature data reported in Carturan et al., (2016) and references therein;

228 <http://www.protezionecivile.tn.it/>.

229

230 We investigated the possible relationship of each variable with the spring-water temperature by means of scatterplots,

231 boxplots, analysis of variance (or Kruskal-Wallis one way analysis of variance on ranks when variances were not

232 homogeneous), Dunn's multiple comparison test, Student's t-test, and regression analysis. We defined spring-water


233 temperature as "the median of all available temperature measurements in the four years", so that we smoothed the

234 interannual variability of water temperature. However, we had also to account for the different number of measurements
235 available for each spring (from one up to four), and in particular for the possible low representativeness of springs
236 measured only once. In this case, there is the possibility of having measured an extreme value, far from the typical
237 conditions of those springs. To evaluate the impact of extreme values, we computed the absolute difference between each
238 single-year spring water measurement and the median of all available measurements at the same spring. The mean of
239 these absolute differences was 0.12°C, the median was 0.05°C, whereas the minimum and the maximum were 0 and
240 0.7°C, respectively, and 89% of values was below 0.3°C. These results indicate a low impact of extreme temperatures
241 and the suitability of using the median of all available measurements (regardless of their number) in statistical analyses.
242 For springs with temperature measured only once, we retained the single value if runoff was >0.1 l/s.
243

244 **3.4 Geophysical investigations**

245 Electrical resistivity tomography (ERT) surveys were performed on 13-14 July 2022 at two neighbouring rock glaciers,
246 classified as relict in the inventory of Seppi et al. (2012), [to constrain spring-water temperature results at local scale](#).
247 These rock glaciers were selected considering their different characteristics (spring-water temperature, vegetation cover,
248 elevation) and the easy access. [In addition, they have uniform lithology, which minimises the uncertainty in the](#)
249 [interpretation of gradients in electrical resistivity](#). The Preghena Rock Glacier has a mean elevation of 2196 m a.s.l., is
250 mainly free of vegetation (although shrubs and trees are present) and its spring-water temperature ranged between 1.6 and
251 1.8°C throughout the late summer during the measuring period. The Bordolona Rock Glacier has a mean elevation of
252 1967 m a.s.l., is completely covered by vegetation and its spring-water temperature ranged between 3.5 and 3.7°C in the
253 late summer during the measuring period. Both rock glaciers are northeast oriented (Fig. 3).

254



255
256 Figure 3: Location of ERT lines (black solid lines) performed on the **a)** Bordolona and **b)** Preghena rock glaciers in July
257 2022. The green dots in the upper panels a) and b) indicate the sampled springs. The white dots **in c) and d)** indicate the
258 lower edge of **Bordolona and Preghena** rock glaciers, **respectively**. The green dots in the upper panels indicate the
259 sampled springs.

260
261 Geophysical surveys were carried out with a Syscal Pro georesistivimeter (Iris Instruments), using arrays of 72 (Line 1
262 Preghena and Line 3 Bordolona) or 48 (Line 2 Preghena) electrodes, with 3-meter electrodes spacing (Fig. 3). A total
263 length of 346 and 216 m was investigated at the Preghena and Bordolona rock glaciers, respectively. A dipole-dipole
264 scheme was used, with two different skips of 0 and 4 electrodes. This configuration ensured relatively high resolution at
265 the surface, and at the same time enough penetration depth. Measurements were carried out with a stack of 3 to 6, imposing
266 an acceptable error threshold of 5%. To estimate a more reliable experimental error for the acquired datasets (Binley,
267 2015), direct and reciprocal measurements were acquired by exchanging injecting and potential dipoles for each

268 quadrupole. To partially overcome the high contact resistances between the electrodes and boulders/debris (Hauck and
269 Kneisell, 2008), the electrodes were inserted between the boulders using sponges soaked with saltwater (Pavoni et al.,
270 2023). Nevertheless, at the blocky surface of the Preghena Rock Glacier the contact resistances remained steadily above
271 $10^{50} \text{ k}\Omega\text{m}$, due to dry environmental conditions. The organic soil at the Bordolona Rock Glacier guaranteed low contact
272 resistances ($<10^4 \text{ k}\Omega\text{m}$).

273 The inversion process of the acquired datasets has been performed with the Python-based software ResIPy (Blanchy et
274 al., 2020), based on the Occam's inversion method (Binley and Kemna, 2005). In each dataset, quadrupoles with a
275 stacking error higher than 5% were removed, and the expected data error was defined using the reciprocal check (Day-
276 Lewis et al., 2008, Pavoni et al., 2023), giving values of 20% and 5% for the Preghena and Bordolona Rock Glacier,
277 respectively. ~~The acquired data were of lower quality at the Preghena Rock Glacier, due to the high contact resistance.~~

278

279 **3.5 Calculation of ice storage in the rock glaciers and glaciers**

280 In order to estimate and compare the ice content of rock glaciers and glaciers in Val di Sole, we applied an approach
281 similar to the one used by Bolch and Marchenko (2009) in the Northern Tien Shan. For glaciers, we estimated residual
282 volumes in 2022 starting from the 2003 ice thickness estimates provided for each glacier in the study area by Farinotti et
283 al., (2019). We first calculated the bedrock topography subtracting the ice thickness from the glacier surface DEM
284 (Farinotti et al., 2019). Then we calculated the 2022 glacier thickness subtracting the bedrock topography from a glacier
285 surface DEM surveyed in September 2022 by the Province of Trento. We finally obtained the glacier volumes multiplying
286 the average thickness by the glacier area, and converted the ice volume into the water volume equivalent using a mean
287 ice density of 900 kg m^{-3} .

288 For rock glaciers, we calculated the total rock glacier volume multiplying their area A by the average thickness provided
289 by the Brenning (2005b) formulation:

$$290 T = cA^\gamma \quad (1)$$

291 where T is the average thickness of rock glaciers, and c and γ are constants equal to 50 and 0.2, respectively. To account
292 for the different geometry of active/transitional and relict rock glaciers, we assumed that the volumetric ice content of
293 active/transitional rock glaciers averages 50% (Jones et al., 2018, and references therein), and therefore that T_r for (true)
294 relict rock glaciers is half that of active/transitional rock glaciers (i.e., they are composed only of debris and all the ice
295 melted away). For pseudo-relict rock glaciers we tested various hypotheses of percent ice content, ranging between 5%
296 and 20%, calculating the average thickness T_{pr} as follows:

$$297 T_{pr} = T_r + T_{ice} \quad (2)$$

298 where T_{ice} is the average ice thickness, calculated in function of the volumetric percent ice content $\%_{ice}$ as:

$$299 T_{ice} = \frac{\%_{ice} T_r}{(1 - \%_{ice})} \quad (3)$$

300

301 **4. Results**

Formattato: Apice

Formattato: Apice

302 **4.1 Spatial variability of spring-water temperature**

303 Water temperature of the 131 springs ranged between 0.0 and 8.5 °C, with a mean of 3.6 °C and a median of 3.4°C (Table
 304 4). The frequency distribution of the spring elevation (i.e., the minimum elevation of catchments) is symmetrical and
 305 normally distributed around a sample mean of 2384 m a.s.l. The lowermost spring was sampled at 1698 m a.s.l., and the
 306 uppermost spring was sampled at 3039 m a.s.l.

307 The mean elevation of spring catchments varies between 2104 and 3151 m a.s.l., whereas the maximum elevation ranges
 308 between 2241 and 3352 m a.s.l. The mean and maximum elevation average 2539 and 2694 m a.s.l., respectively. Both
 309 are also symmetrical around the sample mean and normally distributed.

310 The planimetric length of spring catchments varies between 83 and 2621 m, with a mean of 610 m. The skewness and
 311 kurtosis indicate that the planimetric length is right skewed and leptokurtic.

312

313

314

315

316

317


318 Table 4: Descriptive statistics for spring-water temperature measurements and quantitative variables relative to spring
 319 catchments (as defined in Table 3).

N = 131	Median temperature (T_{Med})	Catchment minimum elevation (m)	Catchment maximum elevation (m)	Catchment mean elevation (m)	Catchment planimetric length (m)
Minimum	0.0	1698	2241	2104	83
Median	3.4	2367	2641	2495	539
Maximum	8.5	3039	3352	3151	2621
Range	8.5	1341	1111	1047	2538
Mean	3.6	2384	2694	2539	610
Standard error of the mean	0.2	22.6	21.9	21.0	34.9
Standard deviation	1.8	259.2	251.1	240.8	399.3
Coefficient of variation	0.500	0.109	0.093	0.095	0.655
Skewness	0.392	0.179	0.446	0.419	2.070
Kurtosis	-0.261	-0.328	-0.107	-0.391	6.095

320

321


322 Spring-water temperature is significantly correlated with the mean elevation of the catchments (Fig. 4a) for all three
 323 aspect classes defined in Table 3. Linear regressions are significant ($p < 0.001$) for south ($R^2 = 0.30$) and for east-west
 324 facing catchments ($R^2 = 0.35$). For the north facing catchments, there is a low significant relation ($R^2 = 0.25$, $p < 0.05$)
 325 between water temperature and elevation. In all three cases, the low R^2 suggests that other factors should affect water
 326 temperature, as well. Similar results were obtained using spring elevation rather than mean catchment elevation (Fig. 5).



327

328 Figure 4: Relationship between spring-water temperature and a) mean catchment elevation (clustered in three classes of
 329 mean catchment aspect), b) mean catchment aspect, c) mean catchment elevation (clustered in seven classes of upslope
 330 area geomorphology), and d) upslope area geomorphology. Acronyms and their meanings are reported in Table 3. Boxes
 331 in b) and d) indicate the 25th and 75th percentile, whiskers indicate the 10th and 90th percentile, whereas the horizontal
 332 solid and dashed lines within the box mark the median and the mean, respectively. Different letters above the boxplots
 333 indicate groups with significantly different ($p < 0.05$) water temperatures based on Dunn's multiple comparison test
 334 (applied after the Kruskal-Wallis test).

335
336
337
338



339

340 Figure 5: Relationship between spring-water temperature and minimum (spring) elevation, clustered in three classes of
341 mean catchment aspect, as in Fig. 4a.

342 As expected, there is a negative relationship between water temperature and elevation (Fig. 4a and 5), but also a large
343 overlap of water temperature among the three aspect classes. NW-NE facing catchments have significantly colder springs
344 compared to SE-SW facing catchments ($p < 0.05$, Dunn's multiple comparison test, applied after the Kruskal-Wallis test),
345 whereas catchments facing NE-SE and SW-NW have water temperature that do not differ significantly from the other
346 two classes (Fig. 4b). NW-NE facing catchments show a lower variability in spring-water temperature compared to the
347 other two classes.

348 Figures 4c and 4d highlight that springs with upslope areas dominated by the presence of rock glaciers (irrespective of
349 their activity) and bedrock outcrops are significantly colder than other springs ($p < 0.05$, Dunn's multiple comparison test,
350 applied after the Kruskal-Wallis test).


351

352 4.2 Temperature of springs downslope of rock glaciers

353 4.2.1 Comparison between ~~inactive~~-active/transitional and relict rock glaciers

354 The spring-water temperature is significantly different for rock glaciers with different degrees of activity (Fig. 6a). Relict
355 rock glaciers have a much warmer spring temperature compared to ~~inactive~~-active/transitional rock glaciers (Student's t-
356 test, $p < 0.001$), and the variability of water temperature is larger for relict rock glaciers. There is a substantial overlap

357 between the two groups, which extended between 1.2 and 3.0°C. This range of water temperature represents 54% of all
 358 springs downslope of rock glaciers (53% of ~~inactive~~ active/transitional rock glaciers and 54% of relict rock glaciers). Almost
 359 half of rock glaciers classified as relict has spring-water temperature similar to rock glaciers classified as
 360 ~~inactive~~ active/transitional.
 361 The two groups of rock glaciers have significantly different minimum elevations (Fig. 6b, Student's t-test, $p < 0.001$), but
 362 there is a wide elevation band, comprised between 2406 and 2630 m a.s.l., where they overlap.



363 **Figure 6:** Spring-water temperature (a) and minimum elevation (b) of rock glaciers sampled in the study area. Boxes
 364 indicate the 25th and 75th percentile, whiskers indicate the 10th and 90th percentile, whereas the horizontal solid and dashed
 365 lines within the box mark the median and the mean, respectively. Maximum and minimum values are represented by dots.
 366 Sample size (n) is reported in a).

368
 369 **4.2.2 Spring-water temperature of relict rock glaciers**

370 The relationship between water temperature and the mean catchment elevation is rather weak for springs fed by relict
 371 rock glaciers (Fig. 7a). The linear regression is significant ($p < 0.05$) only for catchments facing NE-SE and SW-NW, but
 372 the relation is weak ($R^2 = 0.20$). At the same elevation, catchments facing NW-NE have colder springs compared to the
 373 other two aspect classes. The spring-water temperature of catchments facing north is similar to that of catchments facing
 374 east, south and west located 300-400 m above.

375 Relict rock glacier springs with open work deposits in their upslope areas are colder than springs without open work
 376 deposits (Fig. 7b). For the first group, the water temperature is not related to the mean catchment elevation, whereas for
 377 the second group there is a weak but significant relation ($p < 0.05$, $R^2 = 0.15$). Consequently, the difference in water
 378 temperature of the two groups increase towards low elevations, which suggest that open work deposits may have a cooling
 379 effect particularly marked at elevations < 2500 m a.s.l.


380 Similar considerations can be done for rock glacier front characteristics (Fig. 7c) and for rock glacier vegetation cover
 381 (Fig. 7d). Relict rock glaciers with scarce and cold-adapted vegetation cover have colder springs compared to relict rock
 382 glaciers with abundant vegetation cover on their bodies and fronts. However, for all classes of rock-glacier front

Formattato: Tipo di carattere: (Predefinito) Times New Roman, 10 pt

383 characteristics and vegetation cover (Table 3) there is no significant relation between water temperature and mean
384 catchment elevation.

385 Despite the large overlap among the analysed classes (Fig. 7), we found a significant effect of vegetation cover (Student's
386 t-test, $p < 0.001$), open work deposits (Student's t-test, $p < 0.001$) and front characteristics (Student's t-test applied to classes
387 III and IV, $p < 0.01$) on the water temperature of springs downslope of relict rock glaciers. We did not detect any significant
388 influence of the mean aspect of the catchment, the mean elevation of rock glaciers, their length, and the presence or
389 absence of a subdued topography on water temperature.

390



391

392 Figure 7: Relationship between spring-water temperature of relict rock glaciers and mean catchment elevation clustered
393 in a) three classes of mean catchment aspect, b) two classes of open work deposits in the spring upslope area, c) three
394 classes of rock glacier front characteristics, and d) two classes of rock glacier vegetation cover. Classes are described in
395 Table 3.

396

397 4.3 Geophysical investigations

398 Figures 8a and b show the inverted resistivity sections obtained for the investigation Lines 1 and 2 acquired on the
399 Preghena Rock Glacier. High values of resistivity ($>8 \cdot 10^4 \text{ k}\Omega\text{m}$) were found in the uppermost layer, down to about 7-8
400 meters of depth, associated to the dry conditions during ERT soundings and to the air-filled voids among coarse debris
401 and blocks, typical of rock glacier environments. Below this uppermost layer, the resistivity values rapidly decrease ($<10^4$
402 $\text{k}\Omega\text{m}$) indicating a plausible decrease of porosity and grain size in the deposit, and a possible increase in water content.
403 This low resistivity layer develops almost continuously down to the bottom of the models. An increase in resistivity is
404 found at the lower end of line 1 and at the upper end of line 2, in the area where they overlap and at a depth of about 12-
405 13 m, reaching $1.50-2.0 \cdot 10^3 \text{ k}\Omega\text{m}$. This area of increased resistivity can be interpreted as a deep frozen body, providing
406 evidence of probable permafrost inside this rock glacier.

Formattato: Apice

Formattato: Apice

Formattato: Apice

407 Figure 8c shows the inverted resistivity section obtained for the investigation Line 3 acquired on the Bordolona Rock
408 Glacier. In the shallowest layers the resistivity is comprised between $5 \cdot 10^3$ and $10^4 \text{ k}\Omega\text{m}$, significantly lower than the
409 shallow layer of the Preghena Rock Glacier, even if air-filled voids are common on this rock glacier as well.

Formattato: Apice


Formattato: Apice

410 Below this layer, a sharp increase in resistivity is detected along the entire investigation line, with frequent regions
411 exceeding $2 \cdot 10^4 \text{ k}\Omega\text{m}$. The highest resistivity (about $6 \cdot 10^4 \text{ k}\Omega\text{m}$) is found towards the upper end of the ERT line, where
412 a younger rock glacier lobe overlies the main body. This high resistivity layer reaches about 15 meters of depth and can
413 be interpreted as a frozen layer. The bottom of the high-resistivity layer, which seems discontinuous in the lower part and
414 more continuous and thicker in the upper part of the ERT line, is highlighted by a strong decrease in resistivity, below
415 $5 \cdot 10^3 \text{ k}\Omega\text{m}$. This lowermost layer is probably unfrozen and is characterised by an increase in water content and fine
416 sediments.

Formattato: Apice

Formattato: Apice

Formattato: Apice



417

418 Figure 8: Inverted resistivity section of the investigation Line 1 (a) and 2 (b) on the Preghena Rock Glacier, and of the
419 investigation line 3 (c) on the Bordolona Rock Glacier.

420

421 4.4 Ice storage in the rock glaciers and glaciers

422 A total glacier ice volume of $251 \times 10^6 \text{ m}^3$, and a corresponding $226 \times 10^6 \text{ m}^3$ water volume equivalent was calculated for
423 Val di Sole in 2022. In comparison, the water volume equivalent of ~~inactive~~active/transitional rock glaciers is 42.7×10^6
424 m^3 .
425 A water volume equivalent between ~~3.74.4~~ and ~~17.720.9~~ $\times 10^6 \text{ m}^3$, averaging ~~10~~ $12.7 \times 10^6 \text{ m}^3$, can be estimated
426 assuming that ~~38~~50% of the total area of relict rock glaciers contains permafrost (rounded value, based on results
427 reported in Section 4.2.1), and that the average ice content ranges between 5% and 20% in volume.

428

429 5. Discussion

430 5.1 Permafrost distribution and spring-water temperature in the study area

431 Measurements of spring-water temperature collected in this study outside the rock-glacier influence have a high spatial
432 variability and do not show a significant relationship with elevation ($p > 0.05$). Among springs outside the rock glacier
433 influence, only those above 2800 m a.s.l. have a water temperature $\leq 2.2^\circ\text{C}$, which is the upper limit reported in the
434 literature for 'possible permafrost' (Carturan et al., 2016).

435 This result lines up well with mean annual air temperature (MAAT) indications. Indeed, based on the MAAT of -0.9°C
436 measured between 1961 and 2010 at the Careser Diga weather station (2605 m a.s.l., in the northern part of the Val di
437 Sole), the theoretical lower limit of discontinuous permafrost in Val di Sole, corresponding to a MAAT of -2°C (Haeberli,
438 1985), should be comprised between 2700 and 2800 m a.s.l..

439 Similarly, the alpine permafrost index map (APIM, Boeckli et al., 2012) indicates a lower limit of "permafrost mostly in
440 cold conditions" ranging between 2500 and 2900 m outside rock glaciers and coarse-block deposits, varying upon terrain
441 aspect and averaging 2700 m a.s.l.. Based on the mean elevation of ~~inactive~~active/transitional rock glaciers in the study
442 area, Seppi et al. (2012) calculated a present-day lower limit of permafrost at 2720 m a.s.l..

443 As expected, springs draining north-facing catchments are significantly colder compared to springs draining south-facing
444 catchments. On average, there is a difference of about 3°C between springs draining catchments at similar elevation and
445 with opposite aspect. On average, the same spring temperature is found 500-600 m higher on south-facing catchments
446 than on north-facing ones (Fig. 5). This result quantifies the influence of terrain exposure on the ground temperature
447 regime and permafrost distribution in the study area, which are direct consequences of shortwave radiation inputs and
448 related effects on snow cover and surface albedo (Boeckli et al., 2012; Wagner et al., 2019; Amschwand et al., 2024).

449 In our study, at all elevations, springs draining rock glaciers are the coldest, irrespective of the rock glacier activity state
450 (Fig. 4c). This is in agreement with findings of studies in the European Alps and in other mountain chains reporting rock-
451 glacier spring-water temperatures, regardless of their activity state. For example, in the Canadian Rockies, spring-water
452 temperature from an inactive rock glacier hosting small portions of permafrost reached a maximum of 2.2°C , exercising
453 a substantial cooling effect on the creek downstream (Harrington et al., 2018). Interestingly, cold conditions and high
454 daily variability in spring-water temperature in summertime has been recorded in a rock glacier in Norway that shows
455 characteristics favourable to the presence of permafrost, but with minor ice bodies (Lilleøren et al., 2022). In the Austrian
456 Alps, spring-water from a relict rock glacier was monitored for 6 years, showing a mean temperature of 2.2°C , with small
457 seasonal variation (between 1.9 and 2.5°C) and a decrease of the water temperature after precipitation events, attributed
458 to the potential presence of ice lenses in the lower part of the rock glacier (Winkler et al., 2016).


459 Our results align as well with those of studies reconstructing permafrost distribution by empirical modelling in the Alps
460 and at other mountain locations worldwide. A logistic regression model used in the Dry Andes of Argentina accounting
461 for mean annual air temperature, terrain ruggedness, and potential incoming solar radiation suggests that permafrost may
462 occur in several types of coarse blocky deposits, including rock glaciers, even under unfavourable climatic conditions
463 (Tapia Baldis and Trombotto-Liaudat, 2020). A similar empirical-statistical model applied in the Austrian Alps shows
464 that permafrost can be expected above 2500 m a.s.l. in northerly exposed slopes and above 3000 m a.s.l. in southerly
465 exposed slopes (Schrott et al., 2012), providing an elevation difference of about 500 m between south and north exposures,
466 which agrees well with our spring-water temperature results.

467 5.2 Rock glacier classification based on spring-water temperature

468 Although springs draining ~~intact-active/transitional~~ rock glaciers are significantly colder than springs draining relict rock
469 glaciers, there is a remarkable ~50% overlap in the water temperature range of the two rock glacier groups (Fig. 6a).
470 Based on published thresholds (Haerberli, 1975; Frauenfelder et al., 1998; Scapozza, 2009; Carturan et al., 2016), 12 out
471 of the 52 relict rock glaciers sampled in Val di Sole (23%) can be included in the ‘possible permafrost’ category (water
472 temperature between 1 ± 0.2 and 2 ± 0.2 °C), and none of them in the ‘probable permafrost’ category (water temperature <
473 1 ± 0.2 °C). However, the relatively warm water temperature measured downstream of ~~intact-active/transitional~~ rock
474 glaciers (maximum = 3°C, 90th percentile = 2.4°C), and downstream of areas with permafrost evidence (maximum =
475 3.5°C, 90th percentile = 2.2°C), suggest that the upper limit of spring-water temperature for possible permafrost may be
476 higher. Here, the 90th percentile accounts for possible misclassification of ~~intact-active/transitional~~ rock glaciers and other
477 issues affecting spring-water temperature measurements (Sect. 5.3).

478 Assuming a (rounded) upper limit of 2.5°C for spring-water temperature with possible permafrost influence leads to
479 include 19 (38%) relict rock glaciers in the possible permafrost category. This estimate looks more conservative than the
480 ~50% obtained by a mere comparison of water temperature ranges of ~~intact-active/transitional~~ and relict rock glaciers
481 (Fig. 6a). These findings might suggest that permafrost in rock glaciers classified as relict is widespread in Val di Sole,
482 and that a large fraction of them is actually pseudo-relict, or transitional landforms, containing patches of permafrost and
483 reaching an elevation below the tree line (2000-2200 m a.s.l.). Compared to the rock glacier classification performed by
484 Seppi et al. (2012), which was based on remote-sensing geomorphometric evidence combined to field observations
485 (topographic surveys and ground surface temperature measurements for few rock glaciers), spring-water temperature
486 suggests the need for a reclassification of a large fraction of rock glaciers categorised as relict into pseudo-relict.

487 Examples of spring-water temperature downstream of rock glaciers in Val di Sole are shown in Fig. 9. Cold springs
488 draining rock glaciers classified as relict are associated to the presence of open work deposits and scarce vegetation cover
489 (Fig. 7 and 9). These two explaining variables are often correlated, because vegetation tend to be scarce over coarse
490 deposits without fine infill among blocks, and vice versa. The relationship between cold spring temperature (as permafrost
491 evidence) and these two surface characteristics was expected in our case study, based on the existing literature (e.g.,
492 Guglielmin, 1997, and references therein). This relationship is statistically significant only for rock glaciers classified as
493 relict, whereas for ~~intact-active/transitional~~ rock glaciers sampled in the study area it does not exist (Fig. A1).



494
495 Figure 9: Examples of spring-water temperature downstream of rock glaciers in Val di Sole: a) ~~intact~~(active) rock glacier
496 with cold spring at 2950 m a.s.l.; b) ~~intact~~(inactive)transitional rock glacier with relatively warm spring at 2727 m a.s.l.;
497 c) relict rock glacier with cold spring at 2304 m a.s.l., whose surface is open-work and presents scarce vegetation cover;
498 d) relict rock glacier with warm spring at 2266 m a.s.l., whose surface is entirely covered by vegetation.

499 The long-term preservation of permafrost within open work blocky deposits results from overcooling and thermal
500 decoupling of the frozen core from the external climate (Harris and Pedersen, 1998; Morard et al., 2008; Jones et al.,
501 2019). The low thermal conductivity in coarse open work deposits brings to lower ground temperatures compared to fine-
502 grain material (Juliussen and Humlum, 2008; Jones et al., 2019). Soil development over the surficial blocks and boulders
503 can prevent these cooling effects (Ikeda and Matsuoka, 2002). However, if fine-grain infilling does not occur, ground
504 cooling effect goes undisturbed. In central Europe, these processes enable the existence of permafrost much below its
505 regional limit and reaching elevations lower than 1000 m a.s.l. (Gude et al., 2003; Delaloye et al., 2003). According to
506 Delaloye and Lambiel (2005), thousand-year-old permafrost might be potentially preserved in these types of deposits.

507 Open work deposits and/or scarce vegetation cover can be potentially employed to distinguish rock glaciers with or
508 without permafrost, as both can be mapped based on remote-sensing imagery. However, open work deposits and
509 vegetation cover do not enable a full distinction of 'cold' and 'warm' springs affected by relict rock glaciers (Fig. 7b, c
510 and d). Individual non-open-work rock glaciers widely covered by vegetation can have spring-water temperature as low
511 as 1.4°, and rock glaciers almost free of vegetation with blocky surface can have spring-water temperature up to 3.9°C.

512 Other variables considered in this study, such as aspect, elevation, size and the presence or absence of a subdued
513 topography on rock glaciers (Delaloye et al., 2003; Delaloye, 2004), are not related to spring-water temperature. Figure
514 7 suggests the existence of a group of cold springs at low elevations on north-facing catchments, even though water

515 temperature is not significantly different from the temperatures of springs in the other two aspect classes. This result
516 might be due to the small sample size of the NW-NE aspect class.

517

518 **5.3 Limitations and uncertainties in the spring-water temperature approach**

519 The results of this study might be affected by limitations in the experimental design, assumptions, and uncertainties. First,
520 the main assumption of this study is that spring-water temperature provides indication of permafrost occurrence at
521 investigated rock glaciers and spring upslope areas, and can be used as a stand-alone pilot method to rapidly explore the
522 activity state of rock glaciers in a wide area. This approach applies spring-water temperature to the catchment scale,
523 beyond its general use as an ancillary method to other techniques such as InSAR analyses, ground surface temperature
524 measurements and/or geophysics.

525 We base our assumption on previously published work and well-known temperature thresholds for permafrost probability
526 categories (e.g., Haeberli, 1975; Frauenfelder et al., 1998; Scapozza, 2009) and on our first successful application at the
527 catchment scale (Carturan et al., 2016). Data collected in Val di Sole are in line with literature thresholds, provided that
528 the 10% largest spring-water temperature values are excluded (Sect. 5.2). Including these extreme values leads to about
529 1.5°C larger temperature thresholds for possible permafrost compared to literature.

530 The reason behind this discrepancy lies in the uncertainty in the classification of rock glacier activity, which was based
531 on vegetation and geomorphological characteristics, assessed mainly from remote-sensing images (Seppe et al., 2012). In
532 the wide elevation band where ~~intact-active/transitional~~ and relict rock glaciers coexist (minimum elevation between 2406
533 and 2630 m), landforms with similar vegetation cover and surface geomorphology have been classified based on the
534 authors' experience and judgement, implying a certain degree of subjectivity.

535 The distinction between ~~intact-active/transitional~~ and relict rock glaciers is a theoretical concept, and there is a continuum
536 between ~~(true) intact-transitional~~ and (true) relict rock glaciers, ~~with the existence of transitional landforms~~ (Kääb, 2013).
537 In absence of other evidence, this continuum hampers to distinguish unambiguously ~~intact-transitional~~ and relict
538 landforms, in particular if they have similar surface characteristics. In addition, the mentioned transition is a dynamic
539 concept, which depends on the characteristics of individual landforms, their topo-climatic setting, and their response to
540 climatic variations (Kääb, 2013).

541 Another source of uncertainty is related to the distance between the permafrost body and the measured springs. Water
542 temperature is a non-conservative tracer, and if the main permafrost body is distant (e.g., ~~ten~~more than one hundred of
543 meters) from the rock glacier front, ~~or if permafrost is patchy and not in contact with groundwater paths~~, water temperature
544 can ~~be largely influenced by~~significantly increase along the flow paths before reaching the spring, due to the contact with
545 unfrozen sediments and/or mixing with other water sources (e.g., Kellerer-Pirklbauer et al., 2017). This is the case of the
546 Bordolona Rock Glacier (Fig. 8c), where the rather warm spring-water temperature (3.5-3.7°C) would have led to exclude
547 the occurrence of permafrost in absence of geophysical evidence.

548 For smaller distances, we have checked the impact of spring location downstream of rock glacier fronts at three
549 measurement sites, where the same stream emerged briefly at the rock glacier front and a few tens meters downstream.
550 Measurements confirmed that there was negligible warming (from 0.0 to 0.1°C) of the water downstream of the rock
551 glacier front, at least as long as the water remained below the surface.

552 Seasonal ice formed in the topmost ground layer during winter and spring, in areas without permafrost, might cool down
553 spring-water temperature leading to false-positives in permafrost detection. We think that making measurements in late

554 [summer, as proposed by the literature \(e.g. Haeberli, 1975\), prevents this seasonal ice from affecting spring-water](#)
555 [temperature measurements, or at least strongly minimizes its effect. Possible influence from seasonal ground ice formation](#)
556 [should be largest after cool/short summer seasons, but this was not the case in the years analysed.](#)
557 [Depending on the measurement time, which was between 8.00 and 18.00 CET, any variation of temperature during the](#)
558 [day might also influence the results. Hourly records of spring-water temperature collected by Seppi \(2006\) lead us to](#)
559 [exclude significant variation of spring-water temperature during the day, at least for springs with runoff higher than 0.1](#)
560 [l/s.](#)

561 Several authors are cautious when discussing about cold springs downslope of relict rock glaciers. For example, Winkler
562 et al. (2016) do not exclude the presence of remaining ice lenses inside the relict Schöneben Rock Glacier (Niedere Tauern
563 Range, Austria), as a possible explanation for the rapid cooling of the spring water after recharge events, during
564 summertime. However, the authors mention the cold thermal regime beneath coarse blocky materials as a possible
565 explanation, which does not necessarily imply permafrost occurrence, and conclude that additional research is required
566 for the identification of the cooling source.

567 We agree that additional research is required to confirm inference from spring-water temperature. With this study we add
568 that spring-water temperature can be as high as 1.8°C for rock glaciers where permafrost occurrence is confirmed by
569 geophysics or ground surface temperature measurements, and can exceed 3.5°C where the permafrost body is far from
570 the rock glacier front and spring, such as at the Bordolona Rock Glacier. Even if the collected data seem to suggest that
571 temperature thresholds might be slightly higher than those reported in the literature, further investigations are necessary
572 for better constraining them and for defining their range of uncertainty.

573 A last source of uncertainty is represented by the sampling design adopted for Val di Sole, with its particular topographic
574 and geological characteristics. The dominant southward aspect of the investigated rock glaciers, and their spatial
575 clustering, can explain the lack of correlation between water temperature and the aspect of rock glaciers. We tried to
576 minimise the spatial clustering of measured springs, visiting as many headwater catchments as possible, and taking
577 measurements at the largest number of springs on each catchment. However, due to logistic constraints and inherent
578 characteristics of the study area, a certain degree of spatial clustering was unavoidable. For this reason, the role of terrain
579 aspect as a possible controlling factor on spring-water temperature requires additional investigation.

580

581 **5.4 Geophysics**

582 The inverted resistivity sections obtained for the Preghena Rock Glacier (Fig. 8a and b) show results compatible with the
583 presence of permafrost patches. Even considering the high contact resistance due to the dry weather conditions preceding
584 the survey, and the location of the high resistivity body in the areas known to be the least sensitive of the model (the bed
585 and margins, Binley, 2015), we observe that the obtained resistivity values are typical of frozen materials (Hauck and
586 Kneisel, 2008). [The high resistive area is highlighted by both ERT lines in the overlapping area \(\$x < 70\$ m in Line 1 and](#)
587 [\$x > 100\$ m in Line2, Fig. 8\). The data error of 20% applied in the inversion process was defined using the reciprocal](#)
588 [analysis, which minimise possible inversion artifacts compared to the more commonly used stacking error \(Binley, 2015\).](#)
589 This result agrees with the low temperature of the Preghena Rock Glacier spring, which fluctuates between 1.6 and 1.8°C
590 throughout summer, and it suggests that this rock glacier should be classified as a pseudo-relict rock glacier.

591 In the Bordolona Rock Glacier (Fig. 8c), the frozen layer looks discontinuous in the lower section of the ERT Line, and
592 more continuous and thicker in the upper part, where a younger lobe superposes the main body of the rock glacier. The

593 different resistivity detected in the lower and upper sections of the ERT line can be related to a different percent ice
 594 content in the frozen layers, and/or a different temperature of the ice (Hilbich et al., 2008). These results suggest the
 595 probable presence of permafrost also inside the Bordolona Rock Glacier, which was considered a ‘true’ relict rock glacier
 596 due to its abundant vegetation cover, spring-water temperature above 3°C, and low mean elevation. Based on geophysical
 597 investigations, the Bordolona Rock Glacier too should be classified as a pseudo-relict rock glacier.

598 ~~The acquired data were of lower quality at the Preghena Rock Glacier, due to the high contact resistance. The acquired~~
 599 ~~data were of lower quality at the Preghena Rock Glacier, due to the high contact resistance.~~ More conclusive results
 600 should be obtained by repeating the geophysical surveys under moister conditions, especially at the Preghena Rock
 601 Glacier, and possibly coupling ERT to seismic refraction measurements in order to obtain a reliable estimate of the percent
 602 ice content inside these rock glaciers (Hauck et al., 2011, Wagner et al., 2019, Pavoni et al. 2023).

603

604 5.5 Ice storage in the rock glaciers and glaciers of Val di Sole

605 ~~Given the very different response time of glaciers and rock glaciers to projected atmospheric warming, their relative~~
 606 ~~importance has relevant implications for the current and future hydrological cycle. In the study area, glaciers are shrinking~~
 607 ~~fast, and in 2022 they covered 7.18 km², which is 59% of the 2003 area (area loss rate = 2.6% y⁻¹). For these reasons, and~~
 608 ~~in light of the results of this work, it is interesting to estimate and compare the ice content of rock glaciers and glaciers in~~
 609 ~~Val di Sole, similarly to what was done, for example, by Bolch and Marchenko (2009) in the Northern Tien Shan.~~

610 ~~For glaciers, we estimated residual volumes in 2022 starting from the ice thickness estimates provided for each glacier in~~
 611 ~~the study area by Farinotti et al., (2019). We first calculated the bedrock topography subtracting the ice thickness from~~
 612 ~~the glacier surface DEM (Farinotti et al., 2019). Then we calculated the 2002 glacier thickness subtracting the bedrock~~
 613 ~~topography from a glacier surface DEM surveyed in September 2022 by the Province of Trento. We finally obtained the~~
 614 ~~glacier volumes multiplying the average thickness by the glacier area, and converted the ice volume into the water volume~~
 615 ~~equivalent using a mean ice density of 900 kg m⁻³.~~

616 ~~For rock glaciers, we calculated the total rock glacier volume multiplying their area A by the average thickness provided~~
 617 ~~by the Brenning (2005b) formulation:~~

$$618 \quad T = cA^\gamma \quad (1)$$

619 ~~where T is the average thickness of rock glaciers, and c and γ are constants equal to 50 and 0.2, respectively. To account~~
 620 ~~for the different geometry of intact and relict rock glaciers, we assumed that the volumetric ice content of intact rock~~
 621 ~~glaciers averages 50% (Jones et al., 2018, and references therein), and therefore that T_r for (true) relict rock glaciers is~~
 622 ~~half that of intact rock glaciers (i.e., they are composed only of debris and all the ice melted away). For pseudo-relict rock~~
 623 ~~glaciers we tested various hypotheses of percent ice content, calculating the average thickness T_{pr} as follows:~~

$$624 \quad T_{pr} = T_r + T_{ice} \quad (2)$$

625 ~~where T_{ice} is the average ice thickness, calculated in function of the volumetric percent ice content $\%_{ice}$ as:~~

$$626 \quad T_{ice} = \frac{\%_{ice} T_r}{(1 - \%_{ice})} \quad (3)$$

627 A total glacier ice volume of $251 \times 10^6 \text{ m}^3$, and a corresponding $226 \times 10^6 \text{ m}^3$ water volume equivalent was calculated for
628 Val di Sole in 2022. In comparison, the water volume equivalent of intact rock glaciers is $42.7 \times 10^6 \text{ m}^3$.

629 A water volume equivalent between 3.7 and $17.7 \times 10^6 \text{ m}^3$, averaging $10.7 \times 10^6 \text{ m}^3$, can be estimated assuming that 38%
630 of the total area of relict rock glaciers contains permafrost, and that the average ice content ranges between 5% and 20%
631 in volume. ~~Calculations of the ice contained in the pseudo-relict rock glaciers of the study area assumed that 50% of the~~
632 ~~total area of relict rock glaciers contains permafrost (Section 4.2.1) and that the average ice content ranges between 5%~~
633 ~~and 20% in volume.~~ This range is a first hypothesis based on the few geophysical data available at pseudo-relict rock
634 glaciers (Delaloye, 2004; Colucci et al., 2019; Pavoni et al., 2023; this work). To our knowledge, the amount of ice in
635 pseudo-relict rock glaciers has yet to be quantified.

636 Even if preliminary and affected by significant uncertainty, these estimates provide an order of magnitude of water stored
637 as ice in the rock glaciers of Val di Sole. The water equivalent ratio for rock glacier ice versus glacier ice averages 1:4.2
638 1 and ranges between 1:3.7-6 and 1:4.98, considering minimum and maximum estimates reported above. Importantly,
639 based on these calculations, 2023% of the total rock glacier water volume would be stored inside pseudo-relict rock
640 glaciers. Even assuming the lower bound of percent ice content (5%), pseudo-relict rock glaciers would contribute to a
641 significant 89% of the total rock glacier water volume.

642 ~~Based on the more conservative estimate reported in Section 5.2 for the frequency of pseudo-relict rock glaciers (38%~~
643 ~~instead of 50% of the total area covered by rock glaciers classified as relict), the water equivalent ratio for rock glacier~~
644 ~~ice versus glacier ice would average 1:4.3 and would range between 1:3.9 and 1:4.9, with 18% of the total rock glacier~~
645 ~~water volume stored inside pseudo-relict rock glaciers. Even if a little smaller, these numbers do not change significantly~~
646 ~~the meaning of the results.~~

647

648 6. Concluding remarks

649 We have surveyed spring-water temperature in an area of 795 km^2 in Val di Sole, to understand the influence of
650 topographic and geomorphological factors, and to test if it can be used to preliminary differentiate ~~intact-active/transitional~~
651 and relict rock glaciers. Spring-water temperature measurements enabled to characterise a large number of rock glaciers,
652 and to provide a first estimate of the frequency of pseudo-relict rock glaciers in this area. Overall, our results point to a
653 significant hydrological importance of rock glaciers classified as relict in the study area, which is expected to increase in
654 the future due to atmospheric warming.

655 In general, we have found that the spatial variability of spring-water temperature is controlled by elevation, aspect and
656 the presence of rock glaciers in the upslope area. Compared to other landforms in the upslope area, rock glaciers have
657 colder springs, irrespective of their activity state.

658 The spring-water temperature of rock glaciers classified so far as relict is higher and with larger spatial variability
659 compared to ~~intact-active/transitional~~ rock glaciers. However, there is a remarkable ~50% (38% excluding extremes)
660 overlap in the spring temperature range of the two rock glacier groups. Relict rock glaciers tend to have colder springs if
661 their surface is blocky and scarcely covered by (cold-adapted) vegetation.

662 The spring-water temperature data suggest that one third of rock glaciers classified as relict might be actually pseudo-
663 relict, thus containing permafrost. The exact percentage cannot be derived unambiguously from spring-water temperature

664 because i) other evidence is required to confirm inference from water temperature, ii) there is uncertainty in the
665 classification of the activity state of rock glaciers, iii) there is geophysical evidence that rock glaciers containing
666 permafrost may have 'warm' springs (up to 3.7°C), and consequently iv) there is uncertainty in the definition of the
667 thresholds for differentiate among absent/possible/probable permafrost categories.

668 Despite these uncertainties, our study shows that rock-glacier spring-water temperature can provide a pilot approach to
669 estimate the spatial distribution of permafrost in vast areas, and an auxiliary element to the classification of rock glaciers,
670 whose permafrost content might otherwise go underestimated. This method can be applied in other mountainous regions,
671 with the possible exception of arid/semi-arid regions where the presence of springs is scarce.

672 Geophysics applied to two rock glaciers classified as relict enabled to detect the presence of permafrost. While the blocky
673 Preghena Rock Glacier, whose spring temperature was < 1.8°C throughout the summer, was expected to contain
674 permafrost, its occurrence in the Bordolona Rock Glacier was not expected, because it is entirely covered by dense
675 vegetation and its spring temperature reached 3.7°C in late summer.

676 Preliminary calculations of water resources stored as ice inside the rock glaciers of Val di Sole reveal that they amount to
677 ~24% of the water volume equivalent stored in glaciers, which are disappearing very fast. Remarkably, 20% of the total
678 rock glacier water volume is stored inside rock glaciers classified as relict.

679 This study highlights the need for additional investigations and improved understanding of these periglacial landforms.
680 In particular, the possible presence of permafrost in a large fraction of rock glaciers classified as relict poses critical
681 questions regarding the origin, preservation, current behaviour, seasonal dynamics, and future evolution of this
682 permafrost. Thorough study of pseudo-relict rock glaciers is required for understanding the transitionevolution between
683 intact-active, transitional and relict landforms, which is important in view of current and projected climate change.

684

685


686

687

688

689 **Appendix A**

690



691

692 Figure A1: Spring-water temperature for intact rock glaciers with and without open work deposits on their surface. Boxes
 693 indicate the 25th and 75th percentile, whiskers indicate the 10th and 90th percentile, whereas the horizontal solid and dashed
 694 lines within the box mark the median and the mean, respectively. Maximum and minimum values are represented by dots.
 695 Sample size (n) is reported above the boxplots.

696

697

698 **Data availability**

699 Data are available from the corresponding author upon reasonable request.

700

701 **Author contributions**

702 LC designed the methodological approach and carried out the sampling campaigns with the support of AA, RS, MT, TZ
 703 and GZ. MP and JB carried out the geophysical surveys in cooperation with LC, CM and MZ and interpreted the results.
 704 GZ, LC and AA performed the statistical analyses of the dataset. LC prepared the first draft of the manuscript with
 705 contributions from GZ, MP and CM. All authors contributed to the editing of the manuscript.

706

707 **Competing interests**

708 The contact author has declared that none of the authors has any competing interests.

709

710 **Acknowledgments**

711 The authors acknowledge the editor and reviewers for their comments and suggestions.

712

713 **Financial support**

714 This study was carried out within the RETURN Extended Partnership and received funding from the European Union
715 Next-GenerationEU (National Recovery and Resilience Plan – NRRP, Mission 4, Component 2, Investment 1.3 – D.D.
716 1243 2/8/2022, PE0000005). LC, GZ, JB and RS also acknowledge the support of the project PRIN 2022
717 “SUBSURFACE – Ecohydrological and environmental significance of subsurface ice in alpine catchments” (code:
718 2022AL7WKC) (National Recovery and Resilience Plan – NRRP, Mission 4, Component 2, Investment 1.1 – D. D. 104
719 2/2/2022).

720

721 References

722 [Amschwand, D., Scherler, M., Hoelzle, M., Krummenacher, B., Haberkorn, A., Kienholz, C., and Gubler, H.: Surface](https://doi.org/10.5194/tc-18-2103-2024)
723 [heat fluxes at coarse blocky Murtèl rock glacier \(Engadine, eastern Swiss Alps\), *The Cryosphere*, 18, 2103–2139,](https://doi.org/10.5194/tc-18-2103-2024)
724 [https://doi.org/10.5194/tc-18-2103-2024, 2024.](https://doi.org/10.5194/tc-18-2103-2024)

725 Barsch, D.: Rockglaciers: indicators for the present and former geocology in high mountain environments, Springer
726 Berlin Heidelberg, Berlin, Heidelberg, 218 pp., <https://doi.org/10.2307/3060377>, 1996.

727 Bertone, A., Barboux, C., Bodin, X., Bolch, T., Brardinoni, F., Caduff, R., Christiansen, H. H., Darrow, M. M., Delaloye,
728 R., Etzelmüller, B., Humlum, O., Lambiel, C., Lilleøren, K. S., Mair, V., Pellegrinon, G., Rouyet, L., Ruiz, L., and Strozzi,
729 T.: Incorporating InSAR kinematics into rock glacier inventories: insights from 11 regions worldwide, *The Cryosphere*,
730 16, 2769–2792, <https://doi.org/10.5194/tc-16-2769-2022>, 2022.

731 Binley, A.: Tools and Techniques: Electrical Methods, in: *Treatise on Geophysics: Second Edition*, vol. 11, Elsevier,
732 233–259, <https://doi.org/10.1016/B978-0-444-53802-4.00192-5>, 2015.

733 Binley, A. and Kemna, A.: DC Resistivity and Induced Polarization Methods, in: *Hydrogeophysics*, Springer Netherlands,
734 Dordrecht, 129–156, https://doi.org/10.1007/1-4020-3102-5_5, 2005.

735 Blanchy, G., Saneiyani, S., Boyd, J., McLachlan, P., and Binley, A.: ResIPy, an intuitive open source software for complex
736 geoelectrical inversion/modeling, *Comput. Geosci.*, 137, 104423, <https://doi.org/10.1016/j.cageo.2020.104423>, 2020.

737 Boeckli, L., Brenning, A., Gruber, S., and Noetzli, J.: Permafrost distribution in the European Alps: Calculation and
738 evaluation of an index map and summary statistics, *Cryosphere*, 6, 807–820, <https://doi.org/10.5194/tc-6-807-2012>, 2012.

739 Bolch, T. and Marchenko, S.: Significance of glaciers, rockglaciers and ice-rich permafrost in the Northern Tien Shan as
740 water towers under climate change conditions. In: Braun, Ludwig N; Hagg, Wilfried; Severskiy, Igor V; Young, Gordon.
741 *Assessment of Snow, Glacier and Water Resources in Asia: Selected papers from the Workshop in Almaty, Kazakhstan,*
742 2006. Koblenz: IHP UNESCO, 132-144, 2009.

743 Bollati, I. M., Cerrato, R., Lenz, B. C., Vezzola, L., Giaccone, E., Viani, C., Zanoner, T., Azzoni, R. S., Masseroli, A.,
744 Pellegrini, M., Scapozza, C., Zerboni, A., and Guglielmin, M.: Geomorphological map of the Val Viola Pass (Italy-
745 Switzerland), *Geogr. Fis. e Din. Quat.*, 41, 105–114, <https://doi.org/10.4461/GFDQ.2018.41.16>, 2018.

746 Brenning, A.: Geomorphological, hydrological and climatic significance of rock glaciers in the Andes of Central Chile
747 (33–35 S). *Permafr. Periglac. Process.*, 16(3), 231-240, <https://doi.org/10.1002/ppp.528>, 2005a.

748 Brenning, A.: Climatic and geomorphological controls of rock glaciers in the Andes of Central Chile: combining statistical
749 modelling and field mapping, Ph.D thesis, Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche
750 Fakultät II, 2005b.

751 Brighenti, S., Tolotti, M., Bruno, M. C., Engel, M., Wharton, G., Cerasino, L., Mair, V., and Bertoldi, W.: After the peak
752 water: the increasing influence of rock glaciers on alpine river systems, *Hydrol. Process.*, 33, 2804–2823,
753 <https://doi.org/10.1002/hyp.13533>, 2019.

754 Brighenti, S., Hotaling, S., Finn, D. S., Fountain, A. G., Hayashi, M., Herbst, D., Saros, J. E., Tronstad, L. M., and Millar,
755 C. I.: Rock glaciers and related cold rocky landforms: Overlooked climate refugia for mountain biodiversity, *Glob. Chang.*
756 *Biol.*, 27, 1504–1517, <https://doi.org/10.1111/gcb.15510>, 2021.

757 Carturan, L., Fontana, G. D., and Borga, M.: Estimation of winter precipitation in a high-altitude catchment of the Eastern
758 Italian Alps: Validation by means of glacier mass balance observations, *Geogr. Fis. e Din. Quat.*, 35, 37–48,
759 <https://doi.org/10.4461/GFDQ.2012.35.4>, 2012.

760 Carturan, L., Zuecco, G., Seppi, R., Zanoner, T., Borga, M., Carton, A., and Dalla Fontana, G.: Catchment-Scale
761 Permafrost Mapping using Spring Water Characteristics, *Permafr. Periglac. Process.*, 27, 253–270,
762 <https://doi.org/10.1002/ppp.1875>, 2016.

763 Carturan, L., De Blasi, F., Cazorzi, F., Zoccatelli, D., Bonato, P., Borga, M., and Dalla Fontana, G.: Relevance and Scale
764 Dependence of Hydrological Changes in Glacierized Catchments: Insights from Historical Data Series in the Eastern
765 Italian Alps, *Water*, 11, 89, <https://doi.org/10.3390/w11010089>, 2019.

766 Charton, J., Verfaillie, D., Jomelli, V., and Francou, B.: Early Holocene rock glacier stabilisation at col du Lautaret
767 (French Alps): Palaeoclimatic implications, *Geomorphology*, 394, 107962,
768 <https://doi.org/10.1016/j.geomorph.2021.107962>, 2021.

769 Chen, J. and Ohmura, A.: Estimation of Alpine glacier water resources and their change since the 1870s, *IAHS publ*, 193,
770 127–135, 1990.

771 [Chiesa, S., Micheli, P., Cariboni, M., Tognini, P., Motta, D., Longhin, M., Zambotti, G., Marcato, E., and Ferrario, A.:](#)
772 [Note illustrative della Carta Geologica d'Italia alla scala 1:50.000. Foglio 41 Ponte di Legno., Servizio Geologico d'Italia,](#)
773 [ISPRA, 160 pp., 2010.](#)

774 Colucci, R. R., Forte, E., Žebre, M., Maset, E., Zanettini, C., and Guglielmin, M.: Is that a relict rock glacier?,
775 *Geomorphology*, 330, 177–189, <https://doi.org/10.1016/j.geomorph.2019.02.002>, 2019.

776 Cossart, E., Perrier, R., Schwarz, M., and Houee, S.: Mapping permafrost at a regional scale: interpolation of field data
777 by GIS application in the Upper Durance catchment (Southern French Alps), *GeoFocus*, 205–224, 2008.

778 [Dal Piaz, G., Castellarin, A., Martin, S., Selli, L., Carton, A., Pellegrini, G., Casolari, E., Daminato, F., Montresor, L.,](#)
779 [Picotti, V., Prosser, G., Santuliana, E., and Cantelli, L.: Note Illustrative Della Carta Geologica Alla Scala 1:50.000,](#)
780 [Foglio 042 - Malè, Servizio Geologico d'Italia, ISPRA, 143 pp., 2007.](#)

781 Day-Lewis, F. D., Johnson, C. D., Singha, K., and Lane, J. W. J.: Best practices in electrical resistivity imaging: Data
782 collection and processing, and application to data from Corinna, Maine, EPA report, Boston, MA, 2008.

Formattato: Italiano (Italia)

Formattato: Italiano (Italia)

783 Delaloye, R.: Contribution à l'étude du pergélisol de montagne en zone marginale, 244 pp., 2004.

784 Delaloye, R., and Lambiel, C.: Evidence of winter ascending air circulation throughout talus slopes and rock glaciers
785 situated in the lower belt of alpine discontinuous permafrost (Swiss Alps). *Norsk Geografisk Tidsskrift-Norwegian*
786 *Journal of Geography*, 59(2), 194-203, 2005.

787 Delaloye, R., Reynard, E., Lambiel, C., Marescot, L., and Monnet, R.: Thermal anomaly in a cold scree slope (Creux du
788 Van, Switzerland), in: *Proceedings of the Eighth International Conference of Permafrost*, Zürich, Switzerland., 175–180,
789 2003.

790 Dlabáčková, T., Engel, Z., Uxa, T., Braucher, R., and Team, A.: ¹⁰Be exposure ages and paleoenvironmental significance
791 of rock glaciers in the Western Tatra Mts., Western Carpathians, *Quat. Sci. Rev.*, 312, 108147,
792 <https://doi.org/10.1016/j.quascirev.2023.108147>, 2023.

793 Engelhardt, M., Schuler, T. V., and Andreassen, L. M.: Contribution of snow and glacier melt to discharge for highly
794 glacierised catchments in Norway, *Hydrol. Earth Syst. Sci.*, 18, 511–523, <https://doi.org/10.5194/hess-18-511-2014>,
795 2014.

796 Frauenfelder, R., Allgöwer, B., Haeberli, W., and Hoelzle, M.: Permafrost Investigations With GIS - A Case Study in the
797 Fletschhorn Area , Wallis , Swiss Alps, in: *Seventh International Conference on Permafrost*, 291–295, 1998.

798 Frauenfelder, R., Haeberli, W., Hoelzle, M., and Maisch, M.: Using relict rockglaciers in GIS-based modelling to
799 reconstruct Younger Dryas permafrost distribution patterns in the Err-Julier area, Swiss Alps, *Nor. Geogr. Tidsskr.*, 55,
800 195–202, <https://doi.org/10.1080/00291950152746522>, 2001.

801 Frei, C., and Schär, C.: A precipitation climatology of the Alps from high-resolution rain-gauge observations. *Int. J.*
802 *Climatol.*, 18(8), 873-900, [https://doi.org/10.1002/\(SICI\)1097-0088\(19980630\)18:8<873::AID-JOC255>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1097-0088(19980630)18:8<873::AID-JOC255>3.0.CO;2-9),
803 1998.

804 Gude, M., Dietrich, S., Mausbacher, R., Hauck, C., Molenda, R., Ruzicka, V., and Zacharda, M.: Probable occurrence of
805 sporadic permafrost in non-alpine scree slopes in central Europe, in: *Proceedings 8th International Conference on*
806 *Permafrost*, 331–336, 2003.

807 Guglielmin, M.: Il permafrost alpino: concetti, morfologia e metodi di individuazione (con tre indagini esemplificate in
808 alta Valtellina) / di Mauro Guglielmin; con contributi di Adalberto Notarpietro, Centro di studio per la geodinamica
809 alpina e quaternaria, Milano, 1997.

810 Haeberli, W.: Untersuchungen Zur Verbreitung Von Permafrost Zwischen Flueelapass Und Piz Grialetsch
811 (Graubuenden)., *Versuchsanstalt für Wasserbau, Hydrologie und Glaziologie an der ETH*, 1975.

812 Haeberli, W.: *Creep of Mountain Permafrost: Internal Structure and Flow of Alpine Rock Glaciers.*, 1985.

813 Haeberli, W., Schaub, Y., and Huggel, C.: Increasing risks related to landslides from degrading permafrost into new lakes
814 in de-glaciating mountain ranges, *Geomorphology*, 293, 405–417, <https://doi.org/10.1016/j.geomorph.2016.02.009>,
815 2017.

816 Harrington, J. S., Mozil, A., Hayashi, M. and Bentley, L. R.: Groundwater flow and storage processes in an inactive rock
817 glacier, *Hydrol. Process.*, 32(20), 3070-3088, <https://doi.org/10.1002/hyp.13248>, 2018.

Codice campo modificato

- 818 Harris, S. A. and Pedersen, D. E.: Thermal regimes beneath coarse blocky materials, *Permafr. Periglac. Process.*, 9, 107–
819 120, [https://doi.org/10.1002/\(SICI\)1099-1530\(199804/06\)9:2<107::AID-PPP277>3.0.CO;2-G](https://doi.org/10.1002/(SICI)1099-1530(199804/06)9:2<107::AID-PPP277>3.0.CO;2-G), 1998.
- 820 Hauck, C. and Kneisel, C.: *Applied Geophysics in Periglacial Environments*, edited by: Hauck, C. and Kneisel, C.,
821 Cambridge University Press, 1–248 pp., <https://doi.org/10.1017/CBO9780511535628>, 2008.
- 822 Hauck, C., Böttcher, M., and Maurer, H.: A new model for estimating subsurface ice content based on combined electrical
823 and seismic data sets, *The Cryosphere*, 5, 453–468, <https://doi.org/10.5194/tc-5-453-2011>, 2011. Hilbich, C., Hauck, C.,
824 Hoelzle, M., Scherler, M., Schudel, L., Völksch, I., Vonder Mühl, D., and Mäusbacher, R.: Monitoring mountain
825 permafrost evolution using electrical resistivity tomography: A 7-year study of seasonal, annual, and long-term variations
826 at Schilthorn, Swiss Alps, *J. Geophys. Res. Earth Surf.*, 113, F01S90, <https://doi.org/10.1029/2007JF000799>, 2008.
- 827 Ikeda, A. and Matsuoka, N.: Degradation of talus-derived rock glaciers in the Upper Engadin, Swiss Alps. *Permafr.*
828 *Periglac. Process.*, 13, 145–161, <https://doi.org/10.1002/ppp.413>, 2002.
- 829 Ilyashuk, B. P., Ilyashuk, E. A., Psenner, R., Tessadri, R., and Koinig, K. A.: Rock glacier outflows may adversely affect
830 lakes: Lessons from the past and present of two neighboring water bodies in a crystalline-rock watershed, *Environ. Sci.*
831 *Technol.*, 48, 6192–6200, <https://doi.org/10.1021/es500180c>, 2014.
- 832 Imhof, M., Pierrehumbert, G., Haerberli, W., and Kienholz, H.: Permafrost investigation in the Schilthorn Massif, Bernese
833 Alps, Switzerland, *Permafr. Periglac. Process.*, 11, 189–206, [https://doi.org/10.1002/1099-1530\(200007/09\)11:3<189::AID-PPP348>3.0.CO;2-N](https://doi.org/10.1002/1099-1530(200007/09)11:3<189::AID-PPP348>3.0.CO;2-N), 2000.
- 835 Isotta, F. A., Frei, C., Weigluni, V., Perčec Tadić, M., Lassègues, P., Rudolf, B., Pavan, V., Cacciamani, C., Antolini, G.,
836 Ratto, S. M., Munari, M., Micheletti, S., Bonati, V., Lussana, C., Ronchi, C., Panettieri, E., Marigo, G., and Vertačnik,
837 G.: The climate of daily precipitation in the Alps: Development and analysis of a high-resolution grid dataset from pan-
838 Alpine rain-gauge data, *Int. J. Climatol.*, 34, 1657–1675, <https://doi.org/10.1002/joc.3794>, 2014.
- 839 Janke, J. R., Bellisario, A. C., and Ferrando, F. A.: Classification of debris-covered glaciers and rock glaciers in the Andes
840 of central Chile, *Geomorphology*, 241, 98–121, <https://doi.org/10.1016/j.geomorph.2015.03.034>, 2015.
- 841 Janke, J. R., Ng, S., and Bellisario, A.: An inventory and estimate of water stored in firn fields, glaciers, debris-covered
842 glaciers, and rock glaciers in the Aconcagua River Basin, Chile, *Geomorphology*, 296, 142–152,
843 <https://doi.org/10.1016/j.geomorph.2017.09.002>, 2017.
- 844 Jones, D. B., Harrison, S., Anderson, K., and Betts, R. A.: Mountain rock glaciers contain globally significant water
845 stores, *Sci. Rep.*, 8, 2834, <https://doi.org/10.1038/s41598-018-21244-w>, 2018.
- 846 Jones, D. B., Harrison, S., Anderson, K., and Whalley, W. B.: Rock glaciers and mountain hydrology: A review, *Earth-*
847 *Science Rev.*, 193, 66–90, <https://doi.org/10.1016/j.earscirev.2019.04.001>, 2019.
- 848 Juliussen, H. and Humlum, O.: Thermal regime of openwork block fields on the mountains Elgåhogna and Sølén, Central-
849 Eastern Norway. *Permafr. Periglac. Process.*, 19 (1), 1-18, <https://doi.org/10.1002/ppp.607>, 2008.
- 850 Kääb, A.: Rock glaciers and protalus forms. In *Encyclopedia of Quaternary Science*, 2nd Edition, Volume 3, SA Elias
851 (editor-in chief). Elsevier: Amsterdam; 535–541, 2013.

Codice campo modificato

- 852 Kellerer-Pirklbauer, A.: Aspects of glacial, paraglacial and periglacial processes and landforms of the Tauern Range,
853 Austria. Doctoral Thesis, University of Graz, 2008.
- 854 Kellerer-Pirklbauer, A.: Long-term monitoring of sporadic permafrost at the eastern margin of the European Alps
855 (Hochreichart, Seckauer Tauern range, Austria), *Permafr. Periglac. Process.*, 30, 260–277,
856 <https://doi.org/10.1002/ppp.2021>, 2019.
- 857 Kellerer-Pirklbauer, A., Lieb, G. K., and Kleinfelchner, H.: A new rock glacier inventory of the eastern European Alps,
858 *Austrian J. Earth Sci.*, 105, 78–93, 2012.
- 859 Kellerer-Pirklbauer, A., Pauritsch, M., Morawetz, R., and Kuehnast, B.: Thickness and internal structure of relict rock
860 glaciers – a challenge for geophysics : Examples from two rock glaciers in the Eastern Alps, in: EGU General Assembly
861 Conference Abstracts, 12581, 2014.
- 862 Kellerer-Pirklbauer, A., Lieb, G. K., and Kaufmann, V.: The dösen rock glacier in central Austria: A key site for
863 multidisciplinary long-term rock glacier monitoring in the eastern alps, *Austrian J. Earth Sci.*, 110, 10–17738,
864 <https://doi.org/10.17738/ajes.2017.0013>, 2017.
- 865 Kofler, C., Steger, S., Mair, V., Zebisch, M., Comiti, F. and Schneiderbauer, S.: An inventory-driven rock glacier status
866 model (intact vs. relict) for South Tyrol, Eastern Italian Alps. *Geomorphology*, 350, 106887,
867 <https://doi.org/10.1016/j.geomorph.2019.106887>, 2020.
- 868 Lambiel, C. and Reynard, E.: Regional modelling of present, past and future potential distribution of discontinuous
869 permafrost based on a rock glacier inventory in the Bagnes-Hérémence area (Western Swiss Alps), *Nor. Geogr. Tidsskr.*
870 - *Nor. J. Geogr.*, 55, 219–223, <https://doi.org/10.1080/00291950152746559>, 2001.
- 871 Lewkowicz, A. G., Etmüller, B., and Smith, S. L.: Characteristics of Discontinuous Permafrost based on Ground
872 Temperature Measurements and Electrical Resistivity Tomography, Southern Yukon, Canada, *Permafr. Periglac.*
873 *Process.*, 22, 320–342, <https://doi.org/10.1002/ppp.703>, 2011.
- 874 Lilleøren, K. S., Etmüller, B., Rouyet, L., Eiken, T., Slinde, G., and Hilbich, C.: Transitional rock glaciers at sea level
875 in northern Norway, *Earth Surf. Dynam.*, 10, 975–996, <https://doi.org/10.5194/esurf-10-975-2022>, 2022. Millar, C.I. and
876 Westfall, R.D.: Geographic, hydrological, and climatic significance of rock glaciers in the Great Basin, USA. *Arct.*
877 *Antarct. Alp. Res.*, 51(1), 232–249, <https://doi.org/10.1080/15230430.2019.1618666>, 2019.
- 878 [Martin, S., Montesor, L., Mair, V., Pellegrini, G., Avanzini, M., Fellin, G., Gambiullara, R., Tumiat, S., Santuliana, E.,](#)
879 [Monopoli, B., Gaspari, D., Sapigni, M., and Surian, N.: Note illustrative della Carta Geologica d'Italia alla scala 1:50.000,](#)
880 [Foglio 025 Rabbi., Servizio Geologico d'Italia, ISPRA, 187 pp., 2009.](#)
- 881 [Montrasio A, Berra F, Cariboni M, Ceriani M, Deichmann N, Ferliga C, Gregnanin A, Guerra S, Guglielmin M, Jadoul](#)
882 [F, Longhin M, Mair V, Mazzoccola D, Sciesa E, and Zappone A: Note illustrative della Carta Geologica d'Italia alla scala](#)
883 [1:50.000 - Foglio 024 Bormio., Servizio Geologico d'Italia, ISPRA, 150 pp., 2012.](#)
- 884 Morard, S., Delaloye, R., and Dorthe, J.: Seasonal thermal regime of a mid-latitude ventilated debris accumulation, in:
885 *Proceedings of the 9th International Conference on Permafrost, Fairbanks, Alaska, 1233–1238, 2008.*

Formattato: Italiano (Italia)

Formattato: Italiano (Italia)

886 Pavoni, M., Boaga, J., Carrera, A., Zuecco, G., Carturan, L., and Zumiani, M.: Brief communication: Mountain permafrost
887 acts as an aquitard during an infiltration experiment monitored with electrical resistivity tomography time-lapse
888 measurements, *Cryosph.*, 17, 1601–1607, <https://doi.org/10.5194/tc-17-1601-2023>, 2023.

889 Popescu, R.: Permafrost investigations in Iezer Mountains, Southern Carpathians, *Rev. Geomorfol.*, 20, 102–122,
890 <https://doi.org/10.21094/rg.2018.033>, 2018.

891 [RGIK: Guidelines for inventorying rock glaciers: baseline and practical concepts \(version 1.0\). IPA Action Group Rock](#)
892 [glacier inventories and kinematics, 25 pp. DOI: 10.51363/unifr.srr.2023.002, 2023.](#)

893 Scotti, R., Brardinoni, F., Alberti, S., Frattini, P. and Crosta, G.B.: A regional inventory of rock glaciers and protalus
894 ramparts in the central Italian Alps. *Geomorphology*, 186, 136-149, <https://doi.org/10.1016/j.geomorph.2012.12.028>,
895 2013.

896 Salvatore, M. C., Zanoner, T., Baroni, C., Carton, A., Banchieri, F. A., Viani, C., Giardino, M., and Perotti, L.: The state
897 of Italian glaciers: A snapshot of the 2006-2007 hydrological period, *Geogr. Fis. e Din. Quat.*, 38, 175–198,
898 <https://doi.org/10.4461/GFDQ.2015.38.16>, 2015.

899 Sannino, C., Borruso, L., Mezzasoma, A., Battistel, D., Ponti, S., Turchetti, B., Buzzini, P. and Guglielmin, M.: Abiotic
900 factors affecting the bacterial and fungal diversity of permafrost in a rock glacier in the Stelvio Pass (Italian Central Alps).
901 *Appl. Soil Ecol.*, 166, 104079, <https://doi.org/10.1016/j.apsoil.2021.104079>, 2021.

902 Scapozza, C.: Contributo dei metodi termici alla prospezione del permafrost montano : esempi dal massiccio della Cima
903 di Gana Bianca (Val Blenio , Svizzera), *Boll. della Soc. Ticin. di Sci. Nat.*, 66, 55–66, 2009.

904 Schaffer, N., MacDonell, S., Réveillet, M., Yáñez, E. and Valois, R.: Rock glaciers as a water resource in a changing
905 climate in the semiarid Chilean Andes. *Reg. Environ. Change*, 19, 1263-1279, [https://doi.org/10.1007/s10113-018-01459-](https://doi.org/10.1007/s10113-018-01459-3)
906 [3](#), 2019.

907 Schrott, L., Otto, J. C., and Keller, F.: Modelling alpine permafrost distribution in the Hohe Tauern region, Austria.
908 *Austrian J. Earth Sci.*, 105(2), 2012.

909 [Seppi, R.: I rock glaciers delle Alpi Centrali come indicatori ambientali \(Gruppo Adamello-Presanella e settore orientale](#)
910 [del Gruppo Ortles-Cevedale\) - Rock glaciers of the Central Alps as environmental indicators \(Adamello-Presanella Group](#)
911 [and eastern sector of the Ortles-Cevedale Group\). Phd Thesis, 199 pp., doi: 10.13140/RG.2.1.1186.5682, 2006.](#)

912 Seppi, R., Carton, A., and Baroni, C.: Rock glacier relitti e antica distribuzione del permafrost nel Gruppo Adamello
913 Presanella (Alpi Centrali), *Alp. Mediterr. Quat.*, 23, 137–144, 2010.

914 Seppi, R., Carton, A., Zumiani, M., Dall'Amico, M., Zampedri, G., and Rigon, R.: Inventory, distribution and topographic
915 features of rock glaciers in the southern region of the Eastern Italian Alps (Trentino), *Geogr. Fis. e Din. Quat.*, 35, 185–
916 197, <https://doi.org/10.4461/GFDQ.2012.35.17>, 2012.

917 Seppi, R., Carturan, L., Carton, A., Zanoner, T., Zumiani, M., Cazorzi, F., Bertone, A., Baroni, C., and Salvatore, M. C.:
918 Decoupled kinematics of two neighbouring permafrost creeping landforms in the Eastern Italian Alps, *Earth Surf. Process.*
919 *Landforms*, 44, 2703–2719, <https://doi.org/10.1002/esp.4698>, 2019.

Formattato: Inglese (Regno Unito)

920 Slangen, A. B. A. and van de Wal, R. S. W.: An assessment of uncertainties in using volume-area modelling for computing
921 the twenty-first century glacier contribution to sea-level change, *Cryosph.*, 5, 673–686, [https://doi.org/10.5194/tc-5-673-](https://doi.org/10.5194/tc-5-673-2011)
922 2011, 2011.

923 [Strobl, B., Etter, S., van Meerveld, I., and Seibert, J.: Accuracy of crowdsourced streamflow and stream level class
924 estimates. *Hydrolog. Sci. J.*, 65\(5\), 823-841, <https://doi.org/10.1080/02626667.2019.1578966>, 2020.](#)

925 Strozzi, T., Kääb, A., and Frauenfelder, R.: Detecting and quantifying mountain permafrost creep from in situ inventory,
926 space-borne radar interferometry and airborne digital photogrammetry, *Int. J. Remote Sens.*, 25, 2919–2931,
927 <https://doi.org/10.1080/0143116042000192330>, 2004.

928 Tapia-Baldis, C. and Trombotto-Liaudat, D.: Permafrost model in coarse-blocky deposits for the Dry Andes, Argentina
929 (28– 33° S), *Cuadernos de Investigación Geográfica*, 46, 33–58, <https://doi.org/10.18172/cig.3802>, 2020.

930 Thies, H., Nickus, U., Tolotti, M., Tessadri, R., and Krainer, K.: Evidence of rock glacier melt impacts on water chemistry
931 and diatoms in high mountain streams, *Cold Reg. Sci. Technol.*, 96, 77–85,
932 <https://doi.org/10.1016/j.coldregions.2013.06.006>, 2013.

933 Wagner, T., Pauritsch, M., and Winkler, G.: Impact of relict rock glaciers on spring and stream flow of alpine watersheds:
934 examples of the Niedere Tauern Range, Eastern Alps (Austria), *Aust. J. Earth Sci.*, 109, 84–98. doi:
935 10.17738/ajes.2016.0006, 2016.

936 [Wagner, T., Pauritsch, M., Mayaud, C., Kellerer-Pirklbauer, A., Thalheim, F., & Winkler, G.: Controlling factors of
937 microclimate in blocky surface layers of two nearby relict rock glaciers \(Niedere Tauern Range, Austria\). *Geogr. Ann.
938 A*, 101\(4\), 310–333, <https://doi.org/10.1080/04353676.2019.1670950>, 2019.](#)

939 Wagner, T., Kainz, S., Helfricht, K., Fischer, A., Avian, M., Krainer, K., and Winkler, G.: Assessment of liquid and solid
940 water storage in rock glaciers versus glacier ice in the Austrian Alps, *Sci. Total Environ.*, 800, 149593,
941 <https://doi.org/10.1016/j.scitotenv.2021.149593>, 2021.

942 Winkler, G., Wagner, T., Pauritsch, M., Birk, S., Kellerer-Pirklbauer, A., Benischke, R., Leis, A., Morawetz, R.,
943 Schreiechener, M. G., and Hergarten, S.: Identification and assessment of groundwater flow and storage components of
944 the relict Schöneben Rock Glacier, Niedere Tauern Range, Eastern Alps (Austria), *Hydrogeol. J.*, 24, 937–953,
945 <https://doi.org/10.1007/s10040-015-1348-9>, 2016.

946 Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Hoelzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlstrøm,
947 A. P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L. N., Cáceres, B. E., Casassa, G., Cobos, G., Dávila, L. R.,
948 Delgado Granados, H., Demuth, M. N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Ove Hagen, J.,
949 Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V. V., Portocarrero, C. A., Prinz, R., Sangewar, C. V.,
950 Severskiy, I., Sigurdsson, O., Soruco, A., Usabaliyev, R., and Vincent, C.: Historically unprecedented global glacier
951 decline in the early 21st century, *J. Glaciol.*, 61, 745–762, <https://doi.org/10.3189/2015JG15J017>, 2015.

952

953

954

