



1 **Spring-water temperature suggests widespread occurrence of**
2 **Alpine permafrost in pseudo-relict rock glaciers**

3 Luca Carturan¹, Giulia Zuecco^{1,2}, Angela Andreotti¹, Jacopo Boaga³, Costanza Morino¹, Mirko
4 Pavoni³, Roberto Seppi⁴, Monica Tolotti⁵, Thomas Zanoner⁴, Matteo Zumiani⁶

5 ¹Department of Land, Environment, Agriculture and Forestry, University of Padova, Legnaro, Italy

6 ²Department of Chemical Sciences, University of Padova, Padova, Italy

7 ³Department of Geosciences, University of Padova, Padova, Italy

8 ⁴Department of Earth and Environmental Sciences, Pavia, Italy

9 ⁵Fondazione Edmund Mach - Istituto Agrario San Michele All'Adige, S. Michele all'Adige, Italy

10 ⁶Geological Service, Autonomous Province of Trento, Trento, Italy

11 *Correspondence to:* Luca Carturan (luca.carturan@unipd.it)

12 **Abstract**

13 Runoff originating from ground ice contained in landforms like rock glaciers and talus slopes represents an important
14 water supply for the lowlands. Pseudo-relict rock glaciers host patchy permafrost, but appear to be visually relict, and
15 therefore can be misinterpreted by using standard classification approaches. Permafrost content, spatial distribution and
16 frequency of this type of rock glaciers are poorly known. Therefore, identifying pseudo-relict rock glaciers that might still
17 host permafrost, and potentially ice, is crucial for understanding their hydrological role in a climate change context.

18 This work analyses rock-glacier spring-water temperature in a 795 km² catchment in the Eastern Italian Alps to understand
19 how many rock glaciers classified as relict could have spring-water temperature comparable to intact rock glaciers, as
20 possible evidence of their pseudo-relict nature. Spring-water temperature, often used as auxiliary to other approaches for
21 specific sites, was used for a preliminary estimate of the permafrost presence in 50 rock glaciers classified as relict. In
22 addition, we present electrical resistivity tomography (ERT) results on two relict rock glaciers with opposing spring-water
23 temperature and surface characteristics to constrain spring-water temperature results at local scale.

24 The results show that about 50% of rock glaciers classified as relict might be pseudo-relict, thus potentially containing
25 permafrost. Both supposedly relict rock glaciers investigated by geophysics contain frozen sediments. The majority of
26 cold springs are mainly associated with rock glaciers with blocky and sparsely vegetated surface, but geophysics suggest
27 that permafrost may also exist in rock glaciers below 2000 m a.s.l., entirely covered by vegetation and with spring-water
28 temperature up to 3.7°C. We estimate that pseudo-relict rock glaciers might contain a significant portion (20%) of all the
29 ice stored in the rock glaciers in the study area. These results highlight the relevance of pseudo-relict rock glaciers in
30 periglacial environments. Even if not a conclusive method, spring-water-temperature analyses can be used to preliminarily
31 distinguish between relict and pseudo-relict rock glaciers in wide regions.

32

33



34 1. Introduction

35 Timings and magnitude of cryosphere runoff have high climatic sensitivity and are impacted by the current changes of
36 Earth's climate (Engelhardt et al., 2014; Zemp et al., 2015; Carturan et al., 2019). Moreover, a deterioration of the water
37 quality has been reported for springs fed by melting permafrost (Thies et al., 2013; Ilyashuk et al., 2014). Due to glacier
38 decline, in the last decades growing attention has been given to other water reservoirs, such as subsurface ice, including
39 debris-covered glacier ice and, in particular, ground ice stored in periglacial landforms such as rock glaciers and glacial-
40 permafrost composite landforms (e.g., Brighenti et al., 2019; Jones et al., 2019; Schaffer et al., 2019; Seppi et al., 2019;
41 Wagner et al., 2021). Projection of ice loss rates indicates that in the second half of the 21st century more subsurface ice
42 may be preserved than glacier surface ice because of their different response times to atmospheric warming (Haeberli et
43 al., 2017). Subsurface ice is therefore expected to significantly contribute to stream runoff under future climate warming
44 (Janke et al., 2015, 2017).

45 Jones et al. (2018) assessed the importance of ice contained in rock glaciers at global scale, estimating that 62.02 ± 12.40
46 Gt of ice is contained in intact rock glaciers. Even though relict rock glaciers should not contain ice (Haeberli, 1985;
47 Barsch, 1996), more recent studies showed that some relict rock glaciers can preserve permafrost and ice far below the
48 regional lower limit of discontinuous permafrost (e.g., Delaloye, 2004; Strozzi et al., 2004; Lewkowicz et al., 2011;
49 Bollati et al., 2018; Colucci et al., 2019).

50 This evidence raises the question whether a significant fraction of rock glaciers classified as relict is actually to be
51 considered 'pseudo-relict', i.e. "*rock glaciers which appear to be visually relict but still contain patches of permafrost*"
52 (Kellerer-Pirklbauer et al., 2012; Kellerer-Pirklbauer, 2008, 2019). This question is relevant because landforms classified
53 as relict in some regions can be up to an order of magnitude larger and more numerous than intact rock glaciers (e.g.,
54 Seppi et al., 2012; Scotti et al., 2013; Kofler et al., 2020), with potentially significant ecological and hydrological impacts
55 (e.g., Brenning, 2005a; Millar and Westfall, 2019; Brighenti et al., 2021; Sannino et al., 2021). According to Jones et al.
56 (2019), identifying and establishing the activity state of rock glaciers is an important initial step in determining their
57 potential hydrological significance.

58 Previous investigations on the possible permafrost content of relict rock glaciers looked at single case studies or small
59 groups of landforms (e.g., Delaloye, 2004; Kellerer-Pirklbauer et al., 2014; Popescu, 2018; Colucci et al., 2019; Pavoni
60 et al., 2023), whereas studies considering a larger number of relict rock glaciers at the regional scale, were mainly focussed
61 on the past distribution of mountain permafrost and on the reconstruction of related paleoclimatic conditions (e.g.,
62 Frauenfelder et al., 2001; Seppi et al., 2010; Charton et al., 2021; Dlabáčková et al., 2023).

63 As a result, the actual distribution, frequency, and ice content of pseudo-relict rock glaciers might be underestimated,
64 with the latter being essential for implementing worldwide estimates of water resources stored in periglacial landforms
65 (e.g., Jones et al., 2018). Detailed geophysical investigation of selected landforms is certainly suitable as a first step
66 towards a better knowledge of pseudo-relict rock glaciers and their ice content. However, due to logistic constraints, this
67 approach cannot be applied to a large number of rock glaciers at the catchment or regional scale. A recent and
68 commendable advance on this topic has been achieved by the proposition of operational guidelines on the InSAR-based
69 kinematic characterization of rock glaciers (Bertone et al., 2022), which can be used for thorough studies of wide areas.
70 However, this approach is not suitable for distinguishing between relict and pseudo-relict rock glaciers, because their
71 surface is motionless.



72 A possible way to investigate the presence of permafrost in these landforms over large areas is by analysing spring-water
73 temperature measured downslope of rock glaciers. Haeberli (1975) proposed the monitoring of spring-water temperature
74 in late summer as useful evidence of permafrost, and various authors employed such method as auxiliary permafrost
75 evidence (e.g., Frauenfelder et al., 1998; Scapozza, 2009; Imhof et al., 2000; Strozzi et al., 2004; Cossart et al., 2008).
76 Carturan et al. (2016) demonstrated that this method can be used successfully for mapping permafrost distribution at the
77 catchment scale. All these works are based on the evidence that, in late summer, spring water affected by permafrost has
78 lower temperature compared to those unaffected, with upper thresholds ranging between 0.9 and 1.1°C for probable
79 permafrost, and between 1.8 and 2.2°C for possible permafrost.

80 In this work, we analyse the spatial variability of spring-water temperature in a 795 km² catchment located in the Eastern
81 Italian Alps, where 338 rock glaciers were inventoried (Seppi et al., 2012), to better understand permafrost distribution.
82 We hypothesise that a significant portion of rock glaciers classified as relict have spring-water temperature comparable
83 to those of intact rock glaciers, as possible evidence of their permafrost content and of their pseudo-relict nature. The
84 specific objectives of this study are to:


- 85 i) analyse the influence of topographic and geomorphological factors on spring-water temperature,
- 86 ii) investigate the main controls on water temperature for springs downslope of rock glaciers, and particularly relict rock
87 glaciers,
- 88 iii) investigate via geophysical analyses the presence of permafrost in two rock glaciers selected for their different spring-
89 water temperature and surface characteristics, to constrain spring-water temperature results at local scale.

90

91 **2. Study area**

92 Val di Sole is located in the upper part of the Noce River catchment, a tributary of the Adige River, which is the main
93 river system in northeastern Italy (Fig. 1). The catchment is 795 km² wide, with elevation ranging between 520 m a.s.l.
94 at the outlet (Mostizzolo) and 3769 m a.s.l. at the summit of Mt. Cevedale, averaging 1705 m a.s.l. (Fig. 1). Metamorphic
95 rocks (mica schists, paragneiss and orthogneiss) prevail in the northern side of the valley, whereas tonalite is found in the
96 southwestern part and dolomites and limestones prevail in the southeastern part.

97



98

99 Figure 1: Geographic location of the study area and of sampled springs. The background is the hillshaded Lidar 2014
100 DEM surveyed by the Provincia Autonoma di Trento (<https://siat.provincia.tn.it/stem/>).

101

102 The catchment includes a glacierised area of 16 km² (in 2006, Salvatore et al., 2015). Bare bedrock and debris are found
103 outside the glaciers down to an elevation of 2700 m, which is the lower regional limit of discontinuous permafrost
104 (Boeckli et al., 2012). A discontinuous cover of alpine meadows and shrubs is present between 2200 m and 2700 m, while
105 below 2000-2200 m forests are dominant. The valley bottom is covered by cultivations and settlements.

106 Val di Sole lies in a transition zone between the “inner dry alpine zone” in the north (Frei and Schär, 1998) and the wetter
107 area under the influence of the Mediterranean Sea in the south. At the valley floor, the annual precipitation averages ~900
108 mm. Precipitation increases with elevation and in the southern part, with a maximum of 1500 mm in the Adamello-
109 Presanella Group (Carturan et al., 2012; Isotta et al., 2014). The mean annual 0°C isotherm is located at 2500 m. The
110 mean annual air temperature variability is dominated by elevation, whereas latitudinal and longitudinal variations are
111 negligible.

112 Seppi et al. (2012) mapped 338 rock glaciers in Val di Sole. The largest part of rock glaciers was classified as relict (229,
113 68% of the total), whereas of the 109 intact rock glaciers only 42 can be classified as active based on multitemporal high-



114 resolution digital elevation models (DEMs). Most rock glaciers (302, 89% of the total) are composed of deposits of
115 metamorphic rocks in the orographic left side of the valley.

116

117 **3. Materials and methods**

118 **3.1 Experimental design**

119 We focussed our investigations on the northern part of Val di Sole because it has a rather homogeneous lithology
120 (metamorphic rocks with predominant micaschists) and mean annual precipitation (1233 mm at 2600 m, Carturan et al.,
121 2016). This was done to minimise the effects of different lithologies and annual precipitation on the spatial variability of
122 spring-water temperature, and to highlight the role of other variables related to their catchment, upslope area or upslope
123 rock glaciers.

124 To obtain statistically meaningful and generalisable results, we designed a sampling scheme for rock-glacier spring-water
125 temperature considering the variability of permafrost-related characteristics in the study area, namely vegetation cover
126 (related to ground temperature and fine debris infill), size (length, area), elevation, slope, aspect, and lithology (Barsch,
127 1996; Haerberli, 1985; Lambiel and Reynard, 2001; Boekli et al., 2012).

128 We inspected these variables, reported for each rock glacier of Val di Sole in the database of Seppi et al. (2012), using a
129 correlation matrix and the Principal Component Analysis. The aim was to evaluate their possible covariance and to
130 optimise the number of variables and their combinations, to be included in the sampling scheme. The analysis revealed
131 high positive covariance between length and area (both related to size). Negative covariance was found between elevation
132 and vegetation cover, and between slope and length/area.

133 Based on these outcomes and considering accessibility, we built a sampling scheme around four variables: i) rock glacier
134 activity, ii) length, iii) mean elevation, and iv) vegetation cover. The last two variables are correlated because intact rock
135 glaciers are at high elevation and almost free from vegetation, and the opposite is true for relict rock glaciers. Vegetation
136 cover is probably one of the few variables that may aid at identifying rock-glacier activity (Ikeda and Matsuoka, 2002;
137 Strozzi et al., 2004, Kofler et al., 2020), and it can vary greatly among rock glaciers at similar elevation. For this reason,
138 we kept both elevation and vegetation, applying a modification to the vegetation-cover classification proposed by Seppi
139 et al. (2012). We distinguish between two classes, namely 'vegetated' and 'non vegetated' for both intact and relict rock
140 glaciers (see Table 1 for threshold values). The vegetation cover was visually estimated in the field and in orthophotos
141 for each rock glacier. Our sampling scheme ensured that at least one rock glacier was sampled for each combination of
142 variables (Table 2). The frequency distribution of rock glacier length and mean elevation was used to identify three
143 terciles, employed for grouping them into short-mid-long rock glaciers and into low-mid-high elevation rock glaciers.
144 Frequency distributions and terciles of intact and relict rock glaciers were calculated separately (Table 2).

145

146

147

148



149 Table 1 - Classification of intact and relict rock glaciers in two different classes of vegetation cover.

| Rock glacier category | Vegetation cover class | Meaning |
|-----------------------|------------------------|-----------------------|
| Intact | Vegetated | Vegetation cover >10% |
| | Non vegetated | Vegetation cover <10% |
| Relict | Vegetated | Vegetation cover >50% |
| | Non vegetated | Vegetation cover <50% |

150

151

152

153 Table 2: Sampling scheme used for water temperature measurements at rock glaciers springs.

| Activity state | Length | Elevation | Vegetation cover | Number of sampled rock glaciers |
|----------------|-------------------------|-------------------------|------------------|---------------------------------|
| Intact | Short (<142 m) | Low (<2634 m) | Non vegetated | 2 |
| | | | Vegetated | none |
| | | Mid (>2634 and <2811 m) | Non vegetated | 2 |
| | | | Vegetated | none |
| | | High (>2811 m) | Non vegetated | 1 |
| | | | Vegetated | none |
| | Mid (>142 and <251 m) | Low (<2596 m) | Non vegetated | 1 |
| | | | Vegetated | none |
| | | Mid (>2596 and <2817 m) | Non vegetated | 1 |
| | | | Vegetated | 3 |
| | | High (>2817 m) | Non vegetated | 2 |
| | | | Vegetated | none |
| Long (>251 m) | Low (<2655 m) | Non vegetated | none | |
| | | Vegetated | 1 | |
| | Mid (>2655 and <2779 m) | Non vegetated | 1 | |
| | | Vegetated | none | |
| | High (>2779 m) | Non vegetated | 3 | |
| | | Vegetated | none | |
| Relict | Short (<180 m) | Low (<2267 m) | Non vegetated | 3 |
| | | | Vegetated | 4 |
| | | Mid (>2267 and <2453 m) | Non vegetated | 1 |
| | | | Vegetated | 2 |
| | High (>2453 m) | Non vegetated | 2 | |
| | | Vegetated | 2 | |
| | | Low (<2255 m) | Non vegetated | 3 |



| | | | | |
|-----------|-------------------------|-------------------------|---------------|---------------|
| | Mid (>180 and <340 m) | | Vegetated | 4 |
| | | Mid (>2255 and <2425 m) | Non vegetated | 1 |
| | | | Vegetated | 2 |
| | | High (>2425 m) | Non vegetated | 2 |
| | | | Vegetated | 3 |
| | | Long (>340 m) | Low (<2222 m) | Non vegetated |
| | Vegetated | | | 4 |
| | Mid (>2222 and <2388 m) | | Non vegetated | 3 |
| | | | Vegetated | 5 |
| | High (>2388 m) | | Non vegetated | 5 |
| Vegetated | | | 3 | |
| | | | Total: | 67 |

154

155

156 3.2 Data collection

157 Water temperature was measured at 220 springs, 133 of which are located downslope of rock glaciers, 81 are located
 158 downslope of other deposits, and 8 are located in bedrock. Springs were sampled from mid-August to mid-October, after
 159 the end of the snowmelt. Most springs have been measured once per year from 2018 to 2020, and a small group of them
 160 was also measured in 2021. In these four years, the total number of single measurements is 540.

161 Based on the sampling scheme (Table 2), we measured spring-water temperature at 17 intact rock glaciers and 50 relict
 162 rock glaciers, which corresponds to 22% of all rock glaciers existing in the study area. All variables' combinations defined
 163 for relict rock glaciers have been sampled, whereas several combinations for intact rock glaciers lack samplings. This was
 164 due to the inexistence of single combinations (e.g., there are no short and vegetated intact rock glaciers at low elevation)
 165 or to the lack of springs and inaccessibility of some rock glaciers.

166 Measurements of spring-water temperature were carried out using a WTW Cond3310 (WTW GmbH, Weilheim,
 167 Germany) and a Testo 110 (Testo AG, Lenzkirch, Germany). These instruments have both 0.1°C resolution, but the WTW
 168 has higher accuracy ($\pm 0.1^\circ\text{C}$) compared to the Testo ($\pm 0.2^\circ\text{C}$), which was used for back-up/validation. Water temperature
 169 measurements were carried out shading the spring from direct sunlight and avoiding probe contact with sediments, rocks,
 170 and vegetation. The calibration of the two instruments was checked at the beginning and at the end of the annual
 171 campaigns using an ice bath. In addition, runoff was visually estimated at each spring.


172

173 3.3 Data analysis

174 Before proceeding with statistical analyses, we preliminary filtered field data to exclude problematic or redundant
 175 measurements. First, we discarded measurements that were clearly affected by very low runoff (< 0.1 l/s). We then selected
 176 one measurement site for each rock glacier and for groups of springs separated less than 10 m from each other. Spring
 177 selection was carried out favouring springs with higher runoff, repeated readings in the four years, closer location to rock
 178 glacier fronts, and with lower interannual temperature variability.



179 After this selection, 131 springs were retained. We characterise the springs using different variables (Table 3), namely
180 the topographic characteristics of the catchments draining to the springs, the activity-state, topographic,
181 geomorphological, and vegetation characteristics of rock glaciers, and the topographic, geomorphological, geological,
182 vegetation and permafrost characteristics of the area immediately upslope of the springs. The latter is defined by the
183 intersection of the catchment perimeter with a circular buffer zone with a radius of 100 m (Fig. 2; Carturan et al., 2016).
184 Details on these variables, the methodology and the data sources (e.g., DEMs, orthophotos, geological maps and literature)
185 employed to derive them are listed and described in Table 3.



186

187 Figure 2: Delimitation of the spring upslope area, defined by the intersection of a circular buffer zone with a radius of
188 100 m over the catchment perimeter. The methodology was introduced and tested in Carturan et al. (2016).

189

190

191

192

193

194

195

196

197

198

199

200

201



202 Table 3: Quantitative and qualitative variables used for characterizing spring areas and for statistical analyses.

| Spatial scale | Variable type | Variable | Classes/acronym | Meaning | |
|---------------------|---------------|---|--|--|---|
| Catchment | Quantitative | Minimum elevation (m a.s.l.) ^a | \ | Spring elevation | |
| | | Maximum elevation (m a.s.l.) ^a | \ | | |
| | | Mean elevation (m a.s.l.) ^a | \ | Half sum of minimum and maximum elevations | |
| | | Planimetric length (m) ^a | \ | | |
| | Qualitative | Mean aspect ^a | 1 | | NW-NE (315° to 45°) |
| | | | 2 | | NE-SE (45° to 135°) and SW-NW (225° to 315°) |
| | | | 3 | | SE-SW (135° to 225°) |
| Spring upslope area | Qualitative | Geomorphology ^{b,g} | VER | | Slope deposit (scree slope or debris cone) |
| | | | GLAC | | Glacial deposit |
| | | | RG | | Rock glacier |
| | | | PR | | Protalus rampart |
| | | | RP | | Bedrock |
| | | | DF | | Debris flow deposit |
| | | | LS | | Solifluction lobe |
| | | Lithology ^b | TTP | | Sillimanite paragneiss (Tonale Unit) |
| | | | TUG | | Granate and cyanite paragneiss (Ultimo Unit) |
| | | | TUO | | Orthogneiss (Ultimo Unit) |
| | | | OME | | Chlorite e sericite micascists (Peio Unit) |
| | | | OMI | | Granate and staurolite micascists (Peio Unit) |
| | | | OOG | | Orthogneiss (Peio Unit) |
| | | | TPN | | Metapegmatites (Tonale Unit) |
| | | TTM | | Marbles (Tonale Unit) | |
| | | Vegetation cover ^c | 1 | | 0-10% covered by vegetation |
| | | | 2 | | 10-50% covered by vegetation |
| | | | 3 | | 50-90% covered by vegetation |
| | | | 4 | | 90-100% covered by vegetation |
| | | Permafrost evidence ^{a,c,h} | weqt | | winter equilibrium temperature measured by temperature data loggers |
| geophys | | | geophysical investigations (this work) | | |



| | | | | |
|---------------------|---------------------|--|---------------|--|
| | | | snow | perennial snowfields |
| | | | movement | surface displacement visible in multi-temporal DEMs |
| | | | none | no evidence available |
| | | APIM^d | Blue | permafrost in nearly all conditions |
| | | | Purple | permafrost mostly in cold conditions |
| | | | Yellow | permafrost only in very favorable conditions |
| | | | White | no permafrost |
| | | Open work deposit^{e,g} | y | present |
| | | | n | absent (includes boulder deposits with fine infill and/or widespread vegetation cover) |
| Rock glacier | Quantitative | Front slope (degrees)^a | \ | |
| | Qualitative | Activity^{f,g} | Intact | Intact rock glacier (active or inactive) |
| | | | Relict | Relict rock glacier |
| | | Length^a | Short | Short rock glacier length class (as defined in Sect. 3) |
| | | | Mid | Mean rock glacier length class (as defined in Sect. 3) |
| | | | Long | Long rock glacier length class (as defined in Sect. 3) |
| | | Elevation^a | Low | Low rock glacier elevation class (as defined in Sect. 3) |
| | | | Mid | Mean rock glacier elevation class (as defined in Sect. 3) |
| | | | High | High rock glacier elevation class (as defined in Sect. 3) |
| | | Vegetation cover^c | Vegetated | Vegetated rock glacier (as defined in Sect. 3, Table 1) |
| | | | Non vegetated | Non vegetated rock glacier (as defined in Sect. 3, Table 1) |
| | | Front characteristics^g | 1 | No vegetation, evidence of recent instability, outcrop of fine material, little or no surface weathering, weathering degree lower than the surface of the rock glacier |
| | | | 2 | Very little or no vegetation (<20%), very little or no fine material, weathering and lichen cover comparable to the surface of the rock glacier |
| | | | 3 | Scarce or discontinuous and cold-adapted vegetation (≤50%), abundant debris, weathering |



| | | | | |
|--|--|--|---|---|
| | | | | similar to the surface of the rock glacier, cold air draining from voids among blocks |
| | | | 4 | Completely vegetated, little outcropping debris, without voids and cold air drainage |
| | | Subdued topography ^{a,g} | y | The lateral and frontal ridges are clearly evident and the central part of the rock glacier is depressed with respect to them (concave contour lines) |
| | | | n | Lateral ridges are absent or evident only in the upper part of the rock glacier, from halfway down the morphology is convex or almost flat |

203

204 ^a Derived from the 2006 and 2014 LiDAR DEM of the Trento Province (siat.provincia.tn.it)

205 ^b Derived from the 1:10000 geological map of the Trento Province (protezionecivile.tn.it)

206 ^c Derived from the 2014 orthophoto of the Trento Province (siat.provincia.tn.it)

207 ^d Derived from the Boeckli et al. (2012) Alpine Permafrost Index Map

208 ^e Derived from the hillshaded 2014 LiDAR DEM of the Trento Province (siat.provincia.tn.it)

209 ^f Derived from the Seppi et al. (2012) rock glacier inventory

210 ^g Derived from field observations

211 ^h Ground surface temperature data reported in Carturan et al., (2016) and references therein;

212 <http://www.protezionecivile.tn.it/>.

213

214 We investigated the possible relationship of each variable with the spring-water temperature by means of scatterplots,
 215 boxplots, analysis of variance (or Kruskal-Wallis one way analysis of variance on ranks when variances were not
 216 homogeneous), Dunn’s multiple comparison test, Student’s t-test, and regression analysis. We defined spring-water
 217 temperature as “the median of all available temperature measurements in the four years”, so that we smoothed the
 218 interannual variability of water temperature. However, we had also to account for the different number of measurements
 219 available for each spring (from one up to four), and in particular for the possible low representativeness of springs
 220 measured only once. In this case, there is the possibility of having measured an extreme value, far from the typical
 221 conditions of those springs. To evaluate the impact of extreme values, we computed the absolute difference between each
 222 single-year spring water measurement and the median of all available measurements at the same spring. The mean of
 223 these absolute differences was 0.12°C, the median was 0.05°C, whereas the minimum and the maximum were 0 and
 224 0.7°C, respectively, and 89% of values was below 0.3°C. These results indicate a low impact of extreme temperatures
 225 and the suitability of using the median of all available measurements (regardless of their number) in statistical analyses.
 226 For springs with temperature measured only once, we retained the single value if runoff was >0.1 l/s.

227


228 3.4 Geophysical investigations

229 Electrical resistivity tomography (ERT) surveys were performed on 13-14 July 2022 at two neighbouring rock glaciers,
 230 classified as relict in the inventory of Seppi et al. (2012). These rock glaciers were selected considering their different



231 characteristics (spring-water temperature, vegetation cover, elevation) and the easy access. The Preghena Rock Glacier
232 has a mean elevation of 2196 m a.s.l., is mainly free of vegetation (although shrubs and trees are present) and its spring-
233 water temperature ranged between 1.6 and 1.8°C throughout the late summer during the measuring period. The Bordolona
234 Rock Glacier has a mean elevation of 1967 m a.s.l., is completely covered by vegetation and its spring-water temperature
235 ranged between 3.5 and 3.7°C in the late summer during the measuring period. Both rock glaciers are northeast oriented
236 (Fig. 3).

237



238

239 Figure 3: Location of ERT lines (black solid lines) performed on the Bordolona and Preghena rock glaciers in July
240 2022. The white dots indicate the lower edge of rock glaciers. The green dots in the upper panels indicate the sampled
241 springs.

242



243 Geophysical surveys were carried out with a Syscal Pro georesistivimeter (Iris Instruments), using arrays of 72 (Line 1
244 Preghena and Line 3 Bordolona) or 48 (Line 2 Preghena) electrodes, with 3-meter electrodes spacing (Fig. 3). A dipole-
245 dipole scheme was used, with two different skips of 0 and 4 electrodes. This configuration ensured relatively high
246 resolution at the surface, and at the same time enough penetration depth. Measurements were carried out with a stack of
247 3 to 6, imposing an acceptable error threshold of 5%. To estimate a more reliable experimental error for the acquired
248 datasets (Binley, 2015), direct and reciprocal measurements were acquired by exchanging injecting and potential dipoles
249 for each quadrupole. To partially overcome the high contact resistances between the electrodes and boulders/debris
250 (Hauck and Kneisell, 2008), the electrodes were inserted between the boulders using sponges soaked with saltwater
251 (Pavoni et al., 2023). Nevertheless, at the blocky surface of the Preghena Rock Glacier the contact resistances remained
252 steadily above 100 k Ω m, due to dry environmental conditions. The organic soil at the Bordolona Rock Glacier guaranteed
253 low contact resistances (<10 k Ω m).

254 The inversion process of the acquired datasets has been performed with the Python-based software ResIPy (Blanchy et
255 al., 2020), based on the Occam's inversion method (Binley and Kemna, 2005). In each dataset, quadrupoles with a
256 stacking error higher than 5% were removed, and the expected data error was defined using the reciprocal check (Day-
257 Lewis et al., 2008, Pavoni et al., 2023), giving values of 20% and 5% for the Preghena and Bordolona Rock Glacier,
258 respectively. The acquired data were of lower quality at the Preghena Rock Glacier, due to the high contact resistance.

259

260 **4. Results**

261 **4.1 Spatial variability of spring-water temperature**

262 Water temperature of the 131 springs ranged between 0.0 and 8.5 °C, with a mean of 3.6 °C and a median of 3.4°C (Table
263 4). The frequency distribution of the spring elevation (i.e., the minimum elevation of catchments) is symmetrical and
264 normally distributed around a sample mean of 2384 m a.s.l. The lowermost spring was sampled at 1698 m a.s.l., and the
265 uppermost spring was sampled at 3039 m a.s.l.

266 The mean elevation of spring catchments varies between 2104 and 3151 m a.s.l., whereas the maximum elevation ranges
267 between 2241 and 3352 m a.s.l. The mean and maximum elevation average 2539 and 2694 m a.s.l., respectively. Both
268 are also symmetrical around the sample mean and normally distributed.

269 The planimetric length of spring catchments varies between 83 and 2621 m, with a mean of 610 m. The skewness and
270 kurtosis indicate that the planimetric length is right skewed and leptokurtic.

271

272

273

274

275

276




277 Table 4: Descriptive statistics for spring-water temperature measurements and quantitative variables relative to spring
 278 catchments (as defined in Table 3).

| N = 131 | Median temperature (T_{Mdn}) | Catchment minimum elevation (m) | Catchment maximum elevation (m) | Catchment mean elevation (m) | Catchment planimetric length (m) |
|---|--|--|--|---|---|
| Minimum | 0.0 | 1698 | 2241 | 2104 | 83 |
| Median | 3.4 | 2367 | 2641 | 2495 | 539 |
| Maximum | 8.5 | 3039 | 3352 | 3151 | 2621 |
| Range | 8.5 | 1341 | 1111 | 1047 | 2538 |
| Mean | 3.6 | 2384 | 2694 | 2539 | 610 |
| Standard error of the mean | 0.2 | 22.6 | 21.9 | 21.0 | 34.9 |
| Standard deviation | 1.8 | 259.2 | 251.1 | 240.8 | 399.3 |
| Coefficient of variation | 0.500 | 0.109 | 0.093 | 0.095 | 0.655 |
| Skewness | 0.392 | 0.179 | 0.446 | 0.419 | 2.070 |
| Kurtosis | -0.261 | -0.328 | -0.107 | -0.391 | 6.095 |

279

280

281 Spring-water temperature is significantly correlated with the mean elevation of the catchments (Fig. 4a) for all three
 282 aspect classes defined in Table 3. Linear regressions are significant ($p < 0.001$) for south ($R^2 = 0.30$) and for east-west
 283 facing catchments ($R^2 = 0.35$). For the north facing catchments, there is a low significant relation ($R^2 = 0.25$, $p < 0.05$)
 284 between water temperature and elevation. In all three cases, the low R^2 suggests that other factors should affect water
 285 temperature, as well. Similar results were obtained using spring elevation rather than mean catchment elevation (Fig. 5).



286


287 Figure 4: Relationship between spring-water temperature and a) mean catchment elevation (clustered in three classes of
 288 mean catchment aspect), b) mean catchment aspect, c) mean catchment elevation (clustered in seven classes of upslope
 289 area geomorphology), and d) upslope area geomorphology. Acronyms and their meanings are reported in Table 3. Boxes
 290 in b) and d) indicate the 25th and 75th percentile, whiskers indicate the 10th and 90th percentile, whereas the horizontal
 291 solid and dashed lines within the box mark the median and the mean, respectively. Different letters above the boxplots
 292 indicate groups with significantly different ($p < 0.05$) water temperatures based on Dunn's multiple comparison test
 293 (applied after the Kruskal-Wallis test).

294

295

296

297



298

299 Figure 5: Relationship between spring-water temperature and minimum (spring) elevation, clustered in three classes of
300 mean catchment aspect, as in Fig. 4a.

301 As expected, there is a negative relationship between water temperature and elevation (Fig. 4a and 5), but also a large
302 overlap of water temperature among the three aspect classes. NW-NE facing catchments have significantly colder springs
303 compared to SE-SW facing catchments ($p < 0.05$, Dunn's multiple comparison test, applied after the Kruskal-Wallis test),
304 whereas catchments facing NE-SE and SW-NW have water temperature that do not differ significantly from the other
305 two classes (Fig. 4b). NW-NE facing catchments show a lower variability in spring-water temperature compared to the
306 other two classes.

307 Figures 4c and 4d highlight that springs with upslope areas dominated by the presence of rock glaciers (irrespective of
308 their activity) and bedrock outcrops are significantly colder than other springs ($p < 0.05$, Dunn's multiple comparison test,
309 applied after the Kruskal-Wallis test).


310

311 4.2 Temperature of springs downslope of rock glaciers

312 4.2.1 Comparison between intact and relict rock glaciers

313 The spring-water temperature is significantly different for rock glaciers with different degrees of activity (Fig. 6a). Relict
314 rock glaciers have a much warmer spring temperature compared to intact rock glaciers (Student's t-test, $p < 0.001$), and
315 the variability of water temperature is larger for relict rock glaciers. There is a substantial overlap between the two groups,
316 which extended between 1.2 and 3°C. This range of water temperature represents 54% of all springs downslope of rock
317 glaciers (53% of intact rock glaciers and 54% of relict rock glaciers). Almost half of rock glaciers classified as relict has
318 spring-water temperature similar to rock glaciers classified as intact.

319 The two groups of rock glaciers have significantly different minimum elevations (Fig. 6b, Student's t-test, $p < 0.001$), but
320 there is a wide elevation band, comprised between 2406 and 2630 m a.s.l., where they overlap.



321

322 Figure 6: Spring-water temperature (a) and minimum elevation (b) of rock glaciers sampled in the study area. Boxes
 323 indicate the 25th and 75th percentile, whiskers indicate the 10th and 90th percentile, whereas the horizontal solid and dashed
 324 lines within the box mark the median and the mean, respectively. Maximum and minimum values are represented by dots.
 325 Sample size (n) is reported in a).

326

327 4.2.2 Spring-water temperature of relict rock glaciers

328 The relationship between water temperature and the mean catchment elevation is rather weak for springs fed by relict
 329 rock glaciers (Fig. 7a). The linear regression is significant ($p < 0.05$) only for catchments facing NE-SE and SW-NW, but
 330 the relation is weak ($R^2 = 0.20$). At the same elevation, catchments facing NW-NE have colder springs compared to the
 331 other two aspect classes. The spring-water temperature of catchments facing north is similar to that of catchments facing
 332 east, south and west located 300-400 m above.


333 Relict rock glacier springs with open work deposits in their upslope areas are colder than springs without open work
 334 deposits (Fig. 7b). For the first group, the water temperature is not related to the mean catchment elevation, whereas for
 335 the second group there is a weak but significant relation ($p < 0.05$, $R^2 = 0.15$). Consequently, the difference in water
 336 temperature of the two groups increase towards low elevations, which suggest that open work deposits may have a cooling
 337 effect particularly marked at elevations < 2500 m a.s.l.

338 Similar considerations can be done for rock glacier front characteristics (Fig. 7c) and for rock glacier vegetation cover
 339 (Fig. 7d). Relict rock glaciers with scarce and cold-adapted vegetation cover have colder springs compared to relict rock
 340 glaciers with abundant vegetation cover on their bodies and fronts. However, for all classes of rock-glacier front
 341 characteristics and vegetation cover (Table 3) there is no significant relation between water temperature and mean
 342 catchment elevation.

343 Despite the large overlap among the analysed classes (Fig. 7), we found a significant effect of vegetation cover (Student's
 344 t-test, $p < 0.001$), open work deposits (Student's t-test, $p < 0.001$) and front characteristics (Student's t-test applied to classes
 345 III and IV, $p < 0.01$) on the water temperature of springs downslope of relict rock glaciers. We did not detect any significant
 346 influence of the mean aspect of the catchment, the mean elevation of rock glaciers, their length, and the presence or
 347 absence of a subdued topography on water temperature.



348



349

350 Figure 7: Relationship between spring-water temperature of relict rock glaciers and mean catchment elevation clustered
 351 in a) three classes of mean catchment aspect, b) two classes of open work deposits in the spring upslope area, c) three
 352 classes of rock glacier front characteristics, and d) two classes of rock glacier vegetation cover. Classes are described in
 353 Table 3.

354

355 4.3 Geophysical investigations


356 Figures 8a and b show the inverted resistivity sections obtained for the investigation Lines 1 and 2 acquired on the
 357 Preghena Rock Glacier. High values of resistivity (>80 kΩm) were found in the uppermost layer, down to about 7-8
 358 meters of depth, associated to the dry conditions during ERT soundings and to the air-filled voids among coarse debris
 359 and blocks, typical of rock glacier environments. Below this uppermost layer, the resistivity values rapidly decrease (<10
 360 kΩm) indicating a plausible decrease of porosity and grain size in the deposit, and a possible increase in water content.
 361 This low resistivity layer develops almost continuously down to the bottom of the models. An increase in resistivity is
 362 found at the lower end of line 1 and at the upper end of line 2, in the area where they overlap and at a depth of about 12-



363 13 m, reaching 150-200 k Ω m. This area of increased resistivity can be interpreted as a deep frozen body, providing
364 evidence of probable permafrost inside this rock glacier.

365 Figure 8c shows the inverted resistivity section obtained for the investigation Line 3 acquired on the Bordolona Rock
366 Glacier. In the shallowest layers the resistivity is comprised between 5 and 10 k Ω m, significantly lower than the shallow
367 layer of the Preghena Rock Glacier, even if air-filled voids are common on this rock glacier as well.

368 Below this layer, a sharp increase in resistivity is detected along the entire investigation line, with frequent regions
369 exceeding 20 k Ω m. The highest resistivity (about 60 k Ω m) is found towards the upper end of the ERT line, where a
370 younger rock glacier lobe overlies the main body. This high resistivity layer reaches about 15 meters of depth and can be
371 interpreted as a frozen layer. The bottom of the high-resistivity layer, which seems discontinuous in the lower part and
372 more continuous and thicker in the upper part of the ERT line, is highlighted by a strong decrease in resistivity, below 5
373 k Ω m. This lowermost layer is probably unfrozen and is characterised by an increase in water content and fine sediments.



374

375 Figure 8: Inverted resistivity section of the investigation Line 1 (a) and 2 (b) on the Preghena Rock Glacier, and of the

376 investigation line 3 (c) on the Bordolona Rock Glacier.

377

378



379 **5. Discussion**

380 **5.1 Permafrost distribution and spring-water temperature in the study area**

381 Measurements of spring-water temperature collected in this study outside the rock-glacier influence have a high spatial
382 variability and do not show a significant relationship with elevation ($p > 0.05$). Among springs outside the rock glacier
383 influence, only those above 2800 m a.s.l. have a water temperature $\leq 2.2^\circ\text{C}$, which is the upper limit reported in the
384 literature for 'possible permafrost' (Carturan et al., 2016).

385 This result lines up well with mean annual air temperature (MAAT) indications. Indeed, based on the MAAT of -0.9°C
386 measured between 1961 and 2010 at the Careser Diga weather station (2605 m a.s.l., in the northern part of the Val di
387 Sole), the theoretical lower limit of discontinuous permafrost in Val di Sole, corresponding to a MAAT of -2°C (Haerberli,
388 1985), should be comprised between 2700 and 2800 m a.s.l..

389 Similarly, the alpine permafrost index map (APIM, Boeckli et al., 2012) indicates a lower limit of "permafrost mostly in
390 cold conditions" ranging between 2500 and 2900 m outside rock glaciers and coarse-block deposits, varying upon terrain
391 aspect and averaging 2700 m a.s.l.. Based on the mean elevation of intact rock glaciers in the study area, Seppi et al.
392 (2012) calculated a present-day lower limit of permafrost at 2720 m a.s.l..

393 As expected, springs draining north-facing catchments are significantly colder compared to springs draining south-facing
394 catchments. On average, there is a difference of about 3°C between springs draining catchments at similar elevation and
395 with opposite aspect. On average, the same spring temperature is found 500-600 m higher on south-facing catchments
396 than on north-facing ones (Fig. 5). This result quantifies the influence of terrain exposure on the ground temperature
397 regime and permafrost distribution in the study area, which are direct consequences of shortwave radiation inputs and
398 related effects on snow cover and surface albedo (Boeckli et al., 2012).

399 In our study, at all elevations, springs draining rock glaciers are the coldest, irrespective of the rock glacier activity state
400 (Fig. 4c). This is in agreement with findings of studies in the European Alps and in other mountain chains reporting rock-
401 glacier spring-water temperatures, regardless of their activity state. For example, in the Canadian Rockies, spring-water
402 temperature from an inactive rock glacier hosting small portions of permafrost reached a maximum of 2.2°C , exercising
403 a substantial cooling effect on the creek downstream (Harrington et al., 2018). Interestingly, cold conditions and high
404 daily variability in spring-water temperature in summertime has been recorded in a rock glacier in Norway that shows
405 characteristics favourable to the presence of permafrost, but with minor ice bodies (Lilleøren et al., 2022). In the Austrian
406 Alps, spring-water from a relict rock glacier was monitored for 6 years, showing a mean temperature of 2.2°C , with small
407 seasonal variation (between 1.9 and 2.5°C) and a decrease of the water temperature after precipitation events, attributed
408 to the potential presence of ice lenses in the lower part of the rock glacier (Winkler et al., 2016).

409 Our results align as well with those of studies reconstructing permafrost distribution by empirical modelling in the Alps
410 and at other mountain locations worldwide. A logistic regression model used in the Dry Andes of Argentina accounting
411 for mean annual air temperature, terrain ruggedness, and potential incoming solar radiation suggests that permafrost may
412 occur in several types of coarse blocky deposits, including rock glaciers, even under unfavourable climatic conditions
413 (Tapia Baldis and Trombotto-Liaudat, 2020). A similar empirical-statistical model applied in the Austrian Alps shows
414 that permafrost can be expected above 2500 m a.s.l. in northerly exposed slopes and above 3000 m a.s.l. in southerly
415 exposed slopes (Schrott et al., 2012), providing an elevation difference of about 500 m between south and north exposures,
416 which agrees well with our spring-water temperature results.




417 **5.2 Rock glacier classification based on spring-water temperature**

418 Although springs draining intact rock glaciers are significantly colder than springs draining relict rock glaciers, there is a
419 remarkable ~50% overlap in the water temperature range of the two rock glacier groups (Fig. 6a). Based on published
420 thresholds (Haerberli, 1975; Frauenfelder et al., 1998; Scapozza, 2009; Carturan et al., 2016), 12 out of the 52 relict rock
421 glaciers sampled in Val di Sole (23%) can be included in the ‘possible permafrost’ category (water temperature between
422 1 ± 0.2 and 2 ± 0.2 °C), and none of them in the ‘probable permafrost’ category (water temperature $< 1\pm 0.2$ °C). However,
423 the relatively warm water temperature measured downstream of intact rock glaciers (maximum = 3°C, 90th percentile =
424 2.4°C), and downstream of areas with permafrost evidence (maximum = 3.5°C, 90th percentile = 2.2°C), suggest that the
425 upper limit for possible permafrost may be higher. Here, the 90th percentile accounts for possible misclassification of
426 intact rock glaciers and other issues affecting spring-water temperature measurements (Sect. 5.3).

427 Assuming a (rounded) upper limit of 2.5°C for spring-water temperature with possible permafrost influence leads to
428 include 19 (38%) relict rock glaciers in the possible permafrost category. This estimate looks more conservative than the
429 ~50% obtained by a mere comparison of water temperature ranges of intact and relict rock glaciers (Fig. 6a). These
430 findings might suggest that permafrost in rock glaciers classified as relict is widespread in Val di Sole, and that a large
431 fraction of them is actually pseudo-relict, or transitional landforms, containing patches of permafrost and reaching an
432 elevation below the tree line (2000-2200 m a.s.l.).

433 Examples of spring-water temperature downstream of rock glaciers in Val di Sole are shown in Fig. 9. Cold springs
434 draining rock glaciers classified as relict are associated to the presence of open work deposits and scarce vegetation cover
435 (Fig. 7 and 9). These two explaining variables are often correlated, because vegetation tend to be scarce over coarse
436 deposits without fine infill among blocks, and vice versa. The relationship between cold spring temperature (as permafrost
437 evidence) and these two surface characteristics was expected in our case study, based on the existing literature (e.g.,
438 Guglielmin, 1997, and references therein). This relationship is statistically significant only for rock glaciers classified as
439 relict, whereas for intact rock glaciers sampled in the study area it does not exist (Fig. A1).



440

441 Figure 9: Examples of spring-water temperature downstream of rock glaciers in Val di Sole: a) intact (active) rock glacier
442 with cold spring at 2950 m a.s.l.; b) intact (inactive) rock glacier with relatively warm spring at 2727 m a.s.l.; c) relict
443 rock glacier with cold spring at 2304 m a.s.l., whose surface is open-work and presents scarce vegetation cover; d) relict
444 rock glacier with warm spring at 2266 m a.s.l., whose surface is entirely covered by vegetation.

445 The long-term preservation of permafrost within open work blocky deposits results from overcooling and thermal
446 decoupling of the frozen core from the external climate (Harris and Pedersen, 1998; Morard et al., 2008; Jones et al.,
447 2019). The low thermal conductivity in coarse open work deposits brings to lower ground temperatures compared to fine-
448 grain material (Juliussen and Humlum, 2008; Jones et al., 2019). Soil development over the surficial blocks and boulders
449 can prevent these cooling effects (Ikeda and Matsuoka, 2002). However, if fine-grain infilling does not occur, ground
450 cooling effect goes undisturbed. In central Europe, these processes enable the existence of permafrost much below its
451 regional limit and reaching elevations lower than 1000 m a.s.l. (Gude et al., 2003; Delaloye et al., 2003). According to
452 Delaloye and Lambiel (2005), thousand-year-old permafrost might be potentially preserved in these types of deposits.

453 Open work deposits and/or scarce vegetation cover can be potentially employed to distinguish rock glaciers with or
454 without permafrost, as both can be mapped based on remote-sensing imagery. However, open work deposits and
455 vegetation cover do not enable a full distinction of ‘cold’ and ‘warm’ springs affected by relict rock glaciers (Fig. 7b, c
456 and d). Individual non-open-work rock glaciers widely covered by vegetation can have spring-water temperature as low
457 as 1.4°, and rock glaciers almost free of vegetation with blocky surface can have spring-water temperature up to 3.9°C.

458 Other variables considered in this study, such as aspect, elevation, size and the presence or absence of a subdued
459 topography on rock glaciers (Delaloye et al., 2003; Delaloye, 2004), are not related to spring-water temperature. Figure
460 7 suggests the existence of a group of cold springs at low elevations on north-facing catchments, even though water



461 temperature is not significantly different from the temperatures of springs in the other two aspect classes. This result
462 might be due to the small sample size of the NW-NE aspect class.

463

464 **5.3 Limitations and uncertainties in the spring-water temperature approach**

465 The results of this study might be affected by limitations in the experimental design, assumptions, and uncertainties. First,
466 the main assumption of this study is that spring-water temperature provides indication of permafrost occurrence at
467 investigated rock glaciers and spring upslope areas, and can be used as a stand-alone pilot method to rapidly explore the
468 activity state of rock glaciers in a wide area. This approach applies spring-water temperature to the catchment scale,
469 beyond its general use as an ancillary method to other techniques such as InSAR analyses, ground surface temperature
470 measurements and/or geophysics.

471 We base our assumption on previously published work and well-known temperature thresholds for permafrost probability
472 categories (e.g., Haeberli, 1975; Frauenfelder et al., 1998; Scapozza, 2009) and on our first successful application at the
473 catchment scale (Carturan et al., 2016). Data collected in Val di Sole are in line with literature thresholds, provided that
474 the 10% largest spring-water temperature values are excluded (Sect. 5.2). Including these extreme values leads to about
475 1.5°C larger temperature thresholds for possible permafrost compared to literature.

476 The reason behind this discrepancy lies in the uncertainty in the classification of rock glacier activity, which was based
477 on vegetation and geomorphological characteristics, assessed mainly from remote-sensing images (Seppi et al., 2012). In
478 the wide elevation band where intact and relict rock glaciers coexist (minimum elevation between 2406 and 2630 m),
479 landforms with similar vegetation cover and surface geomorphology have been classified based on the authors' experience
480 and judgement, implying a certain degree of subjectivity.

481 The distinction between intact and relict rock glaciers is a theoretical concept, and there is a continuum between (true)
482 intact and (true) relict rock glaciers, with the existence of transitional landforms (Kääb, 2013). In absence of other
483 evidence, this continuum hampers to distinguish unambiguously intact and relict landforms, in particular if they have
484 similar surface characteristics. In addition, the mentioned transition is a dynamic concept, which depends on the
485 characteristics of individual landforms, their topo-climatic setting, and their response to climatic variations (Kääb, 2013).

486 Another source of uncertainty is related to the distance between the permafrost body and the measured springs. Water
487 temperature is a non-conservative tracer, and if the main permafrost body is distant (e.g., tens of meters) from the rock
488 glacier front, water temperature can significantly increase along the flow paths before reaching the spring, due to the
489 contact with unfrozen sediments and/or mixing with other water sources (e.g., Kellerer-Pirklbauer et al., 2017). This is
490 the case of the Bordolona Rock Glacier (Fig. 8c), where the rather warm spring-water temperature (3.5-3.7°C) would
491 have led to exclude the occurrence of permafrost in absence of geophysical evidence.

492 Several authors are cautious when discussing about cold springs downslope of relict rock glaciers. For example, Winkler
493 et al. (2016) do not exclude the presence of remaining ice lenses inside the relict Schöneben Rock Glacier (Niedere Tauern
494 Range, Austria), as a possible explanation for the rapid cooling of the spring water after recharge events, during
495 summertime. However, the authors mention the cold thermal regime beneath coarse blocky materials as a possible
496 explanation, which does not necessarily imply permafrost occurrence, and conclude that additional research is required
497 for the identification of the cooling source.

498 We agree that additional research is required to confirm inference from spring-water temperature. With this study we add
499 that spring-water temperature can be as high as 1.8°C for rock glaciers where permafrost occurrence is confirmed by



500 geophysics or ground surface temperature measurements, and can exceed 3.5°C where the permafrost body is far from
501 the rock glacier front and spring, such as at the Bordolona Rock Glacier. Even if the collected data seem to suggest that
502 temperature thresholds might be slightly higher than those reported in the literature, further investigations are necessary
503 for better constraining them and for defining their range of uncertainty.

504 A last source of uncertainty is represented by the sampling design adopted for Val di Sole, with its particular topographic
505 and geological characteristics. The dominant southward aspect of the investigated rock glaciers, and their spatial
506 clustering, can explain the lack of correlation between water temperature and the aspect of rock glaciers. We tried to
507 minimise the spatial clustering of measured springs, visiting as many headwater catchments as possible, and taking
508 measurements at the largest number of springs on each catchment. However, due to logistic constraints and inherent
509 characteristics of the study area, a certain degree of spatial clustering was unavoidable. For this reason, the role of terrain
510 aspect as a possible controlling factor on spring-water temperature requires additional investigation.

511

512 **5.4 Geophysics**

513 The inverted resistivity sections obtained for the Preghena Rock Glacier (Fig. 8a and b) show results compatible with the
514 presence of permafrost patches. Even considering the high contact resistance due to the dry weather conditions preceding
515 the survey, and the location of the high resistivity body in the areas known to be the least sensitive of the model (the bed
516 and margins, Binley, 2015), we observe that the obtained resistivity values are typical of frozen materials (Hauck and
517 Kneisel, 2008). This result agrees with the low temperature of the Preghena Rock Glacier spring, which fluctuates between
518 1.6 and 1.8°C throughout summer, and it suggests that this rock glacier should be classified as a pseudo-relict rock glacier.

519 In the Bordolona Rock Glacier (Fig. 8c), the frozen layer looks discontinuous in the lower section of the ERT Line, and
520 more continuous and thicker in the upper part, where a younger lobe superposes the main body of the rock glacier. The
521 different resistivity detected in the lower and upper sections of the ERT line can be related to a different percent ice
522 content in the frozen layers, and/or a different temperature of the ice (Hilbich et al., 2008). These results suggest the
523 probable presence of permafrost also inside the Bordolona Rock Glacier, which was considered a ‘true’ relict rock glacier
524 due to its abundant vegetation cover, spring-water temperature above 3°C, and low mean elevation. Based on geophysical
525 investigations, the Bordolona Rock Glacier too should be classified as a pseudo-relict rock glacier.

526 More conclusive results should be obtained by repeating the geophysical surveys under moister conditions, especially at
527 the Preghena Rock Glacier, and possibly coupling ERT to seismic refraction measurements in order to obtain a reliable
528 estimate of the percent ice content inside these rock glaciers (Hauck et al., 2011, Wagner et al., 2019, Pavoni et al. 2023).

529

530 **5.5 Ice storage in the rock glaciers and glaciers of Val di Sole**

531 Given the very different response time of glaciers and rock glaciers to projected atmospheric warming, their relative
532 importance has relevant implications for the current and future hydrological cycle. In the study area, glaciers are shrinking
533 fast, and in 2022 they covered 7.18 km², which is 59% of the 2003 area (area loss rate = 2.6% y⁻¹). For these reasons, and
534 in light of the results of this work, it is interesting to estimate and compare the ice content of rock glaciers and glaciers in
535 Val di Sole, similarly to what was done, for example, by Bolch and Marchenko (2009) in the Northern Tien Shan.



536 For glaciers, we estimated residual volumes in 2022 starting from the ice thickness estimates provided for each glacier in
537 the study area by Farinotti et al., (2019). We first calculated the bedrock topography subtracting the ice thickness from
538 the glacier surface DEM (Farinotti et al., 2019). Then we calculated the 2002 glacier thickness subtracting the bedrock
539 topography from a glacier surface DEM surveyed in September 2022 by the Province of Trento. We finally obtained the
540 glacier volumes multiplying the average thickness by the glacier area, and converted the ice volume into the water volume
541 equivalent using a mean ice density of 900 kg m^{-3} .

542 For rock glaciers, we calculated the total rock glacier volume multiplying their area A by the average thickness provided
543 by the Brenning (2005b) formulation:

$$544 \quad T = cA^\gamma \quad (1)$$

545 where T is the average thickness of rock glaciers, and c and γ are constants equal to 50 and 0.2, respectively. To account
546 for the different geometry of intact and relict rock glaciers, we assumed that the volumetric ice content of intact rock
547 glaciers averages 50% (Jones et al., 2018, and references therein), and therefore that T_r for (true) relict rock glaciers is
548 half that of intact rock glaciers (i.e., they are composed only of debris and all the ice melted away). For pseudo-relict rock
549 glaciers we tested various hypotheses of percent ice content, calculating the average thickness T_{pr} as follows:

$$550 \quad T_{pr} = T_r + T_{ice} \quad (2)$$

551 where T_{ice} is the average ice thickness, calculated in function of the volumetric percent ice content $\%_{ice}$ as:

$$552 \quad T_{ice} = \frac{\%_{ice} \cdot T_r}{(1 - \%_{ice})} \quad (3)$$

553 A total glacier ice volume of $251 \times 10^6 \text{ m}^3$, and a corresponding $226 \times 10^6 \text{ m}^3$ water volume equivalent was calculated for
554 Val di Sole in 2022. In comparison, the water volume equivalent of intact rock glaciers is $42.7 \times 10^6 \text{ m}^3$.

555 A water volume equivalent between 3.7 and $17.7 \times 10^6 \text{ m}^3$, averaging $10.7 \times 10^6 \text{ m}^3$, can be estimated assuming that 38%
556 of the total area of relict rock glaciers contains permafrost, and that the average ice content ranges between 5% and 20%
557 in volume. This range is a first hypothesis based on the few geophysical data available at pseudo-relict rock glaciers
558 (Delaloye, 2004; Colucci et al., 2019; Pavoni et al., 2023; this work). To our knowledge, the amount of ice in pseudo-
559 relict rock glaciers has yet to be quantified.

560 Even if preliminary and affected by significant uncertainty, these estimates provide an order of magnitude of water stored
561 as ice in the rock glaciers of Val di Sole. The water equivalent ratio for rock glacier ice versus glacier ice averages 1:4.2
562 and ranges between 1:3.7 and 1:4.9, considering minimum and maximum estimates reported above. Importantly, based
563 on these calculations, 20% of the total rock glacier water volume would be stored inside pseudo-relict rock glaciers. Even
564 assuming the lower bound of percent ice content (5%), pseudo-relict rock glaciers would contribute to a significant 8%
565 of the total rock glacier water volume.

566 **6. Concluding remarks**

567 We have surveyed spring-water temperature in an area of 795 km^2 in Val di Sole, to understand the influence of
568 topographic and geomorphological factors, and to test if it can be used to preliminarily differentiate intact and relict rock
569 glaciers. Spring-water temperature measurements enabled to characterise a large number of rock glaciers, and to provide
570 a first estimate of the frequency of pseudo-relict rock glaciers in this area. Overall, our results point to a significant



571 hydrological importance of rock glaciers classified as relict in the study area, which is expected to increase in the future
572 due to atmospheric warming.

573 In general, we have found that the spatial variability of spring-water temperature is controlled by elevation, aspect and
574 the presence of rock glaciers in the upslope area. Compared to other landforms in the upslope area, rock glaciers have
575 colder springs, irrespective of their activity state.

576 The spring-water temperature of rock glaciers classified so far as relict is higher and with larger spatial variability
577 compared to intact rock glaciers. However, there is a remarkable ~50% (38% excluding extremes) overlap in the spring
578 temperature range of the two rock glacier groups. Relict rock glaciers tend to have colder springs if their surface is blocky
579 and scarcely covered by (cold-adapted) vegetation.

580 The spring-water temperature data suggest that one third of rock glaciers classified as relict might be actually pseudo-
581 relict, thus containing permafrost. The exact percentage cannot be derived unambiguously from spring-water temperature
582 because i) other evidence is required to confirm inference from water temperature, ii) there is uncertainty in the
583 classification of the activity state of rock glaciers, iii) there is geophysical evidence that rock glaciers containing
584 permafrost may have 'warm' springs (up to 3.7°C), and consequently iv) there is uncertainty in the definition of the
585 thresholds for differentiate among absent/possible/probable permafrost categories. Despite these uncertainties, our study
586 shows that rock-glacier spring-water temperature can provide a pilot approach to estimate the spatial distribution of
587 permafrost in vast areas, and an auxiliary element to the classification of rock glaciers, whose permafrost content might
588 otherwise go underestimated.

589 Geophysics applied to two rock glaciers classified as relict enabled to detect the presence of permafrost. While the blocky
590 Preghena Rock Glacier, whose spring temperature was < 1.8°C throughout the summer, was expected to contain
591 permafrost, its occurrence in the Bordolona Rock Glacier was not expected, because it is entirely covered by dense
592 vegetation and its spring temperature reached 3.7°C in late summer.

593 Preliminary calculations of water resources stored as ice inside the rock glaciers of Val di Sole reveal that they amount to
594 ~24% of the water volume equivalent stored in glaciers, which are disappearing very fast. Remarkably, 20% of the total
595 rock glacier water volume is stored inside rock glaciers classified as relict.

596 This study highlights the need for additional investigations and improved understanding of these periglacial landforms.
597 In particular, the possible presence of permafrost in a large fraction of rock glaciers classified as relict poses critical
598 questions regarding the origin, preservation, current behaviour, seasonal dynamics, and future evolution of this
599 permafrost. Thorough study of pseudo-relict rock glaciers is required for understanding the transition between intact and
600 relict landforms, which is important in view of current and projected climate change.

601

602

603


604

605



606 **Appendix A**

607



608

609 Figure A1: Spring-water temperature for intact rock glaciers with and without open work deposits on their surface. Boxes
610 indicate the 25th and 75th percentile, whiskers indicate the 10th and 90th percentile, whereas the horizontal solid and dashed
611 lines within the box mark the median and the mean, respectively. Maximum and minimum values are represented by dots.
612 Sample size (n) is reported above the boxplots.

613

614

615 **Data availability**

616 Data are available from the corresponding author upon reasonable request.

617

618 **Author contributions**

619 LC designed the methodological approach and carried out the sampling campaigns with the support of AA, RS, MT, TZ
620 and GZ. MP and JB carried out the geophysical surveys in cooperation with LC, CM and MZ and interpreted the results.
621 GZ, LC and AA performed the statistical analyses of the dataset. LC prepared the first draft of the manuscript with
622 contributions from GZ, MP and CM. All authors contributed to the editing of the manuscript.

623

624 **Competing interests**

625 The contact author has declared that none of the authors has any competing interests.

626

627 **Acknowledgments**

628 The authors acknowledge the editor and reviewers for their comments and suggestions.



629

630 **Financial support**

631 This study was carried out within the RETURN Extended Partnership and received funding from the European Union
632 Next-GenerationEU (National Recovery and Resilience Plan – NRRP, Mission 4, Component 2, Investment 1.3 – D.D.
633 1243 2/8/2022, PE0000005). LC, GZ, JB and RS also acknowledge the support of the project PRIN 2022
634 “SUBSURFICE – Ecohydrological and environmental significance of subsurface ice in alpine catchments” (code:
635 2022AL7WKC) (National Recovery and Resilience Plan – NRRP, Mission 4, Component 2, Investment 1.1 – D. D. 104
636 2/2/2022).

637

638 **References**

- 639 Barsch, D.: Rockglaciers: indicators for the present and former geocology in high mountain environments, Springer
640 Berlin Heidelberg, Berlin, Heidelberg, 218 pp., <https://doi.org/10.2307/3060377>, 1996.
- 641 Bertone, A., Barboux, C., Bodin, X., Bolch, T., Brardinoni, F., Caduff, R., Christiansen, H. H., Darrow, M. M., Delaloye,
642 R., Etzelmüller, B., Humlum, O., Lambiel, C., Lilleøren, K. S., Mair, V., Pellegrinon, G., Rouyet, L., Ruiz, L., and Strozzi,
643 T.: Incorporating InSAR kinematics into rock glacier inventories: insights from 11 regions worldwide, *The Cryosphere*,
644 16, 2769–2792, <https://doi.org/10.5194/tc-16-2769-2022>, 2022.
- 645 Binley, A.: Tools and Techniques: Electrical Methods, in: *Treatise on Geophysics: Second Edition*, vol. 11, Elsevier,
646 233–259, <https://doi.org/10.1016/B978-0-444-53802-4.00192-5>, 2015.
- 647 Binley, A. and Kemna, A.: DC Resistivity and Induced Polarization Methods, in: *Hydrogeophysics*, Springer Netherlands,
648 Dordrecht, 129–156, https://doi.org/10.1007/1-4020-3102-5_5, 2005.
- 649 Blanchy, G., Saneiyani, S., Boyd, J., McLachlan, P., and Binley, A.: ResIPy, an intuitive open source software for complex
650 geoelectrical inversion/modeling, *Comput. Geosci.*, 137, 104423, <https://doi.org/10.1016/j.cageo.2020.104423>, 2020.
- 651 Boeckli, L., Brenning, A., Gruber, S., and Noetzi, J.: Permafrost distribution in the European Alps: Calculation and
652 evaluation of an index map and summary statistics, *Cryosphere*, 6, 807–820, <https://doi.org/10.5194/tc-6-807-2012>, 2012.
- 653 Bolch, T. and Marchenko, S.: Significance of glaciers, rockglaciers and ice-rich permafrost in the Northern Tien Shan as
654 water towers under climate change conditions. In: Braun, Ludwig N; Hagg, Wilfried; Severskiy, Igor V; Young, Gordon.
655 *Assessment of Snow, Glacier and Water Resources in Asia: Selected papers from the Workshop in Almaty, Kazakhstan*,
656 2006. Koblenz: IHP UNESCO, 132-144, 2009.
- 657 Bollati, I. M., Cerrato, R., Lenz, B. C., Vezzola, L., Giaccone, E., Viani, C., Zanoner, T., Azzoni, R. S., Masseroli, A.,
658 Pellegrini, M., Scapozza, C., Zerboni, A., and Guglielmin, M.: Geomorphological map of the Val Viola Pass (Italy-
659 Switzerland), *Geogr. Fis. e Din. Quat.*, 41, 105–114, <https://doi.org/10.4461/GFDQ.2018.41.16>, 2018.
- 660 Brenning, A.: Geomorphological, hydrological and climatic significance of rock glaciers in the Andes of Central Chile
661 (33–35 S). *Permafrost. Periglacial Process.*, 16(3), 231-240, <https://doi.org/10.1002/ppp.528>, 2005a.



- 662 Brenning, A.: Climatic and geomorphological controls of rock glaciers in the Andes of Central Chile: combining statistical
663 modelling and field mapping, Ph.D thesis, Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche
664 Fakultät II, 2005b.
- 665 Brighenti, S., Tolotti, M., Bruno, M. C., Engel, M., Wharton, G., Cerasino, L., Mair, V., and Bertoldi, W.: After the peak
666 water: the increasing influence of rock glaciers on alpine river systems, *Hydrol. Process.*, 33, 2804–2823,
667 <https://doi.org/10.1002/hyp.13533>, 2019.
- 668 Brighenti, S., Hotaling, S., Finn, D. S., Fountain, A. G., Hayashi, M., Herbst, D., Saros, J. E., Tronstad, L. M., and Millar,
669 C. I.: Rock glaciers and related cold rocky landforms: Overlooked climate refugia for mountain biodiversity, *Glob. Chang.*
670 *Biol.*, 27, 1504–1517, <https://doi.org/10.1111/gcb.15510>, 2021.
- 671 Carturan, L., Fontana, G. D., and Borga, M.: Estimation of winter precipitation in a high-altitude catchment of the Eastern
672 Italian Alps: Validation by means of glacier mass balance observations, *Geogr. Fis. e Din. Quat.*, 35, 37–48,
673 <https://doi.org/10.4461/GFDQ.2012.35.4>, 2012.
- 674 Carturan, L., Zuecco, G., Seppi, R., Zanoner, T., Borga, M., Carton, A., and Dalla Fontana, G.: Catchment-Scale
675 Permafrost Mapping using Spring Water Characteristics, *Permafr. Periglac. Process.*, 27, 253–270,
676 <https://doi.org/10.1002/ppp.1875>, 2016.
- 677 Carturan, L., De Blasi, F., Cazorzi, F., Zoccatelli, D., Bonato, P., Borga, M., and Dalla Fontana, G.: Relevance and Scale
678 Dependence of Hydrological Changes in Glacierized Catchments: Insights from Historical Data Series in the Eastern
679 Italian Alps, *Water*, 11, 89, <https://doi.org/10.3390/w11010089>, 2019.
- 680 Charton, J., Verfaillie, D., Jomelli, V., and Francou, B.: Early Holocene rock glacier stabilisation at col du Lautaret
681 (French Alps): Palaeoclimatic implications, *Geomorphology*, 394, 107962,
682 <https://doi.org/10.1016/j.geomorph.2021.107962>, 2021.
- 683 Chen, J. and Ohmura, A.: Estimation of Alpine glacier water resources and their change since the 1870s, *IAHS publ*, 193,
684 127–135, 1990.
- 685 Colucci, R. R., Forte, E., Žebre, M., Maset, E., Zanettini, C., and Guglielmin, M.: Is that a relict rock glacier?,
686 *Geomorphology*, 330, 177–189, <https://doi.org/10.1016/j.geomorph.2019.02.002>, 2019.
- 687 Cossart, E., Perrier, R., Schwarz, M., and Houee, S.: Mapping permafrost at a regional scale: interpolation of field data
688 by GIS application in the Upper Durance catchment (Southern French Alps), *GeoFocus*, 205–224, 2008.
- 689 Day-Lewis, F. D., Johnson, C. D., Singha, K., and Lane, J. W. J.: Best practices in electrical resistivity imaging: Data
690 collection and processing, and application to data from Corinna, Maine, EPA report, Boston, MA, 2008.
- 691 Delaloye, R.: Contribution à l'étude du pergélisol de montagne en zone marginale, 244 pp., 2004.
- 692 Delaloye, R., and Lambiel, C.: Evidence of winter ascending air circulation throughout talus slopes and rock glaciers
693 situated in the lower belt of alpine discontinuous permafrost (Swiss Alps). *Norsk Geografisk Tidsskrift-Norwegian*
694 *Journal of Geography*, 59(2), 194-203, 2005.



- 695 Delaloye, R., Reynard, E., Lambiel, C., Marescot, L., and Monnet, R.: Thermal anomaly in a cold scree slope (Creux du
696 Van, Switzerland), in: Proceedings of the Eighth International Conference of Permafrost, Zürich, Switzerland., 175–180,
697 2003.
- 698 Dlabáčková, T., Engel, Z., Uxa, T., Braucher, R., and Team, A.: ^{10}Be exposure ages and paleoenvironmental significance
699 of rock glaciers in the Western Tatra Mts., Western Carpathians, *Quat. Sci. Rev.*, 312, 108147,
700 <https://doi.org/10.1016/j.quascirev.2023.108147>, 2023.
- 701 Engelhardt, M., Schuler, T. V., and Andreassen, L. M.: Contribution of snow and glacier melt to discharge for highly
702 glacierised catchments in Norway, *Hydrol. Earth Syst. Sci.*, 18, 511–523, <https://doi.org/10.5194/hess-18-511-2014>,
703 2014.
- 704 Frauenfelder, R., Allgöwer, B., Haeberli, W., and Hoelzle, M.: Permafrost Investigations With GIS - A Case Study in the
705 Fletschhorn Area , Wallis , Swiss Alps, in: Seventh International Conference on Permafrost, 291–295, 1998.
- 706 Frauenfelder, R., Haeberli, W., Hoelzle, M., and Maisch, M.: Using relict rockglaciers in GIS-based modelling to
707 reconstruct Younger Dryas permafrost distribution patterns in the Err-Julier area, Swiss Alps, *Nor. Geogr. Tidsskr.*, 55,
708 195–202, <https://doi.org/10.1080/00291950152746522>, 2001.
- 709 Frei, C., and Schär, C.: A precipitation climatology of the Alps from high-resolution rain-gauge observations. *Int. J.*
710 *Climatol.*, 18(8), 873-900, [https://doi.org/10.1002/\(SICI\)1097-0088\(19980630\)18:8<873::AID-JOC255>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1097-0088(19980630)18:8<873::AID-JOC255>3.0.CO;2-9),
711 1998.
- 712 Gude, M., Dietrich, S., Mausbacher, R., Hauck, C., Molenda, R., Ruzicka, V., and Zacharda, M.: Probable occurrence of
713 sporadic permafrost in non-alpine scree slopes in central Europe, in: Proceedings 8th International Conference on
714 Permafrost, 331–336, 2003.
- 715 Guglielmin, M.: Il permafrost alpino: concetti, morfologia e metodi di individuazione (con tre indagini esemplificate in
716 alta Valtellina) / di Mauro Guglielmin; con contributi di Adalberto Notarpietro, Centro di studio per la geodinamica
717 alpina e quaternaria, Milano, 1997.
- 718 Haeberli, W.: Untersuchungen Zur Verbreitung Von Permafrost Zwischen Flueelapass Und Piz Grialetsch
719 (Graubuenden)., Versuchsanstalt für Wasserbau, Hydrologie und Glaziologie an der ETH, 1975.
- 720 Haeberli, W.: Creep of Mountain Permafrost: Internal Structure and Flow of Alpine Rock Glaciers., 1985.
- 721 Haeberli, W., Schaub, Y., and Huggel, C.: Increasing risks related to landslides from degrading permafrost into new lakes
722 in de-glaciating mountain ranges, *Geomorphology*, 293, 405–417, <https://doi.org/10.1016/j.geomorph.2016.02.009>,
723 2017.
- 724 Harrington, J. S., Mozil, A., Hayashi, M. and Bentley, L. R.: Groundwater flow and storage processes in an inactive rock
725 glacier, *Hydrol. Process.*, 32(20), 3070-3088, <https://doi.org/10.1002/hyp.13248>, 2018.
- 726 Harris, S. A. and Pedersen, D. E.: Thermal regimes beneath coarse blocky materials, *Permafr. Periglac. Process.*, 9, 107–
727 120, [https://doi.org/10.1002/\(SICI\)1099-1530\(199804/06\)9:2<107::AID-PPP277>3.0.CO;2-G](https://doi.org/10.1002/(SICI)1099-1530(199804/06)9:2<107::AID-PPP277>3.0.CO;2-G), 1998.
- 728 Hauck, C. and Kneisel, C.: Applied Geophysics in Periglacial Environments, edited by: Hauck, C. and Kneisel, C.,
729 Cambridge University Press, 1–248 pp., <https://doi.org/10.1017/CBO9780511535628>, 2008.



- 730 Hauck, C., Böttcher, M., and Maurer, H.: A new model for estimating subsurface ice content based on combined electrical
731 and seismic data sets, *The Cryosphere*, 5, 453–468, <https://doi.org/10.5194/tc-5-453-2011>, 2011. Hilbich, C., Hauck, C.,
732 Hoelzle, M., Scherler, M., Schudel, L., Völksch, I., Vonder Mühl, D., and Mäusbacher, R.: Monitoring mountain
733 permafrost evolution using electrical resistivity tomography: A 7-year study of seasonal, annual, and long-term variations
734 at Schilthorn, Swiss Alps, *J. Geophys. Res. Earth Surf.*, 113, F01S90, <https://doi.org/10.1029/2007JF000799>, 2008.
- 735 Ikeda, A. and Matsuoka, N.: Degradation of talus-derived rock glaciers in the Upper Engadin, Swiss Alps. *Permafr.*
736 *Periglac. Process.*, 13, 145–161, <https://doi.org/10.1002/ppp.413>, 2002.
- 737 Ilyashuk, B. P., Ilyashuk, E. A., Psenner, R., Tessadri, R., and Koinig, K. A.: Rock glacier outflows may adversely affect
738 lakes: Lessons from the past and present of two neighboring water bodies in a crystalline-rock watershed, *Environ. Sci.*
739 *Technol.*, 48, 6192–6200, <https://doi.org/10.1021/es500180c>, 2014.
- 740 Imhof, M., Pierrehumbert, G., Haeberli, W., and Kienholz, H.: Permafrost investigation in the Schilthorn Massif, Bernese
741 Alps, Switzerland, *Permafr. Periglac. Process.*, 11, 189–206, [https://doi.org/10.1002/1099-1530\(200007/09\)11:3<189::AID-PPP348>3.0.CO;2-N](https://doi.org/10.1002/1099-1530(200007/09)11:3<189::AID-PPP348>3.0.CO;2-N), 2000.
- 743 Isotta, F. A., Frei, C., Weilguni, V., Perčec Tadić, M., Lassègues, P., Rudolf, B., Pavan, V., Cacciamani, C., Antolini, G.,
744 Ratto, S. M., Munari, M., Micheletti, S., Bonati, V., Lussana, C., Ronchi, C., Panettieri, E., Marigo, G., and Vertačnik,
745 G.: The climate of daily precipitation in the Alps: Development and analysis of a high-resolution grid dataset from pan-
746 Alpine rain-gauge data, *Int. J. Climatol.*, 34, 1657–1675, <https://doi.org/10.1002/joc.3794>, 2014.
- 747 Janke, J. R., Bellisario, A. C., and Ferrando, F. A.: Classification of debris-covered glaciers and rock glaciers in the Andes
748 of central Chile, *Geomorphology*, 241, 98–121, <https://doi.org/10.1016/j.geomorph.2015.03.034>, 2015.
- 749 Janke, J. R., Ng, S., and Bellisario, A.: An inventory and estimate of water stored in firn fields, glaciers, debris-covered
750 glaciers, and rock glaciers in the Aconcagua River Basin, Chile, *Geomorphology*, 296, 142–152,
751 <https://doi.org/10.1016/j.geomorph.2017.09.002>, 2017.
- 752 Jones, D. B., Harrison, S., Anderson, K., and Betts, R. A.: Mountain rock glaciers contain globally significant water
753 stores, *Sci. Rep.*, 8, 2834, <https://doi.org/10.1038/s41598-018-21244-w>, 2018.
- 754 Jones, D. B., Harrison, S., Anderson, K., and Whalley, W. B.: Rock glaciers and mountain hydrology: A review, *Earth-*
755 *Science Rev.*, 193, 66–90, <https://doi.org/10.1016/j.earscirev.2019.04.001>, 2019.
- 756 Juliussen, H. and Humlum, O.: Thermal regime of openwork block fields on the mountains Elgåhogna and Sjølen, Central-
757 Eastern Norway. *Permafr. Periglac. Process.*, 19 (1), 1–18, <https://doi.org/10.1002/ppp.607>, 2008.
- 758 Kääb, A.: Rock glaciers and protalus forms. In *Encyclopedia of Quaternary Science*, 2nd Edition, Volume 3, SA Elias
759 (editor-in chief). Elsevier: Amsterdam; 535–541, 2013.
- 760 Kellerer-Pirklbauer, A.: Aspects of glacial, paraglacial and periglacial processes and landforms of the Tauern Range,
761 Austria. Doctoral Thesis, University of Graz, 2008.
- 762 Kellerer-Pirklbauer, A.: Long-term monitoring of sporadic permafrost at the eastern margin of the European Alps
763 (Hochreichart, Seckauer Tauern range, Austria), *Permafr. Periglac. Process.*, 30, 260–277,
764 <https://doi.org/10.1002/ppp.2021>, 2019.



- 765 Kellerer-Pirklbauer, A., Lieb, G. K., and Kleinfelchner, H.: A new rock glacier inventory of the eastern European Alps,
766 *Austrian J. Earth Sci.*, 105, 78–93, 2012.
- 767 Kellerer-Pirklbauer, A., Pauritsch, M., Morawetz, R., and Kuehnast, B.: Thickness and internal structure of relict rock
768 glaciers – a challenge for geophysics : Examples from two rock glaciers in the Eastern Alps, in: EGU General Assembly
769 Conference Abstracts, 12581, 2014.
- 770 Kellerer-Pirklbauer, A., Lieb, G. K., and Kaufmann, V.: The dösen rock glacier in central Austria: A key site for
771 multidisciplinary long-term rock glacier monitoring in the eastern alps, *Austrian J. Earth Sci.*, 110, 10–17738,
772 <https://doi.org/10.17738/ajes.2017.0013>, 2017.
- 773 Kofler, C., Steger, S., Mair, V., Zebisch, M., Comiti, F. and Schneiderbauer, S.: An inventory-driven rock glacier status
774 model (intact vs. relict) for South Tyrol, Eastern Italian Alps. *Geomorphology*, 350, 106887,
775 <https://doi.org/10.1016/j.geomorph.2019.106887>, 2020.
- 776 Lambiel, C. and Reynard, E.: Regional modelling of present, past and future potential distribution of discontinuous
777 permafrost based on a rock glacier inventory in the Bagnes-Hérémence area (Western Swiss Alps), *Nor. Geogr. Tidsskr.*
778 - *Nor. J. Geogr.*, 55, 219–223, <https://doi.org/10.1080/00291950152746559>, 2001.
- 779 Lewkowicz, A. G., Etmüller, B., and Smith, S. L.: Characteristics of Discontinuous Permafrost based on Ground
780 Temperature Measurements and Electrical Resistivity Tomography, Southern Yukon, Canada, *Permafr. Periglac.*
781 *Process.*, 22, 320–342, <https://doi.org/10.1002/ppp.703>, 2011.
- 782 Lilleøren, K. S., Etmüller, B., Rouyet, L., Eiken, T., Slinde, G., and Hilbich, C.: Transitional rock glaciers at sea level
783 in northern Norway, *Earth Surf. Dynam.*, 10, 975–996, <https://doi.org/10.5194/esurf-10-975-2022>, 2022. Millar, C.I. and
784 Westfall, R.D.: Geographic, hydrological, and climatic significance of rock glaciers in the Great Basin, USA. *Arct.*
785 *Antarct. Alp. Res.*, 51(1), 232-249, <https://doi.org/10.1080/15230430.2019.1618666>, 2019.
- 786 Morard, S., Delaloye, R., and Dorthe, J.: Seasonal thermal regime of a mid-latitude ventilated debris accumulation, in:
787 *Proceedings of the 9th International Conference on Permafrost*, Fairbanks, Alaska, 1233–1238, 2008.
- 788 Pavoni, M., Boaga, J., Carrera, A., Zuecco, G., Carturan, L., and Zumiani, M.: Brief communication: Mountain permafrost
789 acts as an aquitard during an infiltration experiment monitored with electrical resistivity tomography time-lapse
790 measurements, *Cryosph.*, 17, 1601–1607, <https://doi.org/10.5194/tc-17-1601-2023>, 2023.
- 791 Popescu, R.: Permafrost investigations in Iezer Mountains, Southern Carpathians, *Rev. Geomorfol.*, 20, 102–122,
792 <https://doi.org/10.21094/rg.2018.033>, 2018.
- 793 Scotti, R., Brardinoni, F., Alberti, S., Frattini, P. and Crosta, G.B.: A regional inventory of rock glaciers and protalus
794 ramparts in the central Italian Alps. *Geomorphology*, 186, 136-149, <https://doi.org/10.1016/j.geomorph.2012.12.028>,
795 2013.
- 796 Salvatore, M. C., Zanoner, T., Baroni, C., Carton, A., Banchieri, F. A., Viani, C., Giardino, M., and Perotti, L.: The state
797 of Italian glaciers: A snapshot of the 2006–2007 hydrological period, *Geogr. Fis. e Din. Quat.*, 38, 175–198,
798 <https://doi.org/10.4461/GFDQ.2015.38.16>, 2015.



- 799 Sannino, C., Borruso, L., Mezzasoma, A., Battistel, D., Ponti, S., Turchetti, B., Buzzini, P. and Guglielmin, M.: Abiotic
800 factors affecting the bacterial and fungal diversity of permafrost in a rock glacier in the Stelvio Pass (Italian Central Alps).
801 Appl. Soil Ecol., 166, 104079, <https://doi.org/10.1016/j.apsoil.2021.104079>, 2021.
- 802 Scapozza, C.: Contributo dei metodi termici alla prospezione del permafrost montano : esempi dal massiccio della Cima
803 di Gana Bianca (Val Blenio , Svizzera), Boll. della Soc. Ticin. di Sci. Nat., 66, 55–66, 2009.
- 804 Schaffer, N., MacDonell, S., Réveillet, M., Yáñez, E. and Valois, R.: Rock glaciers as a water resource in a changing
805 climate in the semiarid Chilean Andes. Reg. Environ. Change, 19, 1263–1279, [https://doi.org/10.1007/s10113-018-01459-](https://doi.org/10.1007/s10113-018-01459-3)
806 3, 2019.
- 807 Schrott, L., Otto, J. C., and Keller, F.: Modelling alpine permafrost distribution in the Hohe Tauern region, Austria.
808 Austrian J. Earth Sci., 105(2), 2012.
- 809 Seppi, R., Carton, A., and Baroni, C.: Rock glacier relitti e antica distribuzione del permafrost nel Gruppo Adamello
810 Presanella (Alpi Centrali), Alp. Mediterr. Quat., 23, 137–144, 2010.
- 811 Seppi, R., Carton, A., Zumiani, M., Dall’Amico, M., Zampedri, G., and Rigon, R.: Inventory, distribution and topographic
812 features of rock glaciers in the southern region of the Eastern Italian Alps (Trentino), Geogr. Fis. e Din. Quat., 35, 185–
813 197, <https://doi.org/10.4461/GFDQ.2012.35.17>, 2012.
- 814 Seppi, R., Carturan, L., Carton, A., Zanoner, T., Zumiani, M., Cazorzi, F., Bertone, A., Baroni, C., and Salvatore, M. C.:
815 Decoupled kinematics of two neighbouring permafrost creeping landforms in the Eastern Italian Alps, Earth Surf. Process.
816 Landforms, 44, 2703–2719, <https://doi.org/10.1002/esp.4698>, 2019.
- 817 Slangen, A. B. A. and van de Wal, R. S. W.: An assessment of uncertainties in using volume-area modelling for computing
818 the twenty-first century glacier contribution to sea-level change, Cryosph., 5, 673–686, [https://doi.org/10.5194/tc-5-673-](https://doi.org/10.5194/tc-5-673-2011)
819 2011, 2011.
- 820 Strozzi, T., Kääb, A., and Frauenfelder, R.: Detecting and quantifying mountain permafrost creep from in situ inventory,
821 space-borne radar interferometry and airborne digital photogrammetry, Int. J. Remote Sens., 25, 2919–2931,
822 <https://doi.org/10.1080/0143116042000192330>, 2004.
- 823 Tapia-Baldis, C. and Trombotto-Liaudat, D.: Permafrost model in coarse-blocky deposits for the Dry Andes, Argentina
824 (28– 33° S), Cuadernos de Investigación Geográfica, 46, 33–58, <https://doi.org/10.18172/cig.3802>, 2020.
- 825 Thies, H., Nickus, U., Tolotti, M., Tessadri, R., and Krainer, K.: Evidence of rock glacier melt impacts on water chemistry
826 and diatoms in high mountain streams, Cold Reg. Sci. Technol., 96, 77–85,
827 <https://doi.org/10.1016/j.coldregions.2013.06.006>, 2013.
- 828 Wagner, T., Pauritsch, M., and Winkler, G.: Impact of relict rock glaciers on spring and stream flow of alpine watersheds:
829 examples of the Niedere Tauern Range, Eastern Alps (Austria), Aust. J. Earth Sci., 109, 84–98. doi:
830 10.17738/ajes.2016.0006, 2016.
- 831 Wagner, T., Kainz, S., Helfricht, K., Fischer, A., Avian, M., Krainer, K., and Winkler, G.: Assessment of liquid and solid
832 water storage in rock glaciers versus glacier ice in the Austrian Alps, Sci. Total Environ., 800, 149593,
833 <https://doi.org/10.1016/j.scitotenv.2021.149593>, 2021.



834 Winkler, G., Wagner, T., Pauritsch, M., Birk, S., Kellerer-Pirklbauer, A., Benischke, R., Leis, A., Morawetz, R.,
835 Schreilechner, M. G., and Hergarten, S.: Identification and assessment of groundwater flow and storage components of
836 the relict Schöneben Rock Glacier, Niedere Tauern Range, Eastern Alps (Austria), *Hydrogeol. J.*, 24, 937–953,
837 <https://doi.org/10.1007/s10040-015-1348-9>, 2016.

838 Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Hoelzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlstrøm,
839 A. P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L. N., Cáceres, B. E., Casassa, G., Cobos, G., Dávila, L. R.,
840 Delgado Granados, H., Demuth, M. N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Ove Hagen, J.,
841 Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V. V., Portocarrero, C. A., Prinz, R., Sangewar, C. V.,
842 Severskiy, I., Sigurdsson, O., Soruco, A., Usubaliev, R., and Vincent, C.: Historically unprecedented global glacier
843 decline in the early 21st century, *J. Glaciol.*, 61, 745–762, <https://doi.org/10.3189/2015JoG15J017>, 2015.

844

845

846

847