

Brief communication: Identification of 140,000-year-old blue ice in Grove Mountains, East Antarctica, by krypton-81 dating

Zhengyi Hu^a, Wei Jiang^{b,c}, Yuzhen Yan^d, Yan Huang^e, Xueyuan Tang^a, Lin Li^a, Florian Ritterbusch^c, Guo-Min Yang^c, Zheng-Tian Lu^{b,c}, Guitao Shi^{e,f*}

^aMNR Key Laboratory for Polar Science, Polar Research Institute of China, Shanghai, 200062, China

^bHefei National Laboratory, University of Science and Technology of China, Hefei, 230088, China

^cChinese Academy of Sciences Center for Excellence in Quantum Information and Quantum Physics, School of Physical Sciences, University of Science and Technology of China, Hefei 230026, China

^dState Key Laboratory of Marine Geology, Tongji University, Shanghai, 200092, China

^eKey Laboratory of Geographic Information Science, School of Geographic Sciences, East China Normal University, Shanghai, 200241, China

^fState Key Laboratory of Estuarine and Coastal Research, East China Normal University, Shanghai 200241, China

*Correspondence to: Guitao Shi, gtshi@geo.ecnu.edu.cn

Abstract:

The presence of exceptionally old ice and the relative ease of access make Antarctic blue ice areas (BIAs) attractive paleoclimate archives. However, only a handful of BIAs, mostly situated in West Antarctica and along the Trans-Antarctic Mountains, have been investigated for this purpose. Here, we present the age of surface ice from the Grove Mountains BIA in Princess Elizabeth Land, East Antarctica, determined by measuring ⁸¹Kr in the trapped air. Two samples yield an average age of 143^{+33}_{-29} kyr. Together with the reported terrestrial age of a chondrite, we conclude that Grove Mountains BIA holds considerable potential for paleoclimate studies.

1. Introduction


Antarctic ice cores provide a wealth of information about the Earth's past climate and atmospheric composition, especially greenhouse gases (e.g., Petit et al., 1999). International efforts are underway to locate and retrieve an ice core dating back to 1.5 Myr that is in stratigraphic order (Fischer et al., 2013; Passalacqua et al., 2018), e.g., the ongoing Beyond Epica Oldest Ice project at Little Dome C (<https://www.beyondepica.eu/en/about/>). However, such endeavors are expensive and time-consuming. Shallow drilling in blue ice areas (BIAs) in Antarctica has therefore emerged as a complementary approach (Yan et al., 2019). BIAs are regions where ablation exceeds accumulation. The negative mass balance at the surface is maintained by the supplies of crystalline glacial ice from below. In this case, ancient ice that was once buried deep in the ice sheet is brought to the surface and can be readily accessed. The presence of meteorites that have terrestrial ages up to 2 Myr in the BIAs hints at the existence of 300 kyr old ice (Scherer et al., 1997).

In Antarctica, a number of BIAs have been investigated to understand local meteorology, glaciology, and meteorites (e.g., Spaulding et al., 2013; Liu et al., 2010; Scherer et al., 1997). Thus far, debris-free ice samples were recovered from only five blue ice areas for the purpose of paleoclimate studies (Table 1 and Figure 1): Mount Moulton and Patriot Hills in West Antarctica, and Allan Hills, Taylor Glacier, and Larsen Glacier in East Antarctica. All three East Antarctic BIAs are situated in Victoria Land despite considerable presence of BIAs in other parts of East Antarctica, such as near Grove Mountains. In this study, we focus on the Grove Mountains, which consist of a series of nunataks southwest of Princess Elizabeth Land, East Antarctica. They are approximately 400 km from the Antarctic coast, is on a different side of the continent, far away from the cluster of previous sites (Figure 1). We report the age of surface ice in Grove Mountains determined by ^{81}Kr dating and evaluate the potential of Grove Mountains BIA as paleoclimate archives.

Table 1. List of Antarctic blue ice areas as paleoclimate archives

Areas	Location	Age range	References
Grove Mountains	72.99° S, 75.22°E	143 kyr	This study
Mount Moulton	76.7° S, 134.7° W	105-136 kyr	(Korotkikh et al., 2011)
Patriot Hills	80.3° S, 81.4° W	10-80 kyr;130-134 kyr	(Turney et al., 2020)
Allan Hills	76.7° S, 159.4° E	90-250 kyr; >1 Myr	(Spaulding et al., 2013; Yan et al., 2019)

Taylor Glacier	77.8° S, 161.8° E	9-133 kyr	(Buizert et al., 2014)
Larsen Glacier	74.9° S, 161.6° E	9-25 kyr	(Lee et al., 2022)



60

Figure 1 Antarctic blue ice areas (BIAs) and the Grove Mountains. (a) Sites of BIAs explored as paleoclimate archives listed in Table 1 (closed black stars), sites of deep ice coring, and the traverse route of this work from coastal Zhongshan station to Dome A (base map from ESRI). (b) Satellite imagery of the Grove Mountains BIAs with local ice flow lines (purple arrows) and the transect of an airborne ice radar survey (red arrow) (base image is Landsat Image Mosaic of Antarctica). The sampling site for this study is marked with a red triangle. (c) Radar profile of the transect. (d) Surface morphology of the blue ice at the sampling site, the red triangle in panel b (Credit: Z. Hu).

65

70 2. Materials and methods

2.1 Site and sample description

The Grove Mountains BIA is located in the southwest of the Princess Elizabeth Land, East Antarctica (Figure 1). Ice flows from the southeast to the northwest and eventually drains into the Lambert Glacier. Ice flow is blocked by a series of nunataks in this region. The combined
75 effect of glacial flows and topographic barriers leads to the formation of blue ice as well as a large number of crevasses that prevent access for ice drilling. The highest nunatak, Mason Peak, has an elevation of 2365 m and a topographic prominence of ~700 m. The average ice elevation of Grove Mountains is ~2000 m, and past radar surveys revealed steep bedrock topography and thick (>1000 m) ice layers (Figure 1). The annual average temperature in Grove Mountains is
80 ~-30 °C. Similar to other Antarctic BIAs, Grove Mountains BIA is a meteorite concentration site. To date, more than 10000 meteorites have been recovered by the Chinese National Antarctica Research Expedition (CHINARE) from this region. In January 2016, surface blue ice (up to ~50 cm in depth) was collected manually using stainless-steel spades near the mid-Gale Escarpment where the meteorites are concentrated. Surface morphology of the blue ice at
85 the sampling site is shown in Figure 1d. The sampling size is approximately 40 × 40 × 40 cm. The ice samples are irregular blocks with length ranging from about 10 cm to 30 cm. Based on visual inspection, there are no clear cracks and melted layers in the ice at the sampling site. In total, about 40 kg of ice was collected (72.99° S, 75.22° E). A ~5 cm thick layer of surface ice was removed before sampling. The collected ice was kept in clean polyethylene bags and then
90 preserved in insulated cabinets and transported under freezing conditions (-20 °C).

2.2 Analytical methods

The blue ice collected from Grove Mountains was processed in two batches. The first batch (batch A) contains large ice blocks (size ~20 cm or bigger) with a total weight of 26 kg. The
95 second batch (batch B) contains small ice pieces (size 10-20 cm) and weighs about 9 kg. The reason to separate them into two batches is to see if there are more modern air contaminations in the batch with small ice samples. Before melting, the outer 3-5 mm layer of the ice was removed to reduce potential contamination from modern air. The ice was then melted in an evacuated container. The released air was transferred into 1L stainless steel bottles with a
100 compressor. The extraction system typically achieves recoveries > 95% and contamination < 1 %. The gas contents obtained for sample batch A and batch B were 95 mL STP/kg and 86 mL STP/kg, respectively. The difference is probably due to the larger specific surface area of the smaller sample (batch B) compared to the large one (batch A) and hence greater gas losses from smaller ice samples. Nonetheless, these gas contents are comparable to the blue ice

105 retrieved from other BIAs in Antarctica (e.g., Buizert et al., 2014). Krypton (Kr) was separated from the extracted air with a purification system based on titanium gettering and gas chromatography (Jiang et al., 2020), yielding Kr purities and recoveries higher than 90%. 2.2 μL STP and 0.6 μL STP Kr were obtained for the larger and smaller sample, respectively.

The Kr sample was measured with the atom trap trace analysis (ATTA) method (Jiang et al., 110 2020). ^{81}Kr and ^{85}Kr atoms are selectively captured by laser beams into a magneto-optical trap and are counted by detecting their fluorescence. Meanwhile, ^{83}Kr (a stable isotope) is chosen as the reference and its capture rate is also measured. Each analysis took about 5 hours. The measurement is cycled between the atom-counting mode for ^{81}Kr and ^{85}Kr , and the capture-rate mode for ^{83}Kr in order to cancel the slow drifts in the capture efficiencies of the instrument. 115 The anthropogenic ^{85}Kr isotope is analyzed since it has a half-life of 10.7 years, making it a good indicator of cross-sample contamination from the modern reference sample. As the amount of Kr from the ice samples is generally limited, it is important to keep the cross-sample contamination under control. This effect comes from the discharge source in the ATTA instrument. On the one hand, it slowly consumes the Kr sample in the system and makes the 120 effective sample size smaller. On the other hand, Kr from the previous measurement is slowly released from vacuum parts to cause cross-sample contamination. In order to reduce this effect, the vacuum system of the ATTA instrument was washed continuously with a Xe discharge for one week. After the washing, the Kr outgassing rate was about 10^{-3} μL STP/h. For a 5-hour measurement the cross-sample contamination is 1~3%. The residual cross-sample 125 contamination effect was corrected based on the ^{85}Kr measurement.

The measured relative ^{81}Kr abundance was used to calculate the age of the sample based on the radioactive decay law and the atmospheric input function of ^{81}Kr . The uncertainty of the ^{81}Kr age mainly came from the statistical errors for atom counting. The uncertainty caused by the cross-sample contamination correction was included through error propagation. Since the 130 atom counts for both ^{81}Kr and ^{85}Kr were low (10 ~ 100), we adopted the Feldman-Cousins method, which provided a unified approach to treat measurements with small signals (Feldman and Cousins, 1998). Besides the statistical error, there were additional systematic errors due to the uncertainty of the half-life of ^{81}Kr (229 ± 11 ka) and the uncertainty of the atmospheric ^{81}Kr input functions (Zappala et al., 2020). Note that the measurements were performed in 2017. 135 The ATTA instrument has been improved significantly since then. The sample requirement is now less than 2.0 kg and the dating precision is also better (Crotti et al., 2021).

Two pieces of blue ice from batch A were randomly selected for chemical measurements. In a Class 1000 clean room, the surface layer of ~1 cm was washed with ultrapure Milli-Q water

(18.2M Ω) to remove any surface contaminants. Then, the ice was melted under a super clean
140 hood (Class 100) at 20°C for chemical measurements. The major chemical ions, Na⁺, K⁺, Mg²⁺,
Cl⁻, NO₃⁻, and SO₄²⁻, were determined with an ICS-3000 IC system (Dionex, USA). More
details on ion analysis are provided in Shi et al. (2012). The $\delta^{18}\text{O}$ and δD of ice were measured
with a wavelength scanned cavity ring down spectroscopy (WS-CRDS) instrument, Picarro L-
2130i (Picarro Inc., USA), with the respective analysis precision of 0.05‰ and 0.5‰. Details
145 on the water isotope analysis are described in a previous study (Ma et al., 2020).

2.3 Ice radar survey

During CHINARE36 (2019-2020), radar data were collected in the Grove Mountains using an
ice radar system mounted on the Snow Eagle 601 fixed-wing aircraft. The length of the radar
150 line is about 45 km (Figure 1b). During data acquisition, the system emits electromagnetic
waves with a central frequency of 60 MHz, with a peak emission frequency of 8 kW and a
pulse frequency of 6250 Hz. The radar antenna system consists of two flat dipole antennas that
are mounted below the aircraft wing and used for both transmission and reception.

155 3. Results

The results of radiometric Kr dating of two batches of ice from the Grove Mountains BIAs are
shown in Table 2. The isotopic abundances of ⁸¹Kr in the trapped air of the two batches are
statistically indistinguishable from each other. Furthermore, despite the different ice sizes and
weights, both batches show non-zero but similar level of ⁸⁵Kr activities, indicating modern air
160 contaminations possibly due to cracks near the blue ice surface. Similar intrusion of the modern
atmosphere to the blue ice has previously been observed in other BIAs as well (Spaulding et
al., 2013). After correcting for the modern air contamination (assuming an atmospheric ⁸⁵Kr
activity of 70 ± 5 decay per minute per cubic centimeter krypton at STP (dpm/cc) at the
sampling time (25 Jan. 2016); Kersting et al. (2020)), the averaged relative ⁸¹Kr/Kr is 65 ± 6
165 (pMKr), which corresponds to an age of 143⁺³³₋₂₉ kyr. This age is comparable to the previously
reported age of the surface ice in Mount Moulton, Patriot Hills and Allan Hills BIAs (Korotkikh
et al., 2011; Spaulding et al., 2013; Turney et al., 2020). Moreover, a CR chondrite (No. GRV
021710) discovered near the sampling site has a terrestrial age of 260 kyr (Lu, 2008). These
results hint at the presence of even older blue ice in the vicinity of Grove Mountains.

170

Table 2 ^{81}Kr dating of blue ice near Grove Mountains

	Sample size (kg)	Kr extracted (μL STP)	^{85}Kr activity at sampling time (dpm/cc)	$^{81}\text{Kr}/\text{Kr}$ (pMKr)	$^{81}\text{Kr}/\text{Kr}_{\text{corrected}}$ (pMKr)	^{81}Kr age (ka)
Batch A	26	2.2	27 ± 1	76 ± 5	61 ± 8	165^{+48}_{-43}
Batch B	9	0.7	24 ± 1	82 ± 7	73 ± 11	107^{+53}_{-45}
Average	--	--	--	--	65 ± 6	143^{+33}_{-29}

175 The mean concentrations of Na^+ , K^+ , Mg^{2+} , Cl^- , NO_3^- , and SO_4^{2-} in the samples are 34.9, 9.2, 11.1, 84.0, 44.4, and 94.5 ng g^{-1} , respectively, which are similar to the values of surface snow samples collected along the Chinese inland Antarctica traverse route, about $\sim 60\text{km}$ from the study site (Shi et al., 2021). The mean $\delta^{18}\text{O}$ and $\delta^2\text{H}$ of the blue ice are -40.3‰ and -321.2‰ , respectively, also similar to those of the nearby surface snow (Ma et al., 2020). These stable

180 water isotope values are much higher than those of the snow and ice on the East Antarctic plateau today (Xiao et al., 2008 and references therein) (Table S1) and during the Last Interglacial period (e.g., Petit et al., 1999). Assuming no isotopic modifications after snow deposition and during ice flow, the original deposition site of the surface ice at the Grove Mountains BIA today was likely local, but we acknowledge that the exact location remains

185 unknown. Any future attempt to interpret those isotope data in the context of paleoclimate would require a more thorough investigation of the provenance of the blue ice.

4. Discussion and Conclusions

The mean age of two surface ice samples from the Grove Mountains BIA, dated radiometrically

190 by ^{81}Kr , is 143^{+33}_{-29} kyr; that is, the ice dates back to the Last Interglacial, holding important implications for paleoclimate studies. Furthermore, a meteorite (GRV021710) that has a terrestrial age of 260 kyr suggests that Grove Mountains BIA could harbor even older ice. In general, the major chemical ions and stable water isotopes of the ice resemble those of the nearby surface snow and differ from those from Antarctic plateau sites, suggesting that the blue

195 ice originated nearby. There are no previously published deep ice core records from the Princess Elizabeth Land that date back to the Last Interglacial. Consequently, the Grove Mountains BIA holds the potential to provide large-volume ice samples to study the climate variations during the Last Interglacial in the Indian Ocean sector of Antarctica. To obtain old ice, the potential drilling sites in the BIAs are usually located upstream of the ice flow, where

200 the ice stream was blocked by the nunataks, similar to the drilling sites in Allan Hills (Yan et

al., 2019). Accordingly, the potential old ice drilling sites in the Grove Mountains BIA are expected to be around the mid-Gale Escarpment, following the ice flow direction in this region (Figure 1). It is noted that there are a large number of ice crevasses formed on the side of the mid-Gale Escarpment facing the ice flow, making it currently inaccessible from the ground.

205 The radar profile provides the direct observations of deep englacial stratigraphy in this BIA (Figure 1c). However, only some disturbed layers can be imaged, at depths of 200-400m beneath the ice surface. The radar image showed that the internal layers are not well identified at a depth of >500m, which could be the result of complex ice flow patterns around the nunataks. Nonetheless, it can be seen that the subglacial topographic mountains may cause the ice to flow

210 toward the surface, especially near the nunataks, where the ice depth is relatively shallow, at a few hundred meters. These areas are expected to be potential shallow ice core drilling sites, i.e., easier access to the oldest ice. In addition to retrieving ice cores, a synergistic effect of drilling operations in the Grove Mountains is the potential recovery of bedrock samples. Some previous studies suggest the East Antarctic Ice Sheet would retreat beyond Grove Mountains

215 during past warm intervals (Liu et al., 2010). Bedrock samples from the Grove Mountains can help evaluate this hypothesis and improve our understanding of ice sheet behaviors in a fast-warming world.

In addition, the bedrock at the Grove Mountains BIA could reveal important information about the stability of the East Antarctic Ice Sheet in past Interglacials. Given these considerations,

220 we conclude that Grove Mountains are a region with high scientific values. Future drilling operations in the Grove Mountains BIA could also benefit from its close proximity to a nearby Antarctic research base (Chinese Taishan Station; Figure 1a). More systematic glaciological surveys in this region are called for, with the ultimate goal of retrieving ice cores and bedrock samples to study paleoclimate and past ice sheet behaviors.

225

Code and data availability

Data presented in this work are included in the main text.

Author contribution

230 GS and ZH conceived the study. YY, WJ, and GS designed and wrote the manuscript with the support of all co-authors. ZH and GS analyzed ions and stable water isotopes. WJ, FR, GY, and ZL analyzed the Kr data. YH, XT, LL, and GS prepared the figure.

Competing interests

235 The authors declare no conflicts of interest relevant to this study.

Acknowledgments

This work was supported by the National Science Foundation of China (Grant Nos. 42276243, 41922046, 42071306, T2325024, and 41727901), the Innovation Program for
240 Quantum Science and Technology (2021ZD0303101), and the Program of Shanghai Academic Research Leader (Grant No. 20XD1421600). The authors are grateful to CHINARE members for their support and assistance in sample collection.

References

- 245 Buizert, C., Baggenstos, D., Jiang, W., Purtschert, R., Petrenko, V. V., Lu, Z.-T., Müller, P., Kuhl, T., Lee, J., Severinghaus, J. P., and Brook, E. J.: Radiometric ^{81}Kr dating identifies 120,000-year-old ice at Taylor Glacier, Antarctica, *Proc. Natl. Acad. Sci.*, 111, 6876-6881, <https://doi.org/10.1073/pnas.1320329111>, 2014.
- 250 Crotti, I., Landais, A., Stenni, B., Bazin, L., Parrenin, F., Frezzotti, M., Ritterbusch, F., Lu, Z.-T., Jiang, W., Yang, G.-M., Fourné, E., Orsi, A., Jacob, R., Minster, B., Prié, F., Dreossi, G., and Barbante, C.: An extension of the TALDICE ice core age scale reaching back to MIS 10.1, *Quaternary Sci. Rev.*, 266, <https://doi.org/10.1016/j.quascirev.2021.107078>, 2021.
- Feldman, G. J. and Cousins, R. D.: Unified approach to the classical statistical analysis of small signals, *Phys. Rev. D*, 57, 3873-3889, <https://doi.org/10.1103/PhysRevD.57.3873>,
255 1998.
- Fischer, H., Severinghaus, J., Brook, E., Wolff, E., Albert, M., Alemany, O., Arthern, R., Bentley, C., Blankenship, D., Chappellaz, J., Creyts, T., Dahl-Jensen, D., Dinn, M., Frezzotti, M., Fujita, S., Gallee, H., Hindmarsh, R., Hudspeth, D., Jugie, G., Kawamura, K., Lipenkov, V., Miller, H., Mulvaney, R., Parrenin, F., Pattyn, F., Ritz, C., Schwander, J., Steinhage, D.,
260 van Ommen, T., and Wilhelms, F.: Where to find 1.5 million yr old ice for the IPICS "Oldest-Ice" ice core, *Clim. Past*, 9, 2489-2505, <https://doi.org/10.5194/cp-9-2489-2013>, 2013.
- Jiang, W., Hu, S.-M., Lu, Z.-T., Ritterbusch, F., and Yang, G.-m.: Latest development of radiokrypton dating – A tool to find and study paleogroundwater, *Quatern. Int.*, 547, 166-171, <https://doi.org/10.1016/j.quaint.2019.04.025>, 2020.
- 265 Kersting, A., Schlosser, C., Bollhöfer, A., and Suckow, A.: Evaluating 5 decades of atmospheric ^{85}Kr measurements in the southern hemisphere to derive an input function for dating water and ice with implications for interhemispheric circulation and the global ^{85}Kr emission inventory, *J. Environ. Radioactiv.*, 225, 106451, <https://doi.org/10.1016/j.jenvrad.2020.106451>, 2020.
- 270 Korotkikh, E. V., Mayewski, P. A., Handley, M. J., Sneed, S. B., Introne, D. S., Kurbatov, A. V., Dunbar, N. W., and McIntosh, W. C.: The last interglacial as represented in the glaciochemical record from Mount Moulton Blue Ice Area, West Antarctica, *Quaternary Sci. Rev.*, 30, 1940-1947, <https://doi.org/10.1016/j.quascirev.2011.04.020>, 2011.
- 275 Lee, G., Ahn, J., Ju, H., Ritterbusch, F., Oyabu, I., Buizert, C., Kim, S., Moon, J., Ghosh, S., Kawamura, K., Lu, Z.-T., Hong, S., Han, C. H., Hur, S. D., Jiang, W., and Yang, G.-M.: Chronostratigraphy of the Larsen blue-ice area in northern Victoria Land, East Antarctica, and its implications for paleoclimate, *Cryosphere*, 16, 2301-2324, <https://doi.org/10.5194/tc-16-2301-2022>, 2022.

- 280 Liu, X., Huang, F., Kong, P., Fang, A., Li, X., and Ju, Y.: History of ice sheet elevation in East Antarctica: Paleoclimatic implications, *Earth Planet. Sci. Lett.*, 290, 281-288, <https://doi.org/10.1016/j.epsl.2009.12.008>, 2010.
- Lu, R.: Classification and study of cosmogenic ^{10}Be and ^{26}Al of Antarctic meteorites collected from Grove Mountains region, PhD, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang, 91 pp., 2008.
- 285 Ma, T., Li, L., Li, Y., An, C., Yu, J., Ma, H., Jiang, S., and Shi, G.: Stable isotopic composition in snowpack along the traverse from a coastal location to Dome A (East Antarctica): Results from observations and numerical modeling, *Polar Sci.*, 24, 100510, <https://doi.org/10.1016/j.polar.2020.100510>, 2020.
- 290 Passalacqua, O., Cavitte, M., Gagliardini, O., Gillet-Chaulet, F., Parrenin, F., Ritz, C., and Young, D.: Brief communication: Candidate sites of 1.5 Myr old ice 37 km southwest of the Dome C summit, East Antarctica, *Cryosphere*, 12, 2167-2174, <https://doi.org/10.5194/tc-12-2167-2018>, 2018.
- Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J. M., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., 295 Lipenkov, V. Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E., and Stievenard, M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429-436, <https://doi.org/10.1038/20859>, 1999.
- Scherer, P., Schultz, L., Neupert, U., Knauer, M., Neumann, S., Leya, I., Michel, R., Mokos, J., Lipschutz, M. E., Metzler, K., Suter, M., and Kubik, P. W.: Allan Hills 88019: An Antarctic 300 H-chondrite with a very long terrestrial age, *Meteorit. Planet. Sci.*, 32, 769-773, <https://doi.org/10.1111/j.1945-5100.1997.tb01567.x>, 1997.
- Shi, G., Li, Y., Jiang, S., An, C., Ma, H., Sun, B., and Wang, Y.: Large-scale spatial variability of major ions in the atmospheric wet deposition along the China Antarctica transect ($31^{\circ}\text{N}\sim 69^{\circ}\text{S}$), *Tellus B*, 64, 17134, <https://doi.org/10.3402/tellusb.v64i0.17134>, 2012.
- 305 Shi, G., Ma, H., Hu, Z., Chen, Z., An, C., Jiang, S., Li, Y., Ma, T., Yu, J., Wang, D., Lu, S., Sun, B., and Hastings, M. G.: Brief communication: Spatial and temporal variations in surface snow chemistry along a traverse from coastal East Antarctica to the ice sheet summit (Dome A), *Cryosphere*, 15, 1087-1095, <https://doi.org/10.5194/tc-15-1087-2021>, 2021.
- 310 Spaulding, N. E., Higgins, J. A., Kurbatov, A. V., Bender, M. L., Arcone, S. A., Campbell, S., Dunbar, N. W., Chimiak, L. M., Introne, D. S., and Mayewski, P. A.: Climate archives from 90 to 250 ka in horizontal and vertical ice cores from the Allan Hills Blue Ice Area, Antarctica, *Quaternary Res.*, 80, 562-574, <https://doi.org/10.1016/j.yqres.2013.07.004>, 2013.
- 315 Turney, C. S., Fogwill, C. J., Golledge, N. R., McKay, N. P., van Sebille, E., Jones, R. T., Etheridge, D., Rubino, M., Thornton, D. P., and Davies, S. M.: Early Last Interglacial ocean warming drove substantial ice mass loss from Antarctica, *Proc. Natl. Acad. Sci.*, 117, 3996-4006, <https://doi.org/10.1073/pnas.1902469117>, 2020.
- Xiao, C., Li, Y., Hou, S., Allison, I., Bian, L., and Ren, J.: Preliminary evidence indicating Dome A (Antarctica) satisfying preconditions for drilling the oldest ice core, *Chin. Sci. Bull.*, 53, 102-106, <https://doi.org/10.1007/s11434-007-0520-6>, 2008.
- 320 Yan, Y., Bender, M. L., Brook, E. J., Clifford, H. M., Kemeny, P. C., Kurbatov, A. V., Mackay, S., Mayewski, P. A., Ng, J., Severinghaus, J. P., and Higgins, J. A.: Two-million-year-old snapshots of atmospheric gases from Antarctic ice, *Nature*, 574, 663-666, <https://doi.org/10.1038/s41586-019-1692-3>, 2019.
- 325 Zappala, J. C., Baggenstos, D., Gerber, C., Jiang, W., Kennedy, B. M., Lu, Z. T., Masarik, J., Mueller, P., Purtschert, R., and Visser, A.: Atmospheric ^{81}Kr as an Integrator of Cosmic-Ray Flux on the Hundred-Thousand-Year Time Scale, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2019GL086381>, 2020.