

A Semi-direct Effect from Northern California Biomass Burning Aerosols Contributes to Decreased Cloud Cover over the Region

James L. Gomez¹, Robert J. Allen¹, and King-Fai Li²

¹Department of Earth & Planetary Sciences, University of California Riverside, Riverside, USA

²Department of Environmental Sciences, University of California Riverside, Riverside, USA

Correspondence: James Gomez (jgome222@ucr.edu)

Abstract. Wildfires in the southwestern United States, particularly in northern California (nCA), have grown in size and severity in the past decade. As they have grown larger, they have been associated with large emissions of absorbing aerosols and heat into the troposphere. Utilizing satellite observations from MODIS, CERES, and AIRS, as well as reanalysis from MERRA-2, the meteorological effects of aerosols associated with fires during the wildfire season (June-October) were discerned over the nCA-NV (northern California and Nevada) region in the 2003-2022 time period. Wildfires in the region have a higher probability of occurring on days of positive temperature T anomalies and negative relative humidity RH anomalies, making it difficult to discern the radiative effects of aerosols that are concurrent with fires. To better isolate the effects of large fire emissions on meteorological variables, such as clouds and precipitation, variable anomalies on high fire emission days (90th percentile) were compared to low fire emission days (10th percentile) and were further stratified based on whether surface relative humidity RH_s was anomalously high (75th percentile) or low (25th percentile) compared to typical fire season conditions. Comparing the high fire emission/high RH_s data to the low fire emission/high RH_s data, positive tropospheric T anomalies were found to be concurrent with positive AOD anomalies, which was associated with significant negative 850 hPa-300 hPa RH anomalies during both 75th percentile RH_s conditions. Furthermore, high fire emission days under high RH_s conditions are associated with negative CF anomalies that are concurrent with the negative RH anomalies, which is consistent with the traditionally-defined aerosol-cloud semi-direct effect. This negative CF anomaly is associated with a significantly negative regional precipitation anomaly and an positive net top of atmosphere radiative flux anomaly (a warming effect) in certain areas. The T , RH , and CF anomalies under the high fire emission/high RH_s conditions compared to low fire/high RH_s conditions correlate significantly spatially with AOD anomalies. Additionally, the vertical profile of these variables under the same stratification are consistent with positive black carbon mass mixing ratio anomalies from MERRA-2. As a result, large fires in California may create a positive feedback that could intensify fire weather, and therefore extend fire lifetime and impacts.

1 Introduction

As a result of climate change, land use change, and forest management, the frequency of wildfires in California has trended upward from 100 fires per year in the 1920s to 300 fires per year in the late 2010s (Li & Banerjee, 2021). The size of these wildfires has also increased, with total burned area (square distance burned by a fire) increasing from roughly 1000 km² to

25 almost 4000 km² in the same time period (Li & Banerjee, 2021). According to a recent study, frequency of extreme daily wildfire events in the region are projected to increase by 59%-172% in coming years due to climate change (Brown et al., 2023), which is consistent with findings of numerous other studies (Palinkas, 2020; Ager et al., 2021; United Nations Environment Programme, 2022). In both higher and lower CO₂ mitigation scenarios, large wildfire events are projected to become more commonplace by the end of the 21st century worldwide, as well in the southwestern US (United Nations Environment Programme, 2022). Large wildfire events in the late 2010's and early 2020's were associated with more intense "fire weather": high temperatures T , low relative humidity RH , and high surface wind speeds U_s (Varga et al., 2022; Keeley & Slyphard, 2019). These fire weather conditions may be potentially intensified, or alleviated, by the fires themselves. Higher burn severity wildfires, such as the 2020 wildfires in California (CA), have been observed to inject smoke plumes higher into the troposphere than in previous years (Wilmot et al., 2022). These smoke plumes consist of both shortwave (SW) absorbing aerosols such as black carbon BC and reflective aerosols such as organic aerosol (OA), as well as brown carbon, which is absorbing and reflective. Additionally, wildfires have also been associated with emission of other aerosol species through feedbacks. While dust is not emitted from biomass burning, a number of studies have linked fires to concurrent dust emission through creation of convective updrafts (Wagner et al., 2018, 2021) and delayed dust emissions through wildfire clearing of vegetation (Wagenbrenner et al., 2013, 2017; Yu & Ginoux, 2022). The absorbing properties of wildfire smoke and co-emitted dust over the western US, measured using absorbing aerosol optical depth (AAOD), is uncertain. However, a recent study of CA fires indicates that wildfires increase AAOD relative to the annual mean by tenfold (Cho et al., 2022). An injection of absorbing aerosols into the troposphere may cause a local warming affect, altering the hydrological and radiative balance of the atmosphere (Allen & Sherwood, 2010; Thornhill et al., 2018; Allen et al., 2019; Herbert & Stier, 2023). Smoke plumes that reach the upper troposphere (pressures < 500 hPa) may deposit absorbing aerosols that could burn off high clouds, and promote more stable low clouds (Stjern et al., 2017; Smith et al., 2018; Allen et al., 2019), leading to SW and longwave (LW) cooling. Alternatively, if the absorbing aerosols are concurrent with low clouds, the relative humidity of the liquid cloud layer would be decreased, burning off low clouds and leading to a decrease in outgoing SW flux (Koch & Del Genio, 2010; Allen & Sherwood, 2010). These are both examples of aerosol semi-direct effects. Past observations and modelling experiments have shown dust aerosol is associated with semi-direct effects (Tsikerdekis et al., 2019; Amiri-Farahani et al., 2017; Helmert et al., 2007), as dust also has SW absorbing properties (Highwood & Ryder, 2014; Kok et al., 2023). Furthermore, the higher altitude of absorbing aerosol from California fires may alter cloud microphysics, which also has the potential to change the radiative balance of the surface and atmosphere. An influx of aerosols into the troposphere may create an abundance of cloud condensation nuclei (CCN) for droplets to condense onto, decreasing effective radius R_{eff} of the clouds, an effect already observed with smoke (OA and BC) particles in the northwestern US (Twohy et al., 2021). A decrease in R_{eff} would increase the albedo of the clouds, assuming constant water path, which would then increase outgoing SW radiation. This decrease in R_{eff} can also affect liquid water path LWP as the smaller droplets can evaporate much faster than larger droplets, or the smaller droplets can suppress precipitation, which increases LWP by reducing the liquid water leaving the cloud (Goren & Rosenfeld, 2012). The lighter droplets can also be lofted higher in the atmosphere, where they condensate further and release latent heat, then eventually fall from this greater height and evaporate. Therefore, to compensate, polluted clouds have more intense updrafts and downdrafts

60 than pristine clouds (Khain, 2009). SW absorption itself can also decrease precipitation P in other ways, such as reducing SW radiation reaching the surface or through rapid atmospheric adjustments (Sand et al., 2020; Samset, 2022; Allen et al., 2023).

Large Fires are not only limited to the western US. Australia, the Mediterranean Basin, and South America have all experienced an increase in large fire events due to climate change and land management (Shi et al., 2021; Ruffault et al., 2020; Artaxo et al., 2013; Allen et al., 2024a). As the western US, and other parts of the world, enter this new regime of large fires, there comes a need for improved understanding of the effects of aerosols emitted primarily (through biomass burning), secondarily (oxidation of emitted volatiles), or through feedbacks (such as dust emissions concurrent with fires) by wildfires. Models participating in the Coupled Model Intercomparison Project version 6 (CMIP6) (Eyring et al., 2016) do not have parametrizations of biomass burning (BB) aerosol emissions that respond to CO₂ emissions in most of their experiments, including the DECK (Diagnosis, Evaluation, and Characterization of Klima) experiments (Gomez et al., 2023). The models that have interactive BB aerosol emissions tend to parameterize them as a function of fuel flammability (temperature and moisture), fuel density, and plant functional type (Mangeon et al., 2016; Li et al., 2019). Most models participating in CMIP6 do not have dynamic vegetation models (Li et al., 2019), and therefore are incapable of incorporating fire-dust feedbacks. Instead, modelers rely on prescription of BB aerosols in most experiments.

75

Recent modelling experiments have found significant effects of wildfires on regional and global climate scales. Previously, using prescribed aerosol simulations in the Community Earth System Model version 2 (CESM2), it was shown that the large 2019 wildfires in Australia could have intensified that year's La Niña through aerosols directly cooling the ocean surface (Fasullo et al., 2021). Another CMIP6 study observed a similar effect on La Niña as a result of a teleconnection caused by an influx of absorbing aerosols into the atmosphere from South African wildfires (Amiri-Farahani et al., 2020). Biomass burning aerosols may also have other effects on large scale ocean circulation, such as an invigoration of the Atlantic Meridional Overturning Circulation (Allen et al., 2024b). As far as the southwestern US is concerned, a modeling experiment using the WRF/CHEM model was run to analyze the effects of a wildfire event on weather forecasts (Chen et al., 2014). This study found that the BB aerosols suppressed convection, prevented cloud formation, and decreased precipitation. While studies such as these demonstrate that it is possible to model past effects of fires on local and global climate, without parameterization of BB aerosol emission, as well as parametrization of secondary dust aerosol emission from wildfire-cleared vegetation, the radiative forcing of future fires' primary and secondary aerosols will remain a source of uncertainty. Furthermore, there are few, if any studies, that attempt to discern the impacts of large fires over the southwestern US. Twohy et al. (2021) analyzed satellite observations of cloud microphysical properties over part of the region, however this study was conducted during only one wildfire event in 2018. As a result, there is no comprehensive long-term observational study over the southwestern US concerning wildfire aerosol-cloud interaction. Therefore, to further understand the effects of wildfires on the climate of one of the most populated areas in the US, this paper aims to quantify the radiative as well as microphysical effects that these aerosols have in the region under different atmospheric conditions utilizing satellite data.

2 Satellite and Reanalysis Datasets

95 The objective of this study is to quantify the impacts of wildfire aerosol emissions on meteorological parameters, such as clouds and precipitation, over the southwestern US using observations. This includes the Aqua satellite with the MODIS, AIRS and CERES instruments. The Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2) reanalysis project (Randles et al., 2017; Global Modeling And Assimilation Office & Pawson, 2015) is used to obtain daily black carbon mass mixing ratio vertical profiles. Fire dry matter emission data DM are used as a proxy for fire severity, and are derived from
100 the Global Fire Emissions Database version 4 (GFED4) (van der Werf et al., 2017; Randerson et al., 2017). All data sets are globally gridded observational data sets, with the exception of GFED4 and MERRA-2 which are considered globally gridded reanalysis datasets.

2.1 Global Fire Emissions Database (GFED4)

GFED4 DM emissions are calculated in the Carnegie–Ames–Stanford Approach (CASA) model, which requires MODIS
105 burned area data, meteorological data from the ERA-Interim reanalysis dataset, photosynthetically active radiation data based on Advanced Very high-resolution Radiometer satellite instrument retrievals, and vegetation continuous fields data from the MODIS MOD44B dataset (van der Werf et al., 2017). DM is the emission of any gas or aerosol from burned vegetation, and a list of all these types of emission can be found in van der Werf et al. (2017). The CASA model is run using burned area data from combined MODIS-Aqua and MODIS-Terra level 3 data (MCD64A1). Wildfire studies tend to either use fire power
110 (from MODIS or VIIRS) or burned area-based datasets to quantify fire severity. Burned area is determined by MODIS from a time series of burn sensitive vegetation index, which compares daily surface reflectances (Giglio et al., 2018). Fire power is the radiated energy from fires over time, and MODIS determines this quantity by comparing the brightness temperature of a fire pixel to the background brightness temperature (Peterson et al., 2013). Use of a burned area-based dataset is preferable to a fire power dataset for this paper, as cloud cover may obstruct fire power data retrievals, leading to an underestimation of fire
115 size/severity in a given time period. While cloud cover can also block burned area retrievals, burned area can be recorded once cloud cover has been dissipated, unlike fire power. This introduces a temporal uncertainty, however. This temporal uncertainty is ± 1 day for clear sky conditions, ± 5 days under consistent 75% cloud cover, and up to ± 20 days over persistently very cloudy (85% or higher) intervals (Giglio et al., 2013). However, this temporal uncertainty is likely of little significance for this paper, as cloud cover over the western US during the wildfire season is rarely persistently high (aside from "June gloom"
120 in coastal regions), and the lifetime of biomass burning aerosols (roughly 4-12 days) is generally greater than or equal to the temporal uncertainty of clear sky or consistently cloudy burned area data (Cape et al., 2012). The daily underestimation of fire power is demonstrated in **Figure S1**, which indicates that Aqua fire power retrievals, taken from the MYD14A1 dataset (Giglio & Justice, 2015), underestimate daily fire severity compared to DM with 98% of days reporting a lower normalized fire power than normalized DM . Therefore, for fire power to be a more useful metric, a daily combined Aqua/Terra/VIIRS dataset would
125 have to be used, which is not available for the time period of interest. GFED4 fire emissions are also preferred over fire power data and raw burned area data as calculation of fire emissions takes vegetation type and net primary production into account.

Raw burned area and fire power datasets yield information about fire size and intensity, but as aerosol emission also depends on the type of vegetation being burned, use of either dataset over a fire emissions dataset may under-estimate or over-estimate biomass burning aerosol impacts on clouds. However, use of GFED4 data has drawbacks. While use of burned area data reduces the chance of an underestimation of fire impacts, the previously mentioned temporal uncertainty is introduced. Additionally, the CASA model itself is associated with uncertainties. Calculation of net primary production in the model, for example, does not take meteorological variables into account (Liu et al., 2018). As a result, caution must be taken when analyzing the results. To ensure results are robust, the GFED4 *DM* stratification method was verified by analyzing MODIS *AOD* anomalies (see Section 2.2) during large fire events (Section 3.3, Section 4.2), and by performing cross correlations between *AOD* and *DM* (Supplement section 1, **Figure S2**). GFED4 emissions and burned area data are available from 1997-2016. Data for 2017-2022 is also available, but the data is in "beta" and therefore is more limited. Both the complete and the beta data contain total carbon emissions, as well as dry matter emission. GFED4 also estimates the contribution of 6 different types of vegetation biomes (boreal forest, temperate forest, grassland, agriculture, and peat) to the carbon and dry matter emissions. However, the beta dataset only estimates these contributions for *DM*. Therefore, *DM* is used as a proxy for the severity of a given fire's emissions, as it is the only variable that both the complete and beta data contain and speculate. All observational datasets utilized in this study have a 1° resolution, however GFED4 emission data is of a 0.25° resolution. Therefore, this data was regridded to a 1° grid. It should be noted that GFED5 has recently been released (Chen et al., 2023), however this dataset was not used as it does not yet include emissions, only has data available up to 2020, and was released after analysis for this paper had concluded.

2.2 Aqua

MODIS-Aqua: Cloud and aerosol optical depth (*AOD*) data were derived from Moderate Resolution Imaging Spectroradiometer (MODIS) level 3 data. Specifically, the MODIS collection 6.1 1° level 3 product (MYD08_D3) (Platnick et al., 2003; Salomonson et al., 2002; MODIS Atmosphere Science Team, 2017) is utilized, which yields daily retrieval products from the Aqua satellite. The Aqua satellite makes two overpasses for the region of interest: one ascending run from 2-3 PM, and one descending run from 2-3 AM. The descending dataset is used as most MODIS level 3 cloud property products provided are descending (morning) only. For MODIS cloud retrievals during periods of large *AOD*, especially when the aerosols are concurrent with clouds, it is possible for MODIS to misidentify aerosols as clouds (Herbert & Stier, 2023). This may cause errors in cloud property retrievals, as well as an overestimation of cloud fraction *CF*. This may lead to overestimation of *CF* during anomalously large fire events. While the MODIS Dark-Target and Deep Blue *AOD* algorithms are extensively quality controlled and evaluated (Levy et al., 2013; Platnick et al., 2017; Wei et al., 2019), there is still room for errors in *AOD* and cloud retrieval. Additionally, as it is not possible to distinguish wildfire *AOD* from other *AOD*, whenever possible, fire emissions from GFED4 are used to discern the impacts of fires on cloud properties.

AIRS: Data concerning T , water mass mixing ratio M_{H_2O} , *CF*, and *RH* profiles, as well as surface temperature T_s and surface relative humidity RH_s , were derived from Atmospheric Infrared Sounder (AIRS) level 3 daily data (AIRS3STD) (AIRS

160 Science Team & Teixeira, 2013). As with the MODIS data, the descending dataset is used.

CERES: Top of atmosphere as well as in-atmosphere radiative flux data was derived from Clouds and the Earth's Radiant Energy System (*CERES*) level 3 daily 1 degree Synoptic product (SYN1deg-Day) (Doelling, 2016, 2017, 2023). This is a combined Terra and Aqua dataset from 2002-2021, and for 2022 it is a combined Terra and NOAA-20 dataset. This *CERES* dataset combines cloud data from MODIS/VIIIRS, aerosol data from GEOS, and top of atmosphere radiative flux data from *CERES* to produce all-sky, clear-sky, and aerosol-free radiative flux profiles.

2.3 GPCP Combined Precipitation Dataset

P data for this project was derived from the daily Global Precipitation Climatology Project (GPCP daily) Climate Data Record, Version 1.3 dataset (Huffman et al., 2001; Adler et al., 2018). GPCP combines satellite observations as well as rain gauge data to produce 1° daily precipitation amount data.

2.4 MERRA-2 Aerosol Profiles

Daily vertical black carbon aerosol mass mixing ratio profiles are derived from the M2I3NVAER data product (Global Modeling And Assimilation Office & Pawson, 2015; Buchard et al., 2015). This product estimates aerosol profiles by assimilating MODIS *AOD* into the GEOS5 model, which is radiatively coupled to the Goddard Chemistry, Aerosol, Radiation, and Transport (GOCART) aerosol module. The GOCART model includes biomass burning emissions from the NASA Quick Fire Emission Dataset (QFED) version 2.1, which provides daily biomass burning aerosol estimates (Buchard et al., 2015). These profiles were then validated using ground and satellite observations of aerosol profiles. This dataset has been previously used to determine effects of wildfire aerosols in other parts of the world (Raga et al., 2022; Nguyen et al., 2020). The aerosol profiles are archived in a high-resolution hybrid sigma pressure grid, and therefore must be interpolated into 1 degree grid cells, and converted into traditional pressure levels. For the purposes of this paper, only the black carbon variables are analyzed. MERRA-2 separates *BC* into two types: hydrophobic black carbon BC_{pho} and hydrophilic black carbon BC_{phi} .

2.5 CALIPSO

The Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation satellite (CALIPSO) provides observations of aerosol extinction coefficient profiles. MERRA-2 profiles are utilized in the main analysis instead of CALIPSO profiles as CALIPSO data is not temporally consistent with the other datasets utilized in this paper, and because the gridded data is low resolution. More information on CALIPSO can be found in Supplement Section 2.

3 Methods

3.1 Statistics

The bulk of the analysis for this paper involves empirical cumulative distribution functions (CDFs). Empirical distribution functions are calculated for each variable of interest under differing fire and meteorological conditions, and the shift in each distribution is compared. Plotting two CDFs on the same axis allows for comparison on how likely an anomaly is to be positive or negative under differing circumstances, such as how likely a positive/negative anomaly for a certain variable is to occur during a high (90th percentile) fire dry matter emission (*DM90*) or low (10th percentile) fire dry matter emission (*DM10*) event. The 90th percentile is chosen as the purpose of this paper is to analyze the effects of large fire events on climate, not the effects of fires in general. From the calculated normal distributions, the effect size of one variable's distribution on another variable's distribution are estimated using Cohen's *d*. *d* is an approximation of by how many standard deviations σ s the distribution shifts in response to a change in a variable. In this paper, *d* is calculated to determine the effect size of *DM* on other variables. *d* is approximated using

$$d = \frac{\bar{a} - \bar{b}}{0.5\sqrt{\sigma_a^2 + \sigma_b^2}} \quad (1)$$

where \bar{a} is the mean of the (*DM90*) group (group *a*), and \bar{b} is the mean of the (*DM10*) group (group *b*), σ_a is the standard deviation of group *a*, and σ_b is the standard deviation of group *b*. $d=0.2-0.5$ is considered to be a weak effect, $d=0.5-0.8$ is a moderate effect, and $d=0.8$ or higher is classified as a strong effect.

When comparing two data sets, a two-tailed pooled t-test is used to assess significance, where the null hypothesis of a zero difference is evaluated, with n_1+n_2-2 degrees of freedom, where n_1 and n_2 are the number of elements in each data set respectively. Here, the pooled variance

$$s^2 = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2} \quad (2)$$

is used, where S_1 and S_2 are the sample variances. For the purposes of this project, the t-test is evaluated at 90% significance.

3.2 Data Stratification and Comparison

In section 3.1, it was mentioned that CDFs for variable anomalies during anomalously high and low *DM* emission events are generated to discern to what degree fires impact these anomalies. The purpose of this stratification, particularly stratification of days into anomalously high and low fire events, is to isolate the effects of fires on clouds and/or weather. The remainder of this section will detail how data stratification is accomplished. First, a variable is chosen for analysis (such as *CF*). Next, this variable as well as the variable(s) that are used to stratify the variable are filtered to include only the region of interest. As the Aqua satellite does not record data for each gridcell at every time step, wherever a coordinate (latitude,longitude,time) is missing a value for a specific variable, the variable(s) it is being stratified by also has the value at that coordinate replaced

by a missing value (and vice-versa). Next, to focus on potential feedbacks fires may have on land, a land-sea mask is applied. Then, the daily regional anomaly for each variable is taken. Then, the 2003-2022 wildfire seasons are spliced together, which results in a roughly 3060-day time series. From this 3060 day time series, any days with no data are removed. Next, the average
220 of each day of the wildfire season is removed from each data point in the distribution to give a time series of anomalies for each variable. Booleans that filter out days above or below a certain percentile for the stratification variables are then applied simultaneously. For each dataset, an empirical CDF is then calculated. Then, the means are differentiated from each other to determine if the stratification variable (such as *DM*) leads to a significant change in the variable anomaly in question. This process can be applied both for a regional average, or on a gridcell-by-gridcell basis. When this process is performed
225 on a gridcell-by-gridcell basis, the Pearson cross correlation coefficient r is determined by spatially correlating the stratified variables with one another. This helps determine if one change in a variable as a result of fires (or other factors) feedbacks onto another to cause a change in anomaly. **Figure 1** serves as a verification of the stratification method, as well as validation of GFED4 emissions data. Monthly cross correlation analysis (Supplement Section 1, **Figure S2**) as well as previous works (Wilmot et al., 2022; Schlosser et al., 2017; Cho et al., 2022) indicate that during large fire events, *AOD* and/or particulate
230 matter concentration are significantly larger compared to no fire conditions. The significant increase in *AOD* over most of the southwestern US supports the assertion that GFED4 fire emissions are an acceptable indicator of large fire occurrences.

3.3 Regions of Interest

First, the region within the southwestern US in which the most significant fire emissions originate was discerned. Based on what is generally considered to be the time of year in which most wildfires occur in the western US (Urbanski, 2013; Urbanski
235 et al., 2011), data was collected from June 1st-October 31st for the 2003-2022 time period. 2003-2022 was chosen as this is the time period in which Aqua satellite data is available for the fire season. Analysis was limited to fire seasons as opposed to the entire year so that the threshold for what constitutes a 90th percentile fire is increased. First, for each gridcell, the 2003-2022 seasonal average daily *DM* emissions was taken. The portion of the southwestern US that had the largest 2003-2022 seasonal average daily *DM* emissions is the region that shall be referred to as "northern California" (nCA), which is highlighted in the
240 blue box in **Figure 1a**. The reason for limiting *DM* data to this region is again to ensure that the threshold for 90th percentile *DM* is kept high. The nCA region is characterized by temperate forests along the coastline, in the far north, as well as the east. Agricultural lands are scattered throughout about every gridcell in nCA, with higher concentrations in the central valley as well as the coastal north. Grasslands are also found throughout most grid cells in this region, with higher concentrations in central CA. The dominant contributor of *DM* in this region is the temperate forests in the north (**Figure S3**). At this time of
245 year, predominant wind patterns in nCA would favor transportation of smoke from these fires to northern Nevada. During the fire season, northwesterlies tend to blow across nCA towards northern Nevada, and south westerlies blow through the central valley and Sierra Nevada range (Zaremba & Carroll, 1999; LeNoir et al., 1999). Therefore, the expectation is for the majority of wildfire aerosols to be concentrated in nCA, and neighboring northern/central Nevada. In differentiating *AOD* anomalies on high nCA *DM* days and *AOD* anomalies on low nCA *DM* days, *AOD* is found to be anomalously positive in both nCA
250 and Nevada (**Figure 1b**), confirming this hypothesis. However, there are also significant *AOD* anomalies throughout the entire

region. For reasons that will be explained in **Section 4.1**, the main analysis will still be relegated to northern CA and Nevada. From this point forward, the focus will be on the effects of the fires in the blue box in **Figure 1a** (nCA) on the area highlighted in the green box (nCA-NV) in **Figure 1b**.

3.4 Heating Rate

255 Aerosol shortwave heating rate of the atmosphere SWH_{aer} was calculated using

$$\frac{\partial T}{\partial t} = SWH_{aer} = \frac{g}{c_p} \cdot \frac{\Delta F_{aer}}{\Delta p} \quad (3)$$

where t is time in days, g is gravity, c_p is the heat capacity at constant pressure, F_{aer} is the shortwave radiative effect of the aerosols, and p is pressure. F_{aer} itself was derived from the CERES SYN1deg-Day downward and upward shortwave radiative fluxes. F_{aer} between two atmospheric layers is given by

$$260 F_{aer} = SWd_1 - SWu_1 - (SWd_2 - SWu_2) \quad (4)$$

where SWd_1 denotes downward shortwave flux at the higher layer, SWu_1 denotes upward shortwave flux at the higher layer, SWd_2 denotes downward shortwave flux at the lower layer, and SWu_2 denotes upward shortwave flux at the lower layer.

4 Results

4.1 High & Low Surface Relative Humidity Stratification

265 The fingerprints of a traditionally-defined semi-direct effect where aerosols coincide with clouds would entail an anomalous warming of the cloud layer, and a corresponding decrease in RH . However, the meteorological conditions around which fires tend to occur need to be considered. As previously stated, large fires tend to occur during fire weather, which includes hot, dry, and windy conditions (Varga et al., 2022). Hot and dry conditions themselves are associated with high pressure anomalies in this region (**Figure S4**). Therefore, these fire weather conditions need to be "filtered out" to determine any potential semi-
 270 direct effects. Therefore, in addition to DM , variables need to be stratified by a second variable to account for the influence of meteorology on P , CF , and cloud properties. Fire season data was stratified by high (75th percentile) vs low (25th percentile) T_s , RH_s , U_s , and surface pressure to determine which variable was associated with the largest DM , and successfully filtered out fire weather condition anomalies. The 75th/25th percentiles were chosen for the potential second stratification variables as opposed to extremes (90th/10th percentiles) to ensure a robust number of data points, and to have a dataset that is more
 275 representative of common conditions in the region. **Figure 2** depicts CDFs for meteorological conditions and DM under high RH_s extremes (RH_s75) and low RH_s extremes (RH_s25) in the entire southwestern US. RH_s was chosen as the second stratification variable, as stratifying nCA DM by high (RH_s75) and low RH_s conditions (RH_s25) and differentiating the means of these distributions yields a significant DM anomaly of $\Delta DM = -1.04e-4 \pm 3.5e-5 \text{ kg m}^{-2} \text{ day}^{-1}$. The absolute value of this anomaly is an order of magnitude higher than the differences in mean DM between high and low conditions of

280 the other potential stratification variables (surface pressure, T_s , and U_s) (**Figure S4, Figure S5, and Figure S6**). This indicates that fire occurrence/fire emission are more dependent on RH_s than these other fire weather variables. Low RH_s extremes in the southwestern US are associated with significantly higher T throughout the troposphere/surface, significantly reduced RH throughout the troposphere/surface, and significantly lower CF , while high RH_s extremes are associated with the opposite (**Figure 2**). This demonstrates a need to separate the effects of fires from the meteorological effects of low RH_s extremes, as positive DM anomalies are significantly more likely to occur on (RH_s25) days as opposed to (RH_s75) days, as which is expected as moisture, and moist plants, suppress the ability of fires to grow and be maintained (Minnich & Chou, 1997; Ford & Johnson, 2006). The immediate direct effect of BB aerosols tends to be a net cooling of the surface (Sakaeda et al., 2011; Abel et al., 2005). However, certain semi-direct effects, such as the burning off of low clouds, may overpower this effect, leading to a net surface warming. As the meteorological conditions associated with low RH_s days are also hallmarks of a semi-direct effect (**Figure 2**), from here onward data will be stratified into four categories: one with high DM and high RH_s ($DM90, RH_s75$), one with low DM and high RH_s ($DM10, RH_s75$), one with high DM and low RH_s ($DM90, RH_s25$), and one with one with low DM and low RH_s ($DM10, RH_s25$). In differentiating the average of the variables on ($DM90, RH_s75$) days and ($DM10, RH_s75$) days, the effects of fires can be discerned independent of the meteorological conditions that come with high DM extremes. Additionally, comparing the ($DM90, RH_s25$) dataset to the ($DM10, RH_s25$) potentially allows the effect of 295 fires to be discerned during fire weather, as the meteorological conditions in both distributions should not be significantly different from one another. **Figure 3** demonstrates that the effect of fires on AOD under both high and low RH_s stratifications is significantly positive anomalies in the nCA-NV region. The increase in mean AOD is larger under low RH_s at 0.24 ± 0.04 . The corresponding change under high RH_s is 0.13 ± 0.05 . As the AOD is consistently significant only in the nCA-NV region under both stratifications, this region will be the focus of the study.

300 **4.2 Vertical Distribution of Black Carbon and Absorption in nCA-NV Region**

Freshly emitted BC is highly hydrophobic, and as it ages it becomes less resistant to accumulating water droplets (Lohmann et al., 2020). BC has an average lifetime of 1 week (Lohmann et al., 2020), and the aging process begins after 1-2 days (He et al., 2016). Furthermore, in a region with such low fire season wet deposition such as the southwest US, the BC on average can live much longer than one week (Ogren & Charlson, 1983). Therefore, hydrophobic and hydrophilic BC profiles are important to differentiate because they can give an idea of how long the BC stays in the atmosphere, and it hints at how much BC can contribute to indirect and semi-direct effects. **Figure 4** displays high compared to low DM mass mixing ratio anomalies for BC_{phi} , BC_{pho} , and combined BC on high and low RH_s days. Significant positive anomalies of BC mass mixing ratio are present from 950Pa-300 hPa for all types of BC under both ($DM90, RH_s75$) and ($DM90, RH_s25$) conditions compared to the corresponding low fire conditions. The most significant increase in BC is from about 950-600 hPa for the ($DM90, RH_s75$) days, and from 950-550 hPa for the ($DM90, RH_s25$) days. Comparing the MERRA-2 BC profiles to the CALIPSO $DM90$ - $DM10$ months 2006-2021 smoke aerosol daytime and nighttime extinction coefficient profile, MERRA-2 places more absorbing aerosol below 700 hPa, while CALIPSO generally places more absorbing aerosol above 700 hPa (**Figure S7**). Therefore, it is important to note that CALIPSO profiles do not agree with MERRA-2 when it comes to the 310

positioning of the smoke in the troposphere. However, as the MERRA-2 and CALIPSO profiles are not temporally consistent, the comparison between these profiles is not 1-1. Additionally, as the CALIPSO profiles are not temporally consistent with the rest of the data in this paper, their use is not preferred over the MERRA-2 profiles.

There is roughly an equal amount of BC_{phi} and BC_{pho} during both high and low RH_s days, indicating that on these days there is roughly as much fresh and aged aerosol in the troposphere. This is important as the quantity of BC_{pho} indicates that microphysical effects are possible as it suggests a large amount of CCN are present in the troposphere. Additionally, the presence of aged BC indicates that the BC can affect the atmosphere radiatively over the course of multiple days. To estimate the impact of these aerosols on the troposphere over time, a SWH_{aer} profile was created from CERES radiative flux data (**Figure 5**). Shortwave profiles used to generate these heating rate profiles, along with LW profiles, can be found in **Figure S8**. Under both ($DM90, RH_s75$) (**Figure 5a**) and ($DM90, RH_s25$) (**Figure 5b**) RH_s conditions compared to the corresponding low DM conditions, there is a positive SWH_{aer} anomaly from 850 hPa to the next highest pressure level in the CERES dataset, 500 hPa. For high RH_s , this corresponds to a heating rate of $SWH_{aer} = 0.041 \pm 0.016$ K day⁻¹, and for low RH_s this corresponds to a heating rate of $SWH_{aer} = 0.093 \pm 0.019$ K day⁻¹. A highly idealized prior modelling study on the shortwave absorbing properties BC globally found that a heating rate of about 0.8 K day⁻¹ from increasing BC tenfold was associated with around 0.8 K of warming (Stjern et al., 2017). Assuming linearity, the 0.04-0.09 K day⁻¹ heating rate observed in **Figure 5** would therefore correspond to around a 0.4-0.9 K temperature anomaly. Spatially, the 850 hPa-500 hPa heating rate is significant over almost all grid cells in the region of interest where there is data, with the most positive heating rates over eastern nCA and eastern Nevada (**Figure S9**).

4.3 Responses in Temperature, Humidity, & Cloud Profiles

Figure 6 displays 2003-2022 June-October nCA-NV vertical profiles of the fire effect on T (**Figure 6a,e**), and RH (**Figure 6c,g**) profiles. **Figure 6a-d** are stratified by high RH_s , while **Figure 6e-h** are stratified by low RH_s . In both **Figure 6a** and **Figure 6e**, the temperature anomalies in the 850 hPa to 300 hPa pressure level range are consistently significant and positive at around 1 K. Since the aerosol heating rate accounts for approximately 0.4-0.9 K of warming, the semi-direct effect of these aerosols accounts for about 40-90% of this anomaly. Comparing **Figure 6** to **Figure 4**, the positive differences in temperature anomaly are generally consistent with the positive BC anomalies. Under both high RH_s (**Figure 6c**) and low RH_s (**Figure 6g**) conditions, RH anomalies throughout the entire profiles are negative but are only consistently significant during high RH_s extremes. The AIRS CF profile under high RH_s conditions (**Figure 6d**) demonstrates significant negative anomalies from 300 hPa-600 hPa that are consistent with significant negative RH anomalies and significant positive T anomalies. However, there is an increase in CF at 850 hPa (**Figure 6d**). This pressure level corresponds to the highest concentration of BC_{phi} (**Figure 4c**), and perhaps this indicates at this pressure level there is cloud seeding occurring. For the low RH_s profile, there is only a significant negative cloud anomaly close to the surface at 925 hPa (**Figure 6h**).

Aside from temperature, the other potential factor that could affect RH is that of specific humidity, which is analogous to water mass mixing ratio M_{H_2O} . **Figures 6b,f** depicts the effect of fires on M_{H_2O} anomalies under high RH_s and low RH_s conditions respectively. There is no significant anomaly under high or low RH_s conditions but is consistently positive at 700 hPa and below. Furthermore, the changes in the RH profile follow the changes in the T profile as opposed to the M_{H_2O} profile, implying the positive T anomalies generally dominate the change in RH anomalies.

While these profiles provide a general overview of how T , M_{H_2O} , RH , and CF are changing over the region of interest, it is important to determine if these changes are consistent spatially with one another, as well as whether the changes coincide with BC anomalies. As the T , RH , and CF anomalies are strongest during high RH_s days, the focus from here will be on the meteorological effects of high DM on high RH_s days. **Figures 7-11** depict the effect of fires on the spatial distributions of BC , T , M_{H_2O} , RH , and CF anomalies at each AIRS pressure level up to 200 hPa on under high RH_s conditions. The positive MERRA-2 BC anomalies in **Figure 7** correlate positively and significantly with MODIS AOD for each pressure level between 925 hPa-300 hPa (**Figures 7b-h**), and are spatially consistent with positive AIRS T anomalies (**Figure 8**). Shifting attention to **Figure 9**, there appear to be significant negative anomalies in M_{H_2O} in northeastern Nevada from 700 hPa-400 hPa, and significant positive anomalies over grid cells associated with large fires (**Figure 1a**) in the lower troposphere (925 hPa-850 hPa). Comparing these changes in T and M_{H_2O} spatially to changes in RH (**Figure 10**), it appears that changes in T tend to dominate changes in RH over CA, western NV, and southern NV while changes in M_{H_2O} appear to contribute to the negative RH anomaly in northeastern NV. Additionally, the positive M_{H_2O} at 850 hPa appears to mitigate the negative RH anomalies at the same level, which may explain why BC appears to be able to act as a CCN at this level but not others: RH does not decrease enough to prevent clouds from forming. The increase in M_{H_2O} has a myriad of possible explanations. It may be due to emission of moisture from the burned vegetation (Jacobson, 2014; Dickinson et al., 2021), from lofting of water vapor from the surface to higher levels of the atmosphere (Yu et al., 2024), or from moisture advection due to a change in wind vectors from the northeastern part of Nevada towards CA (**Figure S10**). Viewing **Figure 11c**, the increase in CF at 850 hPa appears to be driven predominantly by a few significant and large coastal CF anomalies. This indicates that there is an increase in shallow marine clouds at this pressure level, while clouds at other pressure levels are generally being suppressed. **Figure 11** demonstrates that significant negative CF anomalies are generally spatially consistent with negative RH anomalies from 700-400 hPa. The significant negative CF anomalies in northeastern Nevada that correspond with significant negative RH anomalies, but not significant positive T anomalies, at 700 hPa and higher indicate that the difference in clouds in this region is specific humidity dependent. This may be due to a transport of moisture outside of these grid cells due to anomalously positive southeastern wind speed anomalies in some of these grid cells (**Figure S10b**) that advect moisture towards southern California and southern Nevada, however further scrutiny is warranted to confirm this. It is not known if these wind speed anomalies are related to warming from the BB aerosols, or if these wind speed anomalies in this region are an artifact. Changes in wind vectors are further analyzed in supplement section 3. As the change in T is the more robust signal over all parts of the troposphere, the changes in T will be the focus of the remainder of the paper.

4.4 Changes in Cloud Type, Precipitation, and Shortwave Flux

With AIRS data indicating that large fires are associated with enhanced T , as well as lower RH and CF , it is essential to determine how liquid vs ice clouds are impacted, and what the corresponding impacts on P and radiative balance are. **Figure 12** displays CDFs for nCA-NV regional average variable anomalies during high DM /low RH_s days (solid red), low DM /high RH_s days (dashed red), high DM /high RH_s days (solid blue), and low DM /low RH_s days (dashed blue). **Figure 12a** and **Figure 12b** demonstrate that during high RH_s extreme days, the effect of fires on the liquid water cloud fraction CF_{lw} distribution and cirrus cloud fraction CF_{cir} distribution is a significant shift towards a preference for negative anomalies. The effect of the large fires creates an average -0.04 ± 0.02 CF_{lw} anomaly, and an average -0.05 ± 0.04 CF_{cir} anomaly under high RH_s conditions. In addition, MODIS total CF shifts by -0.07 ± 0.05 under the same stratifications. Precipitation also shifts significantly by -0.3 ± 0.23 mm day^{-1} . However, these shifts are significant only for high RH_s extreme days (**Figure 12**). The explanation why the distribution shifts farther towards negative anomalies when anomalously large fires occur during high RH_s compared to low RH_s extremes lies in **Figure 2**. During low RH_s days, RH throughout the troposphere is already significantly lower than normal conditions (**Figure 2e**), as temperatures throughout the troposphere are already high (**Figure 2c**) and atmospheric water vapor content is low. This creates conditions of negative CF anomalies (**Figure 2f**). Therefore, further increasing the already high T should not lead to significantly lower cloud fraction as RH is already low, and clouds require 100% RH to form. This can also be explained by the RH profile in **Figure 6g**, which demonstrates through most parts of the troposphere that RH is not significantly lowered due to fire effects. However, during low DM /high RH_s days, **Figure 12** demonstrates that conditions are favorable for clouds and rain. This is because during these high RH_s extremes, T is lower and RH is high. Therefore, when anomalously large fires introduce a positive T anomaly, the drop in RH is significant enough to reduce the chances of seeing positive cloud/precipitation anomalies. In response to the higher probability of negative cloud fraction anomaly, the probability that SW radiation will be reflected into space decreases. This reduction in top of atmosphere shortwave flux leads to a net increase in cloud only (all-sky minus clear-sky) top of atmosphere radiative forcing TOA_{cid} (**Figure 12f**). Though it should be noted that this increase is not significant, it is significant and positive over much of the region marked by a decrease in CF (**Figure 13e,h**), with a significant spatial cross correlation of $r = -0.67$. Regional all-sky SW and LW responses can be found in **Figure S11**.

Figure 13 displays composite differences between meteorological variables on high DM /high RH_s and low DM /high RH_s days for each gridcell over the entire southwestern US. **Figures 13a,b** display the composite differences in cloud layer ($850 \text{ hPa} \geq p \geq 300 \text{ hPa}$) temperature T_{CL} and cloud layer relative humidity RH_{CL} . These plots depict that T_{CL} significantly increases almost everywhere across California and Nevada, with the most significant increase in the green box (the nCA-NV region). The differences in T_{CL} correlate significantly with differences in AOD at $r = 0.70$ across the entire southwest. The decreases in RH_{CL} have a very similar spatial distribution to T_{CL} , with the strongest decreases in the nCA-NV region. Again, this correlates significantly with AOD with $r = -0.60$ over the entire southwest. The differences in all these variables across the southwestern US correlate significantly with AOD , supporting the assertion that aerosols concurrent with fires are associ-

415 ated with semi-direct effects. Of note are the changes in T_s and P , which are two variables intrinsically related to fire duration. Spatially correlating P with RH_{CL} yields a significant correlation of $r = 0.44$, implying that the reduction in P is related to the semi-direct effect. However, it should be noted that though the regional P anomaly is significant and negative, that it appears to be dominated by just strong changes in just a few gridcells. T_s correlates significantly with AOD over the southwestern US, with $r = 0.51$, and is generally spatially concurrent with increases in T_{cl} with $r = 0.72$. The equivalent for **Figure**
420 **13** for low RH_s days is given in **Figure S12**. Of note for this supplementary figure is that there are weak, but significant and widespread, negative CF , RH , and P anomalies over nCA and eastern Nevada, despite not being significant in the regional average (**Figure 12c,e**). This implies that the effects of the absorbing aerosols seen during high RH_s days are also prevalent on low RH_s days, but weaker and less widespread due to the lower availability of moisture.

425 While cross correlations indicate that there is a statistically significant relationship between fires and meteorology, practical significance needs to be established as well. The effect sizes of high DM emissions on nCA-NV regional averages of the variables in **Figure 12** and **Figure 13** are depicted in **Figure 14**. For high RH_s extremes (**Figure 14a**), the anomalously large fires have a moderate-to-strong effect size on most of the relevant variables. **Figure 14b** demonstrates that during low RH_s extremes, anomalously large fires have a weak-to-no effect size on the relevant variables, aside from T_{cl} in which fires have a
430 very strong effect size on.

4.5 Cloud Microphysical Effects

Up to this point, we have investigated aerosol direct/semi-direct effects on clouds. Aerosols may also influence clouds via microphysical effects, which are investigated in this section. High fire emissions under high RH_s conditions are associated with
435 non-significant differences in microphysical variables (**Figure 15**). Spatial maps of the fire effect on R_{eff} and LWP under high RH_s conditions show a mix of areas with positive and negative changes, most of which are not significant (**Figure S13**). Although there is a small tendency for negative R_{eff} anomalies to occur in Nevada and a small tendency for negative LWP anomalies to occur in nCA and western NV. Since negative R_{eff} anomalies can affect precipitation, the spatial distribution of R_{eff} anomalies (**Figure S13, Figure S14**) was compared to the spatial distribution of P anomalies (**Figure 13, Figure S12**)
440 under high compared to low DM conditions. Significant negative R_{eff} anomalies were not found to be spatially consistent with significant negative P anomalies under either high or low RH_s conditions. This casts doubt on wildfires in this region creating microphysical suppression of P .

There are significant regional changes in liquid R_{eff} and LWP under low RH_s conditions (**Figure 15, Figure S14**). Liquid
445 R_{eff} significantly increases under these conditions, which is contrary to what one would expect as a response to increased AOD (Twohy et al., 2021; Conrick et al., 2021; Fan et al., 2016). One possible explanation for this increase in R_{eff} is that R_{eff} is directly proportional to temperature (Martins et al., 2011), and perhaps the effects of the T anomaly dominate over the condensation of new droplets onto BC_{phi} . Alternatively, this increase may be driven by changes in atmospheric dynamics,

as increased updraft strength and enhanced turbulence could lead to increased coalescence (Khain, 2009). Coincident with the
450 strongest increase in R_{eff} (at the northernmost coast of California) under these conditions is a significant negative (upward)
pressure velocity anomaly from 1000 hPa-925 hPa, which implies that an increase in upward convection near the surface may
be a factor of the increase in R_{eff} , as an upward pressure velocity should increase droplet lifetime (**Figure S15**). It is also
noted that there are negative pressure velocity anomalies under high RH_s conditions from 1000 hPa-850 hPa (**Figure S16**),
and this corresponds with an increase in R_{eff} near the Bay Area.

455

The fire effect on LWP under low RH_s conditions is significant a decrease (**Figure 15c**). This significant negative LWP
anomaly may be due to the negative RH_{cl} anomaly (**Figure S12b**), as lower saturation of the air would reduce liquid water
within clouds. This decrease in LWP may be of importance, as LWP scales positively with cloud albedo (Han et al., 1998).
Therefore, this decrease in LWP may contribute to an increase in absorbed solar radiation at the surface. In summary, while the
460 nCA fires significantly inject aerosols into the troposphere, these aerosols do not appear to generally act as CCN, and instead
contribute to a positive T anomaly that burns off clouds. This may be because BC is generally more hydrophobic compared to
other aerosols, and instead the radiative effects of the aerosol dominate.

5 Discussion

The results of this paper indicate that large fires in nCA are concurrent with significant amounts of absorbing aerosols and a
465 warmer troposphere. When the fires occur during high RH_s conditions, this increase in T (**Figure 8, Figure 13a**) is associated
with a significant decrease in RH in the low, mid, and high cloud layers (850 hPa-300 hPa) at the 90% confidence interval
(**Figure 10, Figure b**). This decrease in RH is associated with a reduction of clouds, which results in a reduction in CF and P
significantly in the nCA-NV region. This reduction in clouds is then associated with an increase in TOA radiative flux (**Figure**
13h), despite a decrease in CTH (**Figure 13f**). In short, during high RH_s extremes, fires in nCA appear to create a positive
470 feedback that entails emissions of absorbing aerosols that warm the troposphere, creating a semi-direct effect. This semi-direct
effect contributes at least partially to conditions more favorable to fires. Significant reductions in nCA P may prolong the
wildfire season further into autumn (Goss et al., 2020), and increases in T as well as decreases in RH may create conditions
more favorable for more fires to ignite and grow (Varga et al., 2022). Under the low RH_s stratification, regional warming of
the cloud layer and surface becomes larger, more widespread, and less uncertain (**Figure 6e, Figure S12**). This is likely due to
475 a significantly higher amount of AOD/BC present in the atmosphere on these days (**Figure 3, Figure 4**), which corresponds
with a stronger heating rate (**Figure 5b**). However, with a higher likelihood of lower RH_{cl} anomalies during low RH_s condi-
tions (**Figure 2e**), the result is a muted, but significant, CF and P suppression that is limited to only nCA.

The results of this paper are consistent with the findings of Thornhill et al. (2018), who utilize a similar methodology to this
480 paper to analyze the effects of large fires. They ran the Met Office Unified Model using aircraft observations of AOD and BB
aerosol properties. They compared meteorological variables in high vs low fire emission conditions over South America and

found a clear sky shortwave heating rate of the low-to-mid troposphere that is larger (0.2 K day^{-1}), but comparable, to the heating rates calculated in this paper. They found that their heating rate was associated with a stronger semi-direct effect than the one observed in this paper, with CF decreasing throughout all the troposphere, which would be consistent with their stronger heating rate. Though not related to fire, an aircraft observational study of anthropogenic BC over the Bay of Bengal found a BC heating rate of around 0.5 Kday^{-1} (Kant et al., 2023), which demonstrates that either the biomass burning aerosols in this study are associated with a weaker heating rate, or that the shortwave absorption estimated by CERES is underestimated. The heating rate calculated in this paper is consistent with the calculated heating rate of a modeling study that simulate a global $10\times BC$ scenario, which demonstrate a 0.08 Kday^{-1} (Stjern et al., 2017), a number consistent with the heating rate over the region of interest. Furthermore, the results of this study are consistent with numerous other satellite observational studies over the tropics and subtropics that demonstrate that aerosols associated with wildfires can burn off clouds when at a coincident height, resulting in a positive radiative forcing (Wilcox, 2012; Kaufman et al., 2005; Ackerman et al., 2000; Hansen et al., 1997). Additionally, the reduction in CF and P is consistent with the results of Chen et al. (2014), which was a biomass burning aerosol modelling experiment conducted over the United States. However, their proposed mechanism for these decreases was a change in convection. Concerning the increase in M_{H_2O} above sites of fire emission in **Figure 9b,c**, this is consistent with a recent study that found comparable results (Yu et al., 2024), however they found more water vapor higher in the troposphere than this paper. Additionally, it is noted that the observed microphysical effects of the BB aerosols in this paper, namely the lack of a regional decrease in R_{eff} , contrast to another observational study that overlaps with the region of interest in this paper (Twohy et al., 2021). An important note about that study, however, is that it only sampled the 2018 wildfire season while this study focuses on the entire 2003-2022 time span. Finally, one last caveat to this study is that while it has been demonstrated that the T anomalies in **Figure 6, Figure 7, and Figure 8** are associated with biomass burning aerosols during large fire events, this does not imply that the aerosol semi-direct effect is the only contributor to the observed meteorological anomalies. As previously stated, at most 40-90 % of the T anomaly is accounted for by the semi-direct effect. One possible source of noise is wind. **Figure S10** depicts a positive wind speed in northern Nevada that may be influencing cloud cover over that part of the region, and it is unknown if this signal has anything to do with the semi-direct effect. Additionally, wildfires are associated with an increase in sensible heat flux from the combustion of biofuels, which may contribute to the positive T anomaly as well (Dickinson et al., 2021).

The results of this paper are significant, as the potential positive feedbacks described in this paper may prolong wildfires, and therefore also prolong poor air quality conditions inside the southwestern US (Liu & Peng, 2019; O'Neill et al., 2021; Schlosser et al., 2017), as well as other parts of the country (Hung et al., 2020). Additionally, the significant decreases in P and/or increases T_s highlighted in this paper occur in heavily populated regions in the southwestern US, including: the San Francisco Bay Area, Humboldt County in California, and Washoe County in Nevada. Therefore, this study highlights an increased need for a curtailment of CO_2 emissions (Ma et al., 2021; Touma et al., 2021) and better land management practices (DellaSala et al., 2022; Minnich et al., 2000; Minnich, 2001), as climate change and land mismanagement have both contributed to the large fires in nCA in recent years. Furthermore, as large fires are projected to become more commonplace throughout

the 21st century due to these factors (Flannigan et al., 2013; United Nations Environment Programme, 2022), the results of this paper will become more relevant over time as today's 90th percentile fire emission conditions become more common throughout the 21st century. Additionally, this paper highlights the need for more climate models to incorporate feedbacks
520 between wildfires, their aerosols, and semi-direct effects. While advances in fire parameterization within climate models have been made (Li et al., 2019; Li & Lawrence, 2017; Mangeon et al., 2016; Allen et al., 2024a), these interactive fire modules are not widely implemented. This study demonstrates that interactive fire modules are a necessity in climate models, as lifetime of fires, and therefore the size and duration of aerosol emission events, are influenced by the fires themselves.

Code availability. Code used to process satellite data will be made available at the following GitHub repository:

525 <https://github.com/jgome222/Northern-California-large-fires-Associated-with-Decrease-in-Cloud-Cover-Over-the-Southwestern-US>

Data availability. All datasets utilized in this analysis are available online. MODIS datasets are available via the 787 NASA Level-1 and Atmosphere Archive & Distribution System (LAADS) Distributed Active Archive 788 Center (DAAC) at <https://ladsweb.modaps.eosdis.nasa.gov/archive/allData/61/>. CERES datasets can be found at <https://ceres.larc.nasa.gov/>. AIRS data is available via NASA's Earth Science Data 794 extremes (ESDS) program at <https://www.earthdata.nasa.gov/>. CALIPSO datasets are available at the Atmospheric Science Data
530 Center (ASDC) at <https://asdc.larc.nasa.gov/>. GFED4 fire emission data is archived on the GFED4 web page at <https://www.globalfiredata.org/>. MERRA-2 data can be found on the Goddard Earth Sciences Data and Information Services Center (GES DISC) website at <https://disc.gsfc.nasa.gov/datasets?project=MERRA-2>.

Appendix A

Symbol	Definition	Dataset Derived From	Name of Product(s) Used
BC	Black Carbon	MERRA-2	$BCPHILIC$, $BCPHOBIC$
DM	Fire dry matter emissions	GFED4	DM, daily_fraction
AOD	Aerosol Optical Depth	MODIS	Aerosol_Optical_Depth_Land_Ocean_Mean
M_{H_2O}	Water Mass Mixing Ratio	AIRS	H2O_MMR_D
T	Temperature	AIRS	Temperature_D
T_s	Surface Temperature	AIRS	SurfAirTemp_D
RH	Relative Humidity	AIRS	RelHum_D
RH_s	Surface Relative Humidity	AIRS	RelHumSurf_D
CF	Cloud Fraction	MODIS AIRS	Cloud_Fraction_Mean FineCloudFrc_D
CF_{cir}	Cirrus Cloud Fraction	MODIS	Cirrus_Fraction_Infrared
CF_{lw}	Liquid Water Cloud Fraction	MODIS	Cloud_Retrieval_Fraction_Liquid
CTH	Cloud Top Height	MODIS	Cloud_Top_Height_Mean
P	Precipitation	GPCP	precip
SWH_{aer}	Aerosol Shortwave Heating Rate	CERES	adj_all_sw_dn, adj_all_sw_up, adj_naer_sw_dn, adj_naer_sw_up
F_{aer}	Shortwave aerosol radiative forcing	CERES	Same as above variable
TOA_{cld}	Cloud-only Net Top of Atmosphere Flux	CERES	adj_all_sw_dn, adj_all_sw_up, adj_all_lw_up adj_clr_sw_dn, adj_clr_sw_up, adj_clr_lw_up
SW_u	Shortwave aerosol upward flux	CERES	adj_all_sw_up, adj_clr_sw_up
SW_d	Shortwave aerosol downward flux	CERES	adj_all_sw_dn, adj_clr_sw_dn
U_s	Surface Wind speed	CERES/GEOS	sfc_wind_speed
Liquid R_{eff}	Liquid Cloud Effective Radius	MODIS	Cloud_Effective_Radius_Ice_Mean
Ice R_{eff}	Ice Cloud Effective Radius	MODIS	Cloud_Effective_Radius_Liquid_Mean
LWP	Liquid Water Path	MODIS	Cloud_Water_Path_Liquid_Mean
IWP	Ice Water Path	MODIS	Cloud_Water_Path_Ice_Mean

Table A1. Definition of variables that were derived from satellite observational datasets, as well as the instrument and dataset they are derived from.

Symbol	Definition
nCA	Northern California
nCA-NV	Northern California-Nevada
US	United States
BB	Biomass Burning
OA	Organic Aerosol
CA	California
SW	Shortwave
AAOD	Absorbing Aerosol Optical Depth
LW	Longwave
TOA	Top of atmosphere
CCN	Cloud Condensation Nuclei
CDF	Cumulative Distribution Function

Table A2. Definitions of abbreviations found throughout the paper that are not associated with a dataset.

Descriptor	Definition
(<i>DM</i> 90)	Variable stratified by 90th percentile fire dry matter emission anomaly days in nCA
(<i>RH_s</i> 75)	Variable stratified by 75th percentile surface relative humidity anomaly days in nCA-NV
(<i>DM</i> 10)	Variable stratified by 10th percentile fire dry matter emission anomaly days in nCA
(<i>RH_s</i> 25)	Variable stratified by 25th percentile surface relative humidity anomaly days in nCA-NV
(<i>DM</i> 90, <i>RH_s</i> 75)	Variable stratified by 90th percentile fire dry matter emission anomaly days in nCA and 75th percentile surface relative humidity anomaly days in nCA-NV
(<i>DM</i> 10, <i>RH_s</i> 75)	Variable stratified by 10th percentile fire dry matter emission anomaly days in nCA and 75th percentile surface relative humidity anomaly days in nCA-NV
(<i>DM</i> 90, <i>RH_s</i> 25)	Variable stratified by 90th percentile fire dry matter emission anomaly days in nCA and 25th percentile surface relative humidity anomaly days in nCA-NV
(<i>DM</i> 10, <i>RH_s</i> 25)	Variable stratified by 10th percentile fire dry matter emission anomaly days in nCA and 25th percentile surface relative humidity anomaly days in nCA-NV
cl	Cloud layer (850-300 hPa) average of variable
s	Variable measured at the surface
pho	Hydrophobic aerosol
phi	Hydrophilic aerosol
aer	radiative forcing variable calculated from all-sky minus clear sky products (aerosol only)
cld	radiative forcing variable calculated from all-sky minus no aerosol products (cloud only)
Δ	Difference in variable under different fire and/or relative humidity conditions

Table A3. Definitions of subscripts and other descriptors for variables.

Author contributions. J.L.G. conceived the project, designed the study, performed data analysis and wrote the paper. R.J.A. performed
535 analyses, and wrote the paper. K.L. advised on methods.

Competing interests. The authors declare no competing interests.

Acknowledgements. R.J. Allen is supported by NSF grant AGS-2153486.

References

- Abel, S. J., Highwood, E. J., Haywood, J. M., & Stringer, M. A. (2005). The direct radiative effect of biomass burning aerosols over southern Africa. *Atmospheric Chemistry and Physics*, 5(7), 1999–2018.
- Ackerman, A. S., Toon, O. B., Stevens, D. E., Heymsfield, A. J., Ramanathan, V., & Welton, E. J. (2000). Reduction of Tropical Cloudiness by Soot. *Science*, 288(5468), 1042–1047.
- Adler, R. F., Sapiano, M. R. P., Huffman, G. J., Wang, J.-J., Gu, G., Bolvin, D., Chiu, L., Schneider, U., Becker, A., Nelkin, E., Xie, P., Ferraro, R., & Shin, D.-B. (2018). The Global Precipitation Climatology Project (GPCP) Monthly Analysis (New Version 2.3) and a Review of 2017 Global Precipitation. *Atmosphere*, 9(4), 138.
- Ager, A. A., Day, M. A., Alcasena, F. J., Evers, C. R., Short, K. C., & Grenfell, I. (2021). Predicting Paradise: Modeling future wildfire disasters in the western US. *Science of The Total Environment*, 784, 147057.
- AIRS Science Team & Teixeira, J. (2013). Aqua AIRS Level 3 Standard Daily Product using AIRS IR-only V6.
- Allen, R. J., Gomez, J., Horowitz, L. W., & Shevliakova, E. (2024a). Enhanced future vegetation growth with elevated carbon dioxide concentrations could increase fire activity. *Communications Earth & Environment*, 5(1), 1–15. Publisher: Nature Publishing Group.
- Allen, R. J., Hassan, T., Randles, C. A., & Su, H. (2019). Enhanced land–sea warming contrast elevates aerosol pollution in a warmer world. *Nature Climate Change*, 9(4), 300–305.
- Allen, R. J. & Sherwood, S. C. (2010). Aerosol-cloud semi-direct effect and land-sea temperature contrast in a GCM. *Geophysical Research Letters*, 37(7).
- Allen, R. J., Vega, C., Yao, E., & Liu, W. (2024b). Impact of industrial versus biomass burning aerosols on the Atlantic Meridional Overturning Circulation. *npj Climate and Atmospheric Science*, 7(1), 1–16.
- Allen, R. J., Zhao, X., Randles, C. A., Kramer, R. J., Samset, B. H., & Smith, C. J. (2023). Surface warming and wetting due to methane’s long-wave radiative effects muted by short-wave absorption. *Nature Geoscience*, 16(4), 314–320.
- Amiri-Farahani, A., Allen, R. J., Li, K.-F., Nabat, P., & Westervelt, D. M. (2020). A La Niña-Like Climate Response to South African Biomass Burning Aerosol in CESM Simulations. *Journal of Geophysical Research: Atmospheres*, 125(6), e2019JD031832.
- Amiri-Farahani, A., Allen, R. J., Neubauer, D., & Lohmann, U. (2017). Impact of Saharan dust on North Atlantic marine stratocumulus clouds: importance of the semidirect effect. *Atmospheric Chemistry and Physics*, 17(10), 6305–6322.
- Artaxo, P., Rizzo, L. V., Brito, J. F., Barbosa, H. M. J., Arana, A., Sena, E. T., Cirino, G. G., Bastos, W., Martin, S. T., & Andreae, M. O. (2013). Atmospheric aerosols in Amazonia and land use change: from natural biogenic to biomass burning conditions. *Faraday Discussions*, 165, 203–235.
- Brown, P. T., Hanley, H., Mahesh, A., Reed, C., Strenfel, S. J., Davis, S. J., Kochanski, A. K., & Clements, C. B. (2023). Climate warming increases extreme daily wildfire growth risk in California. *Nature*, (pp. 1–7).
- Buchard, V., da Silva, A. M., Colarco, P. R., Darmenov, A., Randles, C. A., Govindaraju, R., Torres, O., Campbell, J., & Spurr, R. (2015). Using the OMI aerosol index and absorption aerosol optical depth to evaluate the NASA MERRA Aerosol Reanalysis. *Atmospheric Chemistry and Physics*, 15(10), 5743–5760.
- Cape, J. N., Coyle, M., & Dumitrescu, P. (2012). The atmospheric lifetime of black carbon. *Atmospheric Environment*, 59, 256–263.
- Chen, D., Liu, Z., Schwartz, C. S., Lin, H.-C., Cetola, J. D., Gu, Y., & Xue, L. (2014). The impact of aerosol optical depth assimilation on aerosol forecasts and radiative effects during a wild fire event over the United States. *Geoscientific Model Development*, 7(6), 2709–2715.

- Chen, Y., Hall, J., van Wees, D., Andela, N., Hantson, S., Giglio, L., van der Werf, G. R., Morton, D. C., & Randerson, J. T. (2023). Multi-
575 decadal trends and variability in burned area from the fifth version of the Global Fire Emissions Database (GFED5). *Earth System Science Data*, 15(11), 5227–5259. Publisher: Copernicus GmbH.
- Cho, C., Kim, S.-W., Choi, W., & Kim, M.-H. (2022). Significant light absorption of brown carbon during the 2020 California wildfires. *Science of The Total Environment*, 813, 152453.
- Conrick, R., Mass, C. F., Boomgard-Zagrodnik, J. P., & Ovens, D. (2021). The Influence of Wildfire Smoke on Cloud Microphysics during
580 the September 2020 Pacific Northwest Wildfires. *Weather and Forecasting*, 36(4), 1519–1536.
- DellaSala, D. A., Baker, B. C., Hanson, C. T., Ruediger, L., & Baker, W. (2022). Have western USA fire suppression and megafire active management approaches become a contemporary Sisyphus? *Biological Conservation*, 268, 109499.
- Dickinson, M. B., Wold, C. E., Butler, B. W., Kremens, R. L., Jimenez, D., Sopko, P., & O'Brien, J. J. (2021). The Wildland Fire Heat Budget—Using Bi-Directional Probes to Measure Sensible Heat Flux and Energy in Surface Fires. *Sensors*, 21(6), 2135.
- 585 Doelling, D. (2016). CERES Level 3 SSF1deg-Day Aqua-MODIS HDF file - Edition 4A.
- Doelling, D. (2017). CERES Level 3 SYN1DEG-DAYTerra+Aqua HDF4 file - Edition 4A.
- Doelling, D. (2023). CERES and GEO-Enhanced TOA, Within-Atmosphere and Surface Fluxes, Clouds and Aerosols Daily Terra-NOAA20 Edition4A.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016). Overview of the Coupled Model
590 Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*, 9(5), 1937–1958.
- Fan, J., Wang, Y., Rosenfeld, D., & Liu, X. (2016). Review of Aerosol–Cloud Interactions: Mechanisms, Significance, and Challenges. *Journal of the Atmospheric Sciences*, 73(11), 4221–4252.
- Fasullo, J. T., Rosenbloom, N., Buchholz, R. R., Danabasoglu, G., Lawrence, D. M., & Lamarque, J.-F. (2021). Coupled Climate Responses to Recent Australian Wildfire and COVID-19 Emissions Anomalies Estimated in CESM2. *Geophysical Research Letters*, 48(15),
595 e2021GL093841.
- Flannigan, M., Cantin, A. S., de Groot, W. J., Wotton, M., Newbery, A., & Gowman, L. M. (2013). Global wildland fire season severity in the 21st century. *Forest Ecology and Management*, 294, 54–61.
- Ford, P. L. & Johnson, G. V. (2006). Effects of dormant- vs. growing-season fire in shortgrass steppe: Biological soil crust and perennial grass responses. *Journal of Arid Environments*, 67(1), 1–14.
- 600 Giglio, L., Boschetti, L., Roy, D. P., Humber, M. L., & Justice, C. O. (2018). The Collection 6 MODIS burned area mapping algorithm and product. *Remote Sensing of Environment*, 217, 72–85.
- Giglio, L. & Justice, C. (2015). MYD14A1 MODIS/Aqua Thermal Anomalies/Fire Daily L3 Global 1km SIN Grid V006.
- Giglio, L., Randerson, J. T., & Van Der Werf, G. R. (2013). Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4). *Journal of Geophysical Research: Biogeosciences*, 118(1), 317–328.
- 605 Global Modeling And Assimilation Office & Pawson, S. (2015). MERRA-2 inst3_3d_aer_nv: 3d,3-Hourly,Instantaneous,Model-Level,Assimilation,Aerosol Mixing Ratio V5.12.4.
- Gomez, J., Allen, R. J., Turnock, S. T., Horowitz, L. W., Tsigaridis, K., Bauer, S. E., Olivié, D., Thomson, E. S., & Ginoux, P. (2023). The projected future degradation in air quality is caused by more abundant natural aerosols in a warmer world. *Communications Earth & Environment*, 4(1), 1–11.
- 610 Goren, T. & Rosenfeld, D. (2012). Satellite observations of ship emission induced transitions from broken to closed cell marine stratocumulus over large areas. *Journal of Geophysical Research: Atmospheres*, 117(D17).

- Goss, M., Swain, D. L., Abatzoglou, J. T., Sarhadi, A., Kolden, C. A., Williams, A. P., & Diffenbaugh, N. S. (2020). Climate change is increasing the likelihood of extreme autumn wildfire conditions across California. *Environmental Research Letters*, 15(9), 094016.
- Han, Q., Rossow, W. B., Chou, J., & Welch, R. M. (1998). Global Survey of the Relationships of Cloud Albedo and Liquid Water Path with Droplet Size Using ISCCP. *Journal of Climate*, 11(7), 1516–1528.
- Hansen, J., Sato, M., & Ruedy, R. (1997). Radiative forcing and climate response. *Journal of Geophysical Research: Atmospheres*, 102(D6), 6831–6864.
- He, C., Li, Q., Liou, K.-N., Qi, L., Tao, S., & Schwarz, J. P. (2016). Microphysics-based black carbon aging in a global CTM: constraints from HIPPO observations and implications for global black carbon budget. *Atmospheric Chemistry and Physics*, 16(5), 3077–3098.
- Helmert, J., Heinold, B., Tegen, I., Hellmuth, O., & Wendisch, M. (2007). On the direct and semidirect effects of Saharan dust over Europe: A modeling study. *Journal of Geophysical Research: Atmospheres*, 112(D13).
- Herbert, R. & Stier, P. (2023). Satellite observations of smoke–cloud–radiation interactions over the Amazon rainforest. *Atmospheric Chemistry and Physics*, 23(7), 4595–4616.
- Highwood, E. J. & Ryder, C. L. (2014). Radiative Effects of Dust. In P. Knippertz & J.-B. W. Stuut (Eds.), *Mineral Dust: A Key Player in the Earth System* (pp. 267–286). Dordrecht: Springer Netherlands.
- Huffman, G. J., Adler, R. F., Morrissey, M. M., Bolvin, D. T., Curtis, S., Joyce, R., McGavock, B., & Susskind, J. (2001). Global Precipitation at One-Degree Daily Resolution from Multisatellite Observations. *Journal of Hydrometeorology*, 2(1), 36–50.
- Hung, W.-T., Lu, C.-H. S., Shrestha, B., Lin, H.-C., Lin, C.-A., Grogan, D., Hong, J., Ahmadov, R., James, E., & Joseph, E. (2020). The impacts of transported wildfire smoke aerosols on surface air quality in New York State: A case study in summer 2018. *Atmospheric Environment*, 227, 117415.
- Jacobson, M. Z. (2014). Effects of biomass burning on climate, accounting for heat and moisture fluxes, black and brown carbon, and cloud absorption effects. *Journal of Geophysical Research: Atmospheres*, 119(14), 8980–9002.
- Kant, S., Sarangi, C., & Wilcox, E. M. (2023). Aerosol processes perturb cloud trends over Bay of Bengal: observational evidence. *npj Climate and Atmospheric Science*, 6(1), 1–8.
- Kaufman, Y. J., Koren, I., Remer, L. A., Rosenfeld, D., & Rudich, Y. (2005). The effect of smoke, dust, and pollution aerosol on shallow cloud development over the Atlantic Ocean. *Proceedings of the National Academy of Sciences of the United States of America*, 102(32), 11207–11212.
- Keeley, J. E. & Syphard, A. D. (2019). Twenty-first century California, USA, wildfires: fuel-dominated vs. wind-dominated fires. *Fire Ecology*, 15(1), 24.
- Khain, A. P. (2009). Notes on state-of-the-art investigations of aerosol effects on precipitation: a critical review. *Environmental Research Letters*, 4(1), 015004.
- Koch, D. & Del Genio, A. D. (2010). Black carbon semi-direct effects on cloud cover: review and synthesis. *Atmospheric Chemistry and Physics*, 10(16), 7685–7696.
- Kok, J. F., Storelmo, T., Karydis, V. A., Adebisi, A. A., Mahowald, N. M., Evan, A. T., He, C., & Leung, D. M. (2023). Mineral dust aerosol impacts on global climate and climate change. *Nature Reviews Earth & Environment*, 4(2), 71–86.
- LeNoir, J. S., McConnell, L. L., Fellers, G. M., Cahill, T. M., & Seiber, J. N. (1999). Summertime transport of current-use pesticides from California’s Central Valley to the Sierra Nevada Mountain Range, USA. *Environmental Toxicology and Chemistry*, 18(12), 2715–2722.
- Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., & Hsu, N. C. (2013). The Collection 6 MODIS aerosol products over land and ocean. *Atmospheric Measurement Techniques*, 6(11), 2989–3034.

- 650 Li, F. & Lawrence, D. M. (2017). Role of Fire in the Global Land Water Budget during the Twentieth Century due to Changing Ecosystems. *Journal of Climate*, 30(6), 1893–1908.
- Li, F., Val Martin, M., Andreae, M. O., Arneth, A., Hantson, S., Kaiser, J. W., Lasslop, G., Yue, C., Bachelet, D., Forrest, M., Kluzek, E., Liu, X., Mangeon, S., Melton, J. R., Ward, D. S., Darnenov, A., Hickler, T., Ichoku, C., Magi, B. I., Sitch, S., van der Werf, G. R., Wiedinmyer, C., & Rabin, S. S. (2019). Historical (1700–2012) global multi-model estimates of the fire emissions from the Fire Modeling
655 Intercomparison Project (FireMIP). *Atmospheric Chemistry and Physics*, 19(19), 12545–12567.
- Li, S. & Banerjee, T. (2021). Spatial and temporal pattern of wildfires in California from 2000 to 2019. *Scientific Reports*, 11(1), 8779.
- Liu, J. C. & Peng, R. D. (2019). The impact of wildfire smoke on compositions of fine particulate matter by ecoregion in the Western US. *Journal of Exposure Science & Environmental Epidemiology*, 29(6), 765–776.
- Liu, Z., Hu, M., Hu, Y., & Wang, G. (2018). Estimation of net primary productivity of forests by modified CASA models and remotely
660 sensed data. *International Journal of Remote Sensing*, 39(4), 1092–1116.
- Lohmann, U., Friebel, F., Kanji, Z. A., Mahrt, F., Mensah, A. A., & Neubauer, D. (2020). Future warming exacerbated by aged-soot effect on cloud formation. *Nature Geoscience*, 13(10), 674–680.
- Ma, W., Zhai, L., Pivovarov, A., Shuman, J., Buotte, P., Ding, J., Christoffersen, B., Knox, R., Moritz, M., Fisher, R. A., Koven, C. D., Kueppers, L., & Xu, C. (2021). Assessing climate change impacts on live fuel moisture and wildfire risk using a hydrodynamic vegetation
665 model. *Biogeosciences*, 18(13), 4005–4020.
- Mangeon, S., Voulgarakis, A., Gilham, R., Harper, A., Sitch, S., & Folberth, G. (2016). INFERNO: a fire and emissions scheme for the UK Met Office's Unified Model. *Geoscientific Model Development*, 9(8), 2685–2700.
- Martins, J. V., Marshak, A., Remer, L. A., Rosenfeld, D., Kaufman, Y. J., Fernandez-Borda, R., Koren, I., Correia, A. L., Zubko, V., & Artaxo, P. (2011). Remote sensing the vertical profile of cloud droplet effective radius, thermodynamic phase, and temperature. *Atmospheric
670 Chemistry and Physics*, 11(18), 9485–9501.
- Minnich, R. A. (2001). An Integrated Model of Two Fire Regimes. *Conservation Biology*, 15(6), 1549–1553.
- Minnich, R. A., Barbour, M. G., Burk, J. H., & Sosa-Ramírez, J. (2000). Californian mixed-conifer forests under unmanaged fire regimes in the Sierra San Pedro Mártir, Baja California, Mexico. *Journal of Biogeography*, 27(1), 105–129.
- Minnich, R. A. & Chou, Y. H. (1997). Wildland Fire Patch Dynamics in the Chaparral of Southern California and Northern Baja California.
675 *International Journal of Wildland Fire*, 7(3), 221–248.
- MODIS Atmosphere Science Team (2017). MYD08_d3 MODIS/Aqua Aerosol Cloud Water Vapor Ozone Daily L3 Global 1Deg CMG.
- Nguyen, L. S. P., Huang, H.-Y., Lei, T. L., Bui, T. T., Wang, S.-H., Chi, K. H., Sheu, G.-R., Lee, C.-T., Ou-Yang, C.-F., & Lin, N.-H. (2020). Characterizing a landmark biomass-burning event and its implication for aging processes during long-range transport. *Atmospheric
680 Environment*, 241, 117766.
- Ogren, J. A. & Charlson, R. J. (1983). Elemental carbon in the atmosphere: cycle and lifetime. *Tellus B*, 35B(4), 241–254.
- O'Neill, S. M., Diao, M., Raffuse, S., Al-Hamdan, M., Barik, M., Jia, Y., Reid, S., Zou, Y., Tong, D., West, J. J., Wilkins, J., Marsha, A., Freedman, F., Vargo, J., Larkin, N. K., Alvarado, E., & Loesche, P. (2021). A multi-analysis approach for estimating regional health impacts from the 2017 Northern California wildfires. *Journal of the Air & Waste Management Association*, 71(7), 791–814.
- Palinkas, L. A. (2020). The California Wildfires. In L. A. Palinkas (Ed.), *Global Climate Change, Population Displacement, and Public
685 Health: The Next Wave of Migration* (pp. 53–67). Cham: Springer International Publishing.
- Peterson, D., Wang, J., Ichoku, C., Hyer, E., & Ambrosia, V. (2013). A sub-pixel-based calculation of fire radiative power from MODIS observations: 1: Algorithm development and initial assessment. *Remote Sensing of Environment*, 129, 262–279.

- Platnick, S., King, M., Ackerman, S., Menzel, W., Baum, B., Riedi, J., & Frey, R. (2003). The MODIS cloud products: algorithms and examples from Terra. *IEEE Transactions on Geoscience and Remote Sensing*, 41(2), 459–473.
- 690 Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Holz, R. E., Yang, P., Ridgway, W. L., & Riedi, J. (2017). The MODIS Cloud Optical and Microphysical Products: Collection 6 Updates and Examples From Terra and Aqua. *IEEE Transactions on Geoscience and Remote Sensing*, 55(1), 502–525.
- Raga, G. B., Baumgardner, D., Rios, B., Díaz-Esteban, Y., Jaramillo, A., Gallagher, M., Sauvage, B., Wolff, P., & Lloyd, G. (2022). High concentrations of ice crystals in upper-tropospheric tropical clouds: is there a link to biomass and fossil fuel combustion? *Atmospheric*
695 *Chemistry and Physics*, 22(4), 2269–2292.
- Randerson, J., van der Werf, G., Giglio, L., Collatz, G., & Kasibhatla, P. (2017). Global Fire Emissions Database, Version 4.1 (GFEDv4). (pp. 1925.7122549999906 MB).
- Randles, C. A., Silva, A. M. d., Buchard, V., Colarco, P. R., Darmenov, A., Govindaraju, R., Smirnov, A., Holben, B., Ferrare, R., Hair, J., Shinozuka, Y., & Flynn, C. J. (2017). The MERRA-2 Aerosol Reanalysis, 1980 Onward. Part I: System Description and Data Assimilation
700 Evaluation. *Journal of Climate*, 30(17), 6823–6850.
- Ruffault, J., Curt, T., Moron, V., Trigo, R. M., Mouillot, F., Koutsias, N., Pimont, F., Martin-StPaul, N., Barbero, R., Dupuy, J.-L., Russo, A., & Belhadj-Khedher, C. (2020). Increased likelihood of heat-induced large wildfires in the Mediterranean Basin. *Scientific Reports*, 10(1), 13790.
- Sakaeda, N., Wood, R., & Rasch, P. J. (2011). Direct and semidirect aerosol effects of southern African biomass burning aerosol. *Journal of*
705 *Geophysical Research: Atmospheres*, 116(D12).
- Salomonson, V., Barnes, W., Xiong, J., Kempler, S., & Masuoka, E. (2002). An overview of the Earth Observing System MODIS instrument and associated data systems performance. In *IEEE International Geoscience and Remote Sensing Symposium*, volume 2 (pp. 1174–1176 vol.2).
- Samset, B. H. (2022). Aerosol absorption has an underappreciated role in historical precipitation change. *Communications Earth & Envi-*
710 *ronment*, 3(1), 1–8.
- Sand, M., Samset, B. H., Tsigaridis, K., Bauer, S. E., & Myhre, G. (2020). Black Carbon and Precipitation: An Energetics Perspective. *Journal of Geophysical Research: Atmospheres*, 125(13), e2019JD032239.
- Schlosser, J. S., Braun, R. A., Bradley, T., Dadashazar, H., MacDonald, A. B., Aldhaif, A. A., Aghdam, M. A., Mardi, A. H., Xian, P., & Sorooshian, A. (2017). Analysis of aerosol composition data for western United States wildfires between 2005 and 2015: Dust emissions,
715 chloride depletion, and most enhanced aerosol constituents. *Journal of Geophysical Research: Atmospheres*, 122(16), 8951–8966.
- Shi, G., Yan, H., Zhang, W., Dodson, J., Heijnis, H., & Burrows, M. (2021). Rapid warming has resulted in more wildfires in northeastern Australia. *Science of The Total Environment*, 771, 144888.
- Smith, C. J., Kramer, R. J., Myhre, G., Forster, P. M., Soden, B. J., Andrews, T., Boucher, O., Faluvegi, G., Fläschner, D., Hodnebrog, Ø., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Mülmenstädt, J., Olivíé, D., Richardson, T., Samset, B. H., Shindell, D., Stier, P.,
720 Takemura, T., Voulgarakis, A., & Watson-Parris, D. (2018). Understanding Rapid Adjustments to Diverse Forcing Agents. *Geophysical Research Letters*, 45(21), 12,023–12,031.
- Stjern, C. W., Samset, B. H., Myhre, G., Forster, P. M., Hodnebrog, Ø., Andrews, T., Boucher, O., Faluvegi, G., Iversen, T., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Olivíé, D., Richardson, T., Shawki, D., Shindell, D., Smith, C. J., Takemura, T., & Voulgarakis, A. (2017). Rapid Adjustments Cause Weak Surface Temperature Response to Increased Black Carbon Concentrations. *Journal of*
725 *Geophysical Research: Atmospheres*, 122(21), 11,462–11,481.

- Thornhill, G. D., Ryder, C. L., Highwood, E. J., Shaffrey, L. C., & Johnson, B. T. (2018). The effect of South American biomass burning aerosol emissions on the regional climate. *Atmospheric Chemistry and Physics*, 18(8), 5321–5342.
- Touma, D., Stevenson, S., Lehner, F., & Coats, S. (2021). Human-driven greenhouse gas and aerosol emissions cause distinct regional impacts on extreme fire weather. *Nature Communications*, 12(1), 212.
- 730 Tsikerdekis, A., Zanis, P., Georgoulas, A. K., Alexandri, G., Katragkou, E., Karacostas, T., & Solmon, F. (2019). Direct and semi-direct radiative effect of North African dust in present and future regional climate simulations. *Climate Dynamics*, 53(7), 4311–4336.
- Twohy, C. H., Toohey, D. W., Levin, E. J. T., DeMott, P. J., Rainwater, B., Garofalo, L. A., Pothier, M. A., Farmer, D. K., Kreidenweis, S. M., Pokhrel, R. P., Murphy, S. M., Reeves, J. M., Moore, K. A., & Fischer, E. V. (2021). Biomass Burning Smoke and Its Influence on Clouds Over the Western U. S. *Geophysical Research Letters*, 48(15), e2021GL094224.
- 735 United Nations Environment Programme (2022). Spreading like Wildfire – The Rising Threat of Extraordinary Landscape Fires.
- Urbanski, S. P. (2013). Combustion efficiency and emission factors for wildfire-season fires in mixed conifer forests of the northern Rocky Mountains, US. *Atmospheric Chemistry and Physics*, 13(14), 7241–7262.
- Urbanski, S. P., Hao, W. M., & Nordgren, B. (2011). The wildland fire emission inventory: western United States emission estimates and an evaluation of uncertainty. *Atmospheric Chemistry and Physics*, 11(24), 12973–13000.
- 740 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., & Kasibhatla, P. S. (2017). Global fire emissions estimates during 1997–2016. *Earth System Science Data*, 9(2), 697–720.
- Varga, K., Jones, C., Trugman, A., Carvalho, L. M. V., McLoughlin, N., Seto, D., Thompson, C., & Daum, K. (2022). Megafires in a Warming World: What Wildfire Risk Factors Led to California’s Largest Recorded Wildfire. *Fire*, 5(1), 16.
- 745 Wagenbrenner, N. S., Chung, S. H., & Lamb, B. K. (2017). A large source of dust missing in Particulate Matter emission inventories? Wind erosion of post-fire landscapes. *Elementa: Science of the Anthropocene*, 5, 2.
- Wagenbrenner, N. S., Germino, M. J., Lamb, B. K., Robichaud, P. R., & Foltz, R. B. (2013). Wind erosion from a sagebrush steppe burned by wildfire: Measurements of PM10 and total horizontal sediment flux. *Aeolian Research*, 10, 25–36.
- Wagner, R., Jähn, M., & Schepanski, K. (2018). Wildfires as a source of airborne mineral dust – revisiting a conceptual model using large-eddy simulation (LES). *Atmospheric Chemistry and Physics*, 18(16), 11863–11884.
- 750 Wagner, R., Schepanski, K., & Klose, M. (2021). The Dust Emission Potential of Agricultural-Like Fires—Theoretical Estimates From Two Conceptually Different Dust Emission Parameterizations. *Journal of Geophysical Research: Atmospheres*, 126(18), e2020JD034355.
- Wei, J., Li, Z., Peng, Y., & Sun, L. (2019). MODIS Collection 6.1 aerosol optical depth products over land and ocean: validation and comparison. *Atmospheric Environment*, 201, 428–440.
- 755 Wilcox, E. M. (2012). Direct and semi-direct radiative forcing of smoke aerosols over clouds. *Atmospheric Chemistry and Physics*, 12(1), 139–149.
- Wilmot, T. Y., Mallia, D. V., Hallar, A. G., & Lin, J. C. (2022). Wildfire plumes in the Western US are reaching greater heights and injecting more aerosols aloft as wildfire activity intensifies. *Scientific Reports*, 12(1), 12400.
- Yu, J., Jiang, X., Zeng, Z.-C., & Yung, Y. L. (2024). Fire monitoring and detection using brightness-temperature difference and water vapor emission from the atmospheric infrared sounder. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 317, 108930.
- 760 Yu, Y. & Ginoux, P. (2022). Enhanced dust emission following large wildfires due to vegetation disturbance. *Nature Geoscience*, 15(11), 878–884.

Zaremba, L. L. & Carroll, J. J. (1999). Summer Wind Flow Regimes over the Sacramento Valley. *Journal of Applied Meteorology and Climatology*, 38(10), 1463–1473.




Figure 1. Distribution of fires and the corresponding aerosol optical depth AOD anomaly impacts during the fire season. (a) 2003-2022 average daily fire dry matter DM emissions for the southwestern United States during the fire season (June-October). Blue box signifies the nCA (northern California) region, where average daily fire emissions are the highest. (b) 2003-2022 June-October daily MODIS Aerosol optical depth (AOD) difference between average AOD on 90th percentile DM ($DM90$) and average AOD on 10th percentile DM ($DM10$) days within the 2003-2022 June-October time period. ΔAOD represents $AOD(DM90) - AOD(DM10)$. Green box symbolizes the nCA-NV (northern California-Nevada) region, where increases in AOD and changes in cloud properties (**Figure 11**) are most significant. Black dots represent statistically significant differences at 90% confidence according to a two-tailed test.




Figure 2. Dependence of meteorological variables on high versus low surface relative humidity RH_s during the fire season. Regional average cumulative distribution functions (CDFs) for variable anomalies stratified by 75th percentile surface relative humidity (RH_s75) days (red) and 25th percentile (RH_s25) (blue) days within the 2003-2022 June-October time period. Variables depicted include (a) northern California (nCA) fire dry matter (DM) emissions, (b) southwestern US surface temperature T_s , (c) nCA-NV cloud layer (850-300 hPa) average temperature T_{cl} , (d) southwestern US surface relative humidity RH_s , (e) southwestern US cloud layer average relative humidity RH_{cl} , (f) southwestern US cloud fraction CF , (g) southwestern US precipitation P , and (h) southwestern US surface wind speed U . Δ represents the difference between the variable's average anomaly for RH_s75 and RH_s25 days.




Figure 3. Difference in AOD anomalies on high and low RH_s days during the fire season. Daily nCA-NV AOD anomalies stratified by nCA-NV RH_s and nCA DM extremes within the 2003-2022 June-October time period. (a) displays cumulative distribution functions for daily June-October 2003-2022 daily northern California-Nevada nCA-NV AOD stratified by high (90th percentile) nCA DM emissions and high nCA-NV RH_s $AOD(DM90, RH_s 75)$ (solid red line), low (10th percentile) DM and high RH_s $AOD(DM10, RH_s 75)$ (dashed red), high DM /low RH_s $AOD(DM90, RH_s 25)$ (solid blue line), and low nCA DM /low RH_s $AOD(DM10, RH_s 25)$ (dashed blue line). The red ΔAOD represents the difference between the solid red and dashed red line $AOD(DM90, RH_s 75) - AOD(DM10, RH_s 75)$ and the blue ΔAOD represents the difference between the solid and dashed blue lines $AOD(DM90, RH_s 25) - AOD(DM10, RH_s 25)$. (b) Depicts a map of $AOD(DM90, RH_s 25) - AOD(DM10, RH_s 25)$. Pearson cross correlation coefficient r between ΔAOD and nCA DM emissions is depicted in the top left corner. (c) Depicts a map of average $AOD(DM90, RH_s 75) - AOD(DM10, RH_s 75)$. Black dots in (b),(c) represent statistically significant differences at the 90% confidence interval according to a two-tailed test.




Figure 4. Difference in MERRA-2 black carbon BC profiles on high vs low fire days stratified by differing RH_s conditions in the nCA-NV region in the 2003-2022 June-October time period. Profiles of both aged hydrophilic black carbon BC_{phi} (a,d) as well as freshly emitted hydrophobic black carbon BC_{pho} (b,e) are depicted in addition to total BC (c,f). All types of BC have significant anomalies from 850-300 hPa under both high RH_s (a-c) as well as low RH_s conditions (d-f).




Figure 5. High minus low DM days regional average aerosol-only shortwave heating rate SWH_{aer} profiles under differing RH_s conditions in the 2003-2022 June-October time period. There is a significant shortwave aerosol heating rate from 850-500 hPa under both high RH_s conditions (a) as well as low RH_s conditions (b).




Figure 6. Responses in AIRS temperature T , water mass mixing ratio M_{H_2O} , relative humidity RH , and cloud fraction CF profiles to large fires under high and low RH_s extremes during the fire season. nCA-NV regional-temporal average differences in T , water mass mixing ratio M_{H_2O} and relative humidity RH profiles for under high minus low DM conditions stratified by RH_s75 (a-d) and RH_s25 (e-h) in the 2002-2023 fire season (June-October) time period. Error bars represent the 90% confidence interval.




Figure 7. High minus low *DM* days MERRA-2 *BC* anomalies at all AIRS pressure levels from 1000 hPa to 200 hPa (a-j) under high RH_s conditions in the 2003-2022 June-October time period. Black dots indicate statistical significance at the 90% confidence interval. r values indicate spatial Pearson cross correlations between total *BC* and MODIS *AOD*.




Figure 8. High minus low DM days AIRS T anomalies at all AIRS pressure levels from 1000 hPa to 200 hPa (a-j) under high RH_s conditions in the 2003-2022 June-October time period. Black dots indicate statistical significance at the 90% confidence interval. r values indicate spatial Pearson cross correlations between T and MODIS AOD .




Figure 9. High minus low *DM* days AIRS M_{H_2O} anomalies at all AIRS pressure levels from 1000 hPa to 200 hPa (a-j) under high RH_s conditions in the 2003-2022 June-October time period. Black dots indicate statistical significance at the 90% confidence interval. r values indicate spatial Pearson cross correlations between total M_{H_2O} and MODIS *AOD*.




Figure 10. High minus low *DM* days AIRS *RH* anomalies at all AIRS pressure levels from 1000 hPa to 200 hPa (a-j) under high RH_s conditions in the 2003-2022 June-October time period. Black dots indicate statistical significance at the 90% confidence interval. r values indicate spatial Pearson cross correlations between *RH* and MODIS *AOD*.




Figure 11. High minus low *DM* days AIRS *CF* anomalies at all AIRS pressure levels from 1000 hPa to 200 hPa (a-j) under high RH_s conditions in the 2003-2022 June-October time period. Black dots indicate statistical significance at the 90% confidence interval. r values indicate spatial Pearson cross correlations between *CF* and MODIS *AOD*.




Figure 12. Dependence of meteorological variables on high versus low RH_s and fires during the fire season. Empirical CDFs for regional average daily anomalies of meteorological variables over the nCA-NV region in the 2003-2022 June-October time period. Solid red line signifies variable anomalies are stratified by high nCA fire dry matter emission DM and high nCA-NV RH_s anomaly days ($DM90, RH_s75$). The dashed red line signifies variable anomalies are stratified by low DM and high RH_s anomaly days ($DM10, RH_s75$). The solid blue line represents variable anomalies are stratified by high DM and low RH_s anomaly days ($DM90, RH_s25$). The dashed blue line symbolizes variable anomalies are stratified by low DM and RH_s anomaly days ($DM10, RH_s25$). Variables depicted include (a) liquid water cloud fraction CF_{lw} , (b) cirrus cloud fraction CF_{cir} , (c) CF , (d) cloud top height CTH , (e) precipitation P , and cloud-only (all-sky minus clear-sky) net top of atmosphere flux TOA_{cld} . The red Δ represents the differences in the mean of the solid red and dashed red lines ($DM90, RH_s75$)-($DM10, RH_s75$). The blue Δ represents the differences in the mean of the solid blue and dashed blue lines ($DM90, RH_s25$)-($DM10, RH_s25$).




Figure 13. Meteorological responses under high versus low nCA *DM* conditions with simultaneously high nCA-NV RH_s during the fire season. Difference between average variable anomalies on high (90th percentile) nCA fire dry matter *DM* emission days and low (10th percentile) nCA *DM* emission days that occur on high nCA-NV RH_s days in the 2003-2022 June-October time period. Variables include (a) 850 hPa-300 hPa average Temperature T_{cl} , (b) 850 hPa-300 hPa average relative humidity RH_{cl} , (c) surface temperature T_s , (d) RH_s , (e) CF , (f) CTH , (g) P , and (h) TOA_{cld} . Black dots represent statistically significant differences at the 90% confidence interval according to a two-tailed test. Pearson cross correlation r values in each plot represent the spatial correlation between MODIS aerosol optical depth AOD anomaly and the variable anomaly depicted in the figure. All values of r are significant at the 90% confidence interval according to a two-tailed test.




Figure 14. Effect size of large fires in nCA on the mean of various meteorological variables during the fire season. 2003-2022 June-October Cohen's d values for the difference between nCA-NV regional averages of variables on high DM days minus low nCA DM emission days that coincide with (a) high RH_s and (b) low RH_s . For Cohen's d , values of 0.2 through 0.5 signify a weak effect size, values of 0.5 through 0.8 represent a moderate effect size, and values greater or equal to 0.8 signify a strong effect size. Red bars represent standard error.




Figure 15. Dependence of microphysical variables to high versus low surface relative humidity RH_s and fires during the fire season. Empirical CDFs for regional average daily anomalies of cloud microphysical variables over the nCA-NV region in the 2003-2022 June-October time period. Solid red line signifies variable anomalies are stratified by $(DM90, RH_s 75)$. The dashed red line signifies variable anomalies are stratified by $(DM10, RH_s 75)$. The solid blue line represents variable anomalies are stratified $(DM90, RH_s 25)$. The dashed blue line symbolizes variable anomalies are stratified by $(DM10, RH_s 25)$. Variables depicted include (a) liquid effective radius R_{eff} , (b) Ice R_{eff} , (c) liquid water path LWP , (d) and ice water path IWP . The red Δ represents the differences in the mean of the solid red and dashed red lines $(DM90, RH_s 75)-(DM10, RH_s 75)$. The blue Δ represents the differences in the mean of the solid blue and dashed blue lines $(DM90, RH_s 25)-(DM10, RH_s 25)$.