

Antarctic sensitivity to oceanic melting parameterizations

Antonio Juarez-Martinez^{1,2}, Javier Blasco³, Alexander Robinson⁴, Marisa Montoya^{1,2}, and Jorge Alvarez-Solas^{1,2}

¹Complutense University of Madrid, Madrid, Spain

²Geoscience Institute CSIC-UCM, Madrid, Spain

³Université Libre de Bruxelles, Brussels, Belgium

⁴Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

Correspondence: Antonio Juarez-Martinez (antjua01@ucm.es)

Abstract. The Antarctic Ice Sheet (AIS) has experienced accelerated loss of ice over the last decades and could become the main contributor to sea-level rise in the coming centuries. However, the associated uncertainty is very large. The main sources of this uncertainty lie in the future scenarios, the climatic forcing and, most notably, the structural uncertainty due to our lack of understanding of ice-ocean interaction processes, in particular, the representation of sub-shelf basal melt. In this study, we use a higher-order ice-sheet model to investigate the impact of these three sources of uncertainty in the contribution of the AIS to sea level in the coming centuries in the context of the Ice Sheet Model Intercomparison Project for CMIP6 (ISMIP6) but extending the projections until 2500. We test the sensitivity of the model to basal melting parameters using several forcings and scenarios simulated in the CMIP5 and CMIP6 ensembles. Results show a strong dependency on the values of the parameter that controls the heat exchange velocity between ice and ocean and also the forcing and scenario. Higher values of the heat exchange parameter lead to higher sea-level rise, with the contribution depending on the forcing-scenario configuration and reaching in some cases more than 3 meters in sea level equivalent by the end of 2500. Idealized simulations considering the individual effects of the atmospheric and the oceanic forcing have been performed, demonstrating that the oceanic forcing plays a dominant role over the western sector of the AIS while the atmospheric forcing is more important for the eastern sector and the interior.

1 Introduction

Antarctica hosts today's largest ice sheet on Earth, the Antarctic Ice Sheet (AIS), with a total volume close to 27 million km³, corresponding to a sea level equivalent (SLE) of about 58 m (Morlighem et al., 2020). Ice flows from the ice-sheet interior within ice streams towards the ice-sheet margin. As it spreads outwards, it thins and eventually starts floating, forming ice shelves. The AIS is fringed by more than 300 of them, representing nearby 2.5% of its volume of ice and collecting over 80% of the outflow from the grounded ice, in its vast majority from the West Antarctic Ice Sheet (WAIS; Fowler and Ng, 2021).

Subshelf melting and calving are the main mechanisms responsible for ice-mass loss of the AIS today (Depoorter et al., 2013). Thus, the WAIS, which is broadly a marine-based ice sheet, experiences the highest melt rates through the critical influence of warmer water in contact with the ice, especially in the Amundsen and Bellinghausen Seas (Rignot et al., 2013).

Enhanced basal melting leads to ice-shelf thinning. Since ice shelves are floating, this does not lead to an increase in sea level
25 per se. However, laterally-confined ice shelves buttress the upstream flow. Therefore their thinning can reduce their buttressing
effect, accelerating inland ice flow and discharge, thereby contributing to sea-level increase (Fürst et al., 2016).

Several areas of the AIS have experienced significant ice mass loss in the past decades, contributing approximately 8 mm to
the sea level rise since 1992 (Bell and Seroussi, 2020 and references therein). Notably, most ice mass loss has originated in the
WAIS. The ice loss in this region has increased by 70% in the last decades (Paolo et al., 2015), ejecting approximately 3331 Gt
30 of ice to the Amundsen Sea (Davison et al., 2023). This is explained by the reinforced output of some major nearby glaciers
such as Pine Island, which has sped-up and reached its maximum velocity at ~ 4000 m/yr in 2009 (?)([Joughin et al., 2021](#)),
and Thwaites Glacier, which is responsible for 15% of the WAIS ice loss (Schmidt et al., 2023).

An increase in the discharge of ice across the grounding line causes it to retreat. If the ice sheet is grounded on a retrograde
slope, this retreat is thought to lead to an intrinsic instability, the marine ice-sheet instability (MISI; Schoof, 2007; Weertman,
35 1974), where a positive feedback is triggered. On a retrograde slope, the retreat of the grounding line implies an increase in the
ice thickness in that area. This enhances the discharge of ice across the grounding line, which drives a further retreat (Pattyn
and Morlighem, 2020). The MISI is an important source of uncertainty in future projections of the AIS (Robel et al., 2019).
Several studies have suggested that the aforementioned sectors of the WAIS that are experiencing fast ice-mass loss could be
already undergoing MISI (Joughin et al., 2014; Rignot et al., 2013). The current stability of the WAIS has been recently studied
40 by Hill et al. (2023), concluding that present-day retreat of Antarctic grounding lines is not yet irreversible or self-sustained.
Nevertheless, in a continuation of this work, Reese et al. (2022) showed that irreversibility could appear under present-day
climate forcing if the grounding lines retreated further inland.

In the future, under global warming, there is the risk of additional mass loss from the AIS both under sustained warming of
the Southern Ocean and the increased tendency to produce upwelling of Circumpolar Deep Waters (CDW) caused by stronger
45 westerly winds (Holland et al., 2020, 2022). Complete disappearance of the WAIS could lead to an estimated sea-level rise
of 1.91-5.08 m (Sun et al., 2020). Nonetheless, there is substantial uncertainty in the future contribution of the AIS to sea-
level rise (Edwards et al., 2021; Seroussi et al., 2020), which ranges from only about ~ 10 cm (Ritz et al., 2015) to above 1
m (DeConto and Pollard, 2016) in this century. Actually, as stated in the latest IPCC report (Masson-Delmotte et al., 2021),
the AIS is the largest source of uncertainty in multi-centennial sea-level projections, with a potential contribution by 2300
50 ranging from -0.14 m to 0.78 m in low emissions scenarios and from -0.27 m to more than 3 m in high emissions scenarios
(IPCC 2021, ?[Masson-Delmotte et al., 2021](#)). The main reason for this is the large uncertainty in the projections of ice-shelf
stability in warming scenarios (van de Wal et al., 2022), which depends critically both on the climate forcing and on ice-ocean
interaction processes that are not well constrained. Observations of large-scale ice sheets and cavities under ice shelves, where
basal melting takes place, are scarce and thus our understanding of the interactions in these places is limited (Jourdain et al.,
55 2020; Schmidt et al., 2023). Accordingly, melt rates generated by the ocean, their calibration based on oceanic conditions taken
outside of ice shelf cavities, and the ice-sheet dynamic response to these oceanic changes are deeply uncertain (Seroussi et al.,
2020).

Quantifying this uncertainty is a difficult task. Ideally, the proper tool to model ice-ocean interactions are coupled ice-ocean models (Gladstone et al., 2020; Park et al., 2023). However these models are computationally expensive and therefore their use for continental-scale ice-sheet studies is rare. An alternative to study the future evolution of the AIS is to use an offline approach by forcing an ice-sheet model with prescribed ocean temperatures, using a basal melt parameterization (e.g., Jourdain et al., 2020). Following this approach, Favier et al. (2019) analyzed the performance of a variety of basal melting laws. Different models use different parameterizations and therefore can lead to different responses of the AIS under the same forcings. The simplest law considers a linear dependency on thermal forcing with constant heat exchange velocity controlling the rate of melting. A further step in complexity is using a quadratic dependency on the thermal forcing. This allows capturing the effect by which melting induces stronger circulation under the shelves with warmer ocean conditions, fostering more melting, resulting in a positive feedback. In the simplest cases, sub-shelf melting parameterizations are local in the sense that basal melting on each point of the ice shelf only depends on the physical properties at that point. Jourdain et al. (2020) developed an extension of the quadratic law that takes into account non-local processes. This quadratic non-local parameterization was the standard parameterization used in the 6th Phase of the Ice-Sheet Modelling Intercomparison Project for CMIP6 (ISMIP6, Jourdain et al., 2020, Nowicki et al., 2020, Seroussi et al., 2020). Using 13 ice-sheets models from different international groups, Seroussi et al. (2020) investigated the evolution of the AIS during the period 2015–2100, showing a wide spread in their results in terms of sea level, covering an interval between -7.8 and 30 cm under the RCP8.5 scenario, with the WAIS adding roughly 18 cm and the East Antarctic Ice Sheet (EAIS) acting with a counterbalance effect gaining ice mass. These results suggest that if the divergence between models and oceanic conditions is significant by 2100, it would be expected to be even greater on longer time-scale simulations.

In fact, as noted by Lowry et al. (2021), the effects of global warming over Antarctica will have consequences in the long term, beyond the 21st century. Hence, an extension of the range period covering the projections in the ISMIP6 project is needed. A few studies have performed such extensions, analyzing different sources of uncertainty to shed light on what could happen in the coming centuries. Precisely, using a statistical emulator and the NorESM1-M model as forcing, Lowry et al. (2021) sampled several parameters relevant for ice-sheet dynamics (the sliding law exponent, the minimum till friction angle and enhancement factors) to estimate the probability distribution of the AIS contribution to sea level by 2300 under the RCP2.6 and RCP8.5 scenarios. The width of the spread in sea-level projections was found to broaden as time surpasses 2100, with an amplitude of several meters in 2300. Bulthuis et al. (2019) showed that the basal sliding law parameters, calving and ocean melt factors are all major sources of uncertainty for projections by the year 3000, with MISI ruling in marine basins. Chambers et al. (2021) also extend their projections until 3000, considering a constant climate after 2100 using different GCMs and ocean conditions, and demonstrating also that uncertainty is inherent to projections. Lipscomb et al. (2021) and Berdahl et al. (2023) made extensions until 2500 (see details in Section 4), but considering a repeated forcing using the two-decadal range 2081-2100 in order to account for climate variability, and an averaged climatology of the period 2090-2100, respectively, with the original ISMIP6-2100 dataset. This leads to an underestimation of the thermal forcing for the period 2100-2500 compared with other ways of extension like the one used in this study (see Table 5). More recently, ~~Seroussi et al. (2024, under review)~~

[Seroussi et al. \(2024\)](#) considered extensions until 2300, on one hand with new GCMs forcings simulations ranging until that same year and, on the other hand, with 2080-2100 conditions.

Here we investigate the sensitivity of our ice-sheet model to the three main sources of uncertainty: future scenarios, climate-
95 model forcings and ice-ocean interactions in stand-alone ice-sheet model simulations by extending the ISMIP6 projections
beyond the 21st century, until 2500 with a similar procedure to Lipscomb et al. (2021) and Berdahl et al. (2023), but using for
our purpose, a subset of the 14-member ensemble of experiments given by the protocol for ISMIP6-2300 (~~Jourdain et al., 2020~~
~~, Nowicki et al., 2020, Seroussi et al., 2020, Seroussi et al., 2024, under review~~)([Seroussi et al., 2024](#)), taking GCMs from the
CMIP5 and CMIP6 ensembles under different scenarios until 2300 extended to 2500 and applying the quadratic non-local
100 parameterization for basal melting in order to study the sensitivity in ice-ocean interactions. Atmospheric and oceanic forcings
are also considered independently in order to detach their effects over the AIS and potentially identify the main driver of the
changes on the ice sheet. Furthermore, we also focus on the timing of the loss of ice in the WAIS in the different simulations.
This work is structured as follows: in Section 2 the ice-sheet model used, the basal melting parameterization chosen and the
setup configuration for the simulations are described, together with the details of the experiments carried out. In Section 3,
105 the results of the experiments are described and particularly analyzed in terms of the sea-level contribution with respect to a
control simulation. In Section 4 we discuss our results in the context of previous research studies and the outlook for future
work, and finally in Section 5 the main conclusions are summarized.

2 Methodology

In this section the ice-sheet model and the basal melting law used in the study together with the experimental setup are
110 described.

2.1 Description of the ice-sheet model

The ice-sheet model used in this study is Yelmo (Robinson et al., 2020). It is a 3D thermomechanical model containing four
main components: topography, material properties, dynamics and thermodynamics. Each of these components correspond to
the calculation of specific state variables. Nevertheless, due to the fact that the equations involving the dynamics in ice are
115 nonlinear and coupled with other parts of the model, these components are in turn tightly coupled with each other internally.
Yelmo is a higher-order model, in other words, it uses methods that take into account longitudinal, lateral and vertical shear
stresses. Recently, it has incorporated one of these methods, known as the depth-integrated effective viscosity approximation
(DIVA), which permits solving the horizontal momentum balance in 2D while maintaining high fidelity to the full Stokes
solution and running fast at higher resolutions (Goldberg, 2011; Robinson et al., 2022). Yelmo has been tested in benchmark
120 experiments such as EISMINT1, EISMINT2 and MISMIP (Robinson et al., 2020). These idealized tests give confidence that
the model performs well for known conditions. It has also recently been used to simulate the AIS (Blasco et al., 2021) and the
Laurentide Ice Sheet at the Last Glacial Maximum (Moreno-Parada et al., 2023).

Regarding resolution, [for this study we have used](#) a regular grid composed of 381 x 381 cells for a horizontal resolution of 16 km and 10 vertical layers ~~has been used for this study~~. Nonetheless, [we have done](#) additional tests with 32-km cells ~~have been made~~ for comparison (see Section 4).

In this study, basal stress τ_b was implemented as a regularized Coulomb power law (Joughin et al., 2019; Schoof, 2005), given in terms of the basal velocity $\mathbf{u}_b = (u_b, v_b)$ as:

$$\tau_b = -c_f N \left(\frac{|\mathbf{u}_b|}{|\mathbf{u}_b| + u_0} \right)^q \frac{\mathbf{u}_b}{|\mathbf{u}_b|} \quad (1)$$

where N is the effective pressure produced by the ice at the base, $q = 0.2$ an exponent and $u_0 = 100$ m/yr an empirical threshold speed (Zoet and Iverson, 2020). c_f is a tunable dimensionless parameter that represents the frictional properties of the bedrock (see Section 2.3). For the effective pressure N , the parameterization by Leguy et al. (2014) is selected

$$N(p) = \rho_i g H \left(1 - \frac{H_f}{H} \right)^p \quad (2)$$

with g being the acceleration due to gravity and H_f the flotation thickness (dependent on ρ_{sw} and bed elevation). The value of the parameter on the exponent p represents the hydrological connectivity of the subglacial drainage system to the ocean and is set to 0.5.

Concerning calving, a von Mises calving law ([Lipscomb et al., 2019](#)) is used with the scaling calving parameter $k_t = 0.0025 \text{ m yr}^{-1} \text{ Pa}^{-1}$ ~~and~~, $w_2 = 25$ for the calving eigenvalue weighting coefficient [and the calving limit in ice thickness is set to 200 m](#), as in the approach by (Lipscomb et al., 2019). For the geothermal heat flow, the reconstruction of Shapiro and Ritzwoller (2004) is used. Furthermore, ~~ELRA—the Elastic Lithosphere Relaxing Asthenosphere (ELRA)~~ is used as GIA model, but in a five-century period the bedrock is not expected to change considerably. Nonetheless, ~~Seroussi et al. (2024, under review)~~ [Seroussi et al. \(2024\)](#) have demonstrated (pers. comm.) that an ELRA model correction in the bedrock elevation could result in differences of around +11-17% per ~ 1.3 m SLE calculated with the volume above flotation method, which should be kept in mind, as we have used this conversion in our study. As Goelzer et al. (2020) pointed out, on short timescales, this conversion can assume a constant area for the ocean when calculating the sea-level contribution.

2.2 Basal melting parameterization

Several studies have suggested that a quadratic melt parameterization is best able to mimic the ice–ocean thermal forcing coupling as compared with other basal melting laws ([Seroussi et al., 2020](#); [Burgard et al., 2022](#)) ([Favier et al., 2019](#); [Burgard et al., 2022](#)). For this reason, [we have chosen](#) the generalized non-local basal melt parameterization described in Jourdain et al. (2020) ~~is chosen~~ in this study. This parameterization takes into account not only the local forcing, but also the average forcing over the cavity beneath a given ice shelf. It translates through γ_0 , the parameter describing the rate at which heat is transferred between ice and ocean, the thermal forcing into basal melting (of floating ice), $B_f(x, y)$ by:

$$B_f(x, y) = \gamma_0 \times \left(\frac{\rho_{sw} c_{pw}}{\rho_i L_f} \right)^2 \times (T_F(x, y, z_{\text{draft}}) + \delta T_{\text{sector}}) \times |\langle T_F \rangle_{\text{draft} \in \text{sector}} + \delta T_{\text{sector}}| \quad (3)$$

where ~~it has been denoted by~~ $T_F(x, y, z_{\text{draft}})$ denotes the thermal forcing at the ice-ocean interface with z_{draft} the ~~thickness~~ depth of the ice shelf base, $\rho_{sw} = 1028 \text{ kg m}^{-3}$ is the density of ocean water, $\rho_i = 918 \text{ kg m}^{-3}$ the density of ice, $L_f = 3.34 \times 10^5$ J kg^{-1} the latent heat of fusion of ice, $c_{pw} = 3974 \text{ J kg}^{-1} \text{K}^{-1}$ the specific heat of the ocean water and δT_{sector} is a temperature correction (thermal forcing correction). The term $\langle T_F \rangle_{\text{draft} \in \text{sector}}$ represents the thermal forcing averaged over all ice shelves contained in each of the 18 drainage basins in which the Antarctic domain is divided (Jourdain et al., 2020). Yelmo includes a marine-shelf interface that allows to interpolate the thermal forcing at the required depths. First, it computes the depth of the ice shelf base by distinguishing between floating and grounded ice. Second, the weights for the different vertical layers are calculated by taking into account the two nearest layers of the depth of the shelf previously calculated in the first step. Finally, the field is computed in terms of these weights as a weighted sum.

Finally, concerning sub-shelf melting in the vicinity of the grounding line, there is a debate as to how strong the latter should be. In the literature two common options are the no melt parameterization (NMP) and the partial melt parameterization (PMP) (Leguy et al., 2021; Seroussi and Morlighem, 2018). Here, we use the PMP parameterization ~~is used~~. In Yelmo, each individual cell of the domain has assigned values between 0 and 1 for the fraction of the area of the cell that is floating ϕ_f and grounded ϕ_g in such a way that $\phi_g = 1 - \phi_f$. Given a value m for basal melting in a partially-grounded cell, the PMP parameterization uses ϕ_f as a weighting factor for m .

2.3 Experimental setup

~~A~~ Here, we performed a subset of five future-projection experiments defined in the protocol for ISMIP6-2300 (~~Jourdain et al., 2020~~, ~~Nowicki et al., 2020~~, ~~Seroussi et al., 2020~~, ~~Seroussi et al., 2024~~, ~~under review~~) is performed here (Table 1) (~~Seroussi et al., 2024~~), namely the extended scenarios to 2300 without repeated forcing or ice-shelf collapse (Table 1). These experiments use different ~~AOGCM models~~ Atmosphere-Ocean General Circulation Models (AOGCMs) from CMIP5 (CCSM4 and HadGEM2) and CMIP6 (CESM2 and UKESM1), Table 1. Regarding the scenario, ~~higher emissions are used in~~ four of the experiments use higher emissions leaving only one experiment with a low emission scenario. In this way, we focus on the structural uncertainties produced by higher emissions. Although in the protocol the ending year is established in 2300, in this study we extend the simulations towards 2500, by forcing from 2300 to 2500 with the average of the last 10-year period (2291-2300) of the experiments in Table 1.

In ISMIP6-2300, there are four experiments considering global warming conditions (from expAE02 to expAE05 in Table 1). Although their atmospheric forcing anomaly shows a similar evolution in time, ranging approximately between 10 and 15 K (Fig. 1a), there is a mismatch in the oceanic forcing observed in CCSM4, which is between 1 and 2 K below the other higher-emission curves. ~~Therefore, the oceanic thermal forcing has an independent reaction compared to the atmospheric forcing. For instance, while UKESM1 and CCSM4 have very similar atmospheric forcings, reaching the end of the period with nearly the same surface temperature anomalies, their oceanic thermal forcings are very different, diverging at the end of the period by nearly 2 K (Fig. 1b).~~

The contrast between the forcings can also be seen in Fig. 1, for the two main regions of the AIS. For the surface temperature anomaly (Fig. 1d), there is no difference in the evolution of the higher-emission scenarios. Meanwhile, for the oceanic

forcing some differences appear, both for fixed (present-day) and evolving grounding lines (Figs. 1e and 1f). With the evolving grounding lines, the WAIS, where the grounding line experiences a retreat, is between 0.25 and 0.75 K warmer than the EAIS on average (Fig. 1f).

190 High-emission scenarios lead to higher surface temperatures all over Antarctica (Fig. 2a). ~~Regions particularly affected are the Ross ice shelf~~ The temperatures over the Ronne-Filchner and Ross ice shelves and the Amundsen Sea ~~, with temperatures rising~~ rise more than 15 K in 2300 with respect to the present day. Also, surface temperature anomalies in the interior of the AIS in CESM2-WACCM surpass 25 K. The low emission case, however, experiences a cooling in some regions, especially in the waters off Victoria Land, the Amundsen and the Weddell seas. Similar differences in thermal forcing are observed in
 195 ice-shelf cavities (Fig. 2b), with increases of nearly 8 K for CESM2 and UKESM1-0-LL in the WAIS for Ronne-Filchner and Ross, but also underneath the EAIS ice shelves (Amery, Shackleton and the Wilkes Land). These changes can also be observed in terms of ~~SMB~~ Surface Mass Balance (SMB) anomaly where in the WAIS, especially the Antarctic Peninsula, and the EAIS predominate the accumulation of ice in the cases of CESM2-WACCM and UKESM1-0-LL with high emissions (Fig. A2).

For each forcing scenario, a range of values for the heat exchange velocity, γ_0 , is considered in the quadratic non-local basal
 200 melting parameterization (Eq. 3) by taking into account the distribution based on circum-Antarctic observations associated with the MeanAnt calibration, proposed by Jourdain et al. (2020) and used in the context of ISMIP6 by Seroussi et al. (2020). The 5th, 50th and 95th percentiles are used yielding an interval ranging from a low to a high value for γ_0 (Table 2). In this way, a 3-member ensemble is obtained for each forcing scenario.

For the initialization of the projections, spin-up simulations running for 20 kyr with a constant reference forcing were carried
 205 out. For the atmospheric fields, ~~Surface Mass Balance (SMB)~~ SMB and 2m-air temperature, we have used monthly values from the Regional Atmospheric Climate Model (RACMO v2.3, Van Wessem et al., 2014) forced by ERA-Interim reanalysis (Dee et al., 2011) averaged over the time period 1981-2010. Regarding the ocean, the fields produced by Jourdain et al. (2020) for ISMIP6 are used. To evaluate the sensitivity with respect to the heat exchange parameter γ_0 , a different spin-up is required for each experiment. During the first 15 kyr of the spin-up, the spatially explicit basal friction coefficient c_f (Eq. 1) is obtained
 210 through an optimization process so that the modeled ice thickness H is as close as possible to the observed present-day state H_{obs} . The observed ~~variables~~ ice thickness and bedrock elevation are taken from the BedMachine Antarctica V2 dataset (Morlighem et al., 2020) and they also represent the initial configuration of the ice sheet before the spin-up is performed. To optimize c_f , the time-evolution differential equation from Lipscomb et al. (2021) is considered:

$$\frac{dc_f}{dt} = -\frac{c_f}{H_0} \left[\frac{H - H_{obs}}{\tau_c} + 2 \frac{dH}{dt} + \frac{H_0}{20} \frac{\log(c_f/c_{f,target})}{\tau_c} \right] \quad (4)$$

215 where $H_0 = 100$ m and $\tau_c = 500$ years are constants for scaling in ice thickness and relaxation time respectively, and $c_{f,target}$ is a target for c_f . The last term ensures that in places where the optimization is less successful, the value of c_f does not saturate to an extreme value (William H. Lipscomb, pers. comm.). $c_{f,target}$ is defined as a function of elevation following Winkelmann et al. (2011). In order to compute the thermal forcing correction, δT_{sector} in Eq. 3, we carried out an equivalent optimization process to the one described for the basal stress constant c_f ~~is carried out~~. A minimum limit was set at -3 K and a maximum at 0.5 K, so
 220 that δT_{sector} values on each sector are consistent with present-day conditions. As pointed out by Lipscomb et al. (2021), this is

Table 1. Description of the five [types of](#) experiments carried out based on the protocol for ISMIP6-2300. The ending year of the forcing in these experiments is 2300 and ice-shelf collapse is not imposed. Note that in this study we extend the simulations towards 2500 by using the average of the last 10-year period (2291-2300) of the experiments.

Experiment	Model	Scenario
ctrlAE	—	—
expAE02	CCSM4	RCP8.5
expAE03	HadGEM2(-ES)	RCP8.5
expAE04	CESM2(-WACCM)	SSP5-8.5
expAE05	UKESM1(-0-LL)	SSP5-8.5
expAE10	UKESM1(-0-LL)	SSP1-2.6

Table 2. Reference names and values of γ_0 used in the experiments. The values are chosen considering the MeanAnt calibration methodology from Jourdain et al. (2020).

Reference name	Percentile	γ_0 (m/yr)
Low	5th	9620
Medium	50th	14500
High	95th	21000

a different perspective to that developed in Jourdain et al. (2020) who consider Antarctica’s basins with fixed thermal forcing values instead. Additional analysis as to how these two optimised parameters affect the initialization of the experiments can be found in Section 4. For the last 5 kyr of the spin-up, the optimised fields are held constant and the simulation is run with the reference climatology towards steady state.

225 A control simulation using the same climate forcing as its spin-up simulation is carried out starting in 2015 and running until 2500. The only difference with respect to the spin-up run is the fact that optimization is not used for neither of the two variables mentioned before. For each value of γ_0 , [we perform](#) a separate spin-up and control simulation ~~is performed~~. The control runs with the different values of γ_0 are all stable during the simulation, with variations of about 5 mm of SLE between 2015 and 2500 in terms of volume above flotation in the three cases. Nevertheless, with increasing values of γ_0 , the ice sheet volume
230 decreases at its initial state (Fig. A3).

2.4 Individual contribution of the atmosphere and the ocean

In order to study the separate effects of the atmospheric and oceanic forcings on the AIS, ~~additional experiments were conducted~~ [we have conducted additional experiments](#): atmosphere-only runs in which the constant oceanic field at 2015 for




Figure 1. Mean evolution of the atmospheric (a), oceanic thermal forcing at the grounding line depth fixed in 2015 (b) and thermal forcing at the evolving grounding lines ~~in the AIS~~ (c) in the AIS for the five ~~experiments carried out~~ models considered for simulation: CCSM4 (RCP8.5), HadGEM2-ES (RCP8.5), CESM2-WACCM (SSP5-8.5), UKESM1-0-LL (SSP5-8.5) and UKESM1-0-LL (SSP1-2.6). For (d), (e) and (f), same as in (a), (b) and (c) respectively but for the WAIS and EAIS. From 2300 to 2500 a constant climate is imposed through the average climatic conditions in the period 2291-2300.

thermal forcing is kept constant during the whole period of simulation and ocean-only simulations with the SMB and surface
 235 temperature fields at 2015 held constant until 2500. These idealized experiments use the same models and scenarios described in Table 1.

2.4 Error analysis in the initialization procedure

In Figure 3, the root mean square error (RMSE) has been computed for ice thickness and ice-surface velocity at the beginning
 of the control run with respect to the observations given by the BedMachine Antarctica V2 dataset (Morlighem et al., 2020) and
 240 the Rignot et al. (2011) dataset, respectively. This RMSE has been ~~computed~~ calculated using two masks, one with the initial
 configuration of the AIS given by our initialization where ice shelves are allowed to expand further than observations, and the
 other with the observations. In the first case, for ice thickness, the deviation from observations is in the range 201 to 216 m,
 decreasing for increasing γ_0 . For ice surface velocity, the corresponding range is between 201 and 275 m/yr. Overall, Yelmo's
~~root-mean-square-errors~~ RMSEs for ice thickness and ice surface velocity are in the medium and upper range, respectively,
 245 of the values obtained in ISMIP6 (see Figure 3 in Seroussi et al., 2020). For the medium value of γ_0 the model overestimates




Figure 2. Surface temperature anomaly (a, in K) between 2300 and the present day as simulated by each of the GCMs selected from ISMIP6-2300 in Table 1, using four high-emission scenarios and a low emission one. Thermal forcing differences over the ice shelves (b, in K) between 2300 and 2015, together with the grounding lines and coastlines at the initial time and the 2300 grounding line.




Figure 3. Root mean square error (RMSE) for ice thickness (in m) and ice surface velocity (in m/yr) between the start of the simulations in 2015 and observations. The different colors represent the values of γ_0 . To mask over the cells containing ice, the ice-sheet margins at the start of the simulation (extended ice shelves) have been used for the columns labelled with (1) and the observations for the ones labelled with (2).

ice thickness, most notably in the ice-sheet margins (Fig. 4a). Only a few regions in the western (Fig. 4c) and eastern sectors show thinner ice than observations. There is a similar pattern in surface velocity anomalies (Fig. 4b), with velocities on the ice shelves generally overestimated. ~~Yet, in, with the exception of the Ronne-Filchner, Ross and Pine Island ice shelves regions, there are ice shelf and~~ a few ice streams showing lower ice surface velocities than observed in the Ross ice shelf and Pine

250 Island (Fig. 4d).




Figure 4. Ice thickness (a, in m) and surface velocity anomalies (b, in m/yr) at 2015 for the medium value of γ_0 , with respect to observations from the BedMachine Antarctica V2 dataset and Rignot et al. (2011) dataset, respectively. Zoom over the WAIS region (c and d) to better appreciate the details at the grounding line.

3 Results

3.1 Sensitivity to climate combined atmospheric-oceanic forcing and γ_0 value

The evolution of the contribution of the AIS to global sea level following the five projections derived from the experiments in Table 1 is shown in Figure 5, together with the spread produced by varying γ_0 . CESM2-WACCM shows the highest long-term contribution to sea level, which is a consequence of this model leading to the maximum forcing both by the atmosphere and the ocean (Fig. 1). For this particular experiment, the medium value of γ_0 yields ca. 3.5 m SLE at 2500, and the difference between using a low or high γ_0 value is close to 1.3 m SLE. Although CCSM4 and the high-emission case for UKESM1-0-LL have similar atmospheric forcings, their contributions by 2500 differ by more than 2 m SLE. In fact, the bounds of the projections given by the different values of γ_0 do not overlap between these two experiments and furthermore the low-emission case is comparable with the CCSM4-driven experiments in the high-emission case. This is explained by the very different oceanic forcing under the AIS ice shelves, as demonstrated by the diverging curves of CCSM4 and the high scenario for UKESM1-0-

255

260




Figure 5. a) Projections of the AIS sea-level contribution (mm SLE) for the five scenarios chosen (Table 1), relative to the control run. In shaded colours the spread resulting from the range of γ_0 values is represented; the solid line corresponds to the medium value. The values in the legend are indicative of the sea-level contribution given by the medium value followed inside the parentheses by the corresponding low and high value. b) Same as a) but for the rate of sea-level contribution (mm SLE/yr).

LL in Figs. 1b and 1c, despite having very similar atmospheric forcing time series. In fact, in the period 2015-2100 the mean thermal forcing at the grounding line increments ~ 1 K for all high-emission scenarios, with CCSM4 and UKESM1-0-LL showing roughly 2 K at 2100. However, between 2100 and 2300, the thermal forcing rate with CCSM4 is approximately 1 K per century, while UKESM1-0-LL doubles that rate. This points to the critical role of the ocean forcing.

Considering the extension from 2300 to 2500 under constant climate forcing, we can see the sea-level contributions slow down in the last two centuries (Fig. 5b). For the medium value of γ_0 , the rate of sea-level contribution for CESM2-WACCM, HadGEM2-ES and UKESM1-0-LL (SSP5-8.5) reaches a peak near the year 2300 with values between 10 to 15 mm SLE/yr. After 2300, these rates go below 10 mm SLE/yr ending up with nearly 5 mm SLE/yr in 2500 with the exception of HadGEM2-ES, close to 10 mm SLE/yr, but no simulation reaches a new equilibrium before the year 2500 even though the forcings are kept constant after the year 2300.

A summary of the results at 2500 in terms of sea level can be found in Figure 6 for each value of the parameter γ_0 , where the effect of increasing basal melting is reflected.




Figure 6. Bar plots for the sea-level contribution (mm SLE) at 2500 for the different GCM forcings from ISMIP6-2300.

Table 3. Total contribution (in mm SLE) and percentage of contribution of the WAIS and the EAIS to the total amount of the AIS volume variations at year 2500 with the five different climate forcings using the medium value of γ_0 in Table 2.

	CCSM4	HadGEM2-ES	CESM2-WACCM	UKESM1-0-LL	UKESM1-0-LL (low)
Total Contribution (mm SLE)	498	3259	3505	2769	61
WAIS (%)	182	74	74	80	290
EAIS (%)	-82	26	26	20	-190

3.1.1 Sensitivity analysis on the WAIS and EAIS

275 ~~On a regional basis~~

Sensitivity analysis on the WAIS and EAIS

Regionally, the WAIS (including the ice sheet over the Antarctic Peninsula, APIS) is the main contributor to the global sea-level rise (Fig. 7), with all simulations showing positive contributions for the medium value of γ_0 and reaching up to more than 2 m in the cases of HadGEM2-ES, CESM2-WACCM and UKESM1-0-LL. For these three forcings, the EAIS also shows a positive contribution to sea level but never exceeding ~ 1 m SLE. The relative contributions of the WAIS (74-80%) and the EAIS (20-26%) are similar across these three models (Table 3), with the WAIS being the major contributor until the year 2500. Meanwhile, for CCSM4 and UKESM1-0-LL (low), the EAIS counterbalances the contribution of the WAIS by lowering sea level through a gain in accumulation.

280

3.2 Spatial patterns in the loss of ice sensitivity to different climatic forcing

285 To gain insight into ~~the reasons behind the behaviour of the time series shown~~ our results, it is useful to compare the ice thickness and ice surface velocity anomalies at the beginning of each century until 2500 with respect to the start of the projections in 2015 for the medium value of γ_0 (Figs. 8 ~~and A4, respectively~~, A4 and A5). The WAIS loses mass in all experiments under high-emission scenarios (Figs. 8 and A5). This can be explained by the warmer waters in the Amundsen and Bellingshausen




Figure 7. Evolution of the sea-level contribution (mm SLE) from the two main regions of Antarctica, the WAIS (including the Antarctic Peninsula, a) and the EAIS (b), relative to the control run, ~~since the start of the simulations in between 2015 to year and 2500.~~

seas sectors (Fig. 2). In addition, as shown in Figure 8, the EAIS also loses ice mass, ~~and the Amery ice shelf serves as an~~
 290 ~~example. The loss of ice takes place~~ particularly in the margins of the ice sheet (~~e.g. the Amery ice shelf~~), while there is gain of
 ice in the interior, ~~especially in the eastern sector~~. At 2100, the experiments with CESM2-WACCM and CCSM4 show a clear
 loss of ice in the Ronne-Filchner and Amery ice shelves. However, for CCSM4 there is a delay in the loss of ice compared with
 the other higher emission experiments in the subsequent centuries. Changes in the position of the grounding line of the Ross
 ice shelf are not evident until 2400 for CCSM4, while for CESM2 the grounding line begins to retreat ~~between 2200 and 2300.~~
 295 ~~prior to 2100~~. Despite the constant climate forcing over the last two centuries, the grounding line keeps retreating in the WAIS
 from 2300 to 2500. The collapse of the ice shelves fringing the WAIS leads to debuttressing of the interior ice flow (Figs. A4
 and A6). Hence nearly all of the grounded ice of the WAIS ends up disappearing in all the experiments with higher emissions
 with the exception of CCSM4.

The loss of ice in the WAIS, reaching anomalies in ice thickness well above 300 m, activates a retreat in the grounding lines,
 300 mainly in the Ronne-Filchner, Ross, Pine Island Glacier and Thwaites Glacier ice shelves, leading to a debuttressing effect that
 amplifies ice flow and permits the ocean to advance towards the interior, until finally the grounding line migrates hundreds of
 kilometers, destabilizing the ice shelves and in turn producing even more loss of ice in those regions. This is in contrast with




Figure 8. Ice thickness anomalies (in m) for the years 2100, 2200, 2300, 2400 and 2500 with respect to the start of the simulations for the set of five experiments carried out with the medium value of γ_0 . The grounding line is represented by black and violet colours for 2500 and 2015 respectively. To show negative anomalies in the WAIS, no masking has been implemented as time evolves, keeping the original coastlines.

the low-emission scenario with UKESM1-0-LL, where the grounding line position remains largely unchanged and even the Ronne-Filchner ice shelf is larger, in the sense that [it](#) is gaining ice with respect to the start of the simulations. However, Pine
 305 Island and the Ross ice shelf are losing mass even under a low-emission forcing. In connection with the loss of ice, there is an acceleration of the ice-stream flow, with the exception of the Ross ice shelf under low emissions. Figure A4 shows how the ice surface velocity is enhanced with respect to 2015. Only some areas of the APIS for HadGEM2-ES, CESM2-WACCM and UKESM1-0-LL with high emissions present a clear slowdown, but for the rest of the AIS, the acceleration is pronounced, reaching anomalies of more than 100 m/yr compared with 2015 by 2200.

The timing of dynamic loss of ice in the WAIS

In order to investigate the sequence of events leading to the general loss of ice, in particular in the WAIS, we assess the evolution of the ice-area extension over this region (Fig. 9a). The total ice area shows a negative trend for all high-emission scenarios, with an abrupt change between 2100 and 2300 that leads to the loss of about 1 million km² in less than 100 years in the cases of HadGEM2-ES, CESM2-WACCM and UKESM1-0-LL. After the sharp ice-area reduction, the ice area continues to decrease at a lower rate. Nevertheless, Figure 9a also shows that this change is directly caused by the change in floating rather than grounded ice, which evolves rather linearly. Yet, from 2300 onward, the floating ice area has almost disappeared, while the grounded area continues losing ice in all high scenarios.

The rate of change of ice volume above flotation in the WAIS is always negative (Fig. 9b). A minimum is reached near the end of the 23rd century. Hence, the rate at which the WAIS loses ice mass (and as a consequence its contribution to sea level) gradually increases until the year 2300, subsequently decreasing. This change in the tendency can be explained by the effect of the constant climatology imposed from 2300 to 2500 and also by the gain in accumulation produced by an increase in the precipitation in a warming climate (Masson-Delmotte et al., 2021). For the EAIS (Fig. 9c), there is also a minimum near 2300 but the rate is much lower in absolute value.

As has been mentioned in Section 3.1, the experiments with CCSM4 show important differences with respect to the other high-emission simulations, in particular CESM2-WACCM, which is the ~~more~~-most extreme example. Figure A7 shows the evolution of ice surface velocity for these two cases since 2100, where the WAIS begins to lose mass until 2300 where the contribution to sea level produced by this region changes its tendency (Fig. 9b). Both experiments differ drastically during the course of the 22nd century. For CESM2-WACCM, both the Ross and Ronne-Filchner ice shelves disappear while for CCSM4 only a retreat of these ice shelves towards the interior is observed. At 2300, once the forcing is set as constant for the remaining 200 years of study, the main shelves finally collapse for CCSM4, while for CESM2-WACCM, the grounding line continues migrating inwards reducing the area in the west (Fig. A7).

3.3 Individual effect of the atmosphere and the ocean

To assess the individual contributions of the atmosphere and ocean, the atmosphere-only and ocean-only runs are ~~assessed~~analyzed. In the first case, the total contribution is less than 1.5 m SLE at the end of 2500 for all models (Fig. 10a). The simulations using UKESM1-0-LL, both with high and low emissions, and CCSM4 actually gain ice mass as a consequence of the enhanced accumulation rates (Fig. 11a) and the absence of ocean melting.

In contrast, for ocean-only runs, all experiments lead to a positive contribution to sea level (Fig. 10b). Uncertainties are roughly 2 m SLE between the low and high values of γ_0 for HadGEM2-ES, CESM2-WACCM and UKESM1-0-LL, with the medium values surpassing 3 m SLE by the end of 2500. For all simulations, the contribution to sea level from the ocean-only runs is above the corresponding run for the atmosphere-only case. Therefore, the ocean is clearly the main driver of the loss of ice and sea-level rise in Antarctica in all our projections. Figure 11a shows that for the atmosphere-only case, the WAIS




Figure 9. Ice area extension (a, in 10^6 km^2) in the WAIS and rate of change in ice volume above flotation (in mm SLE/yr) in the WAIS (b) and EAIS (c) with time for the different GCM models considering the medium value of γ_0 . In a), the grounded area is represented with dashed lines, the floating area with dash-dotted lines and the total area with solid lines.

loses ice as in the general case considering both forcings, but not enough to produce the collapse of the main ice-shelves. The EAIS ice shelves also have their ice mass reduced. In addition, in the interior and eastern areas, there is a gain of ice (Fig. 11a).
 345 Therefore, not considering the effects of the ocean allows the east to gain ice. Regarding the ocean-only case (Fig. 11b), the difference with the start of the simulations is found on the collapse of the main shelves in the WAIS and also in the EAIS. More interestingly, in the interior it is observed that changes are negligible, which can be explained by the fact that maintaining the present-day atmosphere does not lead to increased precipitation in the interior and eastern sectors, so no difference between 2015 and 2500 is found in terms of gain or loss of ice.




Figure 10. Sea-level contribution of the AIS (mm SLE) in the cases where only the forcing of the atmosphere (a) or the ocean (b) are considered. The shading in b) indicates the spread produced by the effect of the values of γ_0 , while in a), as the optimization process leads to similar initial ice sheet states regardless of γ_0 , the ocean fields maintain the present-day state.

350 4 Discussion

We have studied the sensitivity of the AIS to climatic forcing and scenarios on one hand and to the main basal melting parameter on the other with the ice-sheet-shelf model Yelmo. To this end we have chosen an offline approach incorporating the ocean forcing through the basal melting parameterization of Jourdain et al. (2020), applying the PMP parameterization for melting at the grounding line. ~~Based on our analysis of the RMSE for ice thickness and ice surface velocity, we can say that Yelmo is in the same range as other ISMIP6 models.~~

~~We have shown that the higher the value of the heat exchange parameter γ_0 , the larger the future sea-level contribution. For the medium value, the sea-level contribution varies from ~ 0.5 to 3.5 m SLE by 2500 under the effect of high-emission scenarios. The upper-range values correspond to the forcing given by CESM2-WACCM within the SSP5-85 scenario. Moreover, we have shown that higher emission scenarios per se do not automatically lead to a higher sea-level contribution unless the ocean undergoes a strong warming, as illustrated by the simulations using the CCSM4 forcing with a RCP8.5 scenario.~~




Figure 11. Ice-thickness anomalies (in m) for the atmosphere-only (a) and ocean-only (b) cases between the year 2500 and the start of the simulations. Solid lines and dashed lines represent the coastline and grounding line, respectively, at year 2500. Numbers included in the maps represent the total sea level contribution in meters for the medium value of the basal melting parameter.

~~Furthermore, the rate of change of this contribution for higher emission scenarios increases until the end of the 23rd century, when the forcing is kept constant, but sea level continues to rise in the last two centuries at a lower speed.~~

The oceanic forcing is critical, and therefore it is important to represent it correctly. CMIP5 and CMIP6 future projections indicate that Antarctica’s surrounding ocean will warm because of wind-driven circulation changes (they project a poleward
 365 wind shift of $\sim 0.8^\circ$ and a strengthening of $\sim 10\%$ under ~~SSP585~~ SSP5-8.5 by the end of this century), ocean warming far to the north of the coast, and enhanced intrusion of CDW into the cavities under the ice shelves (Purich and England, 2021). However, CMIP6 models do not solve these cavities. Furthermore, mesoscale eddies, which are a source of ocean heat transport onto the shelves, are not well represented under the typical resolution used in CMIP6 models (Bracegirdle et al., 2020). This should improve with the new generation of models for CMIP7 using a higher resolution (Heuzé, 2021).

370 For the case of our research, we have used two models from CMIP6. CESM2-WACCM has strong warm biases, on and off the shelves (Purich and England, 2021). This model is also known to be affected by the representation of cloud feedbacks, which are particularly strong yielding a larger climate sensitivity and therefore an inflated warming (Zhu et al., 2021). UKESM1-0-LL includes positive feedbacks over the Southern Ocean, inducing warming. In fact, for this model 40% of the global heat uptake occurs in the Southern Ocean and 28% of this uptake is transported northward (Williams et al., 2023).

375 The separate effects of the atmosphere and the ocean on the AIS has recently been studied by Coulon et al. (2024) with comparable results to ours. The main driver of future AIS changes will be the ocean, whose interaction with the ice sheet also happens to be the main source of uncertainty. In fact, only the ocean can ~~contribute to sea level by allow~~ for a sea-level increase of more than 3 meters in 2500, affecting the WAIS predominantly. Meanwhile, in the short term, the atmosphere can help to somewhat balance the loss of ice mass through enhanced accumulation of ice. However in the long term it is expected
 380 that it will also contribute to the ice-mass loss through enhanced melting. Using the SICOPOLIS ice-sheet model, the medium value of the basal melting parameter and the original ISMIP6 forcing for 2100, Chambers et al. (2021) presented similar projections but keeping the climate constant from 2100 to 3000, and Greve et al. (2022) proceeded extending the simulations

Table 4. Sea-level contribution (m SLE) at 2300 and 2500 (when data available) among the studies made by Chambers et al. (2021), Greve et al. (2022), Lipscomb et al. (2021) and the results from Yelmo with the medium value of γ_0 .

GCM Model/ Resolution	Chambers et al. (2021)		Greve et al. (2022)	Lipscomb et al. (2021)	Yelmo		Yelmo	
	8 km		8 km	4 km	16 km		32 km	
	2300	2500	2300	2500	2300	2500	2300	2500
CCSM4	0.5	2.1	1.25	0.28	0.1	0.5	0.16	0.6
HadGEM2-ES	1.6	3.6	3.3	0.35	1.5	3.2	1.6	3.5
CESM2	0.45	1.75	0.95	0.26	1.7	3.5	1.9	3.8
UKESM1-0-LL	1	2.6	1.95	0.38	1.2	2.8	1.4	3.1

from 2100 to 2300 with a climate index derived from simulations with the GCM model MIROC under RCP8.5 and RCP4.5 scenarios. Their results for the year 2300 and 2500 are very similar to ours with HadGEM2-ES and UKESM1-0-LL, but with the forcings from CESM2 and CCSM4, there is a clear mismatch (Table 4). Note that for the case of CESM2 in ISMIP6-2300, a different component is used for the atmosphere (WACCM) instead of that used for ISMIP6-2100 (CAM) that could explain this difference ([see comment and references in Section 4](#)). This comparison also makes evident that the procedure used to extend the simulations to the future is an important source of uncertainty.

Lipscomb et al. (2021) considered a very similar procedure to ours (DIVA solver, basal friction law and optimization for thermal forcing, mainly). The main difference consists in the procedure used to extend the simulations until 2500. They ~~use~~ [used](#) the ISMIP6-2100 dataset with a repeated forcing using the two-decadal range 2081-2100 in order to account for climate variability. Instead, we fix the forcing at 2300. As a consequence, their experiments, with the same MeanAnt calibration as ours, contribute just over 400 mm SLE by year 2500, which is a much lower contribution than in our cases (more than 3000 mm SLE in two experiments). Even using the PIGL calibration, based on melt rates on Pine Island Glacier (Jourdain et al., 2020), and which sets a γ_0 value one order of magnitude larger than the MeanAnt, they do not simulate a very high contribution, reaching roughly 1 m SLE by 2500. Therefore, our results indicate that the repeated forcing used in Lipscomb et al. (2021) leads to ~~an underestimation of the AIS response~~ [a smaller AIS response than ours](#). Berdahl et al. (2023) made a similar study to ours ~~but~~, not only varying γ_0 but also the parameter p from Leguy's parameterization, which in our case is constant ($p=0.5$) in all experiments. In addition, except for CESM2, the GCMs they use as forcings are different from our case. For the experiment with CESM2, they found a sea-level rise of approximately 1600 mm SLE by 2500 while in our simulations, it reached more than 3000 mm SLE. This is explained by the same features pointed out for the results of Lipscomb et al. (2021) because they ~~consider~~ [considered](#) an average of the 2090-2100 period to generate the extension towards 2500. Table 5 shows the mean thermal forcing under the ice shelves (between 210 and 810 m of depth, see also Fig. A1) for the ISMIP6-2300 dataset in several periods used to extend the simulations in these studies and ours. Clearly the thermal forcing increases with time, so our setup allows for stronger thermal forcings that using the repeated forcing as in Lipscomb et al. (2021) or the averaged decadal

Table 5. Comparison of the mean thermal forcing ($^{\circ}\text{C}$) from the ISMIP6-2300 dataset on the ice shelves between depth layers of 210 and 810 meters depth, for the different time ranges used to extend the simulations (2081-2100, 2090-2100 and 2291-2300) in Lipscomb et al. (2021), Berdahl et al. (2023) and this work, respectively. Values relative to the reference given by Jourdain et al. (2020).

	CCSM4	HadGEM2-ES	CESM2-WACCM	UKESM1-0-LL	UKESM1-0-LL (low)
Lipscomb et al. (2021) 2081-2100	1.063	1.347	1.309	0.778	0.206
Berdahl et al. (2023) 2090-2100	1.132	1.441	1.453	0.917	0.183
This study 2291-2300	3.397	5.048	4.863	5.494	0.253

period of Berdahl et al. (2023). For instance, for the UKESM1-0-LL case, the differences between our chosen period to extend the forcing and theirs can reach more than 4.5°C . The same happens for CESM2-WACCM and HadGEM2-ES, reaching 3.5°C approximately, and CCSM4 with roughly 2.3°C .

Regarding the sensitivity study on γ_0 described in Berdahl et al. (2023), we reach similar conclusions. When increasing the value of γ_0 , the retreat in the grounding line (especially in the WAIS) is triggered sooner. However, our initial ice-sheet configuration produces larger ice shelves compared to observations. The RMSE of the initial state with respect to observations, is about 200 m/yr and 200 m for ice surface velocity and ice thickness, respectively. Berdahl et al. (2023) obtained lower RMSE values, ~ 129 m/yr and ~ 58 m respectively (for the case they show with $p=0$). Nevertheless, our simulations were also performed with a lower resolution than in these two works with the Community Ice Sheet Model (CISM, Lipscomb et al., 2019) using higher resolutions, but the sensitivity results should still be valid in this case, while the specific numbers could be more uncertain.

We have also tested the same experiments with a 191×191 grid giving a 32-km resolution. Results for sea-level evolution are very similar, varying 7-10% for high-emission scenarios (Fig. A8). Generally, they yield an increase in the contribution but a decrease in the spread between the low and high values of γ_0 . Lipscomb et al. (2021) made the point with 2 and 4 km resolution with sea level variations of $\sim 15\%$ and with 4 and 8 km with $\sim 20\%$, but they find that the higher the resolution, the higher the sea-level contribution. In our case, it is the ~~contrary~~opposite, at least at the resolutions we tested. In the same context, Sutter et al. (2023) argued using the Parallel Ice Sheet Model (PISM; Winkelmann et al., 2011) ~~with~~using 16, 8 and 4 km resolution ~~grids~~ that their 8-km ice-sheet configuration is close to their 16-km runs, ~~nonetheless~~while the rate of retreating at the grounding line is different, starting sooner for 8 and 4 km compared with 16 km, especially ~~observed~~ for Thwaites Glacier. ~~This suggests the MISI instability is triggered earlier when using a higher resolution.~~

A critical point in our study is the initialization, which is based on the optimization during the first 15 kyr of the 20 kyr spin-up run of the thermal forcing correction (δT) and the bed friction coefficient (Section 2). As mentioned before, higher values of γ_0 lead to smaller ice shelves. As a consequence, lower values of γ_0 lead to higher positive values of the thermal forcing correction. To illustrate this phenomenon, the mean value over each drainage basin at the end of the spin-up is indicated in Figure A11a, showing a decrease with increasing γ_0 . For instance, for $\gamma_0=9620$ m/yr the west of the Ronne-Filchner ice

Table 6. Differences (in percentage) in sea level contribution at 2500 for the experiments carried out, between the runs with extended ice shelves and the ones with limited ice shelves.

$\gamma_0(m/yr)$	CCSM4	HadGEM2-ES	CESM2-WACCM	UKESM1-0-LL	UKESM1-0-LL (low)
9620	23.5	-10	0.4	1.2	38
14500	13.5	4.3	1.9	4.7	-48
21000	7.9	3.1	0.9	2.7	-4.9

shelf illustrates the emergence of a positive anomaly, while for $\gamma_0=14500$ and $\gamma_0=21000$ m/yr, the thermal forcing correction is negative. Regarding the bed friction coefficient (Fig. A11b), it does not strongly depend on γ_0 on the EAIS. This is to be expected because the contact between the ice-sheet and ocean is limited in this area. However, the opposite occurs in the WAIS. For the highest γ_0 value, the basal friction coefficient over a wide part of WAIS (particularly around the divide) saturates at a constant value of 1, the imposed internal upper limit in Yelmo (lower limit is 0.001). This increases basal stress in that region, favouring shear-dominated flow and limiting the basal sliding.

Simulating the correct ice extent of ice shelves is a challenging task. Some ice models impose fixed ice-shelf extensions or prevent ice-shelves to grow beyond the observed ice front (Seroussi et al., 2019). In addition, tuning parameters to simulate realistic ice-shelf extensions may differ between different Antarctic embayments (Wilner et al., 2023). In this study, we simulate ice shelves prognostically, which leads to several ice-shelf fronts extended further than observations, particularly the Ronne-Filchner and the Ross ice shelves. For the Amundsen Sea Embayment, the ice shelf advances various kilometers into the ocean and that results resulting in a deceleration of some ice streams flowing towards that sector at the beginning of the simulations (see Fig. 4 for a comparison with observations). Higher values of γ_0 results in a more realistic set up at the beginning (which can be observed in terms of RMSE, Fig. 3), despite having a greater rate of melting. Since ice shelves are allowed to expand, we have done additional simulations considering limited ice shelves with the mask from observations in the BedMachine dataset. In terms of sea-level contribution, results and spreads are very similar for higher-emission scenarios (Fig. A9) with differences ranging between 0.4 and 23.5% in absolute value, but generally being less than 10% (Table 6). In the case of the low-emission simulation this difference is much greater. In general, the prognostic ice-shelf simulations have no considerable effect on the simulated sea level, with a difference below 141 mm SLE at the end of the five-century period for the medium value of γ_0 .

In relation to the two ISMIP6 protocols, we have also tested the goodness of fit of our simulations following ISMIP6-2300 with respect to the simulations made by Seroussi et al. (2020) for ISMIP6-2100, for the period 2015-2100 and the three values of γ_0 (Fig. A10). The spread and mean values produced by the different ice-sheet models ranges approximately between -80 and 260 mm SLE for different forcings and the medium value of γ_0 (Fig. 7 in Seroussi et al., 2020). Our results until the year 2100 fit between the shading produced for the values that we can compare for CMIP5, CCSM4 and HadGEM2-ES (Fig. A10), with both of them in the lower range of the spread. Our model results therefore show a similar low order of magnitude (always less than 100 mm SLE) in comparison to the sea-level contributions estimated in later centuries.

Throughout this work, the WAIS stands out as the main contributor to sea level rise in the future, with some sectors such as the largest ice-shelves, Ronne-Filchner and Ross, and Pine Island Glacier being particularly affected in some experiments (Fig. 8). In these ice shelves, ice thickness is reduced more than 100 m in less than 100 years during the course of the 22nd century for high emission scenarios. Based on our results, we can say that the WAIS has great potential for large sea-level contributions in the future, independent of the associated uncertainty. The rapid WAIS ice loss is due to the debuttressing effect following ice-shelf thinning, which allows the interior ice flow to accelerate (Fig. A4), increasing the discharge. Nonetheless, it has recently been determined that snowfall, which is expected to increase in a warmer climate (Masson-Delmotte et al., 2021), also influences the imbalance in mass loss observed in this well-known region of Antarctica in the last decades, entailing variations in sea-level rise (Davison et al., 2023). The response of the solid Earth beneath Antarctica to the loss of ice, unevenly distributed on the WAIS and EAIS due to mantle properties, influences in long timescales the contribution to sea level as well, due to variations in grounding line locations and the bedrock (Whitehouse et al., 2019). In our simulations, we have used a simplified isostasy module considering constant mantle viscosities for the WAIS and EAIS, that could overestimate and underestimate respectively for each region the contribution to sea level in 2500. This could be better assessed in the future with a more realistic isostasy module that is currently being implemented (FastIsostasy, ~~?~~; ~~under review~~ [Swierczek-Jereczek et al., 2024](#)).

In our simulations for high-emission scenarios, the WAIS seems to start losing ice as soon as the beginning of the 22nd century with an increasing contribution to sea level rise until the end of the 23rd century, when this tendency is balanced with a lower rate despite the continuous loss of ice. Hill et al. (2023) applied a numerical stability analysis to show, using three different ice-sheets models, that under present conditions the AIS is not yet experiencing an irreversible MISI when activating and deactivating perturbations in basal melting. But reaching a certain point of retreat in the positions of the grounding lines, even with the current climate, could trigger an instability in a range of 300-500 years in some regions of the WAIS like the Amundsen Sea Embayment (Reese et al., 2022).

Further work is needed to better understand the structural uncertainty in future AIS mass loss projections. Our study has just focused on the parametric uncertainty that arises within one basal melt parameterization, the quadratic non-local basal melting law of Jourdain et al. (2020). Other basal melting parameterizations include the Potsdam Ice-shelf Cavity mOdel (PICO; Reese et al., 2018). In Burgard et al. (2022), most existing parameterizations have been assessed considering the ice-shelf slope relative to the horizontal too. Also it could be tried another procedure to guess the rates of basal melting and it has been proved in recent times that machine and deep learning have come in handy in that matter (Rosier et al., 2023). Finally, it is important to note that our offline approach can affect projections of the AIS contribution to sea level because of a misrepresentation of the feedbacks between warm water intrusions, basal melting and other coupling processes (Park et al., 2023). In fact, as shown by Golledge et al. (2019), the freshwater fluxes from AIS melting could confine warm water in the subsurface of the Antarctic Ocean leading to higher basal melting, therefore affecting sea-level projections in coupled ocean-ice-sheet models. This meltwater impacts the ocean's overturning circulation, precipitation patterns and temperature variability across the world (Bracegirdle et al., 2020). Eventually these opposing effects should be addressed by the use of coupled ocean-ice models.

490 5 Conclusions

The effect of the uncertainty in the climate forcing in four high-end scenarios and one low-end scenario on the response of AIS to future climate change has been explored in this work. First, we have shown that oceanic forcing is the main driver of mass loss in the WAIS and thus the AIS overall. Thus, simulations driven by similar atmospheric forcing but different oceanic conditions can produce very different sea-level contributions for the future. This is illustrated by the UKESM1-0-LL and CCSM4 GCM models that under a high-emission scenario are associated with contributions of almost 2.7 m and 0.5 m respectively by 2500 due to the very different oceanic forcing. In fact, we demonstrated that a low emission scenario could be comparable to the case of CCSM4. Therefore, at least on short timescales (hundreds of years), a correct representation of the ocean response to future climate change is critical.

We also analyzed the parametric uncertainty arising from the basal melt law used to force our ice-sheet model, the heat exchange velocity γ_0 between ice and ocean on the ice shelves. Our results show that uncertainty in Antarctica projections is profoundly linked to the selection of this parameter and to the effect of the ocean rather than the atmosphere. Differences between choosing a low and a high value could alter sea level more than 2 m in 2500 in some experiments. Nevertheless, this contribution to sea level is unevenly distributed spatially along the ice sheet. The west region is robustly projected to generate a positive contribution to the rise of sea level, due to its marine ice-sheet condition and the physical effect of the buttressing of the main ice shelves. Meanwhile accumulation zones are identified in the stable east part, which gain ice mass, but most notably in some cases even this region has a positive contribution to sea-level rise contributing to the sea level through the weakening of some ice shelves.

Regarding the WAIS, this region experiences a continuous loss of ice starting roughly in the middle of the 22nd century and extending beyond 2300 when most of the ice disappeared including the main ice shelves, Ross and Ronne-Filchner. Furthermore, the main reason for these changes can be attributed directly to the effect that the ocean has over the AIS.

Code availability. The Yelmo ice sheet model code is additionally available at <https://github.com/palma-ice/yelmo> (Robinson et al., 2020)

Data availability. The netCDF files containing all the results from simulations together with a sample of the namelist of parameters used are available at this Zenodo repository: [10.5281/zenodo.10657938](https://zenodo.org/record/10657938)

Appendix A: Additional figures




Figure A1. Depth-averaged thermal forcing under the ice shelves (K) between 210 and 810 meters in (a) 2015 and in (b) 2300.




Figure A2. SMB ($\text{kg m}^{-2} \text{yr}^{-1}$) anomaly for the different GCM models between 2300 and 2015.




Figure A3. Evolution of the ice volume above flotation (m SLE) in the control runs for the AIS, with the different values of γ_0 .




Figure A4. As Figure 8 but for ice surface velocity anomalies (in m/yr). Note that only the cells where there is ice (floating and grounded) at the specific times are represented.




Figure A5. Ice thickness anomalies (in m) in the WAIS (defined as in Fig. 4a) for the years 2100, 2200, 2300, 2400 and 2500 with respect to the start of the simulations for the set of five experiments carried out with the medium value of γ_0 . The grounding line is represented by black and violet colours for 2500 and 2015, respectively. To show negative anomalies in the WAIS, no masking has been implemented as time evolves, keeping the original coastlines.




Figure A6. As Figure A5 but for ice surface velocity anomalies (in m/yr).




Figure A7. Ice surface velocity evolution (in m/yr) at the beginning of each century for the experiments with the GCM models CCSM4 (top) and CESM2-WACCM (bottom). Solid lines represent the coastline and surface elevation levels at the year considered while dashed lines delineate grounding lines.

Sea Level Contribution (mm SLE)




Figure A8. Sea level contribution for the different experiments carried out and spread of the values of γ_0 using resolutions of 16 km (a) and 32 km (b). In c, differences between a and b.

Sea Level Contribution (mm SLE)




Figure A9. Evolution of the sea level contribution until 2500 for the different experiments carried out, together with the spread produced by the different values of γ_0 . In a with the extended ice shelves, in b with the ice shelves limited by a mask from observations in the BedMachine dataset, and in c difference between a and b.




Figure A10. Sea level contribution (mm SLE) for the experiments carried out with five GCM forcings under high-emission scenarios, until the year 2100 for the medium value of γ_0 . Solid lines represent the curve for the values of γ_0 and the shading the overall results from Seroussi et al. (2020) with different ice-sheet models for the forcings given by CCSM4 and HadGEM2-ES.




Figure A11. a) Thermal forcing correction (K) at the beginning of the control simulation for the three values of γ_0 . In grey, basins in which Antarctica is divided and numerical values with the average thermal forcing on each basin at start. b) Bed friction coefficient at the beginning of the simulations. The minimum value is 0.001, set as a limiting parameter in Yelmo.

515 *Author contributions.* A. J-M. carried out the simulations. A. J-M., J. B. and A. R. implemented the ISMIP6 protocol for driving the Yelmo ice-sheet model. All co-authors analyzed the results. A. J-M. prepared the manuscript with contributions and revisions from all co-authors.

Competing interests. At least one of the (co-)authors is a member of the editorial board of The Cryosphere.

Acknowledgements. This research has been supported by the Spanish Ministry of Science and Innovation (project MARINE, grant no. PID2020-117768RB-I00) and by the European Union Horizon 2020 research and innovation programme (grant no. 820970, TiPES). Antonio
520 Juarez-Martinez was funded from the predoctoral grant partnership between the Complutense University of Madrid and Banco Santander (CT58/21-CT59/21). Alexander Robinson received funding from the European Union (ERC, FORCLIMA, 101044247). All simulations were performed in Brigit, the HPC server of the Faculty of Physics of the [UCM Complutense University of Madrid](https://www.ucm.es). We also acknowledge the projects CliC, CMIP5/CMIP6 and ISMIP as being the main sources for our work, and the two reviewers that have contributed with valuable suggestions and revisions to create the final version of this manuscript.

525 **References**

- Bell, R. E. and Seroussi, H.: History, mass loss, structure, and dynamic behavior of the Antarctic Ice Sheet, *Science*, 367, 1321–1325, 2020.
- Berdahl, M., Leguy, G., Lipscomb, W., Urban, N., and Hoffman, M.: Exploring ice sheet model sensitivity to ocean thermal forcing and basal sliding using the Community Ice Sheet Model (CISM), *The Cryosphere*, 17, 1513–1543, <https://doi.org/10.5194/tc-17-1513-2023>, 2023.
- Blasco, J., Alvarez-Solas, J., Robinson, A., and Montoya, M.: Exploring the impact of atmospheric forcing and basal drag on the Antarctic Ice Sheet under Last Glacial Maximum conditions, *The Cryosphere*, 15, 215–231, 2021.
- 530 Bracegirdle, T., Krinner, G., Tonelli, M., Haumann, A., Naughten, K., Rackow, T., Roach, L., and Wainer, I.: Twenty first century changes in Antarctic and Southern Ocean surface climate in CMIP6, *Atmospheric Science Letters*, 21, <https://doi.org/10.1002/asl.984>, 2020.
- Bulthuis, K., Arnst, M., Sun, S., and Pattyn, F.: Uncertainty quantification of the multi-centennial response of the Antarctic ice sheet to climate change, *The Cryosphere*, 13, 1349–1380, <https://doi.org/10.5194/tc-13-1349-2019>, 2019.
- 535 Burgard, C., Jourdain, N., Reese, R., Jenkins, A., and Mathiot, P.: An assessment of basal melt parameterisations for Antarctic ice shelves, *the Cryosphere*, <https://doi.org/10.5194/tc-2022-32>, 2022.
- Chambers, C., Greve, R., Obase, T., Saito, F., and Abe-Ouchi, A.: Mass loss of the Antarctic ice sheet until the year 3000 under a sustained late-21st-century climate, *Journal of Glaciology*, 68, 1–13, <https://doi.org/10.1017/jog.2021.124>, 2021.
- Coulon, V., Klose, A. K., Kittel, C., Edwards, T., Turner, F., Winkelmann, R., and Pattyn, F.: Disentangling the drivers of future Antarctic ice loss with a historically calibrated ice-sheet model, *The Cryosphere*, 18, 653–681, 2024.
- 540 Davison, B., Hogg, A., Rigby, R., Veldhuijsen, S., Wessem, J., Van den Broeke, M., Holland, P., Selley, H., and Dutrieux, P.: Sea level rise from West Antarctic mass loss significantly modified by large snowfall anomalies, *Nature Communications*, 14, <https://doi.org/10.1038/s41467-023-36990-3>, 2023.
- DeConato, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591–597, 2016.
- 545 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, d. P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the royal meteorological society*, 137, 553–597, 2011.
- Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T., Ligtenberg, S. R., van den Broeke, M. R., and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, *Nature*, 502, 89–92, 2013.
- 550 Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E., Smith, C. J., et al.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74–82, 2021.
- Favier, L., Jourdain, N. C., Jenkins, A., Merino, N., Durand, G., Gagliardini, O., Gillet-Chaulet, F., and Mathiot, P.: Assessment of sub-shelf melting parameterisations using the ocean–ice-sheet coupled model NEMO (v3. 6)–Elmer/Ice (v8. 3), *Geoscientific Model Development*, 12, 2255–2283, 2019.
- 555 Fowler, A. and Ng, F.: *Glaciers and Ice Sheets in the Climate System The Karthaus Summer School Lecture Notes: The Karthaus Summer School Lecture Notes*, Springer Nature, ISBN 978-3-030-42582-1, <https://doi.org/10.1007/978-3-030-42584-5>, 2021.
- Fürst, J., Durand, G., Gillet-Chaulet, F., Tavard, L., Rankl, M., Braun, M., and Gagliardini, O.: The safety band of Antarctic ice shelves, *Nature Climate Change*, 6, <https://doi.org/10.1038/nclimate2912>, 2016.
- Gladstone, R., Galton-Fenzi, B., Gwyther, D., Zhou, Q., Hattermann, T., Zhao, C., Jong, L., Xia, Y., Guo, X., Petrakopoulos, K., Zwinger, T., Shapero, D., and Moore, J.: A Framework for Ice Sheet – Ocean Coupling (FISOC) V1.1, *Geoscientific Model Development Discussions*, <https://doi.org/10.5194/gmd-2020-206>, 2020.
- 560

- Goelzer, H., Coulon, V., Pattyn, F., De Boer, B., and Van De Wal, R.: Brief communication: On calculating the sea-level contribution in marine ice-sheet models, *The Cryosphere*, 14, 833–840, 2020.
- Goldberg, D. N.: A variationally derived, depth-integrated approximation to a higher-order glaciological flow model, *Journal of Glaciology*, 57, 157–170, 2011.
- 565 Golledge, N. R., Keller, E. D., Gomez, N., Naughten, K. A., Bernales, J., Trusel, L. D., and Edwards, T. L.: Global environmental consequences of twenty-first-century ice-sheet melt, *Nature*, 566, 65–72, 2019.
- Greve, R., Chambers, C., Obase, T., Saito, F., Chan, W.-L., and Abe-Ouchi, A.: Future projections for the Antarctic ice sheet until the year 2300 with a climate-index method, *Journal of Glaciology*, pp. 1–11, 2022.
- 570 Heuzé, C.: Antarctic bottom water and North Atlantic deep water in CMIP6 models, *Ocean Science*, 17, 59–90, 2021.
- Hill, E. A., Urruty, B., Reese, R., Garbe, J., Gagliardini, O., Durand, G., Gillet-Chaulet, F., Gudmundsson, G. H., Winkelmann, R., Chekki, M., et al.: The stability of present-day Antarctic grounding lines–Part 1: No indication of marine ice sheet instability in the current geometry, *The Cryosphere*, 17, 3739–3759, 2023.
- Holland, D. M., Nicholls, K. W., and Basinski, A.: The Southern Ocean and its interaction with the Antarctic Ice Sheet, *Science*, 367, 1326–1330, 2020.
- 575 Holland, P. R., O’Connor, G. K., Bracegirdle, T. J., Dutrieux, P., Naughten, K. A., Steig, E. J., Schneider, D. P., Jenkins, A., and Smith, J. A.: Anthropogenic and internal drivers of wind changes over the Amundsen Sea, West Antarctica, during the 20th and 21st centuries, *The Cryosphere*, 16, 5085–5105, 2022.
- Joughin, I., Smith, B. E., and Medley, B.: Marine ice sheet collapse potentially under way for the Thwaites Glacier Basin, West Antarctica, *Science*, 344, 735–738, 2014.
- 580 Joughin, I., Smith, B., and Schoof, C.: Regularized Coulomb Friction Laws for Ice Sheet Sliding: Application to Pine Island Glacier, Antarctica, *Geophysical Research Letters*, 46, <https://doi.org/10.1029/2019GL082526>, 2019.
- Joughin, I., Shapero, D., Smith, B., Dutrieux, P., and Barham, M.: Ice-shelf retreat drives recent Pine Island Glacier speedup, *Science Advances*, 7, eabg3080, 2021.
- 585 Jourdain, N. C., Asay-Davis, X., Hattermann, T., Straneo, F., Seroussi, H., Little, C. M., and Nowicki, S.: A protocol for calculating basal melt rates in the ISMIP6 Antarctic ice sheet projections, *The Cryosphere*, 14, 3111–3134, 2020.
- Leguy, G., Asay-Davis, X., and Lipscomb, W.: Parameterization of basal friction near grounding lines in a one-dimensional ice sheet model, *The Cryosphere*, 8, <https://doi.org/10.5194/tc-8-1239-2014>, 2014.
- Leguy, G. R., Lipscomb, W. H., and Asay-Davis, X. S.: Marine ice sheet experiments with the Community Ice Sheet Model, *The Cryosphere*, 15, 3229–3253, 2021.
- 590 Lipscomb, W. H., Price, S. F., Hoffman, M. J., Leguy, G. R., Bennett, A. R., Bradley, S. L., Evans, K. J., Fyke, J. G., Kennedy, J. H., Perego, M., et al.: Description and evaluation of the Community Ice Sheet Model (CISM) v2. 1, *Geoscientific Model Development*, 12, 387–424, 2019.
- Lipscomb, W. H., Leguy, G. R., Jourdain, N. C., Asay-Davis, X., Seroussi, H., and Nowicki, S.: ISMIP6-based projections of ocean-forced Antarctic Ice Sheet evolution using the Community Ice Sheet Model, *The Cryosphere*, 15, 633–661, 2021.
- 595 Lowry, D. P., Krapp, M., Golledge, N. R., and Alevropoulos-Borrill, A.: The influence of emissions scenarios on future Antarctic ice loss is unlikely to emerge this century, *Communications Earth & Environment*, 2, 221, 2021.
- Masson-Delmotte, V., Zhai, P., Pirani, S., Connors, C., Péan, S., Berger, N., Caud, Y., Chen, L., Goldfarb, M., and Scheel Monteiro, P. M.: IPCC, 2021: Summary for Policymakers. in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the*

- 600 Sixth Assessment report of the intergovernmental panel on climate change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.
- Moreno-Parada, D., Alvarez-Solas, J., Blasco, J., Montoya, M., and Robinson, A.: Simulating the Laurentide Ice Sheet of the Last Glacial Maximum, *The Cryosphere*, 17, 2139–2156, 2023.
- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., et al.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature geoscience*, 13, 132–137, 2020.
- 605 Nowicki, S., Goelzer, H., Seroussi, H., Payne, A. J., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Alexander, P., Asay-Davis, X. S., Barthel, A., et al.: Experimental protocol for sea level projections from ISMIP6 stand-alone ice sheet models, *The Cryosphere*, 14, 2331–2368, 2020.
- Paolo, F. S., Fricker, H. A., and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348, 327–331, 2015.
- 610 Park, J.-Y., Schloesser, F., Timmermann, A., Choudhury, D., Lee, J.-Y., and Nellikattil, A. B.: Future sea-level projections with a coupled atmosphere-ocean-ice-sheet model, *Nature Communications*, 14, 636, 2023.
- Pattyn, F. and Morlighem, M.: The uncertain future of the Antarctic Ice Sheet, *Science*, 367, 1331–1335, 2020.
- Purich, A. and England, M. H.: Historical and future projected warming of Antarctic Shelf Bottom Water in CMIP6 models, *Geophysical Research Letters*, 48, e2021GL092752, 2021.
- 615 Reese, R., Albrecht, T., Mengel, M., Asay-Davis, X., and Winkelmann, R.: Antarctic sub-shelf melt rates via PICO, *The Cryosphere*, 12, 1969–1985, 2018.
- Reese, R., Garbe, J., Hill, E. A., Urruty, B., Naughten, K. A., Gagliardini, O., Durand, G., Gillet-Chaulet, F., Chandler, D., Langebroek, P. M., et al.: The stability of present-day Antarctic grounding lines–Part B: Possible commitment of regional collapse under current climate., *Cryosphere Discussions*, 2022.
- 620 Rignot, E., Mouginot, J., and Scheuchl, B.: Ice flow of the Antarctic ice sheet, *Science*, 333, 1427–1430, 2011.
- Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-shelf melting around Antarctica, *Science*, 341, 266–270, 2013.
- Ritz, C., Edwards, T. L., Durand, G., Payne, A. J., Peyaud, V., and Hindmarsh, R. C.: Potential sea-level rise from Antarctic ice-sheet instability constrained by observations, *Nature*, 528, 115–118, 2015.
- Robel, A. A., Seroussi, H., and Roe, G. H.: Marine ice sheet instability amplifies and skews uncertainty in projections of future sea-level rise, *Proceedings of the National Academy of Sciences*, 116, 14887–14892, 2019.
- 625 Robinson, A., Alvarez-Solas, J., Montoya, M., Goelzer, H., Greve, R., and Ritz, C.: Description and validation of the ice-sheet model Yelmo (version 1.0), *Geoscientific Model Development*, 13, 2805–2823, 2020.
- Robinson, A., Goldberg, D., and Lipscomb, W. H.: A comparison of the stability and performance of depth-integrated ice-dynamics solvers, *The Cryosphere*, 16, 689–709, 2022.
- 630 Rosier, S. H., Bull, C., Woo, W. L., and Gudmundsson, G. H.: Predicting ocean-induced ice-shelf melt rates using deep learning, *The Cryosphere*, 17, 499–518, 2023.
- Schmidt, B. E., Washam, P., Davis, P. E., Nicholls, K. W., Holland, D. M., Lawrence, J. D., Riverman, K. L., Smith, J. A., Spears, A., Dichek, D., et al.: Heterogeneous melting near the Thwaites Glacier grounding line, *Nature*, 614, 471–478, 2023.
- Schoof, C.: The effect of cavitation on glacier sliding, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 635, 461, 609–627, 2005.
- Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *Journal of Geophysical Research: Earth Surface*, 112, 2007.

- Seroussi, H. and Morlighem, M.: Representation of basal melting at the grounding line in ice flow models, *The Cryosphere*, 12, 3085–3096, 2018.
- 640 Seroussi, H., Nowicki, S., Simon, E., Abe-Ouchi, A., Albrecht, T., Brondex, J., Cornford, S., Dumas, C., Gillet-Chaulet, F., Goelzer, H., et al.: initMIP-Antarctica: an ice sheet model initialization experiment of ISMIP6, *The Cryosphere*, 13, 1441–1471, 2019.
- Seroussi, H., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel, A., et al.: ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century, *The Cryosphere*, 14, 3033–3070, 2020.
- 645 Seroussi, H. et al.: Evolution of the Antarctic Ice Sheet over the next three centuries from an ISMIP6 model ensemble, *Earth’s Future*, accepted, 2024.
- Shapiro, N. M. and Ritzwoller, M. H.: Inferring surface heat flux distributions guided by a global seismic model: particular application to Antarctica, *Earth and Planetary Science Letters*, 223, 213–224, 2004.
- Sun, S., Pattyn, F., Simon, E. G., Albrecht, T., Cornford, S., Calov, R., Dumas, C., Gillet-Chaulet, F., Goelzer, H., Gollledge, N. R., et al.: Antarctic ice sheet response to sudden and sustained ice-shelf collapse (ABUMIP), *Journal of Glaciology*, 66, 891–904, 2020.
- 650 Sutter, J., Jones, A., Frölicher, T., Wirths, C., and Stocker, T.: Climate intervention on a high-emissions pathway could delay but not prevent West Antarctic Ice Sheet demise, *Nature Climate Change*, 13, 951–960, 2023.
- Swierczek-Jereczek, J., Montoya, M., Latychev, K., Robinson, A., Alvarez-Solas, J., and Mitrovica, J.: FastIsostasy v1. 0—a regional, accelerated 2D glacial isostatic adjustment (GIA) model accounting for the lateral variability of the solid Earth, *Geoscientific Model Development*, 655 17, 5263–5290, 2024.
- van de Wal, R. S., Nicholls, R. J., Behar, D., McInnes, K., Stammer, D., Lowe, J. A., Church, J. A., DeConto, R., Fettweis, X., Goelzer, H., et al.: A High-End Estimate of Sea Level Rise for Practitioners, *Earth’s future*, 10, e2022EF002 751, 2022.
- Van Wessem, J., Reijmer, C., Morlighem, M., Mougnot, J., Rignot, E., Medley, B., Joughin, I., Wouters, B., Depoorter, M., Bamber, J., et al.: Improved representation of East Antarctic surface mass balance in a regional atmospheric climate model, *Journal of Glaciology*, 60, 660 761–770, 2014.
- Weertman, J.: Stability of the junction of an ice sheet and an ice shelf, *Journal of Glaciology*, 13, 3–11, 1974.
- Whitehouse, P. L., Gomez, N., King, M. A., and Wiens, D. A.: Solid Earth change and the evolution of the Antarctic Ice Sheet, *Nature communications*, 10, 503, 2019.
- Williams, R. G., Ceppi, P., Roussenov, V., Katavouta, A., and Meijers, A. J.: The role of the Southern Ocean in the global climate response to carbon emissions, *Philosophical Transactions of the Royal Society A*, 381, 20220 062, 2023.
- 665 Wilner, J. A., Morlighem, M., and Cheng, G.: Evaluation of four calving laws for Antarctic ice shelves, *The Cryosphere Discussions*, 2023, 1–19, 2023.
- Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueller, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK)—Part 1: Model description, *The Cryosphere*, 5, 715–726, 2011.
- 670 Zhu, J., Otto-Bliesner, B. L., Brady, E. C., Poulsen, C. J., Tierney, J. E., Lofverstrom, M., and DiNezio, P.: Assessment of equilibrium climate sensitivity of the Community Earth System Model version 2 through simulation of the Last Glacial Maximum, *Geophysical Research Letters*, 48, e2020GL091 220, 2021.
- Zoet, L. K. and Iverson, N. R.: A slip law for glaciers on deformable beds, *Science*, 368, 76–78, 2020.